CUX2 deficiency causes facilitation of excitatory synaptic transmission onto hippocampus and increased seizure susceptibility to kainate

Toshimitsu Suzuki[1,2]†, Tetsuya Tatsukawa[2]†, Genki Sudo[2]†, Caroline Delandre[3,18], Yun Jin Pai[3], Hiroyuki Miyamoto[2], Matthieu Raveau[2], Atsushi Shimohata[2,19], Iori Ohmori[4], Shin-ichiro Hamano[5], Kazuhiro Haginoya[6], Mitsugu Uematsu[7], Yukitoshi Takahashi[8,9], Masafumi Morimoto[10], Shinji Fujimoto[11,12], Hitoshi Osaka[13], Hirokazu Oguni[14], Makiko Osawa[14], Atsushi Ishii[15], Shinichi Hirose[15], Sunao Kaneko[16,17], Yushi Inoue[9], Adrian Walton Moore[3], & Kazuhiro Yamakawa[1,2]*

[1]Department of Neurodevelopmental Disorder Genetics, Institute of Brain Science, Nagoya City University Graduate School of Medical Science, Aichi, Japan.

[2]Laboratory for Neurogenetics, RIKEN Center for Brain Science, Saitama, Japan.

[3]Laboratory for Genetic Control of Neuronal Architecture, RIKEN Center for Brain Science, Saitama, Japan.

[4]Department of Special Needs Education, Okayama University Graduate School of Education, Okayama, Japan.

[5]Division of Neurology, Saitama Children's Medical Center, Saitama, Japan.

[6]Department of Pediatric Neurology, Miyagi Children's Hospital, Sendai, Japan.

[7]Department of Pediatrics, Tohoku University School of Medicine, Sendai, Japan.

[8]Department of Pediatrics, Gifu Prefectural Gifu Hospital, Gifu, Japan.

[9]National Epilepsy Center, NHO Shizuoka Institute of Epilepsy and Neurological Disorder, Shizuoka, Japan.

[10]Department of Pediatrics, Kyoto Prefectural University of Medicine, Kyoto, Japan.

[11]Department of Pediatrics, Neonatology and Congenital Disorders, Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan.

[12]Tsutsujigaoka Children's Clinic, Aichi, Japan.

[13]Department of Pediatrics, Jichi Medical University, Shimotsuke, Japan.

[14]Department of Pediatrics, Tokyo Women's Medical University, Tokyo, Japan.

[15]Department of Pediatrics, School of Medicine and Research Institute for the Molecular Pathomechanisms of Epilepsy, Fukuoka University, Fukuoka, Japan.

[16]Department of Neuropsychiatry, Hirosaki University School of Medicine, Hirosaki, Japan.

[17]North Tohoku Epilepsy Center, Minato Hospital, Hachinohe, Japan.

[18]Present address: Menzies Institute for Medical Research, University of Tasmania, Tasmania,
Australia.

[19]Present address: Department of Physiology, Nippon Medical School, Tokyo, Japan.

†These authors contributed equally to this work.

* Address correspondence to: Kazuhiro Yamakawa, Ph.D.,

Department of Neurodevelopmental Disorder Genetics, Institute of Brain Science, Nagoya City

University Graduate School of Medical Science,

1 Kawasumi, Mizuho-cho, Mizuho-ku Nagoya, Aichi 467-8601, Japan

Tel: +81-52-851-5612, Email: yamakawa@med.nagoya-cu.ac.jp

ORCID iD: https://orcid.org/0000-0002-1478-4390

1   **Keywords**: *CUX2*, *CUX1*, *CASP*, temporal lobe epilepsy, seizure, kainate

3   **ORCID number**: Toshimitsu Suzuki: https://orcid.org/0000-0003-0585-5692

4   Tetsuya Tatsukawa: https://orcid.org/0000-0001-8533-4576

5   Iori Ohmori: https://orcid.org/0000-0001-6823-2762

6   Yukitoshi Takahashi: https://orcid.org/0000-0002-1922-8967

7   Shinichi Hirose: https://orcid.org/0000-0001-6796-3790

8   Adrian Walton Moore: https://orcid.org/0000-0002-8242-301X

9   Kazuhiro Yamakawa: https://orcid.org/0000-0002-1478-4390

Abstract

CUX2 gene encodes a transcription factor that controls neuronal proliferation, dendrite branching

and synapse formation, locating at the epilepsy-associated chromosomal region 12q24 that we

previously identified by a genome-wide association study (GWAS) in Japanese population. A

CUX2 recurrent *de novo* variant p.E590K has been described in patients with rare epileptic

encephalopathies and the gene is a candidate for the locus, however the mutation may not be

enough to generate the genome-wide significance in the GWAS and whether *CUX2* variants

appear in other types of epilepsies and physiopathological mechanisms are remained to be

investigated. Here in this study, we conducted targeted sequencings of *CUX2,* a paralog *CUX1*

and its short isoform *CASP* harboring a unique C-terminus on 271 Japanese patients with a

variety of epilepsies, and found that multiple *CUX2* missense variants, other than the p.E590K,

and some *CASP* variants including a deletion, predominantly appeared in patients with temporal

lobe epilepsy (TLE). The CUX2 variants showed abnormal localization in human cell culture

analysis. While wild-type CUX2 enhances dendritic arborization in fly neurons, the effect was

compromised by some of the variants. *Cux2*- and *Casp*-specific knockout mice both showed high

susceptibility to kainate, increased excitatory cell number in the entorhinal cortex, and significant

enhancement in glutamatergic synaptic transmission to the hippocampus. CASP and CUX2

proteins physiologically bound to each other and co-expressed in excitatory neurons in brain

regions including the entorhinal cortex. These results suggest that *CUX2* and *CASP* variants

contribute to the TLE pathology through a facilitation of excitatory synaptic transmission from

entorhinal cortex to hippocampus.

Introduction

Cux2 gene encodes a homeobox transcription factor CUX2 that is predominantly expressed in progenitor cells of the subventricular zone in mouse embryos and pyramidal neurons of the upper neocortical layers (II–IV) in adult mice[1]. CUX2 is also expressed in Reelin-positive neurons distributed throughout the layers II–IV in postnatal day 0 (P0) mice[2]. CUX2 controls neuronal proliferation, dendrite branching, spine morphology and synapse formation[3,4]. We recently reported a genome-wide association study (GWAS) on 1,825 Japanese patients with variable epilepsies which identified an associated region at chromosome 12q24.11 – 12q24.13 harboring 24 transcripts including *CUX2* gene[5]. In the region, *CUX2* is only gene which has been reported to be relevant for epilepsy; a recurrent *de novo* variant (c.1768G>A, p.E590K) has been identified in patients with rare epileptic encephalopathies (EEs)[6,7]. *CUX2* is therefore one of the most promising candidate genes in this 12q24 epilepsy-associated region, but the mutation reported in rare EEs may not be enough to explain the association detected in the Japanese GWAS study.

To investigate whether *CUX2* and its paralogues' mutations are involved in other types of epilepsies, here we performed targeted sequencing of *CUX2*, its paralog *CUX1,* and *CASP* which is a short isoform of *CUX1* with a unique C-terminus, in Japanese patients with variable epilepsies including genetic generalized and structural/metabolic epilepsies, and identified their variants predominantly in patients with temporal lobe epilepsy (TLE), the most common but intractable form of epilepsy[8]. Analyses in human cultured cell and transgenic fly showed that the variants have loss-of-function effects. CUX2 and CASP deficiencies in mice increased their seizure susceptibilities to a convulsant, kainate, which has long been used to generate TLE animal models[9]. Histological and electrophysiological analyses revealed increases of excitatory

1   neuron numbers in entorhinal cortex and those in excitatory input to hippocampus in both mice,

2   proposing a circuit mechanism for the pathology of TLE.

Materials and Methods

Subjects

Genomic DNAs from 271 Japanese patients with a variety of epilepsies (Table S1) and 311

healthy Japanese volunteers recruited by cooperating hospitals were used for the targeted

sequencing analyses for *CUX2*, CUX1 and *CASP*. For the frequency calculation of c.3847G>A

(p.E1283K) variant in *CUX2* gene, additional DNA samples from independent 69 Japanese

patients with TLE from 2 additional independent facilities were further recruited (Table S1). The

patients' DNAs analyzed in our GWAS[5] were not used in the present study, because their

epilepsy subtype information [TLE, etc.] were not available for those materials.

Targeted sequencing

Genomic DNAs were extracted from peripheral venous blood samples using QIAamp DNA

Blood Midi Kit (Qiagen). Genomic DNA samples were amplified with the illustra GenomiPhi

V2 DNA Amplification Kit (GE Healthcare). We designed PCR primers to amplify all coding

regions of *CUX2* (NM_015267), *CUX1* (NM_001202543 and first exon of NM_181552), unique

C-terminus region of *CASP* (NM_001913), and amplified genomic DNA by PCR using the

PrimeSTAR HS DNA Polymerase (TaKaRa) or KOD-plus Ver. 2 (TOYOBO). The PCR

products were purified using ExoSAP-IT PCR product Cleanup (Thermo Fisher Scientific) and

analyzed by direct sequencing using the ABI PRISM 3730xl Genetic Analyzer. All novel

variants identified in amplified DNA by GenomiPhi were verified by direct-sequencing of

patients' genomic DNAs. Primer sequences and PCR conditions are available upon request.

Quantification of mRNA – described in the Supplemental Methods.

Domain search

Domain searches in CUX2, CUX1, and CASP amino acid sequences were performed using the SMART database.

Expression constructs and mutagenesis – described in the Supplemental Methods.

Cell imaging – described in the Supplemental Methods.

Drosophila stocks and crosses – described in the Supplemental Methods.

TUNEL assay in flies – described in the Supplemental Methods.

Mice

Cux2 knockout (KO) mouse was obtained from Texas A&M Institute for Genomic Medicine (TIGM) as cryopreserved sperm of heterozygous *Cux2* gene trap mouse (129/SvEv × C57BL/6 background) derived from the gene trapped clone OST440231. Live animals were produced by *in vitro* fertilization at the Research Resources Division (RRD) of the Institute of Physical and Chemical Research (RIKEN) Center for Brain Science. The heterozygous mice were maintained on the C57BL/6J (B6J) background, and the resultant heterozygous mice were interbred to obtain wild-type (WT), heterozygous, and homozygous mice. Genotyping was carried out as described previously[10].

Casp-specific KO mice were generated using the CRISPR/Cas system. Plasmid vector pX330-U6-Chimeric_BB-CBh-hSpCas9 was a gift from Dr. Feng Zhang (Addgene plasmid # 42230). A pair of oligo DNAs corresponding to *Casp*-gRNA (TTTCCATCATCTCCAGCCAA AGG) in exon 17 of *Casp* (NM_198602) was annealed and ligated into pX330-U6-Chimeric_BB-CBh-hSpCas9. For knock-out mouse production, Cas9 mRNAs and *Casp*-gRNA were diluted to 10 ng/µL each. Further, B6J female mice and ICR mouse strains were used as embryo donors and foster mothers, respectively. Genomic DNA from founder mice was extracted, and PCR was performed using gene-specific primers (CRISPR_check_F: 5'-

GGAGCTATTGTAGGACATCACAGA-3' and CRISPR_check_R: 5'-

CCCCAGTGTTCTTTACTTTGAGTT-3'). PCR products were purified using ExoSAP-IT PCR

product Cleanup and analyzed by direct sequencing using the ABI PRISM 3730xl Genetic

Analyzer. The heterozygous mutant mice (c.1514^1515ins.TT, p.S506fs) were crossbred with

B6J mice, and the resultant heterozygous mice were interbred to obtain WT, heterozygous, and

homozygous mutant mice. The frame-shift mutation was confirmed by sequence analysis of

cDNA from mutant mouse brains.

Seizure susceptibility in mice – described in the Supplemental Methods.

CUX2 antibody generation

To generate a rabbit polyclonal antibody to CUX2, a fusion protein was prepared, in which the

glutathione-S-transferase (GST) protein was fused to the a.a. residues 356 to 415 of mouse

CUX2 which has been used in the previous study's antibody generation[11]. Rabbits were injected

with 0.5 mg of purified GST fusion protein in Freund's complete adjuvant, boosted five times

with 0.25 mg of protein, and serum collection at 1 week following the last boost. Polyclonal

antibody was purified by affinity chromatography. The serum was passed through a GST affinity

column ten times, and the flow-through was then applied to a GST-CUX2 (356-415 a.a.) affinity

column to isolate antibodies.

Histological analyses – described in the Supplemental Methods.

In vitro electrophysiology – described in the Supplemental Methods.

Co-immunoprecipitation – described in the Supplemental Methods.

Statistical analysis

In the *in vitro* and *in vivo* experiments, data are presented as Box-and-whisker plots or means ±

s.e.m. The boxes show median, 25th and 75th percentiles, and whiskers represent minimum and

maximum values. P-value for p.E1283K was calculated using the Cochrane-Armitage trend test. One-way or two-way ANOVA, Tukey's multiple comparison test, Chi-square test, or Kolmogorov-Smirnov test were used to assess the data as mentioned in the figure legends. Statistical significance was defined as $P < 0.05$.

Study approval

Human study: The experimental protocols were approved by the Ethical Committee of RIKEN and by the participating hospitals and universities. All human study experimental procedures were performed in accordance with the guidelines of the Ethical Committee of the RIKEN and with the Declaration of Helsinki. Written informed consents were obtained from all individuals and/or their families in compliance with the relevant Japanese regulations.

Animal study: All animal experimental protocols were approved by the Animal Experiment Committee of RIKEN. All animal breeding and experimental procedures were performed in accordance with the ARRIVE guidelines and the guidelines of the Animal Experiment Committee of the RIKEN.

Data availability

All data generated or analyzed during this study are included in this published article and its Supplementary Information File.

Results

CUX2 variants predominantly appeared in Japanese TLE patients

We carried out a targeted sequencing of *CUX2* in 271 Japanese patients with variable epilepsies consisting of 116 genetic generalized epilepsies and 155 structural/metabolic ones (Table S1). Structural/metabolic epilepsy samples contained 68 TLEs, which were further divided to 57 mesial TLE (mTLE) and 11 lateral TLE (lTLE). We identified five *CUX2* heterozygous missense variants in nine unrelated patients (Figure 1A, Table 1 and Supplemental Note). Notably, eight of the nine patients with *CUX2* variants had TLE (one lTLE and seven mTLE patients). All of patients carrying *CUX2* missense variants belonged to the subgroup of structural/metabolic epilepsy. None of patients with genetic generalized epilepsies showed *CUX2* variants except for silent variants. All of the mTLE patients showed hippocampal sclerosis. Three (p.R34W, p.P454L, and p.W958R) out of the five variants were absent or rare (< 0.5%) in the in-house Japanese control individuals (in-house controls) and databases and were also predicted to be damaging (Table 1). The p.E1283K variant, a frequent variant predicted to be less damaging, appeared in Japanese TLE patients at a significantly high ratio [$P = 5.93 \times 10^{-3}$, OR = 6.94, 95% CI = 1.39–34.61 calculated in 137 (above-mentioned 68 + additional independent 69; Table S1) TLE patients vs 311 in-house controls] and therefore we hypothesized it may also be a genetic contributor for TLE. The p.D337N variant appeared in one case with TLE and controls with a similar allele frequency.

Loss-of-function effects of *CUX2* variants

To investigate the functional impacts of *CUX2* variants appeared in patients with epilepsy (Figure 1A, Table 1), we transfected HeLa.S3 cells with expression constructs of wild-type (WT)

and the five variants (p.R34W, p.D337N, p.P454L, p.W958R, and p.E1283K). We calculated

two parameters of abnormality, "leakage to cytoplasm" and "abnormal puncta" (Figure 1B, C and

Figure S1A). Although some but not all variants showed abnormalities in each parameter, the

combined data reached statistical significance except for p.R34W.

CUX2 is an ortholog of *Drosophila melanogaster cut,* which promotes dendritic arbor

morphological complexity[12]. We generated transgenic fly lines to express human WT CUX2 or

variants (p.R34W, p.D337N, p.P454L, p.W958R, and p.E1283K) and analyzed their dendritic

arbor morphology in *Drosophila* larvae (Figure 1D-G and Figure S1B-D). Similar to its

Drosophila orthologue[12], ectopic expression of CUX2 WT strongly increased dendritic arbor

complexity (branch number and length). However, activities to drive arbor complexity in the

variants were significantly decreased, except for p.R34W and p.D337N. RT-qPCR assays in the

adult transgenic flies revealed that expression levels of the *CUX2* variants, p.P454L and

p.W958R, were significantly lower (Figure 1H). All *CUX2* constructs were inserted into the same

genomic site, therefore the lower expression levels of transgenes are not likely to be due to

position effects but most likely due to these variants because these are only the differences in the

constructs used for the analyses of fly. TUNEL of adult fly brains showed that the alleles did not

promote apoptotic cell death (Figure S1E). Together, these observations suggest that the *CUX2*

variants present in patients with epilepsy cause loss-of-function of the protein.

***Cux2*-deficient mice show increased susceptibility to kainate**

Because of the loss-of-function nature of epilepsy-associated *CUX2* variants, we next

investigated *Cux2*-KO mice[10]. The body weight of 2-month-old mice was comparable among

genotypes (Figure S2A). In electrocorticogram analysis, the median of the poly spike and wave

discharges frequency was slightly higher in the primary somatosensory cortex forelimb region of

Cux2(-/-) than WT mice, however the difference did not reach statistical significance (data not

shown). No obvious epileptic behaviors or changes in local field potential recordings in the

hippocampus were noted in *Cux2*(+/-) or *Cux2*(-/-) mice. Although patients with TLE often have

past histories of febrile seizures[13], *Cux2*-KO mice did not show any seizure susceptibility to

increased body temperature (data not shown). Seizure susceptibility to pentylenetetrazole (PTZ),

a GABA-A receptor antagonist, remained unchanged in *Cux2*-KO mice (Figure S2B-F).

Importantly, however, *Cux2*-KO mice had a high susceptibility to kainate, which is commonly

used to generate TLE animal models[9], in frequencies of generalized convulsive seizures (GS)

(Figure 2A) and lethality (Figure 2B). The latencies to onset of GS and death were also

significantly decreased in *Cux2*(-/-) mice (Figure S2G, H). Seizure severity was also significantly

higher in *Cux2*-KO female mice (Figure 2C). These results support the notion that *CUX2* loss-of-

function mutations cause TLE.

Cux2-deficient mice show increased cell number in entorhinal cortex and glutamatergic

input to hippocampus

      Cux2(-/-) mice have been reported to show overgrowth of the neocortical upper layers[3]. In

a Nissl staining, we also found a significantly increased cell number in entorhinal cortex layers

II-III, which projects to hippocampal dentate granule cells and CA3 pyramidal cells (Figure 2D).

In the slice-patch recordings, we further found that perforant path-evoked excitatory postsynaptic

currents (eEPSCs) in the dentate granule cells were significantly higher in *Cux2*(-/-) mice (Figure

2E), indicating that glutamatergic synaptic transmission from the entorhinal cortex layers II–III

onto the hippocampus was significantly facilitated in *Cux2*-KO mice.

At a glance, hippocampal structures in *Cux2*-KO mice were comparable to 2- and 10-

month-old WT mice (Figure 2D, Figure S3). We generated an anti-CUX2 antibody similarly to a

previous study[11] and confirmed the presence of CUX2 immunosignals in WT and absence in

Cux2(-/-) mice (Figure S3A, B). CUX2 immunosignals were dense in the neocortical upper

layers (II-IV) as previously reported[1] and also dense in the entorhinal cortex upper layers (II-III)

(Figure S3C). In WT hippocampus, we only observed CUX2 immunolabeling signals in

inhibitory interneurons, specifically somatostatin (SST)-positive, reelin (RLN)-positive and

parvalbumin (PV)-positive inhibitory, but not in excitatory neurons (Table S2, Figure S4). We

found that there were no significant differences in interneuron cell numbers between genotypes

(Figure S5A-D). Timm staining and immunohistochemistry for c-Fos (Figure S5E, F),

Doublecortin, phospho-Histone H3, Ki67, NeuN, GFAP, and ZnT-3 (data not shown) did not

show differences in the hippocampus between genotypes. There are five subtypes of kainate

receptors (KARs), GLUK1–GLUK5, in primates and rodents. We investigated KARs expression

in the hippocampus of 2-month-old *Cux2*-KO female mice. RT-qPCR assays revealed that the

expression of *GluK1* (formerly named *GluR5*) was significantly higher in *Cux2*(-/-) mice (Figure

2F), which is presumably a homeostatic compensatory reaction to epileptic seizures (see

Discussion). The baseline frequencies of spontaneous inhibitory postsynaptic currents (sIPSCs)

in hippocampal dentate granule cells were significantly higher at 6–7 weeks old, and this

difference was suppressed after bath-application of GYKI and AP5, which are AMPA and

NMDA receptor antagonists, respectively (Figure 2G). Frequency of sIPSC in dentate PV-

positive interneurons was also increased in *Cux2*-KO mice (Figure S6). These results suggest

that, in *Cux2*(-/-) mice, the function of hippocampal inhibitory neurons remained intact, but the

increased excitatory input from the entorhinal cortex to the hippocampus could facilitate firing

activities of inhibitory neurons, which itself would also be a compensatory action to epileptic

activities in mice. In CA3 pyramidal neurons of *Cux2*-KO mice, EPSCs were not significantly

affected (Figure S6), suggesting that the increased excitatory input in the upstream dentate

granule cells may be neutralized by the increased inhibitory input in those cells.

Taken together, these results suggest that the increase in entorhinal cortex layers II–III cell

numbers and the resultant facilitation of glutamatergic synaptic transmission from the entorhinal

cortex layers II–III onto hippocampi are causal factors leading to the increased susceptibility to

kainate of *Cux2*-KO mice. Increases in GLUK1 and facilitated firing of inhibitory neurons in the

mouse hippocampus would rather be compensatory reactions.

***CASP* variants in TLE patients**

CUX1 is a paralog of *CUX2*, and *CASP* is an alternatively spliced short isoform of *CUX1*

harboring a unique C-terminus[14] (Figure 3A, Figure S7). CUX1 and CUX2 proteins have four

DNA binding domains (three CUT repeats and one homeodomain), but CASP lacks all of these

domains and instead contains a transmembrane domain. We performed targeted sequencing

analyses of *CUX1* and *CASP* in the 271 Japanese patients with epilepsy and identified nine

nonsynonymous variants (Figure 3A, Table 1, and Supplemental Note). Among those, one

variant in *CUX1*, c.4172C>T (p.T1391I) and three variants in *CASP*, c.1433C>T (p.A478V),

c.1524delG (p.R509fs), and c.1868_1870delTCT (p.F623del), were completely absent or very

rare in in-house controls and databases (Table 1). No other truncation variants of the *CASP*-

specific sequence were found in these databases, suggesting that *CUX1* and especially *CASP*

variants contribute to epilepsy. Although epilepsies observed in patients with *CUX1* and *CASP*

variants were rather heterogeneous, CASP-p.G563S and p.F623del variants appeared in mTLE

and lTLE patients, respectively (Table 1). The mTLE patient SIZ-060 showed hippocampal

sclerosis (Supplemental Note).

CASP and CUX2 proteins are co-expressed in excitatory neurons of entorhinal cortex

upper layer and physiologically bind to each other

Immunohistochemistry with CUX1 antibodies in 2-month-old WT mice revealed CUX1

immunosignals in excitatory neurons at the neocortical upper layers (II-IV), as previously

reported[3], and those at the entorhinal cortex upper layers (II-III) (Figure 3B), similar to CUX2

(Figure S3C). In contrast in the hippocampus, CUX1 was expressed in SST-positive, RLN-

positive, and PV-positive interneurons, but not in excitatory neurons (Figure 3B and Figure S8),

similar to CUX2 (Figure S4). Using a CASP-specific antibody recognizing 400–650 a.a., we

found that CASP was rather widely expressed in neurons of multiple brain regions, but still

dense in the neocortical and entorhinal cortex upper layers, similar to CUX1 and CUX2 (Figure

3C). In the hippocampus, CASP was dense in hilar and stratum-oriens SST-positive cells that

expressed CUX2 (Figure 3C, D), and more specifically, within the cytoplasm (Figure S9), which

is consistent with CASP expression in the Golgi apparatus[15].

A protein interaction between CUX1 and CASP has been previously reported[14]. Here we

newly found that the CASP protein physically interacts with CUX2 (Figures S10 and S11). All

three tested CUX2 rare variants bound to CASP, and all three tested CASP rare variants bound to

CUX2 (Figures S10 and S11), suggesting that the variants did not affect protein binding between

CASP and CUX2.

***Casp*-deficient mice also show increased cell number in entorhinal cortex and**

glutamatergic input to hippocampus

It has been reported that the number of cortical neurons was significantly increased in

Cux1(-/-); *Cux2*(-/-) double-mutant mice, but this increase was no greater than that in the *Cux2*(-

/-) single mutant; therefore, regulation of the upper layer neuronal number was assumed to be a

unique function of CUX2 and not redundant with CUX1 activities[3]. Because of the low

survivability of *Cux1*-KO mice[16] and our observation that the TLE variants appeared in the

CASP-unique sequence but not in *CUX1* itself, we decided to investigate *Casp*-specific KO

rather than *Cux1*-KO mice for analysis. We generated a *Casp*-specific KO mouse by targeting

exon 17 at the unique C-terminus (Figure S7C). *Casp*(+/-) and *Casp*(-/-) mutant mice were born

at a Mendelian ratio, grew normally, and were fertile. RT-qPCR analyses revealed that the *Casp*

mRNA became half and diminished in *Casp*(+/-) and *Casp*(-/-) mice, respectively, whereas *Cux1*

and *Cux2* mRNA levels remained unchanged (Figure 3E). CASP immunosignals well

disappeared in *Casp*(-/-) mice (Figure S9A), confirming the specificity of the CASP antibody. At

a glance, there were no abnormal localizations and intensities of CUX1 and CUX2 proteins in

Casp-KO mice (Figure S9B, C). The median body weight was comparable among genotypes at 2

months of age (Figure S9D). RT-qPCR assays of KARs mRNA in the hippocampi of 2-month-

old *Casp*-KO mice did not show significant change in KARs expression levels (Figure S9E).

In a Nissl staining of 2-month-old *Casp*-KO mice, no increase of neuron number was

observed in the entorhinal cortex (Figure S9F). However, immunohistochemical staining using

the anti-CUX1 antibody as a marker of neurons at upper layers of the neocortical (II–IV) and the

entorhinal cortex (II–III) showed a tendency of increase in both the neocortex and entorhinal

cortex ($P = 7.57 \times 10^{-2}$ and $P = 4.50 \times 10^{-1}$, respectively) (Figure3F). Furthermore, *Casp*-KO

mice also showed high susceptibility to kainate (Figure 3G, H, Figure S9G-I). After

intraperitoneal application of kainate, a larger number of *Casp*-KO mice showed GS and lethality

(Figure 3G, H). Onset latencies of GS and death were significantly decreased in *Casp*-KO mice

(Figure S9H, I). Seizure severity in *Casp*(-/-) mice was also significantly higher (Figure S9G).

The differences in seizure susceptibility to kainate were seen mainly in male *Casp*-KO mice

(Figure S9), contrary to *Cux2*-KO mice in which the susceptibility is higher in female (Figure

S2). Notably again, perforant path-evoked EPSCs (eEPSCs) in the dentate granule cells were

significantly higher in *Casp*-KO mice (Figure 3I), which is similar to *Cux2*-KO mice (Figure

2E).

All of these observations propose that facilitation of glutamatergic synaptic transmission

from the entorhinal cortex onto hippocampal dentate granule cells is a common mechanism for

TLE caused by *CUX2* and *CASP* variants.

Discussion

In this study, we performed targeted sequencing analyses of *CUX2, CUX1* and *CASP* on 271 Japanese patients with a variety of epilepsies, and found that *CUX2* missense variants predominantly appear in TLE patients, in that eight of 68 TLE patients (12%) had *CUX2* variants. Three variants (p.R34W, p.P454L, and p.W958R) are quite rare or even absent in various databases, and are consistently predicted to have a damaging effect, therefore these would be regarded as causal or large-effect susceptibility variants. Although the p.E1283K is a high-frequent, relatively common variant, its frequency in TLE patients is statistically higher compared with in-house controls and therefore would potentially be a genetic contributor for TLE. *CASP* variants also appeared in two TLE patients. All of these patients with *CUX* family variants showed symptoms of epileptic seizures, suggesting that the variants may contribute to the threshold for the triggering epileptic seizures through a facilitation of excitatory synaptic transmission from entorhinal cortex to hippocampus in epilepsies caused by *CUX* family variants. Our recent GWAS analysis of Japanese patients with variable epilepsies identified a region with genome-wide significance at chromosome 12q24 which harbors *CUX2*[5]. Although a recurrent *de novo CUX2* variant p.E590K has been described in patients with EEs[6,7], a previous whole exome sequencing study for Japanese patients with EEs[17] did not find the *CUX2* pathogenic variant. Therefore, the *CUX2* recurrent variant in EEs may not be enough to explain the genome-wide association with epilepsy at the 12q24 region in Japanese population. In our GWAS study[5], sub-analyses for subtypes of epilepsies further revealed that a polymorphic marker at the 12q24 epilepsy-associated region showed genome-wide significant association with structural/metabolic epilepsy. Hippocampal sclerosis is the major entity for

structural/metabolic epilepsy, and therefore the *CUX2* variants in patients with TLE would

contribute to the association with epilepsy at 12q24 in Japanese population.

Human cell culture and fly dendritic arborization analyses revealed loss-of-function effects

of the *CUX2* variants, which were found in TLE patients. *CASP* also showed variants in epilepsy

patients including TLE at the unique C-terminus and we further found that the CASP physically

binds to CUX2. Although all tested CASP variants did not affect the binding activity to CUX2,

the CASP protein has been reported to play a role in intra-Golgi retrograde transport[18] and

therefore the variants in CASP may still affect the subcellular transport or protein modification

of CUX2.

Cux2- and *Casp*-KO mice did not show spontaneous seizures but showed significantly

elevated seizure susceptibility to kainate, an agent which has been used to establish TLE animal

models[9]. We previously reported a nonsense mutation of the *KCND2* gene encoding a voltage-

gated potassium channel Kv4.2 in a patient with TLE[19]. Similar to *Cux2*- and *Casp*-KO mice,

Kcnd2-KO mice did not show spontaneous epileptic seizures but showed increased susceptibility

to kainate in seizure and mortality[20]. These observations suggest that increased seizure

susceptibility to kainate correlates with the threshold for triggering epileptic seizures. However,

other additional as-yet unknown modifying, genetic, or environmental factors may influence full

expression of TLE symptoms including hippocampal sclerosis.

Cux2-KO mice showed a significant and *Casp*-KO mice showed a tendency of, increases

in entorhinal cortex layer II–III excitatory neuron cell number. Although *Casp*-KO mice did not

show any significant changes in *Cux2* mRNA expression levels and histological or cytological

distributions of CUX2 protein, co-expression of CASP and CUX2 proteins in neurons including

entorhinal cortex projection neurons, and the physiological interaction between CASP and CUX2

proteins still suggest that CASP deficiency may impair CUX2 function through an as-yet

unknown mechanism, consequently leading to increased entorhinal cortex excitatory neuron cell

number in *Casp*-KO mice. Furthermore, both *Cux2*- and *Casp*-KO mice revealed significant

increases in perforant path-evoked EPSCs in dentate granule cells. These results suggest that

facilitation of glutamatergic synaptic transmission from the entorhinal cortex onto hippocampal

dentate granule cells is a causal basis for the significant increase in seizure susceptibilities to

kainate in *Cux2*- and *Casp*-KO mice. In contrast, the observed changes in the hippocampus of

Cux2-KO mice are assumed to be homeostatic compensatory reactions to the epileptic causal

changes. In the hippocampus of WT mice, CUX2 immunolabelling was only observed in

inhibitory interneurons such as SST-positive, RLN-positive, and PV-positive inhibitory, but not

in excitatory neurons. In the *Cux2*-KO mice, although no changes were observed in interneuron

cell numbers, a significant increase in sIPSC frequency was observed in dentate granule cells

similar to patients with TLE[21], which would presumably be a compensatory reaction[22,23,24] to the

increased excitatory input from the entorhinal cortex to the hippocampus. The increased GLUK1

expression in the hippocampus of *Cux2*-KO mice may also be suppressive for epileptic seizures

because *GluK1* is expressed in inhibitory neurons[25], and GLUK1 expression has been assumed to

be protective at least for kainate-induced epilepsy[26]. Taken together, the changes within the

hippocampus of *Cux2*-KO mice may be homeostatic compensatory responses rather than causal

actions to epileptic seizures, and these changes in the hippocampus themselves also support the

occurrence of epileptic causal changes in these mice.

 In summary, our results of mutation analyses of *CUX* family genes in patients with

epilepsies including TLE and the functional and mouse model analyses suggest that *CUX* family

gene deficiency is one of the bases for TLE and that increase of cell number in the entorhinal

1 cortex projection neurons and resultant increase of glutamatergic synaptic transmission to

2 hippocampus is a possible pathological mechanism for TLE. Further investigations using mouse

3 models with heterozygous missense variants, which have been identified in TLE patients are

4 required to clarify whether the variants are true loss-of-function mutations and contribute to the

5 TLE pathology.

References

1. Zimmer, C., Tiveron, M. C., Bodmer, R. & Cremer, H. Dynamics of Cux2 expression suggests that an early pool of SVZ precursors is fated to become upper cortical layer neurons. *Cereb. Cortex* **14**, 1408–1420 (2004).

2. Cubelos, B. *et al*. Cux-1 and Cux-2 control the development of Reelin expressing cortical interneurons. *Dev. Neurobiol.* **68**, 917–925 (2008).

3. Cubelos, B. *et al*. Cux-2 controls the proliferation of neuronal intermediate precursors of the cortical subventricular zone. *Cereb. Cortex* **18**, 1758–1770 (2008).

4. Cubelos, B. *et al*. Cux1 and Cux2 regulate dendritic branching, spine morphology, and synapses of the upper layer neurons of the cortex. *Neuron* **66**, 523–535 (2010).

5. Suzuki, T. *et al*. Genome-wide association study of epilepsy in Japanese population identified an associated region at chromosome 12q24. *Epilepsia* **62**, 1391-1400 (2021).

6. Chatron, N. *et al.* The epilepsy phenotypic spectrum associated with a recurrent CUX2 variant. *Ann. Neurol*. **83**, 926–934 (2018).

7. Barington, M., Risom, L., Ek, J., Uldall, P. & Ostergaard, E. A recurrent de novo CUX2 missense variant associated with intellectual disability, seizures, and autism spectrum disorder. *Eur. J. Hum. Genet.* **26**, 1388–1391 (2018).

8. Téllez-Zenteno, J. F. & Hernández-Ronquillo, L. A review of the epidemiology of temporal lobe epilepsy. *Epilepsy Res. Treat*. 630853 (2012).

9. Lévesque, M. & Avoli, M. The kainic acid model of temporal lobe epilepsy. *Neurosci. Biobehav. Rev.* **37**, 2887–2899 (2013).

10. Iulianella, A., Sharma, M., Durnin, M., Vanden Heuvel, G. B. & Trainor, P. A. Cux2 (Cutl2) integrates neural progenitor development with cell-cycle progression during spinal cord

neurogenesis. *Development* **135**, 729–741 (2008).

11. Gingras, H., Cases, O., Krasilnikova, M., Bérubé, G. & Nepveu, A. Biochemical characterization of the mammalian Cux2 protein. *Gene* **344**, 273–285 (2005).

12. Jinushi-Nakao, S. *et al*. Knot/Collier and cut control different aspects of dendrite cytoskeleton and synergize to define final arbor shape. *Neuron* **56**, 963–978 (2007).

13. Maher, J. & McLachlan, R. S. Febrile convulsions. Is seizure duration the most important predictor of temporal lobe epilepsy? *Brain* **118**, 1521–1528 (1995).

14. Lievens, P. M., Tufarelli, C., Donady, J. J., Stagg, A., Neufeld, E. J. CASP, a novel, highly conserved alternative-splicing product of the CDP/cut/cux gene, lacks cut-repeat and homeo DNA-binding domains, and interacts with full-length CDP in vitro. *Gene* **197**, 73–81 (1997).

15. Gillingham, A. K., Pfeifer, A. C. & Munro, S. CASP, the alternatively spliced product of the gene encoding the CCAAT-displacement protein transcription factor, is a Golgi membrane protein related to giantin. *Mol. Biol. Cell* **13**, 3761–3774 (2002).

16. Luong, M. X. *et al*. Genetic ablation of the CDP/Cux protein C terminus results in hair cycle defects and reduced male fertility. *Mol. Cell Biol*. **22**, 1424–1437 (2002).

17. Takata, A. *et al*. Comprehensive analysis of coding variants highlights genetic complexity in developmental and epileptic encephalopathy. *Nat. Commun.***10**, 2506 (2019).

18. Malsam, J., Satoh, A., Pelletier, L. & Warren, G. Golgin tethers define subpopulations of COPI vesicles. *Science* **307**,1095–1098 (2005).

19. Singh, B. *et al.* A Kv4.2 truncation mutation in a patient with temporal lobe epilepsy. *Neurobiol. Dis*. **24**, 245–253 (2006).

20. Barnwell, L. F. *et al*. Kv4.2 knockout mice demonstrate increased susceptibility to convulsant stimulation. *Epilepsia* **50**, 1741–1751 (2009).

21. Li, J. M. *et al.* Aberrant glutamate receptor 5 expression in temporal lobe epilepsy lesions. *Brain Res.* **1311**, 166–174 (2010).

22. Marksteiner, J., Ortler, M., Bellmann, R. & Sperk, G. Neuropeptide Y biosynthesis is markedly induced in mossy fibers during temporal lobe epilepsy of the rat. *Neurosci Lett* **112**, 143-148.

23. Tonder, N., Kragh, J., Finsen, B. R., Bolwig, T. G. & Zimmer, J. Kindling induces transient changes in neuronal expression of somatostatin, neuropeptide Y, and calbindin in adult rat hippocampus and fascia dentata. *Epilepsia* **35**, 1299-1308.

24. Vezzani, A., Sperk, G. & Colmers, W. F. Neuropeptide Y: emerging evidence for a functional role in seizure modulation. *Trends Neurosci* **22**, 25-30.

25. Bureau, I., Bischoff, S., Heinemann, S. F. & Mulle, C. Kainate receptor-mediated responses in the CA1 field of wild-type and GluR6-deficient mice. *J. Neurosci.* **19**, 653–663 (1999).

26. Fritsch, B., Reis, J., Gasior, M., Kaminski, R. M. & Rogawski, M. A. Role of GluK1 kainate receptors in seizures, epileptic discharges, and epileptogenesis. *J. Neurosci.* **34**, 5765–5775 (2014).

Acknowledgments

We thank the patients, families, volunteers, Lab for Genetic Control of Neuronal Architecture, Lab for Neurogenetics, Research Resources Division, Center for Brain Science, RIKEN. The authors also thank Dr. A. Miyawaki for the mRFP plasmid. This work was partly supported by grants from the Center for Brain Science and President's Discretionary Fund, RIKEN, and the Japan Agency for Medical Research and Development (AMED) under Grant Number JP20ek0109388.

Author contributions

TS and KY designed the experiments; TS, TT, GS, CD, YJP and MR performed statistical analyses; TS, IO, SH, KH, MU, YT, MM, SF, H. Osaka, H. Oguni, MO, AI, SH, SK, YI and KY managed DNA samples; TS, IO, SH, KH, MU, YT, MM, SF, H. Osaka, H. Oguni, MO, AI, SH, SK, YI and KY recruited case and control samples; TS and KY performed targeted sequencing analyses; TS, CD, YJP, AWM, and KY performed functional analyses; TS, GS, TT, HM, AS, and KY performed mouse analyses; and TS, RM and KY wrote the manuscript.

Additional information

Competing Interests

The authors declare no competing interests.

Table Legend

Table 1: *CUX2*, *CUX1*, and *CASP* gene nonsynonymous variants in patients with epilepsy.

mTLE: mesial temporal lobe epilepsy, lTLE: lateral temporal lobe epilepsy, JME: juvenile

myoclonic epilepsy, CAE: childhood absence epilepsy, SGE: symptomatic generalized epilepsy,

FLE: frontal lobe epilepsy, GEFS: generalized epilepsy with febrile seizure plus, M: Male, F:

Female, JP: in-house Japanese control individuals, J-HGVD: Japanese Human Genetic Variation

Database, EVS: Exome Variant Server NHLBI GO Exome Sequencing Project, 1kGP: The 1000

Genomes Project, ExAC: Exome Aggregation Consortium, gnomAD: Genome Aggregation

Database. ++: Disease causing, Probably damaging, Deleterious, or Damaging, +: Possibly

damaging, -: Polymorphism, Benign, Tolerated, or Neutral, NA: not available, NT: not tested,

NR: not registered. The CUX2 reference sequence (NM_015267) has an error at c.4414, and the

correct nucleotide is C. CUX2 nucleotide change c.4414G>C (p.V1472L) (rs6490073 in dbSNP,

NCBI) was observed in all sequences in databases and in our subjects, suggesting that the CUX2

reference sequence (NM_015267) has an error at this position (the correct nucleotide is

c.4414C). * Informations are not described because of more than 100 individuals.

Figure Legends

Figure 1. Loss-of-function effects of TLE variants in *CUX2*. (A) CUX2 protein structure

(NP_056082) with variants appeared in patients with epilepsy. (**B**) Abnormal subcellular

localization of CUX2 variant proteins. CUX2-WT protein (arrows) was limited to, but well

distributed within, nuclei stained with DAPI (cyan), whereas variants showed abnormal

aggregates in nuclei (W958R) or leaked-out to the cytoplasm (E1283K) (arrowheads). Scale bars

= 20 μm. (**C**) Ratio of abnormally localized CUX2 proteins (> 200 cells counted). n = WT: 545,

R34W: 282, D337N: 363, P454L: 239, W958R: 565, and E1283K: 323 cells. (**D-H**) CUX2 WT

accelerated arborization of fly neurons and TLE variants lowered its activity and expression.

Representative images of neurons without CUX2 (D), WT control (E), and W958R (F). Scale

bars = 50 μm. (G) Shortened dendrite length in transgenic fly with mutants (n = 11–25 neurons

per genotypes) and (H) lowered expression of mutants (n = 6). One-way ANOVA Tukey's

multiple comparison test (C, G, H). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Figure 2. Increased kainate susceptibility, entorhinal cortical cell number, and excitatory

input to hippocampal granule cells in *Cux2*-KO mice. (A-C) Seizure-related events in mice

after intraperitoneal injection of kainate (KA). Ratio of animals exhibiting generalized

convulsive seizure (GS) (A), mortality rate (B), and seizure severity scores (C) was significantly

higher in *Cux2*(-/-) female and combined gender mice. **(D)** Number of entorhinal cortex layer II-

III excitatory neurons was significantly increased in *Cux2*(-/-) mice (2-month-old). Scale bar =

100 μm. **(E)** Slice-patch analyses showed that perforant path-evoked EPSCs in dentate granule

cells were significantly increased in *Cux2*(-/-) female. **(F)** RT-qPCR analyses revealed that

GluK1 mRNA was significantly increased in *Cux2*(-/-) mice. **(G)** Basal frequency of sIPSC in

dentate granule cells of *Cux2*(-/-) female was significantly increased, and it was suppressed with

subsequent applications of antagonists for AMPA receptor (GYKI) and NMDA receptor (AP5).

Kainate (KA) increased the sIPSC frequency, which was then suppressed by the GABA-A

receptor antagonist picrtoxin. DG; dentate gyrus, Ent; entorhinal area. Yates' correction after

Pearson's Chi-square (A, B), one-way ANOVA Tukey's test (C, F, G), one-way ANOVA (D), or

two-way ANOVA Tukey's test (E). n: mouse numbers. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Figure 3. *CASP* variants in epileptic patients, CASP distribution, and increases in kainate susceptibility and excitatory input to hippocampal granule cells in *Casp*-KO mice. (A) Locations of *CUX1* and *CASP* variants in patients with epilepsy (see Table 1). Dashed lines define the common region. **(B)** CUX1 immunosignals (brown) in neocortical and entorhinal cortex upper layer excitatory neurons and hippocampal interneurons. **(C)** CASP (brown) expressed more widely in neurons, and intensely expressed in neocortical and entorhinal cortex upper layer excitatory neurons. **(D)** In hippocampus, CASP (brown) was dense in SST-positive (blue) interneurons at hilus and stratum oriens (arrows). d2-d5; magnified images outlined in d1. Scale bars = 100 μm (B, C and d1), 20 μm (d2-d5). so; stratum oriens, sp; stratum pyramidale, sg; stratum granulosum, h; hilus. **(E)** RT-qPCR analyses revealed that *Casp* mRNA was decreased, while *Cux1* and *Cux2* mRNAs remained unchanged, in *Casp*-KO mice. **(F)** Thickness of the CUX1-positive neocortical layer (left), density of CUX1-positive cell in neocortex (middle), and CUX1-positive cell density in entorhinal cortex (right). CUX1-positive cell density tended increase at neocortex and entorhinal cortex in *Casp*(-/-) mice (2-month-old) but not statistically significant. **(G, H)** *Casp*-KO mice showed significantly higher susceptibility to kainate in seizure rate (G), mortality (H). **(I)** Perforant path-evoked EPSCs in dentate granule cells were significantly increased in *Casp*(-/-) male (6–7-week-old). One-way ANOVA Tukey's test (E), Yates' correction after Pearson's Chi-square (G, H), or two-way ANOVA Tukey's test (I). n: mouse numbers. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

 Supplementary Figure Legends

 Figure S1. CUX2 mutants show abnormal subcellular localizations in human. cultured cells

 and decreased branching effects but no apoptosis in fly. (**A**) In HeLa.S3 cells, CUX2 mutant

 proteins (R34W, D337N, and P454L) show aggregates or abnormal leakage-out into cytoplasm

 (arrowheads) and some WT-like distribution (arrows). Nuclei were stained with DAPI (cyan).

 (**B-D**) Mutations lowered the neurite arborization activity of CUX2 in fly neurons.

 Representative images of CUX2-R34W (B) and CUX2-P454L (C). Terminal point numbers were

 significant decreased in CUX2-P454L, CUX2-W958R and E1283K (n=13 ~ 25) (D). Results of

 one-way ANOVA followed by Tukey's test. (**E**) TUNEL assay revealed no apoptosis in adult

 transgenic flies of CUX2-WT and mutants. Scale bars = 20 μm (A) or 50 μm (B, C, and E). * P

 < 0.05, ** $P < 0.01$, *** $P < 0.001$.

 Figure S2. *Cux2*-**deficient mice show seizure susceptibility to kainate but not to PTZ.** (**A**)

 Body weight was similar among genotypes of 2-month-old mice. (**B-F**) Unchanged seizure

 susceptibility of *Cux2*-KO mice to PTZ. There are no significant differences in latency to

 generalized convulsive seizure (GS) (B), latency to death (C), percentage of animals exhibiting

 GS (D), mortality rate (E), and seizure severity score (F) among genotypes. (**G, H**) *Cux2*-KO

 mice show increased seizure susceptibility to kainate. Latency to onset of GS (G) and that to

 death (H) were significantly decreased in *Cux2*(-/-) female and combined gender mice. One-way

 ANOVA (A-C, F-H) or Pearson's Chi-square test (3×2 contingency table) (D, E). mean

(horizontal bars) ± s.e.m. (B, C, G, H). n or numbers in round brackets =mouse numbers. * $P <$

0.05, ** $P < 0.01$.

Figure S3. CUX2 antibody specifically recognizes CUX2 protein. (**A**) Sagittal brain sections

from 10-month-old adult WT mice were stained with an antibody to CUX2. Immunosignals

(arrows) were observed widely in the brain including in neurons at the hippocampus and cerebral

cortex. A2-A5: magnified images outlined in A1. (**B**) CUX2 immunosignals were not observed

in *Cux2*(-/-) mouse. B2-B5: magnified images outlined in B1. (**C**) In 2-month-old wild-type

mouse, CUX2 (brown) is densely expressed in excitatory neurons at neocortical (II–IV) and

entorhinal cortex (II–III) upper layers, but in hippocampal observed in inhibitory but not

excitatory neurons. (**D**) In hippocampus at P15, CUX2 (brown) is expressed in interneurons but

not in excitatory neurons. Scalebars=500 µm (A1 and B1), 100 µm (C, D) or 20 µm (A2-A5 and

B2-B5). CTX; cerebral cortex, DG; dentate gyrus, so; stratum oriens, h; hilus.

Figure S4. CUX2 is expressed in hippocampal SST-positive, RLN-positive or PV-positive

inhibitory neurons. Tissue sections from P15 (**A-C**) or 2-month-old (**E-G**) WT mouse brains

were stained with antibodies to CUX2 (brown) and somatostatin (SST, blue) (A, E), reelin (RLN,

blue) (B, F) or parvalbumin (PV, blue) (C, G). CUX2 expression was observed in SST-positive,

PV-positive and RLN-positive (arrow) interneurons at both stages. Some of the intense CUX2-

positive cells were SST-negative, RLN-negative or PV-negative (white arrow head), and some of

the intense SST-positive or PV-positive cells were CUX2-negative (black arrow head). CUX2-

positive / PV-positive cell number increased at 2-months compared to P15. A2-A4, B2-B4, C2-

C4, E2-E4, F2-F4, G2-G4, A5 and B5: magnified images outlined in A1, B1, C1, E1, F1, G1, A4

and B4, respectively. (**D, H**) Tissue sections from at P15 (D) or 2-month-old (H) WT mouse

brain were stained with antibodies to SST (green) or RLN (magenta) and DAPI (cyan). The SST

expression was observed in RLN-positive interneurons (arrows) at the hilus of hippocampus.

SST/RLN-double positive cells were more frequent in P15 (D) than 2-month-old sections (H).

RLN signals became weaker at 2-months (H) compared to P15 (D). Some of intense SST-

positive cells were RLN-negative (arrow head), and some of intense RLN-positive cells were

SST-negative (asterisk). Scale bar=100 µm (A1, B1, C1, E1, F1, and G1) or 20 µm (A2-A4, B2-

B4, C2-C4, D, E2-E4, F2-F4, G2-G4 and H). h; hilus.

Figure S5. Unchanged cell numbers of hippocampal inhibitory neurons, mossy fibers and

cFos expression in *Cux2*-deficient mice. (**A**) Hippocampus was separated in 7 regions and

immunoreactive cells were counted. Cell densities were determined by average number of cells

in 4 sections / area. (**B-D**) There were no differences in the number of SST-positive (B), RLN-

positive (C) or PV-positive (D) neurons between genotypes. Statistical analyses were performed

using one-way ANOVA ($p > 0.05$) (n = WT: 6, *Cux2*(+/-): 6, *Cux2*(-/-): 6). so; stratum oriens,

sp; stratum pyramidale, sr; stratum radiatum, lm; lacunosum-moleculare, sm; stratum

moleculare, sg; stratum granulosum, h; hilus, w; whole hippocampus. (**E, F**) Timm staining (E)

and c-Fos immunohistochemistry (F) did not show differences in *Cux2*-KO mice. Scale bar =

100 µm (A and E).

Figure S6. IPSCs in PV-positive cells at dentate gyrus were increased, but no differences in

EPSCs in CA3 pyramidal cells of *Cux2*-deficient mice. (**A-D**) In PV-positive cells at dentate

gyrus, IPSCs were measured at 6~7-week-old. Amplitude histograms of sIPSC (A) and mIPSC

(C) from WT mice and *Cux2*(-/-) mice. Cumulative probability plots and average values (inset)

for sIPSC (B) and mIPSC (D) show significant differences in inter-event intervals of sIPSC and

mIPSC and amplitude of mIPSC populations derived from *Cux2*(-/-) and significantly higher

average of frequency of sIPSC, but unchanged averages of amplitude of sIPSC and both

frequency and amplitude of mIPSC in *Cux2*(-/-). (**E-H**) In pyramidal neurons at CA3 region,

EPSCs were measured at 6~7-week-old. Amplitude histograms of sEPSC (E) and mEPSC (G)

from WT, *Cux2*(+/-) and *Cux2*(-/-) mice. Cumulative probability plots and average values (inset)

for sEPSC (F) and mEPSC (H) in WT, *Cux2*(+/-) and *Cux2*(-/-) show no differences in EPSCs

among genotypes. (I) No significant difference on MF-evoked EPSCs in pyramidal neurons at

CA3 region was observed across genotypes at 6~7-week-old. Statistical analyses were performed

using one-way ANOVA followed by Tukey–Kramer Multiple Comparison Test (B, D, F, H, I) or

Kolmogorov-Smirnov (K-S) Test (B, D). * $P < 0.05$, *** $P < 0.001$.

Figure S7. Genome structures of human and mouse *CUX1* and *CASP* in WT and *Casp*-specific knock-out mice. Exons 1~14 are commonly found in human *CUX1* and *CASP* in human

(**A**) and in the corresponding genes in mice (**B**). In *Casp*-KO mice, a mutation

(c.1514^1515insTT, p.S506fs) was inserted in Casp-specific exon 17 (**C**). The diagrams were

reconstructed from that of UCSC Genome Browser. Vertical lines indicate exons.

Figure S8. CUX1 is expressed in hippocampal SST-positive, RLN-positive, or PV-positive interneurons. CUX1 (brown) is expressed in SST-positive (**A**), RLN-positive (**B**) and PV-positive (**C**) interneurons (arrows). Some of intense CUX1-positive cells were SST-negative,

RLN-negative or PV-negative (white arrowheads), and some of SST-positive, RLN-positive or

PV-positive cells were CUX1-negative (black arrowhead and not shown). Most of CUX1-positive cells were PV-negative (white arrowhead) in hilus (C). A2-A4, B2-B4, C2-C4:

magnified images outlined in A1, B1, C1, respectively. Scale bars = 100 μm (A1, B1 and C1) or

20 µm (A2-A4, B2-B4 and C2-C4). sp; stratum pyramidale, so; stratum oriens, sg; stratum

granulosum, h; hilus.

Figure S9. CASP, CUX1 and CUX2 expression and increased seizure susceptibility to kainate with a tendency of increase in excitatory cell number in entorhinal cortex of *Casp*-deficient mice. (**A**) CASP immunosignals were observed in both excitatory and inhibitory neurons in the hippocampus, though the signals in inhibitory neurons were especially intense such as those at hilus (A3) and stratum oriens of CA3 (A5), while signals in excitatory neurons are tiny such as those at stratum granulosum (A4) or pyramidal of CA3 (A6). CASP signals well disappeared in *Casp*(-/-) mice (A7, A8). Nuclei were stained with hematoxylin (blue). (**B, C**) Unaltered expression and subcellular localization of CUX1 and CUX2 in cerebral cortex (B) and hilus of hippocampus (C) of *Casp*-KO mice. (**D**) Body weights were similar among *Casp*-KO mice and WT littermates at 2-months. (**E**) qPCR experiments showed that mRNA expression levels of kainate receptor subunits were not altered in *Casp*-KO mice at 2-month-old. (**F**) In a Nissl staining, number of entorhinal cortex layer II-III neurons was comparable between WT and *Casp*(-/-) mice (2-month-old). (**G**) Seizure severity scores were significantly higher in *Casp*(-/-) male and combined gender mice. (**H**) Latencies to onset of generalized seizures (GS) were significantly decreased in male and combined gender of *Casp*(-/-) mice. (**I**) Latencies until death were also significantly decreased in combined gender of *Casp*(-/-) mice. Mice without GS or death within 3,600 sec were plotted at "no GS" or "no death", respectively. Circles represent individual mice. One-way ANOVA followed by Tukey–Kramer Multiple Comparison Test (D,

E, G-I), or one-way ANOVA test (F). mean (horizontal bars) ± s.e.m. (H, I). n=mouse number.

Scale bars= 50 μm (A1, A2, A7, A8, B and C) or 10 μm (A3-A6). * $P < 0.05$, ** $P < 0.01$.

Figure S10. CASP interacts with CUX2. mRFP-tagged CUX2 was co-immunoprecipitated

with FLAG-tagged CASP. Mutations did not affect the binding. Endophilin: negative control.

Original blots are presented in Supplementary Figure S11.

Figure S11. Full size western blot images. Original scanned western blot data that were used to

generate Figure S10. Dashed rectangles in the images indicate the location of the cropped

images. Images of blots with adequate length and membrane edges could not be provided

because the blots were cut prior to hybridisation with antibodies and scanned at inside of blots,

respectively.

Table 1: *CUX2*, *CUX1*, and *CASP* gene nonsynonymous variants in patients with epilepsy.

Patient ID	Gene	Nucleotide changes	Amino acid substitutions	SNP ID	Onset age (year)	Evaluation age (year)	Sex	Diagnosis	Variant allele count in		p-value	Variant allele count in					Mutation Taster	PolyPhen-2	PROVEAN	SIFT	M-CAP
									case	JP		J-HGVD	EVS	1kGP	ExAC	gnomAD					
SIZ-220		c.100C>T	p.R34W	rs199531850	10	27	M	**mTLE**	1 / 542	0 / 622	0.284	1 / 1,900	1 / 11,778	1 / 5,008	21 / 119,874	62 / 273,306	++	-	++	++	-
SIZ-016		c.1009G>A	p.D337N	rs201601231	16	24	M	**mTLE**	1 / 542	5 / 622	0.140	4 / 2,192	1 / 12,344	2 / 5,008	18 / 117,772	34 / 279,160	++	++	-	-	-
SIZ-296		c.1361C>T	p.P454L	rs768144991	2	15	M	Doose syndrome	1 / 542	0 / 622	0.284	5 / 2,184	NR	NR	1 / 24,448	8 / 167,304	++	+	++	-	+
SIZ-014	*CUX2*	c.2872T>C	p.W958R	NA	3	42	M	**mTLE**	1 / 542	0 / 622	0.284	NR	NR	NR	NR	NR	++	++	++	++	+
SIZ-004					11	37	F	**mTLE**													
SIZ-022					8	29	F	**mTLE**													
SIZ-073		c.3847G>A	p.E1283K	rs61745424	19	27	M	**mTLE**	5 / 542	2 / 622	0.185	9 / 2,126	227 / 12,676	143 / 5,008	3,893 / 120,384	8,188 / 280,456	-	-	-	-	NA
SIZ-079					13	27	F	**lTLE**													
SIZ-190					16	34	F	**mTLE**													
SIZ-784		c.3161C>T	p.S1054L	rs146486358	16	32	M	JME suspected	2 / 542	NT	NT	9 / 1,912	NR	13 / 5,008	101 / 120,970	231 / 282,774	++	++	-	-	NA
SIZ-891					0	8	M	GEFS													
SIZ-575	*CUX1*	c.3281C>T	p.A1094V	rs184337744	3	5	F	FLE	1 / 542	NT	NT	NR	1 / 13,006	1 / 5,008	38 / 120,936	38 / 250,814	-	+	-	-	+
SIZ-669		c.3815G>A	p.R1272Q	NA	0	30	F	SGE	1 / 542	NT	NT	2 / 2,152	NR	NR	NR	NR	++	++	-	++	+
SIZ-456		c.4172C>T	p.T1391I	NA	2	5	F	CAE	1 / 542	0 / 620	0.284	NR	NR	NR	NR	NR	-	-	-	-	+
		c.1390G>A	p.A464T	rs803064	*	*	*	*	247 / 542	NT	NT	991 / 2,210	NR	2,800 / 5,008	68,923 / 121,236	159,104 / 282,310	-	-	-	-	-
SIZ-127		c.1433C>T	p.A478V	NA	14	31	F	JME	1 / 542	0 / 622	0.284	NR	NR	NR	NR	NR	++	+	-	-	+
SIZ-063	*CASP*	c.1524delG	p.R509fs	rs782400087	6	9	F	CAE	1 / 542	0 / 622	0.284	NR	NR	NR	1 / 121,412	1 / 249,562	++	NA	NA	NA	NA
SIZ-068		c.1687G>A	p.G563S	rs187131238	12	28	F	JME	2 / 542	1 / 622	0.484	3 / 2,164	NR	2 / 5,008	4 / 120,438	7 / 247,922	-	-	-	-	-
SIZ-060					18	26	M	**mTLE**													
SIZ-638		c.1868_1870delTCT	p.F623del	NA	15	48	M	**lTLE**	1 / 542	0 / 622	0.284	NR	NR	NR	NR	NR	++	NA	++	NA	NA

mTLE: mesial temporal lobe epilepsy, lTLE: lateral temporal lobe epilepsy, JME: juvenile myoclonic epilepsy, CAE: childhood absence epilepsy, SGE: symptomatic generalized epilepsy, FLE: frontal lobe epilepsy, GEFS: generalized epilepsy with febrile seizure plus, M: Male, F: Female, JP: in-house Japanese control individuals, J-HGVD: Japanese Human Genetic Variation Database, EVS: Exome Variant Server NHLBI GO Exome Sequencing Project, 1kGP: The 1000 Genomes Project, ExAC: Exome Aggregation Consortium, gnomAD: Genome Aggregation Database. ++: Disease causing, Probably damaging, Deleterious, or Damaging, +: Possibly damaging, -: Polymorphism, Benign, Tolerated, or Neutral, NA: not available, NT: not tested, NR: not registered. The *CUX2* reference sequence (NM_015267) has an error at c.4414, and the correct nucleotide is C. *CUX2* nucleotide change c.4414G>C (p.V1472L) (rs6490073 in dbSNP, NCBI) was observed in all sequences in databases and in our subjects, suggesting that the *CUX2* reference sequence (NM_015267) has an error at this position (the correct nucleotide is c.4414C). * Informations are not described because of more than 100 individuals.

Figure 1

Figure 2

Figure 3