Individualised computational modelling of immune mediated disease onset, flare and clearance in psoriasis

Fedor Shmarov[1,2], Graham R. Smith[3], Sophie C. Weatherhead[4], Nick J. Reynolds[5,*], and Paolo Zuliani[1,*]

[1]School of Computing, Newcastle University, UK
[2]Department of Computer Science, University of Manchester, UK
[3]Bioinformatics Support Unit, Newcastle University, UK
[4]Department of Dermatology, Royal Victoria Infirmary, Newcastle Hospitals NHS Foundation Trust, UK
[5]Translational and Clinical Research Institute, Newcastle University, UK

[*]joint senior authors

Abstract

Despite increased understanding about psoriasis pathophysiology, currently there is a lack of predictive computational models. We developed a personalisable ordinary differential equations model of human epidermis that features two stable steady states: healthy skin and psoriasis. In line with experimental data, an immune stimulus initiated transition from healthy skin to psoriasis and apoptosis induced by UVB phototherapy returned the epidermis back to the healthy state. The flexibility of our model permitted the development of a patient-specific "UVB sensitivity" parameter that enabled accurate simulation of individual patients' clinical response trajectory. In a prospective clinical study of 94 patients, serial individual UVB doses and clinical response (Psoriasis Area Severity Index) values collected over the first three weeks of UVB therapy informed estimation of the "UVB sensitivity" parameter and the prediction of patient outcome at the end of phototherapy. Notably, our model was able to distinguish disease flares and offers the potential for clinical application in early assessment of response to UVB therapy outcome, and for individualised optimisation of phototherapy regimes to improve clinical outcome.

Author Summary

We present a new computer model for psoriasis, an immune-mediated disabling skin disease which presents with red, raised scaly plaques that can appear over the whole body. Psoriasis affects millions of people in the UK alone and causes significant impairment to quality of life, and currently has no cure. Only a few treatments (including UVB phototherapy) can induce temporary remission. Despite our increased understanding about psoriasis, treatments are still given on a 'trial and error' basis and there are no reliable computer models that can a) elucidate the mechanisms behind psoriasis onset or flare and b) predict a patient's response to a course of treatment (e.g. phototherapy) and the likelihood of inducing a period of remission. Our computer model addresses both these needs. First, it explicitly describes the interaction between the immune system and skin cells. Second, our model captures response to therapy at the individual patient level and enables personalised prediction of clinical outcomes. Notably, our model also supports prediction of amending individual UVB

phototherapy regimes based on the patient's initial response that include for example personalised delivery schedules (i.e. 3x weekly vs. 5x weekly phototherapy). Therefore, our work is a crucial step towards precision medicine for psoriasis treatment.

1 Introduction

Psoriasis vulgaris is a systemic immune-mediated inflammatory disease characterised by immune cell infiltration, keratinocyte hyperproliferation (up to an eight-fold increase in epidermal cell turnover) [4] and tortuosity of dermal blood vessels. Psoriasis is common (affecting 1-2% of "Western populations") and manifests itself as red scaly plaques distributed over the whole skin, causing significant disability and impairment to quality of life [6]. As well as being associated with inflammatory arthritis in up to 30% of the patients, psoriasis is increasingly linked to metabolic syndrome and cardiovascular disease [31].

The pathophysiology of psoriasis is complex, multifactorial and thought to be triggered through environmental genetic interactions. For example, in response to an environment stimulus (*e.g.,* injury or infection) myeloid dendritic cells and macrophages produce cytokines such as IL-12 and IL-23 that stimulate the activation and proliferation of T-cell subsets, which in turn produce IL-17A, IL-17F, IL-20, IL-22, TNF and IFN-γ [42]. Tc17/Th17 represent the dominant T cell subset in lesional psoriasis, with expanded Tc17/Th17 clonotypes localising to psoriatic epidermis [35]. These cytokines stimulate keratinocyte growth and production of chemokines and keratinocyte-based growth factors further amplifying the immune response and maintaining the hyperplastic and chronic inflammation within psoriasis plaques [16, 25, 26, 27, 30, 8, 1]. The importance of structural cells in regulating the immune response in health and disease is increasingly recognised [23, 21].

Psoriasis is a chronic persistent disease but may undergo periods of remission; currently though there is no cure. Narrow-band ultraviolet B (UVB) phototherapy is one of just a few treatments that can clear psoriasis plaques over 8-12 weeks of therapy and induce a period of remission [22]. As patients attend the hospital department 3 times weekly, this facilitates the regular recording of disease activity (Psoriasis Area and Severity Index [PASI] - a visual examination of psoriasis extent), clinical response (including disease exacerbations - flares) and any adverse events. Nevertheless, relapse is common (occurring in 70% of patients within 6 months from the end of the therapy); patients may then be considered for further UVB or switched to systemic therapy. At present, accurate prediction of individual patient outcome to UVB or systemic therapy is not feasible, although some progress has been achieved [46]. For example, lower rates of psoriasis clearance in the early stages of the UVB therapy were negatively associated with the final clearance outcome [42].

Previous agent-based models [42] and ordinary differential equation (ODE)-based models [48] have simulated psoriasis onset and psoriasis clearance by induction of keratinocyte apoptosis through UVB phototherapy. However, despite the crucial role of acquired and innate immunity in initiating and maintaining psoriasis, these models provide only an implicit representation of the immune system. In this paper, we present a new ordinary differential equation model of epidermis that: 1) explicitly describes the complex interplay between immune cells and keratinocytes; 2) maintains two stable steady states: a *healthy* one and a *psoriatic* one and 3) enables switching between the two steady states through crossing of an unstable "transition" state. This can occur either via the introduction of a sufficiently strong immune stimulus ("healthy" to "psoriatic" transition) or by inducing an appropriate amount of apoptosis of proliferating keratinocytes and immune cells by UVB therapy ("psoriatic" to "healthy" transition) through UVB phototherapy.

Our ODE-based model delineates complex interactions between immune stimuli, keratinocyte growth, differentiation and apoptosis in the onset of psoriasis lesions and resolution during UVB phototherapy. We demonstrate that individual patients' PASI trajectories, recorded in response to UVB therapy, can be well approximated by a designated model species and by estimating a single individualised model parameter (called "UVB sensitivity") that is proportional to the rate of UVB-induced apoptosis. Notably, we show

that the value of the UVB sensitivity parameter can be accurately estimated by the end of the third week of a patient's UVB therapy (given the UVB doses administered and the PASI measurements over that period). We further demonstrate the applicability of our model in defining psoriasis flares. Together these results support the prediction of longer term clinical outcomes based on trajectory of response and the development of individualised modifications to the delivery of phototherapy that can be tested in the clinic.

2 Model and Methods

2.1 Epidermis model

Our model focuses on human epidermis (see Figure 1a) and incorporates the following cell species:

- proliferating keratinocytes: stem cells (SC) and transit-amplifying cells (TA);

- differentiating keratinocytes (D).

The model also incorporates immune cells (T cells) and dendritic cells (DC), which may be located in either the epidermis or dermis but the location is not relevant to our model.

Proliferation and activation of SC and TA cells is mediated by IL-22, IL-17, type I interferons and TNFα cytokines produced by the activated T cells and/or dentritic cells [1]. In response to the increase in proliferation rate, keratinocytes increase their production of growth factors (GF) and chemokines that attract and activate dendritic cells producing more IL-23, which in turn activates more T cells, resulting in a positive feedback and self-sustaining loop.

Our model is defined by the system of ordinary differential equations (ODEs) displayed in Figure 1c, and is designed as a bi-modal switch consisting of three steady states: two are stable (healthy and psoriatic states), and one is unstable (transition state). (The model features 32 steady states overall, but only the three states mentioned above lay in the positive real space: see Section SM.1.) The model design process considered the following main properties: epidermis composition, speed of psoriasis onset, epidermal

turnover time, keratinocytes, apoptosis and desquamation rates. These were built upon measured parameters described in the literature which are detailed below.

Epidermis composition Figure 1d shows the epidermis composition in the three steady states of our model (the precise values are reported in Table S2). The total number of cells per unit area of human epidermis is different from person to person, and varies across the human body. Ref. [5] reports a total cell density of $73,952 \pm 19,426$ $cells/mm^2$ (mean and standard deviation) and $1,394 \pm 321$ $cells/mm^2$ for Langerhans cells (*i.e.,* dendritic cells in epidermis). In line with these data, the total cell and Langerhans cells densities in our model are $79,828$ $cells/mm^2$ and $1,563$ $cells/mm^2$, respectively.

The proportions of cell species in the healthy steady state are consistent with the values found in the literature. Our model features 63.2% of differentiating cells (40-66% is reported by [42]), 26.4% of transit-amplifying cells, 4.7% of stem cells, 1.8% dendritic cells and 1.8% of T cells. Hence, our model suggests that up to 96% of all cells in the epidermis are keratinocytes [42, 37].

Psoriasis onset We model the onset of psoriasis by increasing the activation rate (dc_{act}) of the dendritic cells by a constant amount dc_{stim} for τ_{stim} days. This produces psoriasis onset at different rates: from fast (as quick as 10 days) to slow (as long as 14 weeks). That is consistent with the known time lags between the onset of an immunological stimulus (*e.g.,* streptococcal sore throat) and the first appearance of psoriatic lesions [42, 38, 17].

In the psoriatic state the growth rate of the proliferating keratinocytes (SC and TA cells) and their differentiation rate increase with respect to the healthy state by 9.74 and 9.73 times, respectively (see Section SM.2 for more details). Compared to normal skin, the increased number of proliferating keratinocytes *in vivo* ranges in psoriasis from 2-3 times [7, 35] to 6-8 times [4, 11]. As a result, the cell density in the psoriatic state of our model is $266,012$ $cells/mm^2$, *i.e.,* 3.33 times the density in the healthy state, which is

Figure 1: **Our ODE model of epidermis describes an explicit interaction between the main types of keratinocytes and immune cells mediating psoriasis.** Panels: (a) - mechanism of interaction between the main cell types and cytokines in epidermis; (b) - interactions between the species of our model, where SC - stem cells, TA - transit-amplifying cells, D - differentiating cells, DC - dendritic cells, T - T cells, A - apoptotic cells, $IL_{17/22}$ - interleukin-17/22, IL_{23} - interleukin-23, $TNF\alpha$ - tumour necrosis factor alpha, GF - keratinocyte-derived growth factors, and $\varnothing$ - degradation species; (c) - differential equations describing the interactions between the model species (the parameter values are reported in Table S1); (d) - cell density (cells/mm^2) for every cell type (excluding apoptotic cells) in the healthy, psoriatic and transition steady states.

within the 2-5 times range reported in clinical studies [39, 44].

Epidermal turnover time In psoriasis, despite an increase in epidermis thickness, the epidermal turnover time drops by up to 4-7 times [45, 19]. This implies that the cell growth rate cannot linearly depend on the cell density (see Section SM.3). Thus, in each ODE, we introduced the non-linear term $a \cdot X^p$, where a is a constant, X is the corresponding keratinocyte species, and p is a parameter modelling the nonlinear dependency between the keratinocyte growth rate and the cell density. For simplicity, we assumed $p = 2$, but any $p > 1$ can be used for personalising the model.

For setting the parameter values related to the growth rate, we applied epidermal turnover time values reported in the literature. Turnover times for healthy epidermis vary from 39 days [43], to 47-48 days [20] to 40-56 days [19], while a previously published model [48] predicts 52.5 days.

Similarly to [43, 48], the epidermal turnover time in our model can be calculated as the sum of the turnover time in the proliferating and the differentiating compartments using the formula:

$$\tau = \frac{SC + TA}{(TNF\alpha + IL_{17/22})(sc_2 SC + sc_{2ta} SC + ta_2 TA)} + \frac{D}{ta_{2d}(TNF\alpha + IL_{17/22})TA}$$

where $TNF\alpha$ is the tumour necrosis factor α, and the cell species densities are taken from either the psoriatic state or the healthy state (see Table S2). In our model, the epidermal turnover time in psoriatic epidermis drops almost three-fold in comparison to healthy epidermis, from 41.31 days to 14.25 days, consistent with *in vivo* studies [45, 19].

Apoptosis and desquamation rates Cell apoptosis and desquamation (terminal differentiation) are the two main mechanisms in our model by which keratinocytes can leave the proliferating and the differentiating compartments. It is technically challenging to measure rates and duration of apoptosis in human tissue, and so the parameters used in the model are based on data from mouse skin and *in vitro* studies of human keratinocytes [41, 18, 28, 42, 29]. In our model we assume apoptosis is distributed between 0 and 24 hours and found little difference if the duration was extended to 48 hours. In addition, it is assumed that in healthy epidermis all cells undergo apoptosis at the same rate (relative to their cell mass). In Section SM.4 we provide more details.

2.2 Modelling UVB phototherapy

In our model, psoriasis clearance with UVB is implemented by increasing the rate of apoptosis of proliferating keratinocytes (SC and TA) and immune cells (DC and T) for t_{uvb} hours by $uvb_{dose} \cdot uvb_s \cdot (X - X_H)$, where uvb_{dose} is the administered UVB dose (in J/cm^2), X is the target species cell density, X_H is the target species cell density in the healthy steady state, and uvb_s is the "UVB sensitivity" parameter in ($J^{-1}d^{-1}$). Note that differentiated keratinocytes (D) may undergo apoptosis when UVB irradiated, but at a negligible rate [41, Section 5.3.4]. Hence, our model does not increase the apoptotic rate of differentiated cells during UVB irradiation.

As mentioned earlier, we assume that $t_{uvb} = 24$ hours, as the peak of cell apoptosis is reported to be between 18 and 24 hours following UVB irradiation [42, Figure S2]. Thus, the rate of UVB induced apoptosis is proportional to the applied dose (defined in terms of energy per unit area), current cell density (*i.e.,* the higher the cell density the higher the number of apoptotic cells) and the patient-specific parameter UVB sensitivity (larger sensitivity values result into higher apoptosis rates). The term $(X - X_H)$ is introduced to account for the fact that the thickness of healthy epidermis does not reduce in response to UV [34, 33]. We also restrict our model to operate only between the healthy and psoriatic states thus, the term $(X - X_H)$ is always kept positive.

Standard clinical protocols administer narrow-band UVB three times a week (*e.g.,* Monday, Wednesday and Friday) for up to 12 weeks, with increasing graduated dose increments (on alternate doses) over the course of the treatment. An example of UVB dosimetry is shown in Figure 2c. The changes in cell densities and cytokine concentrations follow-

ing the indicated UVB regime are shown in Figure 2a and Figure 2b; the rate of cell apoptosis is shown in Figure 2d. The rate of cell apoptosis in the model falls back to the basal value very shortly after the end of the 24-hour period. This is because apoptotic cells are removed from the epidermis fairly quickly in our model (mean lifetime of 40 minutes – see Section SM.4). The obtained apoptosis values are consistent with those reported *in vivo* [42, Figure 1e].

Our model predicts complete clearance and eventual remission when the model dynamics drops below the "transition" steady state (*i.e.,* at approximately 90% clearance). These data are consistent with recent findings in a prospective study of 100 patients in which achievement of PASI90 (*i.e.,* at least 90% improvement over their initial PASI) pointed to longer remission [40]. However, not all patients achieving PASI90 progress to complete clearance and/or remission.

Apoptosis *vs.* Growth Arrest An important question is whether growth arrest contributes to clearance for a given rate of apoptosis. There is evidence that UVB induces both in normal skin [3, 32, 24]. In psoriatic skin, UVB induces apoptosis but no significant difference in the number of proliferating cells in untreated *vs.* UVB-treated lesional skin (although genes associated with growth arrest are upregulated in UVB treated skin) [42]. We thus considered whether in our model growth arrest could account for clearance during UVB therapy. We found that growth arrest alone (no apoptosis) could induce clearance with a delay of several weeks after the end of the therapy (see Figure S1 – similar results are reported by [42]). However, in clinical practice psoriasis does not improve after UVB treatment completion, and therefore we do not further consider growth arrest in this paper.

2.3 Modelling PASI

The Psoriasis Area and Severity Index (PASI) is used for assessing the severity of the ongoing disease [47, 2, 36]. It ranges from 0 to 72, and it is calculated as a weighted sum of sub-scores corresponding to four body regions:

$$\text{PASI} = 0.1 \cdot C_h + 0.2 \cdot C_{ul} + 0.3 \cdot C_t + 0.4 \cdot C_{ll},$$

where C_h, C_{ul}, C_t and C_{ll} are the sub-score values for head, upper limbs, trunk and lower limbs, respectively. Each sub-score is obtained as

$$C = (S_{ind} + S_{ery} + S_{desq}) \cdot S_{area},$$

where S_{ind}, S_{ery} and S_{desq} are values between 0 and 4 representing the severity of induration (thickness), erythema (redness) and desquamation (scaling), respectively, for the four body regions; S_{area} is a value between 0 and 6 representing the extent of the affected area.

We mimic patients' PASI by using the species of our ODE model as follows: we use the total cell density $\Sigma = (SC + TA + D + T + DC + A)$ as a proxy for epidermal thickness, T cell species to estimate erythema and non-proliferating cells D to assess scaling.

We use only T cells to represent erythema (inflammation) for simplicity, as all the species modelling the PASI components are scaled between 0 and 1 (as shown below), and the scaled model dynamics for the cytokine species (*i.e.,* IL-17, IL-23 and TNFα) are almost identical to the scaled dynamics of T cells.

In our model we do not take the surface area into account, and we only consider severity of PASI components over a unit area (mm^2). Thus, our model assumes that the affected area stays constant throughout the therapy, while all the other components change (*i.e.,* the plaques "fade away" rather than "shrink" in size).

If we rescale the cell species concentration to the $[0, 1]$ interval (where 0 and 1 represent healthy and psoriatic steady states, respectively), we can model the relative change in each of the psoriasis symptoms S_{ind}, S_{ery} and S_{desq} over a unit area of skin. After weighting the rescaled species we calculate the severity of the symptoms over a unit area for each body region (*i.e.,* an estimate for $S_{ind} + S_{ery} + S_{desq}$) as

$$C^* = S_{ind} \cdot \Sigma_{[0,1]} + S_{ery} \cdot T_{[0,1]} + S_{desq} \cdot D_{[0,1]},$$

where $\Sigma_{[0,1]}$, $T_{[0,1]}$ and $D_{[0,1]}$ are the cell species rescaled to [0,1]. Thus, obtaining C^* for each body

Figure 2: **ODE psoriasis model demonstrates that UVB-induced apoptosis leads to psoriasis clearance.** Simulation of the model response to a 10-week UVB phototherapy. Panels: (a) - cell densities, where SC - stem cells, TA - transit-amplifying cells, D - differentiating cells, DC - dendritic cells, T - T cells, and $totC = SC + TA + D + DC + T$ - total number of cells, (b) - cytokines concentrations, where $IL17/22$ - interleukin-17/22, $IL23$ - interleukin-23, $TNF\alpha$ - tumour necrosis factor alpha, and GF - keratinocyte-derived growth factors, (c) - corresponding UVB irradiation regime, (d) - number of apoptotic cells per 1,000 cells.

region, and assuming that the amount of energy delivered by UVB per unit of area is the same across the entire surface of the body, the resulting PASI value can be modelled by

$$\Psi = 0.1 \cdot C_h^* \cdot S_{h,area} + 0.2 \cdot C_{ul}^* \cdot S_{ul,area} +$$
$$0.3 \cdot C_t^* \cdot S_{t,area} + 0.4 \cdot C_{ll}^* \cdot S_{ll,area} .$$

However, PASI subscores are seldom recorded in clinical practice – only the cumulative score is likely to be available. In order to mitigate this issue, we consider the average behaviour of the components over the entire body instead of modelling each body region separately. As a result, we introduce a new species to simulate a patient's PASI when the PASI subscores are not available:

$$\Psi^* = C_{[0,1]}^* \cdot PASI_0, \qquad (1)$$

where $C_{[0,1]}^* = \frac{C^*}{S_{ind}+S_{ery}+S_{desq}}$ is the value of C^* rescaled to $[0,1]$, and $PASI_0$ is the baseline PASI.

In the data we used in this work the PASI subscores are not available, and we thus used Ψ^* instead of Ψ and we assumed $S_{ind} = S_{ery} = S_{desq} = 1$ in all our experiments.

2.4 Model personalisation

Patient data Our clinical data are derived from a prospective cohort of 94 patients receiving narrow-band UVB therapy for psoriasis, recruited at the Royal Victoria Infirmary, Newcastle upon Tyne (UK). The dataset is described fully in [40] but in brief it includes serial weekly patients' PASI measurements (median 7; range 4-11), corresponding serial UVB doses (median 24; range 4-11), delivered 3 times a week together with data on time to relapse and PASI at relapse, collected for up to 18 months. Out of 94 patients, 6 subjects did not relapse within the 18-month monitoring period, and 26 individuals

were lost to follow-up.

Parameter fitting   We estimate the UVB sensitivity parameter uvb_s to fit the model PASI simulations to the patients' PASI data. The optimal value of uvb_s for every patient is found by minimising the function

$$\Sigma_{i=1}^{t}(PASI_i - \Psi^*(7 \cdot i))^2 \qquad (2)$$

where t is the number of points (weeks) in the patient's PASI trajectory (the model time units are days while patients data are recorded weekly, hence the use of $7 \cdot i$ in Ψ^*). The minimisation of function (2) with respect to uvb_s is performed via an exhaustive search over the $[0, 1]$ interval, starting at 0 with a step of 0.01. For every value of uvb_s we simulate our ODE model (see Figure 1c) with Ψ^* defined by (1) and $\Psi^*(0) = PASI_0$ and all other model species set at the psoriatic steady state. This method also allows minimising the value of uvb_s: from all the parameter values producing the lowest value for the objective function (2) we choose the lowest value for uvb_s. This prevents the model from being ultra responsive to treatments via unrealistically large uvb_s values. Equivalently, we prevent the model from reaching psoriasis clearance with a small number of (possibly unsafe) strong UVB irradiations.

Flares   During longitudinal follow up [15, 10] and during the course of the therapy some patients may experience spontaneous disease exacerbations or flares (*i.e.,* worsening of the symptoms and signs due to undefined environmental/immunological stimuli). We simulate patients' flares by introducing and fitting parameters $dc_{stim,i}$ for $i = 1, \ldots, t$ (where t is the number of PASI values in the patient's trajectory) for every patient. Each such parameter models an immune stimulus of constant strength lasting 7 days (since in our dataset we have at most one PASI reading per week). The parameters $dc_{stim,i}$ are fitted sequentially beginning with $i = 1$, and every parameter is searched incrementally in the interval $[0, 6{,}000]$ starting at 0 with a step size of 60 (*i.e.,* 100 increments). The current parameter value is increased until the objective function value (2) starts rising. This

parameter value is set in the model, and the fitting of the next parameter $dc_{stim,i+1}$ is performed.

Statistical methods   The goodness of fit of our model is assessed by calculating the distribution of the difference $(PASI_i - \Psi^*(7 \cdot i))$ between the patients' PASI trajectories and the model PASI simulation for all but the baseline PASI values. (Our dataset contains 754 PASI values distributed between 0 and 24.1, with mean 3.41, median 2.3 and IQR [1, 4.5]). We obtain the distribution of the PASI estimation error by uniformly sampling $(PASI_i - \Psi_i^*)$ for each patient. We repeat this process 1,000 times and compute mean, median and standard deviation of the resulting cumulative distribution, which are then compared with PASI assessment errors made by formally trained physicians [14].

3   Results

We developed an ODE model of normal and psoriasis skin that would enable direct comparison to patient specific disease trajectories and prediction of outcomes to therapy at an individual patient level. As outlined in Section 2.1, our model's behaviour (including key indicators such as proliferation rates and epidermal turnover time), is consistent with data from clinical studies. In the sections reported below, we systematically studied the dynamic behaviour of our model for predicting the speed of psoriasis onset and the impact of varying UVB therapy regimes and UVB sensitivity to outcome. Finally, we explored the capabilities of our model for personalisation of disease trajectories and for stratification of patients undergoing disease flares.

3.1   ODE model suggests speed of psoriasis onset depends on strength and duration of immune stimulus

As described in Section 2.1, the onset of psoriasis is initiated by an immune stimulus that increases the activation rate of the dendritic cells. Our analysis indicates that the speed of psoriasis onset depends on both the strength and the duration of this immune

stimulus (Figure 3). In addition, Figure 3 demonstrates the model dynamics for two exemplary scenarios simulating slow (14 weeks) and fast (10 days) psoriasis onset. Within the ranges explored for stimulus strength and duration, 14 weeks is the longest delay and 10 days the shortest delay to a *full* psoriasis plaque achievable by our model – clinical observations report psoriasis onset no sooner than 4 days after an immune stimulus [17].

3.2 ODE model simulates individual clinical outcomes and personalised amendments of phototherapy to induce psoriasis clearance and remission

Our model indicates that a minimum number of UVB episodes and appropriate irradiation frequency are necessary for clearing psoriasis and inducing remission, depending on the patient-specific UVB sensitivity parameter uvb_s (*i.e.*, patient-specific UVB sensitivity to phototherapy), and actual UVB doses that will be administered.

Following a given UVB irradiation regime, our model can simulate different outcomes in which the chances of psoriasis clearance increase for higher values of uvb_s. For example, two models with different UVB sensitivity values (modelling two patients) receiving equal amounts of UVB might not reach the same outcome. The heatmaps presented in Figure 4 depict the model outcome as a function of the number of UVB doses (of a given therapy regime) and the UVB sensitivity parameter uvb_s. Figure 4 shows that changing therapy from 3 times to 5 times a week can, overall, increase the likelihood and speed of psoriasis clearance, consistent with results from a (small) clinical trial [9]. However, as in [9], the actual improvement is modest and might not be clinically justifiable because of inconvenience or risk of side effects (*e.g.*, erythema or "sun burn").

If within Figure 4, we consider a patient characterised by a "low" $uvb_s = 0.05$, we can then compare the model simulations of a therapeutic phototherapy course consisting of 30 doses but delivered 3 times *vs.* 5 times a week and how this affects relapse. The

3 times a week simulation results in relapse of psoriasis within a few months, whereas the 5 times a week therapy regime induces a longer duration of remission, thereby showing that some patients might potentially benefit from phototherapy delivery adjustments.

Finally, we note that the uvb_s parameter is associated with the rate of keratinocyte and lymphocyte apoptosis induced by UVB in the model, and it could be potentially inferred from corresponding biomarkers collected before the start of the treatment (*e.g.*, number of apoptotic cells measured from biopsies 24-48 hours after localised delivery of phototherapy). Taken together with the above results, these data provide evidence that our model can simulate personalised response to UVB therapy, individual dosimetry and UVB administration regimes in clinical practice.

3.3 Representation of UVB response by the UVB sensitivity parameter enables closely fitting PASI trajectories

Using the patients' UVB doses and baseline PASI, we fitted the uvb_s parameter in our model to closely reproduce the patients' PASI trajectories. The resulting models are called uvb_s-*personalised*. For example, Figures 5a and 5b show the PASI simulation computed by the uvb_s-personalised models of two different patients of our cohort.

We fitted the uvb_s parameter for all our 94 patients. All of the derived uvb_s values were distributed between 0 and 1, median 0.24 and IQR range [0.16, 0.34]. The distribution (n=754) of the difference between the model simulated PASI Ψ^* (see Eq.(1)) and the patient's actual PASI data is shown in Figure 5c. The resultant model simulations provided a close fit to the patients' data with a mean PASI difference of 0.27 PASI units (recall that PASI ranges between 0 and 72).

Given a patient with a total of t weekly PASI readings, we calculated (for $i = 1, \ldots, t$) the *absolute* (*i.e.*, $|PASI_i - \Psi^*(7 \cdot i)| = AD_i$, see Figure 5d) and the *relative*[1] ($\frac{AD_i}{PASI_i}$, see Figure 5e) PASI differences and

---

[1]Here we only used the values whose corresponding PASIs

Figure 3: **The speed of psoriasis onset in ODE model depends on the strength and the duration of the immune stimulus.** The heat map white area denotes combinations of immune stimulus strength and duration that do not lead to psoriasis; SC - stem cells, TA - transit-amplifying cells, D - differentiating cells, DC - dendritic cells, T - T cells, $totC = SC + TA + D + DC + T$ - total number of cells, $IL17/22$ - interleukin-17/22, $IL23$ - interleukin-23, $TNF\alpha$ - tumour necrosis factor alpha, and GF - keratinocyte-derived growth factors. In this figure psoriasis onset is defined when $totC = totC_H + 0.9(totC_P - totC_H) \approx 247{,}376$ $cells/mm^2$ (*i.e.,* 90% of the difference between the total cell density in the psoriatic and the healthy steady states – see Table S2). This is due to the relatively slow convergence of the model to the psoriatic steady state.

Figure 4: **Our ODE model predicts that the total number of UVB doses, irradiation frequency and the patient's UVB sensitivity (modelled by the uvb_s parameter) are the key factors in designing a personalised therapy for achieving psoriasis clearance and remission.** The green areas contain the values for which the model predicts remission following the therapy. Two examples of model simulation present dynamics of keratinocytes and immune cells for the same UVB sensitivity ($uvb_s = 0.05$) after the same number of UVB doses ($n = 30$) over two regimes with different irradiation frequencies, where SC - stem cells, TA - transit-amplifying cells, D - differentiating cells, DC - dendritic cells, T - T cells, and $totC = SC + TA + D + DC + T$ - total number of cells.

compared them to the PASI assessment discrepancies achieved by trained clinicians. For example, Ref. [14] reports on three formally trained physicians who performed 720 image-based PASI assessments of 120 patients with psoriasis (every physician assessed PASI twice: at week 0 and 4 weeks later). The mean and standard deviation (see Table 1) of the absolute and relative PASI differences of our models were very similar to the errors made by formally trained physicians assessing patients' PASI.

Together, these results suggest that given the patient's baseline PASI, their UVB doses administered during the therapy and the UVB sensitivity parameter our model can simulate PASI outcomes that are indistinguishable from PASI assessments made by trained professionals. We note that the UVB sensitivity parameter could be estimated at baseline (*e.g.,* by measuring the rate of apoptosis in skin biopsies) or early during the therapy, as we explain in Section 3.4. Therefore, our model can be used to predict individual patient response to phototherapy and enables design of personalised UVB regimes to improve outcomes, initially *in silico*, but ultimately in a clinical trial.

3.4 The UVB sensitivity parameter can be reliably estimated at the third week of treatment

We found no statistically significant associations between the derived uvb_s values and the available patients' clinical variables collected at baseline (*i.e.,* BMI, age, sex, smoking status, alcohol consumption, skin type, baseline MED, age of psoriasis onset, previous phototherapies). Therefore, we tested whether the uvb_s parameter could be predicted by using only a portion of a patient's PASI trajectory. We discovered that the earliest reasonable prediction ($R^2 = 0.895$ and adjusted $R^2 = 0.894$; root mean square error = 0.069, Figure 5f) was made by fitting uvb_s with the data available at the end of week 3 of the therapy, *i.e.,* four PASI measurements and nine UVB doses. These data show that we can make a reasonable estimation of a patient's UVB sensitivity parameter value by the end of the third week of the therapy which can

then be used to predict subsequent response to UVB phototherapy. This discovery opens the path to personalisation of therapy.

3.5 The UVB sensitivity parameter and immune stimuli enable stratification of flaring patients

Flares are spontaneous worsening of a patient's psoriasis symptoms and signs that can happen both on and off therapy. For example, Figure 5b illustrates an unexpected and sustained increase in a subject's PASI (outside of observer error range [14]) despite ongoing UVB phototherapy. We hypothesised that our uvb_s-personalised models could be used to distinguish between psoriasis flares and PASI assessment discrepancies. We remark that in case uvb_s is not available from clinical biomarkers, one could assume a baseline value or estimate a value after three weeks of therapy as previously discussed.

For each patient with t weekly PASI measurements we looked at their PASI errors, computed as

$$\Delta_i = PASI_i - \Psi^*(7 \cdot i) \qquad i = 1, \ldots, t,$$

where Ψ^* is the uvb_s-personalised model PASI simulation (see Section 2.3), and the relative PASI errors with respect to the model simulation, calculated as

$$\delta_i = \frac{PASI_i - \Psi^*(7 \cdot i)}{\Psi^*(7 \cdot i)} \qquad i = 1, \ldots, t,$$

because unlike the patients' PASI, the value of $\Psi^*(7 \cdot i)$ is always strictly positive.

Utilising the DBSCAN clustering algorithm [12], we clustered the calculated PASI differences of all patients (n=754 PASI measurements) into two groups: PASI assessment errors and potential flares. We asked DBSCAN to identify two groups (see Figure 6a) within the data: the main cluster (black triangles) and the outliers (red crosses). By considering all positive relative error values (*i.e.,* the δ's) as flares (n=98) we fitted a *threshold curve*

$$\delta = e^{-5 \cdot \Delta} \qquad (3)$$

that separated the outliers (potential flares) from the main cluster (potential PASI assessment errors).

Figure 5: **PASI measurements and UVB doses over the first three weeks of the therapy are sufficient to predict the UVB sensitivity uvb_s parameter, which allows high-accuracy model personalisation.** Panels: (a,b) - results of parameter fitting for two different patients; (c) - distribution (n = 754) of the difference between the model PASI simulation Ψ^* and the patients' actual PASI data; (d) - distribution (n = 754) of the absolute value of the difference between the model simulation and patients' PASI data; (e) - distribution (n = 738) of relative PASI difference calculated as a ratio between the absolute PASI difference and the corresponding PASI value (excluding those values for which corresponding PASI = 0; n = 16), (f) - results of fitting uvb_s after 3 weeks with respect to fitting uvb_s at the end of the therapy.

Table 1: **Simulation of PASI outcomes by our ODE model that are comparable to PASI assessments made by formally trained physicians and nurses.** The mean assessed PASI value was 8.8 and the range was [0.7, 34.8] (the corresponding figures in our dataset are 4.01 and [0, 31.8]), and the best mean absolute PASI difference was (reported for observers 1 and 2) 1.968 with standard deviation of 2.480, while the best mean absolute intra-observer PASI difference (the difference between the PASI assessments for observer 2) was 1.984 with a standard deviation of 2.175. NA – value is not available.

Source	Absolute PASI difference (mean±sd)	Relative PASI difference (mean±sd)
Inter-rater difference [14]	1.968 ± 2.480	24%± NA
Intra-rater difference [14]	1.984 ± 2.175	29.2%±NA
Our model difference	0.842 ± 0.838	$35.4\% \pm 30.1\%$

(The 98 potential flares are distributed over 47 patients out of 94.)

We note that the threshold curve was identified using a large number of PASI measurements in which the sequential nature of the measurements was not retained. This would enable using our uvb_s-personalised models for 'online' detection of flares in the clinic as follows: every time a PASI measurement is obtained one computes both the simple (Δ) error and the relative (δ) error and then decides whether the pair sits above (flare) or below (measurement error) the threshold curve. Hence, as soon as a flare is detected the patient's phototherapy regime can be modified (through increasing the frequency of irradiations and/or adding more UVB doses) to achieve better clearance.

3.6 Flares during the therapy are associated with shorter remission

Our uvb_s-personalised models provide evidence that detecting flares is extremely important not only for improving overall model fit, but also for predicting patients' remission period.

To improve model fit, we applied an immune stimulus (iteratively increasing its strength starting from 0 while the sum of the squared errors (2) decreases) for 7 days for every week which is identified as a flare by the threshold curve (3). The resulting model is called *flare-enabled*. Figure 6b shows an example of a "flare" that was predicted on the basis of a slower than expected psoriasis clearance that probably could not be detected visually. Our model implements a weak but persistent immunological stimulus but this result could also possibly be explained by increasing adaptation and resistance to UVB, which we do not currently model. In contrast, Figure 6c demonstrates a substantial flare in response to larger immunological stimulus.

Next, we studied patients' remission. We dropped therefore patients who were lost to follow-up (n=26) and those who did not relapse within the 18 month period (n=6; our model correctly predicts remission for all these patients). Then we divided the remaining patients (n=62) into two groups based on their simulated post-therapy PASI values: those whose uvb_s-personalised, flare-enabled model simulations predicted relapse were assigned to a *Simulated Relapse* group (n=13), and the rest were placed into a *Simulated Remission* group (n=49). When compared to the actual, observed behaviour (see Figure 6d), the patients in the *Simulated Relapse* group demonstrated a shorter remission period (median value of 3 months *vs.* 5 months, Kruskal-Wallis p-value=0.035) compared to the *Simulated Remission* group. (*Simulated Relapse* group: mean = 4.23, standard deviation = 4.17, range = [1, 16] and IQR = [1.75, 6]; *Simulated Remission* group: mean = 6.45, standard deviation = 4.7, range = [1, 18] and IQR = [3.75, 8.25].)

We conclude that detecting flares is important so that therapy amendments can be applied not only to achieve better clearance at the end of photother-

apy but also with the aim of extending the remission period.

4   Discussion

There is increasing interest in the application of systems biology modelling in medicine and immune-mediated inflammatory disorders (IMIDs) [13]. Focusing on psoriasis as an archetypal IMID, we have developed an ODE model of normal and psoriatic epidermis that is stable in the two distinct steady states and with specific perturbations transitions between the two. Importantly, quantitative read-outs from the model are entirely consistent with data derived from clinical and *in vivo* studies. Although lacking spatial information, our model enabled a) detailed and iterative *in silico* studies of specific parameters (such as the initiating immune stimulus) and led to specific predictions that can be subsequently tested in the clinical setting and b) testing of specific hypotheses including the relative contribution of immune cell and keratinocyte apoptosis to the resolution of psoriasis. A notable advance is the flexibility of the model to be personalised at the individual subject level. During model development the running of simulations allowed the parameters and behavior of the model to be adjusted and compared to clinical outcome data. Comparisons of the model outputs to patient level data provided insight into the potential mechanisms underlying disease flares that occur even during therapy. Application of the model in the clinical setting would facilitate sampling of tissues and blood for biomarker development.

Currently psoriasis treatments are given on a 'trial and error' basis. Our ODE model could be used as a step towards personalised medicine. Identifying poorer responders early in the therapy (by estimating their UVB sensitivity at baseline or after 3 weeks of therapy) means that these patients could be switched to receive more intensive phototherapy treatment (*e.g.,* 5x weekly) or switched to systemic therapy at an earlier stage, saving the patient and hospital valuable resources. Indeed, our model predicts that altering the therapeutic regime to 5x weekly can in some cases improve clinical outcome. A clinical study is being initiated at our centre to investigate whether intensifying phototherapy treatment to 5x weekly, in patients showing a poor initial response, results in improved outcomes.

By incorporating a bimodal switch within the system of differential equations, the model integrated interactions between dentritic cells, T cells and cytokines mediated via specific cytokines and growth factors, in transitioning between states. During this study, we analysed the characteristics of initiating immune stimulus and the consequence of this on the time lag before psoriasis develops and time to reach steady state. We showed that apoptosis rates consistent with those measured in psoriasis during simulated UVB phototherapy (with clinically-relevant dose escalation) [42] were sufficient to induce resolution of the plaque. Furthermore, in line with previous findings [42], our model predicts that growth arrest alone cannot produce the clearance timeline observed in clinical practice.

By applying subject-level, carefully collected, serial clinical measurements (obtained during regular attendance for phototherapy) to our model, we were able to closely simulate the clinical course of individual patients (Figure 5). In certain subjects though there was clear discrepancy between the model and the clinical trajectory – see for example Figure S2. We defined the inflection of the trajectory based on PASI measurement error that allowed us to categorise subjects undergoing flare which has clinically relevant applications. We went onto test the hypothesis that these discrepancies were due to clinical flares initiated by the occurrence of a new immunological stimulus during therapy. Iterative flare-enabled modelling provided insight into the relative strength and duration of the stimulus. Moreover, our *in silico* analysis combined with clinical data sets showed that the occurrence of flares during therapy was associated with shorter remission.

Strengths and Limitations   Our model is computationally efficient: simulations generally take only a few seconds on a standard desktop or laptop computer. Furthermore, with only two tuneable parameters – UVB sensitivity and immune stimulus – our

Figure 6: **Our ODE model enables identification of psoriasis flares.** Panels: (a) - distribution of relative PASI difference with respect to absolute PASI difference computed by DBSCAN with $\epsilon = 0.5$ and minimum number of points 20 (red crosses – outliers, and black triangles – main cluster) and a possible curve separating flares (all the points above the blue line) and PASI measurement errors (all point below the line); (b,c) - examples of fitting immune stimuli for two different patients, (d) - patient groups based on model predictions of psoriasis relapse: our personalised flares-enabled model predicts Remission (n=49) and Relapse (n=13) groups.

model is able to fit with high accuracy real PASI data, including flares during the therapy. The UVB sensitivity parameter could be estimated at baseline by clinical biomarkers or at week 3 during UVB phototherapy by simple PASI readings. Our model supports personalised therapy outcome prediction by being able to include UVB doses administered in the clinic and different delivery patterns (*i.e.,* 3x weekly *vs.* 5x weekly). Finally, our model is highly generaliseable: it already has the foundations necessary to accommodate other therapies that block the immune stimulus (e.g. anti-IL-17 or anti-IL-23 antibodies) or induce growth arrest.

As for limitations, we remark that the PASI errors clustering can vary depending on the utilised clustering algorithm and its hyper-parameters. This underscores the need for further experimental work to identify the biomarkers that are associated with disease flares and such studies can now be guided by our computational model. Although not strictly a limitation of our modelling, one should keep in mind that PASI does not provide an objective, completely reliable assessment of psoriasis severity. By definition, it is at least in part observer dependent, and this could lead to significant differences between predicted and observed outcomes in the clinic. Finally, our model provides a broader picture of the compartments within epidermis, and therefore, it does not take into account certain specific spatial details. For example, it cannot distinguish between the actively dividing and the dormant stem cells and transit-amplifying cells. Also, cell differentiation is modelled as a single step process while in reality it involves multiple stages.

Future Work Although biologic therapy has not been considered in the current work, the explicit representation of immune cells (DC, T) and mediators (TNF, IL-17, IL-23) opens up the possibility of simulating the effects of biologics by adapting the current model. In addition, we suggest that further studies to identify biomarkers (*e.g.,* obtained from blood or RNA-sequencing of biopsies) associated with the proposed UVB sensitivity parameter should be conducted. Data from further clinical studies should also be used to refine and improve the model parameters.

Finally, validating the model in a clinical setting will open up its use for personalised treatment of psoriasis in practice.

Code and Data Availability

Matlab code for our model is available at `https://github.com/pzuliani/psoriasis/`.

References

[1] C. Albanesi, S. Madonna, P. Gisondi, and G. Girolomoni. The interplay between keratinocytes and immune cells in the pathogenesis of psoriasis. *Frontiers in Immunology*, 9:1549, 2018.

[2] A. Armstrong, K. Parsi, C. Schupp, P. Mease, and K. Duffin. Standardizing training for psoriasis measures effectiveness of an online training video on Psoriasis Area and Severity Index assessment by physician and patient raters. *JAMA Dermatology*, 149:1–6, 02 2013.

[3] M. Athar, A. L. Kim, N. Ahmad, H. Mukhtar, J. Gautier, and D. R. Bickers. Mechanism of Ultraviolet B-Induced Cell Cycle Arrest in G2/M Phase in Immortalized Skin Keratinocytes with Defective p53. *Biochemical and Biophysical Research Communications*, 277(1):107–111, 2000.

[4] Z. Bata-Csorgo, C. Hammerberg, J. J. Voorhees, and K. D. Cooper. Flow cytometric identification of proliferative subpopulations within normal human epidermis and the localization of the primary hyperproliferative population in psoriasis. *Journal of Experimental Medicine*, 178(4):1271–1281, 10 1993.

[5] J. Bauer, F. A. Bahmer, J. Wörl, W. Neuhuber, G. Schuler, and M. Fartasch. A strikingly constant ratio exists between langerhans cells and other epidermal cells in human skin. A stereologic study using the optical disector

method and the confocal laser scanning microscope. *Journal of Investigative Dermatology*, 116(2):313 – 318, 2001.

[6] W.-H. Boehncke and M. P. Schon. Psoriasis. *The Lancet*, 386(9997):983 – 94, 2015.

[7] J. B. Cheng, A. J. Sedgewick, A. I. Finnegan, P. Harirchian, J. Lee, S. Kwon, M. S. Fassett, J. Golovato, M. Gray, R. Ghadially, W. Liao, B. E. Perez White, T. M. Mauro, T. Mully, E. A. Kim, H. Sbitany, I. M. Neuhaus, R. C. Grekin, S. S. Yu, J. W. Gray, E. Purdom, R. Paus, C. J. Vaske, S. C. Benz, J. S. Song, and R. J. Cho. Transcriptional programming of normal and inflamed human epidermis at single-cell resolution. *Cell Reports*, 25(4):871–883, 2018.

[8] M. S. Chimenti, F. Sunzini, L. Fiorucci, E. Botti, G. L. Fonti, P. Conigliaro, P. Triggianese, L. Costa, F. Caso, A. Giunta, M. Esposito, L. Bianchi, R. Santucci, and R. Perricone. Potential role of cytochrome c and tryptase in psoriasis and psoriatic arthritis pathogenesis: Focus on resistance to apoptosis and oxidative stress. *Frontiers in Immunology*, 9:2363, 2018.

[9] R. S. Dawe, N. J. Wainwright, H. Cameron, and J. Ferguson. Narrow-band (TL-01) ultraviolet B phototherapy for chronic plaque psoriasis: three times or five times weekly treatment? *British Journal of Dermatology*, 138(5):833–839, 1998.

[10] C. De Jesús-Gil, L. Sans-de San Nicolás, E. Ruiz-Romeu, M. Ferran, L. Soria-Martinez, A. Chiriac, A. Celada, R. M. Pujol, and L. F. Santamaria-Babí. Specific IgA and CLA+ T-Cell IL-17 Response to *Streptococcus pyogenes* in Psoriasis. *Journal of Investigative Dermatology*, 140(7):1364–1370.e1, 2020.

[11] F. K. Doger, E. Dikicioglu, F. Ergin, E. Unal, N. Sendur, and M. Uslu. Nature of cell kinetics in psoriatic epidermis. *Journal of Cutaneous Pathology*, 34(3):257–263, 2007.

[12] M. Ester, H.-P. Kriegel, J. Sander, and X. Xu. A density-based algorithm for discovering clusters in large spatial databases with noise. In *Proceedings of the Second International Conference on Knowledge Discovery and Data Mining*, KDD'96, page 226–231. AAAI Press, 1996.

[13] K. Eyerich, S. J. Brown, B. E. Perez White, R. J. Tanaka, R. Bissonette, S. Dhar, T. Bieber, D. J. Hijnen, E. Guttman-Yassky, A. Irvine, J. P. Thyssen, C. Vestergaard, T. Werfel, A. Wollenberg, A. S. Paller, and N. J. Reynolds. Human and computational models of atopic dermatitis: A review and perspectives by an expert panel of the International Eczema Council. *The Journal of Allergy and Clinical Immunology*, 143(1):36–45, Jan 2019.

[14] C. Fink, C. Alt, L. Uhlmann, C. Klose, A. Enk, and H. Haenssle. Intra- and interobserver variability of image-based PASI assessments in 120 patients suffering from plaque-type psoriasis. *Journal of the European Academy of Dermatology and Venereology*, 32(8):1314–1319, 2018.

[15] L. Fry and B. S. Baker. Triggering psoriasis: the role of infections and medications. *Clinics in Dermatology*, 25(6):606–615, 2007.

[16] C. E. Griffiths and J. N. Barker. Pathogenesis and clinical features of psoriasis. *The Lancet*, 370(9583):263 – 271, 2007.

[17] J. Gudjonsson, A. Thorarinsson, B. Sigurgeirsson, K. Kristinsson, and H. Valdimarsson. Streptococcal throat infections and exacerbation of chronic plaque psoriasis: a prospective study. *British Journal of Dermatology*, 149(3):530–534, 2003.

[18] A. R. Haake and R. R. Polakowska. Cell death by apoptosis in epidermal biology. *Journal of Investigative Dermatology*, 101(2):107 – 112, 1993.

[19] K. M. Halprin. Epidermal "turnover time"—a re-examination. *British Journal of Dermatology*, 86(1):14–19, 1972.

[20] H. Iizuka. Epidermal turnover time. *Journal of Dermatological Science*, 8(3):215 – 217, 1994.

[21] L. Israel and M. Mellett. Clinical and genetic heterogeneity of CARD14 mutations in psoriatic skin disease. *Frontiers in Immunology*, 9:2239, 2018.

[22] S. Kirke, S. Lowder, J. Lloyd, B. Diffey, J. Matthews, and P. Farr. A randomized comparison of selective broadband UVB and narrowband UVB in the treatment of psoriasis. *The Journal of investigative dermatology*, 127:1641–6, 07 2007.

[23] T. Krausgruber, N. Fortelny, V. Fife-Gernedl, M. Senekowitsch, L. C. Schuster, A. Lercher, A. Nemc, C. Schmidl, A. F. Rendeiro, A. Bergthaler, and C. Bock. Structural cells are key regulators of organ-specific immune responses. *Nature*, 583(7815):296–302, Jul 2020.

[24] D. Kulms and T. Schwarz. Molecular mechanisms of UV-induced apoptosis. *Photodermatology, Photoimmunology & Photomedicine*, 16(5):195–201, 2000.

[25] M. Lowes, A. M Bowcock, and J. G Krueger. Pathogenesis and therapy of psoriasis. *Nature*, 445:866–73, 03 2007.

[26] M. A. Lowes, C. B. Russell, D. A. Martin, J. E. Towne, and J. G. Krueger. The IL-23/T17 pathogenic axis in psoriasis is amplified by keratinocyte responses. *Trends in Immunology*, 34(4):174 – 181, 2013.

[27] M. A. Lowes, M. Suárez-Fariñas, and J. G. Krueger. Immunology of psoriasis. *Annual Review of Immunology*, 32(1):227–255, 2014.

[28] K. R. Mesa, P. Rompolas, G. Zito, P. Myung, T. Y. Sun, S. Brown, D. G. Gonzalez, K. B. Blagoev, A. M. Haberman, and V. Greco. Niche-induced cell death and epithelial phagocytosis regulate hair follicle stem cell pool. *Nature*, 522(7554):94–97, Jun 2015.

[29] L. M. Milstone. Epidermal desquamation. *Journal of Dermatological Science*, 36(3):131 – 140, 2004.

[30] K. Nograles, L. Zaba, E. Guttman-Yassky, J. Fuentes-Duculan, M. Suárez-Fariñas, I. Cardinale, A. Khatcherian, J. Gonzalez, K. Pierson, T. White, C. Pensabene, I. Coats, I. Novitskaya, M. Lowes, and J. Krueger. Th17 cytokines interleukin (IL)-17 and IL-22 modulate distinct inflammatory and keratinocyte-response pathways. *British Journal of Dermatology*, 159(5):1092–1102, 2008.

[31] W. H. Organization. *Global report on psoriasis*. World Health Organization, 2016.

[32] S. Pavey, T. Russell, and B. Gabrielli. G2 phase cell cycle arrest in human skin following UV irradiation. *Oncogene*, 20:6103–6110, 2001.

[33] J.-Z. Qin, V. Chaturvedi, M. F. Denning, P. Bacon, J. Panella, D. Choubey, and B. J. Nickoloff. Regulation of apoptosis by p53 in UV-irradiated human epidermis, psoriatic plaques and senescent keratinocytes. *Oncogene*, 21(19):2991–3002, May 2002.

[34] D. Raj, D. E. Brash, and D. Grossman. Keratinocyte apoptosis in epidermal development and disease. *The Journal of investigative dermatology*, 126(2):243–257, Feb 2006.

[35] G. Reynolds, P. Vegh, J. Fletcher, E. F. Poyner, E. Stephenson, I. Goh, R. A. Botting, N. Huang, B. Olabi, A. Dubois, D. Dixon, K. Green, D. Maunder, J. Engelbert, M. Efremova, K. Polański, L. Jardine, C. Jones, T. Ness, D. Horsfall, J. McGrath, C. Carey, D.-M. Popescu, S. Webb, X.-n. Wang, B. Sayer, J.-E. Park, V. A. Negri, D. Belokhvostova, M. Lynch, D. McDonald, A. Filby, T. Hagai, K. B. Meyer, A. Husain, J. Coxhead, R. Vento-Tormo, S. Behjati, S. Lisgo, A.-C. Villani, J. Bacardit, P. Jones, E. A. O'Toole, G. S. Ogg, N. Rajan, N. J. Reynolds, S. A. Teichmann, F. Watt, and M. Haniffa. Developmental cell programs are co-opted in inflammatory skin disease. *Science*, 371(6527), 2021.

[36] A. Robinson, M. Kardos, and A. Kimball. Physician Global Assessment (PGA) and Psoriasis

Area and Severity Index (PASI): Why do both? A systematic analysis of randomized controlled trials of biologic agents for moderate to severe plaque psoriasis. *Journal of the American Academy of Dermatology*, 66:369–75, 03 2012.

[37] R. Sender and R. Milo. The distribution of cellular turnover in the human body. *Nature Medicine*, 27:45–48, 2021.

[38] H. Valdimarsson, B. S. Baker, I. Jónsdóttir, A. Powles, and L. Fry. Psoriasis: a T-cell-mediated autoimmune disease induced by streptococcal superantigens? *Immunology Today*, 16(3):145 – 149, 1995.

[39] E. J. Van Scott and T. M. Ekel. Kinetics of Hyperplasia in Psoriasis. *Archives of Dermatology*, 88(4):373–381, 10 1963.

[40] N. Watson, N. Wilson, F. Shmarov, P. Zuliani, N. J. Reynolds, and S. C. Weatherhead. The use of psoriasis biomarkers, including trajectory of clinical response, to predict clearance and remission duration to UVB phototherapy. *Journal of the European Academy of Dermatology and Venereology*, submitted, 2021.

[41] S. C. Weatherhead. *The Role of Apoptosis in UVB-Induced Clearance of Psoriasis*. PhD thesis, Newcastle University, 2011.

[42] S. C. Weatherhead, P. M. Farr, D. Jamieson, J. S. Hallinan, J. J. Lloyd, A. Wipat, and N. J. Reynolds. Keratinocyte apoptosis in epidermal remodeling and clearance of psoriasis induced by UV radiation. *Journal of Investigative Dermatology*, 131(9):1916 – 1926, 2011.

[43] G. D. Weinstein, J. L. McCullough, and P. Ross. Cell proliferation in normal epidermis. *Journal of Investigative Dermatology*, 82(6):623 – 628, 1984.

[44] G. D. Weinstein, J. L. McCullough, and P. Ross. Cell kinetic basis for pathophysiology of psoriasis. *Journal of Investigative Dermatology*, 85(6):579 – 583, 1985.

[45] G. D. Weinstein and E. J. van Scott. Autoradiographic analysis of turnover times of normal and psoriatic epidermis. *Journal of Investigative Dermatology*, 45(4):257 – 262, 1965.

[46] N. Wilkinson, T. Tsakok, N. Dand, K. Bloem, M. Duckworth, D. Baudry, A. Pushpa-Rajah, C. Griffiths, N. Reynolds, J. Barker, R. Warren, A. Burden, T. Rispens, D. Stocken, and C. Smith. Defining the therapeutic range for adalimumab and predicting response in psoriasis: a multicenter prospective observational cohort study. *Journal of Investigative Dermatology*, 139(1):115–123, 1 2019.

[47] S. W. Youn, C. Choi, B. Kim, and J. Chae. Reduction of inter-rater and intra-rater variability in psoriasis area and severity index assessment by photographic training. *Annals of Dermatology*, 27:557–62, 10 2015.

[48] H. Zhang, W. Hou, L. Henrot, S. Schnebert, M. Dumas, C. Heuséle, and J. Yang. Modelling epidermis homoeostasis and psoriasis pathogenesis. *Journal of the Royal Society, Interface*, 12:20141071, 02 2015.

Competing Interests

N.J.R. reports grants from PSORT industrial partners as listed (http://www.psort.org.uk/); other research grants from Novartis; and other income to Newcastle University from Almirall, Janssen, Lilly, Leo Pharma, Novartis and UCB Pharma Ltd for lectures/attendance at advisory boards. The remaining authors declare no competing interests.

Author Contributions

F.S. developed the model, ran the simulations, analysed the results and wrote a draft manuscript; G.R.S. contributed improvements to the model; N.J.R. and S.C.W. provided the clinical data; P.Z. and N.J.R. conceived the project, supervised the research and

contributed to writing the manuscript. All authors discussed the results and approved the final manuscript.

Acknowledgments

F.S. was supported by a Rosetrees Trust award while at Newcastle University. This study received funding from the National Institute for Health Research (NIHR) Newcastle Biomedical Research Centre, the British Skin Foundation, the British Phototherapy Group. N.J.R. is also supported by the Newcastle NIHR Medtech and In Vitro Diagnostics Co-operative and is a NIHR Senior Investigator. We are grateful to the research and phototherapy nurses who helped throughout this study and to the patients who participated.

SM   Supplementary Material

SM.1   Model details

We used the SMT solver dReal to show that there are only 3 balls of diameter at most $3 \cdot 10^{-9}$ in the domain $[10^{-9}, 10^{32}] \times \cdots \times [10^{-9}, 10^{32}]$ that can contain the solutions. (Although this does not formally guarantee that there are at least 3 solutions.) So if there are more than one solution within each such ball (there are 32 solutions overall), they are at most $3 \cdot 10^{-9}$ far from each other. Below are the intervals where the solutions for each species might be. So basically there are no solutions in the regions $\pm 10^{-9}$ from the right and the left bounds of the intervals below.

Solution intervals group 1:

$$A \in [7.982009560747935, 7.982009560748187]$$
$$D \in [50529.35801361269, 50529.35801361356]$$
$$DC \in [1563.063946375348, 1563.063946375427]$$
$$GF \in [717.0296163340762, 717.0296163341161]$$
$$IL_{17/22} \in [21.31633768900653, 21.31633768900844]$$
$$IL_{23} \in [42.82366976370817, 42.82366976371033]$$
$$SC \in [3947.21384224557, 3947.213842245908]$$
$$T \in [1556.092651297535, 1556.092651297615]$$
$$TA \in [22224.36715394875, 22224.36715394933]$$
$$TNF \in [21.31633768900732, 21.31633768900842]$$

Solution intervals group 2:

$$A \in [9.942521279320136, 9.942521279321062]$$
$$D \in [63458.54551987114, 63458.54551987203]$$
$$DC \in [1905.499517136538, 1905.499517136938]$$
$$GF \in [881.2100494062016, 881.210049406315]$$
$$IL_{17/22} \in [25.98631442285044, 25.9863144228516]$$
$$IL_{23} \in [52.20546622292645, 52.20546622292981]$$
$$SC \in [4855.45800768624, 4855.458007687109]$$
$$T \in [1897.000952867967, 1897.00095286809]$$
$$TA \in [27308.70879564229, 27308.7087956429]$$
$$TNF \in [25.9863144228494, 25.98631442285055]$$

Solution intervals group 3:

$$A \in [26.59853719540063, 26.59853719540152]$$
$$D \in [173504.1395710201, 173504.139571021]$$
$$DC \in [4804.398597448669, 4804.398597449334]$$
$$GF \in [2271.064736922396, 2271.064736922519]$$
$$IL_{17/22} \in [65.52014914891787, 65.52014914891896]$$
$$IL_{23} \in [131.6273588342101, 131.6273588342123]$$
$$SC \in [12544.22751851258, 12544.22751851354]$$
$$T \in [4782.970887871005, 4782.970887871085]$$
$$TA \in [70349.63537915527, 70349.63537915586]$$
$$TNF \in [65.52014914891788, 65.52014914891897]$$

SM.2 Growth and differentiation rate of keratinocytes

The growth rate and the differentiation rate of keratinocytes are the positive terms of the differential equations (see Figure (1c)) for cell species SC, TA, and D. The terms are $(TNF\alpha + IL_{17/22})(sc_2 SC + sc_{2ta} SC + ta_2 TA)$ and $ta_{2d}(TNF\alpha + IL_{17/22})TA$ for the growth rate and differentiation rate, respectively. Thus, the change of keratinocytes growth rate in the two stable states of the model (healthy and psoriasis) can be calculated as the ratio:

$$\Delta_{KC} = $$
$$\frac{(TNF\alpha^P + IL_{17/22}^P)(sc_2 SC^P + sc_{2ta} SC^P + ta_2 TA^P)}{(TNF\alpha^H + IL_{17/22}^H)(sc_2 SC^H + sc_{2ta} SC^H + ta_2 TA^H)}$$

where the superscripts P and H represent the values of the model species in the psoriatic and the healthy steady states, respectively. By utilising the values found in Table S2 we compute $\Delta_{KC} \approx 9.74$.

Similarly, the change in the differentiation rate is $\Delta_D = \frac{ta_{2d}(TNF\alpha^P + IL_{17/22}^P)TA^P}{ta_{2d}(TNF\alpha^H + IL_{17/22}^H)TA^H} \approx 9.73$.

SM.3 Keratinocytes growth *vs.* epidermis cell density

In our model we assume that cell growth rates depend on the cell density. This assumption was made from observing the dependency between epidermal turnover time and epidermal thickness. Turnover time is the amount of time needed to replace all cells in a particular compartment. It can be calculated as the total number of cells in the compartment divided by the rate of cell growth (cell death for systems in homeostasis). It has been reported that the thickness of psoriatic epidermis increases by around 2-5 times [42, SM], while the epidermal turnover time decreases by up to 4 times [42, SM].

Suppose $\tau_H = \frac{\Sigma_H}{\mu_H}$ and $\tau_P = \frac{\Sigma_P}{\mu_P}$ (where Σ_H and Σ_P are total cell densities in healthy and psoriatic epidermis, and μ_H and μ_P are the rates of cell growth in healthy and psoriatic epidermis) are the turnover times for healthy and psoriatic epidermis, respectively. Typically, the growth rate is described by the law of mass action as $k \cdot \Sigma$, where k is the growth rate constant. Essentially, it states that the cell growth rate depends linearly on the cell density. However, we will show why the law of mass action is not directly applicable in our model without making further assumptions (*e.g.,* growth limiting factors, forces, viscosity, limited nutrition, etc.).

Let $n = \frac{\Sigma_P}{\Sigma_H}$ be the ratio between the thickness of psoriatic and healthy epidermis, and let $m = \frac{\tau_H}{\tau_P}$ be the turnover time ratio between psoriatic and healthy epidermis. From the above it follows that $\mu_P = m \cdot n \cdot \mu_H$, which means that epidermal thickness increases n-fold while the cell growth rate increases by $m \cdot n$. This suggests that the growth rates μ_P and μ_H cannot be modelled by the law of mass action as we derive the following contradiction $\Sigma_P = n \cdot \Sigma_H$ (from the thickness ratio formula) and $\Sigma_P = m \cdot n \cdot \Sigma_H$ (from the growth rates equation).

However, if we assume that the cell growth rate is defined by a nonlinear function $\mu = k \cdot \Sigma^p$, where $p > 1$, we get the following system of equations $\Sigma_P = n \cdot \Sigma_H$ and $\Sigma_P^p = m \cdot n \cdot \Sigma_H^p$. Solving it for p yields $p = (\log_n m) + 1$. Thus, we model keratinocyte growth using the following ODE

$$\frac{dX}{dt} = k_1 \cdot X \cdot IL - k_2 \cdot X^p - k_3 \cdot X,$$

where the term $k_1 \cdot X \cdot IL$ describes the growth rate of keratinocytes as a law of mass action with X being the keratinocyte species and IL being the cytokine

Table S1: Model parameters

Name	Description	Value	Units
il_{172}	IL-17 production rate constant by T cells	0.5	d^{-1}
il_{1720}	IL-17 degradation rate constant	36.5	d^{-1}
il_{232}	IL-23 production rate constant by dendritic cells	1	d^{-1}
il_{2320}	IL-23 degradation rate constant	36.5	d^{-1}
tnf_2	TNF production rate constant by T cells	0.5	d^{-1}
tnf_{20}	TNF degradation rate constant	36.5	d^{-1}
t_2	T cells activation rate constant	55	d^{-1}
t_{20}	T cells deactivation rate constant	1.51	d^{-1}
sc_{2gf}	GF production rate constant by SC	1	d^{-1}
ta_{2gf}	GF production rate constant by TA	1	d^{-1}
gf_{20}	GF degradation rate constant	36.5	d^{-1}
sc_2	SC production rate constant	0.0017635	d^{-1}
sc_{2ta}	TA production rate constant by SC	0.0015415	d^{-1}
sc_{20}	Proliferation limiting factor for SC	1.8135e-05	d^{-p}
p	-	2	-
ta_2	TA production rate constant	0.0017562	d^{-1}
ta_{20}	Proliferation limiting factor for TA	3.7321e-06	d^{-p}
ta_{2d}	Differentiation rate constant	0.0018063	d^{-1}
d_{20}	Differentiation limiting factor	5.05e-07	d^{-p}
d_{desq}	Desquamation rate constant	0.0047529	d^{-1}
a_{base}	Basal apoptosis rate	0.0036	d^{-1}
a_{20}	Degradation rate of apoptotic cells	36	d^{-1}
t_{vm}	Maximum activation rate constant of DC	6000	-
t_{kp}		$3 \cdot 10^{12}$	-
dc_{act}	Basal activation rate constant of DC	1880	$cells/d$
dc_{20}	Deactivation rate constant of DC	1.51	d^{-1}
n	-	4	-

Table S2: Steady states of the model

Species	Description	Healthy	Transition	Psoriatic
SC	Stem cells	3,947.21	4,855.46	12,544.23
TA	Transit-amplifying cells	22,224.37	27,308.71	70,349.64
D	Differentiating cells	50,529.36	63,458.55	173,504.14
T	T cells	1,556.09	1,897	4,782.97
DC	Dendritic cells	1,563.06	1,905.5	4,804.4
A	Apoptotic cells	7.98	9.94	26.6
Σ	Total cell count	79,828.07	99,435.16	266,011.98
$IL_{17/22}$	Interleukin 17/22	21.32	25.99	65.52
IL_{23}	Interleukin 23	42.82	52.20	131.63
TNF	Tumor necrosis factor alpha	21.32	25.99	65.52
GF	Keratinocyte-based growth factors	717.03	881.21	2,271.06

species, $k_3 \cdot X$ represents cell degradation also modelled by the law of mass action, and $k_2 \cdot X^p$ describes the limiting factors of keratinocyte growth due to an increase in cell density (this could be caused by nutrients limitation and spatial constraints, for example); we call p the "growth limiting constant". In addition, we assume that $n = m$ (and hence, $p = 2$) in our model for simplicity. However, the epidermal thickness and the cell growth rate can be changed using parameters n and m, which provides modelling flexibility.

SM.4   Apoptosis and Desquamation Rates

The dynamics of apoptotic cells (A) is governed by the equation $\frac{dA}{dt} = a_{base}(SC + TA + DC + T + D) - a_{20}A$. It is assumed that in healthy epidermis all cells undergo apoptosis at the same rate (relative to their cell mass). The rate of degradation of apoptotic cells is governed by the term $a_{20} \cdot A$, hence their mean lifetime is $\tau_{apop} = \frac{1}{a_{20}} = 40\ mins$ ([41, Figure 5-5] reports a median time of 20 minutes from an *in vitro* study, [18, Figure 1] states 1-3 hours for lymphocytes and up to 48-72 hours for keratinocytes), and [28, Figure 1d] demonstrates an apoptotic cell being destroyed within 20-40 minutes in mice). The apoptosis rate in terms of num-

ber of apoptotic cells per 1,000 can be calculated as $\frac{1000 \cdot A}{SC + TA + DC + T + D + A} \approx 0.1$ (the median number of apoptotic cells per 1,000 is reported to be 0.1 in untreated psoriatic skin [42, Figure 1c]). The epidermal desquamation rate in the model can be calculated as $d_{desq} \cdot D$, and in the healthy state it is $d_{desq} \cdot D_H \approx 240.16\ cells/(day \cdot mm^2)$, where D_H is the density of differentiated cells in the healthy steady state. A similar figure for human skin is reported by [29].

SM.5   Supplementary figures

Figure S1: **Model simulation of UVB-induced cell apoptosis *vs.* cell growth arrest.** The Apoptosis model reaches the healthy steady state at the end of the UVB therapy, while the Growth arrest model is still in moderate psoriasis at the end of the therapy and an eventual psoriasis clearance. The latter contradicts clinical observations, and it was thus disregarded.

Figure S2: **Modelling flares by introducing an immune stimulus to the model.** Panels (a) and (c): two patients PASI trajectories and their model simulation without immune stimulus. Panels (b) and (d): the same two patients with their respective models fitted with immune stimuli, allowing a much closer match between patient data and model simulation. Panels (b) and (d) are the same as Figure 6b and Figure 6c, respectively: they are reported here for the reader's convenience.