Comparative analysis of genome-encoded viral sequences reveals the evolutionary

history of flavivirids (family *Flaviviridae*).

Connor G. G. Bamford[1]*, William M. de Souza[2]*, Rhys Parry[3]

and Robert J. Gifford[4]

*Equal contributions

[1] *Wellcome-Wolfson Institute for Experimental Medicine, School of Medicine,*

Dentistry and Biomedical Sciences, Queen's University Belfast, Belfast, UK

[2] *World Reference Center for Emerging Viruses and Arboviruses and Department of*

Microbiology and Immunology, University of Texas Medical Branch, Galveston, TX,

USA

[3] *School of Chemistry and Molecular Biosciences, University of Queensland,*

Brisbane, QLD 4072, Australia

[4] *MRC-University of Glasgow Centre for Virus Research, Glasgow, UK*

Corresponding author:

Robert J. Gifford,

MRC-University of Glasgow Centre for Virus Research,

464 Bearsden Rd, Bearsden, Glasgow G61 1QH

E-mail: robert.gifford@glasgow.ac.uk

Keywords: Flavivirus | Jingmenvirus | Paleovirology | Genomics | Flavivirid |

Evolution | Tamanavirus | Arbovirus | Vector | Mosquito | Tick | Arthropod

ABSTRACT

The flavivirids (family *Flaviviridae*) are a group of positive-strand RNA viruses

that pose serious risks to human and animal health on a global scale. Here we use

flavivirid-derived DNA sequences, identified in animal genomes, to reconstruct the

long-term evolutionary history of family *Flaviviridae*. We demonstrate that flavivirids

are >100 million years old and show that this timing can be combined with dates

inferred from co-phyletic analysis to produce a cohesive overview of their evolution,

distribution and diversity wherein the main flavivirid subgroups originate in early

animals and broadly co-diverge with major animal phyla. In addition, we reveal

evidence that the 'classical flaviviruses' of vertebrates, most of which are transmitted

via blood-feeding arthropod vectors, originally evolved in hematophagous arachnids

and later acquired the capacity to be transmitted by insects. Our findings imply that

the biological properties of flavivirids have been acquired gradually over the course of

animal evolution. Thus, broad-scale comparative analysis will likely reveal

fundamental insights into their biology. We therefore published our results via an

open, extensible, database (Flavivirid-GLUE), which we constructed to facilitate the

wider utilisation of genomic data and evolution-related domain knowledge in flavivirid

research.

INTRODUCTION

The flavivirids (family *Flaviviridae*) are an important group of RNA viruses incorporating numerous pathogens of humans and animals. Currently, four genera are recognised within the family: *Pegivirus, Pestivirus*, *Hepacivirus* and *Flavivirus* [1]. While pegiviruses are not known to be associated with disease, pestiviruses cause serious illness in domestic ungulates such as cattle and pigs [2], and the *Hepacivirus* genus includes the bloodborne hepatitis C virus (HCV), a major cause of chronic liver disease in human populations throughout the world [3, 4]. Moreover, the genus *Flavivirus* includes viruses that are transmitted between vertebrates via blood-feeding arthropod vectors (e.g., mosquitoes, ticks) and cause large-scale outbreaks resulting in millions of human infections every year - e.g., yellow fever virus (YFV), dengue viruses 1-4 (DENV1-4), Zika virus.

The *Pegi-, Pesti-*, *Hepaci-* and *Flavivirus* genera contain viruses with monopartite genomes ~10 kilobases (Kb) in length and encoding one or more large polyproteins that are co- and post-translationally cleaved to generate mature virus proteins. The structural proteins of the virion - capsid (C), premembrane (prM) and envelope (E) - are encoded toward the 5' end of the genome, while genes encoding non-structural (NS) proteins are located further downstream [5]. However, a diverse variety of novel 'flavivirus-like' viruses (flavivirids) have been identified in recent years and these viruses – most of which were identified in invertebrates and have yet to be incorporated into official taxonomy - exhibit a much greater range of variation in genome structure, with genome lengths ranging up to 20Kb [6-10]. Furthermore, one novel group – 'jingmenvirus' – comprises viruses with genomes that are multipartite rather than monopartite [11]. Some tick-associated jingmenviruses have been linked with disease in humans [12, 13].

To prevent the spread of pathogenic viruses, it is helpful to understand their evolutionary history in as much detail as possible, because this can often provide crucial insights into virus biology and host-virus relationships [14]. Flavivirids are a

taxonomically diverse group that have been extensively examined using comparative approaches, revealing uniquely clear correlations between phylogenetic relationships and ecological characteristics [15-18]. The current, rapid accumulation of genome sequence data from novel flavivirids offers unprecedented opportunities to build on these comparative studies. Presently, however, there are two major obstacles to efficient use of flavivirid genome data.

Firstly, there is a general lack of re-use and reproducibility among comparative analysis of virus genomes, especially when deeper evolutionary relationships are being examined [19]. These analyses typically entail the assembly of complex data sets comprised of virus sequences, multiple sequence alignments (MSAs) and phylogenies linked to other, diverse kinds of data (e.g., spatiotemporal coordinates, taxonomy, immunity-related information). In theory, these data sets could be re-used across a wide range of analysis contents while also being collaboratively developed and refined by multiple contributors. This would likely accelerate knowledge discovery and expedite the development of expert systems utilising virus genome data. Unfortunately, however, such practices remain challenging to implement in practice, largely due to a lack of appropriate tools and data standards [20].

Secondly – knowledge of the appropriate evolutionary timescale is critically lacking. So far, most studies of flavivirid evolution have proposed relatively short timelines in which individual genera and sub-groups emerge within the last 10-100 thousand years [15-17, 21]. However, these studies were based on viruses sampled from a relatively restricted range of hosts. By contrast, recent studies utilising metagenomic techniques to sample flavivirid diversity across a broader range of animal species have prompted suggestions of a much longer timeline extending over hundreds of millions of years [7]. Furthermore, for many RNA virus families, robust evidence for ancient origins has come in the form of *endogenous viral elements* (EVEs) - virus-derived sequences found within eukaryotic genomes [22]. These

sequences are thought to originate when infection of germline cells leads to virus-

derived cDNA being incorporated into chromosomal DNA so that integrated viral

genes are not only inherited as host alleles, but also persist in the gene pool over

many generations until they are genetically fixed (i.e., reach a frequency of 100% in

the species gene pool). Genome comparisons show that EVE loci are often present

as orthologs in closely related host species, establishing their ancient origins (since

they were incorporated into the host germline prior to species divergence). EVEs

derived from flavivirids have been identified in a handful of arthropod species [8, 23-

25], but are relatively uncommon compared to EVEs derived from other RNA virus

families [26]. Partly reflecting this scarcity, robust calibrations of the long-term

evolutionary timeline of flavivirids are lacking.

In this study, we calibrate the long-term evolutionary timeline of flavivirids,

making extensive use of EVEs. In addition, we construct a cross-platform, interactive

database called 'Flavivirid-GLUE', which we use to capture the evolution-related

domain knowledge generated in our study in a way that facilitates downstream use.

RESULTS

Creation of open resources for comparative genomic analysis of flavivirids

We previously developed a software framework called GLUE ("Genes Linked

by Underlying Evolution") [27]. Here, we used GLUE to create Flavivirid-GLUE [28] -

a flexible, extensible, and openly accessible resource for comparative analysis of

flavivirid genomes (**Fig. S1a-b**). The Flavivirid-GLUE project includes: (i) a set of 237

reference genome sequences each representing a distinct flavivirid species and

linked to isolate-associated data (**Table S1**) (ii) a standardized set of 81 flavivirid

genome features (**Table S2**); (iii) genome annotations specifying the coordinates of

genome features within selected 'master' reference genome sequences (**Table S3**);

(iv) a set of hierarchically arranged MSAs constructed to represent distinct taxonomic

ranks within the family *Flaviviridae* (**Table 1**, **Fig. 1**).

We used hierarchically linked MSAs to enable standardised genome

sequence comparisons across the entire *Flaviviridae* family (i.e., both within and

between taxonomic ranks). For each taxonomic rank represented in the project, one

reference genome was selected as the constraining 'master' reference that defines

the genomic coordinate space. Within the MSA hierarchy, each MSA is linked to its

child and/or parent MSAs via our chosen set of references. MSAs representing

internal nodes (see **Table 1**) contain only master reference sequences but can be

recursively populated with all taxa contained in child alignments via GLUE's

command layer [27]. Importantly, the use of an MSA hierarchy simplifies analysis of

novel taxa (since new sequences only need be aligned to the most closely related

reference genome to be aligned with all other taxa included in the MSA hierarchy).

Instantiation of the Flavivirid-GLUE project (**Fig. S2**) generates a relational

database that contains the data items required for comparative analysis of flavivirids

and represents the semantic links between them. This allows comparative analyses

to be implemented in a standardised, reproducible way, wherein GLUE's command

layer is used to coordinate interactions between the Flavivirid-GLUE database and

bioinformatics software tools (e.g., see **Fig. S3-S4**). Flavivirid-GLUE can be installed

on all commonly-used computing platforms, and is fully containerised via Docker [29].

Hosting in an openly accessible online version control system (GitHub) provides a

platform for coordinating ongoing development of the resource – e.g., incorporation

of additional taxa and genome annotations - following practices established in the

software industry (**Fig. S1c**) [30].

Mapping the distribution of flavivirid-derived DNA in animal genomes

To identify flavivirid-derived EVEs, we performed systematic *in silico*

screening of whole genome sequence data representing 1075 animal species. This

led to the identification of 374 EVE loci in 36 animal species (**Table 2**, [28]). We reconstructed consensus sequences representing fragments of the genomes of (presumably) extinct flavivirids, utilising EVE sequences putatively derived from a single germline incorporation event (i.e., orthologs, fragments, duplicates) (**Fig. 2**, **Fig. S5**). EVE consensus sequences, MSAs, and all EVE-associated metadata were incorporated into Flavivirid-GLUE [28].

All major flavivirid lineages are represented in the host germline

We reconstructed the evolutionary relationships between EVEs and contemporary flavivirids using maximum likelihood approaches (**Fig. 3**, **Fig. S6**). Bootstrapped phylogenetic trees were reconstructed from MSAs representing a range of taxonomic ranks within the *Flaviviridae* (**Table 1**), utilising EVE sequences often contain mutations acquired through neutral reconstructed phylogenies both with MSAs containing virus sequences only, and MSAs incorporating both virus and EVE sequences [28]. Consistent with official taxonomy and previously published studies [1, 18] phylogenetic reconstructions split flavivirids into two major lineages - 'pegi-hepaci' and 'flavi-pesti' – each of which contains several well-supported subgroups [10] (**Fig. 3a-e, Fig. S3**). However, a divergent, flavivirid-derived EVE identified in the genome of a priapulid worm (*Priapulus caudatus*: Lamarck, 1816) may represent a third sub-lineage (**Fig. 3b**).

Phylogenetic reconstructions revealed that EVEs derived from a broad range of flavivirus lineages and subgroups are represented in the genomic 'fossil record'. Only one EVE derived from the 'pegi-hepaci' lineage was identified. However, it occurs in a marine mollusc - the Eastern emerald elysia (*Elysia chlorotica*: Gould, 1870) - demonstrating that the host range of this major flavivirid group extends to invertebrates (**Fig. 3a**).

The majority of flavivirid EVEs derived from the 'flavi-pesti' lineage, which is comprised of robustly supported 'flavi-like' and 'pesti-like' clades. Our approach to

phylogenetic reconstruction, which entails reconstructing separate phylogenies for

distinct taxonomic ranks (see **Table 1**), supports a clean division of 'flavi-like' viruses

into three monophyletic clades (**Fig. 3d**) corresponding to genus *Flavivirus*, the

'jingmenviruses', and a clade of viruses related to Tamana bat virus (TABV) which

we here refer to as 'tamanaviruses'. We identified several EVEs that grouped

robustly within the diversity of contemporary 'tamanavirus' and 'jingmenvirus'

isolates. We also identified EVEs derived from a more distantly related,

'jingmenvirus-like' lineage – here labelled X2 – with no known contemporary

representatives (**Fig. 3f**). Notably, we identified X2-derived EVEs in a temorid

copepod (*Eurytemora affinis*: Poppe, 1880) as well as in multiple actinopterygiid fish,

indicating that their host range encompasses both vertebrate and arthropod species

(**Fig. 3f, Table 1**). The copepod element exhibited internal duplications and

rearrangements typical of EVEs identified in piRNA clusters (data not shown) [31].

The 'pesti-like' lineage contains two robustly-supported clades – one

comprised of vertebrate viruses (including the canonical members of genus

Pestivirus) while the other contains a diverse assortment of invertebrate-associated,

'large-genome flavivirids' [10]. The invertebrate clade contains two well-supported

subclades, here labelled 'PL1' and 'PL2'. The 'PL2' clade contains EVEs in addition

to viruses (**Fig. 3c**).

Germline incorporation of flavivirid DNA is relatively rare

While flavivirid-derived EVEs were only identified in a small proportion of the

animal species we screened, they occur at relatively high copy number in the

germline of some insect groups, including mosquitoes of genus *Aedes* (Meigen,

1818) as well as bees and sphecoid wasps (superfamily Apoidea: Latreille, 1802)

(**Table 1**).

In the *Aedes* germline, flavivirid EVEs are closely related to contemporary

flaviviruses – specifically the 'classical insect-specific' flaviviruses' (cISFs).

Numerous, distinct loci occur, largely representing distinct regions of the flavivirus genome (**Fig. S5**). However, where they do span homologous regions of the flavivirus genome loci are highly related (i.e., <1% nucleotide sequence divergence) and are frequently arranged as tandemly repeating arrays (data not shown), suggesting recent, intra-germline amplification.

Among *Apoidea* EVEs, copy number was dramatically inflated in certain species, such as the spurred ceratina (*Ceratina calcarata*: Robertson, 1900), which contains at least 86 distinct flavivirid-derived EVE sequences in its published genome sequence. However, confident estimates of EVE copy number in *Apoidea* species could not be obtained, due to the limitations of current genome assemblies.

Major flavivirid lineages originated in the distant evolutionary past

We used a range approaches to calibrate the evolutionary timeline of flavivirids (**Table 3**). Calibrations based on identification of orthologous EVEs were obtained for 'jingmenvirus'-derived EVEs found in midges (*Chironomus*: Meigen, 1803), X2-derived EVEs in ray-finned fish (Class Actinopterygii: Klein, 1885), and a PL2-derived EVE in superfamily Apoidea. Orthologous, PL2-derived EVEs were identified in Apoidea species estimated to have diverged >100 million years ago (Mya) (**Fig. S7**). We also derived age estimates ranging between 3-62 Mya for pairs of putatively duplicated EVE sequences based on the assumption a neutral molecular clock (**Table 3**). These calibrations, combined with the identification of flavivirid-derived EVEs in basal animal lineages such as cnidarians and priapulids, suggested that flavivirids could in fact have truly primordial origins in multicellular animals. Such an extended evolutionary timeline would be consistent with other recent data supporting the ancient origins of RNA virus families [7, 32].

Horizontal transfer of flavivirids between distantly related hosts has clearly occurred – most likely in association with parasitism [33, 34] - but when phylogenies are considered in the light of an evolutionary timescale extending back to the origin of

multicellular animals (i.e., ~500-800 Mya), a credible argument can be made for

codivergence being the more common mode of evolution, at least where higher

taxonomic ranks are concerned. To investigate this, we compared host and virus

phylogenies. Scope for comparisons was limited due to sparse data and relatively

narrow sampling across host species groups (with most flavivirids isolated from

arthropods and vertebrates). We identified several host and virus clades in which the

branching relationships and divergence times among animal lineages are correlated

with the topology and branch lengths found in virus phylogenies (**Fig. 4, Fig. S8**).

However, we were required to make strong assumptions in each case, particularly

regarding the rooting of virus trees (see **Fig. 4** legend), and these deeper calibrations

should therefore be taken as tentative.

Arthropod-vectored flaviviruses likely emerged from an arachnid source

The *Flavivirus* genus includes viruses that are transmitted among vertebrate

hosts by arthropod vectors, as well as viruses that exclusively infect arthropods [35]

and others that have been identified in vertebrates but have no known arthropod

vector [36]. The long history of association between flavivirids and their hosts implied

by our investigation suggests the largely vector-borne 'classical flaviviruses' could

have emerged in association with the evolution of hematophagy (blood-feeding) in

arthropods. Phylogenetic reconstructions using either NS5 (**Fig. 3a**) or NS3 (**Fig. S9**)

show that flaviviruses exclusively associated with insects (Class Insecta: Linnaeus,

1758) are robustly separated from those that infect vertebrates by viruses identified

in crustaceans (Subphylum Crustacea: Brünnich, 1772), sea spiders (Pycnogonida:

Latreille, 1810) [37], and arachnids (Arachnida: Lamarck, 1801). In addition, rooted

phylogenies show the most basal CFV lineages are either tick-associated (Mpulungu

virus and the tick-borne CFVs) or have no known vector [36] (**Fig. 3a**). While much

uncertainty remains, it is interesting to note that, if we parsimoniously assume that

the basal lineages without known vectors are tick-borne (or were originally before

subsequently losing their association with arachnids), phylogenies suggest (i)

emergence of tick-borne viruses from arachnid-specific viruses, followed by (ii)

emergence of insect-borne viruses from tick-borne viruses (**Fig. 3b**). Intriguingly, we

identified synapomorphic amino acid variation in the NS5 protein, wherein ancestral

residues are conserved between ancestral, arachnid-associated flaviviruses but

variable in the more derived, insect-vectored flaviviruses (**Fig. 3a**, **Fig. S10**). While

these patterns can of course be interpreted in alternative ways, they are consistent

with positive selection accompanying the adaptation of ostensibly tick-borne viruses

to newly acquired insect vectors.

DISCUSSION

The flavivirids are a genetically and ecologically diverse group of viruses that

include an unusually large number of taxonomically recognised species, many of

which are associated with disease [18]. As a diverse and highly studied group,

flavivirids offer unique possibilities for researchers interested in applying comparative

approaches to viruses [15, 17, 18]. Species richness in this group to some extent

reflects their historical importance in the development of virus research - YFV being

the first human virus identified [38] - and sampling of flavivirid diversity consequently

shows historical bias toward potential vector/reservoir species [39, 40]. However,

with dramatic advances in DNA sequencing technology it is now possible to

investigate flavivirid distribution and diversity much more broadly, building on

previous comparative investigations [15-18].

In this report we address two important challenges to effective use of flavivirid

sequence data in comparative genomic studies. First, we implemented our analyses

using a computational framework that supports re-use of underlying data sets and

facilitates reproduction of comparative genomic analyses. Second, we calibrated the

long-term evolutionary history of flavivirids through use of the 'genomic fossil record',

thereby providing broad evolutionary context for interpreting their distribution and

diversity.

We identify flavivirid-derived EVEs that are >100 My old, demonstrating that

the evolution of family *Flaviviridae* spans geological eras. Furthermore, we show that

the robust calibrations obtained from EVEs can be combined with more tentative

calibrations based on co-phyletic analysis to produce a cohesive overview of flavivirid

evolution in which the major lineages emerged during the early evolution of

multicellular animals and subsequently evolved together with major animal phyla

(**Fig. 5c**). The timeline suggested by our analysis raises interesting questions about

the ways in which animal evolution may have impacted flaviviruses, since it

encompasses the development of entire organ systems (e.g., the liver and vascular

system) and spans the evolution of fundamental changes in animal physiology, such

as the emergence of endothermy ("warm blood") in vertebrates. The identification of

flavivirid-derived EVEs in animals lacking a circulatory system (e.g., cnidarians and

priapulids) suggests that cell-to-cell transmission via exosomes - as has been

reported for tick-borne flaviviruses [41] - might represent the ancestral mode among

flavivirids.

The vertebrate circulatory system evolved >400 Mya, and is thought to have

been established in its basic form in endothermic vertebrates (i.e., birds and

mammals) by ~200 Mya [42]. Its role in transporting nutrients makes it a highly

attractive target for parasitism, and hematophagous, arthropod parasites of

vertebrates are thought to have evolved on >20 independent occasions [43].

Whenever this occurred, it would have created new, intimate contact networks

between arthropod and vertebrate species so that viruses circulating within each

group could potentially encounter opportunities to expand into the other [33].

Interestingly, we identified closely related, X2-derived EVEs in both copepods and

fish, raising the possibility that copepod parasitism has enabled X2-like arthropod

viruses to expand into vertebrate hosts (**Fig. 3f**).

Both phylogenetic relationships within genus *Flavivirus*, and the extended

evolutionary timeline implied by our study, are consistent with a parsimonious

evolutionary scenario wherein the vector-borne classical flaviviruses originate in

hematophagous arachnids and later acquire the capacity to be transmitted by

hematophagous insects (**Fig. 5a-b**). This 'ticks first' model is quite appealing

because the circumstances of tick feeding provide an obvious opportunity for tick

flaviviruses capable of limited replication in vertebrate cells to emerge. Feeding

entails a relatively long exposure, and ticks are thought to transmit infection to one

another via host blood when multiple individuals feed in proximity [44]. Once tick

flaviviruses had acquired the capacity to generate sustained viremia in vertebrate

hosts, opportunities for hematophagous insects to acquire vector roles would

presumably arise as they become exposed to virus via blood-feeding. Interestingly,

this model of flavivirus evolution is consistent with evidence that the 3' untranslated

region (UTR) of arthropod-borne flaviviruses – which can modulate the viral life cycle

in complex and nuanced ways - evolved via duplication [45, 46]. Duplicated UTR

sequences are best preserved in tick-borne flaviviruses, consistent with an ancestral

origin, while only remnants are present in the more derived, mosquito-borne

flaviviruses [47].

Due to the overall scarcity of flavivirid-derived EVEs, our investigation

provides only limited insight into the evolutionary history of certain flavivirid groups,

such as the *Hepaci*-, *Pegi*- and *Pestivirus* genera. However, our identification of

'pegi-hepaci'-like EVEs in invertebrates, and the recent description of a pestivirus-like

EVE in an insectivore [48], indicates that additional, diverse flavivirid EVEs will be

identified as whole genome sequencing of animal species advances, allowing for

more complete perspective on the long-term evolution of flavivirids. New information

obtained through broader sampling of contemporary flavivirid diversity will also allow

for more rigorous testing of the evolutionary hypotheses presented here.

Phylogenetic reconstructions utilising currently available data suggest that

reorganisation of the flavivirid taxonomy should be considered (**Fig. 3**). Currently,

family *Flaviviridae* is placed in order Amarillovirales, class Flasuviricetes, which

contains no other orders or families. Flasuviricetes could be restructured to contain

groups representing the 'pegi-hepaci' and 'flavi-pesti' lineages and the subgroups

contained therein. This would enable taxonomic classifications to reflect the deep

evolutionary splits between flavivirid subgroups, such as those separating

'jingmenviruses', 'tamanaviruses' and flaviviruses (genus *Flavivirus*) in the 'flavi-pesti'

lineage. The enigmatic 'Tamana bat virus' (TABV), which was isolated in 1973 from

insectivorous bats and has no known arthropod vector, has puzzled flavivirologists

for decades [36, 49, 50]. It is sometimes considered a basal member of genus

Flavivirus but is only distantly related to other flaviviruses and clearly distinct in

signature genomic traits such as nucleotide composition [51]. Notably, however, we

identified an EVE in the genome of the peach blossom jellyfish (*Craspedacusta*

sowerbii: Lankester, 1880) that is more 'flavivirus-like' than TABV (**Fig. 3d**). The

identification of flavivirus-like EVEs in cnidaria (Hatschek: 1888), a basal animal

lineage, combined with new information describing the extensive diversity and broad

host range of TABV-related viruses [8, 10, 52, 53] (**Fig. 3e**), suggests

'tamanaviruses' have distinct evolutionary origins to genus *Flavivirus* and should be

given separate taxonomic status among 'flavi-like' viruses. The same applies to

'jingmenviruses', which presumably became evolutionarily separated from other

'flavi-like' viruses when they evolved multipartite genomes [11].

Given the extent of uncharacterised flavivirid diversity and the accelerating

rate of virus discovery, further expansion and revision of flavivirid taxonomy will no

doubt be required as exploration of the virome proceeds - numerous novel and

divergent flavivirids have been described since we originally submitted this report [34,

54] - the taxonomic system proposed above would provide a relatively open structure

capable of accommodating novel flavivirid diversity.

Our analysis indicates that incorporation of flavivirid-derived DNA into animal

germlines is uncommon. However, in those taxa where flavivirid-derived EVEs

occur, they are frequently multicopy, with some insect groups exhibiting relatively

high copy number (**Table 1**). Notably, these include *Aedes* mosquitoes in which

EVEs have been shown to produce PIWI-interacting RNAs (piRNAs) that limit

infection with related viruses [55]. While recent data indicate a role for EVEs in

antiviral defence [25, 31, 56], it remains unclear whether germline 'capture' of virus

sequences represents a dynamic system of heritable antiviral immunity analogous to

that of CRISPR-Cas. It should be noted that the distribution and diversity of cISF-

derived EVEs in *Aedes* mosquito genomes is consistent with germline integration of

genome-length cDNA, followed by intra-genomic amplification/fragmentation of the

integrated sequences (e.g., mediated by transposable elements). Numerous, similar

examples of amplification of virus-derived DNA within the animal germline have been

described, involving a diverse range of virus families [57-60]. Thus, the presence of

numerous distinct, flavivirid-derived EVE loci in the *Aedes* germline could reflect a

relatively small number of germline colonisation events, rather than a dynamic

process of EVE acquisition in association with immunity.

Controlling the spread of flavivirid-associated diseases is a public health

priority and genomic data have a critical role to play in these endeavours [4, 61].

Here, we used the GLUE software framework to capture flavivirid genome sequence

data and evolution-related domain knowledge in a way that supports their future use.

By following principles of 'data-oriented programming', wherein an explicit separation

is maintained between data and the code that operates on it [62], the GLUE

framework can facilitate the implementation of stable data resources that make no

assumptions with respect to their future usage, so they can be deployed in distinct

analysis contexts [27]. Besides acting as a stable repository of domain knowledge,

Flavivirid-GLUE provides a broad foundation for the rapid development of

tools/services (e.g., epidemiological tracking, variant analysis) focussed on individual

flavivirid taxa (e.g., see [63, 64]). In addition, Flavivirid-GLUE can underpin analytical

procedures that require a broader taxonomic scope, such as sequence-based

classification of newly identified flavivirids, or supporting empirical, laboratory-based

investigations of important flavivirid traits (e.g., the capacity to replicate in both

arthropod and vertebrate hosts). If - as our investigation suggests – flavivirid traits

have been acquired gradually through long-term evolutionary interactions with animal

hosts, comparative studies will likely yield many useful insights into their biology.

MATERIALS & METHODS

Construction of sequence-based resources for comparative genomic analysis

We used the GLUE software framework [27] to create Flavivirid-GLUE [28],

an openly accessible online resource for comparative analysis of flavivirid genomes.

The advantages of GLUE include: (i) the organization of virus sequence data in

relation to their hypothesized evolutionary relationships (an approach that is key to

the practical interpretation of sequences); (ii) integration with wider computer

systems, using standard technologies such as MySQL and JSON; (iii) mechanisms

to customize functionality on a usage-specific basis, for example schema extension

and scripting mechanisms; (iv) features and functionality to streamline rapid

development of bespoke analysis projects.

We extended GLUE's core database schema to capture information specific

to flavivirid reference sequences (e.g., virus isolate names, isolation species, date

and location of sampling) and EVE loci (e.g., species in which they occur, genomic

coordinates within contigs/chromosomes). A library of flavivirus representative

genome sequences (**Table S1**) was obtained from GenBank via reference to the

International Committee for Taxonomy of Viruses website (see [28]). GenBank

sequence entries in XML format were imported into the Flavivirid-GLUE project using

an appropriately configured version of GLUE's GenBank importer module. We

extracted isolate-specific information (e.g., date and location of isolation, isolate

name, host species) from XML files as well standard GenBank fields (e.g.,

submission date). Additional/missing data was loaded from tabular files using

GLUE's TextFilePopulator module. Via reference to previous studies [6, 9, 10, 18, 53]

we assigned all flavivirus sequences included in Flavivirid-GLUE to a taxonomic

group and defined a standard set of genome features for flaviviruds. The coordinates

of genome features (where known) within all master references were recorded within

the Flavivirid-GLUE database. These reference sequences and annotations were

used in combination with a codon-aware, BLAST-based sequence aligner

implemented in GLUE [27, 65] to generate constrained MSAs (i.e., MSAs in which

the coordinate space is constrained to a selected 'master' reference) for each

taxonomic rank within the *Flaviviridae* (**Table 1**). To address genome and gene-

coverage-related issues we used GLUE to generate genome feature coverage data

for member sequence. Constrained MSAs were used to infer the coordinates of

genome features not explicitly defined in GenBank XML via GLUE's 'inherit features'

command [27].

Genome screening in silico

Systematic *in silico* genome screening was performed using the database-

integrated genome screening (DIGS) tool [66] – a PERL-based screening framework

within which the basic local alignment search tool (BLAST) program suite [65] is used

to perform similarity searches while the MySQL relational database management

system (Community Server 8.0.26) is used to record their output. WGS data were

obtained from the National Center for Biotechnology Information (NCBI) genome

database [67] - we obtained all animal genomes available as of March 2020 (see

[28]). Flavivirid reference genomes and coding feature annotations collated in

Flavivirid-GLUE were used to derive polypeptide probes for tBLASTn-based

screening in the DIGS framework. For virus genomes lacking detailed annotations,

we created polypeptide probes based on fragments of the major polyprotein. Via

screening of WGS assemblies using the DIGS tool we generated a non-redundant

database of flavivirid-derived EVE loci [28]. We used DIGS to investigate these loci

and categorise them into: (i) putatively novel EFV elements; (ii) orthologs of

previously characterised EFVs (e.g., copies containing large indels); (iii) non-viral

sequences that cross-matched to flavivirus probes (e.g., retrotransposons). In

applying IDs to EVE sequences (see **Fig. 2**, **Table 1**), we conservatively assumed

that the presence of multiple EVE loci generally reflects intragenomic amplification

rather than multiple independent germline incorporation events.

Phylogenetic and genomic analysis

Flavivirid-GLUE was used to implement an automated process for

reconstructing midpoint-rooted, bootstrapped phylogenies from MSA partitions

representing each rank within the constrained MSA tree. Gene coverage data was

used to condition the way in which taxa were selected into MSA partitions [68].

Phylogenies were reconstructed using the maximum likelihood approach

implemented in RAxML (version 8.2.12) [68]. Protein substitution models were

selected via hierarchical maximum likelihood ratio test using the

PROTAUTOGAMMA option in RAxML. JalView [69] (version 2.11.1.4) and Se-Al

(version 2.0a11) were used to inspect MSAs.

FIGURES

FIGURE LEGENDS

Figure 1. The multiple sequence alignment hierarchy in Flavivirid-GLUE.

The GLUE software framework defines a 'constrained alignment tree' data structure

comprising a set of alignments that are hierarchically linked to reflect taxonomic

relationships [27]. To enable sequence comparisons across the entire *Flaviviridae*,

we implemented a constrained alignment tree data structure in *Flavivirid*-GLUE, as

shown in the cladogram above. Numbers shown adjacent to nodes correspond to

rows in **Table 1. Abbreviations**: cISF = classical insect-specific flaviviruses; NKV =

no known vector; Jingmen = Jingmen tick virus =; PL = 'pestivirus-like'; MBFV =

mosquito borne flavivirus; MSA = multiple sequence alignment.

Figure 2. Genome structures of endogenous flaviviral elements.

Schematic diagrams showing the genomic regions represented by flavivirid-derived

endogenous viral element (EVE) sequences. **(a)** Pegi-hepacivirus-like elements are

shown relative to hepatitis C virus (HCV); **(b)** elements derived from the 'Pestivirus-

like 2' (PL2) lineage of viruses shown relative to Shuangao lacewing virus 2 (SLwV-

2); **(c)** Elements derived from the *Flavivirus* genus, the 'tamanaviruses', and the X1

lineage shown relative to Dengue virus type 1 (DENV-1); **(d)** 'jingmenvirus'-derived

EFVs shown relative to Jingmen tick virus (JMTV). Homologous regions represented

by EFV sequences are shown as grey horizontal bars. Bars to the right show

taxonomic groups. EVE identifiers (IDs) are shown to the left. EVE IDs were

constructed as indicated in the key, following a convention established for

endogenous retroviruses [70]. IDs have three components - the first is the classifier

'EFV' 'endogenous flaviviral element' (can be dropped when implied by context). The

second comprises two subcomponents separated by a period; (i) the name of the

taxonomic group of viruses from which the EFV is thought to derive; (ii) a numeric ID

that uniquely identifies the integration locus. The third component describes the

known taxonomic distribution of orthologous copies of the element among host

species. For EVEs derived from the 'jingmenvirus' lineage, which contains viruses

with multipartite genomes, we used classifiers that specify the gene from which the

EVE is derived, in line with conventions established for EVEs derived from

segmented viruses [71]– endogenous 'jingmenvirus' helicase (EJH) and endogenous

'jingmenvirus' polymerase (EJP). **Abbreviations**: Kb = kilobase. X1= unclassified

flavivirus-like lineage X1. X2 = unclassified flavivirus-like lineage X2. C = Capsid, prM

= Pre-Membrane, E = Envelope, NS = Non-Structural protein, P = Protease, H =

Helicase. HP = 'pegi-hepaci'. ***** Indicates consensus sequence.

Figure 3. Evolutionary relationships between modern and ancient flavivirids.

Bootstrapped maximum likelihood phylogenies (1000 replicates), reconstructed for

viruses and EFVs across a range of taxonomic ranks, as follows: **(a)** Two major

lineages within family *Flaviviridae* ('flavi-pesti' and 'pegi-hepaci') showing placement

of Hepacivirus-derived EVE (122 amino acid (aa) residues in multiple sequence

alignment (MSA) spanning conserved regions in NS3, substitution model=RTREV);

(b) 'Flavi-pesti' lineage (MSA spanning 99 aa residues in NS5, substitution

model=BLOSUM62); **(c)** 'Pesti-like' lineage (104 aa residues in NS5, substitution

model=RTREV); **(d)** 'Flavi-like' lineage (MSA spanning 179 aa residues in NS5,

substitution model=LG likelihood); **(e)** 'Tamanavirus' (MSA spanning 292 aa residues

in NS5, substitution model=BLOSUM62); **(f)** 'Jingmenvirus' and related lineages

(MSA spanning 727 residues in NS5, substitution model=LG likelihood). EFV names

are shown in bold. Only EFVs that had sufficient coverage (>50% of total MSA

length) were included in the analysis. Triangular terminal branches indicate collapsed

clades containing multiple taxa. Asterisks indicate bootstrap support ≥70% (1000

replicates). The scale bar indicates evolutionary distance in substitutions per site.

Brackets to the right indicate genera and sub-lineages. All trees are midpoint rooted

for display purposes. Host and known/suspected vector associations are indicated by

animal silhouettes as shown in the key. **Abbreviations**: cISF = classical insect-

specific flaviviruses. NKV = no known vector. Jingmen = Jingmen tick virus =. PL =

'pesti-like'; MBFV = Mosquito borne flavivirus.

Figure 4. Putative codivergence of flavivirid groups and host phyla.

'Tanglegrams' illustrating matching topologies in animal (left) and flavivirid (right)

phylogenies. Putative tracking of host lineages by viral lineages is indicated on the

host phylogeny. Clades within which the branch lengths of virus phylogenies are

correlated with divergence times in host animal lineages: the invertebrate and

vertebrate splits in the (**a**) 'pegi-hepaci' lineage (R^2=0.5); (**b**) 'pesti-like' lineage

(R^2=0.7); (**c**) the cnidarian-arthropod, chelicerate-hexapoda, and crustacean-insect

splits in the *Flavivirus* genus (R^2=0.3). Note that, due to limited data, all comparisons

rely on strong assumptions regarding virus phylogenies, indicated in the figure by

numbers, as follows: (1) and (2) midpoint rooting was used, and the deepest

divergence in the virus tree was approximately calibrated using the divergence date

of porifera (the most basal animal lineage) in line our hypothesis that major flavivirid

lineages originated in early metazoans; (3) the splits between nematode and

arthropod viruses in "pesti-like 1" (PL1) and "pesti-like 2" (PL2) clades – indicated by

the yellow square - were poorly resolved in our phylogenies; (4) midpoint rooting was

used, and it is assumed that the arthropod-borne classical flaviviruses originated in

arachnids, as proposed in **Fig 5a**. Abbreviations: Mya=Million years ago.

Figure 5. The timeline of flavivirus evolution

(a) Evolution of arthropod-borne flaviviruses. The phylogeny shown on the left

was constructed from an alignment spanning 804 residues of the precursor

polyprotein (substitution model=LG likelihood). The phylogeny on the right is a time-

calibrated phylogeny of arthropod hosts/vectors of classical flaviviruses, obtained via

TimeTree [72]. The figure shows the hypothesised relationships between the host

and virus trees, with virus phylogeny broadly following host phylogeny at higher

taxonomic ranks but diverging dramatically from this pattern when tick-borne viruses

of vertebrates emerge.

(b) Proposed model for the origin and evolution of the vector-borne

flaviviruses. Three stages are shown: (i) an ancestral group of non-vectored tick

viruses is present; (ii) tick hematophagy provides prolonged exposure to vertebrate

blood, thereby affording tick-specific flaviviruses the opportunity, through chance

events and mutation, to acquire a capacity to replicate in vertebrate cells and

ultimately become tick-borne flaviviruses of vertebrates; (iii) the existence of tick-

borne flaviviruses occasions viraemic vertebrate hosts, exposing mosquitoes and

other hematophagous insects to flaviviruses via blood-feeding, and ultimately

allowing them to acquire a vector role.

(c) A model for long-term evolution of flavivirids. A time-scaled tree summarising

the phylogenetic relationships of flavivirids and calibrated using information obtained

from analysis of endogenous viral elements (EVEs), co-phyletic analysis and the

fossil record of hematophagous arthropods. Symbols on the phylogenies indicate

types of calibration as shown in the key**.** Numbers adjacent to symbols link to rows in

Table 3 'Codivergence type A' = codivergence supported by co-phylogeny.

Codivergence type B = potential codivergence-based calibrations without supporting

evidence. Vertical lines to right of the tree show taxonomic groups.

Table 1. Comprehensive mapping of flavivirid homology via hierarchically linked multiple sequence alignments.

# [a]	Taxonomic scope	Name	Parent [b]	Children [c]	Constraining reference [d]	Genome Coverage [e]	Virus count	EVE count [f]
Family								
1	Flaviviridae-like	Flaviviridae	none	2	YFV	~6%	2	
Major lineage								
2	Hepaci/Pegi-like viruses	HepaPegi	Flaviviridae	2	BVDV1	~60%	2	1
3	Flavi/Pesti-like	FlaviPesti	Flaviviridae	2	YFV	~80%	2	1
4	Flavi-like	FlaviLike	FlaviPesti	2	YFV	~35%	2	
5	Pesti-like	PestiLike	FlaviPesti	3	BVDV1	~60%	3	
Minor lineage								
6	Flavi-Tamana-like	FlaviTamana	FlaviLike	2	YFV	~93%		
Genus								
7	*Flavivirus* genus	Flavivirus	FlaviTamana	11	YFV		8	3
8	*Hepacivirus* genus*	Hepacivirus	HepaPegi	0	HCV	~90%	14	
9	*Pegivirus* genus*	Pegivirus	HepaPegi	0	HPgV2	~60%	9	
10	Insect pesti-like*	PL2	PestiLike	0	SLV2	~73%	6	7
11	SCNV+arthropod*	PL1	PestiLike	0	SCNV5	~72%	8	
12	*Pestivirus* genus*	Pestivirus	PestiLike	0	BVDV1	~88%	11	
13	X2-derived EVEs	X2	FlaviLike	0	NS5			9
14	'Jingmenvirus' seg 1	Jingmen_1	FlaviLike	0	JMTV Seg1	Segment 1	7	2
14	'Jingmenvirus' seg 3	Jingmen_3	FlaviLike	0	JMTV Seg3	Segment 3	4	1
15	'Tamanavirus'	'Tamanavirus'	FlaviTamana	0	TABV	100%	6	5
Subgenus								
16	Crustacean	crustacean	Flavivirus	0		100%Δ-genome	2	
17	cISF	cISF	Flavivirus	0	KRV	100%	14	6
18	NKV1	NKV2	Flavivirus	0	APOIV	100%	6	
19	Tick	tick	Flavivirus	0	POWV	~91%	15	
20	dISF	dISF	Flavivirus	0	LAMV	~95%	9	
21	Mosquito1	mosquito-1	Flavivirus	0	DEN1	~97%	35	
22	NKV2	NKV2	Flavivirus	0	SOKV	100%	3	
23	Mosquito2	mosquito-2	Flavivirus	0	YFV	~93%	14	
	Totals						**159**	**35**

Footnote: [a] Numbers correspond to labelled nodes in **Fig. 1**. [b] The parent multiple sequence alignment (MSA) in the hierarchy. [c] Children of the MSA in the hierarchy. [d] Reference sequence that constrains the genomic co-ordinate coordinate space in the MSA. [e] Percentage of the constraining reference genome spanned by the MSA. Phylogenies constructed for each of these alignment partitions are available in Flavivirid-GLUE [28]. [f] Counts reflect the number of consensus EVE sequences included in the alignment – consensuses are linked to child MSAs that contain the sequences of all individual EVE loci used to create the consensus.

Table 2. Flavivirid-derived endogenous viral elements.

Sequence ID[a]	#Species[b]	#Sequences[c]	IORF[d]		Host[e]
Flavivirus					
EFV-cISF.1-AedAeg*	1	92	564	Yellow fever mosquito	(*Aedes aegypti*)
EFV-cISF.2-AedAlb*	1	35	643	Tiger mosquito	(*Aedes albopictus*)
EFV-cISF.3-AnoMin*	1	1	97		(*Anopheles minimus*)
EFV-cISF.4-AnoSin*	1	1	138		(*Anopheles sinensis*)
EFV-cISF.5-TipOle	1	4	226	Marsh crane fly	(*Tipula oleracea*)
EFV-cISF.6-ConPat	1	1	38	Long-legged fly	(*Condylostylus patibulatus*)
EFV-Flavi.1-CraSow	1	13	1182	Peach blossom jellyfish	(*Craspedacusta sowerbyi*)
EFV-Flavi.2-DapMag*	1	2	573	Water flea	(*Daphnia magna*)
EFV-Flavi.3-LepArc*	1	3	281	Tadpole shrimp	(*Lepidurus arcticus*)
'Tamanavirus'					
EFV-Tamana.1-LedTum	1	1	55	Meltwater stonefly	(*Lednia tumana*)
EFV-Tamana.2-LauKoh	1	1	56	Hawaiian cricket	(*Laupala kohalensis*)
EFV-Tamana.3-AmpSul	1	2	44	Spring stonefly	(*Amphinemura sulcicollis*)
EFV-Tamana.4-StyCho	1	1	873	Tube-eye	(*Stylephorus chordatus*)
Jingmenvirus					
EJP-Jingmen.1-Chironomus	2	2	248		*Chironomus sp.*
EJP-Jingmen.2-*Gerris*	1	1	45	Water strider	(*Gerris buenoi*)
EJH-Jingmen.1-*Gerris*	1	1	51	"	"
X1					
EFV-X1.1-PriCau	1	2	347	Penis worm	(*Priapulus caudatus*)
X2					
EFV-X2.1-AusLim	1	2	214	Mangrove killifish	(*Austrofundulus limnaeus*)
EFV-X2.2-StyCho	1	3	873	Tube-eye	(*Stylephorus chordatus*)
EFV-X2.3-EurAff	1	2	190	Copepod	(*Eurytemora affinis*)
EFV-X2.4-Takifugu	3	5	103	Pufferfish	*Takifugu*
EFV-X2.5-Phycis	2	2	129	Phycid hakes	*Phycis*
EFV-X2.6-MorMor	1	1	36	Common mora	(*Mora moro*)
EFV-X2.7-BroBro	1	1	56	Cusk	(*Brosme brosme*)
EFV-X2.8-MacPee	1	1	51	Murray cod	(*Maccullochella peelii*
EFV-X2.9-BolPec	1	1	155	Blue spotted mudhopper	(*Boleophthalmus pectinirostris*)
Pesti-like					
EFV-PL2.1-CalCec	1	5	78	Red-banded hairstreak	(*Calycopis cecrops*)
EFV-PL2.2-EucHer	1	2	187	Stink bug	(*Eucheros histo*)
EFV-PL2.3-Apoidae	11	239	2469	Bees & sphecoid wasps	Superfamily Apoidae
EFV-PL2.4-AndCur	2	6	961	Cynipid gall wasp	Andricus
EFV-PL2.5-AnoGla	1	1	26	Long-horned beetle	(*Anoplophora glabripennis*)
EFV-PL2.6-XenBra	1	16	2232	Grasshopper	(*Xenocatantops brachycerus*)
EFV-PL2.7-OpeBru	1	2	38	Winter moth	(*Operophtera brumata*)
Pegi-Hepaci					
EFV-HepaPegi.1-ElyChl	1	1	60	Eastern emerald elysia	(*Elysia chlorotica*)

Footnote: [a] Flavivirid-derived EVEs have been assigned standard IDs based on conventions established for endogenous retroviruses, wherein information about virus taxonomy and locus orthology are incorporated into the ID itself [70]. The ID comprises of three elements separated by hyphens. For most EVEs characterised here, the first (i.e., leftmost) element is the classifier 'endogenous flavivirid' (EFV). However, for 'jingmenviruses' the classifier component of the ID also specifies the gene it derives from (EJP=endogenous jingmenvirus polymerase; EJH=endogenous jingmenvirus helicase), following conventions established for multipartite viruses and EVEs derived from mRNA sources [71]. The second ID element comprises two subcomponents separated by a period – the first defines the taxonomic position of the EVE in relation to established *Flaviviridae* taxonomy, the second is a numeric ID that uniquely represents an EVE locus. The third ID component defines the known distribution of orthologous insertions among host species. If it is only known from a single species a shortened version of the Latin binomial species name is used. [b] Number of species in which EVE locus was identified. [c] Number of sequences (i.e., distinct insertions) derived from this EVE that were identified via *in silico* screening. [e] Host species or species groups. Asterisks indicate EFVs loci or lineages that have been reported previously. Abbreviations: L-ORF=Longest open reading frame; PL2=Pesti-like 2; cISF=classical insect-specific flaviviruses.

Table 3. Dates and age estimates used to calibrate flavivirid evolution.

#	Virus lineage	Host lineage(s)	MYA	Low	High
Codivergence A (minimum age)					
1	*Flaviviridae*	Animalia	952	757	1147
6	Jingmen-X2	Insecta-Crustacea	794	678	916
22	Mosquito 2 subgenus	Culex-Aedes	40	22	52
10	Jingmenvirus	Arachnida-Insecta	601	568	642
Codivergence B (minimum age)					
2	Pegi-hepaci lineage	Mollusc-vertebrate	794	678	916
3	Pesti-like viruses	Invertebrates-vertebrates	794	678	916
4	Pegi-Hepaci	Chondrichthyes-Mammalia	465	450	497
5	Pestiviruses	Chondrichthyes-Mammalia	465	450	497
7	Flavivirus	Cnidaria-Arthropoda	824	652	973
8	Arthropod flaviviruses	Crustacea-Arachnida	601	568	642
9	Chelicerata flaviviruses	Pycnogonida-Arachnida	553	476	653
Orthology (minimum age)					
13	PL2	Family Apoidea	102	71	148
24	Jingmenvirus	Genus *Chironomus*	*n/k*	*n/k*	*n/k*
23	X2	Genus *Phycis*	16.9	12.1	37.1
Duplicates /Molecular clock (minimum age)					
20	cISF	Tipula		16	40
21	Crustacean FV	Daphnia 2		26	62
19	Cnidarian FV	Craspedacusta 1		17	40
16	X1	Priapulus (NS5)		12	29
17	X2	Eurytemora (NS5)		20	47
18	'Tamanavirus'	Austrofundulus 1 (NS5)		19	43
14	PL2	Operophtera		17	40
Origin hypothesis (maximum age)					
11	All vectored flaviviruses	Tick hematophagy	300	90	400
12	Mosquito-vectored flaviviruses	Mosquito hematophagy		79	100

Footnote: Node numbers correspond to those shown in **Fig. 5c**. Hematophagy estimates obtained from [43]. *Culex-Aedes* divergence date obtained from [73]. All other divergence data estimates were obtained from TimeTree [72].

Data Availability

Data available via GitHub: https://giffordlabcvr.github.io/Flavivirus-GLUE/

Acknowledgements

RJG was funded by the Medical Research Council of the United Kingdom (MC_UU_12014/12). WMS acknowledges support from the Global Virus Network Fellowship. We thank Anna Gatseva, Joseph Hughes, Alain Kohl, Spyros Lytras, Emilie Pondeville, Charles Rice, David Robertson, Greg Towers, Sam J. Wilson, and anonymous reviewers for critical reading of the manuscript.

References

1. Simmonds, P., et al., *ICTV Virus Taxonomy Profile: Flaviviridae.* J Gen Virol, 2017. **98**(1): p. 2-3.
2. de Oliveira, L.G., et al., *Bovine Viral Diarrhea Virus: Recent Findings about Its Occurrence in Pigs.* Viruses, 2020. **12**(6).
3. Manns, M.P., et al., *Hepatitis C virus infection.* Nature Reviews Disease Primers, 2017. **3**(1): p. 17006.
4. Pierson, T.C. and M.S. Diamond, *The continued threat of emerging flaviviruses.* Nat Microbiol, 2020. **5**(6): p. 796-812.
5. Chambers, T.J., et al., *Flavivirus genome organization, expression, and replication.* Annu Rev Microbiol, 1990. **44**: p. 649-88.
6. Shi, M., et al., *Divergent Viruses Discovered in Arthropods and Vertebrates Revise the Evolutionary History of the Flaviviridae and Related Viruses.* J Virol, 2016. **90**(2): p. 659-69.
7. Shi, M., et al., *The evolutionary history of vertebrate RNA viruses.* Nature, 2018. **556**(7700): p. 197-202.
8. Parry, R. and S. Asgari, *Discovery of novel crustacean and cephalopod flaviviruses: insights into evolution and circulation of flaviviruses between marine invertebrate and vertebrate hosts.* J Virol, 2019.
9. Porter, A.F., et al., *Novel hepaci- and pegi-like viruses in native Australian wildlife and non-human primates.* Virus Evol, 2020. **6**(2): p. veaa064.
10. Paraskevopoulou, S., et al., *Viromics of extant insect orders unveil the evolution of the flavi-like superfamily.* Virus Evol, 2021. **7**(1): p. veab030.
11. Qin, X.C., et al., *A tick-borne segmented RNA virus contains genome segments derived from unsegmented viral ancestors.* Proc Natl Acad Sci U S A, 2014. **111**(18): p. 6744-9.
12. Wang, Z.D., et al., *A New Segmented Virus Associated with Human Febrile Illness in China.* N Engl J Med, 2019. **380**(22): p. 2116-2125.
13. Jia, N., et al., *Emergence of human infection with Jingmen tick virus in China: A retrospective study.* EBioMedicine, 2019. **43**: p. 317-324.
14. Geoghegan, J.L. and E.C. Holmes, *Evolutionary Virology at 40.* Genetics, 2018. **210**(4): p. 1151-1162.
15. Zanotto, P.M., et al., *Population dynamics of flaviviruses revealed by molecular phylogenies.* Proc Natl Acad Sci U S A, 1996. **93**(2): p. 548-53.
16. Cook, S. and E.C. Holmes, *A multigene analysis of the phylogenetic relationships among the flaviviruses (Family: Flaviviridae) and the evolution of vector transmission.* Arch Virol, 2006. **151**(2): p. 309-25.
17. Gould, E.A., et al., *Origins, evolution, and vector/host coadaptations within the genus Flavivirus.* Adv Virus Res, 2003. **59**: p. 277-314.
18. Moureau, G., et al., *New insights into flavivirus evolution, taxonomy and biogeographic history, extended by analysis of canonical and alternative coding sequences.* PLoS One, 2015. **10**(2): p. e0117849.
19. Holmes, E.C. and S. Duchêne, *Can Sequence Phylogenies Safely Infer the Origin of the Global Virome?* mBio, 2019. **10**(2).
20. Grüning, B., et al., *Practical Computational Reproducibility in the Life Sciences.* Cell Syst, 2018. **6**(6): p. 631-635.
21. Pettersson, J.H. and O. Fiz-Palacios, *Dating the origin of the genus Flavivirus in the light of Beringian biogeography.* J Gen Virol, 2014. **95**(Pt 9): p. 1969-82.
22. Holmes, E.C., *The evolution of endogenous viral elements.* Cell Host Microbe, 2011. **10**(4): p. 368-77.
23. Katzourakis, A. and R.J. Gifford, *Endogenous viral elements in animal genomes.* PLoS Genet, 2010. **6**(11): p. e1001191.

24.   Lequime, S. and L. Lambrechts, *Discovery of flavivirus-derived endogenous viral elements in Anopheles mosquito genomes supports the existence of Anopheles-associated insect-specific flaviviruses.* Virus Evol, 2017. **3**(1): p. vew035.

25.   Whitfield, Z.J., et al., *The Diversity, Structure, and Function of Heritable Adaptive Immunity Sequences in the Aedes aegypti Genome.* Curr Biol, 2017. **27**(22): p. 3511-3519.e7.

26.   Blair, C.D., K.E. Olson, and M. Bonizzoni, *The Widespread Occurrence and Potential Biological Roles of Endogenous Viral Elements in Insect Genomes.* Curr Issues Mol Biol, 2020. **34**: p. 13-30.

27.   Singer, J.B., et al., *GLUE: a flexible software system for virus sequence data.* BMC Bioinformatics, 2018. **19**(1): p. 532.

28.   Gifford, R.J. *Flavivirid-GLUE.* 2021; Available from: https://giffordlabcvr.github.io/Flavivirus-GLUE.

29.   Merkel, D., *Docker: lightweight linux containers for consistent development and deployment.* Linux Journal, 2014. **239**(2).

30.   Loeliger, J. and M. McCullough, *Version Control with Git: Powerful tools and techniques for collaborative software development.* 2nd ed. 2012, Sebastopol, CA: O'Reilly Media, Inc.

31.   Ophinni, Y., et al., *piRNA-Guided CRISPR-like Immunity in Eukaryotes.* Trends Immunol, 2019. **40**(11): p. 998-1010.

32.   Aiewsakun, P. and A. Katzourakis, *Marine origin of retroviruses in the early Palaeozoic Era.* Nat Commun, 2017. **8**: p. 13954.

33.   Dolja, V.V. and E.V. Koonin, *Metagenomics reshapes the concepts of RNA virus evolution by revealing extensive horizontal virus transfer.* Virus Res, 2018. **244**: p. 36-52.

34.   Dheilly, N.M., et al., *A World of Viruses Nested within Parasites: Unraveling Viral Diversity within Parasitic Flatworms (Platyhelminthes).* Microbiol Spectr, 2022: p. e0013822.

35.   Blitvich, B.J. and A.E. Firth, *Insect-specific flaviviruses: a systematic review of their discovery, host range, mode of transmission, superinfection exclusion potential and genomic organization.* Viruses, 2015. **7**(4): p. 1927-59.

36.   Blitvich, B.J. and A.E. Firth, *A Review of Flaviviruses that Have No Known Arthropod Vector.* Viruses, 2017. **9**(6).

37.   Conway, M.J., *Identification of a Flavivirus Sequence in a Marine Arthropod.* PLoS One, 2015. **10**(12): p. e0146037.

38.   Staples, J.E. and T.P. Monath, *Yellow Fever: 100 Years of Discovery.* JAMA, 2008. **300**(8): p. 960-962.

39.   Rosenberg, R., et al., *Search strategy has influenced the discovery rate of human viruses.* Proc Natl Acad Sci U S A, 2013. **110**(34): p. 13961-4.

40.   Gibb, R., et al., *Mammal virus diversity estimates are unstable due to accelerating discovery effort.* Biol Lett, 2022. **18**(1): p. 20210427.

41.   Zhou, W., et al., *Exosomes serve as novel modes of tick-borne flavivirus transmission from arthropod to human cells and facilitates dissemination of viral RNA and proteins to the vertebrate neuronal cells.* PLoS Pathog, 2018. **14**(1): p. e1006764.

42.   Monahan-Earley, R., A.M. Dvorak, and W.C. Aird, *Evolutionary origins of the blood vascular system and endothelium.* J Thromb Haemost, 2013. **11 Suppl 1**(Suppl 1): p. 46-66.

43.   Mans, B.J., *Evolution of vertebrate hemostatic and inflammatory control mechanisms in blood-feeding arthropods.* J Innate Immun, 2011. **3**(1): p. 41-51.

44.   Hermance, M.E. and S. Thangamani, *Tick☐Virus☐Host Interactions at the Cutaneous Interface: The Nidus of Flavivirus Transmission.* Viruses, 2018. **10**(7).

45.	Gritsun, T.S. and E.A. Gould, *Direct repeats in the flavivirus 3' untranslated region; a strategy for survival in the environment?* Virology, 2007. **358**(2): p. 258-65.

46.	Ochsenreiter, R., I.L. Hofacker, and M.T. Wolfinger, *Functional RNA Structures in the 3'UTR of Tick-Borne, Insect-Specific and No-Known-Vector Flaviviruses.* Viruses, 2019. **11**(3).

47.	Gritsun, D.J., et al., *Molecular archaeology of Flaviviridae untranslated regions: duplicated RNA structures in the replication enhancer of flaviviruses and pestiviruses emerged via convergent evolution.* PLoS One, 2014. **9**(3): p. e92056.

48.	Li, Y., et al., *Discovery of Flaviviridae-derived endogenous viral elements in shrew genomes provide novel insights into Pestivirus ancient history.* bioRxiv, 2022: p. 2022.02.11.480044.

49.	Price, J.L., *Isolation of Rio Bravo and a hitherto undescribed agent, Tamana bat virus, from insectivorous bats in Trinidad, with serological evidence of infection in bats and man.* Am J Trop Med Hyg, 1978. **27**(1 Pt 1): p. 153-61.

50.	Kuno, G., et al., *Phylogeny of the genus Flavivirus.* J Virol, 1998. **72**(1): p. 73-83.

51.	de Lamballerie, X., et al., *Genome sequence analysis of Tamana bat virus and its relationship with the genus Flavivirus.* J Gen Virol, 2002. **83**(Pt 10): p. 2443-54.

52.	Skoge, R.H., et al., *New virus of the family Flaviviridae detected in lumpfish (Cyclopterus lumpus).* Arch Virol, 2018. **163**(3): p. 679-685.

53.	Geoghegan, J.L., et al., *Hidden diversity and evolution of viruses in market fish.* Virus Evol, 2018. **4**(2): p. vey031.

54.	Dong, X., et al., *A Novel Virus of Flaviviridae Associated with Sexual Precocity in Macrobrachium rosenbergii.* mSystems, 2021. **6**(3): p. e0000321.

55.	Suzuki, Y., et al., *Non-retroviral Endogenous Viral Element Limits Cognate Virus Replication in Aedes aegypti Ovaries.* Curr Biol, 2020. **30**(18): p. 3495-3506.e6.

56.	Goic, B., et al., *Virus-derived DNA drives mosquito vector tolerance to arboviral infection.* Nat Commun, 2016. **7**: p. 12410.

57.	Belshaw, R., et al., *High copy number in human endogenous retrovirus families is associated with copying mechanisms in addition to reinfection.* Mol Biol Evol, 2005. **22**(4): p. 814-7.

58.	Fischer, M.G. and C.A. Suttle, *A virophage at the origin of large DNA transposons.* Science, 2011. **332**(6026): p. 231-4.

59.	Inoue, Y., et al., *Fusion of piggyBac-like transposons and herpesviruses occurs frequently in teleosts.* Zoological Lett, 2018. **4**: p. 6.

60.	Lytras, S., G. Arriagada, and R.J. Gifford, *Ancient evolution of hepadnaviral paleoviruses and their impact on host genomes.* Virus Evol, 2021. **7**(1): p. veab012.

61.	Hill, V., et al., *Progress and challenges in virus genomic epidemiology.* Trends Parasitol, 2021. **37**(12): p. 1038-1049.

62.	Sharvit, Y., *Data-Oriented Programming - Unlearning objects.* 2021: Manning.

63.	Singer, J., et al., *CoV-GLUE: A Web Application for Tracking SARS-CoV-2 Genomic Variation.* Preprints, 2020.

64.	Campbell, K., et al., *Making genomic surveillance deliver: A lineage classification and nomenclature system to inform rabies elimination.* PLoS Pathog, 2022. **18**(5): p. e1010023.

65.	Altschul, S.F., et al., *Gapped BLAST and PSI-BLAST: a new generation of protein database search programs.* Nuc. Acids Res., 1997. **25**: p. 3389-3402.

66.	Zhu, H., et al., *Database-integrated genome screening (DIGS): exploring genomes heuristically using sequence similarity search tools and a relational database.* bioRxiv, 2018.

67. Kitts, P.A., et al., *Assembly: a resource for assembled genomes at NCBI.* Nucleic Acids Res, 2016. **44**(D1): p. D73-80.
68. Stamatakis, A., *RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies.* Bioinformatics, 2014. **30**(9): p. 1312-3.
69. Waterhouse, A.M., et al., *Jalview Version 2--a multiple sequence alignment editor and analysis workbench.* Bioinformatics, 2009. **25**(9): p. 1189-91.
70. Gifford, R.J., et al., *Nomenclature for endogenous retrovirus (ERV) loci.* Retrovirology, 2018. **15**(1): p. 59.
71. Kawasaki, J., et al., *100-My history of bornavirus infections hidden in vertebrate genomes.* Proc Natl Acad Sci U S A, 2021. **118**(20).
72. Kumar, S., et al., *TimeTree: A Resource for Timelines, Timetrees, and Divergence Times.* Mol Biol Evol, 2017. **34**(7): p. 1812-1819.
73. Sieglaff, D.H., et al., *Comparative genomics allows the discovery of cis-regulatory elements in mosquitoes.* Proc Natl Acad Sci U S A, 2009. **106**(9): p. 3053-8.