Cryo-electron tomography of ASC signalling sites in pyroptotic cells

Yangci Liu[1,2], Haoming Zhai[1,2], Helen Alemayehu[1,2], Lee J. Hopkins[3,4], Alicia C. Borgeaud[5],

Christina Heroven[5], Jonathan D. Howe[5], Jérôme Boulanger[5],

Clare E. Bryant[3,4,*] & Yorgo Modis[1,2,*]

[1]Molecular Immunity Unit, Department of Medicine, University of Cambridge, MRC Laboratory of

Molecular Biology, Francis Crick Avenue, Cambridge, CB2 0QH, United Kingdom

[2]Cambridge Institute of Therapeutic Immunology & Infectious Disease (CITIID), University of

Cambridge School of Clinical Medicine, Cambridge CB2 0AW, United Kingdom

[3]Department of Medicine, University of Cambridge, Box 157, Level 5, Addenbrooke's Hospital,

Cambridge, CB2 0QQ, United Kingdom

[4]Department of Veterinary Medicine, University of Cambridge, Madingley Road, Cambridge

CB3 0ES, United Kingdom

[5]MRC Laboratory of Molecular Biology, Francis Crick Avenue, Cambridge, CB2 0QH, United

Kingdom

Current addresses: Institute of Biochemistry and Molecular Medicine, University of Bern,

Bühlstrasse 28, 3012 Bern, Switzerland (A.C.B.); Wren Therapeutics, Clarendon House,

Clarendon Road, Cambridge CB2 8FH, UK (L.J.H.).

*Corresponding authors: C.E.B., ceb27@cam.ac.uk; Y.M., ymodis@mrc-lmb.cam.ac.uk

Abstract

Inflammasomes induce cell death in response to infection, chemical entities and cell damage. Human genetics and animal models have identified the NLRP3 inflammasome as a gatekeeper of caspase-1-dependent pyroptosis. NLRP3 activation induces polymerization of ASC into a single, micron-scale perinuclear punctum, where caspase-1 is activated. These puncta have yet to be imaged at sufficient resolution to resolve their ultrastructure. Here, we apply correlative cryo-light microscopy and cryo-electron tomography to visualize ASC/caspase-1 puncta and mitochondrial membrane remodelling within NLRP3-activated cells. The puncta are composed primarily of branched ASC filaments. The N-terminal pyrin domain forms an 8-10 nm-diameter tubular core, which is decorated by the flexibly linked C-terminal caspase recruitment domain. The variable filament density allows ribosomes and *trans*-Golgi-like vesicles to permeate the network. We propose this provides structural integrity while allowing macromolecules and vesicles to diffuse in or bind with the necessary density, timing and localization.

Keywords

NLR family pyrin domain containing 3 (NLRP3); apoptosis-associated speck-like protein containing a CARD (ASC) speck; caspase-1; pyroptosis; correlative light and electron microscopy (CLEM); *In situ* cryo-electron tomography (cryo-ET); focused-ion beam (FIB) milling; inflammasome; caspase activation and recruitment domain (CARD); pyrin domain (PYD)

Running title: Cryo-ET of ASC specks and mitochondria in cell lamellae

Introduction

Inflammasomes are vital innate immune oligomeric signalling complexes that induce proteolytic

activation of caspase-1, leading to secretion of proinflammatory cytokines interleukin (IL)-1β and

IL-18, and cell death by gasdermin D-driven pyroptosis. In the first step of inflammasome

assembly, Nod-like receptors (NLR) sense cytosolic chemical signatures associated with

microbes and cell dysfunction or damage[1]. Ligand binding induces NLR oligomerization and

recruitment of downstream proteins linked to the caspase-1 signalling cascade.

NLRP3 is one of the most studied inflammasome NLRs due to the large number of

NLRP3 genetic variants associated with human diseases and the availability of animal disease

models. At least 250 human NLRP3 protein variants have been identified, over 30 of which have

been associated with disease through gain-of-function[2,3], including autoinflammatory diseases,

degenerative disorders, dermatitis and metabolic disorders[4,5]. NLRP3 inflammasome activation

has been linked to a broad spectrum of cellular processes or cytosolic chemical signatures

including potassium efflux, reactive oxygen species, calcium influx, chloride efflux,

phosphatidylinositol-4-phosphate on Golgi membranes, and mitochondrial stress or damage[6,7].

Cryo-electron microscopy (cryo-EM) single-particle image reconstructions show that ligand-

bound inflammasome NLRs assemble into disk- or split washer-like oligomers[8-11]. In addition to

ligand recognition, activation of the NLRP3 and pyrin inflammasomes requires active transport

along microtubules by dynein to the microtubule organizing centre (MTOC)[12,13]. Upon activation,

NLRP3 recruits the adaptor protein ASC (Apoptosis-associated speck-like protein containing a

caspase recruitment domain (CARD)). NLRP3 and ASC interact through their pyrin domains

(PYDs)[14]. ASC then assembles into a single perinuclear punctum with a micron-scale diameter,

also known as a speck, which functions as a platform for the recruitment and activation of

caspase-1[15,16]. The PYD and CARD of ASC are both required for punctum formation and

signalling[15,17,18]. Structural studies on purified ASC PYD and ASC CARD show that both

domains have a death fold and independently form helical filaments[19-24]. In the cell, fluorescence

microscopy imaging suggests that PYD-PYD interactions drive ASC filament formation whereas

CARD-CARD interactions promote cross-linking and compaction of PYD filaments into

puncta[17,18,23-26]. In addition, ASC recruits procaspase-1 via a CARD-CARD interaction between

the two proteins[15,23]. Super-resolution light microscopy of endogenous ASC specks revealed

punctate structures containing ASC, caspase 1 and NLRP3 within the same macro-assembly[27].

These structures have yet to be imaged inside a cell at sufficient resolution to resolve the ASC

or caspase-1 ultrastructure within puncta[26]. Hence, it remains unclear how the PYD and CARD

contribute to the formation of an ASC punctum inside cells.

Understanding how ASC filament formation contributes to the assembly and activation of

inflammasome puncta requires structural information in the cellular context. Recent advances in

the preparation of frozen vitrified cellular lamellae by focused ion-beam (FIB) milling[28,29] have

allowed the ultrastructure of multimeric protein complexes to be determined in a near-native

context by cryo-electron tomography (cryo-ET). Here, we use *in situ* cryo-ET to obtain three-

dimensional image reconstructions of multimeric signalling complexes in cells with active

NLRP3 inflammasomes. We identify ASC/caspase-1 signalosome puncta by correlative light

and electron microscopy (CLEM) and reveal their ultrastructure and pyroptotic cellular

landscape in immortalised bone marrow derived macrophages (iBMDMs) by cryo-ET image

reconstruction. We find that inflammasome puncta are composed of short, hollow ASC/caspase-

1 filaments, extensively branched, with variable packing density allowing ribosomes and certain

other cellular components to permeate the puncta. We propose that this ultrastructure of the

ASC filament network functions as a signalling platform by providing structural integrity while

allowing downstream signalling molecules to diffuse freely and bind at high density within the

network.

Results

In situ cryo-CLEM of ASC/caspase-1 puncta

High-resolution structures of inflammasome components have been obtained from purified proteins. In a previous study in zebrafish larvae, electron tomography of ASC puncta in fixed larval sections stained with uranyl acetate showed that the puncta contained a dense filament network[26], but the imaging resolution was limited by chemical fixation and staining of the sample. To examine the architecture and cellular interactions of the mammalian NLRP3 signalosome in its physiological environment, we imaged ASC puncta in iBMDMs by correlative fluorescence light microscopy and *in situ* cryo-ET (cryo-CLEM). Cells expressing ASC fused with a C-terminal fluorescent protein have been used extensively to visualize ASC puncta formation[30]. In some experimental systems, ASC overexpression can lead to puncta formation in the absence of activating inflammasome stimuli[30]. Fluorescence microscopy of live iBMDM cells overexpressing ASC-mCerulean used in this study showed that both priming with lipopolysaccharide (LPS) and stimulation with nigericin were still required to induce ASC puncta formation (**Fig. S1a, Movie S1**). Gasdermin D cleavage observed by Western blotting and IL-1β cytokine release measured in ELISAs confirmed that the ASC-mCerulean puncta activated pyroptotic signalling (**Fig. S1b**). Limiting the stimulation time with nigericin to 30 min allowed early-stage inflammasome signalling complexes to be captured, as indicated by colocalization of NLRP3, proteolytically activated caspase-1 and IL-1β with ASC in the puncta, in both wild-type (WT) iBMDMs and ASC-mCerulean iBMDMs (**Fig. S1c-e**). NLRP3 and IL-1β localization was determined by immunofluorescence. Caspase-1 localization was inferred using the carboxyfluorescein-labelled caspase-1 substrate FAM-FLICA. FAM-FLICA or its non-fluorescent analogue Z-VAD-FMK also facilitated imaging by delaying pyroptotic cell death (**Movies S1, S2**).

For high-resolution imaging, cells grown on electron microscopy grids were vitrified by

plunge-freezing in liquid ethane. We used both iBMDMs overexpressing ASC-mCerulean and

WT iBMDMs labelled with FAM-FLICA after NLRP3 inflammasome activation[31]. ASC/caspase-1

signalling complexes were located by cryo-fluorescence light microscopy (cryo-FM; **Fig. S2a**).

Areas of the frozen cells containing an ASC or caspase punctum were FIB-milled down to

lamellae 150-300 nm-thick and 15-20 µm wide[28,29], with micro-expansion joints to reduce

bending[32] (**Fig. S2b,c**). The lamellae were imaged by cryo-CLEM, using cryo-FM to locate any

remaining fluorescent signal from ASC-mCerulean or FAM-FLICA in the lamellae after milling

(**Fig. 1a,b; Fig. S2d,e**). The micron-scale dimensions of the fluorescent puncta allowed cryo-ET

tilt-series acquisition at regions of interest with the necessary targeting precision after

correlation of the cryo-FM images with low-magnification cryo-EM maps (**Fig. S2f,g**)[33,34].

In situ **Cryo-ET reveals a branched-filament ultrastructure in ASC/caspase-1 puncta**

Three-dimensional cryo-ET image reconstructions show a dense network of branched filaments

in areas identified by cryo-CLEM as containing ASC-mCerulean or caspase-1 labelled with

FAM-FLICA (**Figs. 1c-e, 2a, S2g**). Filaments in cells expressing ASC-mCerulean had additional

density at the filament periphery, and a less distinct outline compared to filaments in cells

expressing WT ASC (**Fig. 1d,e**). This suggests the additional density is primarily attributable to

mCerulean, which at 27 kDa is larger than ASC (22 kDa), although other components such as

caspase-1 (which was shown by light microscopy to localize to ASC specks), may also

contribute to this density. A green fluorescent protein tag was similarly found to decorate cryo-

ET reconstructions of amyloid-like poly-Gly-Ala aggregates with additional densities[35]. Since

mCerulean was fused to the C-terminus of ASC, the presence of mCerulean at the filament

periphery supports a model in which the filaments are formed at their core by the N-terminal

PYD of ASC, with the CARD located in between the PYD and mCerulean. Indeed, the cryo-EM

structure of purified recombinant ASC PYD filaments and the NMR structure of full-length ASC,

taken together, suggest that the PYD forms the filament core (9-nm in diameter), with the

flexibly-linked CARD increasing the filament diameter to 16-18 nm[23,24]. In our cryo-ET

reconstructions, filaments containing WT ASC had a diameter of 11-15 nm, which increased to

22-30 nm for filaments containing ASC-mCerulean (**Figs. 1d,e**). Notably, a continuous core of

density 8-10 nm in diameter (**Fig. 2a-c**) was visible within the ASC-mCerulean filaments (for

which the cryo-ET images were of higher quality). Although fluorescence from ASC-mCerulean

and FAM-FLICA-labelled capsase-1 colocalized in iBMDMs (**Fig. S1e**), the resolution of cryo-

CLEM was insufficient to determine whether any of the cryo-ET filament densities corresponded

to caspase-1 or other components. We note, however, that caspase-1 (30 kDa) and

procaspase-1 (45 kDa) are both larger than ASC and would therefore be expected to increase

filament diameter significantly, if present with the same stoichiometry as ASC.

Filament length measurements showed that WT filaments were on average slightly

shorter (191 ± 74 nm, n = 54) than ASC-mCerulean filaments (245 ± 116 nm, n = 402), with

similarly broad length distributions (**Fig. 1f,g**). This difference could potentially be due to the

higher levels of ASC protein expression in iBMDMs expressing ASC-mCerulean, which also

express endogenous WT ASC. Analysis of the branching angles showed that the angle

distributions of untagged and tagged filaments were slightly different, with larger branching

angles more common in mCerulean-tagged filaments, potentially due to steric constraints from

the tag (**Fig. 1h**). Our filament branching angle measurements are likely underestimated as any

filaments with their axis closely aligned with the Z-axis – which would result in a large (70-90˚)

branching angle – would not have been detected due to the missing-wedge effect (from sample-

tilt limitations). Taken together with previously published structural data from purified proteins,

our cryo-ET reconstructions suggest ASC filaments form the backbone of inflammasome-

induced puncta, but that the branching of these filaments results in a structural scaffold that can

recruit and concentrate downstream effector proteins.

Figure 1. Cryo-ET image reconstructions of ASC-caspase 1 puncta in iBMDMs.

(a) Cryo-FM image of a lamella from an iBMDM expressing ASC-mCerulean. ASC-mCerulean

fluorescence is shown (green), overlaid on a bright-field image. Scale bar, 5 µm.

(b) Cryo-FM image shown in (**a**) correlated and overlaid on a transmission cryo-EM map of the

same lamella using the EC-CLEM ICY[36] plugin. Scale bar, 8 µm.

(c) Reconstructed cryo-ET volume (1.36-nm thick virtual tomographic slice), of the ASC-

mCerulean punctum shown in (**a,b**). Scale bar, 250 nm.

(d-e) Galleries of filaments within ASC-mCerulean puncta, (**d**), or FAM-FLICA-labelled caspase-

1 puncta, (**e**), in 4-nm thick virtual tomographic slices of iBMDMs. White arrows indicate

filament axes. Pairs of yellow triangles show filament thickness. Dashed white boxes

highlight ring-shaped densities. Scale bars, 50 nm.

(f-g) Filament length distribution in iBMDMs expressing ASC-mCerulean, (**f**), and in WT

iBMDMs labelled with FAM-FLICA, (**g**).

(h) Distribution of branching angles in ASC-mCerulean puncta (from 2 tomograms), and a

WT ASC punctum (1 tomogram).

For a more complete visualisation of the ASC speck ultrastructure in its cellular

environment, we generated a three-dimensional model of the filament network and other cellular

components including mitochondria, ER and ribosomes with IMOD[37]. This model showed that

puncta consist of several core regions of densely packed filaments separated by sparser

regions, with branching filaments connecting adjacent densely packed regions (**Fig. 2a, Movies**

S3,S4). Ribosomes were abundant within the filament network. Vesicles were also present, but

organelles including mitochondria were excluded.

We next investigated the equilibrium dynamics of ASC-mCerulean filaments by

fluorescence recovery after photobleaching (FRAP) in iBMDMs. We bleached circular regions

containing ASC-mCerulean puncta and imaged the ASC-mCerulean fluorescence intensity

before and after bleaching (**Fig. 2d**). Our data showed that ASC-mCerulean filament assemblies

have a small mobile fraction (6.9%; **Fig. 2e**). The dissociation rate (k_{off}) of ASC-mCerulean from

the imaged area was measured from the rate of fluorescence recovery as 0.023 ± 0.001 s^{-1}

(**Fig. 2d,e; Dataset S1**). Most of the observed fluorescence recovery was at the periphery of

puncta, suggesting that the ASC-mCerulean filaments inside the puncta are largely immobile

and are not in equilibrium with ASC monomers or oligomers in the cytosol (**Fig. 2d**).

Figure 2. Visualization of the filament network and assembly dynamics of ASC puncta

(a) 3-D segmented model of the ASC-mCerulean punctum shown in **Fig. 1c**, generated with

IMOD[37]. Scale bar, 250 nm.

(b) Representative instances of different filament types observed in 4-nm thick virtual

tomographic slices of iBMDMs. Scale bar, 50 nm.

(c) Average normalized density line profile of filament cross-sections, calculated from

projections of 16-nm thick virtual tomographic slices of iBMDMs. Error bars show standard

deviations (ASC-mCerulean, n = 25; WT, n = 13).

(d) Fluorescence recovery after photobleaching (FRAP) of ASC-mCerulean in iBMDMs. Scale

bar, 10 µm.

(e) FRAP curve used to calculate the ASC-mCerulean dissociation rate (k_{off}) from the bleached

area. Shaded area represents ± s.e.m. (n = 8). See **Dataset S1** for source data.

***Trans*-Golgi-like vesicles are present within the ASC filament network**

How NLRP3 is activated by a broad range of different stimuli – including ion fluxes, reactive

oxygen species and mitochondrial damage – remains an open question. A common feature of

NLRP3 activation by these different stimuli is that the *trans*-Golgi network (TGN) is dispersed,

and NLRP3 locally accumulates at foci within the dispersed TGN[7,12]. These NLRP3 foci have

been proposed to serve as nucleation sites for ASC and caspase-1 recruitment and activation[7].

Consistent with previous studies, we observed dispersion of the TGN in WT iBMDM and THP-1

cells, and in iBMDMs expressing ASC-mCerulean (**Figs. 3a,b; S3a**). Moreover, NLRP3 partially

colocalized with the TGN marker TGN38 in WT iBMDMs primed with LPS, with or without

nigericin stimulation (**Fig. 3a**). In stimulated iBMDMs, ASC-mCerulean puncta were located

adjacent to anti-TGN38 fluorescence, but there was little direct overlap between the two

markers (**Fig. 3b**). In our tomographic reconstructions, vesicles with diameters of 50 to 300 nm

were observed within the ASC filament network (**Figs. 3c,d; S2g; S3b; Movies S3, S4**). We

observed similar vesicles within the filament network in chemically fixed ASC-mCerulean puncta

imaged by room temperature electron tomography (**Fig. S3c-e**). Notably, these vesicles had

low-density lumens, a feature characteristic of TGN vesicles[38]. Ribosomes were also found

embedded throughout the ASC filament network. Ring-shaped densities were occasionally

observed within tomograms of ASC puncta, but the resolution of the reconstructions was

insufficient to infer the composition of these features (**Fig. 1d**). Together, our cryo-ET and

fluorescence microscopy data supports the model that the dispersed TGN may serve as a

platform for ASC recruitment to multiple NLRP3 foci, which nucleate ASC polymerization, with

subsequent branching of ASC filaments driving growth into a micron-scale punctum.

Figure 3. TGN dispersion and TGN-like vesicles in ASC puncta after NLRP3 stimulation

(a) Immunofluorescence microscopy of WT iBMDMs primed with LPS or stimulated with

nigericin. NLRP3 partially colocalized with TGN38. The TGN was dispersed following

stimulation. Scale bar, 10 μm.

(b) Immunofluorescence microscopy of iBMDMs expressing ASC-mCerulean before and after

stimulation with nigericin, or nigericin and caspase inhibitor Z-VAD-FMK. ASC-mCerulean

puncta were adjacent to anti-TGN38 fluorescence, but there was little overlap between the

two markers. Scale bar, 10 µm.

(c) 1.36-nm thick virtual tomographic slices of vesicles within the ASC-mCerulean filament

network in an iBMDM. Scale bar, 200 nm. Lower panel, 3-D segmentation model of the

same volume.

(d) Gallery of vesicles from tomographic reconstructions of cryo-FIB-milled ASC-mCerulean

iBMDMs. Scale bar, 100 nm. The histogram shows the vesicle diameter distribution.

Since NLRP3 inflammasome activation was reported to require transport of NLRP3 to

the MTOC[12,13], we examined the localization of ASC/caspase-1 puncta relative to the MTOC.

Fluorescent markers for ASC and caspase-1 did not colocalize with the MTOC components γ-

tubulin, pericentrin or ninein in WT or ASC-mCerulean iBMDMs stimulated with LPS and

nigericin (**Fig. S4a,b**). In stimulated THP-1 cells, ASC/caspase-1 puncta were on average closer

to the MTOC, with one quarter of puncta within 2 µm of the MTOC, but there was little direct

overlap between the fluorescent markers (**Fig. S4c**).

Aberrant mitochondrial morphology during pyroptosis visualized by cryo-ET

Mitochondria function as a nexus for multiple pathogen-sensing and damage-sensing signalling

pathways. Innate immune sensors of viral nucleic acids[39] and mitochondrial damage converge

on the outer mitochondrial membrane, and induce apoptosis or pyroptosis, depending on the

specific danger signal that is sensed[40,41]. We observed fragmentation of the filamentous network

of mitochondria in ASC-mCerulean iBMDMs following stimulation with LPS and nigericin, similar

to AIM-2 dependent mitochondrial remodelling[41], even prior to ASC punctum formation (**Fig.**

4a). Mitochondrial remodelling coincided with dissipation of the membrane potential (**Fig. 4b,**

Fig. S5a). To visualize and analyse the morphology of mitochondria soon after inflammasome

activation we performed cryo-ET reconstructions of mitochondria following LPS priming and

nigericin stimulation to induce ASC speck formation. In LPS primed iBMDMs expressing ASC-

mCerulean, a 30-min stimulation with nigericin resulted in mitochondria with a more rounded

overall shape and smaller size than in unstimulated cells (**Fig. 4c**). Cryo-ET reconstructions and

segmentation models of WT iBMDMs 1-h post-stimulation showed that the mitochondria in

these cells had an abnormal ultrastructure, with the inner membranes forming concentric tubular

structures instead of the regularly spaced lamellar cristae observed in unstimulated cells (**Fig.**

4c,d). To quantify the changes in mitochondrial morphology associated with ASC-dependent

signalling, four parameters were extracted and measured using Fiji[42]: cristae spacing, inner-to-

outer membrane spacing, cristae lumen width and cristae apex angles (**Fig. 5a,b**). We found the

cristae spacing to be two- to fourfold smaller in pyroptotic mitochondria than in healthy

mitochondria. The other three parameters were not significantly different.

a ASC-mCerulean iBMDMs +LPS +Nigericin

t = 30 s   t = 6 min 30 s

Mitoview

b Mitochondrial membrane potential (ΔΨm)

LPS +Nigericin

LPS +ATP

30%

48%

WT iBMDMs (thousands)

Tetramethylrhodamine (TMRM) fluorescence

c

WT iBMDM control

ASC iBMDM +LPS + nigericin (30 min)

WT iBMDM +LPS+ nigericin (1 h)

d

Figure 4. Remodelling of mitochondrial cristae after activation of NLRP3 inflammasome.

(a) Snapshots of live ASC-mCerulean iBMDMs labelled with a MitoView mitochondrial dye. Overlaid fluorescence and bright-field images show soon fragmentation of the mitochondrial network prior to ASC-speck formation. Scale bar 10 μm.

(b) Flow cytometry histogram of iBMDMs stained with tetramethylrhodamine (TMRM), a dye sensitive to mitochondrial membrane potential, ΔΨm. Cells were stimulated with LPS and nigericin (left), or LPS and ATP (right) for 30 min.

280   **(c)** Reconstructed cryo-ET tomographic slices of an unstimulated iBMDM (left); ASC-mCerulean

281   iBMDMs stimulated with LPS and nigericin for 30 min (centre); or WT iBMDMs, stimulated

282   with LPS and nigericin for 1 h (right).

283   **(d)** 3-D segmentation models of the outer and inner envelope membranes (purple) and cristae

284   membranes (pink) in the tomographic slices shown immediately above in (**c**).

Discontinuities in the outer membranes of mitochondria during pyroptosis

During apoptosis, the Bax and Bak proteins create large disruptions in the outer mitochondrial

membrane[40,43]. Our cryo-ET reconstructions showed that mitochondria in pyroptotic iBMDMs

stimulated with LPS/nigericin lacked any such large outer-membrane disruptions. We did,

however, identify small discontinuities in the outer mitochondrial membrane in some of the

stimulated ASC-mCerulean and WT iBMDMs (**Fig. 5c,d**). The size of these gaps varied from 10

to 22 nm (**Fig. S5b**). The presence of an intact inner membrane adjacent to the gap in the outer-

membrane implied that the gap was not due to the missing wedge effect or the anisotropic

resolution of cryo-ET tomograms[44].

The presence of 10-22-nm gaps in the outer membranes of mitochondria during

pyroptosis raises the question of whether these gaps cause mitochondrial outer membrane

permeabilization (MOMP) during inflammasome stimulation. To address this question, we

measured the cryo-EM density in the intermembrane space and in an adjacent cytosolic area,

where outer-membrane gaps were present. Unexpectedly, we found that the density at sites

where outer membrane gaps were present was higher than in the cytosol, indicating an increase

in macromolecular density at gap sites (**Fig. 5e,f**). In contrast, at sites where both membranes

were intact the intermembrane space had a lower density than the cytosol, as expected. The

increased cryo-ET density at gap sites spanned 10-22 nm along the Z-axis, matching the

dimensions of the gaps in X and Y. We conclude that protein-rich disruptions of the outer

mitochondrial membrane 10-22 nm in diameter form during pyroptosis.

Figure 5. Outer mitochondrial membrane discontinuities in pyroptotic cells.

(a) Cryo-ET reconstruction of a mitochondrion in an unstimulated WT iBMDM with closeup

panels defining the following morphological parameters: inner-to-outer membrane spacing,

inter-cristae spacing, crista width and crista apex angle.

(b) Quantitative analysis of the morphological parameters quantified in (**a**) in 3-D segmented

surfaces of the mitochondrial membranes, as shown in (**Fig. 4d**). Error bars represent

standard deviation between measurements. Parameters were measured from four

tomograms for the control and two tomograms for each of the nigericin-stimulated samples.

Error bars represent standard deviation from the mean. For the crista apex angles: "control",

n = 327; "30 min", n = 187; "1 h", n = 96. For other parameters: "control", n =16; "30 min", n

= 8; "1 h", n = 8. Statistical test: two-way ANOVA, ns, $P > 0.05$; ****, $P < 10^{-4}$. See **Dataset**

S3 for source data.

(c) Cryo-ET reconstructions of mitochondria in ASC-mCerulean iBMDMs stimulated with

nigericin for 30 min (left) and in WT iBMDMs stimulated with nigericin for 1 h (centre, right).

Dashed boxes denote outer-membrane discontinuities. Scale bars, 100 nm.

(d) Higher magnification view of a mitochondrial outer-membrane discontinuity (dashed box).

Scale bar, 100 nm.

(e) Z-stack series for the boxed region in (**d**). Arrows mark the outer-membrane discontinuity.

Scale bar, 20 nm.

(f) Quantitative analysis of the cryo-ET density at the outer-membrane (OM) gap site in the Z-

stack series shown in (**e**), and at a site in the same tomogram with intact inner and outer

membranes. The graph plots the average pixel density ratio $\langle\rho(IMS)\rangle/\langle\rho(cyt)\rangle$ for each Z-

slice, where $\langle\rho(IMS)\rangle$ and $\langle\rho(cyt)\rangle$ are the average pixel densities of circular areas 35.4 nm

in diameter, in the mitochondrial intermembrane space (IMS) and an adjacent cytosolic

area, respectively. See **Dataset S2** for source data.

Discussion

NLRP3 inflammasome signalling assemblies have long been known from fluorescence

microscopy imaging to form puncta, or specks, containing ASC and caspase-1. Room

temperature electron tomography of ASC puncta in fixed zebrafish sections[26] and iBMDMs (**Fig.**

S3d) showed a dense filament network within the punctum, but the chemical fixation and

staining agents precluded further ultrastructure from being resolved. Our cryo-ET

reconstructions, obtained in unstained and fully hydrated conditions, show that the speck is

formed of a filamentous network consisting of hollow-tube branched filaments with the

dimensions predicted for ASC filaments, based on structural studies of purified ASC PYD

filaments and full-length monomeric ASC[23,24]. The identification of a filamentous network

containing mCerulean-labelled ASC positively identifies the protein as the principal filament-

forming component in the puncta. These data are supported by analysis of similar structures in

wild-type cells labelled with a fluorescent caspase 1 substrate. The filament structure of the

ASC with an mCerulean tag on the C-terminus supports the model whereby the PYD forms the

filament core, and the CARD decorates the filament. The extensive filament branching visible in

the reconstructions is thought to depend on CARD-CARD interactions because the CARD is

required for puncta formation, whereas the ASC PYD alone forms unbranched filaments that

lack the structural stability that is inherent to a branched network[17,18,23-26].

The density of ASC filaments varied in the cryo-ET reconstructions, with multiple core

regions of densely packed filaments separated by sparser regions. This suggests that the

higher-order structure of the punctum was likely seeded by multiple oligomeric assemblies,

rather than growing concentrically from a single nucleation site. In contrast with huntingtin and

other neurotoxic protein aggregates, which exclude other macromolecules based on cryo-ET

reconstructions[45], the structural organisation of the ASC filament network allows ribosomes and

small TGN-like vesicles to be retained in the puncta, or to permeate through them. Permeability

of the ASC network to macromolecules could be important for efficient recruitment and release

of effector, substrate and product molecules. Permeability to vesicles would allow active

transport of TGN vesicles carrying activated inflammasome seed-oligomers from sites of NLRP3

activation to the site of punctum formation, as recently proposed[12]. Moreover, the ribosomes

within the puncta could potentially contribute to the pyroptotic signalling programme, for

example by expressing proteins required for signalling, or by responding to danger signals.

Indeed, NLRP3 signalling can be activated by translational arrest through direct and indirect

mechanisms[46], including binding of fungal polysaccharides to ribosomes[47].

Our cryo-ET reconstructions, in addition to revealing the novel ultrastructure of ASC

within the cell, also allowed morphometric analysis of mitochondria in the inflammasome

activated cellular environment. The clearest differences between mitochondria in inflammasome

activated cells and unstimulated cells were a two to fourfold reduction in cristae spacing, and

the appearance of protein-rich discontinuities of 10-20 nm in diameter in the outer membrane,

which are distinct from BAK/BAX macropores. Permeabilization of both the plasma membrane

and outer mitochondrial membrane was reported previously in cells undergoing caspase-1-

dependent pyroptosis after inflammasome activation[41]. Pore formation in the plasma membrane

by the N-terminal domain of gasdermin D (GSDMD) after cleavage by caspase-1 is required for

inflammasome-dependent pyroptosis[48]. The outer-membrane discontinuities in the mitochondria

of inflammasome-activated cells had similar dimensions to GSDMD pores, which have an

average inner diameter of 22 nm[48] (or 13-34 nm based on atomic force microscopy[49]). Future

studies are warranted to test the hypothesis that the outer-membrane gaps observed by cryo-

ET may correspond to mitochondrial GSDMD pores.

The filament branching and packing density within ASC puncta provide structural

integrity while allowing downstream signalling molecules to diffuse freely and bind at high

density within the network. Although many structural details of the NLR inflammasome signalling

machinery remain unclear, this study demonstrates the potential for combined cryo-CLEM and

cryo-ET approaches to extract detailed, hypothesis-generating ultrastructural information for

fully assembled innate immune signalling complexes in their cellular context.

Materials and Methods

Mammalian cell culture. Wild-type immortalized primary bone-marrow derived macrophages (WT iBMDMs) and ASC-mCerulean iBMDMs, kindly provided by Eicke Latz (Univ. Bonn), were cultured in Dulbecco's modified Eagle Medium high glucose (DMEM; Gibco, Waltham, MA, USA), supplemented with 10% v/v foetal bovine serum (FBS; Gibco). Human monocytic THP-1 cells (European Collection of Authenticated Cell Cultures) were grown in Roswell Park Memorial Institute (RPMI) 1640 Medium (Gibco), supplemented with heat inactivated 10% v/v FBS, 10 mM HEPES pH 7.4, 1 mM Sodium Pyruvate and 0.05 beta-mercaptoethanol. Cells were checked regularly for mycoplasma contamination by MycoAlert™ mycoplasma detection kit (Lonza) and cells were free of mycoplasma contamination.

Inflammasome stimulation. For stimulation of WT iBMDMs and ASC-mCerulean iBMDMs, cells were primed with 200 ng/ml LPS for 3 h in Opti-MEM® (Gibco) or DMEM supplemented with 10% FBS before NLRP3 stimulus was used. The following stimulus were used: 10 μM nigericin (Sigma or Invivogen) and 5 mM ATP (pH adjusted to 7.4, Sigma) for 30 min or 60 min with pan-caspase inhibitor Z-VAD-FMK (Invivogen) or carboxyfluorescein-labelled inhibitor of caspase-1 YVAD (FAM-FLICA, ImmunoChemistry Technologies) for *in situ* cryo-ET sample preparation unless otherwise specified. For stimulation of THP-1 cells, priming and stimulation were performed in RPMI Medium and conditions were listed above. To stain active caspase-1, the supernatant was removed following inflammasome stimulation; replaced with DMEM supplemented with heat-inactivated 10% v/v FBS containing 0.5x reconstituted FAM-FLICA (ImmunoChemistry Technologies).

Cytokine measurement and immunoblotting. IL-1β secretion was measured by ELISA as previously described[27]. Briefly, supernatants were collected after inflammasome activation and BD BioSciences OptEIA was used to measure mouse IL-1β concentration in cell supernatant according to the manufacturer's instructions.

The cells were lysed in cell lysis buffer (150 mM NaCl, 50 mM Tris-HCl pH 8, 1% Triton

X-100, 1 mM PMSF, 10 µg/mL leupeptin, 1 µg/mL aprotinin) on ice for 10 min before

centrifugation. Supernatant was collected and boiled in SDS sample loading buffer for 5 min.

Following separation by 4-20% gradient SDS-PAGE, the proteins were transferred to a PVDF

membrane. Membrane was blocked in 5% milk in PBS + 0.2% Tween-20. The primary antibody

was rabbit monoclonal anti-mouse GSDMD (Abcam, ab209845, RRID:AB_2783550). The

secondary antibody was goat anti rabbit IgG-HRP (Santa Cruz Biotechnology, sc-2004,

RRID:AB_631746).

Immunofluorescence staining. For immunofluorescence, cells were plated as monolayer on

µ-Slide 8-Well chamber (ibiTreat, Ibidi, 80826) or 12-well chamber (removable, Ibidi, 81201).

Cells were primed and stimulated as above, and then fixed with 4% paraformaldehyde for 5 to

10 min at room temperature. Cells were permeabilised and blocked in 0.1% Saponin (Sigma,

47036) supplemented with 20% FBS (Gibco) or 2% Bovine Serum Albumin (Sigma, A7030) in

PBS with primary antibody overnight. Washing was performed before cells were incubated with

secondary antibodies. The following primary antibodies were used: rabbit anti-ASC rabbit

polyclonal pAb AL177, 1:200 dilution (AdipoGen, AG-25B-0006, RRID:AB_2490440); rabbit

anti-ASC monoclonal ASC/TMS1(D2W8U), 1:800 dilution (Cell Signalling Technology, 67824,

RRID:AB_2799736); goat anti-NLRP3 polyclonal, 1:200 dilution (Abcam, ab4207,

RRID:AB_955792); goat anti-IL-1β polyclonal, 1:500 dilution (R&D Systems, AF-401-NA,

RRID:AB_416684); mouse anti-γ-Tubulin monoclonal, 1:400 dilution (Sigma-Aldrich, T6557,

RRID:AB_477584); rabbit anti-TGN38 polyclonal, 1:250 dilution (Novus Biologicals, NBP1-

03495, RRID:AB_1522533); rabbit anti-TOM20 FL-145 polyclonal, 1:80 dilution (Santa Cruz

Biotechnology, sc-11415, RRID:AB_2207533). The Alexa Fluor secondary antibodies (488, 555,

568 and 647) used were: 488-labelled donkey anti-goat IgG (H+L) (1:500, ThermoFisher,

A11055, RRID:AB_2534102); 555-labelled goat anti-rabbit IgG(H+L) (1:500, ThermoFisher,

A21428 RRID:AB_141784); 568-labelled goat anti-mouse IgG (H+L) (1:500, ThermoFisher,

A11004, RRID:AB_2534072); 647-labelled goat anti-rabbit IgG (H+L) (1:500, ThermoFisher,

A21244, RRID:AB_2535812). After incubation with secondary antibodies at room temperature

for 1 h, cells were washed three times with blocking buffer, one time with PBS and final wash

with water before mounting with ProLong Gold Antifade Mountant with DAPI (ThermoFisher).

Confocal fluorescence imaging. Single point-scanning confocal microscopy was carried out

on a Zeiss (Oberkochen, Germany) LSM 780 or LSM 710 microscope using a 40x/1.3 NA Fluar

or 63x/1.4 NA Plan-apochromat oil immersion objective lens. The microscope was equipped

with 405, 458, 488, 514, 561 and 633 nm laser lines. Multi point-scanning confocal microscopy

was carried out on Visitech (Sunderland, UK) iSIM mounted on a Nikon (Tokyo, Japan) Ti2

microscope stand using a 100x/1.49 NA SR Apo TIRF oil immersion objective lens or a 60x/1.2

NA Plan Apo VC water immersion objective lens. The iSIM was equipped with 405, 445, 488,

561 and 640 nm laser lines and Hamamatsu (Hamamatsu, Japan) ORCA-Flash4.0 V3 sCMOS

cameras. Filter ranges: green (500-545 nm), red (593-624 nm). Live-cell samples were heated

to 37˚C and supplemented with 5% CO_2 using a microscope incubation chamber.

Live-cell imaging and Fluorescence Recovery after Photobleaching (FRAP). ASC-

mCerulean iBMDMs were primed and induced as previously described with the presence of

caspase-1 inhibitor (Z-VAD-FMK). Confocal FRAP was performed on a Zeiss LSM 710

microscope equipped with a 63x/1.4 NA Plan-apochromat oil immersion objective lens and a

458 nm laser line for excitation and bleaching. The sample environment was heated to 37˚C and

supplemented with 5% CO_2 using a microscope incubation chamber. ASC-mCerulean speck

was photobleached with 100% laser power. Images were acquired at 3 s intervals. Images were

collected at three pre-bleach timepoints and for 160 s post-bleaching. Movies were analysed in

ImageJ/Fiji[42] using a customised script (https://github.com/jboulanger/imagej-

macro/tree/main/FRAP_Measure). Briefly, measures were normalised to account for the general

photobleaching caused by image acquisition as well as sample motion over time. Normalised

fluorescence intensity measurements were obtained after background and bleaching

corrections. The timeframe varied for experiments (n = 8). Each measurement was interpolated

using the normalised fluorescence intensity measurements and imported to Python for plotting

of the fluorescence intensity curve.

Quantification of Mitochondrial membrane potential. After 30 min of inflammasome

stimulation as described above, iBMDMs were incubated with 20 nM TMRM (Life Technologies)

for 15 min. iBMDMs were incubated for 3 h in 50 μM of the mitochondrial uncoupler carbonyl

cyanide 3-chlorophenylhydrazone (CCCP; ThermoFisher, M20036). Control samples included:

untreated; TMRM+; TMRM+CCCP+; TMRM+LPS+ iBMDMs. TMRM fluorescence was

quantified by flow cytometry and data were acquired on Eclipse flow cytometer (Sony

Biotechnology). Quantification was set at 100,000 for cell count and a spectrum window of (FL3

(595BP)) was used to detect TMRM signal. Data were analysed and visualised using Flowjo10

(TreeStar, Ashland, Oregon, USA).

Cryo-ET sample preparation. Quantifoil finder grids (R2/1 or R2/2, Au 200 mesh grid,

Quantifoil Micro Tools, Germany) were glow discharged with a 30 mA current for 30 s using

Edwards Sputter Coater (S150B). Then grids were sterilized by UV irradiation for 10 min and

immersed in PBS supplemented with fibronectin solution (10 μg/ml, Sigma) overnight in 8-well

or coculture wells (ibidi). Grids were washed with PBS three times. Cells were seeded in 8-well

or coculture wells (ibidi) and incubated overnight at 37˚C and 5% CO_2. Cells cultivated on grids

were primed, induced with inflammasome stimuli and were plunge-frozen in liquid ethane using

Leica EM GP2 cryo-plunger. Prior to plunging, 4 μl of cell culture medium was added to the cell

side and backside blotting was applied for 6-8 s. The chamber conditions were maintained at

37˚C, 100% humidity during freezing. Grids were stored in liquid nitrogen.

Cryo-fluorescent light microscopy (Cryo-FM) and cryo-focused ion-beam (FIB) milling.

Grids were screened for cells with ASC/caspase-1 speck by light microscopy using a cryo-FM

microscope (Leica EM Cryo CLEM, Leica Microsystems), equipped with a cryo-stage, an

ORCA-Flash4.0 V2 sCMOS camera (Hamamatsu Photonics) and a 50x/0.9 NA HCX PL APO

cryo-objective lens. Montage acquisition of grids was performed with Leica LAS X software,

while recording the following channels: green (L5 filter, 50 ms), far red (Y5 filter, 20 ms), and

brightfield (50 ms). Z-stacks were recorded at 0.5 μm with step size 21 for each grid square to

determine the best focus for the montage. The montage was completed by stitching the best

focus image from the Z-stack with Leica LAS X.

Lamellae were prepared using a Scios Dual Beam FIB scanning electron microscope

(SEM; ThermoFisher) equipped with a Quorum cryo-stage (PP3010T). The milling protocol was

adapted from previously published method[29]. Grids were coated with organometallic platinum

using the gas injection system (GIS; ThermoFisher) operated at RT, for 8 s, at 12 mm working

distance and 25˚ stage tilt. A first rough milling was performed at 25˚ stage tilt with a 30 kV ion

beam voltage and 1 nA current until the lamella thickness reached 10 μm. The micro-expansion

joints were applied to improve lamella stability[32] using the milling parameters listed above. Then

stage was tilted to 20˚ for all subsequent milling steps. Rough milling steps were applied as

following with 30 kV ion beam voltage: a 5 μm lamella thickness was reached with current 0.5

nA; 3 μm lamella thickness was reached with a 0.3 nA current; and 1 μm lamella thickness was

obtained with a 0.1 nA current. Fine milling to a final lamella thickness of approximately 200 nm

was performed with ion beam settings of 30 kV and 50 pA or 10 pA, or 10 kV and 23 pA or 11

pA.

Grids were loaded to a Linkam cryo-stage (CMS196V), and lamellae were imaged on a

Zeiss microscope equipped with Axiocam 503 mono and Colibri 7-illumination module R(G/Y)

CBV-UV. Z-stacks of the lamellae were acquired to identify ASC/caspase-1 specks using a 200-

Cryo-fluorescent light microscopy (Cryo-FM) and cryo-focused ion-beam (FIB) milling.

Grids were screened for cells with ASC/caspase-1 speck by light microscopy using a cryo-FM microscope (Leica EM Cryo CLEM, Leica Microsystems), equipped with a cryo-stage, an ORCA-Flash4.0 V2 sCMOS camera (Hamamatsu Photonics) and a 50x/0.9 NA HCX PL APO cryo-objective lens. Montage acquisition of grids was performed with Leica LAS X software, while recording the following channels: green (L5 filter, 50 ms), far red (Y5 filter, 20 ms), and brightfield (50 ms). Z-stacks were recorded at 0.5 μm with step size 21 for each grid square to determine the best focus for the montage. The montage was completed by stitching the best focus image from the Z-stack with Leica LAS X.

Lamellae were prepared using a Scios Dual Beam FIB scanning electron microscope (SEM; ThermoFisher) equipped with a Quorum cryo-stage (PP3010T). The milling protocol was adapted from previously published method[29]. Grids were coated with organometallic platinum using the gas injection system (GIS; ThermoFisher) operated at RT, for 8 s, at 12 mm working distance and 25˚ stage tilt. A first rough milling was performed at 25˚ stage tilt with a 30 kV ion beam voltage and 1 nA current until the lamella thickness reached 10 μm. The micro-expansion joints were applied to improve lamella stability[32] using the milling parameters listed above. Then stage was tilted to 20˚ for all subsequent milling steps. Rough milling steps were applied as following with 30 kV ion beam voltage: a 5 μm lamella thickness was reached with current 0.5 nA; 3 μm lamella thickness was reached with a 0.3 nA current; and 1 μm lamella thickness was obtained with a 0.1 nA current. Fine milling to a final lamella thickness of approximately 200 nm was performed with ion beam settings of 30 kV and 50 pA or 10 pA, or 10 kV and 23 pA or 11 pA.

Grids were loaded to a Linkam cryo-stage (CMS196V), and lamellae were imaged on a Zeiss microscope equipped with Axiocam 503 mono and Colibri 7-illumination module R(G/Y) CBV-UV. Z-stacks of the lamellae were acquired to identify ASC/caspase-1 specks using a 200-

300 nm step size. A projected cryo-fluorescent image was used to correlate with an SEM image

using the EC-CLEM plugin from ICY[36]. Registration between the lamella map and SEM image

was achieved by transforming coordinates using edges or salient features of the lamella. The

maximum projection Z-stack was then saved for correlation with SerialEM[32,33].

Tilt-series acquisition and tomogram reconstruction. Grids with a fluorescent lamella were

transferred to a 300 kV Titan Krios electron microscope (ThermoFisher) equipped with an

energy filter (Gatan). Movies were acquired with a K2 or K3 direct electron detector with

SerialEM[33,34]. Cryo-FM/EM correlation was then used to locate an area of interest for tilt-series

acquisition. Briefly, a medium-magnification montage EM map (MMM) of the lamella was

generated and a fluorescence map (FM) was loaded into SerialEM. The maps were correlated

through recognition of geometric edges of the lamella. Transformation of coordinates yielded an

overlay map which provided guidance for tilt-series acquisition. Tilt-series were collected at a

nominal 32,000x or 29,000x magnification, resulting in pixel size 3.4 Å or 2.7 Å, over a tilt range

of -60˚ to +60˚ with 1˚, or 2˚ increments, a total dose of 140-240 electrons/A^2 and a nominal

defocus range of -4 to -8 μm. A dose-symmetric scheme was used[50]. A pre-processing script

was used from SubTOM (by Dustin Morado; https://github.com/DustinMorado/subTOM).

Frames were aligned using IMOD[37]. Reconstruction was performed by weighted back-

projection, and segmentation with IMOD. The MATLAB script deconv from Warp[51] was used for

visualisation. Filament tracing was performed manually with IMOD and UCSF Chimera[52].

Membrane segmentation was performed with TomoSegMemTV[53].

Quantifications and statistical analysis. Filament lengths were measured using IMOD[37] from

six tomograms of ASC-mCerulean iBMDMs and one tomogram of WT iBMDMs labelled with

FAM-FLICA. Histograms were plotted with MATLAB. Branching angles were measured with

Fiji[42] and exported to MATLAB for plotting. The filament diameter was determined in Fiji by

measuring density profiles of 20-nm lines perpendicular to the ASC filament (15 pixels in width,

ASC-mCerulean tomogram with13.6 Å/pixel and WT ASC tomogram with 10.78 Å/pixel). The

grey value profiles were normalized to their maximal individual value. Normalized grey profile

were aligned and exported to Python for plotting. Similarly, diameters of trans-Golgi-like vesicles

were measured in Fiji and exported to MATLAB for plotting.

537   Mitochondrial tomograms were loaded in Fiji[42] and ROI manager was used to measure

morphological parameters, which were compared using Two-way ANOVA and plotted with

GraphPad Prism. For the inner-to-outer membrane spacing, the distance between the nearest

edges of the inner and outer membranes was measured. The cristae width was measured

between the inner membranes of the cristae. The spacing between cristae were measured

using the perpendicular distance between the internal border of the neighbouring cristae. The

inner-to-outer membrane spacing, cristae width and inter-cristae spacing were measured

perpendicular to the membranes, evenly spaced for each tomogram, with thirty values per slice,

for four slices per tomogram, from eight tomograms. The cristae apex angles were measured for

every tip visible in ten slices per tomogram. Every value was plotted along with the average. If

the end of a crista was flat, then the measurement of the two angles on either side was

measured. Any measurement was only taken if the internal border of the membrane was clearly

distinguishable from the matrix. All measurements were evenly spaced through each slice, to

give a value representative of the whole slice, using all the mitochondria present in each image.

All measurements were taken using 8 different tomograms: four control, two 30-min nigericin-

treated, and two 60-min nigericin-treated mitochondria.

553   For measurement of the outer-mitochondrial membrane gap density, we calculated the

density ratio as follows. A total of 21 virtual slices were selected in a region containing a visible

outer-membrane gap. The average pixel grey value was calculated from circular areas 13.6 nm

in diameter: three cytosolic areas, three inter-membrane space areas and an outer-membrane

gap area. This was repeated for each of the 21 virtual slices. The ratio of average outer-

membrane gap site to cytosolic grey value, and the ratio of average inter-membrane space site

to cytosolic grey value were calculated for each virtual slice. The grey area ratio was then

plotted with Igor (WaveMetrics, Inc.) to show the change of protein density in Z direction.

Data availability

Representative electron tomograms were deposited in EMDB with codes EMDB with codes

EMD-13585, EMD-13586, EMD-135XX and EMD-135XX. Other data are available from the

corresponding authors upon request.

Code availability

SubTOM scripts and instructions are available from Github

[https://github.com/DustinMorado/subTOM]. The FRAP ImageJ macro is available on Github

[https://github.com/jboulanger/imagej-macro/tree/main/FRAP_Measure].

References

1   Xue, Y., Enosi Tuipulotu, D., Tan, W. H., Kay, C. & Man, S. M. Emerging Activators and

Regulators of Inflammasomes and Pyroptosis. *Trends Immunol* **40**, 1035-1052,

doi:10.1016/j.it.2019.09.005 (2019).

2   Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in

141,456 humans. *Nature* **581**, 434-443, doi:10.1038/s41586-020-2308-7 (2020).

3   Landrum, M. J. *et al.* ClinVar: improvements to accessing data. *Nucleic Acids Res* **48**,

D835-D844, doi:10.1093/nar/gkz972 (2020).

4   Gong, T., Liu, L., Jiang, W. & Zhou, R. DAMP-sensing receptors in sterile inflammation and

inflammatory diseases. *Nature Reviews Immunology* **20**, 95-112, doi:10.1038/s41577-019-

0215-7 (2020).

581 5 Heneka, M. T., McManus, R. M. & Latz, E. Inflammasome signalling in brain function and

582  neurodegenerative disease. *Nature Reviews Neuroscience* **19**, 610-621,

583  doi:10.1038/s41583-018-0055-7 (2018).

584 6 Swanson, K. V., Deng, M. & Ting, J. P. The NLRP3 inflammasome: molecular activation

585  and regulation to therapeutics. *Nat Rev Immunol* **19**, 477-489, doi:10.1038/s41577-019-

586  0165-0 (2019).

587 7 Chen, J. & Chen, Z. J. PtdIns4P on dispersed trans-Golgi network mediates NLRP3

588  inflammasome activation. *Nature* **564**, 71-76, doi:10.1038/s41586-018-0761-3 (2018).

589 8 Shen, C., Sharif, H., Xia, S. & Wu, H. Structural and mechanistic elucidation of

590  inflammasome signaling by cryo-EM. *Current Opinion in Structural Biology* **58**, 18-25,

591  doi:10.1016/j.sbi.2019.03.033 (2019).

592 9 Diebolder, C. A., Halff, E. F., Koster, A. J., Huizinga, E. G. & Koning, R. I. Cryoelectron

593  Tomography of the NAIP5/NLRC4 Inflammasome: Implications for NLR Activation.

594  *Structure* **23**, 2349-2357, doi:10.1016/j.str.2015.10.001 (2015).

595 10 Zhang, L. *et al.* Cryo-EM structure of the activated NAIP2-NLRC4 inflammasome reveals

596  nucleated polymerization. *Science* **350**, 404-409, doi:10.1126/science.aac5789 (2015).

597 11 Hu, Z. *et al.* Structural and biochemical basis for induced self-propagation of NLRC4.

598  *Science* **350**, 399-404, doi:10.1126/science.aac5489 (2015).

599 12 Magupalli, V. G. *et al.* HDAC6 mediates an aggresome-like mechanism for NLRP3 and

600  pyrin inflammasome activation. *Science* **369**, doi:10.1126/science.aas8995 (2020).

601 13 Li, X. *et al.* MARK4 regulates NLRP3 positioning and inflammasome activation through a

602  microtubule-dependent mechanism. *Nat Commun* **8**, 15986, doi:10.1038/ncomms15986

603  (2017).

604 14 Oroz, J., Barrera-Vilarmau, S., Alfonso, C., Rivas, G. & de Alba, E. ASC Pyrin Domain Self-

605  associates and Binds NLRP3 Protein Using Equivalent Binding Interfaces. *Journal of*

606  *Biological Chemistry* **291**, 19487-19501, doi:10.1074/jbc.M116.741082 (2016).

15   Stehlik, C. *et al.* Apoptosis-associated speck-like protein containing a caspase recruitment

domain is a regulator of procaspase-1 activation. *J Immunol* **171**, 6154-6163,

doi:10.4049/jimmunol.171.11.6154 (2003).

16   Proell, M., Gerlic, M., Mace, P. D., Reed, J. C. & Riedl, S. J. The CARD plays a critical role

in ASC foci formation and inflammasome signalling. *The Biochemical journal* **449**, 613-621,

doi:10.1042/BJ20121198 (2013).

17   Dick, M. S., Sborgi, L., Rühl, S., Hiller, S. & Broz, P. ASC filament formation serves as a

signal amplification mechanism for inflammasomes. *Nature Communications* **7**, 11929,

doi:10.1038/ncomms11929 (2016).

18   Sahillioglu, A. C., Sumbul, F., Ozoren, N. & Haliloglu, T. Structural and dynamics aspects of

ASC speck assembly. *Structure* **22**, 1722-1734, doi:10.1016/j.str.2014.09.011 (2014).

19   Matyszewski, M. *et al.* Cryo-EM structure of the NLRC4(CARD) filament provides insights

into how symmetric and asymmetric supramolecular structures drive inflammasome

assembly. *J Biol Chem* **293**, 20240-20248, doi:10.1074/jbc.RA118.006050 (2018).

20   Gong, Q. *et al.* Structural basis for distinct inflammasome complex assembly by human

NLRP1 and CARD8. *Nat Commun* **12**, 188, doi:10.1038/s41467-020-20319-5 (2021).

21   Li, Y. *et al.* Cryo-EM structures of ASC and NLRC4 CARD filaments reveal a unified

mechanism of nucleation and activation of caspase-1. *Proc Natl Acad Sci U S A* **115**,

10845-10852, doi:10.1073/pnas.1810524115 (2018).

22   Sborgi, L. *et al.* Structure and assembly of the mouse ASC inflammasome by combined

NMR spectroscopy and cryo-electron microscopy. *Proc Natl Acad Sci U S A* **112**, 13237-

13242, doi:10.1073/pnas.1507579112 (2015).

23   Lu, A. *et al.* Unified polymerization mechanism for the assembly of ASC-dependent

inflammasomes. *Cell* **156**, 1193-1206, doi:10.1016/j.cell.2014.02.008 (2014).

24   de Alba, E. Structure and interdomain dynamics of apoptosis-associated speck-like protein

containing a CARD (ASC). *J Biol Chem* **284**, 32932-32941, doi:10.1074/jbc.M109.024273

(2009).

25   Nambayan, R. J. T., Sandin, S. I., Quint, D. A., Satyadi, D. M. & de Alba, E. The

inflammasome adapter ASC assembles into filaments with integral participation of its two

Death Domains, PYD and CARD. *J Biol Chem* **294**, 439-452,

doi:10.1074/jbc.RA118.004407 (2019).

26   Kuri, P. *et al.* Dynamics of in vivo ASC speck formation. *J Cell Biol* **216**, 2891-2909,

doi:10.1083/jcb.201703103 (2017).

27   Man, S. M. *et al.* Inflammasome activation causes dual recruitment of NLRC4 and NLRP3

to the same macromolecular complex. *Proc Natl Acad Sci U S A* **111**, 7403-7408,

doi:10.1073/pnas.1402911111 (2014).

28   Rigort, A. *et al.* Focused ion beam micromachining of eukaryotic cells for cryoelectron

tomography. *Proc Natl Acad Sci U S A* **109**, 4449-4454, doi:10.1073/pnas.1201333109

(2012).

29   Schaffer, M. *et al.* Cryo-focused Ion Beam Sample Preparation for Imaging Vitreous Cells

by Cryo-electron Tomography. *Bio Protoc* **5**, doi:10.21769/bioprotoc.1575 (2015).

30   Stutz, A., Horvath, G. L., Monks, B. G. & Latz, E. ASC speck formation as a readout for

inflammasome activation. *Methods Mol Biol* **1040**, 91-101, doi:10.1007/978-1-62703-523-

1_8 (2013).

31   Smolewski, P. *et al.* Detection of caspases activation by fluorochrome-labeled inhibitors:

Multiparameter analysis by laser scanning cytometry. *Cytometry* **44**, 73-82,

doi:10.1002/1097-0320(20010501)44:1<73::aid-cyto1084>3.0.co;2-s (2001).

32   Wolff, G. *et al.* Mind the gap: Micro-expansion joints drastically decrease the bending of

FIB-milled cryo-lamellae. *Journal of Structural Biology* **208**, 107389,

doi:10.1016/j.jsb.2019.09.006 (2019).

33 Schellenberger, P. *et al.* High-precision correlative fluorescence and electron cryo

microscopy using two independent alignment markers. *Ultramicroscopy* **143**, 41-51,

doi:10.1016/j.ultramic.2013.10.011 (2014).

34 Mastronarde, D. N. Automated electron microscope tomography using robust prediction of

specimen movements. *J Struct Biol* **152**, 36-51, doi:10.1016/j.jsb.2005.07.007 (2005).

35 Guo, Q. *et al.* In Situ Structure of Neuronal C9orf72 Poly-GA Aggregates Reveals

Proteasome Recruitment. *Cell* **172**, 696-705 e612, doi:10.1016/j.cell.2017.12.030 (2018).

36 Paul-Gilloteaux, P. *et al.* eC-CLEM: flexible multidimensional registration software for

correlative microscopies. *Nat Methods* **14**, 102-103, doi:10.1038/nmeth.4170 (2017).

37 Kremer, J. R., Mastronarde, D. N. & McIntosh, J. R. Computer Visualization of Three-

Dimensional Image Data Using IMOD. *Journal of Structural Biology* **116**, 71-76,

doi:10.1006/jsbi.1996.0013 (1996).

38 Donohoe, B. S. *et al.* Cis-Golgi cisternal assembly and biosynthetic activation occur

sequentially in plants and algae. *Traffic* **14**, 551-567, doi:10.1111/tra.12052 (2013).

39 Hwang, M. S. *et al.* MAVS polymers smaller than 80 nm induce mitochondrial membrane

remodeling and interferon signaling. *The FEBS journal*, doi:10.1111/febs.14772 (2019).

40 Ader, N. R. *et al.* Molecular and topological reorganizations in mitochondrial architecture

interplay during Bax-mediated steps of apoptosis. *eLife* **8**, doi:10.7554/eLife.40712 (2019).

41 Yu, J. *et al.* Inflammasome activation leads to Caspase-1-dependent mitochondrial damage

and block of mitophagy. *Proc Natl Acad Sci U S A* **111**, 15514-15519,

doi:10.1073/pnas.1414859111 (2014).

42 Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods*

9, 676-682, doi:10.1038/nmeth.2019 (2012).

43 McArthur, K. *et al.* BAK/BAX macropores facilitate mitochondrial herniation and mtDNA

efflux during apoptosis. *Science* **359**, doi:10.1126/science.aao6047 (2018).

44   Koning, R. I., Koster, A. J. & Sharp, T. H. Advances in cryo-electron tomography for biology

and medicine. *Ann Anat* **217**, 82-96, doi:10.1016/j.aanat.2018.02.004 (2018).

45   Bauerlein, F. J. B., Fernandez-Busnadiego, R. & Baumeister, W. Investigating the Structure

of Neurotoxic Protein Aggregates Inside Cells. *Trends Cell Biol* **30**, 951-966,

doi:10.1016/j.tcb.2020.08.007 (2020).

46   Vyleta, M. L., Wong, J. & Magun, B. E. Suppression of ribosomal function triggers innate

immune signaling through activation of the NLRP3 inflammasome. *PLoS One* **7**, e36044,

doi:10.1371/journal.pone.0036044 (2012).

47   Briard, B. *et al.* Galactosaminogalactan activates the inflammasome to provide host

protection. *Nature* **588**, 688-692, doi:10.1038/s41586-020-2996-z (2020).

48   Xia, S. *et al.* Gasdermin D pore structure reveals preferential release of mature interleukin-

1. *Nature*, doi:10.1038/s41586-021-03478-3 (2021).

49   Mulvihill, E. *et al.* Mechanism of membrane pore formation by human gasdermin-D. *EMBO

J* **37**, doi:10.15252/embj.201798321 (2018).

50   Hagen, W. J. H., Wan, W. & Briggs, J. A. G. Implementation of a cryo-electron tomography

tilt-scheme optimized for high resolution subtomogram averaging. *J Struct Biol* **197**, 191-

198, doi:10.1016/j.jsb.2016.06.007 (2017).

51   Tegunov, D. & Cramer, P. Real-time cryo-electron microscopy data preprocessing with

Warp. *Nat Methods* **16**, 1146-1152, doi:10.1038/s41592-019-0580-y (2019).

52   Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and

analysis. *J Comput Chem* **25**, 1605-1612 (2004).

53   Martinez-Sanchez, A., Garcia, I., Asano, S., Lucic, V. & Fernandez, J. J. Robust membrane

detection based on tensor voting for electron tomography. *J Struct Biol* **186**, 49-61,

doi:10.1016/j.jsb.2014.02.015 (2014).

54   Ader, N. R. & Kukulski, W. triCLEM: combining high-precision, room temperature CLEM

with cryo-fluorescence microscopy to identify very rare events. *Method Cell Biol* **140**, 303-

320, doi:10.1016/bs.mcb.2017.03.009 (2017).

55   Kukulski, W. *et al.* Correlated fluorescence and 3D electron microscopy with high sensitivity

and spatial precision. *Journal of Cell Biology* **192**, 111-119, doi:10.1083/jcb.201009037

(2011).

Acknowledgements

We acknowledge the MRC Laboratory of Molecular Biology (LMB) Electron Microscopy (EM)

Facility for access and support in EM sample preparation and data collection, and the Flow

Cytometry Facility for access and training. We are grateful to Dustin Morado for assisting in

cryo-ET data collection and image processing. We acknowledge the MRC LMB Light

Microscopy Facility, in particular Jonathan Howe, for access and support imaging data

collection. We are grateful for access to the MRC-LMB Scientific Computing facilities. We thank

Kendra E. Leigh, Wanda Kukulski, Emma Jones, Robert Pickering, Maria Daly, Victoria L. Hale

and Zunlong Ke for helpful advice and discussions. This work was supported by Wellcome

Senior Research Fellowship 217191/Z/19/Z to Y.M.; a PhD studentship to Y.L. from the China

Scholarship Council and Cambridge Trust; and Wellcome Investigator Award 108045/Z/15/Z to

C.E.B.

Author contributions

Y.L., L.J.H., C.E.B and Y.M. conceived the experiments. Y.L., H.Z. and J.H. collected light

microscopy data. Y.L. and J.B. analysed light microscopy data. Y.L., A.C.B. and C.H. prepared

cryo-EM samples, performed CLEM imaging and collected cryo-ET data. Y.L. processed the

cryo-ET data. Y.L. and H.A. analysed the cryo-ET data and flow-cytometry analysis. Y.L. and

L.J.H. performed cell-based assays. Y.L, C.E.B and Y.M. wrote the manuscript with input from

all authors. Project supervision, administration and funding acquisition, C.E.B. and Y.M.

Competing interests

C.E.B are Y.M. are consultants for Related Sciences LLC and have profits interests in Danger

Bio LLC. CEB is on the SAB of NodThera and Lightcast.

Cryo-electron tomography of ASC sites in pyroptotic cells

Liu *et al.*
740
741
742 **Supplementary Information**
743
744
745 **Supplementary Materials and Methods**
746
747 **Figures S1-S5**
748

Supplementary Materials and Methods

Correlative fluorescence microscopy and electron tomography of resin-embedded cells.

Correlative fluorescence microscopy and electron tomography (RT-CLEM) of resin-embedded

cells was performed as described[40,54]. Briefly, ASC-mCerulean cells were grown on carbon-

coated 3 mm sapphire discs (Wohlwend GmbH) in two-well chambers (ibidi) for 24 h before

priming and NLRPC-driven inflammasome stimulation as described above. Cells were stained

with MitoView far-red and FLICA before high pressure freezing with an HPM100 high pressure

freezing system (Leica Microsystems). Grids were imaged with a cryo-fluorescence microscope

(Leica EM Cryo CLEM, Leica Microsystems) to localise cells containing ASC-mCerulean

specks. Freeze substitution was preformed using 0.008% uranyl acetate in acetone and

embedded in Lowicryl HM20 (Polysciences) using an AFS2 (Leica Microsystems). Blocks were

sectioned into 300 nm thin sections using a microtome (Leica Microsystems) equipped with a

diamond knife (Diatome). The sections were collected on 200 mesh/300 mesh copper grids with

carbon support (Agar Scientific Ltd). TetraSpeck 100-nm microspheres were diluted 1:100 in

PBS and applied to the sections for use as fiducial markers for correlation. Sections were

imaged on a Nikon Ti2 wide field microscope equipped with a Niji LED light source (Bluebox

Optics, Blackwood, UK), a x100/1.49 NA Apo TIRF oil immersion objective lens and a Neo

sCMOS DC-152Q-C00-FI camera (Andor Technology, Belfast, UK). Filters: mCerulean (89006

filter set, Chroma Technology, Bellows Falls, USA), fluorescein (49002 filter set, Chroma),

MitoView Far Red (49006 filter set, Chroma). EM images were collected using a Tecnai F20

electron microscope (ThermoFisher) operated at 200 kV and a high-tilt tomography holder

(Fischione Instruments; Model 2020). An image montage of regions of interest was acquired on

a BM-Orius detector using TEM mode at 150-200 µm defocus using SerialEM v3.8.0[34] at a pixel

size of 1.1 nm. The correlation between the montaged map and fluorescent map was performed

by image transformation of registered fiducial markers in both image modalities as previously

774 described[55]. Tilt-series were acquired from approximately -60 to +60 with 1˚ increment at a pixel

775 size of 1.1 nm. Samples were rotated 90˚ to acquire dual axis tilt-series. Tomograms were

776 reconstructed and visualised with IMOD[37].

777
778
779
780

Figure S1. Tracking NLRP3 inflammasome activation in iBMDMs with fluorescence microscopy and biochemical assays.

(a) Snapshots of live iBMDMs expressing ASC-mCerulean. ASC specks formed within 10 min of NLRP3 inflammasome activation with LPS and nigericin. Merge, overlay of fluorescence and bright-field images. Scale bar, 10 μm.

(b) Biochemical assays of inflammasome activation in ASC-mCerulean iBMDMs: upper panel, ELISA measuring IL-1β secretion; lower panel, Western blot showing proteolytic cleavage of GSDMD following different stimuli. FL, full-length. NTD, N-terminal domain.

(c) Fluorescence microscopy of WT iBMDMs stimulated with LPS and either nigericin or ATP. Caspase-1 was detected with FAM-FLICA. NLRP3 and ASC were detected by immunostaining. Scale bars, 5 μm.

(d) Immunofluorescence microscopy of WT iBMDMs stimulated with LPS and either nigericin or ATP, with staining for IL-1β and ASC. Scale bars, 5 μm.

(e) Snapshots of live ASC-mCerulean iBMDMs stained with FAM-FLICA after nigericin stimulation showing ASC-mCerulean colocalization with caspase-1 and FAM-FLICA fluorescence intensity increasing over time. Scale bar, 5 μm.

Figure S2. Workflow for correlative cryo-light and electron microscopy (cryo-CLEM).
799
(a) Cryo-fluorescence microscopy (cryo-FM) of ASC-mCerulean iBMDMs, with brightfield image
800
overlaid. Cells were cultured on finder grids, primed with LPS, stimulated with nigericin for
801
30 min, labelled with MitoView and FAM-FLICA, vitrified, and then imaged. The white circle
802
denotes one area of interest. Scale bar: 300 µm.
803
(b) Grids were transferred to a scanning electron microscope (SEM) fitted with a focused-ion-
804
beam (FIB). The cell containing the area of interest, identified using the finder-grid markers,
805
was imaged by SEM. Scale bar, 40 µm.
806
(c) SEM images of the area of interest before and after FIB milling: Upper panel, before FIB
807
milling with initial target milling windows shown in brown (scale bar, 10 µm); lower panel,
808
side-on view of the lamella after FIB milling (scale bar, 4 µm).
809
(d) Cryo-FM image of a lamella containing an ASC speck with brightfield image overlaid. Scale
810
bar, 10 µm.
811
(e) Low magnification transmission cryo-EM map of the lamella shown in (**d**). Scale bar, 7 µm.
812
(f) Medium magnification cryo-EM map overlaid with the cryo-FM map after registration and
813
transformation. The region of interest for tilt-series acquisition is boxed. Scale bar, 1 µm.
814
(g) Reconstructed tomographic slice acquired in the area marked in (**f**) containing the
815
fluorescent signal. Several short filaments and small trans-Golgi-like vesicles were visible.
816
Scale bar, 300 nm. See **Movie S4** for a tomogram of this region.
817
818
819
820

Figure S3. Imaging of *trans*-Golgi-like vesicles within ASC puncta.

(a) Fluorescence micrographs of THP-1 cells with and without stimulation with LPS and
nigericin, with TGN (anti-TGN38) immunostaining. Scale bar, 10 µm.

(b) Two tomographic slices of ASC-mCerulean speck showing vesicles adjacent to filaments.
Scale bars, 100 nm.

(c-e) CLEM of ASC-mCerulean iBMDMs stimulated with nigericin for 30 min, high-pressure
frozen, chemically fixed, sectioned and imaged at room temperature (see Supplementary
Materials and Methods). (**c**) Low magnification EM map overlaid with ASC-mCerulean
fluorescence. Scale bar, 1 µm. (**d**) 5.5 nm-thick tomographic slice of in the region boxed in
(c). Scale bar, 200 nm. (**e**) Tomographic slices showing vesicles with low-density lumens
within ASC-mCerulean puncta. Scale bar, 100 nm. The histogram shows the vesicle
diameter distribution.

834
Figure S4. Localization of ASC and caspase-1 relative to MTOC-associated components.
(a) Fluorescence micrographs of ASC-mCerulean iBMDMs stimulated with LPS and nigericin,
with γ-tubulin immunostaining. Scale bar, 10 μm. Right, distribution of the distance between
ASC-mCerulean and anti-γ-tubulin fluorescence.
(b) Fluorescence micrographs of WT iBMDMs stimulated with LPS and nigericin. Cells were
stained with FAM-FLICA, and with antibodies against γ-tubulin and ASC. Scale bar, 10 μm.
Right, distribution of the distance between ASC/FAM-FLICA and anti-γ-tubulin fluorescence.
(c) Fluorescence micrographs of THP-1 cells stimulated with LPS and nigericin. Cells were
stained with FAM-FLICA, and with antibodies against ninein and ASC. Scale bar, 10 μm.
Right, distribution of the distance between ASC/FAM-FLICA and anti-ninein fluorescence.

b	iBMDM type	Nigericin stimulation (min)	Gap width (nm)	Estimated gap depth (nm)
	ASC-mCerulean	30	11	11
	ASC-mCerulean	30	15	12
	ASC-mCerulean	30	18	14
	ASC-mCerulean	30	11	14
	WT	60	20	-
	WT	60	17	17

Figure S5. Functional and ultrastructural analyses of mitochondria in iBMDMs stimulated
with nigericin.
(a) Additional flow cytometry histograms of iBMDMs. The following controls are shown to
accompany **Fig. 4b**: cells with no TMRM staining; cells stained with TMRM and a potent
mitochondrial uncoupling chemical, carbonyl cyanide 3- chlorophenylhydrazone (CCCP);
cells stained with TMRM only; and cells primed with LPS and stained with TMRM.
(b) Measured sizes of outer-membrane discontinuities in cryo-ET reconstructions of
mitochondria in WT and ASC-mCerulean iBMDMs, as partly shown in **Fig. 5c-e**.