1 Exposure to the gut microbiota from cigarette smoke-exposed mice exacerbates cigarette smoke extract-
2 induced inflammation in zebrafish larvae
3

4 Simone Morris[1]†, Kathryn Wright[1], Vamshikrishna Malyla[2,3], Warwick J Britton[1,4], Philip M Hansbro[2],
5 Pradeep Manuneedhi Cholan[1] *†, Stefan H Oehlers[1,5] *
6
Affiliations
1 Tuberculosis Research Program at the Centenary Institute, The University of Sydney, Camperdown, NSW,
Australia.
2 Centre for Inflammation, Centenary Institute and University of Technology Sydney, Faculty of Science,
School of Life Sciences, Sydney, NSW, Australia.
3 Discipline of Pharmacy, Graduate School of Health, University of Technology Sydney, Sydney, NSW,
Australia.
4 Department of Clinical Immunology, Royal Prince Alfred Hospital, Camperdown, Australia.
5 The University of Sydney, Faculty of Medicine and Health, Camperdown, NSW, Australia
† Co-first authors, contributed equally
* Co-corresponding authors: pradeep.manuneedhicholan@mq.edu.au or s.oehlers@centenary.org.au

18
**Abstract**

Cigarette smoke (CS)-induced inflammation leads to a range of diseases including chronic obstructive
pulmonary disease and cancer. The gut microbiota is a major modifying environmental factor that
determine the severity of cigarette smoke-induced pathology. Microbiomes and metabolites from CS-
exposed mice exacerbate lung inflammation via the gut-lung axis of shared mucosal immunity in mice but
these systems are expensive to establish and analyse. Zebrafish embryos and larvae have been used to
model the effects of cigarette smoking on a range of physiological processes and offer an amenable
platform for screening modifiers of cigarette smoke-induced pathologies with key features of low cost and
rapid visual readouts. Here we exposed zebrafish larvae to cigarette smoke extract (CSE) and characterised
a CSE-induced leukocytic inflammatory phenotype with increased neutrophilic and macrophage
inflammation in the gut. The CSE-induced phenotype was exacerbated by co-exposure to microbiota from
the faeces of CS-exposed mice, but not control mice. Microbiota could be recovered from the gut of
zebrafish and studied in isolation in a screening setting. This demonstrates the utility of the zebrafish-CSE
exposure platform for identifying environmental modifiers of cigarette smoking-associated pathology and
demonstrates that the CS-exposed mouse gut microbiota potentiates the inflammatory effects of CSE
across host species.

## Introduction

Smoking tobacco is one of the greatest public health challenges, accounting for more than 8 million deaths worldwide, per year (1). It ranks as the third leading risk factor for illness and death, including from diseases such as cancers and inflammatory disorders like inflammatory bowel disease (IBD) (1-4). Cigarette smoke (CS) contains a multitude of toxic compounds affecting cell function, foetal development, gut function and microbiome, neural activity, inflammation, and lung cancer (3, 5-12).

The toxicity of CS has further been demonstrated to alter the microbiome of smokers, resulting in respiratory and gut microbiome dysbiosis, which in turn is associated with increased peptic ulcers, IBD and gastrointestinal cancers via the gut-lung axis of shared mucosal immunity (3, 4, 6, 11, 13). This is suggested to occur through alterations to the immune system (5), mucin production (6) or through direct anti-microbial effects of the toxins within the smoke (14). Shifts in gut microbiome and the profile of gut metabolites and pathology are associated with chronic obstructive pulmonary disease (COPD) in humans and CS-induced emphysema in mice (4, 10, 11, 15, 16). Microbiomes and metabolites associated with COPD and CS exacerbate lung inflammation via the gut-lung axis of shared mucosal immunity contributing to lung pathology in mammals (4, 11, 17).

Zebrafish have been used to study the developmental defects caused by CS exposure. These studies demonstrated stunted embryonic development upon exposure to cigarette smoke extract (CSE) and smoke condensate (18-20). Exposure of adult zebrafish to CSE reduced the capacity for fin regeneration (19), as well as behavioural abnormalities in response to nicotine (7). A study by Progatzky *et al.* further modelled the respiratory effects of CS using adult zebrafish gills demonstrating increased pro-inflammatory cytokine production and reduced immune cell recruitment (21). Massarsky *et al.* used total particulate matter (TPM) from cigarettes to model the effects of CS in zebrafish, and showed cardiac malfunction and defects in brain vascularisation (22) as well as behavioural alterations (23).

Thus far, the effect of CS exposure on immune function has been unexplored in zebrafish embryos and larvae (19, 21). This transparent stage of development provides the opportunity to investigate the inflammatory effects of CSE by live imaging of fluorescently tagged leukocytes, and is amenable to simplified experimental manipulation by gene editing or treatment with small molecules by immersion (24). Here we define the pro-inflammatory effects of CSE on larval zebrafish and find a transferrable effect of mammalian CS-associated microbiota on CSE-induced inflammation in larval zebrafish.

## Methods

### Zebrafish husbandry

Zebrafish embryos were obtained through natural spawning (Sydney Local Health District Animal Welfare Committee Approval 17-036). The transgenic strains used to visualise leukocytes were $Tg(lyzC:DsRed)^{nz50}$, where neutrophils are labelled by DsRed2 expression (25), and $Tg(mfap4:turquoise)^{xt27}$, where macrophages are labelled by Turquoise2 (26). Embryos were raised in E3 media supplemented with methylene blue in a dark incubator at 28°C until 1 day post fertilisation (dpf). Cleaned 1 dpf embryos were transferred to E3 without methylene blue and reared (dark incubator, 28°C) until used.

Zebrafish larvae were anaesthetised in 250 µg/mL MS-222 (Tricaine, Sigma) and euthanised either by fixation in 4% paraformaldehyde, lysis in Trizol, or disposed of in bleach as per biosafety protocols.

### Preparation of CSE

Cigarette smoke extract (CSE) was prepared by using a custom-built smoking device where reference cigarette 3R4F (Kentucky Tobacco Research & Development Centre) was lit using a lighter and CS was bubbled directly in to 10 mL of PBS until the whole cigarette was exhausted. This 10 mL of CSE was filtered using a 0.22-micron filter to remove CS-derived debris. This preparation of the smoke of one cigarette per 10 mL PBS is considered 100% CSE and was diluted as indicated. CSE was prepared fresh before treatment.

### Exposure of zebrafish to CSE

89    Embryos/larvae were exposed to CSE at a density of approximately 1 larva per mL of E3 in plastic Petri
90    dishes at the ages indicated, either 3 or 5 dpf, and maintained in a dark incubator at 28°C. The stock
91    solution of 100% CSE in PBS was diluted directly into E3 zebrafish media to a final concentration of 1%, 2%,
92    or 3%. Filter sterilised PBS was used as vehicle control.
93
<u>Zebrafish behavioural analysis</u>
Zebrafish larvae were transferred to a 24-well plate with 1 embryo per well in their CSE-supplemented E3
media. The plate was transferred to a ZebraBox (Viewpoint) and locomotion was recorded for 15 minutes.
Video files were analysed using EthoVision XT 11 (Noldus) using the following parameters: sensitivity 110,
video pixel smoothing low, subject size 4-4065. Locomotion analysis was calculated as the total distance
travelled (mm) of individual larva over the recording window.
100
101   <u>Tail transection assays</u>
102   Caudal fin amputations were performed on 5 dpf larvae. Larvae were anesthetised with 2.5% (v/v) ethyl-3-
103   aminobenzoate methanesulfonate (tricaine) (Sigma, E10521), wounded posterior to the notochord using a
104   sterile scalpel and kept in a 28°C incubator to recover as previously described (27). Wounded larvae were
105   imaged at 6 hours post wounding.
106
107   <u>Imaging and image analysis</u>
108   Larvae were imaged using a Leica M205FA fluorescent microscope. ImageJ software was used to quantify
109   fluorescent pixel counts as previously described (27).
110
<u>Generation of microbiome-depleted zebrafish embryos</u>
Microbiome-depleted zebrafish were created and maintained as previously described (28). Briefly, freshly
laid embryos were rinsed with 0.003% v/v bleach in sterile E3 and rinsed 3 times with sterile E3. Bleached
embryos were raised in sterile E3 supplemented with 50 µg/mL ampicillin (Sigma), 5 µg/mL kanamycin
(Sigma) and 250 ng/mL amphotericin B (Sigma) in sterile tissue culture flasks. Dead embryos and chorions
were aseptically removed at 1 and 3 dpf, respectively.
117
<u>Generation of mouse faecal microbiota specimens</u>
Mice were housed at the Centenary Institute (Sydney Local Health District Animal Welfare Committee
Approval 2020-003). C57BL/6J mice were housed in a pathogen-free and temperature-controlled
environment, with 12 hours of light and 12 hours of darkness, and free access to food and water. Mice
were smoked with 12x 3R4F cigarettes for 75 minutes, twice a day, 5 days per week for 10 weeks by using a
custom-designed and purpose-built nose-only, directed flow inhalation and smoke-exposure system (CH
Technologies, NJ) (10, 29). This is representative of a pack-a-day smoker (2).
125
126   Faecal pellets were collected from mice that were housed in different cages. Individual faecal pellets were
127   collected into sterile 1.7 mL microcentrifuge tubes and homogenised in 1 mL of sterile E3 by pipetting.
128   Homogenised specimens were centrifuged (500 x$g$, 2 minutes) to sediment fibrous material and
129   supernatant were collected then pooled. Pooled supernatants were supplemented with glycerol to a final
130   concentration of 25% v/v, aliquoted, and frozen at -80°C before use.
131
132   Mice were not anaesthetised for these procedures and were used for other experiments as per Approval
133   2020-003 where they were eventually sacrificed by CO2 asphyxiation.
134
135   <u>Exposure of zebrafish larvae to mouse faecal microbiota</u>
136   Microbiome-depleted zebrafish were rinsed at 3 dpf with sterile E3, without antibiotic and amphotericin B
137   supplementation, and colonised by the addition of 200 µL thawed faecal homogenate supernatant into 30
138   mL sterile E3 delivering a final concentration of ~5x10$^4$ CFU per mL. Enumeration of culturable bacterial
139   load was carried out by culture under aerobic conditions at 28°C on LB agar (Thermofisher).
140
<u>Detection of gene expression</u>

Groups of at least 10 larvae were homogenised in Trizol (Thermofisher) and processed for RNA extraction
as per manufacturer's instructions. cDNA was reverse transcribed using the Applied Biosystems High
Capacity cDNA synthesis kit (Thermofisher) from 2 µg of total RNA. Quantitative PCR was carried out on a
CFX Connect machine (Biorad) with Powerup SYBR reagent (Thermofisher), primer sets were (5'-3'): *il1b* Fw
ATCAAACCCCAATCCACAGAGT, Rv GGCACTGAAGACACCACGTT; *18s* Fw TCGCTAGTTGGCATCGTTTATG, Rv
CGGAGGTTCGAAGACGATCA.
148
149 Statistical analysis
150 Statistical testing was carried out by ANOVA with post hoc multiple comparisons as appropriate using
151 GraphPad Prism 9 (GraphPad). Data are expressed as mean ± SD, calculated P values are provided in
152 graphs. Outliers were excluded using a 1% robust regression and outlier removal (ROUT) test. Every
153 datapoint represents a single embryo unless otherwise noted as for qPCR analyses.
154
**Results**
Chronic exposure to CSE is toxic to zebrafish post-hatching larvae
To establish a survivable dose of CSE that did not cause gross pathology, we exposed 3 dpf zebrafish larvae
to a range of CSE doses and monitored survival. We found exposure at up to 2% CSE was tolerated by
zebrafish larvae (Figure 1A), with no overt changes to morphology (Figure 1B). Exposure to 3% CSE caused
significant mortality within 1 day post exposure.
161
162 To verify that the CSE was bioactive at 1% and 2% concentrations, we tracked larval motility to detect
163 nicotine-induced increases in motility at 5 dpf, or 2 days post exposure (30). Exposure to 1% or 2% CSE
164 increased the total distance travelled of zebrafish larvae compared to mock exposed larvae in a 15 minute
165 recording period indicating components of the CSE were still bioactive after 2 days of immersion exposure
166 in E3 media (Figure 1C).
167
We next used *Tg(lyzC:DsRed)*[nz50] larvae to visualise neutrophils and *Tg(mfap4:turquoise)*[xt27] to visualise
macrophages (25, 26). Immune cell fluorescent area was used as a proxy to estimate changes to immune
cell number by non-invasive live imaging. We performed tail transection on 5 dpf larvae that had been
exposed to CSE from 3 dpf and assessed leukocyte recruitment to the wound at 6 hours post wounding.
While neutrophil recruitment was slightly but statistically significantly reduced by 1% CSE, there was no
effect of 2% CSE exposure compared to vehicle control and overlap between neutrophil recruitment in the
1% and 2% CSE exposure groups (Figure 1D). There was significantly more macrophage recruitment to
wounds in 2% CSE-exposed larvae compared to control and 1% CSE larvae (Figure 1E).
176
Acute exposure to CSE induces myelopoiesis in zebrafish larvae
To examine the acute effects of CSE exposure on larvae after major organogenesis is complete, we exposed
5 dpf larvae to CSE for 24 hours. Immersion in 1% and 2% CSE caused a gradated increase in total
neutrophil fluorescent area in zebrafish larvae compared to PBS vehicle (Figure 2A and 2B). This
inflammatory effect of CSE appeared stronger on total macrophage fluorescent area with a statistically
significant increase in total macrophages in 2% CSE-exposed compared to control larvae (Figure 2C and 2D).
183
Acute exposure to CSE induces an inflammatory response in the gut of zebrafish larvae
The effects that we observed suggested that CSE induced an inflammatory haematopoietic response. Since
the larval midgut and hindgut is an important site of immune response to chemical irritants, we next
quantified leukocyte recruitment to the larval intestine of 6 dpf larvae that had been exposed to 2% CSE
from 5 dpf (31). Leukocyte fluorescence was increased in the gut compared to either control or 1% CSE-
exposed larvae (Figure 3A and 3B). However, the estimation of gut macrophages by fluorescent pixel count
in our *Tg(mfap4:turquoise)*[xt27] larvae was potentially confounded by luminal autofluorescence that was
more noticeable in CSE-exposed larvae (Figure 2B).
192
The enhanced effects of CSE on innate immune cell numbers in the larval gut led us to hypothesise that the
microbiota may interact with CSE to drive inflammation at zebrafish larval mucosal surfaces as observed
with other chemical irritants (31, 32). Thus, we next examined the effect of CSE and the role of changes in

the endogenous microbiota by depleting bacteria with broad spectrum antibiotics prior to CSE exposure
from 5 to 6 dpf. Antibiotic treatment did not affect the ability of 2% CSE to increase neutrophil or
macrophage fluorescence in the larval gut suggesting that the natural gut microbiota is not necessary to
drive the acute inflammatory effects of 2% CSE (Figure 3A and 3B).
200
<u>Microbiota from CS-exposed mice increases the neutrophilic inflammatory effects of CSE</u>
We next assessed if the microbiota of chronically CS-exposed mice could impart increases in sensitivity to
CSE exposure by exposing 3 dpf microbiota-depleted larval zebrafish to faecal homogenate from mice
exposed to CS/air controls for 10 weeks and CSE. We used the suboptimal concentration of 1% CSE, which
did not induce intestinal inflammation in our acute exposure assay, to detect enhancement of inflammation
by microbiota derived from the faeces of CS-exposed mice.
207
208 The total count of neutrophil fluorescence was unaffected in 5 dpf larvae immersed with microbiota from
209 CS- or normal air-exposed control mice, and the addition of 1% CSE (Figure 4A and 4B).
210
Exposure to microbiota from CS-exposed or control mice did not affect neutrophil fluorescence in the gut of
wild-type 5 dpf larvae compared to germ-free reared larvae (Figure 4C and 4D). Addition of 1% CSE did not
affect neutrophil fluorescence in the gut in 5 dpf larvae raised microbiota-depleted or in those exposed to
microbiota from control non-smoked mice. However, 5 dpf larvae that had been exposed to microbiota
from smoked mice and 1% CSE had significantly more neutrophil fluorescence in the gut than any of the
control groups. This pattern was replicated by *il1b* transcription levels quantified by RT-qPCR (Figure 4E).
217
<u>CS-exposed mouse microbiota increases the macrophage inflammatory effects of CSE</u>
Analysis of *Tg(mfap4:turquoise)*[xt27] macrophage reporter larvae revealed increased total macrophage
fluorescence in 5 dpf larvae that that had been exposed to microbiota from CS-exposed mice and 1% CSE
compared to any of the control groups, except for the air-exposed mouse microbiota and 1% CSE control
group (Figure 5A and 5B). Quantification of gut macrophage fluorescence was again confounded by luminal
autofluorescence in CSE-exposed larvae, appearing to be elevated in all CSE-exposed groups (Figure 5C and
5D).
225
**CS-exposed faecal microbiota contains culturable bacteria that do not affect inflammatory responses**
To investigate if any individual species of bacteria from CS-exposed mouse microbiota were able to
recapitulate the effects of exposure to the bulk microbiota homogenate, we recovered bacteria from the
homogenised CS-exposed mouse microbiota-exposed larvae onto rich LB agar. Four species were
subcultured to purity by passage on LB agar and genotyped by 16s sequencing as *Acinetobacter*
*radioresistens*, *Bacteroides vulgatus*, *Enterobacter cancerogenus*, and *Stenotrophomonas maltophilia*
(Supplementary File). We then screened these individual bacteria by colonising 3 dpf microbiota-depleted
larvae in combination with 1% CSE exposure and found no effect on total or gut leukocyte fluorescent area
(Figure 6).
235
**Discussion**
Here we found the immune system of zebrafish larvae is sensitive to the pro-inflammatory neutrophilic and
macrophage effects of CSE. Exposing zebrafish larvae to microbiota derived from the faeces of CS-exposed
mice demonstrate the presence of microbes (including culturable *Acinetobacter radioresistens*, *Bacteroides*
*vulgatus*, *Enterobacter cancerogenus*, and *Stenotrophomonas maltophilia*) or metabolites capable of
amplifying the inflammatory effects of CSE across species. Together, these data establish zebrafish larvae as
a platform for studying the environmental interactions that affect cigarette- and tobacco-induced
inflammatory damage.
244
Our finding that CSE concentrations of >2% are toxic to zebrafish larvae are in line with previous studies by
Alvarez *et al.* who showed larvae can survive in a solution of approximately 1% CSE and Progatzky *et al.*
who utilised a 1% CSE challenge with adult zebrafish (19, 21). Exposure of adult zebrafish to CSE by Alvarez
*et al.* demonstrated an impairment in post-amputation fin regeneration (19). Our acute wound leukocyte

response data adds to this observation by suggesting that increased numbers of wound-associated
macrophages may impede physiological regeneration in the presence of CSE.
251
Our data demonstrate that exposure of zebrafish larvae to 2% CSE causes an expansion of leukocytes
(neutrophils and macrophages) and leukocytic inflammation in the gut. Despite the increased leukocyte
fluorescent area, we did not observe the mass redistribution of leukocytes to the skin mucosal surfaces as
we have previously documented with the enterocolitis-inducing chemical irritants dextran sodium sulfate
and 2,4,6-trinitrobenzene sulfonic acid (31, 32). Exposure of zebrafish larvae to a dose of 1% CSE did not
cause detectable changes to inflammatory markers in conventionally reared larvae. Our results are
comparable to work using adult zebrafish that found only a subtle inflammatory response to 1% CSE
exposure in the gill tissue of adult zebrafish with upregulation of inflammatory gene expression but
decreased or unchanged immune cell infiltration in the gills depending on the length of exposure (21).
261
Our system of using a sub-optimal dose of 1% CSE for two days to stimulate a below measurable amount of
inflammation proved sensitive to positive modulation by microbiota samples from CS-exposed mouse
faeces. Zebrafish have been used to as hosts for trans-species microbiota transplants (33, 34), and our
study expands this knowledge by demonstrating sensitivity of recipient zebrafish host inflammation to a
microbiota-environment interaction. These proof-of-concept data suggest that zebrafish larvae may be a
feasible platform with which human microbiota-CS interactions can be screened to uncover potential
missing links in microbiota-environment interactions that appear to predispose some smokers to the
development of COPD (4, 9, 11).
270
The failure of single bacterial species that could be cultured from the guts of zebrafish to recapitulate the
pro-inflammatory effects of CS-exposed mouse microbiota in the context of 1% CSE co-exposure suggest
either the presence of multiple microbial species and/or other microbiota or factors present in the faecal
homogenate. This was not unexpected as the aerobic conditions of the larval zebrafish gut does not
facilitate sustained colonisation by some members of the mammalian microbiota (33), and our method of
recovering microbes on LB agar in atmospheric conditions will further bias recovery of only species present
at a high proportion in the gut of colonised larvae.
278
Recent mouse studies have demonstrated the importance of the metabolome, including short chain fatty
acid output, in susceptibility to smoking-induced pathology (15, 16, 35). Given the lack of butyrate
production found in conventionally raised zebrafish larvae (27), we think it is unlikely that exposure to
control mouse faecal microbiota produces anti-inflammatory short chain fatty acids as the protective
mechanism against 1% CSE exposure compared to the pro-inflammatory CS-exposed mouse faecal
microbiota. Instead, we propose that the presence of additional toxins or pro-inflammatory factors either
already present or produced by complex mixtures of microbes harvested from the supernatant of faecal
homogenates drives inflammatory responses.
287
The dose response effects of CSE exposure on inflammation in larval zebrafish suggest this platform could
be adapted for use as a screening platform to identify positive and negative regulators of inflammation
(36). In this study we have demonstrated the utility of a low CSE challenge to identify CS-associated
microbiota as factors that increase the susceptibility of zebrafish larvae to CSE-induced inflammation.
Follow-up studies would identify other soluble environmental factors that increase the inflammatory
potential of CSE. Conversely, a higher dose of 2% CSE challenge could be used as a platform to screen for
inhibitors of CSE-induced inflammation. Since we have recently demonstrated the sensitivity of zebrafish
larval inflammation to the short chain fatty acid butyrate (27), it would be interesting to determine if high
dose CSE-induced inflammation in zebrafish larvae can also be ameliorated by butyrate supplementation.
297
298  In summary, we have demonstrated the suitability of zebrafish larvae for studying the inflammatory effects
299  of cigarette smoke-environment interactions and show that the CS-exposed mouse gut microbiota
300  potentiates the inflammatory effects of CSE across host species.
301
302  **DATA AVAILABILITY**

303	Raw data is available upon reasonable request to the corresponding authors.
304

305	**ACKNOWLEDGEMENTS**
306	We thank other members of the Oehlers lab at the Centenary Institute for their input into this work and
307	maintenance of the zebrafish aquarium; Dr Angela Fontaine and Sydney Cytometry for assistance with
308	imaging; Dr Diana Quan and Ms Trixie Wang for collection of mouse faeces; Dr Kamal Dua for supervision of
309	VM.
310

**FUNDING**
This work was supported by a Centenary Institute Summer Scholarship to S.M.; University of Sydney
Fellowship [grant number G197581] and NSW Ministry of Health under the NSW Health Early-Mid Career
Fellowships Scheme [grant number H18/31086] to S.H.O.; the NHMRC Centre of Research Excellence in
Tuberculosis Control (APP1153493) to W.J.B.; NHMRC Fellowship (APP1175134) UTS, Cancer Council NSW
(1157073), and the Rainbow Foundation to P.M.H.; Graduate School of Health and University of Technology
Sydney (UTS President's scholarship and UTS International Research Scholarship) to V.M..

318
319

320	**References**
321

1.	WHO. Tobacco 2021 [Available from: https://www.who.int/news-room/fact-sheets/detail/tobacco.
2.	Jones B, Donovan C, Liu G, Gomez HM, Chimankar V, Harrison CL, et al. Animal models of COPD:
What do they tell us? Respirology. 2017;22(1):21-32.
3.	Keely S, Talley NJ, Hansbro PM. Pulmonary-intestinal cross-talk in mucosal inflammatory disease.
Mucosal Immunol. 2012;5(1):7-18.
4.	Budden KF, Gellatly SL, Wood DL, Cooper MA, Morrison M, Hugenholtz P, et al. Emerging
pathogenic links between microbiota and the gut-lung axis. Nat Rev Microbiol. 2017;15(1):55-63.
5.	Verschuere S, Bracke KR, Demoor T, Plantinga M, Verbrugghe P, Ferdinande L, et al. Cigarette
smoking alters epithelial apoptosis and immune composition in murine GALT. Laboratory Investigation.
2011;91(7):1056-67.
6.	Allais L, Kerckhof FM, Verschuere S, Bracke KR, De Smet R, Laukens D, et al. Chronic cigarette
smoke exposure induces microbial and inflammatory shifts and mucin changes in the murine gut.
Environmental microbiology. 2016;18(5):1352-63.
7.	Wronikowska O, Michalak A, Skalicka-Woźniak K, Crawford AD, Budzyńska B. Fishing for a deeper
understanding of nicotine effects using zebrafish behavioural models. Progress in Neuro-
Psychopharmacology and Biological Psychiatry. 2020;98:109826.
8.	Hoyt ATM, Canfield MAPD, Romitti PAPD, Botto LDMD, Anderka MTSD, Krikov SVM, et al.
Associations between Maternal Periconceptional Exposure to Secondhand Tobacco Smoke and Major Birth
Defects. American journal of obstetrics and gynecology. 2016;215(5):613.e1-.e11.
9.	Budden KF, Shukla SD, Rehman SF, Bowerman KL, Keely S, Hugenholtz P, et al. Functional effects of
the microbiota in chronic respiratory disease. Lancet Respir Med. 2019;7(10):907-20.
10.	Fricker M, Goggins BJ, Mateer S, Jones B, Kim RY, Gellatly SL, et al. Chronic cigarette smoke
exposure induces systemic hypoxia that drives intestinal dysfunction. JCI Insight. 2018;3(3).
11.	Bowerman KL, Rehman SF, Vaughan A, Lachner N, Budden KF, Kim RY, et al. Disease-associated gut
microbiome and metabolome changes in patients with chronic obstructive pulmonary disease. Nat
Commun. 2020;11(1):5886.
12.	Malyla V, Paudel KR, Shukla SD, Donovan C, Wadhwa R, Pickles S, et al. Recent advances in
experimental animal models of lung cancer. Future Med Chem. 2020;12(7):567-70.
13.	Shanahan ER, Shah A, Koloski N, Walker MM, Talley NJ, Morrison M, et al. Influence of cigarette
smoking on the human duodenal mucosa-associated microbiota. Microbiome. 2018;6(1):150.
14.	Pavia CS, Pierre A, Nowakowski J. Journal of medical microbiology. 2000;49(7):675-6.
15.	Jang YO, Kim OH, Kim SJ, Lee SH, Yun S, Lim SE, et al. High-fiber diets attenuate emphysema
development via modulation of gut microbiota and metabolism. Scientific reports. 2021;11(1):7008.

16.      Jang YO, Lee SH, Choi JJ, Kim DH, Choi JM, Kang MJ, et al. Fecal microbial transplantation and a high
fiber diet attenuates emphysema development by suppressing inflammation and apoptosis. Exp Mol Med.
2020;52(7):1128-39.
17.      Gosens R, Hiemstra PS, Adcock IM, Bracke KR, Dickson RP, Hansbro PM, et al. Host-microbe cross-
talk in the lung microenvironment: implications for understanding and treating chronic lung disease. Eur
Respir J. 2020;56(2).
18.      Ellis LD, Soo EC, Achenbach JC, Morash MG, Soanes KH. Use of the zebrafish larvae as a model to
study cigarette smoke condensate toxicity. PLoS ONE. 2014;9(12):e115305.
19.      Alvarez M, Chávez MN, Miranda M, Aedo G, Allende ML, Egaña JT. A Novel In Vivo Model to Study
Impaired Tissue Regeneration Mediated by Cigarette Smoke. Scientific reports. 2018;8(1):10926.
20.      Palpant NJ, Hofsteen P, Pabon L, Reinecke H, Murry CE. Cardiac Development in Zebrafish and
Human Embryonic Stem Cells Is Inhibited by Exposure to Tobacco Cigarettes and E-Cigarettes. PLoS ONE.
2015;10(5):e0126259.
21.      Progatzky F, Cook H, Lamb J, Bugeon L, Dallman M. Mucosal inflammation at the respiratory
interface: A zebrafish model. American journal of physiology Lung cellular and molecular physiology.
2015;310:ajplung.00323.2015.
22.      Massarsky A, Prasad GL, Di Giulio RT. Total particulate matter from cigarette smoke disrupts
vascular development in zebrafish brain (Danio rerio). Toxicology and applied pharmacology. 2018;339:85-
96.
23.      Massarsky A, Jayasundara N, Glazer L, Levin ED, Prasad GL, Di Giulio RT. Outcomes of
developmental exposure to total particulate matter from cigarette smoke in zebrafish (Danio rerio).
Neurotoxicology. 2018;68:101-14.
24.      Renshaw SA, Trede NS. A model 450 million years in the making: zebrafish and vertebrate
immunity. Disease models & mechanisms. 2012;5(1):38-47.
25.      Hall C, Flores MV, Storm T, Crosier K, Crosier P. The zebrafish lysozyme C promoter drives myeloid-
specific expression in transgenic fish. BMC Dev Biol. 2007;7(1):42.
26.      Walton EM, Cronan MR, Beerman RW, Tobin DM. The Macrophage-Specific Promoter mfap4 Allows
Live, Long-Term Analysis of Macrophage Behavior during Mycobacterial Infection in Zebrafish. PLoS ONE.
2015;10(10):e0138949.
27.      Cholan PM, Han A, Woodie BR, Watchon M, Kurz AR, Laird AS, et al. Conserved anti-inflammatory
effects and sensing of butyrate in zebrafish. Gut Microbes. 2020;12(1):1-11.
28.      Melancon E, Gomez De La Torre Canny S, Sichel S, Kelly M, Wiles TJ, Rawls JF, et al. Best practices
for germ-free derivation and gnotobiotic zebrafish husbandry. Methods in cell biology. 2017;138:61-100.
29.      Beckett EL, Stevens RL, Jarnicki AG, Kim RY, Hanish I, Hansbro NG, et al. A new short-term mouse
model of chronic obstructive pulmonary disease identifies a role for mast cell tryptase in pathogenesis. J
Allergy Clin Immunol. 2013;131(3):752-62.
30.      Petzold AM, Balciunas D, Sivasubbu S, Clark KJ, Bedell VM, Westcot SE, et al. Nicotine response
genetics in the zebrafish. Proc Natl Acad Sci U S A. 2009;106(44):18662-7.
31.      Oehlers SH, Flores MV, Okuda KS, Hall CJ, Crosier KE, Crosier PS. A chemical enterocolitis model in
zebrafish larvae that is dependent on microbiota and responsive to pharmacological agents. Dev Dyn.
2011;240(1):288-98.
32.      Oehlers SH, Flores MV, Hall CJ, Crosier KE, Crosier PS. Retinoic acid suppresses intestinal mucus
production and exacerbates experimental enterocolitis. Dis Model Mech. 2012;5(4):457-67.
33.      Rawls JF, Mahowald MA, Ley RE, Gordon JI. Reciprocal gut microbiota transplants from zebrafish
and mice to germ-free recipients reveal host habitat selection. Cell. 2006;127(2):423-33.
34.      Cholan PM, Morris S, Luo K, Chen J, Boland JA, McCaughan GW, et al. Transplantation of high fat
fed mouse microbiota into zebrafish embryos identifies pathobiont species. bioRxiv.
2021:2021.04.28.441900.
35.      Rutting S, Xenaki D, Malouf M, Horvat JC, Wood LG, Hansbro PM, et al. Short-chain fatty acids
increase TNFalpha-induced inflammation in primary human lung mesenchymal cells through the activation
of p38 MAPK. American journal of physiology Lung cellular and molecular physiology. 2019;316(1):L157-
L74.
36.      Oehlers SH, Flores MV, Hall CJ, Wang L, Ko DC, Crosier KE, et al. A whole animal chemical screen
approach to identify modifiers of intestinal neutrophilic inflammation. FEBS J. 2017;284(3):402-13.

409
410

**FIGURE LEGENDS**

**Figure 1: Cigarette smoke extract is toxic to zebrafish larvae, and causes increased embryo motility, and greater macrophage influx into wounds**

A. Survival of 3 dpf larvae after exposure to CSE. Individual Log-rank comparison scores: P(0% vs 1%) = 0.3173, P(0% vs 2%) = 0.0798, P(0% vs 3%) <0.0001, P(1 vs 3) <0.0001, P(2% vs 3%) <0.0001. Starting n per group: 0 = 45 larvae, 1% = 45 larvae, 2% = 45 larvae, 3% = 36 larvae.

B. Representative images of 5 dpf larvae exposed to CSE from 3 dpf. Scale bar = 1 mm.

C. Total distance moved in 15 minutes by 5 dpf larvae exposed to CSE from 3 dpf. Statistical comparison by ANOVA, data are representative of two biological replicates.

D. Quantification of neutrophil fluorescent pixel area within 100 µm of the wound margin at 6 hpw in 5 dpf larvae exposed to CSE from 3 dpf. Statistical comparison by ANOVA, data are normalised and combined from two biological replicates.

E. Quantification of macrophage fluorescent pixel area within 100 µm of the wound margin at 6 hpw in 5 dpf larvae exposed to CSE at doses indicated from 3 dpf. Statistical comparison by ANOVA, data are normalised and combined from two biological replicates.

**Figure 2: Acute exposure to CSE induces myelopoiesis in zebrafish larvae**

A. Representative images of 6 dpf *Tg(lyzC:DsRed)*[nz50] larvae exposed to CSE for 24 hours from 5 dpf.

B. Quantification of total neutrophil fluorescent pixel area in 6 dpf larva exposed to CSE for 24 hours from 5 dpf.

C. Representative images of 6 dpf *Tg(mfap4:turquoise)*[xt27] larvae exposed to CSE for 24 hours from 5 dpf.

D. Quantification of total macrophage fluorescent pixel area in 6 dpf larva exposed to CSE for 24 hours from 5 dpf.

A & C: Scale bars = 1 mm. Bounded area in insets are used to measure gut leukocytes in Figure 3 onwards.

B & D: Statistical comparison by ANOVA, data are representative of two biological replicates.

**Figure 3 Acute exposure to CSE induces an inflammatory response in the gut of zebrafish larvae**

A. Quantification of gut neutrophil fluorescent pixel area in 6 dpf larvae treated with broad spectrum antibiotics (Abx) ampicillin and kanamycin and exposed to CSE from 5 dpf.

B. Quantification of gut macrophage fluorescent pixel area in 6 dpf larvae treated with broad spectrum antibiotics (Abx) ampicillin and kanamycin and exposed to CSE from 5 dpf.

Statistical comparisons by ANOVA, data are representative of two biological replicates.

**Figure 4 CS-exposed mouse microbiota increases the neutrophilic inflammatory effects of CSE**

A. Representative images of 5 dpf microbiota-depleted *Tg(lyzC:DsRed)*[nz50] larvae exposed to combinations of CS- or control normal air-exposed mouse microbiota with or without 1% CSE from 3 dpf.

B. Quantification of total neutrophil fluorescent pixel area per larva exposed to combinations of CS- or control normal air-exposed mouse microbiota with or without 1% CSE from 3 dpf. Statistical comparison by ANOVA, data are representative of two biological replicates.

C. Representative images of the posterior guts of microbiome depleted *Tg(lyzC:DsRed)*[nz50] larvae exposed to combinations of CS- or control normal air-exposed mouse microbiota with or without 1% CSE from 3 dpf. Area used for analysis demarcated by yellow box.

D. Quantification of gut neutrophil fluorescent pixel area per larva exposed to combinations of CS- or control normal air-exposed mouse microbiota with or without 1% CSE from 3 dpf. Statistical comparison by ANOVA, data are representative of two biological replicates.

E. Quantification of *il1b* transcripts by qPCR in larvae exposed to combinations of CS- or control normal air-exposed mouse microbiota with or without 1% CSE from 3 dpf. Statistical comparison by ANOVA, each data point represents a biological replicate with at least 10 larvae per group.

Scale bars = 1 mm.

**Figure 5 CS-exposed mouse microbiota increases the macrophage inflammatory effects of CSE**

A. Representative images of 5 dpf microbiota-depleted *Tg(mfap4:turquoise)*[xt27] larvae exposed to combinations of CS- or control normal air-exposed mouse microbiota with or without 1% CSE from 3 dpf.

B. Quantification of total macrophage fluorescent pixel area per larva exposed to combinations of CS- or
control normal air-exposed mouse microbiota with or without 1% CSE. Statistical comparison by ANOVA,
data are representative of two biological replicates.
C. Representative images of the posterior guts of microbiome depleted *Tg(mfap4:turquoise)$^{xt27}$* larvae
exposed to combinations of CS- or control normal air-exposed mouse microbiota with or without 1% CSE.
Area used for analysis demarcated by yellow box.
D. Quantification of gut macrophage fluorescent pixel area per larva exposed to combinations of CS- or
control normal air-exposed mouse microbiota with or without 1% CSE. Statistical comparison by ANOVA,
data are representative of two biological replicates.
Scale bars indicate 1 mm.
474
**Figure 6 CS-exposed faecal microbiota contain culturable bacteria that do not affect inflammatory**
**responses**
A. Quantification of total neutrophil fluorescent pixel area per larva exposed to bacterial isolates with or
without 1% CSE.
B. Quantification of gut neutrophil fluorescent pixel area per larva exposed to bacterial isolates with or
without 1% CSE.
C. Quantification of total macrophage fluorescent pixel area per larva exposed to bacterial isolates with or
without 1% CSE.
D. Quantification of gut macrophage fluorescent pixel area per larva exposed to bacterial isolates with or
without 1% CSE. Statistical comparisons by ANOVA.
485
486  **Supplementary File: Identification of isolated bacterial strains**
487  Top NCBI BLAST hit strain information and sequencing data used for BLAST.
488

**A**

Microbiome depleted

Microbiome depleted + 1% CSE

Control mouse microbiota

Control mouse microbiota + 1% CSE

Smoked mouse microbiota

Smoked mouse microbiota + 1% CSE

**B**

Neutrophil fluorescent pixel count (whole body)

0.9918
0.9594
0.7063
0.7394
0.9797

| | | | | | |
|---|---|---|---|---|---|
| 1% CSE | − | + | − | + | − | + |
| Control mouse microbiota | − | − | + | + | − | − |
| Smoked mouse microbiota | − | − | − | − | + | + |

**C**

+ 1% CSE

Microbiome depleted

Control mouse microbiota

Smoked mouse microbiota

**D**

Neutrophil fluorescent pixel count (gut)

0.0005
0.0025
0.0137
0.0060
0.0293

| | | | | | |
|---|---|---|---|---|---|
| 1% CSE | − | + | − | + | − | + |
| Control mouse microbiota | − | − | + | + | − | − |
| Smoked mouse microbiota | − | − | − | − | + | + |

**E**

Relative quantity of *il1b* transcripts

0.0016
0.0049
0.0016
0.0014
0.0043

| | | | | | |
|---|---|---|---|---|---|
| 1% CSE | − | + | − | + | − | + |
| Control mouse microbiota | − | − | + | + | − | − |
| Smoked mouse microbiota | − | − | − | − | + | + |