Title: Using all gene families vastly expands data available for phylogenomic inference in primates

Authors: Megan L. Smith[1], Dan Vanderpool[1], and Matthew W. Hahn[1]

[1]Department of Biology and Department of Computer Science, Indiana University, Bloomington, Indiana, USA

Corresponding Author: mls16@indiana.edu

Abstract

Traditionally, single-copy orthologs have been the gold standard in phylogenomics. Most

phylogenomic studies identify putative single-copy orthologs by using clustering approaches and

retaining families with a single sequence from each species. However, this approach can severely

limit the amount of data available by excluding larger families. Recent methodological advances

have suggested several ways to include data from larger families. For instance, tree-based

decomposition methods facilitate the extraction of orthologs from large families. Additionally,

several popular methods for species tree inference appear to be robust to the inclusion of

paralogs, and hence could use all of the data from larger families. Here, we explore the effects of

using all families for phylogenetic inference using genomes from 26 primate species. We

compare single-copy families, orthologs extracted using tree-based decomposition approaches,

and all families with all data (i.e., including orthologs and paralogs). We explore several species

tree inference methods, finding that across all nodes of the tree except one, identical trees are

returned across nearly all datasets and methods. As in previous studies, the relationships among

Platyrrhini remain contentious; however, the tree inference methods matter more than the dataset

used. We also assess the effects of each dataset on branch length estimates, measures of

phylogenetic uncertainty and concordance, and in detecting introgression. Our results

demonstrate that using data from larger gene families drastically increases the number of genes

available for phylogenetic inference and leads to consistent estimates of branch lengths, nodal

certainty and concordance, and inferences of introgression.

Keywords: phylogenetics, orthologs, paralogs, concatenation, coalescence

Introduction

Advances in sequencing technology have led to the availability of more genomic data than ever before, and the promise of phylogenomics is the application of this data to infer species relationships (Scornavacca et al. 2020). Essential to the application of genomic data to phylogenetic inference is the identification of homologous genes, or genes that share a common ancestor. Homologous genes may share a common ancestor due to speciation (orthologs) or duplication (paralogs). Since the terms ortholog and paralog were coined (Fitch 1970), orthologs have been considered the appropriate genes for phylogenetic inference because they are related only through speciation events, and therefore are thought to best reflect species relationships. Thus, identifying orthologs is a central part of most phylogenomic pipelines.

Nearly all pipelines for extracting putative orthologs from genomic data begin with a clustering step (Figure 1). Clustering approaches aim to identify sets of homologous genes. While the details vary, these approaches generally begin with pairwise comparisons of all sequences across genomes, identify putative pairwise homologs, and then use clustering approaches to attempt to group many sets of these genes together (reviewed in Altenhoff et al. 2019). The end-products of graph-based clustering approaches are clusters of orthologs and paralogs—i.e., gene families. Since most phylogenetic methods were designed for use with orthologs (and a single sequence per taxon), these groups must be further processed for downstream phylogenetic inference.

Three primary approaches have been used to process families for downstream inference (Figure 1; Step 1). The first and most common is to extract clusters with only a single copy in each species—these represent putative single-copy orthologs. Using single-copy families is generally seen as a conservative approach in phylogenomics, as these genes are likely to be orthologs; this choice also limits the amount of further downstream processing needed. However, the number of genes that are single-copy in all sampled species decreases sharply as additional species are included in the analyses (Emms and Kelly 2018), limiting the usefulness of this approach in many phylogenetic contexts.

In lieu of relying only on single-copy clusters, tree-based decomposition approaches (Willson et al. 2021) for orthology detection can be applied to extract orthologous genes from clusters that may have more than one copy in one or more species (Figure 1; Step 2). Tree-based decomposition approaches attempt to infer whether nodes in gene trees represent duplication or

speciation events, followed by the extraction of orthologs based on these node labels (reviewed

in Altenhoff et al. 2019). Early tree-based approaches relied on gene tree reconciliation to a

known species tree (e.g., Goodman et al. 1979), limiting their utility in cases where the species

tree is unknown or uncertain. However, recent approaches have relaxed these requirements. For

example, the method LOFT relies on a species overlap approach to identify duplication nodes in

gene trees (van der Heijden et al. 2007). Similarly, the software package Agalma (Dunn et al.

2013), the methods of Yang and Smith (2014), and DISCO (Willson et al. 2021) extract subtrees

without duplicates to generate sets of orthologs. While the exact implementations vary, in

general tree-based decomposition approaches aim to extract orthologous genes from families of

any size. Tree-based approaches allow researchers to vastly increase the number of genes

retained compared to using only the single-copy clusters. However, these approaches require that

users construct gene trees and perform ortholog extraction for each gene family, and they are

therefore more computationally intensive than relying on single-copy clusters alone (Figure 1).

Finally, families containing both orthologs and paralogs could be used for phylogenetic

inference. Although orthologs have traditionally been considered the appropriate genes for

phylogenetics, methods for estimating phylogenies from data including paralogs were introduced

more than forty years ago (Goodman et al. 1979; reviewed in Smith and Hahn 2021a). Recently,

several popular methods for species tree estimation have been shown to be robust to the presence

of paralogs (Hill et al. 2020; Legried et al. 2020; Markin and Eulenstein 2020; Yan et al. 2021).

Of particular interest, quartet-based methods, such as ASTRAL (Zhang et al. 2018), should be

robust to the inclusion of paralogs because the most common quartet is still expected to match

the species tree even in the presence of gene duplication and loss. Given that all ortholog

extraction methods may erroneously lead to the inclusion of paralogs, using methods that are

robust to their inclusion is likely a good strategy no matter the method employed to process the

output of clustering methods.

Though there have been several empirical comparisons between ortholog-detection

methods (e.g., Fernández et al. 2018; Kallal et al. 2018; Altenhoff et al. 2019), along with several

simulation-based (e.g., Legried et al. 2020; Zhang et al. 2020; Yan et al. 2021) and empirical

(e.g., Yan et al. 2021) studies evaluating the effects of paralog inclusion on phylogenetic

inference, several questions remain. First, a comparison of inference on single-copy clusters to

tree-based decomposition methods and methods that use all of the data (i.e., use orthologs and

paralogs for phylogenetic inference) would shed light on the advantages of the three approaches.

In addition, the joint effects of dataset, missing data requirements, and gene and species tree

inference method on species tree topology will provide information on the importance of each.

Finally, questions remain about the effects of the dataset used on branch length estimates,

measures of nodal support, and tests for introgression.

To address these questions, we reanalyze a recently published phylogenomic dataset that

includes 26 species of primates and 3 outgroups (Vanderpool et al. 2020). The data consist of

whole genomes from all 29 species. In the original study, Vanderpool et al. restricted inference

to 1,730 single-copy clusters present in 27 of the 29 studied species, a relatively small proportion

of the >20,000 genes available from each species; the species tree was inferred using

concatenated maximum likelihood, concatenated maximum parsimony, and quartet-based

approaches applied to gene trees inferred using both maximum likelihood and maximum

parsimony. The authors found robust relationships among all species except the Platyrrhini

("New World Monkeys"), for which inferences differed across species-tree and gene-tree

inference methods. In this paper, we compare inferences from three major types of dataset:

single-copy families, orthologs extracted from larger families using tree-based decomposition

approaches, and all families including all data (orthologs + paralogs). These datasets are then

compared in three different phylogenetic applications. First, we compare the species trees

inferred from these datasets using several methods, including concatenation-based and gene-tree-

based approaches. Second, we compare several measures of nodal support and nodal

consistency, as well as branch length estimates across datasets. Finally, we perform tests of

introgression and compare results across different datasets. Our results suggest minimal effects

of the dataset used on downstream phylogenetic inference, while highlighting the fact that both

tree-based decomposition approaches and approaches using both orthologs and paralogs greatly

expand the amount of data available.

Results

Using all gene families vastly expands the data available for phylogenetics

We compared three types of datasets produced by clustering approaches: single-copy

clusters, orthologs extracted from all clusters using tree-based decomposition approaches, and all

clusters (orthologs + paralogs) (Figure 1). For all datasets we considered both a stringent

missing-data threshold (only those genes present in at least 27 of the 29 sampled species;

MIN27) and a relaxed missing-data threshold (only those genes present in at least 4 of the 29

sampled species; MIN4). First, we considered only clusters that included a single gene from each

species (single-copy clusters; SCCs). While these genes are not guaranteed to be orthologs—due

to the potential inclusion of pseudoorthologs (Doolittle and Brown 1994; Koonin 2005)—this is

considered a safe approach and is often employed in phylogenomics. As expected, this dataset

included the fewest genes (Table 1).

Tree-based decomposition approaches aim to extract orthologous genes from any

cluster/family. We constructed gene trees for all clusters, and then used several tree-based

approaches to extract orthologous genes. First, we considered those clusters in which all

duplications were specific to a single lineage and kept a single gene-copy from this lineage.

When duplications are restricted to a single lineage, choosing one of the copies as the ortholog

cannot mislead phylogenetic inference regardless of which sequence is retained (see Figure 1d

from Smith and Hahn 2021a; Figure S1a). This dataset ("lineage-specific duplicates"; LSDs)

included more than 4X as many genes as the SCC dataset (Table 1). Next, we further expanded

our criteria to include those clusters with duplications specific to a pair of lineages ("two-species

duplicates"; TSDs; Figure S1b). Such duplications also cannot mislead topological inference,

though picking a non-orthologous pair could lead to longer branches. It is straightforward to pick

the most closely related pair of genes from the two species, which should not mislead either

topological or branch length inferences; including these genes further expanded the dataset

compared to the LSD dataset (Table 1).

We considered two tree-based decomposition approaches from Yang and Smith (Yang

and Smith 2014): minimum inclusion (MI) and monophyletic outgroups (MO). The MI approach

takes a gene tree and iteratively extracts subtrees with the highest number of taxa without taxon

duplication, until it cannot extract anymore subtrees with the minimum number of taxa. The MO

approach considers only those gene trees with a monophyletic outgroup, roots the tree, and infers

gene duplications from the root to the tips, pruning at nodes with duplications. These two

approaches were each applied to three datasets: the original gene trees, the original gene trees

trimmed to remove lineage-specific duplicates, and the original gene trees trimmed to remove

both lineage-specific and two-lineage duplicates. We explored the effects of additional filtering

and alternative parameters for the MI approach; as these changes had minimal effects, the results

are presented in the Supplementary Material (Appendix A). We also considered a new tree-based

decomposition approach: subtree extraction (SE). In this approach, we midpoint-root gene trees,

trimming away lineage-specific and two-lineage duplicates. We then extract subtrees that include

a single representative from each taxon (i.e., subtrees with no duplicates) and keep those trees

that meet minimum taxon-sampling thresholds (Figure S1c, d).

All tree-based approaches further expanded the amount of data available (Table 1). Since

the SE and MI approaches are highly similar (neither requires an outgroup, and both aim to

extract subtrees with no duplication events), we further examined the genes extracted using the

two approaches. We compared the MI dataset with TSDs trimmed and a minimum of 27 taxa to

the SE dataset with a minimum of 27 taxa sampled (this method trims TSDs internally). The

number of trees extracted using the two approaches was very similar (12,046 vs 12,198 genes in

the MI and SE datasets, respectively). For the 12,046 trees in the MI dataset, there was no analog

in the SE dataset for 2.4%, there was an identical tree in the SE dataset for 92.7%, and there was

a similar tree in the SE dataset for 4.8% (median Robinson-Foulds distance of these trees=2.0).

Thus, the MI and SE approaches extract very similar subsets of trees from the original clusters.

Finally, we considered two approaches that made no attempt to remove paralogs from the

dataset. We considered one dataset in which all orthologs and paralogs were included ("All

Paralogs"). This dataset was the most complete, as, even though it had fewer gene trees than

some tree-based approaches, the gene trees from these tree-based approaches are subtrees

extracted from this full dataset. This dataset cannot be analyzed using concatenation methods

because these approaches require an alignment that includes a single sequence for each species.

To address this, and to evaluate the effects of stochastic sampling of paralogs, we also included a

dataset in which a single gene (without regard to whether it was an ortholog or paralog) was

sampled at random from each species ("One Paralogs").

In total, we considered 20 datasets each with MIN4 and MIN27 taxon sampling. The

number of gene families ranged from 1,820 to 27,900 (Table 1). Clearly, considering only SCCs

drastically restricts the amount of data available, in terms of the number of gene trees (Table 1),

the number of decisive sites for each branch of the species tree (Figure 2A), and the number of

gene trees informative about each branch of the species tree (Figure 2B). All other datasets are

subsets of the All Paralogs dataset, and thus this dataset is necessarily the most informative.

Apart from the All Paralogs dataset, including a randomly sampled paralog (One Paralogs) leads

to the most decisive sites (Figure 2A), though they are not necessarily the most accurate sites

(see below and Figure 3). MI and SE lead to the most informative gene trees (Figure 2B).

Species tree inference is largely consistent across datasets

We inferred species trees using six approaches: ASTRAL (Sayyari and Mirarab 2016;

Zhang et al. 2018; Rabiee et al. 2019) on maximum likelihood (ML) gene trees, ASTRAL on

maximum parsimony (MP) gene trees, ASTRID (Vachaspati and Warnow 2015) on ML gene

trees, ASTRID on MP gene trees, concatenated ML inference in IQ-Tree (Nguyen et al. 2015),

concatenated MP inference in PAUP* (Swofford 2001), SVDQuartets (Chifman and Kubatko

2014), and ASTRAL-Pro (Zhang et al. 2020) on MP and ML gene trees. ML gene trees were

inferred in IQ-Tree, while MP gene trees were inferred in PAUP*. ASTRAL, ASTRID,

concatenated ML, and concatenated MP were all developed with orthologs in mind, but

ASTRAL has subsequently been demonstrated to be statistically consistent under models of gene

duplication and loss (Hill et al. 2020; Legried et al. 2020; Markin and Eulenstein 2020).

ASTRAL-Pro, on the other hand, was designed with paralogs in mind, and was only applied to

the All Paralogs datasets.

Across all nodes of the species tree, except for the relationships among the Platyrrhini

(discussed below), an identical phylogeny was recovered across all datasets and species tree

inference methods (Figure 3) with two exceptions. When concatenated MP or SVDQuartets was

used to infer a species tree from the One Paralogs dataset (MIN27), *Macaca fasciularis* was

recovered as sister to *M. nemestrina* rather than *M. mulatta*, as in all other datasets and previous

studies (e.g., Vanderpool et al. 2020), but bootstrap support for this relationship was low (55%)

in the SVDQuartets analysis. Additionally, when SVDQuartets was used to infer a species tree

from the One Paralogs (MIN4) dataset, *Mandrillus leucophaeus* was recovered as sister to a

clade containing *Cerocebus atys, Papio anubis,* and *Theropithecus gelada*, rather than sister to

Cerocebus atys as in other analyses and previous studies, but bootstrap support for this

relationship was also low (< 50%).

Branch support values were also highly similar across filtering methods. Local posterior

probabilities were 1.0 in ASTRAL for all datasets and for all nodes except the contentious node

in the Platyrrhini. All bootstrap support values in the concatenated ML analyses were 100, and

all bootstrap support values were 100 in the concatenated MP analyses except for in the One

Paralogs (MIN27) dataset, which also had topological issues among macaques as mentioned

above. Similarly, in all SVDQuartets analyses bootstrap values were 99 or 100 except in the

Platyrrhini and One Paralogs datasets.

In addition to branch support values, we calculated measures of genealogical

discordance: gene and site concordance factors (Minh et al. 2020). These analyses were carried

out for all datasets except All Paralogs, because it is not possible to calculate these statistics for

this dataset. For all datasets except the One Paralogs dataset, site and gene concordance factors

were highly similar across datasets (Figure 3A, B, C). Concordance in the One Paralogs dataset

was consistently lower, as would be expected from the random sampling of homologs. In some

cases, gene concordance factors were slightly lower for the SCC and MO dataset than for the

other datasets (Figure 3B); this seems to be due to more genes that fall into the 'paraphyly'

category (i.e., genes for which at least one of the reference clades for a particular branch is not

monophyletic), rather than for more genes supporting either of the two minor topologies. gCFs

and sCFs for the MIN4 datasets are shown in Supplemental Figure S2.

Resolution of the Platyrrhini radiation varies across species tree and gene tree inference

methods

As in Vanderpool et al. (2020), we found uncertainty around relationships among the Platyrrhini.

Concatenated ML analyses and gene-tree based analyses that relied on gene trees inferred using

ML preferred a symmetric tree, with *Saimairi boliviensis* and *Cebus capucinus imitator* as sister

species and *Callithrix jacchus and Aotus nancymaae* as sister species (topology 1 in Figure 4A).

However, concatenated MP and gene-tree based analyses that relied on gene trees inferred using

MP preferred an asymmetric topology, with *Saimiri boliviensis* and *Cebus capucinus imitator*

sister and *Aotus nancymaae* sister to these two (topology 2 in Figure 4A). Finally, SVDQuartets

preferred a third topology that placed *Callithrix* sister to *Saimairi boliviensis* and *Cebus*

capucinus (topology 3 in Figure 4A).

Gene and site concordance factors clarify these results. A slight majority of ML gene

trees prefer topology 1 (Figure 4B), a majority of MP gene trees prefer topology 2 (Figure 4B),

while slightly more sites support topology 2 than support topology 1 (Figure 4C). While the

results from SVDQuartets may seem counterintuitive at first, SVDQuartets relies on symmetry

between the two minor topologies to infer the third topology as the correct topology. Since there

are relatively equal numbers of sites supporting topologies 1 and 2, it is therefore expected that

SVDQuartets would prefer topology 3, even though fewer sites support this topology. Results for

the MIN4 dataset are similar and are shown in Figure S3.

To further investigate the causes of disagreement among these taxa, we focused on the

SCC dataset with MIN27 filtering to compare ML and MP gene trees. For each gene, we

recorded the ML and MP gene tree topology and the site concordance factor with respect to the

focal node, as well as various summary statistics about each locus (number of site patterns,

number of parsimony informative sites, tree length, etc.). The percentage of sites supporting the

best topology was highest when ML and MP gene trees agreed (Supplemental Figure S6a,c).

Additionally, there was more variance in sCFs within a gene (i.e., the number of sites supporting

each topology differed more) when ML and MP gene trees agreed (Supplemental Figure S6a,b).

This suggests that for genes with similar numbers of sites supporting multiple topologies, ML

and MP were more likely to infer conflicting gene trees. Notably, 17.6 percent of gene trees

supported Tree 1 under both ML and MP inference, while 18.8 percent of the gene trees

supported Tree 2 under both ML and MP inference.

Branch length estimates are largely consistent across datasets

We inferred branch lengths using two approaches. In general, our results suggest that all

methods that extract orthologs perform similarly and should lead to reliable estimates of branch

lengths. First, we estimated branch lengths in units of substitutions per site using concatenated

ML (i.e., site-based branch lengths). We expect that the inclusion of paralogs will lead to

overestimates of site-based branch lengths, since the divergence times of paralogs should pre-

date divergence times of orthologs. As expected, estimated site-based branch lengths for the One

Paralogs dataset are longer than those estimated for the SCC dataset (Figure 5a, b). For all other

MIN27 datasets, estimated site-based branch lengths were highly similar to those from the SCC

dataset (Figure 5 c, d). However, there are some inconsistencies with site-based branch lengths

for terminal branches (Figure 5d), and all site-based branch lengths are more variable for the

MIN4 datasets (Supplemental Figure S4).

We also inferred discordance-based branch lengths in coalescent units using ASTRAL

for the ML gene tree datasets. We expect that the inclusion of paralogs will lead to

underestimated discordance-based branch lengths, because datasets with paralogs should have

higher levels of discordance. As expected, estimated discordance-based branch lengths from the

All and One Paralogs dataset using ASTRAL are shorter than those estimated from the All

Paralogs dataset using ASTRAL-Pro, a method that accounts for the extra discordance caused by

the inclusion of paralogs (Figure 5 e, f, g). In general, across all datasets except the two including

paralogs (All and One), discordance-based branch lengths were highly similar to those estimated

in ASTRAL-Pro (Figure 5g). However, there were some surprising results. Specifically, the SCC

and MO datasets led to slightly shorter discordance-based branch length estimates than both

ASTRAL-Pro and the datasets from other tree-based decomposition methods (Figure 5g). In

addition, all discordance-based branch length estimates are relatively short, which could be

explained by difficulties estimating the lengths of longer branches with very little gene tree

discordance (i.e., for which all (or most) genes support a single topology) in ASTRAL.

Tests for introgression are consistent across datasets

To test for introgression, we looked for a deviation from the expected number of alternate

gene tree topologies using the statistic Δ (Huson et al. 2005; Vanderpool et al. 2020). We used

only the ML gene trees from each dataset for this analysis. There was evidence of introgression

across several branches of the primate phylogeny (Figure 6A), and values of Δ were similar

across datasets (Figure 6B). Notably, there was evidence of introgression in a majority of tests at

the contentious node in the Platyrrhini, which may explain difficulties inferring the species tree

topology at this node. There was also evidence of introgression in the macaques, as found by

Vanderpool et al. (2020). Deeper in the tree, results were more suspect, with tests on some

datasets suggesting introgression while others did not (Figure 6B). Results of introgression tests

were similar with less stringent missing-data filters (Supplemental Figure S5).

Discussion

Our results demonstrate that, regardless of the dataset used, the inferred species tree

topology is largely stable across datasets. Regardless of whether all families, families with only a

single copy per species, or large families from which orthologs are extracted were used, the only

disagreements between trees were with respect to relationships among the Platyrrhini; in this

case the species tree inference method was a larger determinant of results than the particular

dataset (Figure 4). Despite the overall similarity among results, when a single gene was

randomly sampled per species results were unstable in two cases, suggesting—unsurprisingly—

that such a sampling strategy is not ideal. Based on the results presented here, when whole

genome sequence data are available, using all of families output by clustering methods followed

by the application of gene-tree decomposition methods can greatly expand the data available

without sacrificing the accuracy of inference. These results align with several recent simulation

and theoretical studies focusing on gene-tree based methods and demonstrating their robustness

to paralog inclusion (Legried et al. 2020; Yan et al. 2021).

Despite the stability of inference across most of the tree, there remains disagreement

about relationships amongst the Platyrrhini, a notably contentious node (Perelman et al. 2011;

Springer et al. 2012; Perez et al. 2013; Jameson Kiesling et al. 2015; Schrago and Seuánez 2019;

Wang et al. 2019; Vanderpool et al. 2020). As in Vanderpool et al. (2020), we find that both

concatenated ML and ASTRAL based on ML gene trees favor a symmetrical topology (tree 1 in

Figure 4A). A bias towards the symmetrical 4-taxon tree is expected when using ML in the

presence of recombination and when the time between speciation events is short (Kubatko and

Degnan 2007; Roch and Steel 2015). Although the bias on ML under these conditions is often

linked to concatenation methods, if the gene trees themselves are inaccurate due to the

concatenation of multiple unique histories (e.g., among exons; Mendes et al. 2019), then the

same bias in inferred trees can occur. Bias in the gene trees can then lead to bias in the methods

that they are used as input to (e.g., ASTRAL). Note that this bias does not affect inferences under

maximum parsimony (Mendes and Hahn 2018). Furthermore, there are nearly equal numbers of

trees supporting the two best-supported topologies in the primate data (Figure 4B), which

suggests two things: first, choosing the best topology will be difficult no matter what method is

used, as the evidence in favor of one topology over the other is minimal. Second, there is likely

some introgression, since we would otherwise expect equal numbers of the two minor

topologies. We do not see equal numbers of the two minor topologies, as confirmed by

significant tests for introgression in this clade (Figure 6). Finally, a detailed comparison of SCC

gene trees inferred by both ML and MP suggests that genes whose topologies disagreed across

the two approaches did not support either topology as strongly as genes for which ML and MP

agreed (Figure S6). Of the gene trees that agreed across ML and MP inference, more supported

Tree 2 than supported Tree 1 (Figure 4A). Thus, of the genes for which methods agree, more

support the asymmetric topology than the symmetric topology (as in Vanderpool et al. 2020).

We also compared branch length estimates and tests for introgression across datasets.

Branch length estimates are largely consistent across datasets, with the exception of datasets that

explicitly include paralogs, which led to biases in expected directions for both discordance-based

and site-based branch lengths. Site-based branch lengths are very consistent across all datasets

except the One Paralogs dataset when stringent filters for missing data are applied. When

paralogs are included, site-based branch lengths are overestimated, as expected (e.g., Siu-Ting et

al. 2019). Discordance-based branch lengths (i.e., those estimated in ASTRAL) are

underestimated for datasets including paralogs, because these datasets have higher levels of

discordance. These methods accommodate increased discordance by positing a shorter time

between speciation events. Otherwise, discordance-based branch lengths are largely similar

across datasets, though the SCC and MO datasets appear to have slightly shorter estimated

branch lengths than all other methods (Figure 5e). Given the consistency of results across tree-

based decomposition methods, as well as ASTRAL-Pro, and the vastly larger number of gene

trees used in these cases, we suggest that discordance-based branch lengths may actually be

underestimated for the SCC and MO datasets. This result is consistent with lower gCFs in these

datasets (Figure 3b), and suggests that branch lengths estimated from these datasets may be

inaccurate because they include pseudoorthologs.

To our knowledge, this is the first evaluation of the effects of including more than just

single-copy families on tests for introgression based on the asymmetry in minor topology

frequencies. We expected that the inclusion of paralogs would not bias such tests, because under

models that include duplication and loss the two minor topologies should occur in equal

frequencies (Smith and Hahn 2021a; Smith and Hahn 2021b). Our results largely confirm these

expectations: although there is variation in whether or not tests are significant across datasets,

estimates of Δ are very similar (Figure 6B). At some nodes there is consistent evidence for

introgression across datasets, suggesting a strong signal of asymmetry: for example in the

macaques and among the Platyrrhini. Deeper in the tree, there may be more gene tree error (e.g.,

due to long-branch attraction), since introgression is detected for some datasets and not for others

(Figure 6B).

Phylogenetics based on whole-genome sequences almost always begins by identifying

homologous genes via clustering. The clustering process operationally defines gene families,

using clustering methods that range from very simple to very complex. While the single-copy

clusters output by any one of these methods have most often been used in phylogenetics, there is

nothing inherently more suitable about these clusters. First, SCCs may not be orthologs, due to

the presence of pseudoorthologs—paralogs that are mistaken as orthologs due to differential

patterns of gene duplication and loss (Doolittle and Brown 1994; Koonin 2005). In other words,

having only a single representative sequence in each species does not guarantee that all the

sampled genes are orthologs. Second, and more importantly, the size of clusters identified by

clustering approaches is determined by parameters set by the user. For example, in OrthoMCL

(Li 2003) the inflation parameter determines the size of output clusters: by changing this

parameter, users can identify larger or smaller clusters. Because genes are related to all other

genes via a long history of duplication and divergence (with a few exceptions; Knowles and

McLysaght 2009; Zhao et al. 2014), there is no single level of similarity that uniquely identifies

gene families (Demuth and Hahn 2009). However, users can choose the value of the inflation

parameter that identifies more, smaller clusters in order to find more single-copy clusters; this

does not mean these genes do not have paralogs, only that more distant paralogs were not

included at this clustering threshold. Many clustering methods aim to form groups of genes that

descend from a single common ancestor in the studied taxa (e.g., Emms and Kelly 2015), though

this does not ensure a lack of duplication events since the common ancestor. While tree-based

decomposition approaches still rely on the clustering step to initially identify the homologs from

which gene trees are built, their output is directly related to the definitions of orthologs and

paralogs, and is more easily interpreted in a phylogenetic context. By applying these

decomposition approaches to larger clusters, researchers can avoid arbitrary determinants of

which clusters are single-copy and can instead attempt to extract as many sets of orthologs as

possible. Not only does this approach increase the amount of data available, but it also uses

criteria more directly linked to the evolutionary history of gene families.

In conclusion, our results suggest that methods for species tree inference are accurate

across datasets whether single-copy clusters are used, tree-based decomposition methods are

used to extract orthologs from larger clusters, or all clusters (i.e., orthologs and paralogs) are

included. Our results highlight the benefits of using all gene families by showing that the amount

of data used can be increased by an order of magnitude (Table 1; Figure 2). While even the

smallest dataset was sufficient for accurate species tree inference in this case study, that is not

always the case (e.g., Emms and Kelly 2018; Thomas et al. 2020). Finally, more data facilitates

inferences beyond species tree topology, including branch length estimates and the detection of

introgression.

Materials and Methods

Dataset and alignment

The full sets of protein-encoding genes for 26 primates and 3 non-primates were obtained

as in (Vanderpool et al. 2020), and clusters were obtained as in that study. Briefly, an all-by-all

BLASTP search (Altschul et al. 1990; Camacho et al. 2009) was executed, and the longest

isoform of each protein-coding gene from each species was used. Then, the mcl algorithm (Van

Dongen 2000) as implemented in FastOrtho (Wattam et al. 2014) with an inflation parameter of

5 was used to cluster the BLASTP output. CDSs for each cluster that included samples from at

least four species were aligned, cleaned and trimmed as in (Vanderpool et al. 2020). Sequences

were aligned by codon using GUIDANCE2 (Sela et al. 2015) with MAFFT v7.407 (Katoh and

Standley 2013) with 60 bootstrap replicates. Sequence residues with GUIDANCE scores < 0.93

were converted to gaps and sites with >50% gaps were removed using Trimalv1.4rev22

(Capella-Gutiérrez et al. 2009). Alignments shorter than 200 bp and alignments that were

invariant or contained no parsimony informative characters were removed from further analyses.

A subset of alignments that could not be aligned by codon were aligned by nucleotide, and

subsequent steps were as with the codon-aligned dataset. In total 18,484 alignments were used in

downstream analyses.

Gene tree inference

We inferred gene trees from all alignments with at least four species (18,484 alignments)

in IQ-TREE v2.0.6 (Nguyen et al. 2015) with nucleotide substitution models selected using

ModelFinder (Kalyaanamoorthy et al. 2017) as implemented in IQ-TREE. The full IQ-TREE

command used on each alignment was 'iqtree2 -s *alignment name* -m MFP -c 1 -pre *alignment*

name'. We also inferred gene trees from all 18,484 alignments using the maximum parsimony

criterion in PAUP* v 4.0a (Swofford 2001). We treated gaps as missing data, obtained a starting

tree via random stepwise addition, held a single tree at each step, and used the TBR branch-

swapping algorithm with a reconnection limit of 8. We kept a maximum of 1000 trees and did

not collapse zero-length branches.

Filtering

We considered three major groups of filtering methods:

1) Single-copy clusters (SCCs): We considered a dataset that consisted only of those clusters
that included a single gene copy from each species.

2) Tree-based decomposition approaches: We considered several methods that involved
trimming branches of gene trees to extract orthologs. All custom branch-cutting operations
were written in python3 and used the python package ete3 (Huerta-Cepas et al. 2016) to read,
traverse, trim, and output gene trees and modified sequence alignments. We used postorder
node traversal when traversing trees, and prior to custom trimming operations, we midpoint-
rooted gene trees.

i. Lineage-specific duplicates: In this dataset we identified gene duplications that were
specific to a single species. For such lineage-specific duplicates, we selected the sequence
copy that was closest in length to the median length of sequences in the alignment, kept
that copy, and trimmed the other copy or copies from both the alignment and the gene tree.

ii. Two-lineage duplicates: To expand our data beyond lineage-specific duplicates, in addition
to trimming lineage-specific duplicates, we identified gene duplications specific to a pair of
species. For such duplicates, we selected the two sequence copies with the minimum
branch distance separating them and trimmed the remaining copies from the tree and the
alignment.

iii. Minimum Inclusion: We applied the minimum inclusion (MI) approach described in (Yang
and Smith 2014) to trim gene trees. We used the python script provided by Yang and Smith
(2014; prune_paralogs_MI.py) and used as input one of three sets of gene trees: the original
18,484 gene trees, the original 18,484 gene trees with lineage-specific duplicates trimmed,
and the original 18,484 gene trees with lineage-specific and two-lineage duplicates
trimmed. For the MI approach, branches longer than a specified threshold are trimmed to
remove potential pseudoorthologs; we used the following branch-length cutoffs: 0.4
substitutions per site for the ML gene trees and 500 changes for MP trees. We explored
additional cutoffs in the Supplementary Material (Appendix A).

iv. Monophyletic Outgroups: We also applied the monophyletic outgroups (MO) approach
described in (Yang and Smith 2014) to trim gene trees. We used the python script provided

by Yang and Smith (2014; prune_paralogs_MO.py) and used as input one of three sets of

gene trees: the original 18,484 gene trees, the original 18,484 gene trees with lineage-

specific duplicates trimmed, and the original 18,484 gene trees with lineage-specific and

two-lineage specific duplicates trimmed.

v. Subtree Extraction: Finally, we evaluated a new tree-based decomposition approach

introduced here (subtree extraction; SE). In this approach we start by midpoint-rooting

gene trees, followed by trimming lineage-specific and two-lineage duplicates. We then

extract subtrees with a single representative from each taxon (i.e., subtrees with no

duplicates) and keep those subtrees that meet minimum taxon-sampling thresholds.

3) Paralog methods: We considered two approaches that included paralogs in addition to

orthologs. First, we included all genes (All Paralogs). Additionally, we randomly sampled a

single gene (without regard to orthology) per species (One Paralogs).

For all datasets, we considered a stringent (minimum of 27 of 29 taxa) and relaxed (minimum of

4 of 29 taxa) missing data threshold.

Species tree inference

We inferred species trees using six methods. Three methods inferred species trees from

concatenated datasets: maximum parsimony, maximum likelihood, and SVDQuartets. To infer

an MP tree from the concatenated datasets we used PAUP* v4.0a (build 168) (Swofford 2001).

We set the criterion to parsimony, and used 500 bootstrap replicates to assess nodal support. For

all other options we used PAUP* defaults. To infer an ML tree from the concatenated dataset we

used IQ-TREE v2.0.6 (Nguyen et al. 2015) with nucleotide substitution models selected using

ModelFinder (Kalyaanamoorthy et al. 2017) as implemented in IQ-TREE. We used an edge-

linked, proportional partition model (Chernomor et al. 2016) and 1000 ultrafast bootstrap

replicates (Hoang et al. 2018). The full IQ-TREE command used on each alignment was 'iqtree2

-s *alignment name* -p *partition file name* -c 1 -pre *alignment name -B 1000*'. For three

alignments, IQTree v2.0.6 failed to run, and, based on a suggestion from the developers, we

reverted to IQTree v.1.6.12 to infer the species trees for these alignments. For these three

alignments, the full IQ-TREE command used was 'iqtree -s *alignment name* -spp *partition file*

name -pre *alignment name* -bb 1000 -nt 4'. Finally, to infer a species tree from the concatenated

alignments using SVDQuartets, we used PAUP* v4.0a (build 168) (Swofford 2001). We

evaluated all quartets and treated ambiguous sites as missing to infer the species tree topology.

To assess nodal support, we evaluated 10,000 random quartets for each of 100 bootstrap

replicates. We used the multilocus bootstrapping option and again treated ambiguous sites as

missing.

In addition to the three concatenation-based methods, we inferred species trees using

three gene-tree based methods. Prior to inferring species trees or estimating discordance (see

below) from filtered gene trees, we collapsed all zero-length branches. For each gene tree, we did

the following: First, we midpoint-rooted the gene tree. Then, we calculated site concordance

factors using IQ-Tree v2.0.6 (Minh et al. 2020) for the alignment with the rooted gene tree as the

reference tree. We used 100 randomly sampled quartets to compute the sCF, collapsing any

nodes where $sN == 0$; in other words, any nodes for which no sites were informative.

We inferred a species tree using ASTRAL-III v5.7.3 (Sayyari and Mirarab 2016; Zhang

et al. 2018; Rabiee et al. 2019). ASTRAL infers a species tree from a set of gene trees by

extracting quartets and finding the species tree that maximizes the number of shared quartet

trees. It has been demonstrated to be consistent under the multispecies coalescent (MSC) model

(Mirarab et al. 2014) and under models of gene duplication and loss (Legried et al. 2020). Gene

trees obtained using ML and MP, from all datasets described above, and with zero-length

branches collapsed, were used as input to ASTRAL; local posterior probabilities were used to

assess nodal support. For the All Paralogs dataset, we used the mapping file and treated each

gene copy as a separate individual. Additionally, we inferred species trees using ASTRID v2.2.1

(Vachaspati and Warnow 2015), again using the filtered and zero-length collapsed ML and MP

gene trees as input. ASTRID is a distance-based approach that estimates species trees using

internode distances and is statistically consistent under the MSC model (Vachaspati and Warnow

2015). As in ASTRAL, for the All Paralogs dataset we treated gene copies from the same species

as individuals using the mapping file. Finally, we inferred species trees from the All Paralogs

datasets using ASTRAL-Pro (Zhang et al. 2020). ASTRAL-Pro uses an internal rooting-and-

tagging algorithm to label nodes as duplication or speciation nodes, and then infers quartets

using only speciation nodes before finding the species tree that maximizes the number of shared

quartet trees. ASTRAL-Pro has been shown to be statistically consistent under a model of gene

duplication and loss (Zhang et al. 2020).

Assessing discordance

To assess levels of discordance across datasets we calculated site and gene concordance

factors in IQ-Tree v2.0.6 (Minh et al. 2020). We used as the reference tree the tree shown in

Figure 3, and to estimate sCFs, we used 1000 randomly sampled quartets. gCFs were estimated

for filtered ML and MP gene trees after zero-length branches were collapsed. sCFs were

estimated for the alignments that resulted from filtering the ML gene trees.

Testing for introgression

We used the approach used in (Vanderpool et al. 2020) to test for introgression. Briefly,

the introgression test assesses whether there is a deviation from the expected equal numbers of

alternative tree topologies (under the MSC model without gene flow) using the statistic Δ (Huson

et al. 2005), where

$$\Delta = \frac{Number\ of\ DF1\ trees - Number\ of\ DF2\ trees}{Number\ of\ DF1\ trees + Number\ of\ DF2\ trees}$$

DF1 represents the most common minor topology, and *DF2* represents the least common minor

topology. In the absence of introgression, Δ is expected to be equal to zero. To test whether

deviations from zero were significant, we followed the procedure of (Vanderpool et al. 2020) and

used 2,000 datasets generated by resampling gene trees with replacement, considering only those

nodes where more than five percent of trees were discordant. This distribution was used to

calculate *Z*-scores and *p*-values for the observed Δ statistic, and for each filtered dataset, we

corrected for multiple comparisons using the Dunn-Sidák correction (Dunn 1959; Šidák 1967).

Acknowledgements

This work was supported by a National Science Foundation postdoctoral fellowship to MLS

(DBI-2009989) and an NSF grant to MWH (DEB-1936187).

Data Availability

Scripts used for filtering gene trees are available on GitHub

(github.com/meganlsmith/Primate_Paralogs). Alignments, gene trees, and species trees are

available from FigShare (doi: 10.6084/m9.figshare.16653025).

Tables

Filter	MIN4	MIN27
Single-copy clusters	5771	1820
Lineage-specific duplicates (LSD)	13627	7693
Two-species duplicates (TSD)	14931	8719
Minimum Inclusion	27880	4849
Minimum Inclusion (LSD)	22360	11479
Minimum Inclusion (TSD)	21793	12046
Monophyletic Outgroups	9724	4805
Monophyletic Outgroups (LSD)	16962	10222
Monophyletic Outgroups (TSD)	17104	10254
Subtree Extraction	20562	12198
All Paralogs	18484	11555
One Paralogs	18484	11555

567 **Table 1: Number of genes trees included with different filtering approaches.** LSD and TSD

568 indicate when lineage-specific and both lineage-specific and two-lineage specific duplicates were

569 trimmed; the subtree extraction method trims these automatically. The MIN4 dataset required a

570 minimum of 4 taxa (out of 29 total), while the MIN27 dataset required a minimum of 27 taxa.

572　**Figures**

573
574

575　**Figure 1.** Conceptual overview of methods for inferring species trees from genomic data. We
576　begin with All Genes, clustering them into gene families. We can then use single-copy ortholog
577　clusters for inference (Dataset 1), use tree-based decomposition approaches to extract orthologs
578　from all clusters (Dataset 2), or infer species trees from all clusters (i.e., from datasets including
579　orthologs and paralogs; Dataset 3).

A

B

Figure 2. Numbers of informative genes and sites across datasets using the MIN27 datasets. A)

The distribution of the number of decisive sites (across branches) as calculated in IQ-Tree.

Decisive sites are defined in Minh et al. (2020). B) The distribution of the number of decisive

gene trees (across branches) as calculated in IQ-Tree. Decisive gene trees are defined in Minh et

al. (2020). SCC=single-copy clusters; LSD=lineage-specific duplicates; TSD=two-species

duplicates; MO=monophyletic outgroup; MI=minimum inclusion; SE=subtree extraction;

ONE=one paralogs.

Figure 3. Gene (gCF) and site (sCF) concordance factors among primate datasets using ML gene trees (MIN27). A) Primate phylogeny from ASTRAL using the ML gene trees (all input datasets give the same topology). Nodes show Node ID: gCF values from the SCC dataset. B) Distribution of gCF values across datasets. C) Distribution of sCF values across datasets. Node IDs correspond to the numbers displayed on the tree in panel A. SCC=single-copy clusters; LSD=lineage-specific duplicates; TSD=two-species duplicates; MO=monophyletic outgroup; MI=minimum inclusion; SE=subtree extraction; ONE=one paralogs.

Figure 4. Alternative resolutions of Platyrrhini relationships. A) The three most common tree
topologies. Below each resolution, inference methods and filtering approaches that supported the
topology are listed. B) The percentage of gene trees supporting Tree 1 minus the percentage of
gene trees supporting Tree 2 for ML and MP gene trees across datasets. C) The percentage of
sites supporting Tree 1 minus the percentage of sites supporting Tree 2 across datasets.
SCC=single-copy clusters; LSD=lineage-specific duplicates; TSD=two-species duplicates;
MO=monophyletic outgroup; MI=minimum inclusion; SE=subtree extraction; ONE=one
paralogs. Results in B and C from MIN27 datasets.

Figure 5. Branch lengths across datasets and species tree inference methods. Site-based branch

lengths estimated using concatenated ML when A) SCCs and B) one randomly selected paralog

per species are used for inference. Note the different scales in panels A and B. C) Difference

between site-based branch lengths for internal branches from the SCC dataset and all other

datasets, normalized by SCC branch length. D) Same as in panel C, but for terminal branches.

Discordance-based branch lengths calculated on the All Paralogs dataset when E) ASTRAL-Pro

and F) ASTRAL are used for inference. Note that terminal branch lengths are arbitrary in these

panels. G) Difference between discordance-based branch lengths estimated with ASTRAL-Pro

(APro) and all other methods, normalized by APro branch length. Colors represent different

filtering methods, and each row is a different branch. SCC=single-copy clusters; LSD=lineage-

specific duplicates; TSD=two-species duplicates; MO=monophyletic outgroup; MI=minimum

inclusion; SE=subtree extraction; ONE=one paralogs. Results from MIN27 datasets.

Figure 6. Results of introgression tests on MIN27 ML gene trees. A) Pie charts are shown for branches with any significant introgression tests. Numbers are node numbers. B) For all branches with some significant tests, we show the number of informative genes versus Δ. Observations are colored by filtering method, and shapes indicate whether a particular test was significant. SCC=single-copy clusters; LSD=lineage-specific duplicates; TSD=two-species duplicates; MO=monophyletic outgroup; MI=minimum inclusion; SE=subtree extraction; ONE=one paralogs.

Reference

Altenhoff AM, Glover NM, Dessimoz C. 2019. Inferring orthology and paralogy. In: Anisimova M, editor. Evolutionary genomics: Statistical and computational methods. New York, NY: Springer. p. 149–175. Available from: https://doi.org/10.1007/978-1-4939-9074-0_5

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* 215:403–410.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25:1972–1973.

Chernomor O, von Haeseler A, Minh BQ. 2016. Terrace aware data structure for phylogenomic inference from supermatrices. *Syst Biol* 65:997–1008.

Chifman J, Kubatko L. 2014. Quartet inference from SNP data under the coalescent model. *Bioinformatics* 30:3317–3324.

Demuth JP, Hahn MW. 2009. The life and death of gene families. *BioEssays* 31:29–39.

Doolittle WF, Brown JR. 1994. Tempo, mode, the progenote, and the universal root. *PNAS* 91:6721–6728.

Dunn CW, Howison M, Zapata F. 2013. Agalma: an automated phylogenomics workflow. *BMC Bioinformatics* 14:330.

Dunn OJ. 1959. Confidence intervals for the means of dependent, normally distributed variables. *J Am Stat Assoc* 54:613–621.

Emms DM, Kelly S. 2015. OrthoFinder: solving fundamental biases in whole genome comparisons dramatically improves orthogroup inference accuracy. *Genome Biol* 16:157.

Emms DM, Kelly S. 2018. STAG: Species Tree Inference from All Genes. bioRxiv:10.1101/267914.

Fernández R, Kallal RJ, Dimitrov D, Ballesteros JA, Arnedo MA, Giribet G, Hormiga G. 2018.
Phylogenomics, diversification dynamics, and comparative transcriptomics across the
spider tree of life. *Curr Biol* 28:1489–1497.

Fitch WM. 1970. Distinguishing homologous from analogous proteins. *Syst Zool* 19:99–113.

Goodman M, Czelusniak J, Moore GW, Romero-Herrera AE, Matsuda G. 1979. Fitting the gene
lineage into its species lineage, a parsimony strategy illustrated by cladograms
constructed from globin sequences. *Syst Biol* 28:132–163.

Hill M, Legried B, Roch S. 2020. Species tree estimation under joint modeling of coalescence
and duplication: sample complexity of quartet methods. arXiv:2007.06697.

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving the
ultrafast bootstrap approximation. *Mol Biol Evol* 35:518–522.

Huerta-Cepas J, Serra F, Bork P. 2016. ETE 3: reconstruction, analysis, and visualization of
phylogenomic data. *Mol Biol Evol* 33:1635–1638.

Huson DH, Klöpper T, Lockhart PJ, Steel MA. 2005. Reconstruction of reticulate networks from
gene trees. In: Miyano S, Mesirov J, Kasif S, Istrail S, Pevzner PA, Waterman M, editors.
Research in Computational Molecular Biology. Vol. 3500. Lecture Notes in Computer
Science. Berlin, Heidelberg: Springer Berlin Heidelberg. p. 233–249. Available from:
http://link.springer.com/10.1007/11415770_18

Jameson Kiesling NM, Yi SV, Xu K, Gianluca Sperone F, Wildman DE. 2015. The tempo and
mode of New World monkey evolution and biogeography in the context of phylogenomic
analysis. *Mol Phylogenet Evol* 82:386–399.

Kallal RJ, Fernández R, Giribet G, Hormiga G. 2018. A phylotranscriptomic backbone of the
orb-weaving spider family Araneidae (Arachnida, Araneae) supported by multiple
methodological approaches. *Mol Phylogenet Evol* 126:129–140.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder:
fast model selection for accurate phylogenetic estimates. *Nature Methods* 14:587–589.

Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Mol Biol Evol* 30:772–780.

Knowles DG, McLysaght A. 2009. Recent de novo origin of human protein-coding genes.

Genome Res 19:1752–1759.

Koonin EV. 2005. Orthologs, paralogs, and evolutionary genomics. *Annu Rev Genet* 39:309–

338.

Kubatko LS, Degnan JH. 2007. Inconsistency of phylogenetic estimates from concatenated data

under coalescence. *Syst Biol* 56:17–24.

Legried B, Molloy EK, Warnow T, Roch S. 2020. Polynomial-time statistical estimation of

species trees under gene duplication and loss. *J Comput Biol* 28:452-468.

Li L. 2003. OrthoMCL: Identification of ortholog groups for eukaryotic genomes. *Genome Res*

13:2178–2189.

Markin A, Eulenstein O. 2020. Quartet-Based inference methods are statistically consistent

under the unified duplication-loss-coalescence model. *arXiv*:2004.04299.

Mendes FK, Livera AP, Hahn MW. 2019. The perils of intralocus recombination for inferences

of molecular convergence. *Philos Trans R Soc London [Biol]* 374:20180244.

Minh BQ, Hahn MW, Lanfear R. 2020. New methods to calculate concordance factors for

phylogenomic datasets. *Mol Biol Evol* 37:2727–2733.

Mirarab S, Reaz R, Bayzid MdS, Zimmermann T, Swenson MS, Warnow T. 2014. ASTRAL:

genome-scale coalescent-based species tree estimation. *Bioinformatics* 30:i541–i548.

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol Evol*

32:268–274.

Perelman P, Johnson WE, Roos C, Seuánez HN, Horvath JE, Moreira MAM, Kessing B, Pontius
J, Roelke M, Rumpler Y, et al. 2011. A Molecular Phylogeny of Living Primates. *PLOS*
Genetics 7:e1001342.

Perez SI, Tejedor MF, Novo NM, Aristide L. 2013. Divergence Times and the Evolutionary
Radiation of New World Monkeys (Platyrrhini, Primates): An Analysis of Fossil and
Molecular Data. *PLOS ONE* 8:e68029.

Rabiee M, Sayyari E, Mirarab S. 2019. Multi-allele species reconstruction using ASTRAL. *Mol*
Phylogenet Evol 130:286–296.

Roch S, Steel M. 2015. Likelihood-based tree reconstruction on a concatenation of aligned
sequence data sets can be statistically inconsistent. *Theor Popul Biol* 100:56–62.

Sayyari E, Mirarab S. 2016. Fast coalescent-based computation of local branch support from
quartet frequencies. *Mol Biol Evol* 33:1654–1668.

Schrago CG, Seuánez HN. 2019. Large ancestral effective population size explains the difficult
phylogenetic placement of owl monkeys. *Am J Primatol* 81:e22955.

Scornavacca C, Delsuc F, Galtier N. 2020. Phylogenetics in the genomic era. Open access book
available from https://hal.inria.fr/PGE/.

Sela I, Ashkenazy H, Katoh K, Pupko T. 2015. GUIDANCE2: accurate detection of unreliable
alignment regions accounting for the uncertainty of multiple parameters. *Nucleic Acids*
Res 43:W7–W14.

Šidák Z. 1967. Rectangular confidence regions for the means of multivariate normal
distributions. *J Am Stat Assoc* 62:626–633.

Siu-Ting K, Torres-Sánchez M, San Mauro D, Wilcockson D, Wilkinson M, Pisani D,
O'Connell MJ, Creevey CJ. 2019. Inadvertent paralog inclusion drives artifactual
topologies and timetree estimates in phylogenomics. *Mol Biol Evol* 36:1344–1356.

Smith ML, Hahn MW. 2021a. New approaches for inferring phylogenies in the presence of
paralogs. *TIG* 37:174–187.

Smith ML, Hahn MW. 2021b. The frequency and topology of pseudoorthologs.
bioRxiv:2021.02.17.431499.

Springer MS, Meredith RW, Gatesy J, Emerling CA, Park J, Rabosky DL, Stadler T, Steiner C,
Ryder OA, Janečka JE, et al. 2012. Macroevolutionary dynamics and historical
biogeography of primate diversification inferred from a species supermatrix. *PLOS ONE*
7:e49521.

Swofford DL. 2001. Paup*: Phylogenetic analysis using parsimony (and other methods) 4.0. B5.

Thomas GWC, Dohmen E, Hughes DST, Murali SC, Poelchau M, Glastad K, Anstead CA,
Ayoub NA, Batterham P, Bellair M, et al. 2020. Gene content evolution in the
arthropods. *Genome Biol* 21:15.

Vachaspati P, Warnow T. 2015. ASTRID: Accurate species trees from internode distances. *BMC*
Genomics 16:S3.

van der Heijden RT, Snel B, van Noort V, Huynen MA. 2007. Orthology prediction at scalable
resolution by phylogenetic tree analysis. *BMC Bioinformatics* 8:83.

Van Dongen SM. 2000. Graph clustering by flow simulation.

Vanderpool D, Minh BQ, Lanfear R, Hughes D, Murali S, Harris RA, Raveendran M, Muzny
DM, Hibbins MS, Williamson RJ, et al. 2020. Primate phylogenomics uncovers multiple
rapid radiations and ancient interspecific introgression. *PLOS Biology* 18:e3000954.

Wang X, Lim BK, Ting N, Hu J, Liang Y, Roos C, Yu L. 2019. Reconstructing the phylogeny of
new world monkeys (platyrrhini): evidence from multiple non-coding loci. *Curr Zool*
65:579–588.

Wattam AR, Abraham D, Dalay O, Disz TL, Driscoll T, Gabbard JL, Gillespie JJ, Gough R, Hix
D, Kenyon R, et al. 2014. PATRIC, the bacterial bioinformatics database and analysis
resource. *Nucleic Acids Res* 42:D581–D591.

Willson J, Roddur MS, Liu B, Zaharias P, Warnow T. 2021. DISCO: Species tree inference
using multi-copy gene family tree decomposition. *Syst Biol* syab070.

Yan Z, Smith ML, Du P, Hahn MW, Nakhleh L. 2021. Species tree inference on data with
paralogs is accurate using methods intended to deal with incomplete lineage sorting. *Syst
Biol* syab056.

Yang Y, Smith SA. 2014. Orthology inference in nonmodel organisms using transcriptomes and
low-coverage genomes: Improving accuracy and matrix occupancy for phylogenomics.
Mol Biol Evol 31:3081–3092.

Zhang C, Rabiee M, Sayyari E, Mirarab S. 2018. ASTRAL-III: polynomial time species tree
reconstruction from partially resolved gene trees. *BMC Bioinformatics* 19:153.

Zhang C, Scornavacca C, Molloy EK, Mirarab S. 2020. ASTRAL-Pro: quartet-based species-tree
inference despite paralogy. *Mol Biol Evol* 37:3292–3307.

Zhao L, Saelao P, Jones CD, Begun DJ. 2014. Origin and spread of de novo genes in *Drosophila
melanogaster* populations. *Science* 343:769–772.