Title: Hyperconcentrated Mucus Unifies Submucosal Gland and Superficial Airway Dysfunction in Cystic Fibrosis

Authors: Takafumi Kato[1,2], Giorgia Radicioni[1], Micah J. Papanikolas[3], Georgi V. Stoychev[1], Matthew R. Markovetz[1], Kazuhiro Aoki[4], Mindy Porterfield[4], Kenichi Okuda[1], Selene M. Barbosa Cardenas[1], Rodney C. Gilmore[1], Cameron B. Morrison[1], Camille Ehre[1], Kimberlie A. Burns[1], Kristen K. White[5], Henry P. Goodell[1], Holly Thacker[1], Henry T. Loznev[1], Lawrence J. Forsberg[6], Takahide Nagase[2], Michael Rubinstein[7], Scott H. Randell[1], Michael Tiemeyer[4], David B. Hill[1], Mehmet Kesimer[1], Wanda K. O'Neal[1], Stephen T. Ballard[8], Ronit Freeman[3], Brian Button[1]*, Richard C. Boucher[1]*

Affiliations:

[1]Marsico Lung Institute, University of North Carolina at Chapel Hill; Chapel Hill, NC, USA

[2]Department of Respiratory Medicine, University of Tokyo; Tokyo, Japan

[3]Department of Applied Physical Sciences, University of North Carolina at Chapel Hill; Chapel Hill, NC, USA

[4]Complex Carbohydrate Research Center, University of Georgia; Athens, GA, USA

[5]Department of Pathology and Laboratory Medicine, University of North Carolina at Chapel Hill; Chapel Hill, NC, USA

[6]Protein Expression and Purification Core, University of North Carolina at Chapel Hill; Chapel Hill, NC, USA

[7]Department of Mechanical Engineering and Materials Science, Duke University; Durham, NC, USA

[8]Department of Physics and Astronomy, University of North Carolina at Chapel Hill; Chapel Hill, NC, USA

[9]Department of Physiology and Cell Biology, University of South Alabama; Mobile, AL, USA

*Co-corresponding authors. Email: richard_boucher@med.unc.edu (R.C.B.), brian_button@med.unc.edu (B.B.)

One Sentence Summary: Hyperconcentrated Mucus Unifies Submucosal Gland and Superficial Airway Dysfunction in Cystic Fibrosis

Abstract:

Cystic fibrosis (CF) is characterized by abnormal transepithelial ion transport. However, a description of CF lung disease pathophysiology unifying superficial epithelial and submucosal gland (SMG) dysfunctions has remained elusive. We hypothesized that biophysical abnormalities associated with CF mucus hyperconcentration provide a unifying mechanism. Studies of the anion secretion-inhibited pig airway CF model revealed elevated SMG mucus concentrations, osmotic pressures, and SMG mucus accumulation. Human airway studies revealed hyperconcentrated CF SMG mucus with raised osmotic pressures and cohesive forces predicted to limit SMG mucus secretion/release. Utilizing proline-rich protein 4 (PRR4) as a biomarker of SMG secretion, proteomics analyses of CF sputum revealed markedly lower PRR4 levels compared to healthy and bronchiectasis controls, consistent with a failure of CF SMGs to secrete mucus onto airway surfaces. Raised mucus osmotic/cohesive forces, reflecting mucus hyperconcentration, provide a unifying mechanism that describes disease-initiating mucus accumulation on airway surfaces and within SMGs of the CF lung.

Main text:

INTRODUCTION

Cystic fibrosis (CF) is characterized by abnormal transepithelial ion transport due to genetic mutations of the cystic fibrosis transmembrane regulator (CFTR) (*1-3*). *In vitro* and *in vivo* studies of the superficial epithelia lining CF airway surfaces suggest dysfunctional CFTR-mediated ion transport produces hyperconcentrated mucus (*4-8*). Mucus hyperconcentration, defined as an increased mucus percent solids content (% solids) and/or mucin concentration, generates abnormal biophysical properties, *e.g.*, raised osmotic pressures and increased cohesive

forces (*8, 9*). Osmotic compression of the periciliary layer (PCL) by the mucus layer abolishes mucociliary transport, results in mucus accumulation on distal airway surfaces, and initiates the obstruction, infection, and inflammation that produces bronchiectasis in CF(*9-11*).

However, there is controversy with respect to the contribution of CF submucosal gland (SMG) dysfunction to the proximal airway features of CF lung disease. Studies in CF pig tracheas described secretion of SMG mucus strands in response to maximal cholinergic stimulation (a mimic of cough) that are postulated to adhere to proximal CF airway surfaces, accumulate, and become a nidus for persistent bacterial infection (*12-17*). These data are juxtaposed to other studies that suggest CF SMG dysfunction reflects a failure to secrete SMG mucus with its robust antimicrobial components onto airway surfaces (*18-22*). Biophysical or clinical measurements that distinguish between these hypotheses have not been reported.

We hypothesized that SMG mucus has distinct characteristics from airway surface mucus and that CF SMG mucus is altered, hindering appropriate secretion onto airway lumens. We utilized an established method for SMG mucus collection from freshly excised porcine and human airway tissues and measured multiple properties of SMG mucus and investigated how it is altered in CF and different from airway surface mucus. We also identified a biomarker selective for SMG secretion and utilized it to discuss how SMG secretion is altered in CF.

RESULTS

SMG mucus from CF-mimicking maneuvers exhibits concentration-dependent high osmotic pressures

We first studied the biophysical properties of normal vs CF-like SMG mucus in the anion transport inhibitor-treated wild-type (WT) pig model of CF (*23-26*). Because polymer

biophysical properties are higher-order powers of polymer concentration, mucus concentration was also measured in parallel (*5, 27*). Maximal acetylcholine stimulation, block of Cl$^-$ secretion [bumetanide (Bum)], block of bicarbonate secretion [dimethylamiloride (DMA)], combined Bum/DMA block, or vehicle, in WT pig SMG mucus reproduced evidence of anion-transport inhibitor-mediated block of SMG fluid and hence increased SMG mucus concentrations (% solids) (Fig.1A). Osmotic pressures exhibited similar trends as % solids (Fig.1B) suggesting the increase is dominated by mucus concentration (Fig.1C).

CF-mimicking maneuvers produce mucus retention and compressed cilia in SMG ducts

We investigated SMG ducts of the control (Ach) and CF-mimicking (Ach+Bum+DMA) maneuvers using transmission electron microscopy (TEM). Cilia in glands treated with Bum/DMA were compressed (Figs.1D,E), consistent with findings in CF (*5*). In addition, SMG mucus retention was observed in Bum/DMA (low gland fluid flow/anion secretion inhibition) conditions (Fig.1E).

Collectively, these data suggest that anion/fluid secretion inhibitors produced a hyperconcentrated pig SMG mucus that, via the osmotic compression onto ductal surface PCL, was retained in SMGs.

Human SMG mucus exhibits concentration-dependent elevated osmotic pressures and cohesive strength

We performed complementary studies of SMG mucus collected from non-CF and CF airway tissues. Consistent with the absence of CFTR anion/fluid secretory function in CF, CF SMG mucus is hyperconcentrated with increased osmotic pressures compared to non-CF SMG mucus

(Fig.2A,B). Mucus concentrations correlated with osmotic pressure over ranges bracketed by CF and non-CF SMG mucus (Fig.2C). Studies designed to measure the forces required to "tear" mucus from gland orifices for transport up airway surfaces, *i.e.*, cohesive forces (*28*), identified elevated cohesive forces in CF SMG mucus compared to non-CF (Fig.2D).

CF SMGs have mucus retention in ducts and compressed cilia

Consistent with the findings in CF-mimicking pig (Fig.1D,E), human CF SMG ducts exhibited cilial compression and mucus retention as observed by TEM (Figs.2E,F). These findings are consistent with previous reports of increased CF SMG mucus viscosity (*29*), and predict that CF SMG mucus secretion is hindered by both SMG mucus osmotic compression onto ductal surfaces and increased cohesive forces that limit mucus separation from ductal orifices.

SMG mucus shows less solubility and forms strands

We next compared the properties of airway surface [human bronchial epithelial (HBE) cell culture] and SMG mucus. Because SMG mucus in porcine models has been reported to contain "strands" that maintain their structure after secretion onto airway surfaces rather than dissolve into the ambient surface mucus layer (*12-15*), we investigated whether strand formation was unique to SMG mucus, *i.e.*, not a feature of superficial epithelial mucus, and whether CF SMGs produced abnormal strands. While superficial HBE mucus dissolved fully into phosphate buffered saline (PBS), leaving behind no strand-like structures, ~50% of non-CF human and pig SMG mucus failed to dissolve (Fig.3A). However, addition of a non-ionic, non-denaturing detergent (IGEPAL) to the mucus promoted dissolution of the insoluble SMG mucus component (Table.S1). Light scattering for the PBS-soluble SMG mucins demonstrated that SMG mucus

exhibited similar size (Rg) compared to HBE mucins (Fig.3B). Light scattering of the insoluble SMG mucins was not possible. Instead, we immunostained the insoluble components of SMG mucus and identified that strands in SMG mucus were comprised of MUC5B but not MUC5AC (Fig.3C).

Glycomics revealed SMG mucus contain shorter, less-branched glycans

Biochemical properties of SMG vs superficial epithelial MUC5B mucins were next studied. HBE and SMG mucus were collected from three matched donors. SMG mucin-associated glycans consisted of larger proportion of shorter, less-branched (Cores 1 and 3) carbohydrate structures than for HBE mucins (Figs.4A,B,S1,Data.S1). Anionic O-glycans, in which the negative charge is imparted either as sulfate or sialic acid, were also measured. Sulfation of extended O-glycans, but not sialic acid, was reduced in SMG relative to HBE MUC5B (Figs.S2,Data.S2).

CF SMG mucus strands exhibit distinct features

We visualized strands in the insoluble SMG mucus from non-CF and CF subjects using scanning electron microscopy (SEM). Consistent with CF SMG mucus hyperconcentration (Fig.2A), strand density was higher in CF (Fig.5A). The width and length of SMG strands were similar in non-CF and CF (Figs.5B,C). Alignment analyses demonstrated that non-CF SMG mucus strands contained extraordinarily long arrays of aligned bundles (Fig.5D), whereas CF SMG mucus strands exhibited randomly entangled bundles (Fig.5E). Quantitation of orientational order revealed less order, *i.e.*, more heterogeneity, in CF strands (Figs.5F,G).

Proline-rich protein 4 (PRR4) is a SMG-selective biomarker

Biomarkers selective for SMG secretion were investigated in human airway tissues. Four proteins, lactoferrin (LTF), zinc-α2-glycoprotein (AZGP1/ZAG), lysozyme (LYZ), and PRR4, were screened based on previous reports of selective SMG expression (*30-32*). RNA *in situ* hybridization (RNA-ISH) and immunohistochemistry (IHC) demonstrated that only PRR4 was selectively expressed in SMG serous cells and not in superficial epithelia lining large or small airways in healthy or diseased subjects (Figs.6A,B,S3). In contrast, LTF, AZGP1 and LYZ were detected in some surface epithelium or infiltrating cells (Fig.S4). qPCR data from freshly dissected specimens from different pulmonary regions confirmed the specificity of PRR4 for SMGs (Fig.S5). Co-staining for PRR4 and MUC5B distinguished SMG serous and mucous acini, respectively (Fig.6C). SMG mucus passing through gland ducts exhibited co-positivity for PRR4 and MUC5B (Fig.6C). In addition, SMG mucus strands exhibit co-positivity for PRR4 and MUC5B (Fig.6D). Accordingly, PRR4 was selected as a biomarker to identify the contribution of SMGs to human airway secretions.

For the co-staining of PRR4 and MUC5B, we utilized anti-MUC5B mouse antisera raised in-house (M-25), which has not been published. We therefore performed a validatory assay by staining with an established anti-MUC5B antibody (H-300, Santa Cruz) and confirmed identical positive staining (Fig.S6).

Sputum PRR4 level is reduced in CF but not in other diseases

Mass spectrometry (MS)-based proteomics analyses were performed on induced sputum samples collected from healthy controls and subjects with chronic airway diseases to investigate if disease status affects SMG contribution to oveall airway secretion. While non-CF bronchiectasis

(NCFB) and primary ciliary dyskinesia (PCD) subjects exhibited similar or increased sputum PRR4 levels compared to healthy controls, CF sputum exhibited strikingly decreased PRR4 concentrations compared to both healthy (~ 30-fold reduction) and the bronchiectatic disease controls (~ 300-fold reduction) (Fig.7A). This result did not reflect PRR4 degradation by human neutrophil elastase (*33*), as sputum HNE levels were similar in CF and NCFB/PCD subjects (Fig.7B), and PRR4 was not degraded by co-incubation with HNE *in vitro* (Fig.S7).

Mucus flakes from CF subjects show reduced PRR4

Mucous flakes, insoluble material observed in bronchoalveolar lavage fluid (BALF) (*10*), were co-stained with PRR4 and wheat germ agglutinin (WGA) ,which stains lectins. PRR4 was localized in mucus flakes from both healthy control and CF subjects. However, quantification of these signals revealed that PRR4 was reduced in CF compred to healthy control (Fig.7C).

Healthy and CF airways have similar serous vs mucous cell proportion in SMG

Proportion of serous vs mucous cells of SMGs were quantified by the images of PRR4 and MUC5B co-staining. Healthy and CF tissues exhibited similar ratio of serous and mucous cells (Fig.8).

DISCUSSION

CF airway surface epithelium exhibits hyperconcentrated mucus with raised osmotic pressures and cohesive strength (*4, 5, 9*). However, there is controversy how SMG mucus is altered in CF and if loss of CFTR affects the contribution of SMG secretions to overall airway secretion. In

this study, we sought to elucidate whether mucus hyperconcentration is a unified disease phenotype in both airway surface epithelium and SMGs in CF.

We observed that SMG mucus from pigs treated with CF-mimicking maneuvers were hyperconcentrated, consistent with previous reports by Ballard *et al* (*26, 34*) and Joo *et al* (Fig.1A) (*18*). CF SMG mucus also showed hyperconcentration compared to non-CF (Fig.2A), which fits previous studies by Martens *et al (35)* and Xie *et al* (*17*). We observed elevated SMG mucus osmotic pressures in CF-mimicking pig glands and CF subjects (Figs.1B,C,2B,C). Importantly, elevated osmotic pressure was dominated by mucus concentration (Fig.1C), consistent with observations in airway surface mucus (*8, 36*). Elevated osmotic pressure of airway surface mucus results in dehydration and osmotic compression of PCL, causing deficient ciliary transport (*5*). Cohesive forces of human CF SMG mucus were also significantly elevated compared to control (Fig.2D). TEM investigations revealed that SMG ducts of CF-mimicking pig and human CF tissues exhibited both mucus retention and cilial compression (Fig.1D,E,2E,F). These observations are consistent with previous reports of reduced airway surface fluid in CF pigs (*37*) and of increased CF SMG mucus viscosity (*29*), and predict that CF SMG mucus secretion is hindered by both SMG mucus osmotic compression onto ductal surfaces and increased cohesive forces that limit mucus separation from ductal orifices.

SMG mucus flows as a liquid but has been reported to contain "strands" that maintain their structure after secretion onto airway surfaces rather than dissolve into the ambient surface mucus layer (*14, 15*). We investigated whether strand formation was unique to SMG mucus, *i.e.*, not a feature of superficial epithelial mucus, and whether CF SMGs produced abnormal strands.

Whereas, superficial HBE mucus dissolved fully into PBS, leaving behind no strand-like structures, ~50% of non-CF human and pig SMG mucus failed to dissolve into excess PBS (Fig.3A).

Glycomics analyses for the HBE vs SMG mucus from matched donors revealed insights into SMG MUC5B strand formation. First, SMG mucin glycans consisted of larger proportion of shorter, less-branched (Cores 1 and 3) carbohydrate structures than HBE mucins (Fig.4,Data.S1). Mucin glycosylation with shorter glycans presents fewer hydroxyls for capture of water molecules through hydrogen bonding, a feature predicted to promote hydrophobic interactions between SMG MUC5B mucins and strand formation (*38*). The observation that a non-ionic, non-denaturing detergent (IGEPAL) promoted dissolution of the insoluble SMG mucus component is consistent with this prediction (Table.S1). Second, anionic O-glycans, in which the negative charge is imparted either as sulfate or sialic acid, were also measured (*38*). Sulfation of extended O-glycans, but not sialic acid, was reduced in SMG relative to HBE MUC5B, also consistent with a more hydrophobic, strand-prone SMG MUC5B mucin (Fig.S1,S2,Data.S1).

Finally, the co-localization of a unique SMG-secreted protein with SMG strands, *i.e.*, PRR4 (discussed below), suggests a role for protein cross-linkers in strand formation (Fig.6D).

Collectively, these data suggest specialized functions for surface vs SMG mucus in normal lung physiology. Superficial epithelial mucins likely dissolve rapidly upon secretion to replenish mucus cleared by transport. In contrast, strands secreted by SMGs maintain their structure on airway surfaces, *i.e.*, are "permanent" gels, but may aggregate post-secretion with inhaled particles to form collections (masses) of sufficient size to promote cough clearance in times of stress (*28*).

We also identified that CF SMG strands exhibited distinct features (Fig.5). Consistent with CF SMG mucus hyperconcentration, strand density was higher in CF SMG mucus (Fig.5A). The width and length of SMG strands were similar in non-CF and CF mucus and width approximated the inner diameter of mucus tubules (~ 15µm diameter), suggesting that strand dimensions are defined by distal SMG duct structures (Figs.5B,C) (*39*). Notably, SEM demonstrated that non-CF human SMG mucus strands contained extraordinarily long arrays of aligned bundles (Fig.5D), whereas CF SMG mucus strands exhibited randomly entangled bundles (Fig.5E). Quantitation of orientational order revealed less order, *i.e.*, more heterogeneity, in CF strands (Fig.5F,G). We speculate that decreased alignment of CF SMG strands reflects abnormal strand formation in a flow-limited, hyperconcentrated environment.

Finally, clinical investigations were performed to distinguish between the hypotheses that CF SMGs fail to secrete mucus vs secrete a strand-laden mucus that accumulates on airway surfaces. PRR4 was identified as a biomarker selective for SMG secretion (Figs.6A-C,S3,S4) to identify and evaluate the contribution of SMG to airway secretions. While NCFB and PCD subjects exhibited similar or increased sputum PRR4 levels (measured by MS) compared to healthy control subjects, CF sputum exhibited strikingly decreased PRR4 concentrations compared to both normal subjects and bronchiectatic disease controls. The former indicates hypertrophy of SMG and hypersecretion of SMG mucus in non-CF chronic airway diseases (*40*). Reduced PRR4 levels in CF sputum suggests that CF subjects do not have proper SMG secretion even though they have hypertrophied glands. Consistent results were obtained from mucus flakes in BALF harvested from healthy control and CF subjects stained with PRR4. These results did not reflect

PRR4 degradation by HNE (Fig.7B,S7) or diminished PRR4 productive/secretitary capacity (Fig.4A,B,8). Collectively, these data suggest that the SMG contribution to CF pathogenesis reflects reduced CF SMG secretion onto, not accumulation of SMG mucus strands on, proximal airway surfaces.

Our study has several limitations. Because of the limited availability of fresh human airway tissues, we included specimens from non-CF diseased (*e.g.*, smokers, chronic bronchitis) subjects in some studies (labeled as "Non-CF"). While, we believe inclusion of non-CF diseased specimens did not affect the final conclusion that CF SMG mucus exhibits distinct features, more detailed exploration is necessary to fully elucidate the SMG mucus properties between health and disease. Another limitation is that, although we identified PRR4 as a SMG-selective maker, we have not elucidated any function of PRR4. Investigation of function of PRR4 and other proline-rich protein families (Fig.S5) may identify how these proteins contribute the SMG-unique mucus properties.

In summary, SMG mucus exhibits unique characteristics configured for host defense, *e.g.*, strand formation suitable for cough clearance. SMG mucus from human CF and CF-mimicking porcine airways is hyperconcentrated compared to non-CF SMG mucus. Hyperconcentrated SMG mucus exhibits increased osmotic pressures and cohesive forces predicted to retain mucus in SMG ducts and reduce secretion. The concentration of a SMG-selective secretory protein (PRR4) was lower in CF sputum than healthy controls, and strikingly lower than in subjects with non-CF bronchiectatic diseases, suggesting that CF SMG mucus secretion onto proximal CF airway surfaces is impaired. Accordingly, restoration of SMG host defense proteins/peptides on CF

airway surfaces early in life may be therapeutic (*41*). Finally, these data suggest that increased

mucus osmotic/cohesive forces, a product of abnormal CFTR-dependent mucus hydration

(hyperconcentration), provide a unifying mechanism to describe the muco-obstructive distal and

proximal airway components of CF lung disease.

MATERIALS AND METHODS

Human airway dissection

Human tissues were provided by the University of North Carolina (UNC) Tissue and Cell

Culture Procurement Core under protocols approved by UNC Office of Research Ethics

Biomedical Institutional Review Board (IRB number 03-1396). Non-CF airway tissues were

obtained from lungs of donors that could not be used for transplant. CF airways were obtained

from lungs resected from CF patients undergoing transplantation at UNC Hospitals. Excised

tissues were transported in saline on ice, and all studies were conducted within 12 to 36 hours of

lung removal. Proximal airways (2^{nd} to 4^{th} generation bronchi) were manually dissected.

Intraluminal mucus was removed by washing with Ringer's bicarbonate solution until the

flushed solution appeared clear, and residual fluid was collected as completely as possible with

positive-pressure pipettes.

Pig airway dissection

All procedures with swine performed at the University of South Alabama were conducted under

protocols approved by the University of South Alabama Institutional Animal Care and Use

Committee in compliance to the US Public Health Service *Policy on Humane Care and Use of

Laboratory Animals*. Pig bronchial dissection was performed as previously described (*26*). For

the collection of SMG mucus for solubility and light scattering assays performed at UNC, freshly excised porcine airway tissues were obtained from a local slaughterhouse. Main stem or segmental bronchi were dissected in the lab.

Gland mucus collection

Pig and human airway SMG mucus collection was performed using established protocols with minor modifications (*26, 35*). The proximal end of an isolated airway was cannulated with polyethylene tubing and ligated with sutures. All distal branches of the excised preparation were tightly ligated with suture material to physically separate the airway lumen from the incubation buffer, enabling SMG mucus to be secreted into the lumen without contamination by incubation buffer (Krebs-Ringer bicarbonate, pH 7.4, under continuous gassing with 95% O_2 and 5% CO_2 at 37°C). For the porcine main stem to segmental bronchi, both proximal and distal edges were sealed with silicon plugs. Acetylcholine chloride (A625, Sigma) was added to the incubation solution at a concentration of 10μM to stimulate SMG secretion. For porcine experiments, dimethylamiloride (A4562, Sigma) and/or bumetanide (B3023, Sigma) were used together to block bicarbonate and/or chloride secretion, respectively. After 2-hour incubation interval with acetylcholine with or without anion transport inhibitors, the bronchial tissues were removed from the buffer solution and decannulated. After wiping the outer surface of the preparation to remove buffer solution, the airway was opened longitudinally. Mucus secreted into the lumen was collected by aspiration with a 10μl positive pressure pipette. In each experiment, % solids concentration was measured immediately (described below), and the remaining sample was kept frozen at -80°C until further investigation.

% solids measurement

Percent solids concentrations (% solids) of human and porcine SMG mucus (3 to 5 μl) were measured gravimetrically as described previously (42). The mucus aliquots were weighed on a pre-weighed piece of aluminum foil. The samples were heated at 80□ in an oven overnight to allow the liquid content to evaporate completely. The final weight of the dried foil and mucus sample was determined and solid weight% (% solids) was calculated.

Osmotic pressure measurement

Osmotic pressures of human and pig SMG mucus (20-30 ul) were measured with a custom-designed osmometer incorporating a 10-kDa membrane as previously reported (5). Twenty to 30 μl of SMG mucus was placed into the fluid chamber, and the steady-state osmotic pressure of mucus was acquired by the pressure transducer placed on the other side of the osmotic membrane.

Cohesion measurements

The cohesive strength of human SMG mucus samples (20-40 ul) was measured using a custom-designed peel test device (28, 43). Twenty to 40 μl of a SMG mucus sample was placed on a mucus-binding mesh and the peeling-force to tear the mucus apart was measured.

TEM for porcine and human airway tissues

Airway tissues were immersed in 2% paraformaldehyde/2.5% glutaraldehyde/0.15M sodium phosphate, pH 7.4 and stored for several days at 4□. After post-fixation for 1 hour with buffered potassium ferrocyanide-reduced osmium (1% osmium tetroxide/1.25% potassium

ferrocyanide/0.15M sodium phosphate) the samples were washed in deionized water, then

dehydrated through a graded ethanol and propylene oxide (*44*). The samples were infiltrated with

a 1:1, then 1:2 mixture of propylene oxide and Polybed 812 epoxy resin for 4 hours and

infiltrated overnight in 100% resin (08792-1, Polysciences). Samples were transferred to

embedding molds and polymerized at 60□ overnight. Using a diamond knife, 1µm semi-thin

sections were cut, mounted on slides, and stained with 1% toluidine blue to examine by light

microscopy and isolate regions of interest. Ultrathin sections (70-80 nm) were cut with a

diamond knife and mounted on 200 mesh copper grids followed by staining with 4% aqueous

uranyl acetate for 12 minutes and Reynold's lead citrate for 8 minutes (*45*). The samples were

observed using a JEOL JEM-1230 transmission electron microscope operating at 80kV (JEOL

USA) and images were acquired with a Gatan Orius SC1000 CCD Digital Camera and Gatan

Microscopy Suite 3.0 software (Gatan). Quantitative measurement of cilial height and semi-

quantitative scoring of mucus retention in SMG ducts were performed by a blinded observer.

Cilial height from the ductal cell surface to the tip of the cilia was quantified. Mucus retention

was scored from 0 to 3 with 0.5 intervals.

SEM of SMG mucus

SMG mucus was diluted 10-fold in PBS and mixed for two hours at room temperature (RT).

Samples were allowed to sediment onto an APTES-coated surface for 15 minutes before being

fixed with 2% paraformaldehyde, 2.5% glutaraldehyde in PBS for another 15 minutes. The

samples were washed three times with sterile water and dehydrated into ethanol using increments

of 10% in ten-minute increments. The samples were then critically point dried and mounted onto

a SEM specimen holder using copper tape, coated with 5nm AuPd (108 Auto, Cressington) and

imaged on SEM (FEI Helios Nanolab 600) using 5kV accelerating voltage. Analysis of orientation was performed using OrientationJ (http://bigwww.epfl.ch/demo/orientation/) using a method adapted from Nerger *et al* (*46*).

Human airway surface mucus production

HBE cells were isolated from non-smoker donors without respiratory diseases by the UNC Tissue and Cell Culture Procurement Core as described (*47*). For glycomics comparison of SMG vs HBE mucus, isolated HBE primary (passage 0) cells, donor-matched to tissue sections from which SMG mucus was collected, were seeded on collagen (5005, Advanced BioMatrix)-coated Transwell inserts (3460, Corning) at a density of 0.3 million cells per insert. For solubility and light scattering studies, HBE primary cells were expanded as previously described (*48*). Passage 1 or 2 cells were seeded on collagen-coated Transwell inserts at a density of 0.3 million cells per insert. The cells were cultured under submerged condition until confluence followed by air-liquid interface (ALI) culture with UNC ALI media (*49*). After a full differentiation confirmed by ciliation under light microscopic observation, secreted mucus was allowed to accumulate on apical surface for up to three weeks and collected by washing with PBS. The HBE mucus was pooled from multiple codes and then concentrated with ultrafiltration using 50K cutoff filter (UFC905096, Millipore) to the %solids of 4%.

Solubility measurements

Collected SMG and HBE mucus was diluted 40-fold with PBS and kept at 4°C for 4 days to allow the mucus to solubilize. After the 4-days, HBE mucus was completely dissolved, whereas the SMG mucus dissolved only partly in the PBS. Undissolved portions were diluted in PBS and

%solids of the insoluble part was calculated. The SMG supernatant, containing the soluble component of SMG mucus, was collected for light scattering. The PBS-insoluble SMG mucus was exposed to surfactant (1 mM in PBS, IGEPAL CA-630, Sigma) and solubility was reexamined 48 hours later.

Light scattering

Static light scattering for soluble components of SMG and HBE mucus were performed on a research goniometer (BI-200SM, Brookhaven Instruments), equipped with a 640nm laser. A series of solutions with different concentrations were measured at scattering angles between 40^{o} and 150^{o}. The data were plotted as a Zimm plot, from which the radius of gyration (R_g) of the solutes were extracted. The lognormal size distribution of the solutes was obtained by cumulant analysis of the correlation function between 100 nsec and 10 msec. Prior to measurements, all solutions were filtered through a 0.45 μm hydrophilic PVDF syringe filter (Titan 3, Thermo Scientific) to remove dust particles and large cell debris.

Analysis of mucin O-glycosylation

O-glycans from HBE and SMG mucus were released by reductive β-elimination and analyzed by nano-spray ionization multi-dimensional mass spectrometry (NSI-MSn) as described (*50, 51*). Briefly, mucus samples were precipitated from cold 80% aqueous acetone, dried, and resuspended for reductive β-elimination. Following release and clean-up, O-glycans were permethylated and recovered by water-dichloromethane (DCM) extraction. Non-sulfated and sulfated O-glycans were recovered in the DCM and aqueous phases, respectively. The two phases were separated, dried, and resulting glycans were analyzed by NSI-MSn. MS data was

acquired in both negative and positive ion modes to detect sulfated and non-sulfated glycans, respectively. For quantification of permethylated, non-sulfated glycans, a known amount of maltotetraose permethylated with isotopically heavy methyliodide was added to each O-glycan sample, and mucin glycan amounts were measured in reference to the standard (*52*). For sulfated glycans, the same quantification was not possible due to ionization suppression caused by the sulfate moiety. Therefore, sulfated glycans were quantified based on the signal intensity measured in the mass spectrometer. Graphical representations of monosaccharide residues are presented in accordance with the broadly accepted Symbolic Nomenclature For Glycans (SNFG) and O-glycan analysis was performed according to the MIRAGE guidelines (*53*).

Tissue slide preparation for RNA-ISH and IHC

Human airway and lung tissues were provided by the UNC Tissue and Cell Culture Procurement Core. The donors included healthy controls (non-smokers without any history of respiratory diseases), CF, asthma, NCFB and PCD. Tissues were collected from multiple generations, *i.e.*, trachea, main bronchi, and distal lungs with bronchioles. The dissected tissues were fixed with 10% neutral buffered formalin for 24-36 hours and embedded in paraffin. The formalin-fixed paraffin-embedded (FFPE) blocks were cut to 5 μm.

RNA-ISH

RNA-ISH was performed using the RNAscope® 2.5 HD Reagent Kit-RED (322350, Advanced Cell Diagnostics) according to the manufacturer's instruction and published protocols (*48*). Probes targeting *PRR4* (522791), *LTF* (425101), *AZGP1* (401861) and *LYZ* (421441), in addition to positive and negative controls (310041 and 310043, respectively), were hybridized at 40□ for

2 hours in the HybEZ oven (241000) followed by signal amplification and washing. Signals were visualized by Fast Red followed by counterstaining with hematoxylin. The probe targeting human housekeeping gene Ubiquitin C (*UBC*, 310041) served as a positive control to evaluate RNA quality. A bacterial gene, Bacillus subtilis dihydrodipicolinate reductase (*DapB*, 310043) served as negative control . The images were acquired using an Olympus VS200 slide scanner light microscope with a 60x 1.42 N.A. objective.

Immunostaining

FFPE tissue sections were baked at 60°C for 2 hours, deparaffinized with xylene for 10 minutes three times, and rehydrated with graded ethanol. Antigen retrieval was performed by boiling slides in 0.1 M sodium citrate pH 6.0 (1200W microwave settings: 100% power for 6.5 minutes, followed by 60% power for 6 minutes twice). After cooling, quenching of endogenous peroxidase was performed with 0.5% hydrogen peroxide in methanol for 15 minutes. Then slides were blocked with Blocking One (03953-95, Nacalai Tesque) at RT for 30 minutes. The primary antibodies rabbit anti-PRR4 (PA5-59883, Invitrogen, 1:2500), anti-LTF (10933-1-AP, Proteintech, 1:1000), anti-ZAG (13399-1-AP, Proteintech ,1:1000) and anti-LYZ (MA5-32154, Invitrogen, 1:1000) were applied to slides and incubated at 4 overnight. Sections were washed with PBS and incubated with biotinylated donkey anti-rabbit IgG secondary antibodies (711-065-152, Jackson ImmunoResearch, 1:400) at RT for 1 hour. Slides were treated with Vectastain Kit (PK-4000, Vector Laboratories) according to manufacturer's instruction. Sections were stained with 3,3'-diamino benzidine and counterstained with Fast Red. Images were acquired as described above. For immunofluorescent co-staining, tissue sections were treated with rabbit anti-PRR4 antibody and mouse anti-human MUC5B antiserum (raised in-

house, M-25, 1:1000) at 4□ overnight followed by fluorescent staining with Alexa Fluor Plus

647-conjugated donkey anti-rabbit IgG (A32795, Invitrogen, 1:1000) and Alexa Fluor® Plus

555-conjugated donkey anti-mouse antibody (A32773, Invitrogen, 1:1000) at RT for 30 minutes.

Arter autofluorescence quenching (SP-8400-15, Vector Laboratories), tissue was mounted with

DAPI-containing medium (P36931, Invitrogen). Fluorescent images were acquired using

Olympus VS200 scanner with DAPI (488 nm), Cy3 (555 nm) and Cy5 (647 nm) filters. Isotype

controls were prepared using rabbit and/or mouse IgG (011-000-003 and 015-000-003, Jackson

ImmunoResearch). Volume ratio of serous (PRR4 positive) cells to mucous (MUC5B positive)

cells volume in SMGs was quantified with Olyvia (Olympus) and VisioPharm (VisioPharm)

software.

Immunocytochemistry of SMG mucus

SMG mucus was diluted 10-fold in PBS and mixed for 2 hours at RT. Samples were allowed to

sediment onto an APTES-coated surface for 15 minutes before being fixed with 2% PFA for

another 15 minutes. Samples were washed 3 times with PBS. Samples were then incubated with

PRR4 antibody and MUC5B antisera or MUC5B antibody (H-300, Santa Cruz, 1:1000) and

MUC5AC antibody (45M1, Invitrogen, 1:1000) at 4□ overnight followed by washing 3 times

with PBS and incubated with secondary antibodies overnight at 4□. Samples were washed again

and stained with DAPI (D9542-5MG, Sigma, 1:10,000) for 15 minutes, and then washed and

mounted onto slides (9990402, Fisher Scientific). Images were acquired with a Zeiss 880

confocal microscope. To view co-localization of MUC5B and MUC5AC, Airyscan acquisition

mode was used for imaging (Fig.3C).

RNA isolation from multiple regions of human airways

Freshly excised airway tissues from healthy non-smoker donors were dissected under light microscopy. Small (~ 1 cm^2) sections that contained surface and submucosal tissue were dissected from trachea and main bronchus. Small bronchioles were carefully dissected from surrounding alveolar tissue. Distal lung parenchyma sections of 0.5 to 1cm^2 in size were cut as alveolar tissues. Dissected specimens were minced with a homogenizer, and total RNA extracted with TRI reagent (T9424, Sigma) and an RNA extraction kit (R2052, Zymo Research).

cDNA synthesis and RT-qPCR

200ng of RNA was used for synthesizing cDNA (1708840, Bio-Rad) according to the manufacturer's instructions. qPCR was performed with QuantStadio 6 (Thermo Fisher). Taqman PCR probes (Life Technologies) used for the investigation of SMG markers included *PRR4* (Hs00200615_m1), *PRB1* (Hs00818764_m1), *PRB3* (Hs00818925_m1), *PRB4* (Hs00864002_m1), *PRH2* (Hs00818765-mH), *LTF* (Hs00914334_m1), *AZGP1* (Hs00426651_m1) and *LYZ* (Hs00426232_m1). A marker for distal bronchioles and alveoli, *SFTPB* (Hs00167036_m1) was measured to assure airway regions were obtained from. *TBP* (Hs00427620_m1) was chosen for the internal control. All probes were designed to span exons.

Label-free quantification mass spectrometry for induced sputum

Induced sputum samples were obtained from the UNC sample repositories (IRB numbers 16-3142 (NCFB), 15-2431 (healthy controls and CF) and 02-0948 (PCD)). One hundred µl of samples were prepared with the Filter Aided Sample Preparation (FASP) method. Briefly, samples were denatured with 800 µl of 6M GuHCl (pH 8.0) and reduced by adding DTT to a

final concentration of 20 mM for 1 hour at 65□. After reduction, samples were alkylated with

iodoacetamide (I1149, Sigma) at a final concentration of 50 mM for 1 hour at 25□ in the dark.

Samples were centrifuged at 14,000 g for 10 min, and the filter was washed twice with50 mM

ammonium hydrogen carbonate (NH_4HCO_3). 0.5 µg modified trypsin (proteomics grade, T6567,

Sigma) was added and samples were incubated for 18 hours at 37□. The peptides solutions were

concentrated by vacuum centrifugation and dissolved in 30 µL 0.1 % formic acid. Mass

spectrometry was performed with a Dionex ultimate 3000 RSLCnano system coupled to a hybrid

quadrupole orbitrap mass spectrometer with a Nano spray source (Thermo Scientific Q-

Exactive). Samples (3 µL) were loaded into a trap column (Acclaim PepMap 100, 100 µm x 2

cm, nanoViper C18 5 µm 100 Å), at 5 µL/min with aqueous solution containing 0.1 % (v/v)

trifluoroacetic acid and 2 % acetonitrile. The column used for peptides separation was an

Acclaim PepMap RSLC, 75 µm x 15 cm, nanoViper C18 2 µm 100 Å. LC – MS runs were 120

minutes long, with two liquid phases: 97% water 3% Acetonitrile 0.1% Formic Acid (buffer A)

and Acetonitrile 0.1% Formic Acid (buffer B). The gradients (% of buffer B) of the run (120

minutes) applied were; 0 to 7 minutes: 4%; 7 to 90 minutes: 30%; 90 to 105 minutes: 80%; and

105 to 120 minutes: 4%.

Data were acquired at a resolution of 70,000 at m/z 200, target AGC value of 5e5, maximum fill

times of 200 ms, a multiplex degree of 6 with an isolation width of 3 m/z.

Proteins were identified by searching against the most current human database with Proteome

Discover 1.4, setting a maximum of two missed cleavage sites for trypsin, oxidation of

methionine (+15.995 Da) as a dynamic modification, and carbamidomethylation of cysteine

(+57.0210 Da) as static a modification. Proteins were quantified using Scaffold (Proteome

Software), using the normalized total precursor intensity, with 95% probability by the Scaffold Local FDR algorithm.

PRR4 and WGA staining for airway mucus flakes

Mucus flakes were obtained from BALF from healthy non-smoker adult subjects and school-age CF subjects as previouswily reported under IRB-approved protocol at UNC Chapel Hill (*54*). All studies were approved by the UNC Ethics Committee and informed consents from the subjects or parents obtained prior to any study procedures. BALF (10 µl) was applied to glass microscopy slides using a cytocentrifuge (StatSpin CytoFuge 2, Beckman Coulter). Slides were fixed with 10% neutral buffered formalin, washed with PBS, and blocked with 3% bovine serum albumin at RT for 1 hour. PRR4 and mucus were immunohistochemically stained by PRR4 antibody (1:1000) and biotinylated WGA (BK-1000, Vector Laboratories, 1:1000), respectively, at 4°C overnight. Slides were washed with PBS three times, followed by incubation with secondary antibodies (Alexa Fluor® Plus 488-conjugated donkey anti-rabbit IgG (A32795, Invitrogen, 1:1000) and 594-conjugated streptavidin (U0291, Abnova, 1:1000)) and DAPI (D3571, Invitrogen, 1:1000) at RT for 1 hour. The slides were then washed with PBS and mounted with Fluorsave mounting media (345789, Millipore). The staining images were obtained with an Olympus FV1000 confocal microscope using a 20x objective and uniform settings. Average PRR4 intensity was normalized to overall mucus signal by dividing by the average WGA intensity.

Validation of mouse anti-MUC5B antiserum for immunostaining

The mouse anti-MUC5B antiserum raised in-house was not previously published. As a validation, we performed immunohistochemistry of MUC5B for human tracheal serial sections with the antiserum (M-25, 1:1000) and previously established commercial rabbit anti-MUC5B antibody (H-300, Santa Cruz, 1:1000). Staining with these antibody and antiserum appeared identical, confirming the sensitivity and specificity of the novel antiserum (Fig.E4B).

Production of recombinant PRR4 protein

PRR4 cDNA was obtained from ORF clone plasmid (OHu15631, GenScript). PRR4 cDNA without a signal peptide sequence (corresponding to amino acids 17 to 134) was cloned into the pM-secSUMOstar Vector (7121, LifeSensors). SUMOstar-PRR4 vectors were transfected into Expi293 cells (1 mg of DNA per liter of transfection) per manufacturer instructions, with culture supernatants harvested and sterile-filtered on day 3 post-transfection. Supernatants were then concentrated and buffer-exchanged by tangential-flow filtration into TALON-binding buffer (50 mM sodium phosphate pH 7.4, 500 mM NaCl, 5 mM imidazole, 0.05% sodium azide, 10% glycerol), then loaded onto hand-packed 1 ml TALON (635507, Takara Bio) columns recirculating overnight at 1.5 ml/min. Twenty-four hours later, columns were changed to flow-through, and run until all supernatant was loaded. Columns were then washed with 20 column volumes binding buffer, followed by binding buffer with 10 mM imidazole. Proteins were then eluted with 20 CV binding buffer supplemented with 150 mM imidazole. Fractions were assessed by SDS-PAGE, and those containing bands corresponding to His-SUMO-tagged PRR4 (10 mM wash and elution) were pooled. Pooled fractions were then dialyzed against PBS with 2 mM DTT (3.5 kDa MWCO). SUMOstar protease (SP4110, LifeSensors; 10 units/mg substrate) was added to dialyzed protein and the reaction was incubated overnight at 4°C to cleave the His-

SUMO tags. Cleaved proteins were then passed back over the 1 ml TALON columns to separate the His-SUMO tags from the cleaved PRR4 proteins. Subtractive columns were washed and eluted as described above. Fractions were again assessed by SDS-PAGE to confirm separation, with flow-through and 5 mM imidazole fractions pooled, concentrated, and dialyzed into PBS (3.5 kDa MWCO). The final purified protein was confirmed to be PRR4 by both western blotting and mass spectrometry.

Confirmation of no degradative effect for PRR4 by neutrophil elastase

To quantitate the potential effects of proteolysis on the proteomics results, endotracheal tube (ETT) mucus samples were treated with designated quantities of HNE (10-50 ug/ml, 324681, Sigma). ETT mucus was obtained as previously reported from endotracheal tube tips following post-surgical extubation at UNC General Hospital under UNC IRB approved protocols (55). Aliquots of ETT mucus (500uL) were treated with 10, 20 and 50 ug of HNE and rotated/mixed for 30 minutes, and then incubated at 37□ for overnight. PBS was added to the control. PRR4, MUC5B and MUC5AC peptide intensities were measured by mass spectrometry as described above. (Fig.E4b).

Statistical analyses

Statistical analyses and figure creation were performed with GraphPad Prism 9.0.2. Each single measured/calculated value and mean ± SEM were plotted in the graphs unless otherwise specified. Statistical methods used were described in the Figure Captions. Statistical tests were two-tailed whenever appropriate.

Supplementary Materials

References

1. R. C. Boucher, Muco-Obstructive Lung Diseases. *The New England journal of medicine* **380**, 1941-1953 (2019).
2. D. A. Stoltz, D. K. Meyerholz, M. J. Welsh, Origins of cystic fibrosis lung disease. *The New England journal of medicine* **372**, 351-362 (2015).
3. S. M. Rowe, S. Miller, E. J. Sorscher, Cystic fibrosis. *The New England journal of medicine* **352**, 1992-2001 (2005).
4. H. Matsui, B. R. Grubb, R. Tarran, S. H. Randell, J. T. Gatzy, C. W. Davis, R. C. Boucher, Evidence for periciliary liquid layer depletion, not abnormal ion composition, in the pathogenesis of cystic fibrosis airways disease. *Cell* **95**, 1005-1015 (1998).
5. B. Button, L. H. Cai, C. Ehre, M. Kesimer, D. B. Hill, J. K. Sheehan, R. C. Boucher, M. Rubinstein, A periciliary brush promotes the lung health by separating the mucus layer from airway epithelia. *Science (New York, N.Y.)* **337**, 937-941 (2012).
6. H. M. Leung, S. E. Birket, C. Hyun, T. N. Ford, D. Cui, G. M. Solomon, R. J. Shei, A. T. Adewale, A. R. Lenzie, C. M. Fernandez-Petty, H. Zheng, J. H. Palermo, D. Y. Cho, B. A. Woodworth, L. M. Yonker, B. P. Hurley, S. M. Rowe, G. J. Tearney, Intranasal micro-optical coherence tomography imaging for cystic fibrosis studies. *Science translational medicine* **11**, (2019).
7. S. E. Birket, K. K. Chu, L. Liu, G. H. Houser, B. J. Diephuis, E. J. Wilsterman, G. Dierksen, M. Mazur, S. Shastry, Y. Li, J. D. Watson, A. T. Smith, B. S. Schuster, J. Hanes, W. E. Grizzle, E. J. Sorscher, G. J. Tearney, S. M. Rowe, A functional anatomic defect of the cystic fibrosis airway. *American journal of respiratory and critical care medicine* **190**, 421-432 (2014).
8. D. B. Hill, R. F. Long, W. J. Kissner, E. Atieh, Ian C. Garbarine, M. R. Markovetz, N. C. Fontana, M. Christy, M. Habibpour, R. Tarran, M. G. Forest, R. C. Boucher, B. Button, Pathological mucus and impaired mucus clearance in cystic fibrosis patients result from increased concentration, not altered pH. *European Respiratory Journal* **52**, 1801297 (2018).
9. A. G. Henderson, C. Ehre, B. Button, L. H. Abdullah, L. H. Cai, M. W. Leigh, G. C. DeMaria, H. Matsui, S. H. Donaldson, C. W. Davis, J. K. Sheehan, R. C. Boucher, M. Kesimer, Cystic fibrosis airway secretions exhibit mucin hyperconcentration and increased osmotic pressure. *The Journal of clinical investigation* **124**, 3047-3060 (2014).

10. C. R. Esther, Jr., M. S. Muhlebach, C. Ehre, D. B. Hill, M. C. Wolfgang, M. Kesimer, K. A. Ramsey, M. R. Markovetz, I. C. Garbarine, M. G. Forest, I. Seim, B. Zorn, C. B. Morrison, M. F. Delion, W. R. Thelin, D. Villalon, J. R. Sabater, L. Turkovic, S. Ranganathan, S. M. Stick, R. C. Boucher, Mucus accumulation in the lungs precedes structural changes and infection in children with cystic fibrosis. *Science translational medicine* **11**, (2019).

11. M. Boon, S. E. Verleden, B. Bosch, E. J. Lammertyn, J. E. McDonough, C. Mai, J. Verschakelen, M. Kemner-van de Corput, H. A. Tiddens, M. Proesmans, F. L. Vermeulen, E. K. Verbeken, J. Cooper, D. E. Van Raemdonck, M. Decramer, G. M. Verleden, J. C. Hogg, L. J. Dupont, B. M. Vanaudenaerde, K. De Boeck, Morphometric Analysis of Explant Lungs in Cystic Fibrosis. *American journal of respiratory and critical care medicine* **193**, 516-526 (2016).

12. A. Ermund, L. N. Meiss, B. Dolan, A. Bähr, N. Klymiuk, G. C. Hansson, The mucus bundles responsible for airway cleaning are retained in cystic fibrosis and by cholinergic stimulation. *Eur Respir J* **52**, 1800457 (2018).

13. A. J. Fischer, M. I. Pino-Argumedo, B. M. Hilkin, C. R. Shanrock, N. D. Gansemer, A. L. Chaly, K. Zarei, P. D. Allen, L. S. Ostedgaard, E. A. Hoffman, D. A. Stoltz, M. J. Welsh, M. H. Abou Alaiwa, Mucus strands from submucosal glands initiate mucociliary transport of large particles. *JCI insight* **4**, (2019).

14. A. Ermund, L. N. Meiss, A. M. Rodriguez-Pineiro, A. Bahr, H. E. Nilsson, S. Trillo-Muyo, C. Ridley, D. J. Thornton, J. J. Wine, H. Hebert, N. Klymiuk, G. C. Hansson, The normal trachea is cleaned by MUC5B mucin bundles from the submucosal glands coated with the MUC5AC mucin. *Biochemical and biophysical research communications* **492**, 331-337 (2017).

15. L. S. Ostedgaard, T. O. Moninger, J. D. McMenimen, N. M. Sawin, C. P. Parker, I. M. Thornell, L. S. Powers, N. D. Gansemer, D. C. Bouzek, D. P. Cook, D. K. Meyerholz, M. H. Abou Alaiwa, D. A. Stoltz, M. J. Welsh, Gel-forming mucins form distinct morphologic structures in airways. *Proceedings of the National Academy of Sciences* **114**, 6842-6847 (2017).

16. M. J. Hoegger, A. J. Fischer, J. D. McMenimen, L. S. Ostedgaard, A. J. Tucker, M. A. Awadalla, T. O. Moninger, A. S. Michalski, E. A. Hoffman, J. Zabner, D. A. Stoltz, M. J. Welsh, Impaired mucus detachment disrupts mucociliary transport in a piglet model of cystic fibrosis. *Science (New York, N.Y.)* **345**, 818-822 (2014).

17. Y. Xie, L. Lu, X. X. Tang, T. O. Moninger, T. J. Huang, D. A. Stoltz, M. J. Welsh, Acidic Submucosal Gland pH and Elevated Protein Concentration Produce Abnormal Cystic Fibrosis Mucus. *Developmental cell*, (2020).

18. N. S. Joo, H.-J. Cho, M. Khansaheb, J. J. Wine, Hyposecretion of fluid from tracheal submucosal glands of CFTR-deficient pigs. *The Journal of clinical investigation* **120**, 3161-3166 (2010).

19. N. S. Joo, T. Irokawa, R. C. Robbins, J. J. Wine, Hyposecretion, not hyperabsorption, is the basic defect of cystic fibrosis airway glands. *J Biol Chem* **281**, 7392-7398 (2006).

20. N. S. Joo, D. J. Lee, K. M. Winges, A. Rustagi, J. J. Wine, Regulation of antiprotease and antimicrobial protein secretion by airway submucosal gland serous cells. *J Biol Chem* **279**, 38854-38860 (2004).

21. N. S. Joo, T. Irokawa, J. V. Wu, R. C. Robbins, R. I. Whyte, J. J. Wine, Absent secretion to vasoactive intestinal peptide in cystic fibrosis airway glands. *J Biol Chem* **277**, 50710-50715 (2002).

22. J. H. Widdicombe, J. J. Wine, Airway Gland Structure and Function. *Physiological reviews* **95**, 1241-1319 (2015).

23. S. K. Inglis, M. R. Corboz, A. E. Taylor, S. T. Ballard, Effect of anion transport inhibition on mucus secretion by airway submucosal glands. *Am J Physiol* **272**, L372-377 (1997).

24. J. L. Cooper, P. M. Quinton, S. T. Ballard, Mucociliary transport in porcine trachea: differential effects of inhibiting chloride and bicarbonate secretion. *Am J Physiol Lung Cell Mol Physiol* **304**, L184-190 (2013).

25. S. K. Inglis, M. R. Corboz, S. T. Ballard, Effect of anion secretion inhibitors on mucin content of airway submucosal gland ducts. *Am J Physiol* **274**, L762-766 (1998).

26. S. T. Ballard, L. Trout, Z. Bebok, E. J. Sorscher, A. Crews, CFTR involvement in chloride, bicarbonate, and liquid secretion by airway submucosal glands. *The American journal of physiology* **277**, L694-699 (1999).

27. M. Rubinstein, R. H. Colby, *Polymer Physics*. (Oxford University Press, Oxford, 2003).

28. B. Button, H. P. Goodell, E. Atieh, Y. C. Chen, R. Williams, S. Shenoy, E. Lackey, N. T. Shenkute, L. H. Cai, R. G. Dennis, R. C. Boucher, M. Rubinstein, Roles of mucus adhesion and cohesion in cough clearance. *Proceedings of the National Academy of Sciences of the United States of America* **115**, 12501-12506 (2018).

29. S. Jayaraman, N. S. Joo, B. Reitz, J. J. Wine, A. S. Verkman, Submucosal gland secretions in airways from cystic fibrosis patients have normal [Na(+)] and pH but elevated viscosity. *Proceedings of the National Academy of Sciences of the United States of America* **98**, 8119-8123 (2001).

30. A. J. Fischer, K. L. Goss, T. E. Scheetz, C. L. Wohlford-Lenane, J. M. Snyder, P. B. McCray, Jr., Differential gene expression in human conducting airway surface epithelia and submucosal glands. *American journal of respiratory cell and molecular biology* **40**, 189-199 (2009).

31. T. F. Warner, E. A. Azen, Proline-rich proteins are present in serous cells of submucosal glands in the respiratory tract. *The American review of respiratory disease* **130**, 115-118 (1984).

32. N. S. Joo, I. A. Evans, H. J. Cho, I. H. Park, J. F. Engelhardt, J. J. Wine, Proteomic analysis of pure human airway gland mucus reveals a large component of protective proteins. *PloS one* **10**, e0116756 (2015).

33. C. Jenkinson, J. Earl, P. Ghaneh, C. Halloran, A. Carrato, W. Greenhalf, J. Neoptolemos, E. Costello, Biomarkers for early diagnosis of pancreatic cancer. *Expert Review of Gastroenterology & Hepatology* **9**, 305-315 (2015).

34. L. Trout, M. King, W. Feng, S. K. Inglis, S. T. Ballard, Inhibition of airway liquid secretion and its effect on the physical properties of airway mucus. *The American journal of physiology* **274**, L258-L263 (1998).

35. C. J. Martens, S. K. Inglis, V. G. Valentine, J. Garrison, G. E. Conner, S. T. Ballard, Mucous solids and liquid secretion by airways: studies with normal pig, cystic fibrosis human, and non-cystic fibrosis human bronchi. *American journal of physiology. Lung cellular and molecular physiology* **301**, L236-246 (2011).

36. L. Trout, J. T. Gatzy, S. T. Ballard, Acetylcholine-induced liquid secretion by bronchial epithelium: role of Cl- and HCO3- transport. *The American journal of physiology* **275**, L1095-1099 (1998).

37. X. Luan, G. Belev, J. S. Tam, S. Jagadeeshan, N. Hassan, P. Gioino, N. Grishchenko, Y. Huang, J. L. Carmalt, T. Duke, T. Jones, B. Monson, M. Burmester, T. Simovich, O. Yilmaz, V. A. Campanucci, T. E. Machen, L. D. Chapman, J. P. Ianowski, Cystic fibrosis swine fail to secrete airway surface liquid in response to inhalation of pathogens. *Nature communications* **8**, 786 (2017).

38. A. McShane, J. Bath, A. M. Jaramillo, C. Ridley, A. A. Walsh, C. M. Evans, D. J. Thornton, K. Ribbeck, Mucus. *Curr Biol* **31**, R938-r945 (2021).

39. S. K. Inglis, M. R. Corboz, A. E. Taylor, S. T. Ballard, In situ visualization of bronchial submucosal glands and their secretory response to acetylcholine. *The American journal of physiology* **272**, L203-210 (1997).

40. L. Reid, Measurement of the bronchial mucous gland layer: a diagnostic yardstick in chronic bronchitis. *Thorax* **15**, 132-141 (1960).

41. L. S. Ostedgaard, M. P. Price, K. M. Whitworth, M. H. Abou Alaiwa, A. J. Fischer, A. Warrier, M. Samuel, L. D. Spate, P. D. Allen, B. M. Hilkin, G. S. Romano Ibarra, M. E. Ortiz Bezara, B. J. Goodell, S. E. Mather, L. S. Powers, M. R. Stroik, N. D. Gansemer, C. E. Hippee, K. Zarei, J. A. Goeken, T. R. Businga, E. A. Hoffman, D. K. Meyerholz, R. S. Prather, D. A. Stoltz, M. J. Welsh, Lack of airway submucosal glands impairs respiratory host defenses. *eLife* **9**, (2020).

42. D. B. Hill, P. A. Vasquez, J. Mellnik, S. A. McKinley, A. Vose, F. Mu, A. G. Henderson, S. H. Donaldson, N. E. Alexis, R. C. Boucher, M. G. Forest, A biophysical basis for mucus solids concentration as a candidate biomarker for airways disease. *PloS one* **9**, e87681 (2014).

43. H. P. Goodell, S. K. Shenoy, N. T. Shenkute, E. Lackey, R. G. Dennis, B. Button, Adhesive and Cohesive Peel Force Measurement of Human Airway Mucus. *Bio Protoc* **9**, e3287 (2019).

44. L. Russel, S. Burguet, Ultrastructure of leydig cells as revealed by secondary tissue treatment with a ferrocyanide-osmium mixture. *Tissue Cell* **9**, 751-766 (1977).

45. E. S. Reynolds, The use of lead citrate at high pH as an electron-opaque stain in electron microscopy. *J Cell Biol* **17**, 208-212 (1963).

46. B. A. Nerger, P. T. Brun, C. M. Nelson, Marangoni flows drive the alignment of fibrillar cell-laden hydrogels. *Science advances* **6**, eaaz7748 (2020).

47. M. L. Fulcher, S. Gabriel, K. A. Burns, J. R. Yankaskas, S. H. Randell, Well-differentiated human airway epithelial cell cultures. *Methods in molecular medicine* **107**, 183-206 (2005).

48. K. Okuda, G. Chen, D. B. Subramani, M. Wolf, R. C. Gilmore, T. Kato, G. Radicioni, M. Kesimer, M. Chua, H. Dang, A. Livraghi-Butrico, C. Ehre, C. M. Doerschuk, S. H. Randell, H. Matsui, T. Nagase, W. K. O'Neal, R. C. Boucher, Localization of Secretory Mucins MUC5AC and MUC5B in Normal/Healthy Human Airways. *American journal of respiratory and critical care medicine* **199**, 715-727 (2019).

49. M. L. Fulcher, S. H. Randell, Human nasal and tracheo-bronchial respiratory epithelial cell culture. *Methods in molecular biology (Clifton, N.J.)* **945**, 109-121 (2013).

50.   K. Aoki, M. Porterfield, S. S. Lee, B. Dong, K. Nguyen, K. H. McGlamry, M. Tiemeyer, The diversity of O-linked glycans expressed during Drosophila melanogaster development reflects stage- and tissue-specific requirements for cell signaling. *The Journal of biological chemistry* **283**, 30385-30400 (2008).

51.   T. Kumagai, T. Katoh, D. B. Nix, M. Tiemeyer, K. Aoki, In-gel β-elimination and aqueous-organic partition for improved O- and sulfoglycomics. *Analytical chemistry* **85**, 8692-8699 (2013).

52.   N. Mehta, M. Porterfield, W. B. Struwe, C. Heiss, P. Azadi, P. M. Rudd, M. Tiemeyer, K. Aoki, Mass Spectrometric Quantification of N-Linked Glycans by Reference to Exogenous Standards. *Journal of proteome research* **15**, 2969-2980 (2016).

53.   W. S. York, S. Agravat, K. F. Aoki-Kinoshita, R. McBride, M. P. Campbell, C. E. Costello, A. Dell, T. Feizi, S. M. Haslam, N. Karlsson, K. H. Khoo, D. Kolarich, Y. Liu, M. Novotny, N. H. Packer, J. C. Paulson, E. Rapp, R. Ranzinger, P. M. Rudd, D. F. Smith, W. B. Struwe, M. Tiemeyer, L. Wells, J. Zaia, C. Kettner, MIRAGE: the minimum information required for a glycomics experiment. *Glycobiology* **24**, 402-406 (2014).

54.   T. L. Noah, H. R. Black, P. W. Cheng, R. E. Wood, M. W. Leigh, Nasal and bronchoalveolar lavage fluid cytokines in early cystic fibrosis. *The Journal of infectious diseases* **175**, 638-647 (1997).

55.   M. R. Markovetz, D. B. Subramani, W. J. Kissner, C. B. Morrison, I. C. Garbarine, A. Ghio, K. A. Ramsey, H. Arora, P. Kumar, D. B. Nix, T. Kumagai, T. M. Krunkosky, D. C. Krause, G. Radicioni, N. E. Alexis, M. Kesimer, M. Tiemeyer, R. C. Boucher, C. Ehre, D. B. Hill, Endotracheal tube mucus as a source of airway mucus for rheological study. *American journal of physiology. Lung cellular and molecular physiology* **317**, L498-l509 (2019).

Acknowledgements

The authors thank the UNC CF Center Tissue Procurement and Cell Culture Core for providing human airway tissues and primary cells, and the UNC Animal Histology Core for paraffin embedding and sectioning the tissues. We also thank Dr. Hong Dang for statistical analyses and Eric Roe for editing the manuscript.

Fundings

This research was supported by grants from National Institutes of Health (NIDDK UH3 HL123645, P01 HL110873, R01 HL136961, P30 DK 065988, P01 HL108808 and R01HL125280), and Cystic Fibrosis Foundation (BOUCHE15R0 to R.C.B., BUTTON19G0 to B.B., FREEMA19G0 to R.F, OKUDA19I0 and OKUDA20G0 to K.O., KATO20I0 to T.K. and HILL20Y2-OUT and HILL19G0 to D.B.H) and a research grant from Cystic Fibrosis Research Incorporation to K.O.. T.K. was supported by the Senior Research Training Fellowship by the American Lung Association (RT-575362). M.R.M. was supported by the Postdoctoral Research

Fellowship by the Cystic Fibrosis Foundation (MARKOV18F0). Glycomic analysis of airway mucins was supported by grants from the National Institutes of Health (NIGMS P41GM103490 and Common Fund U01GM125267 to M.T.). SEM of SMG mucus strands was performed at the Chapel Hill Analytical and Nanofabrication Laboratory, CHANL, a member of the North Carolina Research Triangle Nanotechnology Network, RTNN, which was supported by the National Science Foundation, Grant ECCS-1542015, as part of the National Nanotechnology Coordinated Infrastructure, NNCI. The UNC Hooker Imaging Core Facility, the Microscopy Services Laboratory, and the Protein Expression & Purification Core Facility were all supported in part by P30 CA016086 Cancer Center Core Support Grant to the UNC Lineberger Comprehensive Cancer Center.

Author contributions

T.K., S.T.B., B.B. and R.C.B. conceived the project and designed experiments. S.H.R. provided human airway tissues and primary cells. T.K., G.V.S. and S.T.B. collected pig SMG mucus. T.K. collected human SMG mucus. T.K., H.P.G, H.T., H.T.L and B.B. performed %solids, osmotic pressures, and cohesion measurements. G.V.S. performed solubility and light scattering assays. K.A., M.P. and M.T. performed glycomics analyses. M.J.P. and R.F. acquired SEM and immunofluorescent images of SMG mucus strands and characterized their structural properties. K.A.B. and K.K.W. acquired EM images of porcine and human airway tissues. K.O. prepared human histology sections and RNA samples from human airway tissues. R.C.G performed RNA *in situ* hybridization. T.K. and S.M.B.C performed immunostaining and RT-qPCR. G.R. and M.K. performed proteomic analysis. M.R.M, C.B.M. and D.B.H performed analyses of mucus flakes. C.B.M and C.E. provided mouse anti-MUC5B antiserum. L.J.F. performed PRR4 overexpression and purification. T.N. and M.R. provided feedback to overall work. T.K., B.B.,

W.K.O. and R.C.B. analyzed and interpreted the data. B.B. and R.C.B. supervised the project. T.K., R.F., W.K.O, B.B. and R.C.B drafted and finalized the manuscript.

Competing interests

Authors declare that they have no competing interests.

Data and materials availability

All data are available in the main text or the supplementary materials.

Figures

Figure 1. Studies of anion transport-inhibited porcine airways

 (A) Solids concentration (% solids) of porcine submucosal gland (SMG) mucus after treatment with acetylcholine (Ach) without or with dimethylamiloride (DMA), bumetanide (Bum), or the combination of DMA and Bum. N = 6 to 8 per group. Welch's ANOVA test followed by Dunnett's multiple comparison test. **(B)** Osmotic pressure (in pascals, Pa) of porcine SMG mucus after treatment with Ach without or with DMA, Bum, or DMA/Bum. N = 6 to 8 per group. Welch's ANOVA test followed by Dunnett's multiple comparison test. **(C)** Linear correlation between % solids and osmotic pressure of pig SMG mucus. N = 29. **(D)** Transmission electron microscope (TEM) images of porcine SMG ducts after Ach treatment without or with DMA and Bum. **(E)** Quantitation of SMG (**i**) cilia height and (**ii**) mucus retention in ducts for each treatment condition. (**i**) N = 4 to 10 ducts from 3 airway specimens for each condition from 2 pigs. One to 8 regions of cilia height per duct were measured. Mean of cilia height for each duct was used for plot and statistical analyses. Welch's ANOVA test followed by Dunnett's

multiple comparison test. (**ii**) N = 4 to 10 ducts from 3 airway specimens for each condition from 2 pigs. One to 8 regions of mucus retention per duct were evaluated. Each score was used for plot and statistical analyses. N = 72 regions from 25 ducts in total. Kruskal-Wallis test followed by Dunn's multiple comparison.

Figure 2. Studies of non-CF and CF human submucosal glands (SMGs)

(**A**) Solids concentration (% solids) of human SMG mucus from non-CF (N = 30) and CF (N = 9) subjects. Welch's t-test. (**B**) Osmotic pressure (Pa) of human SMG mucus from non-CF (N = 11) and CF (N = 9) subjects. Welch's t-test. (**C**) Linear correlation between % solids and osmotic pressure of human SMG mucus. (**D**) Cohesive strength (mN/cm) of human SMG mucus from non-CF (N = 8) and CF (N = 5) subjects. Welch's t-test. (**E**) SMG ducts of healthy control vs CF airways after acetylcholine treatment captured by transmission electron microscopy (TEM). Scale bar = 2μm. (**F**) Quantitation of human SMG (**i**) cilia height and (**ii**) mucus ductal retention in healthy control vs CF groups. (**i**) N = 15 ducts from 3 healthy control donors and N = 12 ducts from 3 CF donors. One to 7 regions of cilia height per duct were measured. Mean of cilia height for each duct was used for plot and statistical analyses. Welch's t-test. (**ii**) N = 37 regions from 15 healthy control ducts and N = 42 regions from 12 CF ducts. One to 8 regions of mucus retention per duct were evaluated depending on TEM image availability. Each score was used for plot and statistical analyses. Wilcoxon's signed rank test.

Figure 3. Submucosal gland mucus characteristics

(**A**) Solubility of human bronchial epithelial (HBE) cell culture mucus and human and porcine submucosal gland (SMG) mucus in phosphate buffered saline (PBS). (**B**) Radius of gyration (R_g)

of the soluble fraction of SMG mucus and HBE mucus determined by light-scattering. N = 10-15

for HBE, N=3 each for human and pig SMG mucus. Welch's ANOVA test (P = 0.1003). **(C)** (**i**)

Immunofluorescent staining of MUC5B and MUC5AC in non-CF SMG mucus strands. Scale

bar = 10μm. (**ii**) Intensity of MUC5B and MUC5AC signals on the yellow line in the left panel.

Figure 4. Glycomics analyses of HBE vs SMG mucus

(A) Glycan abundance of less complex (Core 1 and 3) glycans in HBE vs SMG mucus. Unpaired

t-test. N = 3 for each. (**B**) Abundance of glycans with shorter (≤ 3) monosaccharides in HBE vs

SMG mucus. Umpaired t-test. N = 3 for each.

Figure 5. Comparison of SMG mucus strands from non-CF and CF subjects

(A) Relative number of strands found within SMG samples. Values were normalized by sample

volume, imaged, and divided by the number of mean healthy strands. **(B)** Mean strand width of

non-CF vs CF SMG mucus. **(C)** Mean strand length of non-CF vs CF SMG mucus. **(D)** Scanning

electron microscopy (SEM) images of SMG strands from non-CF subjects. Inset falsely colored

according to angular alignment of pixels quantified through OrientationJ. **(E)** SEM images of CF

SMG strands. Inset falsely colored according to angular alignment of pixels. Samples from N = 9

(non-CF) and N = 4 (CF) donors. One to 5 strands per donor were measured. Mean of each donor

was plotted. (**F,G**) Alignment of non-CF vs CF SMG mucus. **F** shows the histogram of relative

alignment angles of SEM images in panels **D** and **E**. Non-CF sample shows increased alignment

of surface features whereas CF strand does not. **G** shows the quantification of alignment fraction

for each clinical sample based on decreased numbers of bins of panels **D** and **E** to account for

slight deviations in angle from strand directionality. N = 3 each for non-CF and CF. (**A**) Mann-Whitney test. (**B**, **C**, and **G**) Unpaired t-test.

Mann-Whitney test was used for the data for Fig.5A. Samples from N = 9 (non-CF) and N = 4 (CF) donors. P = 0.0238. Unpaired t-test was used for the data for Fig.5B. One to 5 strands per donor was measured for N = 8 (non-CF) and N=4 (CF) donors. Mean of each donor was plotted. P = 0.4817. Unpaired t-test was used for the data for Fig.5C. One to 5 strands per donor was measured for N = 8 (non-CF) and N=4 (CF) donors. Mean of each donor was plotted. P = 0.4809. Unpaired t-test was used for the data for Fig.5G. One to 4 strands per donor was measured for N = 3 (non-CF) and N = 3 (CF) donors. Mean of each donor was plotted. P = 0.0262.

Figure 6. Distribution of PRR4 expression in airways

(**A, B**) RNA *in situ* hybridization (RNA-ISH) and immunohistochemistry (IHC) for proline-rich protein 4 (PRR4) in (**A**) healthy control and (**B**) cystic fibrosis (CF) human airways. Bars: 500μm (low power images) and 200μm (magnified images). (**C**) Immunofluorescent co-staining of PRR4 and MUC5B in human proximal airways. Bars = 200μm. (**D**) Immunofluorescent co-staining of PRR4 and MUC5B in human SMG mucus strands.

Figure 7. PRR4 levels in induced sputum and BALF samples

(**A,B**) Peptide intensities of (**A**) PRR4 and (**B**) human neutrophil elastase in induced sputum samples from healthy donors (N = 18), CF (N = 20), non-CF bronchiectasis (NCFB) (N = 10)

and primary ciliary dyskinesia (PCD) (N = 9) measured by label-free quantitation (LFQ) utilizing mass-spectrometry. Kruskal-Wallis test followed by Dunn's multiple comparison. **(C)** PRR4 content in BAL-obtained flakes. (**i**) Immunofluorescent co-staining of WGA and PRR4 for healthy control and CF mucus flakes (Bars = 100μm) and (**ii**) ratio of staining intensity of PRR4 to WGA in mucus flakes. N = 3 for healthy control, N = 6 for CF. Mann-Whitney test.

Figure 8. Identification and quantitation of serous and mucous cells of SMGs

(**i**) Immunofluorescent staining of PRR4 and MUC5B for SMGs in healthy control vs CF tissues. Scale bar = 100μm. (**ii**) Quantification of serous cell (PRR4) vs mucous cell (MUC5B) signals in SMG in healthy control vs CF. N = 9 for healthy control, N = 6 for CF. Welch's t-test.

Figure S1. Glycomics analyses of human bronchial epithelial (HBE) and submucosal gland (SMG) mucus

(A) Glycan types detected in HBE and SMG mucus samples. **(B)** Relative abundance of each glycan type in HBE vs SMG mucus from the matched donors. **(C)** Abundance of glycans with 2 to 10 monosaccharides per glycan in HBE vs SMG mucus from matched donors. **(D)** Abundance of more complex (Core 2 and 4) in HBE vs SMG mucus. N = 3 for each. Unpaired t-test.

Figure S2. Sulfated glycan analyses of human bronchial epithelial (HBE) and submucosal gland (SMG) mucus

(A) Sulfated glycan types detected in HBE and SMG mucus samples. **(B)** Relative abundance of each sulfated glycan type in HBE vs SMG mucus from the matched donors. N = 2.

Figure S3. PRR4 distribution in disease airways

Immunohistochemistry for PRR4 in airways from (**A**) non-CF bronchiectasis (NCFB), (**B**) primary ciliary dyskinesia (PCD) and (**C**) asthma subjects. Bars: 500μm (low power images) and 200μm (magnified images).

Figure S4. Region-specific localization of submucosal gland candidate biomarkers

RNA *in situ* hybridization (RNA-ISH) and immunohistochemistry (IHC) for submucosal gland markers: (**A,D**) lactoferrin (LTF); (**B,E**) zinc-α2-glycoprotein (ZAGP1/ZAG); and (**C,F**) lysozyme (LYZ) in (**A-C**) normal and (**D-F**) CF human airways. Bars: 500μm (low power images) and 200μm (magnified images).

Figure S5. Region specific mRNA expression of submucosal gland (SMG) markers in healthy airways

Quantitation of mRNA expression of candidate SMG marker genes including proline-rich protein family genes (*PRR4*, *PRB1*, *PRB2*, *PRB4* and *PRH2*), *LTF*, *AZGP1* and *LYZ*. A distal airway marker (*SFTPB*) was measured as a validation for airway region specificity. N = 9 donors. Friedman tests followed by Dunn's multiple comparison tests.

Figure S6. Validatory experiment results for anti-MUC5B antisera raised in-house

Immunohistochemistry for MUC5B with mouse antisera raised in-house (M-25) and rabbit polyclonal antibody (H-300) in healthy human trachea. Bar = 200μm.

Figure S7. Absence of PRR4 degradation by co-incubation with human neutrophoil elastase

Integrated intensity of MUC5B, MUC5AC and PRR4 peptides after incubation with human neutrophil elastase at designated concentrations for overnight as measured by mass spectroscopy, showing PRR4 resistance to HNE.

Supplementary Table 1. Solubility of SMG mucus after addition of IGEPAL

All the PBS-insoluble components of human and pig SMG mucus exhibited full or partial solubility after adding IGEPAL.

Figure 1.

Figure 2.

Figure 3.

A

B

C
i
| MUC5B | MUC5AC |

Merge

ii

Figure 4.

A

B

Figure 5.

Figure 6.

A PRR4 Healthy control   RNA-ISH   IHC

B PRR4 CF   RNA-ISH   IHC

C PRR4   MUC5B   Merge
DAPI / PRR4 / MUC5B

D PRR4   MUC5B   Merge
PRR4 / MUC5B

Figure 7.

Figure 8.

i

Healthy control      CF

DAPI / PRR4 / MUC5B

ii

Figure S1.

A

B

Figure S2.

Figure S3.

A PRR4
   NCFB

B PRR4
   PCD

C PRR4
   Asthma

Figure S4.

Figure S5.

Figure S6.

M-25   H-300

Figure S7.