Altered synaptic connectivity in an *in vitro* human model of STXBP1 encephalopathy.

Faye McLeod[1], Anna Dimtsi[1], David Lewis-Smith[1,2], Rhys Thomas[1,2], Gavin J Clowry[1*] and Andrew J Trevelyan[1*]

Author affiliations:

1 Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, NE2 4HH, United Kingdom

2 Department of Clinical Neurosciences, Royal Victoria Infirmary, Newcastle Upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, NE1 4LP, United Kingdom

Correspondence to: Faye McLeod

* Equal contribution

Full address: Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne, NE2 4HH, United Kingdom

E-mail: faye.mcleod@newcastle.ac.uk

Running title: STXBP1 encephalopathy *in vitro* model

Keywords: Human organotypic slices; synapses; STXBP1; subplate; neurodevelopment.

Abbreviations: HDBR = Human Developmental Biology Resource; STXBP1 = Syntaxin Binding protein 1; MUNC18-1 = Mammalian uncoordinated-18; pcw = post conception week; DIV = days *in vitro*; AAV = adeno-associated virus; shRNA = short hairpin RNA.

Abstract

Early infantile epileptic encephalopathies are devastating conditions, generally of genetic origin, but the pathological mechanisms often remain obscure. A major obstacle in this field of research is the difficulty of studying cortical brain development in humans, *in utero*. To address this, we established an *in vitro* assay to study the impact of gene variants on the developing human brain, using living organotypic cultures of the human subplate and neighbouring cortical regions, prepared from ethically sourced, 14-17 post conception week brain tissue (www.hdbr.org). We were able to maintain cultures for several months, during which time, the gross anatomical structures of the cortical plate, subplate and marginal zone persisted, while neurons continued to develop morphologically, and form new synaptic networks. This preparation thus permits the study of genetic manipulations, and their downstream effects, upon an intact developing human cortical network. We focused upon STXBP1 haploinsufficiency, which is among the most common genetic causes of infantile epileptic encephalopathy. This was induced using shRNA interference, leading to impaired synaptic function and a drop in the number of glutamatergic synapses. We thereby provide a critical proof-of-principle for how to study the impact of any gene of interest on the development of the human cortex.

Introduction

Approximately 1 in 2000 children in the UK develop epilepsy before three years of age from a monogenic cause[1]; of these, about a third will continue to have seizures into adulthood[2]. Knowing the genetic cause can aid clinical management, yet the pathological mechanisms remain obscure, at least in part because the aetiology often starts *in utero*, making their study extremely difficult. A critical phase of prenatal neocortical development is the transient appearance, and subsequent dissolution, of the subplate. This cortical region contains the earliest cortical synaptic network and forms the scaffold for subsequent neocortical development[3, 4]. It is therefore the site where synaptic malfunction will first impact.

Of particular interest is *Syntaxin-binding protein 1 (STXBP1)*, an essential pre-synaptic protein also called MUNC18-1[5], that is highly expressed in the prenatal developing brain[6]. Loss of function variants in *STXBP1*, typically termed 'STXBP1 encephalopathies', result in poor cognitive function and epilepsy in early development[7-9], but the underlying pathological mechanisms are still not fully understood. It is probable that the effects of STXBP1

haploinsufficiency arise prenatally, supported by the fact that STXBP1 is highly expressed in the developing subplate[6]. For early human development, however, the study of functional prenatal interactions presents a major practical challenge.

STXBP1 haploinsufficiency has been replicated in transgenic mouse models and is associated with a postnatal deficit in synaptic neurotransmission[10, 11]. Rodent models of this condition, though, are limited in one important respect, which is that the rodent subplate is very simple in organisation, compared with the human subplate[12], calling into question their relevance to human development. Likewise, human cerebral organoids are similarly limited, because, to date, there have been no descriptions of any organoid structures that accurately resembles the subplate as it exists *in utero*[13].

To overcome this critical gap in our experimental arsenal, we have developed a protocol for preparing living organotypic cultures of human neocortex, and its underlying subplate, from ethically approved samples of human fetal brain tissue, provided by the MRC-Wellcome Trust Human Developmental Biology Resource (HDBR, www.hdbr.org). Over time, neurons within the culture display evolving morphology, while the culture itself retains its gross anatomical structure. Importantly, the cultures are amenable to genetic manipulation, including knock-down of STXBP1. This was achieved using shRNA interference, delivered by AAV viral vectors, to mimic the disease-causing haploinsufficient state, and using viral vectors carrying a scrambled STXBP1 sequence as a control. STXBP1 knock-down resulted in altered synaptic stability and impaired spontaneous pre-synaptic neurotransmitter release, and changes in the numbers of synaptic terminals. We have thus developed a translatable model of early neocortical development for the study of critical genes that are suspected of causing neurodevelopmental conditions in humans.

Materials and methods

Detailed methods are provided in the supplementary material.

Human organotypic slice cultures

Human foetal brain tissue was acquired from the HDBR; (https://www.hdbr.org/) in Newcastle upon Tyne (Project 200428). Full consent was given in line with ethical approval from the Newcastle and North Tyneside NHS Health Authority Joint Ethics Committee (REC reference: 18/NE/0290). Cultures were prepared from 10 different foetal brains (three from 14 post-conception week (pcw), four 16 pcw and three 17 pcw). Slices were sectioned on a vibratome (280μm), placed inside 6-transwell plates containing culture media, which was changed every two days and the plates maintained at 37°C, 5% CO_2 in ambient O_2 and 90% humidity.

Viral vectors

Three adeno-associated viral (AAV) vectors were used (VectorBuilder):

1. Scrambled control shRNA - packaged into an AAV9 vector expressing the cytomegalovirus (CMV) promoter and enhanced green fluorescent protein (eGFP) to achieve a final titre of $5x10^{12}$ GC/ml. The scrambled target sequence was 5'-CCTAAGGTTAAGTCGCCCTCG-3'. Infected cells were referred to as 'control'.
2. STXBP1 shRNA - packaged into an AAV9 vector expressing the CMV promoter and eGFP to achieve a final titre of $5x10^{12}$ GC/ml. The STXBP1 target sequence was 5'TACTGAAGCACAAACATATAT-3'.
3. AAV9-CMV-GFP (final titre of $1x10^{13}$ GC/ml).

Infected cells were identified by their eGFP/GFP tag. At 0 DIV, 3-5 μl of viral vectors were added to the top of each slice. All experiments were conducted 1-4 weeks later.

Live imaging

Live imaging was performed using an inverted Zeiss LSM800 Airyscan confocal microscope. FM4-64 dye was utilised as previously described[14] and analysis was performed using Fiji software (NIH, RRID:SCR_002285).

Whole cell-patch clamp recordings

Slices were transferred into the recording chamber of an upright Leica DMLFSA fluorescent microscope fitted with Micro Control Instruments micromanipulators and continuously perfused at 34°C with oxygenated ACSF. AMPA receptor-mediated miniature glutamatergic post-synaptic currents were recorded at -60 mV in the presence of 100 nM TTX, 10 μM bicuculline, 50 μM AP-5. Miniature GABAergic post-synaptic currents were recorded at 0 mV in the presence of 100 nM TTX and 50 μM AP-5.

Immunohistochemistry

Staining was performed using the protocol optimised specifically for acute slices[15].

Image acquisition and analysis

Images were acquired on a Zeiss LSM800 Airyscan confocal microscope with fixed settings. The x5 objective (NA = 0.16) was used to acquire image stacks (z-step = 5μm) of general slice anatomy. The x10 objective was used to aid identification of the subplate. This region has an easily identifiable anatomy with a sparse cell population in comparison to the neighbouring cortical plate. Subsequently, x20 objective (NA = 0.8) with x2 zoom was used to acquire image stacks (z-step = 2μm) of the infected and non-infected subplate neurons and the x63 objective (NA = 1.4) to acquire image stacks (z-step = 0.25μm) of synaptic components. Three to four image stacks in the upper layer of the subplate were acquired per slice in a randomised manner. A final image resolution of 1024x1024 pixels was obtained.

The number as well as the length of primary subplate neuron processes were measured using Fiji software. Synaptic counts and integrated density (area × mean intensity) measurements were conducted using Zen 3.1 software (Zeiss; RRID:SCR_013672).

Western blot

Equal homogenate amounts were run on 10% SDS-PAGE gel and transferred onto PVDF membranes using BIO-RAD western blot apparatus.

Statistical analysis

Statistical analyses were performed on GraphPad Prism 9 (RRID:SCR_002798). We assessed data normality using Kolmogorov-Smirnov tests. The appropriate parametric test was used for all normally distributed data, including paired/unpaired t-tests, one-way ANOVAs and two-way ANOVAs with Tukey's multiple comparisons test and repeated measures. Statistical significance was accepted as * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$, and non-significant data were indicated as ns. All data in the results section are presented as $\pm$ SEM.

Data availability

All data generated or analysed and used in this study are available upon reasonable request.

Results

STXBP1 expression in the developing human brain

We created organotypic cortical slice cultures from 10 samples of fresh human foetal brain tissue, aged between 14-17 post-conception weeks (pcw) at the time of preparation. In all cases, the organotypic cultures retained the distinct and identifiable anatomical regions of the subplate, its neighbouring cortical plate, and the marginal zone (Figure 1A-B). Synapsin1 immunoreactivity revealed that pre-synaptic terminals were predominantly located in the upper layer of the subplate, at this stage of development (Figure 1A). The subplate was smallest in the 14 pcw samples (n = 3 biological samples; average depth = 308 $\pm$ 57 µm), becoming progressively larger in the cultures prepared from 16 (n = 4; 525 $\pm$ 11 µm) and 17 pcw (n = 3; 651 $\pm$ 33 µm) ($p<0.05$, one-way ANOVA; Figure 1B). STXBP1 showed a higher expression at 16-17 pcw, in comparison to 14 pcw (Figure 1B), and was observed near spine-like structures and along putative axons of subplate neurons (Figure 1D). Collectively these data show that the subplate becomes more prominent, and enriched in pre-synaptic proteins including STXBP1, over the 14-17 pcw period.

Synapse maturation in human cortical slice cultures

Given the relatively higher expression of STXBP1 in the subplate of 16–17 pcw tissue, we focused on this age for subsequent analysis of how the organotypic cultures continued to develop *in vitro*. To examine how neuronal morphology developed in the human cortical slice

cultures, we infected cultures with viral vectors carrying the eGFP construct. By 7 DIV, we were able to visualise the neurites of individual labelled neurons, and could subsequently identify the same neurons at 21 and 27 DIV. By doing this, we recorded a marked elaboration of the neurite processes over this time (Figure 1D).

The number of STXBP1 immunoreactive puncta in the subplate more than doubled, between 0 DIV and 27 DIV (16 pcw cultures increase from 4.1 ± 1.6 to 9.0 ± 0.4 puncta / 100 μm^3; 17 pcw cultures increase from 4.2 ± 0.6 to 13.3 ± 1.1 puncta / 100 μm^3; Figure 2A). Consistent with this, we found significant increases also in both glutamatergic synapses (vGlut1/Homer immunoreactivity; 16 pcw cultures increase from 2.7 ± 0.5 to 21.1 ± 1.7 puncta / 100 μm^3; 17 pcw cultures increase from 9.2 ± 0.9 to 30.3 ± 2.1 puncta / 100 μm^3; Figure 2B), and GABAergic synapses (vGAT/gephyrin immunoreactivity; 16 pcw, increase from 1.0 ± 0.4 to 3.7 ± 0.6 puncta / 100 μm^3; 17 pcw, increase from 0.6 ± 0.2 to 4.8 ± 0.4 puncta / 100 μm^3; Figure 2C). These data show there is maturation of neurons and a continued production of both glutamatergic and GABAergic synapses in the subplate of human cortical slice cultures at both 16 and 17 pcw with time spent *in vitro*.

Divergent regulation of synapses with reduced STXBP1 levels.

We next investigated the effects of manipulating the expression of STXBP1 in the human cortical slice cultures. We created a generic haploinsufficiency phenotype for STXBP1 ("knock-down") in our human cortical slice cultures using an AAV packaged *STXBP1* shRNA in 16 pcw tissue from two human cases (Figure 3). To control for any effects of the viral infection, we performed similar infections with a scrambled STXBP1 sequence, in age matched brain slices ("scrambled controls"). The shRNA achieved about 50% downregulation of STXBP1 at 14 DIV, compared with the scrambled control cultures prepared from the same brain tissue sample (Figure 3A). Within the subplate, STXBP1 immunoreactivity was reduced by approximately 30%, although there was no overall change in the absolute number of STXBP1 puncta (Supplementary Figure 1). Since each punctum was presumed to represent a synaptic terminal, this suggests that our shRNA design effectively reduces endogenous STXBP1 protein levels, without affecting the total number of STXBP1-positive synaptic terminals.

We found no difference in the size of the subplate between control (n = 10 slices, average depth = 529 ± 25 μm) and knock-down cultures (n = 10, average depth = 549 ± 21 μm; p>0.05, unpaired-test). Furthermore, the number and length of neurite processes in subplate neurons

measured at 14 DIV was unaffected (Figure 3B). Nonetheless, approximately 86% reduction in the number of glutamatergic synapses was found in *STXBP1* shRNA conditions within the subplate (Figure 3C). To evaluate the functional effects of these anatomical changes, we recorded glutamatergic currents. The frequency of synaptic events was decreased by 50% in cells expressing *STXBP1* shRNA (Figure 3D). In contrast, there was nearly an 80% increase in GABAergic synapse count in *STXBP1* shRNA conditions within the subplate (Figure 3E), compared to scrambled control with a 70% increase in the frequency of GABAergic post-synaptic currents (Figure 3F). Amplitude measurements of glutamatergic and GABAergic currents were unchanged (Figures 3D and 3F). Overall, these data imply that reduced STXBP1 levels does not alter the overall stability of primary subplate neuron neurites but modifies the synaptic stability during early subplate development at 16 pcw.

Impaired vesicle recycling with loss of STXBP1

STXBP1 plays a critical role in synaptic vesicle recycling[16, 17]; to assess this function, we assayed depletion of fluorescence in tissue that had been loaded with the lipophilic dye, FM4-64. Synaptic vesicles loaded with FM4-64 were identified along isolated axons in the subplate in 16 pcw tissue (Figure 4A). Under baseline conditions, signal intensity was reduced by 30% from 0 to 12 seconds in controls but not in *STXBP1* shRNA conditions (Figure 4B), indicating an impairment in spontaneous neurotransmission at synapses with reduced STXBP1 levels.

To assess evoked exocytosis, slices were additionally perfused with a depolarisation inducing concentration of potassium ions. The intensity of the signal started to decrease 120 seconds after the start of stimulation until the end of the experiment (overall 50% loss) in both scrambled controls and *STXBP1* shRNA, similarly (Figure 4B). This indicates that reduced STXBP1 does not affect evoked exocytosis in the subplate under these experimental conditions.

Discussion

We present a means of investigating genetic manipulations in human cortical tissue at developmental stages equivalent to the 14[th] pcw and onwards, through the prenatal period. In these initial, proof-of-principle studies, we demonstrate a knock-down of STXBP1, thereby mimicking the haploinsufficient state, which has previously been reported to underlie many cases of infantile epileptic encephalopathy[10].

Reduced levels of STXBP1 do not affect the gross morphology of subplate neurons at early developmental stages, consistent with other studies: first, absence of UNC-18, the ortholog gene of *STXBP1* in C. elegans, does not affect neurite length or maintenance[17]; second, heterozygous mice or Stxbp1 haploinsufficiency mouse models display no difference in dendrite length[16] or cortical neuron density[10] postnatally. Collectively, these findings indicate that reduced STXBP1 does not affect subplate neuronal architecture.

Deficiencies in STXBP1 can result in glutamatergic synaptic neurotransmission dysfunction in rodent models[10] and neurons derived from human embryonic stem cells[18]. Similarly, we find reduced glutamatergic post-synaptic current frequency and synapse number in the human subplate following knock-down of STXBP1. Interestingly, we observed an increase both in the frequency of GABAergic synaptic events, and in the density of GABAergic synapses. STXBP1 is normally present at GABAergic and glutamatergic synapses, so the divergent effects of STXBP1 knock-down on the two types of synapses was unexpected; given that the two synaptic classes are likely to be regulated in parallel, this divergence may represent a compensatory change manifest in one class, subsequent to a primary effect in the other.

The large decrease in frequency of glutamatergic currents, without a change in amplitude, is indicative of an impairment in pre-synaptic neurotransmitter release. Reliable and sustainable neurotransmitter release is essential for effective neuronal communication. Both spontaneous, and evoked neurotransmitter release is affected by lowering the levels of STXBP1 in many animal models[16-19]. These results are consistent with the observation that at mature synapses STXBP1 is crucial for the assembly of the SNARE complex, a collection of proteins involved in the fusion of vesicles with the plasma membrane[20-22], variants in several of which are implicated in neurodevelopmental disorders and epilepsy (termed SNAREopathies[9]). We observe impairments in spontaneous neurotransmitter release, but not evoked exocytosis. Interestingly, a similar dissociation between impaired spontaneous, but normal evoked release has previously been reported in mouse neurons expressing the human disease variant C180Y[11]. Our results may reflect some compensation from other cooperative proteins known to function together to chaperone SNARE assembly, including MUNC13-1[23, 24]. STXBP1 could also have a more important role in spontaneous than evoked exocytosis at early developmental stages because spontaneous neuronal activity is the predominant driving force controlling network formation[25].

In summary, the complex interplay highlighted here between GABAergic and glutamatergic elements is crucial for many critical stages of brain development, including setting homeostatic thresholds[26]. The subplate synaptic networks, which are far more developed in the human brain are proposed to guide neurogenesis, neuron migration, and formation of thalamocortical and cortico-cortical connectivity[4, 27, 28]. Therefore, alterations in subplate synaptic activity may alter the trajectory of development leading to the deficits typical of early infantile epileptic encephalopathies later in development. Using our model system to study other pre-synaptic proteins, such as neuronal SNAREs associated with neurodevelopmental disorders, could reveal integrative or divergent pathological mechanisms with STXBP1.

Acknowledgements

This work would not have been possible without the tissue donation to the HDBR and the efforts of everyone within the biobank and the help of the Bioimaging Unit at Newcastle University.

Author contributions

F.M. performed all experimental procedures in the manuscript, design of experiments and conceptualising the project. A.D assisted with the immunohistochemistry and image analysis. D.L.-S. and R.T were involved with conceptualisation of the project. G.J.C. and A.J.T. conceived the overall project, helped design the experiments and analyses. Funding for the project was obtained by A.J.T. All authors participated in the interpretation of data and writing of the manuscript.

Funding

The research to conduct the experiments was funded by the Wellcome Trust (Grant numbers 102037 and 203914/Z/16/Z) and Engineering and Physical Sciences Research Council (A000026). The work of the HDBR was funded by the Wellcome Trust and the Medical Research Council (MR/R006237/1). For open access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission.

Competing interests

Authors report no competing interests.

References

1. Symonds JD, Zuberi SM, Stewart K, et al. Incidence and phenotypes of childhood-onset genetic epilepsies: a prospective population-based national cohort. Brain. 2019;142(8):2303-18.

2. Lewis-Smith D, Ellis CA, Helbig I, et al. Early-onset genetic epilepsies reaching adult clinics. Brain. 2020;143(3):e19.

3. Luhmann HJ, Kirischuk S, Kilb W. The Superior Function of the Subplate in Early Neocortical Development. Front Neuroanat. 2018;12:97.

4. Molnar Z, Luhmann HJ, Kanold PO. Transient cortical circuits match spontaneous and sensory-driven activity during development. Science. 2020;370(6514).

5. Rizo J, Xu J. The Synaptic Vesicle Release Machinery. Annu Rev Biophys. 2015;44:339-67.

6. Polioudakis D, de la Torre-Ubieta L, Langerman J, et al. A Single-Cell Transcriptomic Atlas of Human Neocortical Development during Mid-gestation. Neuron. 2019;103(5):785-801 e8.

7. Stamberger H, Nikanorova M, Willemsen MH, et al. STXBP1 encephalopathy: A neurodevelopmental disorder including epilepsy. Neurology. 2016;86(10):954-62.

8. O'Brien S, Ng-Cordell E, Study DDD, et al. STXBP1-associated neurodevelopmental disorder: a comparative study of behavioural characteristics. J Neurodev Disord. 2019;11(1):17.

9. Verhage M, Sorensen JB. SNAREopathies: Diversity in Mechanisms and Symptoms. Neuron. 2020;107(1):22-37.

10. Chen W, Cai ZL, Chao ES, et al. Stxbp1/Munc18-1 haploinsufficiency impairs inhibition and mediates key neurological features of STXBP1 encephalopathy. Elife. 2020;9.

11. Kovacevic J, Maroteaux G, Schut D, et al. Protein instability, haploinsufficiency, and cortical hyper-excitability underlie STXBP1 encephalopathy. Brain. 2018;141(5):1350-74.

12. Wang WZ, Hoerder-Suabedissen A, Oeschger FM, et al. Subplate in the developing cortex of mouse and human. J Anat. 2010;217(4):368-80.

13. Di Lullo E, Kriegstein AR. The use of brain organoids to investigate neural development and disease. Nat Rev Neurosci. 2017;18(10):573-84.

14. Cousin MA. Use of FM1-43 and other derivatives to investigate neuronal function. Curr Protoc Neurosci. 2008;Chapter 2:Unit 2 6.

15. McLeod F, Marzo A, Podpolny M, et al. Evaluation of Synapse Density in Hippocampal Rodent Brain Slices. J Vis Exp. 2017(128).

16. Toonen RF, Wierda K, Sons MS, et al. Munc18-1 expression levels control synapse recovery by regulating readily releasable pool size. Proc Natl Acad Sci U S A. 2006;103(48):18332-7.

17. Weimer RM, Richmond JE, Davis WS, et al. Defects in synaptic vesicle docking in unc-18 mutants. Nat Neurosci. 2003;6(10):1023-30.

18. Patzke C, Han Y, Covy J, et al. Analysis of conditional heterozygous STXBP1 mutations in human neurons. J Clin Invest. 2015;125(9):3560-71.

19. Verhage M, Maia AS, Plomp JJ, et al. Synaptic assembly of the brain in the absence of neurotransmitter secretion. Science. 2000;287(5454):864-9.

20. Deak F, Xu Y, Chang WP, et al. Munc18-1 binding to the neuronal SNARE complex controls synaptic vesicle priming. J Cell Biol. 2009;184(5):751-64.

21. Parisotto D, Pfau M, Scheutzow A, et al. An extended helical conformation in domain 3a of Munc18-1 provides a template for SNARE (soluble N-ethylmaleimide-sensitive factor attachment protein receptor) complex assembly. J Biol Chem. 2014;289(14):9639-50.

22. Rizo J, Sudhof TC. Snares and Munc18 in synaptic vesicle fusion. Nat Rev Neurosci. 2002;3(8):641-53.

23. He E, Wierda K, van Westen R, et al. Munc13-1 and Munc18-1 together prevent NSF-dependent de-priming of synaptic vesicles. Nat Commun. 2017;8:15915.

24. Shu T, Jin H, Rothman JE, et al. Munc13-1 MUN domain and Munc18-1 cooperatively chaperone SNARE assembly through a tetrameric complex. Proc Natl Acad Sci U S A. 2020;117(2):1036-41.

25. Luhmann HJ, Sinning A, Yang JW, et al. Spontaneous Neuronal Activity in Developing Neocortical Networks: From Single Cells to Large-Scale Interactions. Front Neural Circuits. 2016;10:40.

26. Baines RA. Neuronal homeostasis through translational control. Mol Neurobiol. 2005;32(2):113-21.

27. Kostovic I. The enigmatic fetal subplate compartment forms an early tangential cortical nexus and provides the framework for construction of cortical connectivity. Prog Neurobiol. 2020;194:101883.

28. Ozair MZ, Kirst C, van den Berg BL, et al. hPSC Modeling Reveals that Fate Selection of Cortical Deep Projection Neurons Occurs in the Subplate. Cell Stem Cell. 2018;23(1):60-73 e6.

Figures

Figure 1 Human organotypic cortical brain slices cultures with developing subplate neurons. (**A**) Schematic and low magnification photomicrographs of organotypic cultures prepared from human foetal brain tissue. Cultures are maintained for 1–4 weeks following infection with and without different viral vectors. Micrograph image represents a 16 pcw sample, 2 weeks in culture with clearly identifiable cortical regions including the marginal zone (MZ), cortical plate (CP) and subplate (SP). DAPI labels all cell nuclei and Synapsin1 immunohistochemistry labels reveals pre-synaptic terminals predominantly located in SP but also CP and MZ. (**B**) Representative images from 14, 16 and 17 pcw cultures (0 DIV). STXBP1 (red) and NeuN (green, all neurons) immunofluorescence reveals location of the SP. Note the enrichment of STXBP1 within the SP at 16–17 pcw (*n = 3* human samples/age). (**C**) SP neuron (green, eGFP transduced cell) with STXBP1 (red) present on neurites along thin putative axons extending from the cell body (i) and spine-like structures (ii) indicated by the white arrows. (**D**) Photomicrographs of a 16 pcw culture infected with AAV9-CMV-GFP and taken at 7, 21 and 27 DIV, showing the same neuron (highlighted by the masked images). Note the dynamic nature and elaboration of neurites over time. Coloured outlines highlight reference cells – red and green are stable with a minor orientation change in the blue cell at early culture stages.

Figure 2 Synapse maturation in human cortical slice cultures. (A) Left: schematic displaying the pre-synaptic location of STXBP1. Middle: photomicrographs of STXBP1 (red) in the subplate of 16–17 pcw cultures at 0 and 27 DIV. Right: quantification of the increase in STXBP1 over time at both ages. **(B)** Left-middle: co-localisation of the pre- and post-synaptic glutamatergic markers, vGlut1 (green) and Homer1 (red), respectively, in the subplate at 0 and 27 DIV. Areas, demarcated by dotted boxes in the upper micrographs, are shown enlarged below. All cultures were prepared at 16–17 pcw. **(C)** Similar analyses performed for GABAergic synapses based upon co-localisation of vGAT (pre-synaptic in green) and Gephyrin (post-synaptic in red). Analyses for panels (A) to (C) were done with a two-way ANOVA with repeated measures from comparisons made at 0 DIV for each age, $n = 6$–8 slices per time point from 2–3 human samples/age. Error bars indicate SEM.

Figure 3 Divergent regulation of glutamatergic and GABAergic synapses in the subplate following knock-down of STXBP1. (A) Left: western blot showing knock-down of STXBP1 protein in a whole slice at 16 pcw following 2 weeks with scrambled control (Ctr) or *STXBP1* shRNA. Right: quantification of band intensity showing approximately 50% reduction in STXBP1 levels (unpaired t-test, taken from *n = 5* slices/condition from *2* human samples). **(B)** Left: example subplate neurons infected with scrambled control or *STXBP1* shRNA for 2 weeks at 16 pcw. Note the putative axons highlighted with the arrows. Right: quantification of primary neurite number and length reveals no gross morphology changes in both conditions (*n = 15* cells from *10* slices and *2* human samples). **(C-E)** Quantification of glutamatergic synapses (white arrows) (C) and GABAergic (E) synapses (white arrows) in the subplate at 16 pcw following two weeks of scrambled control or *STXBP1* shRNA by colocalisation of pre- and post-synaptic markers. Analysis performed using an unpaired t-test, *n = 6–8* slices per time point from *2* human samples. **(D-F)** Representative miniature glutamatergic post-synaptic currents recorded at -60 mV (D) and miniature GABAergic post-synaptic currents recorded at 0 mV (F) from 16 pcw subplate cells in both genetic conditions. The frequency and amplitude of events (starred) were quantified (unpaired t-test, *n = 8* cells per time point from *2* human samples). All data represented as mean ± SEM

Figure 4 Impaired spontaneous vesicle release with reduced STXBP1 levels. (A) Isolated axons tracked from FM4-64 loaded subplate neurons in scrambled control and STXBP1 shRNA slices (16 pcw). Note the presence of spontaneous release from vesicles loaded with FM4-64 (white arrows). **(B)** Quantification of spontaneous exocytosis (left; *n = 8* slices from *2* human samples) and evoked exocytosis (right; *n = 6* slices from *2* human samples) following brief potassium chloride (KCl) stimulation in both genetic conditions. Analysis was performed using a two-way ANOVA with repeated measures. All error bars represented as SEM.

Supplementary Figures

Supplementary Figure 1: Genetic knock-down of STXBP1 in the subplate of human cortical slice cultures. (A) Photomicrographs of STXBP1 (red) in the subplate in the subplate at 16 pcw following 2 weeks with scrambled control or *STXBP1* shRNA **(B)** Quantification of STXBP1 puncta count and integrated density levels in the subplate. Note the comparable terminal numbers (STXBP1 count) but less protein (STXBP1 levels). Analysis performed using an unpaired t-test, *n = 8* slices/condition from *2* human samples). All data represented as mean ± SEM.

Supplementary Methods

Human organotypic slice cultures

Tissue was transferred to oxygenated ice-cold high sucrose ACSF containing (in mM): 180 sucrose, 25 $NaHCO_3$, 10 glucose, 10 $MgSO_4$, 2.5 KCl, 2 N-acetylcysteine, 1.25 NaH_2PO_4, 1 taurine, 1 ascorbic acid, 0.5 $CaCl_2$, 0.1 aminoguanidine hydrochloride, 0.044 indomethacin, 0.044 ethyl pyruvate and adjusted to 300–310 mOsM and 7.4 pH. Subsequently 280μm slices were cut using a 5100mz Vibratome (Camden Instruments). Cortical slices were placed inside 6-transwell plates containing BrainPhys (5790; Stemcell Technologies) culture media supplemented with 1 x N2, 1 x B27, 40ng/ml Brain-derived Neutrophic Factor (BDNF), 20ng/ml Glia-derived Neurotrophic Factor (GDNF), 30ng/ml Wnt7a, 200 nM ascorbic acid, 1 mM dibutyryl cyclic AMP and 1 μg/ml laminin.

Live imaging

Live imaging was performed using an inverted Zeiss LSM800 Airyscan confocal microscope. AAV9-CMV-GFP infected slices were placed in culture media and maintained at 37°C, 5% CO_2. Images were acquired in the subplate using a x20 objective (Numerical Aperture; NA = 0.8) every 30 seconds for 15 minutes from the same slice for 4 weeks generating final image resolutions of 512 x 512 pixels.

FM4-64 dye was utilised as previously described[14]. Briefly, infected slices were transferred to control solution containing (in mM, pH 7.3): 120 NaCl, 10 HEPES, 10 Glucose, 2 $CaCl_2$ and 1 $MgCl_2$ to equilibrate. After 5 minutes, they were placed in control solution, FM4-64 dye (7.5 μM) and potassium chloride (KCl; 50 mM with osmolarity compensated) for 90 seconds to stimulate and load the dye in vesicles. Subsequently, slices were placed for 60 seconds in control solution and FM4-64 dye. Finally, slices were transferred to control solution to wash off any excess non-internalised dye.

Consecutive images were acquired of FM4-64 loaded vesicles using a 40x objective (NA = 1.3) at room temperature (RT). Slices were firstly perfused with control solution and axonal regions loaded with FM4-64 vesicles identified for imaging. After 10 minutes, vesicle exocytosis (FM4-64 dye unloading) was stimulated by perfusion with control solution (substituted with 50mM KCl) for 150 seconds, before reperfusion with control solution for a further 3 minutes. Simultaneous images of the GFP-positive axons and FM4-64 loaded vesicles

were acquired just before, during and after stimulation, with a 3 second delay between each frame.

Analysis was performed using Fiji software (NIH, RRID:SCR_002285). Regions of interest (ROIs) representing the vesicles, were identified for each slice. One ROI on an area with no signal (representing the background) and one ROI on an area with a stable signal (to account for the FM4-64 bleaching) were selected. The ROIs remained the same throughout the videos and their mean fluorescent measurements were quantified. Mean background values were subtracted from the measured signal, and each value was divided by the mean stable signal value. Finally, these values were normalised to the average ROI values of all frames 60 seconds before stimulation to calculate the fraction of loaded vesicle pool. Evoked exocytosis was averaged every 9 seconds. For spontaneous exocytosis (just before stimulation), ROI values were normalised to the average of first five frames.

Whole cell-patch clamp recordings

Slices were transferred into the recording chamber of an upright Leica DMLFSA fluorescent microscope fitted with Micro Control Instruments micromanipulators and continuously perfused at 34°C with oxygenated ACSF containing (in mM): 126 NaCl, 3 KCl, 1.25 NaH_2PO_4, 24 $NaHCO_3$, 10 Glucose, 1.2 $CaCl_2$ and 1 $MgSO_4$. Cells were voltage-clamped in whole cell configuration using borosilicate glass patch electrodes (5–8 MΩ) filled with an intracellular solution containing (in mM): 125 K-methyl-SO_4, 10 HEPES, 2.5 Mg-ATP, 6 NaCl, 290 mOsM and pH 7.35. All data were collected using an Axopatch 200B amplifier, filtered (1 kHz) and digitised (10 kHz). Miniature currents were monitored and analysed using WinEDR and WinWCP software (http://spider.science.strath.ac.uk/sipbs/software_ses.htm).

Immunohistochemistry

Slices were fixed in 4% paraformaldehyde (PFA)/4% sucrose in phosphate buffered saline (PBS; 0.12 M, pH 7.4; PBS) for 20 minutes at room temperature (RT). Staining was performed using the protocol optimised specifically for acute slices[15]. In brief, slices were incubated in blocking solution (10% donkey serum, 1% Triton X-100 in PBS) for ~ 4-6 hours at RT on a vibrating shaker. Primary antibodies against Synapsin1 (1:500; Millipore, AB1543P), vGlut1 (1:2000; Millipore, AB5905), Homer1 (1:1000; Synaptic Systems, 160006)), vGAT (1:1000; Synaptic Systems, 131003), Gephyrin (1:500; Synaptic Systems, 147318), NeuN (1:1000; Abcam, AB104224), STXBP1 (1:500; Abcam, ab109023), GAD67 (1:1000; Millipore,

MAB5406), MAP2 (1:2000, Abcam, AB5392) or GFP (1:5000; Abcam, ab290) were applied overnight at 4°C. Slices were subsequently incubated with fluorophore-bound secondary antibodies (1:500; Life Technologies) for 2-3 hours at RT on a vibrating shaker before DAPI (1:5000) applied and mounted on glass slides in Fluoromount-G (SouthernBiotech).

Image analysis

The number as well as the length of primary subplate neuron processes was measured using Fiji software. Synaptic counts and integrated density (area × mean intensity) measurements were conducted using Zen 3.1 software (Zeiss; RRID:SCR_013672). These were optimised with a range of filters and thresholds to specifically identify the ROI for pre- and post-synaptic markers. Settings for markers of interest were kept consistent throughout all replicates. A synapse was defined as the colocalisation of specific pre- and post-synaptic markers with a minimum of 2 pixels overlapping. The counts of individual and colocalised ROIs were normalised to the volume of each image then represented per $100 \ \mu m^3$.

Western blot

Membranes were blocked then probed overnight at 4°C on a vibrating shaker with anti STXBP1 antibody (1:1000; Abcam) and anti β-actin (1:5000; Cell Signalling). Appropriate species specific HRP-conjugated secondary antibodies were added (BIO-RAD) at RT. Visualisation of protein bands was performed using a GBOX-Chemi-XRQ gel system (Syngene).