Oscillatory calcium release and sustained store-operated oscillatory calcium signaling prevents

differentiation of human oligodendrocyte progenitor cells

Richard A. Seidman[1], Heba Khattab[2], Jessie J. Polanco[1], Jacqueline E. Broome[2], Fraser J. Sim[1,2]

Neuroscience Program[1], Department of Pharmacology and Toxicology[2], Jacobs School of Medicine

and Biomedical Sciences, University at Buffalo, Buffalo, NY.

Address correspondence to:

Fraser Sim, Ph.D.

Department of Pharmacology and Toxicology

Jacob's School of Medicine and Biomedical Sciences

University at Buffalo

955 Main Street

Buffalo, NY 14203

716-829-2151

fjsim@buffalo.edu

Keywords: human, lentivirus, STIM1, STIM2, muscarinic receptor, metabotropic glutamate receptor

Running title: STIM-dependent Ca-signaling blocks hOPC differentiation.

Abstract, 242 words; Introduction, 862 words; Discussion, 1753 words.

Figures: 5; Tables: 2

Acknowledgements:

This work was supported by grants from the NINDS (R01NS104021), the National Multiple

Sclerosis Society (RG 5505-A-2, RG 5110A1/1), the Kalec Multiple Sclerosis Foundation, the Change

MS Foundation, and the Skarlow Memorial Trust. JJP received support from NIGMS (R25GM09545902),

NCATS (UL1TR001412-S1). JJP and RAS received support from NYSTEM Stem Cells in Regenerative

Medicine Fellowship (#C302090G). In addition, we would like to acknowledge the assistance of the

Confocal Microscope and Flow Cytometry Facility in the School of Medicine and Biomedical Sciences,

University at Buffalo and thank Dr. Pablo Paez for advice with calcium imaging of OPCs.

ABSTRACT

Endogenous remyelination in demyelinating diseases such as multiple sclerosis is contingent upon the successful differentiation of oligodendrocyte progenitor cells (OPCs). Signaling via the Gα$_q$-coupled muscarinic receptor (M$_{1/3}$R) inhibits human OPC differentiation and impairs endogenous remyelination in experimental models. We hypothesized that calcium release following Gα$_q$-coupled receptor (G$_q$R) activation directly regulates human OPC (hOPC) cell fate. In this study, we show that specific G$_q$R agonists activating muscarinic and metabotropic glutamate receptors induce characteristic oscillatory calcium release in hOPCs and that these agonists similarly block hOPC maturation *in vitro*. Both agonists induce calcium release from endoplasmic reticulum (ER) stores and store operated calcium entry (SOCE) likely via STIM/ORAI-based channels. siRNA mediated knockdown (KD) of obligate calcium sensors STIM1 and STIM2 decreased the magnitude of muscarinic agonist induced oscillatory calcium release and attenuated SOCE in hOPCs. In addition, STIM2 expression was necessary to maintain the frequency of calcium oscillations and STIM2 KD reduced spontaneous OPC differentiation. Furthermore, STIM2 siRNA prevented the effects of muscarinic agonist treatment on OPC differentiation suggesting that SOCE is necessary for the anti-differentiative action of muscarinic receptor-dependent signaling. Finally, using a gain-of-function approach with an optogenetic STIM lentivirus, we demonstrate that independent activation of SOCE was sufficient to significantly block hOPC differentiation and this occurred in a frequency dependent manner while increasing hOPC proliferation. These findings suggest that intracellular calcium oscillations directly regulate hOPC fate and that modulation of calcium oscillation frequency may overcome inhibitory Gα$_q$-coupled signaling that impairs myelin repair.

Significance Statement

In this study, Seidman et al. show that SOCE is a common component of ligand-based Gα$_q$-coupled signaling in hOPCs and that SOCE alone is sufficient to block hOPC differentiation and drive proliferation. Therefore, SOCE blocks differentiation and pathological SOCE could contribute to myelin disease.

INTRODUCTION

Efficient central nervous system (CNS) remyelination is contingent upon successful intercellular

signaling between neurons and glia that together orchestrate the processes of oligodendrocyte progenitor

cell (OPC) recruitment, differentiation, and maturation into myelin-forming oligodendrocytes (OL) [1]. In

demyelinating diseases such as multiple sclerosis (MS), the pathological environment contributes to an

inhibition of OPC differentiation that limits remyelination and may contribute to subsequent permanent

neurodegeneration [2]. In support of this hypothesis, numerous quiescent OPCs are observed in chronic

MS lesions that appear to be unable to undergo OL differentiation [3,4]. Following demyelination, OPCs are

exposed to a complex cellular microenvironment comprising extracellular matrix and soluble factors

released by a variety of sources including infiltrating immune cells, reactive glia, and endothelial cells as

well as injured axons that release both soluble growth factors and neurotransmitters. Neurotransmitters

activate multiple and functionally distinct receptors expressed on OPCs and OLs that may contribute to

activity-dependent or so-called adaptive myelination (reviewed in [5]). Following demyelination, neuronal

activity promotes remyelination [6] and this is regulated in part by AMPA and NMDA receptor signaling [7].

As such, intracellular calcium represents a common target for these activity-dependent pathways and

suggests that a diverse set of neurotransmitters can induce intracellular calcium increase in glial

progenitors [8].

Intracellular calcium release occurs in OPCs following activation of ionotropic voltage and ligand

gated channels and/or following activation of metabotropic receptors [9]. Muscarinic acetylcholine

receptors M_1 and M_3 are $G\alpha_q$-coupled receptors (G_qR) which also cause intracellular calcium release via

phospholipase-Cβ (PLCβ) and inositol trisphosphate (IP$_3$) second messengers [10]. Genetic deletion of $M_{1/3}$

receptors in OPCs delays differentiation and remyelination in animal models of experimental

demyelination [11,12]. Furthermore, pharmacological inhibition of muscarinic receptors results in accelerated

OL differentiation and remyelination in the brain and spinal cord following both toxin-induced

demyelination and experimental autoimmune encephalitis (EAE) [11,13-15]. In primary human OPCs

(hOPCs), $M_{1/3}R$ signaling delays differentiation *in vitro* and reduces myelin production following

transplantation into hypomyelinated *shiverer* mice [12]. OPCs have been shown to express numerous other

$G\alpha_q$-coupled receptors including endothelin (ET) subtype B (ET$_B$; ENDRB) receptors [16], purinergic type

2Y (P2Y) receptors [17], histamine type 1 (H1) receptors [18], metabotropic glutamate receptors type 3

(mGluR$_3$) and type 5 (mGluR$_5$) [19], 5-HT serotonin receptors [20], sphingosine-1-phosphate receptors

(S1PR) [21], calcium-sensing receptors (CaSR) [22], and adenosine 2B receptors [23]. While the precise

function of these receptors remains unclear, previous reports suggest that activation of at least some of

these receptors can paradoxically promote OPC differentiation. For example, *in vitro* findings suggest

that ET receptor agonists increase OPC migration and block proliferation [16]. Stimulation of myelin basic

protein (MBP) expression occurs via ET$_A$ [24] receptor activity, and activation of ET$_B$ receptors has been

found to promote remyelination in cerebellar slice culture [25]. Likewise, conditional deletion of S1PR

causes increased OPC proliferation and reduced oligodendrocyte differentiation [26], while CaSR null mice

exhibit reduced MBP expression during development [22]. Despite these findings, the key signaling

components downstream of muscarinic receptor activation that act to delay OPC differentiation and

distinguish M$_{1/3}$R from other Gα$_q$-receptors have not yet been established.

Following activation of Gα$_q$-coupled receptors and calcium release from the endoplasmic

reticulum (ER), calcium sensing STIM proteins in the ER membrane oligomerize and complex with ORAI

channel proteins located in the plasma membrane. These calcium release activated calcium (CRAC)

channels mediate store operated calcium entry (SOCE) that replenish depleted ER-stores through

sarcoplasmic/endoplasmic reticulum ATPase (SERCA) pumps [27]. Successive activation of ER-calcium

release and CRAC channel influx events result in characteristic oscillatory calcium transients [28].

Intriguingly, the frequency encoding of oscillatory calcium signaling is associated with cell-type specific

activation of distinct downstream signaling and gene expression [28,29]. In rodent OPCs, *in vitro* application

of a pharmacological SOCE antagonist limits platelet derived growth factor (PDGF)-dependent

proliferation [30]. However, the mechanisms of action of these drugs are complex with additional actions

on IP$_3$ receptors, gap junction permeability, and transient receptor potential channels. Furthermore, while

ligand-induced calcium oscillations have been observed in OL lineage cells [19,31,32], a specific role of

calcium oscillations in human OPCs has not yet been determined.

In this study, we hypothesized that the specific oscillatory signature of ligand activated Gα$_q$-

receptor mediated calcium responses may be responsible for the regulation of hOPC differentiation. To

this end, we identified several Gα$_q$-coupled receptors expressed by hOPCs and characterized the calcium

response following application of receptor subtype-specific ligands. Only activation of muscarinic

acetylcholine M$_{1/3}$ receptors or metabotropic type 5 glutamate receptors (mGluR$_5$) resulted in oscillatory

transients in a large fraction of OPCs and persisted for several minutes following application of the ligand.

Intriguingly, these ligands also reduced hOPC differentiation. We demonstrate that SOCE downstream

of M$_{1/3}$R ligand activation is dependent on ER calcium sensors STIM1/2. More specifically, we found that

both STIM proteins contribute to the magnitude of SOCE in human OPCs and that STIM2 regulates the

oscillatory frequency of M$_{1/3}$R-induced calcium responses. Knockdown of STIM2 resulted in reduced OL

differentiation and prevented the anti-differentiative effects of M$_{1/3}$R signaling in hOPCs. Lastly, using

optogenetic activation of CRAC-mediated calcium flux, we demonstrate that SOCE induction is sufficient

to inhibit hOPC differentiation in the absence of G$_q$R signaling and that modulation of OPC proliferation

and differentiation was dependent on SOCE frequency.

MATERIALS AND METHODS

Human CD140a/PDGFαR cell preparation and culture with Gα$_q$-agonists

Fetal brain tissue samples, between 17 and 22 weeks of gestational age, were obtained from

Advanced Bioscience Resources (Alameda, CA) with informed consent obtained from all donors. All

research was performed in accordance with relevant guidelines/regulations. Following review by the

University at Buffalo Research Subjects Institutional Review Board the tissue acquisition and research

was not deemed to involve human subjects as defined under HHS regulations 45 CFR 46, 102 (f).

Forebrain samples were minced and dissociated using papain and DNase as previously described [33].

Magnetic sorting of CD140a/PDGFαR positive cells was performed as described [34].

Plasmid construction, lentiviral generation, and titration

EF1α:GCaMP6s was described previously [12]. EF1α:jRCaMP1a was cloned as follows. pGP-CMV-

NES-jRCaMP1a was obtained from AddGene (#61562) [35]. The nuclear export signal (NES) and

jRCaMP1a coding region was PCR amplified: SpeI, forward–5'

AAAACTAGTGAACCGTCAGATCCGCTAG-3';PspXI (Thermo Fisher Scientific), reverse, 5'-

AAAGCTCGAGCTCTACAAATGTGGTATGGCTG-3' (Thermo Fisher Scientific). PCR products were

purified (Monarch PCR Cleanup Kit (NEB, #T1030S)) and restriction digested with unique 5' SpeI and 3'

PspXI sites, and purified again. NES-jRCaMP1a digested PCR fragments were then ligated into lentiviral

pTRIP-EF1α vector (a gift of Abdel Benraiss, University of Rochester)[36]. The plasmid pLenti-EF1α-

OptoSTIM was a gift from Taeyoon Kyung (KAIST, DAEJEON, Republic of Korea) [37]. Lentiviruses were

generated as described [38] and titration of EF1α:GCaMP6s and EF1α:OptoSTIM was performed as

previously described [39]. Optimized multiplicity of infection (MOI) in hOPCs was determined for

EF1α:GCaMP6s and EF1α:jRCaMP1a by infecting hOPCs and quantifying proportion of responsive cells

following administration of 25µM Oxo-M. Optimal MOI for EF1α:OptoSTIM lentivirus was determined

through analysis of optimal GFP expression in infected hOPCs.

Calcium imaging using LV EF1a:GCaMP6s

Calcium imaging was performed as described previously [12]. GCaMP6s infected hOPCs were

seeded at 4-5 x 10^4 cells/mL and imaged 24 hours after seeding. All experimental data were generated

from independent experiments using individual naïve cultures of hOPCs from separate patient fetal

samples. To avoid possible confounds relating to sequential application of small molecules, all cultures

were used only once for imaging calcium responses and only a single dose was tested in each well, with

two technical replicate imaging fields imaged simultaneous per well for each dose or condition tested. In

dose response experiments, all doses per each drug were tested on the same experimental day for each

fetal sample preparation. Bright field and fluorescence images were acquired (10x; Olympus IX51

microscope with a Exfo X-Cite or Xylis illumination and Hamamatsu ORCA-FLASHv4 camera) using

µManager [40]. All drugs were reconstituted and aliquoted within 1-2 weeks and thawed immediately prior

to use. Region of interest (ROI) generation and image processing were performed using ImageJ, and

analysis of calcium transients was performed in R as described [12].

Briefly, per cell calcium responses (raw pixel intensity) were normalized to baseline fluorescence

calculated prior to addition of Gα_q-receptor ligand (F_0, average 5-10 frames). To define calcium peaks

and characterize the properties of ligand-induced calcium oscillations, individual cell traces were loess

smoothed prior to the identification of local minima and maxima in 16 second windows. Calcium peaks

were defined as maxima whose amplitude increased ≥0.35 fold-change above the preceding local

minima. Peak amplitude was defined as the difference in fluorescence values from the calcium peak to

the preceding local minima. The number of oscillations was defined as the number of peaks identified

during the recording session. A responsive cell was defined as any cell with at least one peak, and an

oscillatory cell as any cell with two or more peaks. Response duration was measured from the local

minima prior to the first peak oscillation to the local minima following the last peak oscillation, or to the

end of the time-lapse recording if an on-going calcium peak had not yet reached baseline. For oscillatory

cells, frequency was calculated by dividing the total number of peak oscillations by the response duration.

Area   under   the   curve   (AUC)   was   calculated   (*Mess*   package,   v0.5.6;   https://cran.r-

project.org/web/packages/MESS/index.html) by subtraction of F_0, and then analyzed as integral pixel

intensity above baseline beginning from the local minima preceding the first identified peak on the

normalized calcium curve following ligand stimulation for the duration of the imaged response.

Raw data measurements of individual cell calcium responses were assessed for normality. Per-

cell AUC and frequency measurements exhibit significant skew and were log-transformed to achieve a

near normal distribution. Unless otherwise stated, a linear model was used for one-way ANOVA using

drug dose as a covariate, as well as human sample to consider individual sample variability. Non-linear

regression was performed using GraphPad Prism (8.0e) using a variable slope-four parameter logistic

equation to model and calculate EC$_{50}$ for each drug. For STIM KD experiments, GCaMP6s cells were

transfected with STIM siRNA and imaged 24-48h post transfection.

Assessment of store-operated calcium entry

For direct assessment of SOCE, hOPCs were infected with LV-GCaMP6s (1 MOI) and 5 x 10^4

cells/mL were seeding in 48-well plates and imaged the following day. To remove extracellular calcium,

15 minutes prior to imaging media was replaced with HBSS without phenol red, calcium, and magnesium

(Corning, #21-022) (HBSS-). For all conditions, fields were imaged with a 5 second interval for at least

15 minutes (488nm excitation, 400ms exposure). Addition of the SERCA blocker Thapsigargin (Tg) or

193 the muscarinic agonist Oxotremorine-Methiodide (Oxo-M) with or without store operated calcium entry

194 (SOCE) channel antagonists occurred at ~1-2 minutes after the start of imaging. Calcium [1.2mM]) was

195 re-introduced 10-15 minutes following addition of Tg or Oxo-M. Final concentrations of compounds were

196 Tg [10µM] or Oxo-M [25µM], and 2-APB [50µM] or MRS-1845 [10µM] (final DMSO concentration did not

197 exceed 0.15%, which did not influence [Ca^{2+}]$_i$). For quantification of SOCE, per cell calcium response

198 curves were normalized to baseline fluorescence F_0 (average of 5-10 frames prior to addition of Oxo-M

199 or Tg). Normalized traces were loess smoothed to identify responsive cells exhibiting ≥0.35 fold increase

200 in GCaMP6s fluorescence from baseline following initial Oxo-M or Tg ER-store depletion. Area under the

201 curve (AUC) was measured for ER depletion for 10 minutes following Oxo-M or Tg ER-store depletion,

202 and AUC for SOCE was measured for 5-8 minutes (**Fig. 1H**) or 8 minutes (**Fig. 2E**) following calcium re-

203 addition. AUC durations were matched between conditions and were quantified. For analysis of SOCE

204 following STIM KO, AUC of SOCE was normalized to the ER-release of control cells in matched biological

205 replicates and quantified using repeated measures (RM) one-way ANOVA with Holm-Sidak's post-hoc

206 test. Maximum amplitude of ER depletion was identified as the maximal peak response >0.35 fold-change

207 increase from baseline following Oxo-M or Tg addition and following calcium re-addition respectively.

208 Cells which did not exhibit an initial ER-store depletion response (≥0.35 fold-change from baseline

209 following Tg or Oxo-M addition and prior to calcium re-addition) were excluded from further analyses.

210 *siRNA transfection*

211  hOPCs were seeded at 5 x 10^4 cells/mL and Lipofectamine RNAi max (Invitrogen, #13778-150)

212 was used to transfect cells according to the manufacturer's protocol with minor alterations. Briefly, stealth

213 RNAi negative control duplex medium GC content (Invitrogen, #12935300) or three combined stealth

214 RNAis against STIM1 (Invitrogen, ID#'s: HSS110308, HSS110309, HSS186144) and/or STIM2

215 (Invitrogen, ID#'s: HSS183972, HSS183973, HSS183974) were pooled and used for transfection. Stealth

216 RNAi (100 nM total RNAi per well) was used to transfect hOPCs per each condition with 1.5% v/v

217 Lipofectamine. For calcium imaging, cells were plated in 48-well plates and imaged 24 hours post-

218 transfection. In differentiation experiments, mitogen containing media was removed 24 hours post-

219 transfection (experimental day 0) and replaced with media without mitogens. Oxo-M [25µM] was added

220 on experimental day 0 upon removal of growth factors and on experimental day 2 during a half media

221 change. hOPCs were fixed on experimental day 3-4 following live staining with mouse x O4 Hybridoma

222 primary antibody [1:25]. Immuno-cytochemical analysis was performed as described previously [33]. For

223 analysis of STIM expression, cells were plated at 5 x 10^4 cells/mL 6-well plates and transfected after 24

224 hours. RNA was extracted 24-48 hours post-transfection for cDNA synthesis and quantitative RT-PCR

225 analysis.

Quantitative RT-PCR analyses

Total RNA was extracted (Omega Biotek) and cDNA was synthesized using SuperScript III

reverse transcriptase (Invitrogen, Carlsbad, CA). Human-specific primers for SYBR green-based PCR

were designed (**Table 1**). Samples were run in triplicate and gene expression calculated by $\Delta\Delta C_t$ method

using GAPDH as a reference. Statistical significance was tested on log$_2$-transformed data normalized to

control using a one-sample t-test assuming equal SD and compared to a hypothetical value of 100

(GraphPad Prism 8.0e).

Assessment of long-term OptoSTIM-induced SOCE

For all long-term experiments, hOPCs were seeded at a density of 2.5-3.0 x 10^5 cells/mL into

poly-ornithine and laminin coated ibidi tissue culture treated μ-slide VI 0.4 (iBIDI, #80606) in mitogen

containing Livelight ND media to reduce phototoxicity [41]. Cells were then imaged in an on-stage ibidi

environmentally controlled chamber to maintain 37ºC, 5.0% CO2, and >60% humidity for the duration of

each imaging experiment. In long-term calcium imaging experiments, hOPCs were infected with

jRCaMP1a (1 MOI) and OptoSTIM (1 MOI) lentiviruses. Two fields per each of 3 μ-slide channels were

then imaged at 10x with 594 nm excitation every 20 seconds for 3 hours to measure jRCaMP1a

fluorescence calcium responses. After a 30-minute no-stimulation period and for the remainder of

imaging, hOPCs were stimulated with 488 nm excitation (every 5 or 30 minutes with 330 and 2000 ms

illumination times respectively) to induce SOCE calcium responses. A custom beanshell script was

designed and implemented to facilitate multi-channel and multi-position imaging and provide an interface

between the motorized stage, fluorescent shutter (Prior), and μManager software. The complete imaging

script is available upon request. Analysis of OptoSTIM-induced calcium responses was conducted by

implementing the automated tracking plugin TrackMate in ImageJ [42]. Mean fluorescence intensity data

within the region of interest for all cells tracked across all frames was exported and subsequently

analyzed in R. Briefly, a baseline curve was generated by fitting minima points under each calcium trace

using the "spc-Rubberband" function (*hyperSpec* package). Traces were first normalized by division of

the baseline curve and loess fitted to identify responsive cells with ≥12.5% increase in jRCaMP1a

fluorescence within 2 minutes following the first OptoSTIM 488nm pulse. For quantification of AUC, each

tracked cell's mean jRCaMP1a fluorescence values per each time-point were normalized by subtraction

of the baseline curve and AUC was calculated for the duration of the stimulation period. Calcium traces

are presented as fold change in jRCaMP1a fluorescence from baseline, averaged across all cells per

each time-point with SEM calculated. In differentiation and proliferation experiments, OptoSTIM infected

hOPCs (1 MOI) were seeded into μ-slides and bright field imaged for 2–3 days. Following a 10-12 hour

baseline period, OptoSTIM activation was initiated with 488nm stimulation at different frequencies (every

5 or 30m with 330 or 2000ms illumination times respectively) for the remainder of imaging. After imaging,

cells were immunostained for O4 and the number of O4$^+$ and GFP$^+$ cells were counted and analyzed.

Repeated measures (RM) one-way ANOVA was employed with Holm-Sidak's post-hoc test for

quantification of the proportion of O4$^+$ to GFP$^+$ cells and quantification of morphological branching. In

proliferation experiments, cell density was counted at the onset of blue light stimulation and mitotic events

were identified and counted throughout the stimulation period. RM one-way ANOVA with Holm-Sidak's

post-hoc test was implemented to quantify significance in proliferative events normalized to initial

densities for each condition.

RESULTS

Gα$_q$-receptor activation induces ER-Ca^{2+} responses and SOCE in hOPC.

We hypothesized that Gα$_q$-coupled receptors would exert distinct effects on intracellular calcium

responses in hOPCs and that these responses would correlate with their functional effects. Using RNA-

seq of human primary PDGFRα-isolated OPCs, we determined that hOPCs expressed a diverse set of

122 GPCRs including 14 GPCRs that couple via Gα$_q$ proteins (GPCRs "GO:0004930" and PLC activating

Gα$_q$-coupled "GO:0007200", respectively). To determine the patterns of Gα$_q$-mediated calcium responses

in hOPCs, we used lentivirus to express the intracellular calcium reporter GCaMP6s hOPCs. We next

exposed independent cultures to receptor selective ligands at 10-fold the reported EC$_{50}$ concentration

and measured resultant calcium responses (**Fig. 1** and **Table 2**). All ligands induced a detectable

increase in intracellular calcium concentration measured by GCaMP6s fluorescence. However,

consistent with receptor heterogeneity, the majority of ligands tested increased calcium in only a small

fraction of GCaMP6-expressing hOPCs (between 1-30%). In contrast, activation of M$_{1/3}$R and mGluR$_5$

receptors by Oxo-M and CHPG, respectively, elicited robust responses in >46% of hOPCs (n = 3 human

samples). Both agonists induced a dose-dependent increase in the proportion of responding cells (**Fig.**

1C-D). The extent of total calcium flux was determined by measuring the area-under-curve (AUC). AUC

was defined as the area of the response above baseline beginning at the initial response until the end of

the imaging session (see methods). Raw AUC values were log$_{10}$-normalized to achieve normality and

displayed as mean ± SEM of pooled responses from >250 cells quantified per each dose from three

independent experiments using three independent human samples. Both agonists induced a clear dose-

dependent intracellular calcium response with EC$_{50}$ determined as 2.2 μM and 1.3 mM for Oxo-M and

CHPG respectively (**Fig. 1E-F**).

The calcium responses to Gα$_q$-specific agonists were further classified into monotonic and

oscillatory responses (**Table 2**). Intriguingly, of those ligands which resulted in calcium transients in <30%

of cells, we often observed that the majority of responses were monotonic and typically short-lived

returning to baseline within 3 mins following agonist application. In contrast, agonists for M$_{1/3}$R and

mGluR$_5$ receptors resulted in prolonged oscillatory calcium transients which persisted for over nine

minutes (9.5 ± 2.0 and 11.0 ± 0.6 mins, for Oxo-M and CHPG respectively; mean ± SEM, n=3 human

samples). For all doses tested, Oxo-M elicited oscillatory calcium signals in ~50% of all responsive cells

(**Supplementary Fig. S1A**). In comparison, cells responsive to CHPG treatment exhibited a significant

increase in the percentage of oscillatory responsive cells at higher concentrations (75μM vs 2300μM, RM

one way ANOVA with Holm-Sidak's post-hoc test, p<0.05) (**Supplementary Fig. S1B**). The number of

oscillations did not differ significantly between Oxo-M nor CHPG concentrations tested (**Supplementary**

Fig. S1C, D). There was a significant increase in maximum amplitude with Oxo-M dose (main effect,

p=0.03), which was not observed in CHPG treated cells (**Supplementary Fig. S1E, F**). Both ligands

induced oscillatory calcium signals with an average frequency of ~10 mHz (**Supplementary Fig. S1G,**

H). As such, the calcium responses following activation of Gα_q-coupled receptors were distinguishable

from one another, and the similarity of prolonged oscillations in the calcium response following mGluR$_5$

agonist treatment to M$_{1/3}$R suggested that a common downstream signaling pathway may be engaged

compared to other ligands.

As M$_{1/3}$R activation leads to a profound blockade of hOPC differentiation [12], we hypothesized that

the prolonged oscillatory calcium response may underlie this effect and that mGluR$_5$ activation would

therefore lead to a similar inhibition of differentiation. In addition, we predicted that EDNRB activation,

which instead elicits a minimal and transient calcium response, would not influence hOPC differentiation.

To test this hypothesis, we treated hOPCs with ET-1 or CHPG in the absence of mitogens and determined

the effect on O4$^+$ oligodendrocyte (OL) differentiation after 3-4 days (**Supplementary Fig. S2**). 150 nM

and 1.5 µM ET-1 did not influence the proportion of O4$^+$ OLs (RM One-way ANOVA with Holm-Sidak's

post-hoc test, p=0.64, n = 2 human samples) (**Supplemental Fig. S2A, B**). Furthermore, ET-1 did not

alter the generation of complex branched O4$^+$ oligodendrocytes (p=0.59) (**Supplemental Fig. S2C**). In

contrast, treatment with the mGluR$_5$ agonist CHPG [1.3mM] significantly attenuated O4$^+$ OL

differentiation from 42 ± 4% to 32 ± 5% (**Supplemental Fig. S2D, E**)**;** RM One-way ANOVA with Holm-

Sidak's post-hoc test, p=0.01, n=5 human samples). Furthermore, CHPG treatment significantly reduced

complex branch formation in differentiated O4$^+$ immature oligodendrocytes from 27 ± 4% to 18 ± 5%

(p=0.006) suggesting blockade of OL maturation (**Supplemental Fig. S2F**). These data suggest that

distinct patterns of calcium signaling occur following Gα_q-receptor activation and that calcium signaling

downstream of Gα_q-receptor activation may act to block hOPC differentiation and maturation.

SOCE is activated following M$_{1/3}$R agonist stimulation.

Oscillatory calcium responses can be sustained following endoplasmic reticulum (ER) store

release by several mechanisms including calcium recycling and, importantly, by store-operated calcium

entry (SOCE) through the activation of calcium release activated channels (CRAC) (for review see[27]).

We hypothesized that CRAC channels were activated following M$_{1/3}$R Gα_q-receptor activation and store

depletion in hOPCs. Calcium responses in hOPCs were measured following infection with lentivirus

encoding calcium reporter GCaMP6s [12]. To investigate whether SOCE occurs following ER Ca^{2+}

depletion in hOPCs, we treated cells in the absence of extracellular Ca^{2+} with thapsigargin (Tg) [10µM],

a sarco-endoplasmic reticulum Ca^{2+}-ATPases inhibitor (SERCA) blocker [43] (**Fig. 1G & Supplemental**

Fig. S3A). After the initial peak in calcium due to store depletion, we observed a clear SOCE associated

peak following the reintroduction of calcium into the media. Pre-incubation with CRAC channel inhibitors

2-APB [44] or MRS-1845 [45] substantially reduced SOCE in hOPCs (**Fig. 1G & Supplemental Fig. S3C,**

E). We quantified SOCE as the area under the calcium curve (AUC) following calcium re-addition (n = 2

independent human sample preparations with ≥ 44 cells quantified per condition/replicate). We observed

substantial SOCE following ER depletion with Tg (254.5 ± 51.3 AUC arbitrary units) which was nearly

abolished by pre-incubation with 2-APB or MRS-1845 (**Fig. 1H;** RM one-way ANOVA with Holm-Sidak's

post-hoc test, p<0.05; 2-APB: 0.2 ± 0.02; MRS: 7.1± 4.3**).** There were no differences in the initial cytosolic

calcium release following Tg induced ER depletion in each condition (RM one-way ANOVA, AUC,

p=0.42). Together these results were consistent with robust SOCE in hOPCs following ER-depletion that

could be blocked by pharmacological CRAC channel antagonists.

To test the hypothesis that M$_{1/3}$R activation leads directly to SOCE, we assessed SOCE following

muscarinic agonist treatment. Similar to Tg treatment, Oxo-M [25µM] induced robust SOCE following

calcium reintroduction (**Fig. 1G-H & Supplemental Fig. S3B;** 326 ± 67.72 AUC, n = 3 human sample,

with ≥48 cells quantified per condition/replicate). Likewise, pre-incubation with 2-APB similarly resulted

in a substantial reduction of SOCE (**Fig. 1G-H & Supplemental Fig. S3D;** RM one-way ANOVA with

Holm-Sidak's post-hoc test, p=0.037; Oxo-M+2-APB: 9.8 ± 3.9). While not significant, we also observed

a ~50% reduction in SOCE following pre-incubation with MRS (**Fig. 1G-H & Supplemental Fig. S3F,**

Oxo-M+MRS: 169.3 ± 53.8, p=0.1). As above, there were no differences in the magnitude of intracellular

calcium release following ER-depletion between all Oxo-M conditions tested (RM one-way ANOVA,

p=0.2). Together, these results show that robust SOCE occurs following M$_{1/3}$R induced signaling in

hOPCs.

STIM1/2 expression is necessary for muscarinic receptor induced SOCE.

The permeability of CRAC channels are regulated by STIM calcium sensor proteins [27]. RNA-seq

analyses of human OPCs revealed expression of both calcium sensors STIM1 and STIM2 (STIM1: 8.69

± 0.004 and STIM2: 30.0 ± 0.56; FPKM ± SEM, n=2) and was similar to that observed previously in

mouse OPCs [46] (Stim1: 13.9 ± 1.4 and Stim2: 14.9 ± 0.7). We hypothesized that the calcium sensors

STIM1 and STIM2 would be directly involved in regulating calcium influx through CRAC channels

following Gα$_q$-receptor activation in hOPCs. To test this hypothesis, we used siRNA targeting STIM1

and/or STIM2 coupled with calcium imaging. We observed a 70-85% decrease in mRNA expression of

STIM1 (**Fig. 2A;** one sample t-test p<0.0001) and STIM2 (**Fig. 2B;** STIM2 & STIM1/2: p<0.0001)

following STIM knockdown (KD) in hOPCs. Interestingly, we also observed a small but significant

decrease in STIM1 expression (p<0.05) following STIM2 siRNA transfection, as well as a reduction in

STIM2 expression following STIM1 siRNA (p<0.01). To directly assess the role of STIM proteins in

muscarinic receptor-induced SOCE, we assessed calcium influx following extracellular calcium removal

and re-addition in the context of STIM1/2 KD and Oxo-M treatment. As shown above we identified robust

SOCE downstream Oxo-M stimulation which was effectively blocked by CRAC antagonist 2-APB (n=4

independent human fetal sample preparations, RM one-way ANOVA with Holm-Sidak's post-hoc test,

p=0.003). STIM1 and STIM1/2 KD significantly reduced SOCE by ~40% (STIM1: p=0.047; STIM1/2: p =

0.041) while STIM2 had no effect (p=0.19) (**Fig. 2C-E**). These results suggest that STIM1 contributes to

muscarinic-induced SOCE and is a key component of the endogenous hOPC CRAC apparatus. As

STIM1/2 silencing may influence endogenous ER-store calcium content, we analyzed the total calcium

flux (AUC) and maximum amplitude of ER calcium store depletion in response to Oxo-M following

STIM1/2 KD in the absence of extracellular calcium. RM one-way ANOVA revealed no significant

differences in either the AUC (**Supplemental Fig. S4A**, Holm-Sidak's post-hoc tests, p=0.1) or maximum

amplitude (**Supplemental Fig. S4B**, p>0.4) of ER store release following STIM KD compared to control

cells, indicating that STIM silencing did not influence basal ER calcium store content or the extent of ER-

calcium depletion following G$_q$ stimulation.

We next sought to investigate the effects of STIM1 and STIM2 KD and their roles in regulation of

ligand induced Ca^{2+} oscillations in the presence of extracellular calcium. While STIM1 KD induced a

greater reduction in Ca^{2+} release following Oxo-M treatment, knockdown of STIM1 or STIM2 significantly

attenuated the magnitude of Ca^{2+} release (**Fig. 2F**; linear model with Tukey's HSD posttest, p<0.0001,

n=4 human samples). Intriguingly, STIM2 KD also resulted in a significant reduction in the frequency of

Ca^{2+} oscillations (**Fig. 2G**; STIM2: p= 0.002; STIM1/2: p=0.0323) while STIM1 KD had no effect (STIM1:

p = 0.7236). We also observed that neither STIM1 KD nor STIM2 KD altered the percentage of oscillatory

responsive cells (**Supplemental Fig. S5A,** RM one-way ANOVA, p=0.45) or the maximum oscillatory

peak amplitude (**Supplemental Fig. S5B,** p=0.1). Interestingly, the oscillatory response duration of Oxo-

M induced calcium responses increased following STIM2 KD (p<0.01) and 2-APB treatment (p<0.0001)

compared to control cells (**Supplemental Fig. S5C**). In summary, these results suggest that STIM1 is

required for Gα$_q$-induced SOCE while STIM2 specifically modulates the frequency of Ca^{2+} oscillations

following ligand-mediated SOCE.

STIM2 is required for human oligodendrocyte progenitor differentiation.

To determine whether STIM-dependent calcium oscillations may directly contribute to muscarinic

M$_{1/3}$R-mediated inhibition of hOPC differentiation, we assessed the independent effects of STIM1 and

STIM2 siRNA treatment and their interactions with Oxo-M treatment on hOPC differentiation *in vitro* by

three-way ANOVA (**Fig. 3A**). As shown previously[15], Oxo-M treatment significantly blocked differentiation

into O4$^+$ immature OLs (F(1,3)=14.31, p=0.032, n=4 human samples). Both siRNA control and STIM1

siRNA transfected hOPCs showed reduced differentiation in response to Oxo-M (**Fig. 3B;** Holm-Sidak's

post-hoc test, control: p = 0.006; STIM1: p=0.015). Interestingly, STIM2 siRNA transfection alone or in

combination with STIM1 abolished the anti-differentiative effects of Oxo-M (p > 0.2). This is consistent

with a role of STIM2 downstream of Oxo-M mediated signaling.

In addition to Oxo-M dependent effects, STIM2 siRNA alone caused a significant reduction in O4$^+$

OL differentiation (RM three-way ANOVA, STIM2 main effect $F(1,3)$ = 28.64, p = 0.013, n=4 human

samples) while STIM1 siRNA had no effect (p = 0.42). In the absence of Oxo-M, we observed a significant

decrease in O4$^+$ OL differentiation cells following STIM2 siRNA transfection relative to siControl

(p=0.002). Interestingly, this effect was not observed when STIM1/2 siRNA were combined (p>0.05).

These results suggested that STIM2 signaling in the absence of Gα_q agonist regulates hOPC

differentiation and is required for the anti-differentiative effects of muscarinic agonism. We further

examined the effect of STIM1/2 on morphological maturation by assessing the proportion of O4$^+$ cells

with simple or complex process elaboration as described [47] (**Supplemental Fig. S6A**). STIM1/2 KD did

not influence the percentage of simple O4$^+$ oligodendrocytes (3-way ANOVA, p > 0.3). Oxo-M reduced

the proportion of complex O4$^+$ cells following transfection with either control or STIM1 siRNA but did not

influence complex O4$^+$ cell number following STIM2 KD (Holm-Sidak's post-hoc test, p < 0.05)

(**Supplemental Fig. S6C**). Interestingly, in the absence of Oxo-M, STIM2 KD reduced the percentage of

complex O4$^+$ cells (Holm-Sidak's post-hoc test, p =0.0013), while STIM1 KD had no effect (p=0.61).

As differentiation and OPC proliferation are intimately linked, we next examined whether ER

calcium sensors STIM1/2 influenced hOPC proliferation. We quantified the proportion of dividing Ki67$^+$

OPCs following transfection with STIM1/2 siRNA or control siRNA (**Fig. 3C**). Similar to previous

observations [15], Oxo-M had no effect on hOPC proliferation. STIM1 or STIM2 siRNA treatment did not

significantly influence proliferation (n = 3 human samples; three-way ANOVA, STIM1 $F(1, 2)$ = 12.57, p

= 0.071, STIM2 $F(1, 2)$ = 6.27, p = 0.12). Taken together, these data suggest that STIM2 appears to be

necessary for OL differentiation and morphological maturation and may be required for the effects of

muscarinic agonist-mediated inhibition.

Optogenetically-induced SOCE blocks hOPC differentiation and promotes proliferation.

Our previous results suggest that STIM1/2 mediated SOCE may be required for sustained calcium

release following Gα_q stimulation and thereby contribute to muscarinic inhibition of differentiation. We

therefore sought to test whether oscillatory calcium SOCE directly regulates hOPC development. To this

end, we expressed an optogenetically controllable STIM protein in hOPCs to enable blue-light control of

SOCE in the absence of Gα_q-ligand stimulation [37]. Following infection with lentiviral OptoSTIM, hOPCs

were transduced with lentiviral jRCaMP1a, a red-shifted calcium indicator, to simultaneously measure

intracellular Ca^{2+} while induced SOCE [35]. Blue light activation of OptoSTIM resulted in a consistent and

robust oscillatory calcium influx which could be sustained for several hours in a frequency-specific

manner (**Fig. 4A-B**). To avoid summation of OptoSTIM-induced Ca^{2+} release and ensure a distinct

oscillatory wave form, based on the dissociation kinetics of OptoSTIM [37], we selected an upper frequency

of 3.33 mHz to model that of 10mHz waves observed following muscarinic receptor activation. As a

control, we selected a lower frequency of 0.55 mHz which is below the lower threshold associated with

frequency-dependent decoding in other non-excitable cells [28,48]. We matched the total blue-light exposure

over time between frequencies by pulsing for 330 and 2000 ms seconds at 3.33 and 0.55 mHz

respectively. Based on jRCaMP1a imaging, oscillatory SOCE could be maintained for at least three hours

in hOPCs *in vitro* and was clearly observable at 18 hours with 2000 ms pulses without loss of signal (**Fig.**

4B, n = 3 human samples). Importantly, we found that the total calcium release defined by AUC over the

2.5 h period was equivalent between groups (**Fig. 4C**; RM one-way ANOVA, Holm-Sidak's post-hoc test,

p = 0.49). As such, we could assess frequency dependent SOCE effects while controlling for total calcium

flux.

We examined the effect of optogenetically-induced oscillatory SOCE on hOPC proliferation and

differentiation (**Fig. 5**). Using time lapse microscopy and concurrent intermittent blue-light stimulation to

induce SOCE, we assessed the number of mitotic events (**Fig. 5A**) that occurred in the presence of both

high and low frequency OptoSTIM1-induced SOCE. Induction of high frequency (3.33 mHz) oscillatory

calcium entry produced a clear increase in the number of cumulative mitotic events at 24h and 36h post

stimulation respectively (**Fig. 5B**, quantified in **C**; RM one-way ANOVA with Holm-Sidak's post-hoc test,

p = 0.006, n = 4 human samples). After 36 hours of stimulation, high frequency SOCE resulted in a 40%

increase rate of cell division relative to unstimulated control hOPCs (3.33 mHz: 25.6 ± 4.9% vs. control:

19.1 ± 4.7%, p = 0.014) (**Fig. 5C**). In contrast, low frequency SOCE had no effect on proliferation (0.55

mHz; 17.6 ± 5.9%, p = 0.38). Next, we investigated the effects of long term OptoSTIM induction on hOPC

differentiation (**Fig. 5D**). We observed no gross alterations in transduced cell number following blue-light

stimulation (**Fig. 5E**; RM one-way ANOVA with Holm-Sidak's post-hoc test, p = 0.90, n = 4 human

samples). Importantly, induction of high frequency SOCE substantially impaired hOPC differentiation by

50% relative to unstimulated control hOPCs (**Fig. 5F**; 3.33 mHz: 7.5 ± 1.7% O4$^+$/GFP$^+$ cells vs. control:

16.6 ± 3.4, p = 0.02). Differentiation by hOPCs which received induction of low frequency SOCE was not

significantly different from unstimulated control cells (0.55 mHz: 12.83 ± 3.3%). Moreover, there was a

significant reduction in the number of O4$^+$ cells with complex branch morphology (>3 processes) following

3.33 mHz stimulation compared to control cells (**Fig. 5G;** RM one-way ANOVA with Holm-Sidak's post-

hoc test, p = 0.04). This suggests that induction of oscillatory SOCE was sufficient to inhibit hOPC

morphological maturation. Collectively these results indicate that sustained high frequency oscillatory

SOCE was sufficient to both block hOPC differentiation and augment hOPC proliferation.

DISCUSSION

In the present study, we demonstrate that a subset of Gα$_q$-receptors expressed by human primary

OPCs, including muscarinic M$_1$ and M$_3$ receptors (M$_{1/3}$R) and metabotropic-glutamate-receptor-5

(mGluR5), induced prolonged oscillatory calcium signaling and that the ability to induce oscillatory

calcium signaling correlated with their capacity to block differentiation. We found that a shared

mechanism downstream muscarinic Gα$_q$-receptor induced oscillatory signaling was the activation of store

operated calcium entry (SOCE). Inhibition with SOCE antagonists or knockdown of ER-localized SOCE

calcium sensors STIM1/2 reduced the magnitude of calcium signaling induced by muscarinic agonist

stimulation. STIM2 ablation selectively reduced calcium oscillation frequency and spontaneous OL

differentiation *in vitro* and, unlike control OPCs, treatment with muscarinic ligand did not further reduce

differentiation following STIM2 knockdown. While SOCE exerted a major influence on calcium

oscillations, as these experiments were performed using constant Gα$_q$-receptor agonist exposure, we

cannot exclude the requirement of IP$_3$-mediated release for the maintenance of prolonged calcium

oscillations. Together, these data suggested a functional linkage between muscarinic receptor signaling

and SOCE. Using an optogenetic approach to selectively activate SOCE in OPCs, we found that SOCE

was sufficient to modulate OL differentiation in the absence of Gα$_q$ dependent signaling suggesting that

SOCE is a key facet that impairs OL differentiation and remyelination. Furthermore, the frequency of

SOCE rather than the overall amount of calcium influx due to SOCE was found to be vital in mediating

the effect of calcium entry on differentiation and proliferation. This suggests that the calcium pathway

includes frequency-dependent 'decoders' which regulate OL differentiation and that modulation of

calcium frequency might represent an important mechanism by which diverse signals are integrated by

parenchymal OPCs.

In drug development, G-protein coupled receptors (GPCRs) are attractive candidates for

therapeutic intervention and recently many small molecules that target GPCRs have been identified that

modulate OPC function and OL differentiation to promote myelin repair [49]. For example, GPR17 which

largely couples via Gα$_{i/o}$ was found to negatively regulate OPC differentiation while knockout was

observed to promote remyelination [50]. Muscarinic receptor antagonists have displayed a remarkable

capacity to improve OL differentiation and remyelination in various animal models. Using genetic

approaches, the effects of these drugs have been localized to M$_1$R and M$_3$R. Both of these receptors

represent canonical Gα$_q$-coupled receptors. Indeed, when Gα coupling was systematically analyzed

across different subunits, M$_3$R displayed exquisite selectivity for rapid activation for Gα$_q$/Gα$_{11}$ coupling [51].

In this study, we focused on Gα$_q$-coupled receptors to determine if a shared characteristic of Gα$_q$

receptors could be used to predict their ability to prevent efficient OL differentiation and thereby impair

remyelination. We analyzed RNA-seq analysis of primary hOPC to identity those Gα$_q$-coupled receptors

expressed at the mRNA level and hypothesized that receptor-specific ligand activation would produce

canonical oscillatory calcium signaling and block hOPC differentiation. Surprisingly, only a small subset

of pharmacological Gα_q ligands were capable of inducing prolonged calcium oscillation in hOPCs.

Intriguingly, this correlated with the ability of ligands to effectively block OL differentiation. CHPG, which

activates mGluR$_5$ in a highly selective manner, was the only other ligand capable of inducing calcium

oscillations in a large fraction of hOPCs. The specificity of Gα_q-receptor mediated signaling is consistent

with previous results which have shown that while endothelin (ET) agonist treatment induces a monotonic

calcium response [24] while metabotropic glutamate receptor agonists induce calcium oscillations [19].

mGluR$_5$ couples primarily via Gα_q and does not have appreciable activity at Gα_i. In a similar manner to

M$_3$R, mGluR$_5$ mRNA was relatively highly expressed in rodent OPCs and down-regulated following

differentiation [19]. Herein, we found that treatment with CHPG resulted in a dose-dependent blockade of

hOPC differentiation.

In contrast, the other Gα_q couple receptors that elicited calcium responses in OPCs induced

calcium release that was either considerably shorter in duration and/or were largely monotonic in nature.

These differences could be due to differences in overall level of expression and/or differences in receptor

coupling. From a large-scale screen of G-protein coupling [52], we observed that many of these receptors

were promiscuously coupled with other Gα subunits. For example, sphingosine-1-phosphate-receptor-1

(S1PR1) was observed as primarily coupled via Gα_i and Gα_o with little evidence of Gα_q coupling. This

difference corresponded with the observed function difference following activation of S1PR1 which results

in OL differentiation from human OPCs [53]. Furthermore, endothelin-B receptor (ENDRB), thrombin

receptor II (F2R) and purinergic receptor Y1 (P2RY1) also show promiscuous coupling with multiple Gα

proteins upon activation. Together this suggests that selectively in coupling may influence the net

influence on downstream signaling and the nature of the resultant calcium response. As the coupling of

GPCRs to specific Gα proteins are not absolute and in many cases is promiscuous, we anticipate that

biased ligands or antagonists capable of promoting or blocking coupling along specific pathways may

represent optimal approaches to promote OPC differentiation.

The mechanisms which shape the calcium-response following ligand application are complex and

likely dependent on multiple interacting proteins. Among these, Gα subunits form specific complexes with

their cognate G$\beta\gamma$ proteins which possess their own signaling capability and modulate the signaling of Gα

subunits they interact with [54]. Among these, specific isoforms of phosphoinositide 3-kinase (PI3K) and

adenylate cyclase, voltage-operated calcium channels (VOCC) and inwardly rectifying K$^+$ channel can

be regulated by G$\beta\gamma$. Activation of PI3K/Akt [55], adenylate cyclase [56], and VOCC [57] are associated with

promotion of OPC differentiation suggesting that G$\beta\gamma$ activation may counteract oscillatory calcium

signaling. For muscarinic receptor signaling, in rat pituitary cells G$_{\beta3}$ is necessary for the carbachol-

mediated calcium current [58], while in a rat leukemia cell line G$_{\beta 1}$ and G$_{\beta 4}$ were more important in the

regulation of this current [59]. In primary hOPCs, G$_{\beta 4}$ ablation did not influence calcium signaling following

Oxo-M treatment [60], though redundancy with G$_{\beta 1}$ and G$_{\beta 2}$ is possible as these subunits are expressed at

substantially higher levels in human, rat, and mouse OPCs.

In addition to the G$_{\beta\gamma}$ subunit specific regulation, receptor-specific activation of regulators of G-

protein signaling (RGS) proteins provide specificity for G-protein coupling and modulate the kinetics of

second messenger activation [61]. RGS proteins promote termination of GPCR signaling via GTPase

activation and prevent GPCR signaling in the absence of ligand. Following ligand activation and elevated

Ca^{2+}, RGS proteins exhibit a cyclical pattern of association with phosphatidylinositol-3,4,5-trisphosphate

(PIP$_3$) and activated calcium/calmodulin (Ca^{2+}/CaM) followed by re-association with the GPCR complex

after Ca^{2+} levels return to baseline. This allows a second cycle to start and initiate another calcium spike.

The kinetics of these steps, especially the slow dissociation of RGS protein from Ca^{2+}/CaM, contributes

to the resultant frequency of calcium oscillations [62]. While the role of specific RGS proteins in the OPC

and OL lineage is poorly understood, several RGS proteins are expressed in OPCs [46]. Among these,

RGS4 has been shown to be a potent negative regulator of muscarinic receptor signaling in insulinoma

cells [63]. Also noteworthy, the R7 family of RGS proteins (RGS6/7/9/11) form heterodimers with G-protein

β subunit Gβ5 (GNB5) which similarly attenuates M$_3$R-dependent Ca^{2+} signaling [64].

Once activated, SOCE is associated with many cellular functions in neural progenitors such as

proliferation [65] and differentiation [66,67]. In OPCs, SOCE can modulate PDGF-AA induced OPC

proliferation [30]. However, given the known role of PI3K/Akt signaling downstream of PDGF receptor,

SOCE is unlikely to be a critical component of PDGF signaling. Likewise, the cellular role of calcium

oscillations in OPC soma is not yet known. Our data using optogenetic induction of SOCE strongly

suggest that oscillatory SOCE alone is sufficient to modulate both OPC proliferation and differentiation.

While we were limited by the dissociation kinetics of OptoSTIM to an upper limit of 3.33 mHz stimulation,

this frequency fits within the known frequencies of calcium oscillations in non-excitable cells and may

recruit NFAT, NFκB and other downstream pathways [48]. By matching total calcium influx during the period

of exposure, we were able to demonstrate frequency dependence as a substantially lower frequency

(0.55 mHz) was unable to modulate proliferation or differentiation. New approaches will be required to

determine whether higher frequency SOCE can further alter these critical cellular processes.

SOCE itself is dependent on the association of CRAC subunit comprising STIM ER-calcium

sensors and ORAI channel subunits. While STIM1 and STIM2 contributed to different aspects of

muscarinic-induced SOCE and sustained oscillatory calcium signaling in hOPCs, we found that only

STIM2 knockdown reduced spontaneous OL differentiation. Previous reports have found that apart from

its role in agonist mediated SOCE, STIM2 signaling often persists to a lesser degree in the absence of

an ER-release stimulus. STIM2 responds to small fluctuations in ER calcium levels, and drives small local

SOCE calcium entry when local calcium levels fall below baseline (reviewed in [68]). It is also known that

STIM2 can regulate endogenous differentiation of other cell types through modulation of cytoplasmic

calcium levels following STIM1-induced oscillatory SOCE [69]. Furthermore, small spontaneous SOCE

calcium transients in the absence of agonist have been observed in mouse OPCs, and have been

implicated in the regulation of differentiation [70]. It is possible, therefore, that STIM2 is required for these

spontaneous transients and thus STIM2 KD may inhibit differentiation. In contrast, our data is consistent

with prolonged oscillatory SOCE, either produced optogenetically or in response to M$_3$R and mGluR$_5$

agonists, causing a blockade of hOPC differentiation via global calcium-mediated changes. Additional

studies will be necessary to tease apart the specific mechanisms by which STIM2 directly influences

oligodendrocyte differentiation.

The ER-calcium sensors STIM1/2 physically and functionally interact with one another to regulate

SOCE [68]. The relative abundance of STIM1 and STIM2 vary greatly between tissues and individual cell

types. In human OPCs, STIM2 is more than 3-fold more abundant than STIM1 mRNA but both genes are

relatively highly expressed (>10 FPKM). In contrast, both STIM1 and STIM2 are expressed at equivalent

levels in mouse OPCs [46]. As the differences in the relative expression of STIM1/2 influences the resultant

properties of SOCE in response to store depletion (reviewed in [71]), the differences in STIM1:STIM2 ratio

in mouse and human may therefore underlie some of the species-specific functional differences in OPC

behavior [72]. Both STIM1 and STIM2 have been shown to contribute to oscillatory SOCE in a context

dependent manner [73,74]. Our data support a specific role of STIM2 in the regulation of OPC calcium

oscillations and suggest that expression of STIM2 is required for muscarinic receptor induced

differentiation blockade in hOPCs. Together with the direct regulation of OPC differentiation via

optogenetically-gated SOCE, these data suggest that SOCE mediated via STIM activation will likely play

an important role in mediating the negative effects of Gα$_q$-coupled receptors capable of blocking

oligodendrocyte differentiation in health and disease.

REFERENCES

Franklin, R. J. M. & Ffrench-Constant, C. Regenerating CNS myelin - from mechanisms to experimental medicines. *Nature reviews. Neuroscience* **18**, 753-769, doi:10.1038/nrn.2017.136 (2017).

Faissner, S., Plemel, J. R., Gold, R. & Yong, V. W. Progressive multiple sclerosis: from pathophysiology to therapeutic strategies. *Nat Rev Drug Discov* **18**, 905-922, doi:10.1038/s41573-019-0035-2 (2019).

Kuhlmann, T. *et al.* Differentiation block of oligodendroglial progenitor cells as a cause for remyelination failure in chronic multiple sclerosis. *Brain : a journal of neurology* **131**, 1749-1758, doi:10.1093/brain/awn096 (2008).

Wolswijk, G. Chronic stage multiple sclerosis lesions contain a relatively quiescent population of oligodendrocyte precursor cells. *J Neurosci* **18**, 601-609 (1998).

Thornton, M. A. & Hughes, E. G. Neuron-oligodendroglia interactions: Activity-dependent regulation of cellular signaling. *Neuroscience letters* **727**, 134916, doi:10.1016/j.neulet.2020.134916 (2020).

Ortiz, F. C. *et al.* Neuronal activity in vivo enhances functional myelin repair. *JCI Insight* **5**, doi:10.1172/jci.insight.123434 (2019).

Gautier, H. O. *et al.* Neuronal activity regulates remyelination via glutamate signalling to oligodendrocyte progenitors. *Nat Commun* **6**, 8518, doi:10.1038/ncomms9518 (2015).

Bernstein, M., Lyons, S. A., Moller, T. & Kettenmann, H. Receptor-mediated calcium signalling in glial cells from mouse corpus callosum slices. *Journal of neuroscience research* **46**, 152-163, doi:10.1002/(SICI)1097-4547(19961015)46:2<152::AID-JNR3>3.0.CO;2-G (1996).

Paez, P. M. & Lyons, D. A. Calcium Signaling in the Oligodendrocyte Lineage: Regulators and Consequences. *Annual review of neuroscience* **43**, 163-186, doi:10.1146/annurev-neuro-100719-093305 (2020).

Berridge, M. J. & Irvine, R. F. Inositol phosphates and cell signalling. *Nature* **341**, 197-205, doi:10.1038/341197a0 (1989).

Mei, F. *et al.* Accelerated remyelination during inflammatory demyelination prevents axonal loss and improves functional recovery. *Elife* **5**, e18246, doi:10.7554/eLife.18246 (2016).

Welliver, R. R. *et al.* Muscarinic Receptor M3R Signaling Prevents Efficient Remyelination by Human and Mouse Oligodendrocyte Progenitor Cells. *J Neurosci* **38**, 6921-6932, doi:10.1523/JNEUROSCI.1862-17.2018 (2018).

Mei, F. *et al.* Micropillar arrays as a high-throughput screening platform for therapeutics in multiple sclerosis. *Nature medicine* **20**, 954-960, doi:10.1038/nm.3618 (2014).

631   14   Deshmukh, V. A. *et al.* A regenerative approach to the treatment of multiple sclerosis. *Nature*
632   **502**, 327-332, doi:10.1038/nature12647 (2013).
633   15   Abiraman, K. *et al.* Anti-muscarinic adjunct therapy accelerates functional human
634   oligodendrocyte repair. *J Neurosci* **35**, 3676-3688, doi:10.1523/JNEUROSCI.3510-14.2015
635   (2015).
636   16   Gadea, A., Aguirre, A., Haydar, T. F. & Gallo, V. Endothelin-1 regulates oligodendrocyte
637   development. *J Neurosci* **29**, 10047-10062, doi:10.1523/JNEUROSCI.0822-09.2009 (2009).
638   17   Agresti, C. *et al.* ATP regulates oligodendrocyte progenitor migration, proliferation, and
639   differentiation: involvement of metabotropic P2 receptors. *Brain Res Brain Res Rev* **48**, 157-
640   165, doi:10.1016/j.brainresrev.2004.12.005 (2005).
641   18   Takeda, M., Nelson, D. J. & Soliven, B. Calcium signaling in cultured rat oligodendrocytes. *Glia*
642   **14**, 225-236, doi:10.1002/glia.440140308 (1995).
643   19   Luyt, K., Varadi, A., Durant, C. F. & Molnar, E. Oligodendroglial metabotropic glutamate
644   receptors are developmentally regulated and involved in the prevention of apoptosis. *J*
645   *Neurochem* **99**, 641-656, doi:10.1111/j.1471-4159.2006.04103.x (2006).
646   20   Fan, L. W. *et al.* Exposure to serotonin adversely affects oligodendrocyte development and
647   myelination in vitro. *J Neurochem* **133**, 532-543, doi:10.1111/jnc.12988 (2015).
648   21   Jung, C. G. *et al.* Functional consequences of S1P receptor modulation in rat oligodendroglial
649   lineage cells. *Glia* **55**, 1656-1667, doi:10.1002/glia.20576 (2007).
650   22   Chattopadhyay, N. *et al.* Calcium receptor expression and function in oligodendrocyte
651   commitment and lineage progression: potential impact on reduced myelin basic protein in CaR-
652   null mice. *Journal of neuroscience research* **86**, 2159-2167, doi:10.1002/jnr.21662 (2008).
653   23   Coppi, E. *et al.* Adenosine A2B receptors inhibit K(+) currents and cell differentiation in cultured
654   oligodendrocyte precursor cells and modulate sphingosine-1-phosphate signaling pathway.
655   *Biochem Pharmacol* **177**, 113956, doi:10.1016/j.bcp.2020.113956 (2020).
656   24   Jung, K. J. *et al.* The role of endothelin receptor A during myelination of developing
657   oligodendrocytes. *J Korean Med Sci* **26**, 92-99, doi:10.3346/jkms.2011.26.1.92 (2011).
658   25   Yuen, T. J. *et al.* Identification of endothelin 2 as an inflammatory factor that promotes central
659   nervous system remyelination. *Brain : a journal of neurology* **136**, 1035-1047,
660   doi:10.1093/brain/awt024 (2013).
661   26   Dukala, D. E. & Soliven, B. S1P1 deletion in oligodendroglial lineage cells: Effect on
662   differentiation and myelination. *Glia* **64**, 570-582, doi:10.1002/glia.22949 (2016).
663   27   Prakriya, M. & Lewis, R. S. Store-Operated Calcium Channels. *Physiol Rev* **95**, 1383-1436,
664   doi:10.1152/physrev.00020.2014 (2015).

665   28   Smedler, E. & Uhlen, P. Frequency decoding of calcium oscillations. *Biochim Biophys Acta*
666   **1840**, 964-969, doi:10.1016/j.bbagen.2013.11.015 (2014).

667   29   Samanta, K. & Parekh, A. B. Spatial Ca(2+) profiling: decrypting the universal cytosolic Ca(2+)
668   oscillation. *J Physiol* **595**, 3053-3062, doi:10.1113/JP272860 (2017).

669   30   Paez, P. M. *et al.* Regulation of store-operated and voltage-operated Ca2+ channels in the
670   proliferation and death of oligodendrocyte precursor cells by golli proteins. *ASN Neuro* **1**,
671   doi:10.1042/AN20090003 (2009).

672   31   Schmidt, C., Ohlemeyer, C., Kettenmann, H., Reutter, W. & Horstkorte, R. Incorporation of N-
673   propanoylneuraminic acid leads to calcium oscillations in oligodendrocytes upon the application
674   of GABA. *Febs Letters* **478**, 276-280, doi:Doi 10.1016/S0014-5793(00)01868-8 (2000).

675   32   Paez, P. M., Fulton, D., Spreuer, V., Handley, V. & Campagnoni, A. T. Modulation of canonical
676   transient receptor potential channel 1 in the proliferation of oligodendrocyte precursor cells by
677   the golli products of the myelin basic protein gene. *J Neurosci* **31**, 3625-3637,
678   doi:10.1523/JNEUROSCI.4424-10.2011 (2011).

679   33   Conway, G. D., O'Bara, M. A., Vedia, B. H., Pol, S. U. & Sim, F. J. Histone deacetylase activity
680   is required for human oligodendrocyte progenitor differentiation. *Glia* **60**, 1944-1953,
681   doi:10.1002/glia.22410 (2012).

682   34   Pol, S. U. *et al.* Sox10-MCS5 enhancer dynamically tracks human oligodendrocyte progenitor
683   fate. *Experimental neurology* **247**, 694-702, doi:10.1016/j.expneurol.2013.03.010 (2013).

684   35   Dana, H. *et al.* Sensitive red protein calcium indicators for imaging neural activity. *Elife* **5**,
685   doi:10.7554/eLife.12727 (2016).

686   36   Sevin, C. *et al.* Intracerebral adeno-associated virus-mediated gene transfer in rapidly
687   progressive forms of metachromatic leukodystrophy. *Hum Mol Genet* **15**, 53-64,
688   doi:10.1093/hmg/ddi425 (2006).

689   37   Kyung, T. *et al.* Optogenetic control of endogenous Ca(2+) channels in vivo. *Nature*
690   *biotechnology* **33**, 1092-1096, doi:10.1038/nbt.3350 (2015).

691   38   Wang, J. *et al.* Paired Related Homeobox Protein 1 Regulates Quiescence in Human
692   Oligodendrocyte Progenitors. *Cell reports* **25**, 3435-3450 e3436,
693   doi:10.1016/j.celrep.2018.11.068 (2018).

694   39   Geraerts, M., Willems, S., Baekelandt, V., Debyser, Z. & Gijsbers, R. Comparison of lentiviral
695   vector titration methods. *BMC biotechnology* **6**, 34, doi:10.1186/1472-6750-6-34 (2006).

696   40   Edelstein, A., Amodaj, N., Hoover, K., Vale, R. & Stuurman, N. Computer control of
697   microscopes using microManager. *Current protocols in molecular biology* **Chapter 14**,
698   Unit14.20, doi:10.1002/0471142727.mb1420s92 (2010).

41   Stockley, J. H. *et al.* Surpassing light-induced cell damage in vitro with novel cell culture media.
Sci Rep **7**, 849, doi:10.1038/s41598-017-00829-x (2017).

42   Tinevez, J. Y. *et al.* TrackMate: An open and extensible platform for single-particle tracking.
Methods **115**, 80-90, doi:10.1016/j.ymeth.2016.09.016 (2017).

43   Treiman, M., Caspersen, C. & Christensen, S. B. A tool coming of age: thapsigargin as an
inhibitor of sarco-endoplasmic reticulum Ca(2+)-ATPases. *Trends in pharmacological sciences*
19, 131-135, doi:10.1016/s0165-6147(98)01184-5 (1998).

44   DeHaven, W. I., Smyth, J. T., Boyles, R. R., Bird, G. S. & Putney, J. W., Jr. Complex actions of
2-aminoethyldiphenyl borate on store-operated calcium entry. *The Journal of biological*
chemistry **283**, 19265-19273, doi:10.1074/jbc.M801535200 (2008).

45   Harper, J. L. *et al.* Dihydropyridines as inhibitors of capacitative calcium entry in leukemic HL-60
cells. *Biochem Pharmacol* **65**, 329-338, doi:10.1016/s0006-2952(02)01488-0 (2003).

46   Zhang, Y. *et al.* An RNA-sequencing transcriptome and splicing database of glia, neurons, and
vascular cells of the cerebral cortex. *J Neurosci* **34**, 11929-11947,
doi:10.1523/JNEUROSCI.1860-14.2014 (2014).

47   Huang, J. K. *et al.* Retinoid X receptor gamma signaling accelerates CNS remyelination. *Nat*
Neurosci **14**, 45-53, doi:10.1038/nn.2702 (2011).

48   Boulware, M. J. & Marchant, J. S. Timing in cellular Ca2+ signaling. *Curr Biol* **18**, R769-R776,
doi:10.1016/j.cub.2008.07.018 (2008).

49   Mogha, A., D'Rozario, M. & Monk, K. R. G Protein-Coupled Receptors in Myelinating Glia.
Trends in pharmacological sciences **37**, 977-987, doi:10.1016/j.tips.2016.09.002 (2016).

50   Ou, Z. *et al.* Olig2-Targeted G-Protein-Coupled Receptor Gpr17 Regulates Oligodendrocyte
Survival in Response to Lysolecithin-Induced Demyelination. *J Neurosci* **36**, 10560-10573,
doi:10.1523/JNEUROSCI.0898-16.2016 (2016).

51   Masuho, I. *et al.* Distinct profiles of functional discrimination among G proteins determine the
actions of G protein-coupled receptors. *Sci Signal* **8**, ra123, doi:10.1126/scisignal.aab4068
(2015).

52   Inoue, A. *et al.* Illuminating G-Protein-Coupling Selectivity of GPCRs. *Cell* **177**, 1933-1947
e1925, doi:10.1016/j.cell.2019.04.044 (2019).

53   Cui, Q. L., Fang, J., Kennedy, T. E., Almazan, G. & Antel, J. P. Role of p38MAPK in S1P
receptor-mediated differentiation of human oligodendrocyte progenitors. *Glia* **62**, 1361-1375,
doi:https://doi.org/10.1002/glia.22688 (2014).

54 Khan, S. M. *et al.* The expanding roles of Gbetagamma subunits in G protein-coupled receptor
signaling and drug action. *Pharmacological reviews* **65**, 545-577, doi:10.1124/pr.111.005603
(2013).
55 Goebbels, S. *et al.* Elevated phosphatidylinositol 3,4,5-trisphosphate in glia triggers cell-
autonomous membrane wrapping and myelination. *J Neurosci* **30**, 8953-8964,
doi:10.1523/jneurosci.0219-10.2010 (2010).
56 Raible, D. W. & McMorris, F. A. Induction of oligodendrocyte differentiation by activators of
adenylate cyclase. *Journal of neuroscience research* **27**, 43-46,
doi:https://doi.org/10.1002/jnr.490270107 (1990).
57 Cheli, V. T. *et al.* Conditional Deletion of the L-Type Calcium Channel Cav1.2 in
Oligodendrocyte Progenitor Cells Affects Postnatal Myelination in Mice. *J Neurosci* **36**, 10853-
10869, doi:10.1523/JNEUROSCI.1770-16.2016 (2016).
58 Kleuss, C., Scherübl, H., Hescheler, J., Schultz, G. & Wittig, B. Different β-subunits determine
G-protein interaction with transmembrane receptors. *Nature* **358**, 424-426,
doi:10.1038/358424a0 (1992).
59 Dippel, E., Kalkbrenner, F., Wittig, B. & Schultz, G. A heterotrimeric G protein complex couples
the muscarinic m1 receptor to phospholipase C-beta. *Proceedings of the National Academy of*
Sciences **93**, 1391-1396, doi:10.1073/pnas.93.4.1391 (1996).
60 Pol, S. U. *et al.* Network-Based Genomic Analysis of Human Oligodendrocyte Progenitor
Differentiation. *Stem cell reports* **9**, 710-723, doi:10.1016/j.stemcr.2017.07.007 (2017).
61 Kach, J., Sethakorn, N. & Dulin, N. O. A finer tuning of G-protein signaling through regulated
control of RGS proteins. *Am J Physiol Heart Circ Physiol* **303**, H19-35,
doi:10.1152/ajpheart.00764.2011 (2012).
62 Luo, X., Popov, S., Bera, A. K., Wilkie, T. M. & Muallem, S. RGS Proteins Provide Biochemical
Control of Agonist-Evoked [Ca2+]i Oscillations. *Molecular cell* **7**, 651-660,
doi:https://doi.org/10.1016/S1097-2765(01)00211-8 (2001).
63 Ruiz de Azua, I. *et al.* RGS4 is a negative regulator of insulin release from pancreatic β-cells in
vitro and in vivo. *Proceedings of the National Academy of Sciences* **107**, 7999-8004,
doi:10.1073/pnas.1003655107 (2010).
64 Karpinsky-Semper, D., Volmar, C. H., Brothers, S. P. & Slepak, V. Z. Differential effects of the
Gβ5-RGS7 complex on muscarinic M3 receptor-induced Ca2+ influx and release. *Mol*
Pharmacol **85**, 758-768, doi:10.1124/mol.114.091843 (2014).
65 Domenichini, F. *et al.* Store-Operated Calcium Entries Control Neural Stem Cell Self-Renewal in
the Adult Brain Subventricular Zone. *Stem Cells* **36**, 761-774, doi:10.1002/stem.2786 (2018).

66   Forostyak, O. *et al.* Physiology of Ca(2+) signalling in stem cells of different origins and
differentiation stages. *Cell Calcium* **59**, 57-66, doi:10.1016/j.ceca.2016.02.001 (2016).

67   Gopurappilly, R., Deb, B. K., Chakraborty, P. & Hasan, G. Stable STIM1 Knockdown in Self-
Renewing Human Neural Precursors Promotes Premature Neural Differentiation. *Front Mol*
Neurosci **11**, 178, doi:10.3389/fnmol.2018.00178 (2018).

68   López, E., Salido, G. M., Rosado, J. A. & Berna-Erro, A. Unraveling STIM2 function. *J Physiol*
Biochem **68**, 619-633, doi:10.1007/s13105-012-0163-1 (2012).

69   Darbellay, B. *et al.* Human muscle economy myoblast differentiation and excitation-contraction
coupling use the same molecular partners, STIM1 and STIM2. *J Biol Chem* **285**, 22437-22447,
doi:10.1074/jbc.M110.118984 (2010).

70   Rui, Y., Pollitt, S. L., Myers, K. R., Feng, Y. & Zheng, J. Q. Spontaneous Local Calcium
Transients Regulate Oligodendrocyte Development in Culture through Store-Operated Ca(2+)
Entry and Release. *eNeuro* **7**, doi:10.1523/ENEURO.0347-19.2020 (2020).

71   Berna-Erro, A., Jardin, I., Salido, G. M. & Rosado, J. A. Role of STIM2 in cell function and
physiopathology. *J Physiol* **595**, 3111-3128, doi:10.1113/jp273889 (2017).

72   Dietz, K. C., Polanco, J. J., Pol, S. U. & Sim, F. J. Targeting human oligodendrocyte progenitors
for myelin repair. *Experimental neurology* **283**, 489-500, doi:10.1016/j.expneurol.2016.03.017
(2016).

73   Kar, P., Bakowski, D., Di Capite, J., Nelson, C. & Parekh, A. B. Different agonists recruit
different stromal interaction molecule proteins to support cytoplasmic Ca2+ oscillations and
gene expression. *Proc Natl Acad Sci U S A* **109**, 6969-6974, doi:10.1073/pnas.1201204109
(2012).

74   Thiel, M., Lis, A. & Penner, R. STIM2 drives Ca2+ oscillations through store-operated Ca2+
entry caused by mild store depletion. *J Physiol* **591**, 1433-1445,
doi:10.1113/jphysiol.2012.245399 (2013).

75   Harriss, D. R., Marsh, K. A., Birmingham, A. T. & Hill, S. J. Expression of muscarinic M3-
receptors coupled to inositol phospholipid hydrolysis in human detrusor cultured smooth muscle
cells. *J Urol* **154**, 1241-1245 (1995).

76   Doherty, A. J., Palmer, M. J., Henley, J. M., Collingridge, G. L. & Jane, D. E. (RS)-2-chloro-5-
hydroxyphenylglycine (CHPG) activates mGlu5, but no mGlu1, receptors expressed in CHO
cells and potentiates NMDA responses in the hippocampus. *Neuropharmacology* **36**, 265-267,
doi:10.1016/s0028-3908(97)00001-4 (1997).

77   Pennington, L. D. *et al.* 4-Methoxy-N-[2-(trifluoromethyl)biphenyl-4-ylcarbamoyl]nicotinamide: A
Potent and Selective Agonist of S1P1. *ACS Med Chem Lett* **2**, 752-757, doi:10.1021/ml2001399
(2011).

78   Vassallo, R. R., Jr., Kieber-Emmons, T., Cichowski, K. & Brass, L. F. Structure-function
relationships in the activation of platelet thrombin receptors by receptor-derived peptides. *The*
Journal of biological chemistry **267**, 6081-6085 (1992).

79   Calleri, E. *et al.* Frontal affinity chromatography-mass spectrometry useful for characterization of
new ligands for GPR17 receptor. *J Med Chem* **53**, 3489-3501, doi:10.1021/jm901691y (2010).

80   Kawanabe, Y., Okamoto, Y., Enoki, T., Hashimoto, N. & Masaki, T. Ca(2+) channels activated
by endothelin-1 in CHO cells expressing endothelin-A or endothelin-B receptors. *American*
journal of physiology. Cell physiology **281**, C1676-1685, doi:10.1152/ajpcell.2001.281.5.C1676
(2001).

81   Chhatriwala, M. *et al.* Induction of novel agonist selectivity for the ADP-activated P2Y1 receptor
versus the ADP-activated P2Y12 and P2Y13 receptors by conformational constraint of an ADP
analog. *The Journal of pharmacology and experimental therapeutics* **311**, 1038-1043,
doi:10.1124/jpet.104.068650 (2004).

82   McGuire, J. J., Saifeddine, M., Triggle, C. R., Sun, K. & Hollenberg, M. D. 2-furoyl-LIGRLO-
amide: a potent and selective proteinase-activated receptor 2 agonist. *The Journal of*
pharmacology and experimental therapeutics **309**, 1124-1131, doi:10.1124/jpet.103.064584
(2004).

83   Tabarean, I. V. Functional pharmacology of H1 histamine receptors expressed in mouse
preoptic/anterior hypothalamic neurons. *Br J Pharmacol* **170**, 415-425, doi:10.1111/bph.12286
(2013).

84   Console-Bram, L., Brailoiu, E., Brailoiu, G. C., Sharir, H. & Abood, M. E. Activation of GPR18 by
cannabinoid compounds: a tale of biased agonism. *Br J Pharmacol* **171**, 3908-3917,
doi:10.1111/bph.12746 (2014).

85   Hinz, S., Lacher, S. K., Seibt, B. F. & Muller, C. E. BAY60-6583 acts as a partial agonist at
adenosine A2B receptors. *The Journal of pharmacology and experimental therapeutics* **349**,
427-436, doi:10.1124/jpet.113.210849 (2014).

86   Meis, S. *et al.* NF546 [4,4'-(carbonylbis(imino-3,1-phenylene-carbonylimino-3,1-(4-methyl-
phenylene)-car bonylimino))-bis(1,3-xylene-alpha,alpha'-diphosphonic acid) tetrasodium salt] is
a non-nucleotide P2Y11 agonist and stimulates release of interleukin-8 from human monocyte-
derived dendritic cells. *J Pharmacol Exp Ther* **332**, 238-247, doi:10.1124/jpet.109.157750
(2010).

TABLES

Primer	Sequence 5'-3'
WPRE	Forward: CGCTGCTTTAATGCCTTTGTAT
	Reverse: GGGCCACAACTCCTCATAAA
GAPDH	Forward: GTGAAGGTCGGAGTCAACGG
	Reverse: CCTGGAAGATGGTGATGGGA
STIM1	Forward: CCTCTCTTGACTCGCCATAATC
	Reverse: CTTGGAGTAACGGTTCTGGATATAG
STIM2	Forward: CAACACACACCACACTCCTT
	Reverse: GCCTCTTCCTCCTCTTCATTTC

833 **Table 1: qPCR primers.**

Gene Symbol	FPKM (hOPC; mOPCs)	Agonist/ Doses tested	Reported EC_{50}	% Responsive	Maximum amplitude	Duration (min)	% Oscillatory
CHRM3	16.3 ± 2.0; 0.27 ± 0.11	Oxo-M [25 µM]	2.5 µM [75]	67.6 ± 10.0	2.17 ± 0.06	6.1 ± 0.8	58.3 ± 20.5
CHRM1	4.0 ± 0.9; 2.06 ± 0.66						
GRM5	1.2 ± 0.004; 13.23 ± 3.63	CHPG [7.5 mM]	750µM [76]	46.9 ± 9.7	0.87 ± 0.03	6.7 ± 0.1	48.6 ± 4.1
S1PR1	3.1 ± 0.3; 34.32 ± 8.80	TC-G 1006 [350 nM]	35 nM [77]	27.8 ± 2.3	0.6 ± 0.02	2.3 ± 0.2	24.8 ± 14.1
F2R	26.7 ± 0.1; 23.54 ± 3.31	TRAP-6 [8.0 µM]	800 nM [78]	21.2 ± 13.3	1.0 ± 0.10	2.8 ± 0.5	38.9 ± 7.3
GPR17	3.6 ± 1.4; 440.98 ± 42.13	PF9 [400 pM]	36 pM [79]	19.3 ± 0.7	0.6 ± 0.01	2.3 ± 0.3	17 ± 10.3
EDNRB	17.3 ± 01; 292.72 ± 10.54	ET-1 [150 nM]	10 nM [80]	16.8 ± 4.3	1.03 ± 0.09	2.4 ± 0.3	28.8 ± 5.6
P2RY1	1.0 ± 0.2; 4.94 ± 1.46	MRS 2365 [12 nM]	1.2 nM [81]	14.6 ± 8.4	2.9 ± 0.20	9.3 ± 0.9	48.3 ± 0.3
HTR2B	0.8 ± 0.2; 0.12 ± 0.02	5-HT [100 µM]	* 10 µM [20]	13.0 ± 12.4	0.7 ± 0.08	2.1 ± 0.1	14.3 ± 14.3
F2RL1	1.3 ± 0.02; 0.10 ± 0.10	2-FLY [1.0 µM]	100 nM [82]	8.5 ± 1.3	1.6 ± 0.16	11.2 ± 0.2	63.7 ± 7.0
HRH1	0.4 ± 0.1; 5.01 ± 0.49	2-PED [560 µM]	56 µM [83]	7.4 ± 3.7	0.9 ± 0.06	3.1 ± 0.1	29.2 ± 4.2
GPR18	0.9 ± 0.4; 0.10 ± 0.10	NAGly [10 µM]	1 µM [84]	5.0 ± 3.9	1.3 ± 0.23	6.3 ± 3.5	21.8 ± 21.8
ADORA2B	0.6 ± 0.2; 8.79 ± 0.88	Bay [5.8 µM]	580 nM [85]	2.7 ± 2.7	0.4 ± 0.40	1.4 ± 1.4	12.5 ± 12.5
P2RY11	0.3 ± 0.1; –	NF 546 [5.0 µM]	500 nM [86]	1.1 ± 1.1	0.28 ± 0.28	0.9 ± 0.9	0.0 ± 0.0

Table 2: Human OPC-expressed $G\alpha_q$-coupled receptors differentially regulated oscillatory
intracellular calcium fluctuations. RNA-seq data from human and mouse [46] was used to identify $G\alpha_q$-
coupled receptors expressed by OPCs. FPKM values shown (mean ± SEM, n = 2 independent human
fetal sample preparations). Ligands activating each receptor were identified to determine the calcium
response characteristics in hOPCs. hOPCs were infected with GCaMP6s-expressing virus and exposed
to various $G\alpha_q$-coupled receptor ligands at doses approximately 10-fold larger than reported EC_{50} values
(or previously effective doses in rodent OPCs*). Calcium responses were recorded for at least 10 minutes
and analyzed as described. The percentage of responding cells (at least 1 peak response ≥ 0.35 fold
increase from baseline on normalized calcium trace following ligand addition), maximum response

amplitude (fold change from baseline), duration of response (minutes) and percentage of responding
cells displaying an oscillatory calcium response (cells with ≥ 2 identified peak responses) were
determined. While Gα$_q$-coupled LPA receptors LPAR1 and LPAR4 were detected in hOPCs, due to the
lack of specific ligands we did not test these receptors. Mean ± SEM of total % responsive cells from n=2-
3 independent human fetal sample culture preparations (Oxo-M, CHPG and ET-1 data: n=3 independent
fetal sample culture preparations; data for all additional ligands: n=2 individual fetal sample culture
preparations).

Abbreviations: **2-FLY**, 2-furoy-Leu-Ile-Gly-Arg-Leu-Orn-NH2; **2-PED**, 2-pyridylethylamide
dihydrochloride; **5-HT**, 5-hydroxytryptamine; **Bay**, Bay 60-6583 2-[[6-Amino-3 5-dicyano-4-[4-
cyclopropylmethoxy phenyl]-2-pyridinyl]thio]-acetamide; **CHPG**, 2-Chloro-5-hydroxyphenylglycine
sodium salt; **ET-1**, Endothelin-1; **MRS 2365**, [[(1R 2R 3S 4R 5S)-4-[6-Amino-2-(methylthio)-9H-purin-9-
yl]-2 3-dihydroxybicyclo[3.1.0]hex-1yl]methyl] diphosphoric acid mono ester trisodium salt; **NAGly**, N-
Arachidonoylglycine; **NF 546**, 4 4'-(Carbonylbis(imino-3 1-phenylene-carbonylimino-3 1-(4-methyl-
phenylene)carbonylimino))-bis(1 3-xylene-alpha alpha'-diphosphonic acid; **Oxo-M**, Oxotremorine
methiodide; **PF9**, 2-(Phenylethynyl)adenosine-5'-triphosphate tetrasodium salt; **TC-G**, 4-Methoxy-N-[[[2-
trifluoromethyl)[1 1'-biphenyl]-4-yl]amino]carbonyl]-3-pyridinecarboxamide; **TRAP-6**, Thrombin receptor
activating peptide 6.

FIGURE LEGENDS

Figure 1. Gα$_q$-receptor activation induces ER-Ca^{2+} responses and SOCE in primary human OPC.

Primary fetal brain-derived PDGFαR$^+$ hOPCs were infected with intracellular Ca^{2+} reporter GCaMP6s

expressing lentivirus. Time-lapse microscopy of GCaMP6s fluorescence following Gα$_q$-receptor agonist

treatment was recorded and analyzed. **A,** pseudo-colored GCaMP6s fluorescence time-lapse images of

oscillatory calcium transients following stimulation with Oxotremorine-M (Oxo-M) [2.5µM] (top) and CHPG

[2.3mM] (bottom). **B**, shows corresponding normalized Ca^{2+} traces. Drug stimulation occurs at 0s (black

arrow in B) and time following drug addition is indicated in each panel. Red and green dots indicated

defined local maxima and minima, respectively. Gold dots indicated defined peaks. Images shown in A

after drug addition correspond to calcium peaks (red) or post-peak minima (blue) and are indicated as

arrows in B. Blue lines below traces indicate the calculated response duration. **C, D,** dose-response

curves of the percentage of hOPCs responding to Gα$_q$ ligand treatment following treatment with either

oxotremorine–M (Oxo-M) or CHPG (mean ± SEM, n=3 human fetal samples, >240 cells per biological

replicate). **E, F,** dose-response curve for the log$_{10}$-transformed overall area under the curve (AUC) for

[Ca^{2+}]$_i$ release per cell (mean ± SEM shown, n > 250 cells per dose, obtained from n=3 human fetal

sample). EC$_{50}$ for AUC was calculated by non-linear regression for each compound (variable slope, four

parameters). **G**, to directly assess SOCE, GCaMP6s-expressing hOPCs were cultured in calcium-free

conditions for 5-10 minutes prior to time-lapse microscopy. Following a 1-2m baseline, cells were treated

with thapsigargin (Tg) or Oxo-M, followed by re-addition of calcium containing solution [1.2mM] to

stimulate SOCE (green dashed lines) in the presence (middle and bottom traces) or absence (top trace)

of SOCE antagonists. Calcium traces showing mean pixel intensity ± SEM (blue shading) of ≥ 44 cells

from a single biological replicate. Timings for addition of Tg, Oxo-M and Ca^{2+}-containing solution are

indicated by horizontal bars above each plot. **H**, quantification of SOCE was performed by analysis of

AUC of calcium influx following calcium re-addition (mean ± SEM; Tg: n=2 human fetal sample

preparations; OxoM: n=3 human fetal sample preparations). * p<0.05, RM one-way ANOVA with Holm-

Sidak's post-hoc test. Scale: 25µm.

Figure 2. STIM1/2 calcium sensors are required for muscarinic-induced SOCE and [Ca^{2+}]$_i$

response in human OPCs. hOPCs were transfected with STIM1 and/or STIM2, or scrambled siRNA,

cultured in mitogen-containing media, and RNA was extracted after 24-48 hours. For calcium imaging

experiments, hOPCs were initially infected with GCaMP6s lentivirus and then transfected with STIM RNAi

or scrambled control siRNA prior to time-lapse microscopy. RT-qPCR analysis of STIM1 (**A)** or STIM2

(**B**) mRNA expression relative to scrambled control transfected hOPCs following STIM1, STIM2 or

STIM1/2 RNAi knockdown (STIM KD) (mean ± SEM, n=7-8 human samples). Control normalized mean

± SEM, one-sample t-test, *p<0.05, **p<0.01, ****p<0.0001. **C**, representative pseudo-colored

fluorescence time-lapse images of peak Oxo-M-induced calcium responses in siRNA transfected hOPCs

cultured in calcium-free conditions. Images represent peak Oxo-M [25µM] induced ER-store depletion

(red text and arrows in **C** and **D** respectively) and SOCE following calcium re-addition (green text and

arrows in **C** and **D** respectively). **D**, calcium traces of Oxo-M induced calcium response following siRNA

transfection showing normalized mean pixel intensity ± SEM (blue shading) of n ≥ 30 cells per condition

from one representative biological replicate. Timings for addition of Oxo-M and Ca^{2+}-containing solution

are indicated by horizontal bars above plots. **E**, SOCE was measured as the total area under the curve

following calcium re-addition normalized to ER-release of control cells in the matched biological replicate.

STIM1/2 KD and pharmacological SOCE inhibition both resulted in a significant reduction in muscarinic-

induced SOCE relative to control hOPCs. Cells were analyzed across 1-2 imaging fields in a single well

per each condition and all responses were averaged per condition for each biological replicate. Data are

presented as mean ± SEM of averaged cell responses per each of four independent experiments, with

>140 total cells quantified per each condition, n=4 independent human fetal sample preparations. RM

one-way ANOVA with Holm-Sidak's post-hoc test, * p<0.05, ** p<0.01). **F–G**, the net effect of

pharmacological SOCE inhibition or STIM KD on Oxo-M-induced increased [Ca^{2+}]$_i$ was determined in

calcium-containing normal growth media by time-lapse microscopy. STIM1 KD or STIM2 KD hOPCs

displayed a reduction in Oxo-M-induced calcium responses as denoted by reduced average AUC (**F**) as

well as a reduction in oscillatory frequency relative to control hOPCs (**G**). Cells were analyzed across two

imaging fields in a single well per each condition and all responses were averaged per condition for each

biological replicate. Data represent log$_{10}$-transformed mean ± SEM of per cell responses from four

independent experiments (n=4 independent human fetal sample preparations), with >100 total cells

quantified per each condition. * p<0.05, ** p<0.01, *** p <0.001, ****p<0.0001, linear model with Tukey's

HSD posttest. Scale: 50 µm.

Figure 3. Calcium sensor STIM2 regulates human OPC differentiation and is required for the anti-

differentiative effect of muscarinic agonists. hOPCs were transfected with STIM1 and/or STIM2

siRNA, or control non-targeting siRNA, and 24 h later PDGF-AA and NT-3 were removed to induce

oligodendrocyte differentiation. **A**, representative images of immature O4$^+$ oligodendrocytes (green) in

the presence or absence of muscarinic agonist, Oxo-M [40µM]. **B**, quantification of the percentage of O4$^+$

oligodendrocyte differentiation among DAPI$^+$ cells (blue). Mean ± SEM shown (n=4 human fetal sample

culture). Differences in O4% were analyzed by RM three-way ANOVA using OxoM, STIM1, and STIM2

as factors and repeated measures across individual patient samples. Both Oxo-M and STIM2 siRNA

treatment resulted in a significant effect on O4$^+$ oligodendrocyte differentiation (OxoM: F (1, 3) = 14.31,

p = 0.032 and STIM2: F (1, 3) = 28.64, p = 0.0128). * p<0.05 and ** p<0.01 indicate pairwise Holm-Sidak

post-hoc tests. **C**, the percentage of dividing OPCs defined as KI-67$^+$/DAPI$^+$ cells following siRNA

transfection was determined in each group (n = 3 human samples). Neither STIM1/2 siRNA nor Oxo-M

treatment significantly influenced OPC proliferation (RM three-way ANOVA, main effects p > 0.05). Scale:

50 μm.

Figure 4. OptoSTIM drives oscillatory calcium responses in hOPCs in the absence of Gα$_q$-coupled

signaling. hOPCs were infected with OptoSTIM and jRCaMP1a lentiviruses to induce SOCE and

observe calcium responses simultaneously. Cells were cultured in a temperature/CO$_2$ controlled chamber

during time-lapse fluorescence microscopy. Blue light (488nm) excitation was used to induce SOCE at

distinct frequencies using 5 minute (3.33 mHz) or 30 minute (0.55 mHz) stimulation intervals respectively,

Blue light exposures were matched to ensure an equivalent amount of blue light stimulation across

conditions (330 and 2000 ms per stimulus, respectively). Changes in intracellular Ca^{2+} concentration

were assessed by jRCaMP1a fluorescence every 20 seconds. **A**, representative integrated calcium

response to OptoSTIM-induced SOCE in the absence of Gα$_q$-receptor activation. Images shown

represent total fluorescence across 2.5 hours of stimulation following a 30-minute baseline period. **B**,

jRCaMP1a traces of mean ± SEM (blue shading) normalized calcium responses to 0.55 mHz and 3.33

mHz frequencies over the course of 2.5 hours following a 30m baseline with no stimulation (n≥50 cells

from a single representative biological replicate). SOCE was robustly induced throughout the imaging

period and directly corresponded with the timing of blue light stimulation (vertical blue lines in middle and

lower traces). **C**, quantification of integrated calcium responses at different stimulation frequencies

measured for the duration of the stimulation period. Mean ± SEM of results from three independent

experiments (n=3 independent human fetal sample preparations) with >50 cells analyzed per condition/

replicate. * p<0.05, RM one-way ANOVA with Holm-Sidak's post-hoc test. Scale: 50 μm.

Figure 5. SOCE is sufficient to block hOPC differentiation and promote hOPC mitosis. Matched

cultures of primary hOPCs were infected with OptoSTIM lentivirus and imaged for 2-3 days in an on-

stage temperature/CO$_2$ controlled chamber. Bright field images were acquired every 5 minutes.

OptoSTIM-mediated SOCE was induced with intermittent 488nm excitation (5m or 30m interval for 330ms

and 2000ms duration, respectively). **A**, hOPC proliferation was quantified by counting mitotic events

(white arrowheads). **B**, the cumulative proportion of cells undergoing mitosis was quantified over 36 h

following the onset of blue light stimulation. **C,** proliferation rates were normalized to the matched

unstimulated condition. High frequency (3.33 mHz) OptoSTIM induction of SOCE resulted in a significant

increase in hOPC mitotic events compared to control unstimulated and low frequency (0.55 mHz)

stimulated hOPCs (n=4 independent human fetal sample culture preparations). **D**, the effect of

OptoSTIM-induced SOCE on hOPC differentiation was determined by O4 immunocytochemistry (red) at

48 hrs post-stimulation. OptoSTIM transduced cells were identified by GFP expression which was uniform

across experimental conditions (**E**). **F**, quantification of O4$^+$ oligodendrocyte differentiation following

OptoSTIM-induced SOCE (% O4$^+$/GFP$^+$). **G**, analysis of O4$^+$ morphological maturation. High frequency

(3.33 mHz) SOCE induced a significant reduction in O4$^+$ cells as well as a significant reduction in complex

branching morphology relative to control hOPCs. * p<0.05, Holm-Sidak's post-hoc test. Mean ± SEM

shown for each panel (n=4 independent human fetal samples). Scale: 25 µm.

Figure 2

Figure 3

Figure 4

Figure 5