1

Prolactin and the shared regulation of parental care and cooperative helping in white-browed sparrow weaver societies

Lindsay A. Walker[1], Linda Tschirren[2], Jennifer E. York[1,3], Peter J. Sharp[4],

Simone L. Meddle[4], Andrew J. Young[1*]

[1] Centre for Ecology and Conservation, University of Exeter Cornwall Campus, Penryn, Cornwall TR10 9EZ, UK

[2] ZHAW School of Life Sciences and Facility Management, Institute of Natural Resource Sciences, Grüental, 8820 Wädenswil

[3] Department of Zoology, University of Cambridge, Downing Street, Cambridge, CB2 3EJ, UK

[4] The Roslin Institute, The Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter Bush, Midlothian EH25 9RG, Scotland, UK

[*] Corresponding author

A.J.Young@exeter.ac.uk

Centre for Ecology and Conservation

University of Exeter, Cornwall Campus

TR10 9EZ

UK

KEY WORDS: Cooperation, mechanism, evolution, hormone, helpers, parental care, alloparental care

ABSTRACT

In many cooperatively breeding species non-breeding individuals help to rear the offspring of breeders. The physiological mechanisms that regulate such cooperative helping behavior are poorly understood, but may have been co-opted, during the evolution of cooperative breeding, from pre-existing mechanisms that regulated parental care. Key among these may be a role for prolactin. Here we investigate whether natural variation in circulating prolactin levels predicts both parental and helper contributions to nestling provisioning in cooperatively breeding white-browed sparrow weavers, *Plocepasser mahali*. In sparrow weaver groups, a single dominant pair monopolize reproduction and non-breeding subordinates help with nestling feeding. We show that: (i) among parents, dominant females feed nestlings at higher rates, make longer provisioning visits, and have higher prolactin levels than dominant males; and (ii) among subordinates, engaged in cooperative helping behavior, those within their natal groups feed nestlings at higher rates and have higher prolactin levels than immigrants. Accordingly, continuous variation in prolactin levels positively predicts nestling-provisioning rates and mean provisioning visit durations when all bird classes are combined. These relationships are principally driven by differences among bird classes in both circulating prolactin levels and provisioning traits. The more limited within-class variation in prolactin and provisioning traits were not evidently correlated, highlighting a likely role for additional mechanisms in the fine-scale regulation of care. Our findings broadly support the hypothesis that parental care and cooperative helping behavior are regulated by a common underlying mechanism and highlight the need for experimentation to now establish the causality of any role for prolactin.

[250 words]

INTRODUCTION

In many cooperatively breeding species, non-breeding helpers assist with the rearing of parents' young,

via cooperative contributions to diverse forms of care (e.g. incubation, babysitting and offspring

provisioning; Solomon and French, 1997; Koenig and Dickinson, 2004; 2016). The majority of research

on such alloparental 'helping behavior' has sought to explain its evolution, by identifying the effects of

helping on recipients and the means by which these yield fitness benefits to helpers (Cockburn, 1998;

Dickinson and Hatchwell, 2004; Koenig and Dickinson, 2016; Capilla-Lasheras et al., 2021). By contrast,

our understanding of the proximate physiological mechanisms that regulate the expression of

cooperative behavior is less advanced (Schoech et al., 2004; Soares et al., 2010; Sanderson et al.,

2014; Dantzer et al., 2017; Vullioud et al., 2021), despite a surge of interest in the origins of consistent

individual differences in both cooperative behavior and endocrine traits (English et al., 2010; Sanderson

et al., 2015; Dantzer et al., 2019; Houslay et al., 2019; 2022). As cooperatively breeding species typically

evolved from species in which parental care was already well developed in both sexes (Cornwallis et

al., 2010; Lukas and Clutton-Brock, 2012), it seems likely that the physiological mechanisms that

regulate cooperative helping behavior among non-breeders were co-opted from the pre-existing

mechanisms that regulated parental care among breeders. Attempts to identify the proximate

mechanisms that regulate helping behavior may therefore be well served by testing candidate

mechanisms already identified for the regulation of parental care in non-cooperative species (Ziegler,

2000; Schoech et al., 2004; Carlson et al., 2006a; 2006b). One such mechanism is the neuroendocrine

pathway involving the anterior pituitary gland hormone prolactin (Buntin, 1996; Sharp et al., 1998;

Ziegler, 2000; Carlson et al., 2006a; 2006b; Angelier et al., 2016).

Numerous studies suggest that prolactin can play a causal role in the expression of parental care,

though its precise role is not clear and seems likely to vary across taxa (Buntin, 1996; Sharp et al., 1998;

Angelier et al., 2016). In birds, prolactin is thought to play a causal role in the onset and maintenance of

parental care, but it is less clear whether variation in circulating prolactin levels is also involved in the

quantitative regulation of contributions to care once caring behavior has begun (Boos et al., 2007;

Angelier et al., 2016; Smiley and Adkins-Regan, 2018). The transition from sexual activity to parenting

is typically associated with an increase in circulating prolactin levels, which peak during the care period

(Buntin, 1996; Sharp et al., 1998; Angelier et al., 2016; e.g. Schoech et al., 1996; Badyaev and

Duckworth, 2005). Evidence that naturally low prolactin levels are commonly associated with breeding

attempt abandonment and/or failure (e.g. Chastel and Lormee, 2002; Chastel et al., 2005), and that

experimental reductions in circulating prolactin levels can disrupt incubation behavior (e.g. Thierry et al.,

2013) and the expression of post-natal nestling care (e.g. Smiley and Adkins-Regan, 2018), suggest

that these elevated prolactin levels are necessary for the onset and/or maintenance of both pre- and

post-natal parental care. Indeed, experimental elevations of circulating prolactin suggest that elevated

prolactin levels can promote the onset of both incubation behavior (e.g. Sockman et al., 2000) and

nestling provisioning behavior (e.g. Badyaev and Duckworth, 2005; Buntin et al., 1991). Positive

associations between continuous variation in circulating prolactin levels and the rates at which parents

provision their offspring (e.g. Duckworth et al., 2003; Ouyang et al., 2011) highlight the possibility that

prolactin levels also regulate the *amount* of care that an actively caring parent provides to its offspring.

However, such positive associations could arise instead via effects of provisioning activity on a bird's

circulating prolactin levels, as parental contact with offspring cues can increase prolactin secretion (Hall,

1987; Sharp et al., 1998). Causal links between prolactin and provisioning rates could therefore exist in

both directions. Indeed, such a feedback loop (in which offspring cues stimulate prolactin secretion that

in turn maintains and/or elevates the expression of parental care) could conceivably both maintain

parental care while offspring survive, and regulate its expression according to offspring vigor and need

(Angelier et al., 2016).

A number of studies of cooperatively breeding species have now begun to investigate the relationships

between prolactin and care-giving behavior, both among parents and non-breeding helpers (Ziegler,

2000; Schoech et al., 2004; Soares et al., 2010). Prolactin levels have been shown to rise in parents

and non-breeding helpers during the transition to incubation and nestling care in at least three species

of cooperatively breeding bird (Schoech et al., 1996; Brown and Vleck, 1998; Khan et al., 2001; see

also Vleck et al., 1991). While few studies have investigated specifically whether variation in circulating

prolactin levels predicts variation in cooperative contributions to helping, studies of at least two

cooperative breeders have yielded compelling evidence in this regard. In Florida scrub jays

(*Aphelocoma coerulescens*), breeders fed offspring at higher rates than non-breeders and showed

higher circulating prolactin levels (Schoech et al., 1996; see also Vleck et al., 1991), and those non-

breeders that helped to feed offspring showed higher prolactin levels than those that did not (Schoech

et al., 1996). Continuous variation in circulating prolactin levels also predicted continuous variation in

feeding contributions, both among all birds combined and specifically among non-breeders (Schoech et

al., 1996). Similarly, in meerkat (*Suricata suricatta*) societies, continuous variation in the prolactin levels

of helpers positively predicted their cooperative contributions to both babysitting and pup-feeding

(Carlson et al., 2006a; 2006b). In the pup-feeding study, prolactin levels only predicted the cooperative

pup-feeding rates of helpers in statistical models that did not allow for an independent positive effect of

circulating cortisol levels on the helper's pup-feeding rates (Carlson et al., 2006a). Experimental work

since highlights that this putative positive effect of cortisol on pup-feeding rates may not have been

causal, however, as glucocorticoid receptor blockade increased rather than decreased pup-feeding

rates among meerkat helpers (Dantzer et al., 2017). Such relationships between circulating prolactin

levels and helping behavior are not always apparent. For example, prolactin levels did not predict the

offspring provisioning rates of helpers in red-cockaded woodpecker (*Picoides borealis*) groups (Khan et

al., 2001), and the pituitary gland prolactin mRNA levels of a cooperatively breeding fish were not

evidently related to care-giving behavior (Bender et al., 2008); though the relevant sample sizes in both

studies were modest. Attempts to identify correlations between natural continuous variation in prolactin

levels and care-giving behavior are expected to be complicated, however, by the existence of other

neuroendocrine pathways that may also modulate the expression of care-giving (e.g. glucocorticoids

may play a role in its state-dependent modulation; Sanderson et al., 2014; Dantzer et al., 2017; see

also Angelier and Chastel, 2009; Angelier et al., 2016), potentially independent of circulating prolactin

levels (Schoech et al., 1998; Ziegler, 2000; Young et al., 2005; Carlson et al., 2006a). Nevertheless, the

promising findings to date highlight the need for further studies to investigate whether natural circulating

levels of prolactin predict variation in contributions to both parental care and helping behavior in

cooperatively breeding species, and, ultimately, the use of experimental manipulations to test the

causality and nature of any hormone-behavior relationships detected (Sockman et al., 2000; Carlson et

al., 2003; Badyaev and Duckworth, 2005; Smiley and Adkins-Regan, 2018).

Here we investigate whether natural variation in circulating prolactin levels positively predicts the

nestling provisioning behavior of both parents and non-breeding helpers in a wild cooperatively breeding

bird, the white-browed sparrow-weaver (*Plocepasser mahali*). White-browed sparrow weavers are rain-

dependent breeders that live in year-round territorial groups throughout the semi-arid regions of sub-

Saharan Africa (Lewis, 1982; Wood et al., 2021). Within each social group, a single dominant male and

female completely monopolize within-group reproduction and up to 10 non-breeding subordinates of

both sexes help to feed their nestlings (Harrison et al., 2013a; 2013b; Capilla-Lasheras et al., 2021;

2023). Subordinates are typically offspring from previous broods that have delayed dispersal from their

natal group (and so are helping to rear their parents' young), but subordinate immigrants of both sexes

do also occur (Harrison et al., 2013a; 2013b; Harrison et al., 2014). Subordinates contribute to several

cooperative activities year-round, including territorial defense, roost construction and anti-predator

vigilance (Lewis, 1982; Walker et al., 2016; York et al., 2019), and during breeding periods they

contribute substantially to nestling provisioning (Cram et al., 2015a; Capilla-Lasheras et al., 2021; 2023).

Helping behavior by subordinates in this species appears to have positive effects on both the helped

offspring being fed and their parents. First, helping behavior has a causal positive effect on the overall

rate at which nestlings are fed (Capilla-Lasheras et al., 2021), and, accordingly, helper numbers

positively predict nestling survival to fledging during dry periods, reducing the environmentally-induced

variance in the reproductive success of the dominant pair (Capilla-Lasheras et al., 2021). Evidence

suggestive of positive helper effects on offspring telomere lengths suggest that helpers may also

improve the downstream performance of surviving young (Wood, 2017; Wood and Young, 2019).

Second, helping behavior appears to lighten the post-natal provisioning workload of the dominant female

(Capilla-Lasheras et al., in press), which may explain why mothers with more help increase their pre-

natal investment in the egg (Capilla-Lasheras et al., in press) and show higher overwinter survival

(O'Callaghan, 2021). While the neuroendocrine correlates of white-browed sparrow weaver

reproduction, aggression and song production have been investigated (e.g. Wingfield and Lewis, 1993;

Voigt et al., 2007; York, 2012; York et al., 2016), the regulation of parental and helper contributions to

offspring provisioning remains unexplored.

We test three predictions of the hypothesis that prolactin promotes the expression of both parental care

(among dominants) and cooperative helping behavior (among non-breeding subordinates) in

cooperatively breeding societies. First, with regard to parental care, we predict that differences between

the nestling provisioning rates of dominant females and dominant males will be mirrored by parallel

differences in their mean circulating prolactin levels. Dominant females are expected to provision

nestlings at higher rates (and to have higher prolactin levels) than dominant males, as the 12-18%

incidence of extra-group paternity in this population leaves dominant females more closely related than

dominant males, on average, to the offspring that they rear (Harrison et al., 2013a). Second, with regard

to alloparental helping behavior, we predict that differences between the nestling provisioning rates of

subordinates still residing within their natal group (hereafter 'natal subordinates') and immigrant

subordinates, will also be mirrored by parallel differences in their mean circulating prolactin levels. Natal

subordinates are expected to provision nestlings at higher rates (and to have higher prolactin levels)

than immigrant subordinates, as while the former are typically rearing future generations of siblings born

to their parents, the latter are typically unrelated to the nestlings in their group (Harrison et al., 2013a).

Finally, we predict that continuous variation in circulating prolactin levels will positively predict

continuous variation in the nestling provisioning rates of birds, and that this relationship will be apparent

(i) at the population level (when all four of the bird classes above are combined), and (ii) within bird

classes, having factored out the among-class differences in prolactin levels and provisioning rates.

METHODS

General field methods

Data were collected in the context of a long-term research project that monitors ~40 cooperative groups

of white-browed sparrow weavers at Tswalu Kalahari Reserve, South Africa (27°160S, 22°250E), and

at a similar time in two separate breeding seasons (January to February 2013, and January to March

2014). White-browed sparrow weavers in this population may breed at any time from September through

to May (the Southern summer), depending on the timing of unpredictable summer rainfall (Wood et al.,

2021; Capilla-Lasheras et al., 2021). Each bird within our study population is fitted with a metal ring and

three color rings, providing a unique ring combination for identification in the field (SAFRING license

1444). From around six months of age, males and females of the focal subspecies (*Plocepasser mahali*

mahali) can be distinguished by their bill color; males have a dark brown bill while females have a paler

grey-to-pink bill (Leitner et al., 2009). Dominance status and social group compositions were determined

via regular (at least twice weekly) group visits. Social dominance was assigned based on the monitoring

of key dominance-related behaviors: the dominant pair routinely displace other group members and

produce synchronized duet song, the dominant female is the sole incubator, and the dominant male

consistently produces dawn song during breeding periods (Harrison et al., 2013a; Cram et al., 2015b;

York et al., 2016). The dispersal status (natal or immigrant) of subordinate birds was determined via the

continuous monitoring of the study population since 2007. Based on this information, four classes of

birds were assigned: Dominant Females; Dominant Males; Natal Subordinates and Immigrant

Subordinates. Social group size was defined as the number of adult (> 1 year of age) birds consistently

seen foraging and roosting together at the time of the focal breeding attempt. The breeding status of

each group was determined by monitoring the contents of all woven nest structures within each group's

territory, at least every other day throughout the two study periods. When one or more eggs were newly

detected, the active nest was visited daily in the afternoon until no new eggs were detected (the birds

lay one egg per day in the morning, and typically lay clutches of 2 eggs (range 1-4); Harrison et al.,

2013a). To determine hatch dates, daily monitoring of the active nest resumed 14 days after the

detection of the first egg (as incubation lasts 14-19 days; Harrison et al., 2013a). This method yielded

accurate information on the day on which the first nestling in each clutch hatched, which was termed

'Day 1' of the nestling provisioning period for the focal breeding attempt. All protocols were approved by

the Ethics Committees of the Universities of Exeter and Pretoria and complied with regulations stipulated

in The Association for the Study of Animal Behaviour (ASAB) Guidelines for Use of Animals in Research.

Monitoring provisioning behavior

To identify individuals during the recording of nestling provisioning events, group members were

captured from their roost chambers (details below) during the incubation period and marked on the vent

with a unique dye-mark. The dominant female was left unmarked to minimize disturbance during

incubation, but could still be distinguished from other group members by being the only unmarked bird

within her group (only resident group members provision offspring). To record provisioning events, a

Panasonic SDR-S50 camcorder attached to a tripod (approximately 0.5 meters in height) was placed

on the ground beneath the entrance to the active nest two days before recording commenced (to allow

the birds to habituate to it). On the days of provisioning monitoring, the recordings (approximately 3

hours in the duration) were started between 06:15 and 07:54, with this start time being adjusted through

the season to maintain an approximately constant time offset from sunrise. Provisioning videos were

collected in this way for all focal breeding attempts (n = 37 broods across 30 social groups) on two

mornings between Days 6 and 9 inclusive of the nestling provisioning period (typically for the two

consecutive mornings of Days 7 and 8; nestlings fledged from day 20). This approach yielded a mean

total duration of provisioning video of 6.08 hours (range 4.05 – 8.12 hours) per breeding attempt.

Video recordings were transcribed using VLC Media Player version 2.2, with the observer recording, for

each provisioning visit, the identity of the bird visiting the nest (determined via their distinct dye mark

and bill color, which reveals their sex) and the duration of time that they spent within the nest (the time

elapsed between passing in and out of the enclosed nest structure; hereafter 'Provisioning visit

duration'). Prior work on this study population using within-nest cameras has shown that all nest visits

during the nestling age window studied here entail the delivery of a single prey item to the brood, unless

the visiting bird is carrying a feather or grass in which case no food is delivered (Walker, 2016). We

therefore excluded such feather- or grass-carrying nest visits from our provisioning visit records. From

the transcribed data for each focal brood we then calculated two provisioning trait values for each adult

group member: (i) 'Provisioning rate' (feeds / hr) was calculated as the total number of provisioning visits

that the bird conducted over the two monitored mornings divided by the total duration of video collected

over those two mornings, and (ii) 'Mean Visit Duration' (minutes) was calculated as the mean duration

of all provisioning visits conducted by the focal bird over the two monitored mornings.

Bird capture and blood sampling

To obtain a matched blood sample for prolactin measurement, we attempted to capture and blood

sample all adult (> 1 year old at the time of sampling) birds within the monitored brood's social group on

the evening of the second day of provisioning behavior recording. Birds were captured individually at

night from the woven roost chambers within their group's territory (Cram et al., 2015a) by flushing

individuals into a custom-made capture bag. All captures, dye-marking and blood sampling were

conducted by a single investigator (LW). Birds were then immediately returned to a roost chamber within

their territory to pass the remainder of the night.

Upon capture, a blood sample (*c.* 140 µL) was taken from the brachial vein of the bird using a 26g

needle and heparinized capillary tubes. Captures occurred soon after dusk, once the birds were roosting

in their woven chambers. Time of capture was recorded (to allow us to fit the time lag from sunset to

capture as a covariate predictor in our prolactin analyses, in case of diel variation and/or effects of the

time elapsed since roosting on a bird's prolactin levels), along with the time lag between capture and

the completion of blood sampling (mean ± standard deviation [S.D.] = 3.12 ± 0.73 minutes [range 1.65

– 4.78 minutes]; to allow us to control for potential effects of capture stress on prolactin levels in our

statistical models). Blood samples were immediately centrifuged in the field (12,000 g for 3 minutes;

Haematospin 1400; Hawksley Medical and Laboratory Equipment, Lancing, UK) and the plasma was

drawn off and stored in a cryovial on ice until it could be transferred to liquid nitrogen on return from the

field (mean ± S.D. time lag from sample collection to storage on liquid nitrogen = 148 ± 63 min). At the

end of the field season, samples were transferred to the UK on dry ice and then stored at -80 degrees

Celsius until analysis for prolactin.

Prolactin Radioimmunoassay

The prolactin assay was carried out in July 2014 at the Roslin Institute (University of Edinburgh, Easter

Bush, Midlothian, Scotland, UK). Plasma prolactin levels were measured using a highly specific

heterologous micro-radioimmunoassay of donkey anti-rabbit serum to European starling (*Sturnus*

vulgaris) prolactin (Sharp antibody code 44/2). Prolactin was radiolabeled with iodine[125] using

chloramine-T. 168 (out of a total of 208) samples were assayed in duplicate, and the remaining 40

samples were assayed as singletons (not all samples assayed were for use in this study). All samples

were measured in a single assay, in which the intra-assay coefficient of variation for the duplicate

samples was 3.31%.

Statistical methods

The above methods yielded a final data set of matched provisioning trait data (estimated from the focal

bird's average performance over two mornings of provisioning recordings; see above) and circulating

prolactin levels (when sampled on the evening of the second day of provisioning monitoring) for 70

different adult birds, each sampled once (for all traits), while feeding a total of 37 broods across 30 social

groups. While some analyses utilized the whole data set (i.e. n = 70 adults birds each sampled once for

all traits), others used subsets of it (e.g. when focusing only on dominants engaged in parental care or

subordinates engaged in cooperative helping behavior), and so the sample sizes for each analysis are

reported within the relevant results section and model output table. As mean provisioning visit duration

data were only available for birds that had a non-zero provisioning rate, the sample sizes for mean visit

duration analyses were sometimes smaller than those for provisioning rate (see results).

All statistical models and visualizations were carried out in R (version 4.1.0; R Core Team). Mixed effects

modelling was conducted using the R package 'lme4' (Bates et al., 2015) We conducted our statistical

modelling using a full model approach, in which we (i) specified a full model containing both the primary

fixed effect predictors of interest and covariates whose potential effects we also wished to control for,

and then (ii) tested the effects of these predictors in that context, without any model selection or

simplification. The statistical significance of a given predictor was assessed by using a likelihood ratio

test to determine the significance of the change in the explanatory power of the full model when the

focal predictor was dropped from the full model. This conservative approach ensures that the

significance of all predictors is assessed while controlling for the potential effects of the other predictors

specified in the full model, regardless of whether those other predictors themselves have significant

effects. The specific modelling exercises conducted for each results section are described below.

1. Are differences in the provisioning behavior of dominant females and males (engaged in

parental care) mirrored by differences in their circulating prolactin levels?

We used two separate mixed effects models with Gaussian error structure to model the causes of

variation in (i) the provisioning rates and (ii) the mean provisioning visit durations of dominant birds (i.e.

engaged in parental care). The two modeling exercises began with an identical full model structure. In

addition to the primary predictor of interest, 'parent class' (dominant female or dominant male), we fitted

the following terms as fixed effect predictors: brood size (the brood size being fed), adult group size (the

number of adult group members during the focal nestling provisioning period) and year (a two-level

factor capturing the year in which sampling occurred; 2013 or 2014). We fitted both social group ID and

brood ID (the identity of the brood being fed) as random effects, retaining them in the model structure

regardless of the degree of variance that they explained.

We then used a third mixed effects model with Gaussian error structure to model the causes of variation

in the circulating prolactin levels of these same dominant birds, starting with a full model structure

containing the same fixed and random effect predictors as the provisioning trait models just described,

but with the addition of two further fixed effects to account for potential methodological effects on

prolactin concentrations: (i) the time lag from sunset to the bird's capture for blood sampling (to allow

for the possibility of diel variation in prolactin levels and/or changes in prolactin levels once the birds

entered their roosts) and (ii) the time lag from capture to the completion of blood sampling (to allow for

possible effects of capture stress on circulating prolactin levels).

2. Are differences in the provisioning behavior of natal and immigrant subordinates (engaged in

alloparental helping behavior) mirrored by differences in their circulating prolactin levels?

We then used two mixed effects models with Gaussian error structure to model the causes of variation

in the provisioning rates and circulating prolactin levels of subordinate birds engaged in alloparental

helping behavior. We fitted the same set of fixed and random effect predictors to these models as were

fitted to the corresponding provisioning rate and prolactin level models conducted for dominant birds

(see above) with two exceptions: (i) in place of 'parent class' we fitted 'helper class', reflecting whether

the focal bird was a natal subordinate or an immigrant subordinate, and (ii) here only brood ID was fitted

as a random effect (social group ID was not, as all subordinate birds from any given social group were

sampled while feeding the same single brood, leaving brood ID and social group ID with identical

structure). We did not model the causes of variation in the mean provisioning visit durations of

subordinates as too few immigrant subordinates actually provisioned the focal broods, leaving us with

an insufficient sample size of measures of the provisioning visit durations of this bird class.

3. Does continuous variation in prolactin levels predict variation in provisioning behaviour?

To investigate whether natural variation in prolactin levels predicted continuous variation in the birds'

nestling provisioning rates and mean provisioning visit durations at the population level (i.e. when all

bird classes were combined) we conducted two mixed effect models (one for each provisioning trait

response term), with circulating prolactin concentration as the sole fixed effect predictor (as we have

not hypothesized specific mechanisms by which other variables might impact provisioning traits

independent of prolactin levels) and social group ID and brood ID as random effects. Mean provisioning

visit duration was logarithm transformed for analysis, to normalize model residuals.

Inspection of the patterns of the mean prolactin levels and provisioning trait values of the four different

bird classes (i.e. dominant females, dominant males, natal subordinates and immigrant subordinates)

suggested that any such continuous relationship between prolactin levels and provisioning trait values

at the population level across all bird classes could be driven principally by the variation in these traits

among the bird classes (Figures 3a & 3c). In order to then investigate whether the more limited variation

in circulating prolactin levels *within* bird classes predicted the *within*-class variation in provisioning trait

values, we first mean-centered each birds' prolactin level and provisioning trait values (the log

transformed values in the case of mean provisioning visit duration) around the mean value of the focal

trait for birds of their class (by subtracting from it the mean value of the focal trait for their bird class).

We then conducted two mixed effects models (one for each mean-centered provisioning trait response

term), with mean-centered circulating prolactin concentration as the sole fixed effect predictor and social

group ID and clutch ID as random effects.

RESULTS

1. Are differences in the provisioning behavior of dominant females and males (engaged in
parental care) mirrored by differences in their circulating prolactin levels?

Analyzing the provisioning behavior of dominant birds engaged in parental care (n = 46 dominants, 20
females and 26 males, feeding 31 broods at 28 social groups) revealed that dominant females feed
offspring at significantly higher rates than dominant males (Figure 1a; parent class effect, for dominant
male relative to dominant female, ± S.E. = +5.11 ± 0.57; P < 0.001; Table 1) and also show significantly
longer mean provisioning visit durations (Figure 1b; parent class effect, for dominant male relative to
dominant female, ± S.E. = +1.29 ± 0.31; P < 0.001; Table 2). There was also evidence that dominant
birds fed larger broods at significantly higher rates (brood size effect ± S.E. = +1.08 ± 0.50; P = 0.035;
Table 1). There was no compelling evidence that the provisioning rates or mean visit durations of
dominant birds were associated with either the year of study or adult group size (Tables 1 & 2).

Analyzing the circulating prolactin levels of dominant birds during the provisioning periods analyzed
above (again, n = 46 dominants, 20 females and 26 males, feeding 31 broods at 28 social groups)
revealed evidence that dominant females also have significantly higher circulating prolactin levels than
dominant males (Figure 1c; parent class effect, for dominant male relative to dominant female, ± S.E. =
+1.69 ± 0.26; P < 0.001; Table 3). There was no compelling evidence that the prolactin levels of
dominants were associated with group size, brood size, the time lag from sunset to capture or the time
lag from capture to blood sampling (Table 3). Note that our full model approach ensured that any effects
of these latter predictors were controlled (regardless of their significance) when assessing the effects of
other terms.

Figure 1. The (a) provisioning rates, (b) mean provisioning visit durations, and (c) circulating prolactin

concentrations of dominant females (Dom Fem) and dominant males (Dom Male) engaged in parental

care. Squares present the predicted means (± S.E.) from the full model for the relevant trait (Tables 1,

2 and 3 respectively), while controlling for the effects of all other variables in the full model. The predicted

means were calculated with all continuous predictors set to their mean values and for the 2013 category

of the Season factor. The points show the raw data points.

Table 1: Modelling the effects of **Parental Sex on Nestling Provisioning Rate during Parental care**.

N = 46 dominant birds, 20 female and 26 male, feeding 31 broods at 28 social groups. The 'Parent class'

effect size is for the dominant female relative to the dominant male. The 'Year' effect size is for the 2014

field season relative to the 2013 field season. Significant effect sizes (on the basis of likelihood ratio

tests when comparing the full model to the full model without the focal term) are highlighted in bold. SE

= Standard Error; LRT = Likelihood ratio test and associated P value.

	Estimate	SE	t	LRT	P
(Intercept)	-0.461	1.252	-0.368		
Parent class (female > male)	**5.106**	**0.568**	**8.984**	**46.611**	**<0.001**
Year	1.027	0.638	1.608	2.516	0.113
Group size	0.050	0.276	0.181	0.033	0.856
Brood size	**1.084**	**0.501**	**2.164**	**4.461**	**0.035**

Table 2: Modelling the effects of **Parental Sex on Mean Nestling Provisioning Visit Duration during**
Parental care. N = 44 dominant birds, 20 female and 24 male, feeding 31 broods at 28 social groups;
the sample size for this analysis was slightly lower than that for the Provisioning Rate analysis (Table
1) as 2 dominant males did not provision their brood, leaving us without a measure of their mean visit
duration. The 'Parent class' effect size is for the dominant female relative to the dominant male. The
'Year' effect size is for the 2014 field season relative to the 2013 field season. Significant effect sizes
(on the basis of likelihood ratio tests when comparing the full model to the full model without the focal
term) are highlighted in bold. SE = Standard Error; LRT = Likelihood ratio test and associated P value.

	Estimate	SE	t	LRT	P
Intercept	0.960	0.720	1.333		
Parent class (female > male)	**1.292**	**0.314**	**4.119**	**14.351**	**<0.001**
Year	-0.091	0.348	-0.262	0.069	0.793
Group size	0.032	0.153	0.210	0.044	0.834
Brood size	-0.253	0.285	-0.887	0.780	0.377

Table 3: Modelling the effects of **Parental Sex on Circulating Prolactin level during Parental care**.
N = 46 dominant birds, 20 female and 26 male, feeding 31 broods at 28 social groups (identical to the
nestling provisioning rate analysis in Table 1, as all birds were sampled for both traits in the same
contexts). The 'Parent class' effect size is for the dominant female relative to the dominant male. The
'Year' effect size is for the 2014 field season relative to the 2013 field season. Significant effect sizes
(on the basis of likelihood ratio tests when comparing the full model to the full model without the focal
term) are highlighted in bold. SE = Standard Error; LRT = Likelihood ratio test and associated P value.
Sunset to capture lag = time elapsed between sunset and capture (to account for potential circadian
variation in prolactin levels). Capture to bleed lag = time elapsed between first contact with the roost
chamber and blood sample (to allow for the possibility of a prolactin stress response).

	Estimate	SE	t	LRT	P
(Intercept)	0.731	1.199	0.610		
Parent class (female > male)	**1.691**	**0.261**	**6.487**	**23.196**	**<0.001**
Year	0.328	0.446	0.735	0.378	0.539
Group size	0.271	0.180	1.509	2.217	0.137
Brood size	0.298	0.320	0.931	0.783	0.376
Sunset to capture lag	0.000	0.003	-0.002	0.000	0.998
Capture to bleed lag	-0.001	0.004	-0.295	0.087	0.768

2. Are differences in the provisioning behavior of natal and immigrant subordinates (engaged in alloparental helping behavior) mirrored by differences in their circulating prolactin levels?

Analyzing the provisioning behavior of subordinate birds engaged in cooperative helping behavior (n = 24 subordinates, 17 natal and 7 immigrant, feeding 16 broods at 16 social groups) revealed evidence that subordinates within their natal groups feed offspring at higher rates than immigrant subordinates (Figure 2a; helper class effect, for natal subordinates relative to immigrant subordinates, ± S.E. = +1.61 ± 0.33; P < 0.001; Table 4). There was also evidence that helpers fed larger broods at significantly higher rates (brood size effect ± S.E. = +1.06 ± 0.35; P = 0.007; Table 4). There was no compelling evidence that helper provisioning rates were associated with either group size or the year of study (Table 4). No analysis of the provisioning visit durations of subordinates was conducted as an insufficient number of subordinate immigrants ever provisioned the broods (see Figure 2a).

Analyzing the circulating prolactin levels of subordinate birds during the provisioning periods analyzed above (again, n = 24 subordinates, 17 natal and 7 immigrant, feeding 16 broods at 16 social groups) revealed that natal subordinates also have significantly higher prolactin levels than immigrant subordinates (Figure 2b; helper class effect, for natal subordinates relative to immigrant subordinates, ± S.E. = +0.86 ± 0.28; P = 0.005; Table 5). There was also evidence that subordinate prolactin levels were significantly higher in the second year of study (year effect, for 2014 relative to 2013, ± S.E. = +0.85 ± 0.37; P = 0.029; Table 5) and in smaller groups (Group size effect ± S.E. = -0.37 ± 0.13; P = 0.007; Table 5). There was no compelling evidence that subordinate prolactin levels were associated with brood size or the time lags from sunset to capture and from capture to blood sampling (Table 5). Note that our full model approach ensured that any effects of these latter predictors were controlled (regardless of their significance) when assessing the effects of other terms.

Figure 2. The (a) provisioning rates, and (b) circulating prolactin levels of natal subordinates (Sub Natal)

and immigrant subordinates (Sub Imm) engaged in cooperative helping behavior, feeding the broods of

the dominant male and female. Y axis scales match those in Figure 1 to facilitate comparison. Squares

present the predicted means (± S.E.) from the full model for the relevant trait (Tables 4 and 5

respectively), while controlling for the effects of all other variables in the full model. The predicted means

were calculated with all continuous predictors set to their mean values and for the 2013 category of the

Season factor. The points show the raw data points. No analysis of provisioning visit durations was

conducted as an insufficient number of subordinate immigrants ever provisioned the broods (see panel

2a).

Table 4: Modelling the effects of **Helper Class on Nestling Provisioning Rate during Alloparental**
Helping Behaviour. N = 24 subordinate birds, 17 natal and 7 immigrant, feeding 16 broods at 16 social
groups. The 'Helper class' effect size is for natal subordinates (those residing within their natal groups)
relative to immigrant subordinates (those that have dispersed to another group). The 'year' effect size
is for the 2014 field season relative to the 2013 field season. Significant effect sizes (on the basis of
likelihood ratio tests when comparing the full model to the full model without the focal term) are
highlighted in bold. SE = Standard Error; LRT = Likelihood ratio test and associated P value.
469

	Estimate	SE	t	LRT	P
Intercept	-0.297	1.097	-0.271		
Helper class **(natal > immigrant)**	**1.610**	**0.334**	**4.826**	**15.858**	**<0.001**
Year	-0.810	0.435	-1.861	3.173	0.075
Group size	-0.274	0.226	-1.212	1.415	0.234
Brood size	**1.057**	**0.354**	**2.986**	**7.166**	**0.007**

Table 5: Modelling the effects of **Helper Class on Circulating Prolactin level during Alloparental**
Helping Behaviour. N = 24 subordinate birds, 17 natal and 7 immigrant, feeding 16 broods at 16 social
groups (identical to the nestling provisioning rate analysis in Table 4, as all birds were sampled for both
traits in the same contexts). The 'Helper class' effect size is for natal subordinates (those residing within
their natal groups) relative to immigrant subordinates (those that have dispersed to another group). The
'year' effect size is for the 2014 field season relative to the 2013 field season. Significant effect sizes
(on the basis of likelihood ratio tests when comparing the full model to the full model without the focal
term) are highlighted in bold. SE = Standard Error; LRT = Likelihood ratio test and associated P value.
Sunset to capture lag = time elapsed between sunset and capture (to account for potential circadian
variation in prolactin levels). Capture to bleed lag = time elapsed between first contact with the roost
chamber and blood sample (to allow for the possibility of a prolactin stress response).
482

	Estimate	SE	t	LRT	P
Intercept	1.255	0.910	1.378		
Helper class **(natal > immigrant)**	**0.860**	**0.281**	**3.054**	**7.881**	**0.005**
Year **(2014 > 2013)**	**0.848**	**0.370**	**2.294**	**4.757**	**0.029**
Group size	**-0.365**	**0.126**	**-2.901**	**7.152**	**0.007**
Brood size	-0.175	0.227	-0.770	0.586	0.444
Sunset to capture lag	0.002	0.003	0.709	0.498	0.480
Capture to bleed lag	0.004	0.003	1.561	2.320	0.128

3. Does continuous variation in prolactin levels predict variation in provisioning behaviour?

Our analysis at the population level, including birds of all classes, revealed that a bird's circulating

prolactin level strongly and significantly positively predicts both (i) its provisioning rate (Figure 3a; effect

size ± S.E. = 1.20 ± 0.23; P < 0.001; n = 70 birds feeding 37 broods at 30 social groups) and (ii) its mean

provisioning visit duration (Figure 3c; effect size ± S.E. = 0.32 ± 0.076; P < 0.001; n = 59 birds feeding

36 broods at 30 social groups). Plotting out the mean prolactin levels and provisioning trait values of the

different bird classes (Figure 3a & 3c), reveals that both of these population-level relationships between

prolactin and provisioning traits are driven in large part by the among-bird-class differences in prolactin

levels being mirrored by parallel among-bird-class differences in mean provisioning rate (Figure 3a) and

mean provisioning visit duration (Figure 3c). Indeed, after mean-centering each bird's prolactin level and

provisioning trait values around the focal trait's mean value for their bird class, we found no evidence

that within-bird-class variation in prolactin levels predicted within-bird-class variation in either

provisioning rate (Figure 3b; effect size ± S.E. = -0.15 ± 0.21; P = 0.50) or mean provisioning visit

duration (Figure 3d; effect size ± S.E. = 0.049 ± 0.081; P = 0.55).

Figure 3. At the population level, considering all bird classes together, natural variation in circulating prolactin levels predicts variation in both (**a**) provisioning rate and (**c**) mean provisioning visit duration. These relationships are driven principally by differences among the mean trait values of the different focal bird classes (presented ± S.E. by the squares and diamonds within panels **a** and **c**; see legend within panel **a**). Follow-up analyses revealed no evidence that variation in prolactin levels *within* these bird classes predicted within-bird-class variation in either (**b**) provisioning rate or (**d**) mean provisioning visit duration. In panels **a** and **c** the line and shaded ribbon present the predicted mean relationship and its standard error, while the shaded circular points within all panels present the raw data points.

DISCUSSION

This study investigated the hypothesis that prolactin plays a role in the regulation of nestling

provisioning, both among dominant birds (engaged in parental care) and non-breeding subordinate birds

(engaged in cooperative helping behavior), in cooperatively breeding white-browed sparrow weaver

societies. Among dominants engaged in parental care, we found that the dominant female (the mother)

fed offspring at higher rates, made longer provisioning visits and had higher circulating prolactin levels

than the dominant male (typically the father). Among subordinates, we found that natal subordinates

helped to feed offspring at higher rates and had higher circulating prolactin levels than immigrant

subordinates. Indeed, when all bird classes were combined, we found that continuous variation in the

circulating prolactin levels of the birds predicted continuous variation in their provisioning rates and

mean provisioning visit durations. These patterns appear to be driven principally by correlated

differences among the four different bird classes in their prolactin levels and provisioning traits. We

found no evidence that the more limited variation in circulating prolactin levels within the different bird

classes predicted the more limited within-class variation in their provisioning traits. Together, these

findings are broadly consistent with the hypothesis that parental care and cooperative helping behavior

are regulated by a common underlying mechanism and that prolactin plays a role in that pathway, and

highlight the need for experimental studies to now probe the causality and nature of any role for prolactin.

Below, we discuss potential explanations for these findings, the different roles that prolactin could

conceivably play in the regulation of parenting and cooperative helping in this species, and the wider

implications of our findings for mechanistic and evolutionary research on cooperative behavior.

While our findings are consistent with the hypothesis that parental care and helping behavior are

regulated by a common mechanism in which prolactin plays a role, the lack of a relationship between

within-class variation in prolactin levels and provisioning traits, coupled with the correlative nature of our

findings, leave it important to consider the range of possible roles that prolactin could play in the

regulation of provisioning behavior in this species. At least three main possibilities exist, which will

require careful experimentation to tease apart. First, it is possible that circulating prolactin is one key

regulator of continuous variation in individual contributions to offspring provisioning, both among parents

and helpers. While most of our findings are consistent with this hypothesis, the absence of evident

relationships between within-class variation in prolactin levels and provisioning traits complicates this

view. However, the lack of evident *within*-class relationships could be attributable simply to a major

source of variation in both traits (*among*-class variation) having been factored out at this stage of the

analysis, leaving these within-class analyses seeking relationships between the more limited *within*-

class variation in both traits, which could be readily obscured by a number of mechanisms. First,

difficulties with the synchronous and accurate assessment of both prolactin levels and provisioning rates

could have yielded noise in the data set that precluded the detection of these more subtle prolactin-

provisioning relationships. While we sampled birds for prolactin on the evening following the morning

provisioning-monitoring session (a time lag comparable to, or shorter than, those of similar studies; e.g.

Duckworth et al., 2003; Ouyang et al., 2011), individuals may have differed in the way that their prolactin

levels changed during the day, leaving their evening prolactin levels only a modest proxy for those while

provisioning. The focal birds also varied in the timing of blood sampling, and while our analyses did not

detect any overall effects on prolactin levels of the time lag from capture to sampling, any individual

variation in the prolactin stress response (if this species shows one; Krause et al., 2015) could have

further decoupled the assessed prolactin levels from those during provisioning. Second, even if prolactin

levels were a key regulator of continuous variation in provisioning rates, alternative mechanisms are

also expected to impact provisioning rates potentially independent of circulating prolactin levels, leaving

the relationship between natural variation in prolactin levels and provisioning behavior potentially weak

in the first place (Schoech et al., 1998; Angelier et al., 2016). Key among these could be (i) variation in

other components of a prolactin-mediated pathway (such as inter-individual and temporal variation in

the density of prolactin receptors; Zhou et al., 1996; Ohkubo et al., 1998; Angelier et al., 2016), as well

as (ii) mechanisms that may impact provisioning behavior via prolactin-independent pathways (e.g. the

effects of circulating testosterone; Schoech et al., 1998; Angelier et al., 2016). Third, even if prolactin

levels alone determined provisioning 'motivation', the extent to which variation in provisioning motivation

578 was reflected in provisioning rates would depend upon the prey capture skills of the focal bird and the

579 environmental availability of prey. Indeed, all points considered, it is arguably mechanistically naïve to

580 expect particularly fine-grained associations between the levels of a single hormone and behavior to be

581 evident in natural populations even where a causal link exists between the two. To now robustly test the

582 hypothesis that prolactin regulates continuous variation in the magnitude of both parental and helper

583 contributions to offspring provisioning, there is a need to experimentally elevate the circulating prolactin

584 levels of actively provisioning birds whose natural prolactin levels are not at the upper end of the

585 physiological range (dominant males and natal subordinates may serve this purpose well; Figure 3a).

586 This manipulation would allow one to test the key prediction that an increase in the prolactin levels of

587 an actively provisioning bird will cause it to increase its provisioning rate; a prediction that to our

588 knowledge has yet to be tested in either a parenting or helping context (the few experimental elevations

589 of endogenous prolactin secretion in a provisioning context to date have focussed on the establishment

590 of provisioning in non-provisioning birds rather than its quantitative variation within actively provisioning

591 birds; e.g. Badyaev and Duckworth, 2005).

593 A second potential explanation for the balance of our findings is that prolactin could instead play a causal

594 role in the onset and maintenance of provisioning behavior among parents and helpers, without playing

595 a role in the quantitative regulation of contributions to provisioning among actively provisioning birds

596 (Angelier et al., 2016). For example, a threshold level of prolactin may be required for the onset and/or

597 maintenance of provisioning behavior (Angelier et al., 2006; Boos et al., 2007). Under this scenario, the

598 higher prolactin levels of natal subordinates and dominant birds, relative to immigrant subordinates,

599 could be causally responsible for the former bird classes engaging in provisioning while the latter

600 typically does not. This could be the case without prolactin playing any causal role in regulating

601 continuous variation in the provisioning rates of actively provisioning birds; a scenario that could account

602 for the lack of within-class correlations between prolactin levels and provisioning behavior. The elevated

603 prolactin levels of dominant females (relative to dominant males and natal subordinates) could

conceivably be a downstream consequence of either a role for prolactin in incubation (Buntin, 1996;

Sharp et al., 1998; Khan et al., 2001; as dominant females are the sole incubator in this species) and/or

their differential exposure to offspring cues during the nestling period (which can increase prolactin

secretion; Hall, 1987; Sharp et al., 1998), given their markedly higher provisioning rates and mean visit

durations than other classes. The hypothesis that prolactin maintains provisioning behavior but does

not quantitatively regulate contributions to it could now be tested by (i) experimentally elevating the

prolactin levels of subordinate immigrants, to test the prediction that this would cause these typically

non-provisioning birds to commence provisioning behavior (e.g. see Badyaev and Duckworth (2005) for

a demonstration of this transition in the context of parental nestling feeding), (ii) experimentally reducing

the prolactin levels of the actively-provisioning classes to test whether this eliminates provisioning

behavior (e.g. Smiley and Adkins-Regan, 2018), and (iii) experimentally elevating the prolactin levels of

actively provisioning dominant males and/or natal subordinates (the manipulation proposed in the

previous paragraph), as doing so should *not* increase their provisioning rates if prolactin merely

maintains provisioning behavior without regulating contributions to it.

Given the correlative nature of our findings, it is also conceivable that prolactin plays no causal role in

the onset, maintenance or quantitative regulation of parenting and/or cooperative helping in this species

(despite experimental evidence of causal effects on parenting in other species; see Introduction;

Angelier et al., 2016). In this scenario, one might attribute the evident associations between prolactin

and provisioning to a 'reverse causal' relationship, in which provisioning interactions with offspring

stimulate prolactin release (Hall, 1987; Sharp et al., 1998). However, such a reverse causal argument

alone cannot readily account for our findings in their entirety, as within-class variation in provisioning

rates and mean provisioning visit durations were not evidently associated with prolactin levels (though,

again, the lack of such an association could be attributable to challenges with accurately and

simultaneously quantifying hormone and behavior; see above). When considering whether our findings

could be attributable solely to effects of provisioning on prolactin levels (i.e. in the absence of any effect

of prolactin on provisioning), it is worth considering why selection would have left prolactin levels

sensitive to offspring interactions in the first place. Arguably the most plausible explanation is that this

mechanism plays a role in a feedback loop in which a causal relationship exists in both directions: if

prolactin did establish, maintain and/or regulate care, selection may have favored regulating prolactin

secretion according to offspring interactions in order to maintain care as long as offspring survive and/or

regulate care according to offspring viability or need (Hall, 1987; Sharp et al., 1998; Angelier et al.,

2016). As such, where offspring cues do stimulate prolactin release, such a relationship might generally

be expected to occur alongside causal effects of prolactin on care. While the experiments outlined above

would shed light on the causality of the prolactin-provisioning associations detected here, wider

investigations are also needed to probe the role, if any, that such a feedback loop (with causal

relationships in both directions) may play in the maintenance and/or regulation of cooperative care.

While experimental tests of causality are needed, our findings are broadly consistent with the hypothesis

that pre-existing mechanisms that regulated parental care in ancestral bi-parental species were co-

opted for the regulation of cooperative helping behavior on the evolution of cooperative breeding. The

often-overlooked possibility that parenting and cooperative helping are indeed regulated by a common

mechanism has important evolutionary implications. Explanations for the evolution, maintenance and

optimization of cooperative behavior typically focus on the roles of the fitness benefits and costs of

cooperation *per se* (Hamilton, 1964; Cockburn, 1998; West et al., 2007; Capilla-Lasheras et al., 2021).

However, if cooperation and parenting are regulated by a common underlying mechanism, it is

conceivable that this shared regulatory architecture for care giving is shaped as much by the payoffs

from its outcomes in a parental context as by the payoffs from its outcomes in a cooperative helping

context. While selection might independently optimize parental and non-breeding helper caring

strategies (e.g. via the evolution of an entirely context-dependent caring strategy), it is conceivable that

mechanistic constraints preclude their independent optimization. For example, genetic variants that

modified sensitivity to begging could conceivably impact the expression of both parental care and

cooperative helping, yielding scope for intra-locus genetic conflict to constrain the independent

optimization of both parental care and cooperative helping (Pennell et al., 2018; see also the conceptual

parallels with sexual conflict: Stewart et al., 2010; Pennell and Morrow, 2013). Where this is the case,

attempts to understand the evolutionary origins, maintenance and optimization of cooperative behavior

may require attention to the extent to which genetic correlations exist between parental and cooperative

behavior. Notably, our findings suggest that cooperative helping behavior in sparrow-weaver societies

is not maintained by selection *solely* because a genetic correlation with parenting has precluded the

evolution of 'non-helping' (see Brown and Vleck, 1998 for a similar debate), because a context-

dependent helping strategy does appear to have evolved. Subordinates routinely help while within their

natal groups (where they are closely related to the broods that they help to rear; Harrison et al., 2013a),

but typically cease to do so following immigration into another group (where they are typically unrelated

to broods, reducing the potential indirect fitness payoff from helping; Harrison et al., 2013a). While

endocrine research on cooperative breeders has historically focused principally on the proximate causes

of the rank-related reproductive disparities that typify such societies (Schoech et al., 2004; Young et al.,

2006; 2008), a renewed focus on the endocrinology of care in cooperative breeders (Schoech et al.,

2004; Soares et al., 2010; Dantzer et al., 2017; 2019) would now help to shed light on the extent to

which parenting and cooperation are indeed regulated by shared underlying pathways.

CONCLUSION

Our findings lend new support to the hypotheses that helping behavior in cooperatively breeding

societies has shared mechanistic underpinnings with parental care, and that prolactin plays a key role

in this pathway (see also Vleck et al., 1991; Schoech et al., 1996; Khan et al., 2001; Carlson et al.,

2006). Our findings and their complexity highlight the need for experimental studies to investigate both

the causality and nature of the relationship between prolactin and provisioning in this species, in both

parental and cooperative helping contexts. Our findings also highlight that attempts to understand the

evolution of cooperative helping may benefit from attention to the possibility of constraints on the

independent optimization of cooperative helping and parenting. Our study has implications too for the

growing interest in the mechanistic origins of consistent individual differences in cooperative helping

behavior (Sanderson et al., 2015; Dantzer et al., 2019). Specifically, our findings highlight that such

differences could arise from consistent individual differences within the pathway by which prolactin acts

(e.g. via differences in prolactin secretion and/or reception; Ohkubo et al., 1998; Zhou et al., 1996).

ACKNOWLEDGMENTS

We dedicate this manuscript to our co-author Peter Sharp (P.J.S.), whose openness and interest in this

project were key to its inception, and whose wider contributions to avian endocrinology we celebrate.

Any errors are our own, as he did not have a chance to comment on the final manuscript. We are grateful

to D. O'Connell for support with data collection, D. L. Cram for assistance in training in bird capture and

blood sampling, N.C. Bennett for invaluable logistical support, J.L. McDonald and K.J. Walker for

comments on an earlier draft, and P. Capilla-Lasheras for statistical advice. We also thank E.O. and

Son and all at Tswalu Kalahari Reserve for access to our field site and exceptional logistical support

(with particular gratitude to Cornelia le Roux and Dylan Smith), and the Northern Cape Conservation

Authority for granting permission to conduct our research. L.A.W. was funded by a NERC PhD

studentship (NE/J500185/1). J.E.Y. was funded by the BBSRC (BB/H022716/1). S.L.M. was funded by

Roslin Institute Strategic Grant funding from the BBSRC (BB/P013759/1). A.J.Y. was funded by BBSRC

David Phillips and NERC Blue Skies Research Fellowships (BB/H022716/1 and NE/E013481/1).

REFERENCES
703

Angelier, F., Barbraud, C., Lormee, H., Prud'homme, F., and Chastel, O. (2006). Kidnapping of chicks
in emperor penguins: a hormonal by-product? *Journal of Experimental Biology* 209(8), 1413-
1420.

Angelier, F., and Chastel, O. (2009). Stress, prolactin and parental investment in birds: A review.
General and Comparative Endocrinology 163(1-2), 142-148. doi: Doi
10.1016/J.Ygcen.2009.03.028.

Angelier, F., Wingfield, J.C., Tartu, S., and Chastel, O. (2016). Does prolactin mediate parental and
life-history decisions in response to environmental conditions in birds? A review. *Horm Behav*
77**,** 18-29. doi: 10.1016/j.yhbeh.2015.07.014.

Badyaev, A., and Duckworth, R.A. (2005). "Evolution of plasticity in hormonally-integrated parental
tactcis: An example with the house finch," in *Functional Avian Endocrinology,* eds. A. Dawson
& P.J. Sharp. (New Delhi, India: Narosa Publishing House), 375-386.

Bates, D., Machler, M., Bolker, B.M., and Walker, S.C. (2015). Fitting Linear Mixed-Effects Models
Using lme4. *Journal of Statistical Software* 67(1)**,** 1-48.

Bender, N., Taborsky, M., and Power, D.M. (2008). The role of prolactin in the regulation of brood care
in the cooperatively breeding fish Neolamprologus pulcher. *J Exp Zool A Ecol Genet Physiol*
309(9)**,** 515-524. doi: 10.1002/jez.482.

Boos, M., Zimmer, C., Carriere, A., Robin, J.P., and Petit, O. (2007). Post-hatching parental care
behaviour and hormonal status in a precocial bird. *Behavioural Processes* 76(3)**,** 206-214.

Brown, J.L., and Vleck, C.M. (1998). Prolactin and helping in birds: has natural selection strengthened
helping behavior? *Behavioral Ecology* 9(6)**,** 541-545.

Buntin, J.D. (1996). Neural and hormonal control of parental behavior in birds. *Advances in the Study*
of Behavior 25**,** 161-213.

Buntin, J.D., Becker, G.M., and Ruzycki, E. (1991). Facilitation of Parental Behavior in Ring Doves by
Systemic or Intracranial Injections of Prolactin. *Hormones and Behavior* 25(3)**,** 424-444.

Capilla-Lasheras, P., Bircher, N., Brown, A.M., Harrison, X., Reed, T.E., York, J.E., et al. (2023).
Evolution of sex differences in cooperation: the role of trade-offs with dispersal. *BioRxiv*. doi:
https://doi.org/10.1101/2023.09.13.557370

Capilla-Lasheras, P., Wilson, A.J., and Young, A.J. (in press). Mothers in a cooperatively breeding bird
increase investment per offspring at the pre-natal stage when they will have help with post-
natal care. *PLOS Biology*. doi: https://doi.org/10.1101/2021.11.11.468195

Capilla-Lasheras, P., Wood, E.M., Harrison, X., Wilson, A.J., and Young, A.J. (2021). Altruistic bet-
hedging and the evolution of cooperation in a Kalahari bird. *Science Advances* 7**,** eabe8980.

Carlson, A.A., Manser, M.B., Young, A.J., Russell, A.F., Jordan, N.R., McNeilly, A.S., et al. (2006a).
Cortisol levels are positively associated with pup-feeding rates in male meerkats. *Proceedings*
of the Royal Society of London Series B-Biological Sciences 273(1586)**,** 571-577.

Carlson, A.A., Nicol, L., Young, A.J., Parlow, A.F., and McNeilly, A.S. (2003). Radioimmunoassay of
prolactin for the meerkat (Suricata suricatta), a cooperatively breeding carnivore. *General and*
Comparative Endocrinology 130(2)**,** 148-156.

Carlson, A.A., Russell, A.F., Young, A.J., Jordan, N.R., McNeilly, A.S., Parlow, A.F., et al. (2006b).
Elevated prolactin levels immediately precede decisions to babysit by male meerkat helpers.
Hormones and Behavior 50(1)**,** 94-100.

Chastel, O., Lacroix, A., Weimerskirch, H., and Gabrielsen, G.W. (2005). Modulation of prolactin but
not corticosterone responses to stress in relation to parental effort in a long-lived bird.
Hormones and Behavior 47(4)**,** 459-466.
Chastel, O., and Lormee, H. (2002). Patterns of prolactin secretion in relation to incubation failure in a
tropical seabird, the red-footed booby. *Condor* 104(4)**,** 873-876.
Cockburn, A. (1998). Evolution of helping behavior in cooperatively breeding birds. *Annual Review of*
Ecology and Systematics 29**,** 141-177.
Cornwallis, C.K., West, S.A., Davis, K.E., and Griffin, A.S. (2010). Promiscuity and the evolutionary
transition to complex societies. *Nature* 466(7309)**,** 969-U991. doi: 10.1038/nature09335.
Cram, D.L., Blount, J.D., and Young, A.J. (2015a). The oxidative costs of reproduction are group-size
dependent in a wild cooperative breeder. *Proceedings of the Royal Society B-Biological*
Sciences 282(1819). doi: https://doi.org/10.1098/rspb.2015.2031.
Cram, D.L., Blount, J.D., and Young, A.J. (2015b). Oxidative status and social dominance in a wild
cooperative breeder. *Functional Ecology* 29(2)**,** 229-238. doi: 10.1111/1365-2435.12317.
Dantzer, B., Dubuc, C., Goncalves, I.B., Cram, D.L., Bennett, N.C., Ganswindt, A., et al. (2019). The
development of individual differences in cooperative behaviour: maternal glucocorticoid
hormones alter helping behaviour of offspring in wild meerkats. *Philos Trans R Soc Lond B Biol*
Sci 374(1770)**,** 20180117. doi: 10.1098/rstb.2018.0117.
Dantzer, B., Goncalves, I.B., Spence-Jones, H.C., Bennett, N.C., Heistermann, M., Ganswindt, A., et
al. (2017). The influence of stress hormones and aggression on cooperative behaviour in
subordinate meerkats. *Proc Biol Sci* 284(1863). doi: 10.1098/rspb.2017.1248.
Dickinson, J.L., and Hatchwell, B.J. (2004). "Fitness consequences of helping," in *Ecology and*
Evolution of Cooperative Breeding in Birds, eds. W.D. Koenig & J.L. Dickinson. (Cambridge:
Cambridge University Press), 48-66.
Duckworth, R.A., Badyaev, A.V., and Parlow, A.F. (2003). Elaborately ornamented males avoid costly
parental care in the house finch (Carpodacus mexicanus): a proximate perspective. *Behavioral*
Ecology and Sociobiology 55(2)**,** 176-183.
English, S., Nakagawa, S., and Clutton-Brock, T.H. (2010). Consistent individual differences in
cooperative behaviour in meerkats (Suricata suricatta). *Journal of Evolutionary Biology* 23(8)**,**
1597-1604. doi: 10.1111/j.1420-9101.2010.02025.x.
Hall, M.R. (1987). External stimuli affecting incubation behavior and prolactin secretion in the duck
(Anas platyrhynchos). *Horm Behav* 21(3)**,** 269-287. doi: 10.1016/0018-506x(87)90015-8.
Hamilton, W.D. (1964). The genetical evolution of social behaviour. Parts I and II. *Journal of*
Theoretical Biology 7**,** 1-52.
Harrison, X., York, J., Cram, D., Hares, M., and Young, A. (2013a). Complete reproductive skew within
white-browed sparrow weaver groups despite outbreeding opportunities for subordinates of

both sexes. *Behavioral Ecology and Sociobiology* 67(12)**,** 1915-1929. doi: 10.1007/s00265-
013-1599-1.

Harrison, X.A., York, J.E., Cram, D.L., and Young, A.J. (2013b). Extra-group mating increases
inbreeding risk in a cooperatively breeding bird. *Molecular Ecology* 22(22)**,** 5700-5715. doi:
10.1111/mec.12505.

Harrison, X.A., York, J.E., and Young, A.J. (2014). Population genetic structure and direct
observations reveal sex-reversed patterns of dispersal in a cooperative bird. *Molecular Ecology*
23(23)**,** 5740-5755. doi: 10.1111/mec.12978.

Houslay, T.M., Earley, R.L., White, S.J., Lammers, W., Grimmer, A.J., Travers, L.M., et al. (2022).
Genetic integration of behavioural and endocrine components of the stress response. *Elife* 11**,**
e67126. doi: https://doi.org/10.7554/eLife.67126

Houslay, T.M., Earley, R.L., Young, A.J., and Wilson, A.J. (2019). Habituation and individual variation
in the endocrine stress response in the Trinidadian guppy (Poecilia reticulata). *Gen Comp*
Endocrinol 270**,** 113-122. doi: 10.1016/j.ygcen.2018.10.013.

Khan, M.Z., McNabb, F.M.A., Walters, J.R., and Sharp, P.J. (2001). Patterns of testosterone and
prolactin concentrations and reproductive behavior of helpers and breeders in the
cooperatively breeding red-cockaded woodpecker (Picoides borealis). *Hormones and Behavior*
40(1)**,** 1-13.

Koenig, W.D., and Dickinson, J.L. (eds.). (2004). *Ecology and Evolution of Cooperative Breeding in*
Birds. Cambridge: Cambridge University Press.

Koenig, W.D., and Dickinson, J.L. (eds.). (2016). *Cooperative breeding in vertebrates: studies of*
ecology, evolution and behavior. Cambridge University Press.

Krause, J.S., Meddle, S.L., and Wingfield, J.C. (2015). The Effects of Acute Restraint Stress on
Plasma Levels of Prolactin and Corticosterone across Life-History Stages in a Short-Lived
Bird: Gambel's White-Crowned Sparrow (Zonotrichia leucophrys gambelii). *Physiol Biochem*
Zool 88(6)**,** 589-598. doi: 10.1086/683321.

Leitner, S., Mundy, P., and Voigt, C. (2009). Morphometrics of White-browed Sparrow-Weavers
Plocepasser mahali in south-western Zimbabwe. *Ostrich* 80(2)**,** 99-102. doi: Doi
10.2989/Ostrich.2009.80.2.6.833.

Lewis, D.M. (1982). Cooperative Breeding in a Population of White-Browed Sparrow Weavers
Plocepasser-Mahali. *Ibis* 124(4)**,** 511-522.

Lukas, D., and Clutton-Brock, T. (2012). Cooperative breeding and monogamy in mammalian
societies. *Proceedings of the Royal Society B-Biological Sciences* 279(1736)**,** 2151-2156. doi:
10.1098/rspb.2011.2468.

O'Callaghan, O. (2021). *Social effects on adult survival in a wild cooperative bird.* MbyRes, University
of Exeter.

Ohkubo, T., Tanaka, M., Nakashima, K., and Sharp, P.J. (1998). Relationship between prolactin
819   receptor mRNA in the anterior pituitary gland and hypothalamus and reproductive state in male
820   and female bantams (Gallus domesticus). *General and Comparative Endocrinology* 111(2)**,**
821   167-176.

822   Ouyang, J.Q., Sharp, P.J., Dawson, A., Quetting, M., and Hau, M. (2011). Hormone levels predict
823   individual differences in reproductive success in a passerine bird. *Proceedings of the Royal*
824   *Society B-Biological Sciences* 278(1717)**,** 2537-2545. doi: Doi 10.1098/Rspb.2010.2490.

825   Pennell, T.M., Holman, L., Morrow, E.H., and Field, J. (2018). Building a new research framework for
826   social evolution: intralocus caste antagonism. *Biological Reviews* 93(2)**,** 1251-1268.

827   Pennell, T.M., and Morrow, E.H. (2013). Two sexes, one genome: the evolutionary dynamics of
828   intralocus sexual conflict. *Ecology and Evolution* 3(6)**,** 1819-1834.

829   Sanderson, J.L., Stott, I., Young, A.J., Vitikainen, E.I.K., Hodge, S.J., and Cant, M.A. (2015). The
830   origins of consistent individual differences in cooperation in wild banded mongooses, Mungos
831   mungo. *Animal Behaviour* 107**,** 193-200. doi: 10.1016/j.anbehav.2015.06.022.

832   Sanderson, J.L., Young, A.J., Hodge, S.J., Kyabulima, S., Walker, S.L., and Cant, M.A. (2014).
833   Hormonal mediation of a carry-over effect in a wild cooperative mammal. *Functional Ecology*
834   28(6)**,** 1377-1386. doi: 10.1111/1365-2435.12307.

835   Schoech, S.J., Ketterson, E.D., Nolan, V., Sharp, P.J., and Buntin, J.D. (1998). The effect of
836   exogenous testosterone on parental behavior, plasma prolactin, and prolactin binding sites in
837   dark-eyed juncos. *Hormones and Behavior* 34(1)**,** 1-10.

838   Schoech, S.J., Mumme, R.L., and Wingfield, J.C. (1996). Prolactin and helping behaviour in the
839   cooperatively breeding Florida scrub-jay, *Aphelocoma coerulescens*. *Animal Behaviour* 52**,**
840   445-456.

841   Schoech, S.J., Reynolds, S.J., and Boughton, R.K. (2004). "Endocrinology," in *Ecology and Evolution*
842   *of Cooperative Breeding in Birds,* eds. W.D. Koenig & J.L. Dickinson. (Cambridge: Cambridge
843   University Press), 128-141.

844   Sharp, P.J., Dawson, A., and Lea, R.W. (1998). Control of luteinizing hormone and prolactin secretion
845   in birds. *Comparative Biochemistry and Physiology C-Pharmacology Toxicology &*
846   *Endocrinology* 119(3)**,** 275-282.

847   Smiley, K.O., and Adkins-Regan, E. (2018). Lowering prolactin reduces post-hatch parental care in
848   male and female zebra finches (Taeniopygia guttata). *Horm Behav* 98**,** 103-114. doi:
849   10.1016/j.yhbeh.2017.12.011.

850   Soares, M.C., Bshary, R., Fusani, L., Goymann, W., Hau, M., Hirschenhauser, K., et al. (2010).
851   Hormonal mechanisms of cooperative behaviour. *Philosophical Transactions of the Royal*
852   *Society B-Biological Sciences* 365(1553)**,** 2737-2750. doi: 10.1098/rstb.2010.0151.

853   Sockman, K.W., Schwabl, H., and Sharp, P.J. (2000). The role of prolactin in the regulation of clutch
854   size and onset of incubation behavior in the American kestrel. *Hormones and Behavior* 38(3)**,**
855   168-176.

856   Solomon, N.G., and French, J.A. (eds.). (1997). *Cooperative Breeding in Mammals.* Cambridge:
857   Cambridge University Press.

858   Stewart, A.D., Pischedda, A., and Rice, W.R. (2010). Resolving Intralocus Sexual Conflict: Genetic
859   Mechanisms and Time Frame. *Journal of Heredity* 101**,** S94-S99.

860   Thierry, A.M., Brajon, S., Massemin, S., Handrich, Y., Chastel, O., and Raclot, T. (2013). Decreased
861   prolactin levels reduce parental commitment, egg temperatures, and breeding success of
862   incubating male Adelie penguins. *Horm Behav* 64(4)**,** 737-747. doi:
863   10.1016/j.yhbeh.2013.06.003.

864   Vleck, C.M., Mays, N.A., Dawson, J.W., and Goldsmith, A.R. (1991). Hormonal Correlates of Parental
865   and Helping-Behavior in Cooperatively Breeding Harris Hawks (Parabuteo-Unicinctus). *Auk*
866   108(3)**,** 638-648.

867   Voigt, C., Leitner, S., and Gahr, M. (2007). Socially induced brain differentiation in a cooperatively
868   breeding songbird. *Proceedings of the Royal Society of London Series B-Biological Sciences*
869   274**,** 2645-2652. doi: 10.1098/rspb.2007.0858|ISSN 0962-8452.

870   Vullioud, P., Mendonca, R., Glauser, G., Bennett, N., Zottl, M., Katlein, N., et al. (2021). Increases in
871   glucocorticoids are sufficient but not necessary to increase cooperative burrowing in
872   Damaraland mole-rats. *Horm Behav* 135**,** 105034. doi: 10.1016/j.yhbeh.2021.105034.

873   Walker, L.A. (2016). *The evolution and regulation of cooperation in the wild.* Ph.D., University of
874   Exeter.

875   Walker, L.A., York, J.E., and Young, A.J. (2016). Sexually selected sentinels? Evidence of a role for
876   intrasexual competition in sentinel behavior. *Behavioral Ecology* 27(5)**,** 1461-1470. doi:
877   10.1093/beheco/arw064.

878   West, S.A., Griffin, A.S., and Gardner, A. (2007). Evolutionary explanations for cooperation. *Current*
879   *Biology* 17(16)**,** R661-R672.

880   Wingfield, J.C., and Lewis, D.M. (1993). Hormonal and Behavioral-Responses to Simulated Territorial
881   Intrusion in the Cooperatively Breeding White-Browed Sparrow Weaver, Plocepasser-Mahali.
882   *Animal Behaviour* 45(1)**,** 1-11.

883   Wood, E.M. (2017). *Causes and fitness consequences of telomere dynamics in a social bird.* PhD,
884   University of Exeter.

885   Wood, E.M., Capilla-Lasheras, P., Cram, D.L., Walker, L.A., York, J.E., Lange, A., et al. (2021). Social
886   dominance and rainfall predict telomere dynamics in a cooperative arid-zone bird. *Mol Ecol*.
887   doi: 10.1111/mec.15868.

Wood, E.M., and Young, A.J. (2019). Telomere attrition predicts reduced survival in a wild social bird,
but short telomeres do not. *Mol Ecol* 28(16)**,** 3669-3680. doi: 10.1111/mec.15181.

York, J.E. (2012). *The evolution and regulation of dawn chorus performance in white-browed sparrow*
weaver society. Ph.D., University of Bristol.

York, J.E., Radford, A.N., de Vries, B., Groothuis, T.G., and Young, A.J. (2016). Dominance-related
seasonal song production is unrelated to circulating testosterone in a subtropical songbird.
General and Comparative Endocrinology 233**,** 43-52. doi: 10.1016/j.ygcen.2016.05.011.

York, J.E., Wells, K.J., and Young, A.J. (2019). Dominance-related contributions to collective territory
defence are adjusted according to the threat. *Animal Behaviour* 158**,** 25-34.

Young, A.J., Carlson, A.A., and Clutton-Brock, T.H. (2005). Trade-offs between extra-territorial
prospecting and helping in a cooperative mammal. *Animal Behaviour* 70**,** 829-837.

Young, A.J., Carlson, A.A., Monfort, S.L., Russell, A.F., Bennett, N.C., and Clutton-Brock, T. (2006).
Stress and the suppression of subordinate reproduction in cooperatively breeding meerkats.
Proceedings of the National Academy of Sciences of the United States of America 103(32)**,**
12005-12010.

Young, A.J., Monfort, S.L., and Clutton-Brock, T.H. (2008). The causes of physiological suppression
among female meerkats: a role for subordinate restraint due to the threat of infanticide?
Hormones and Behavior 53**,** 131-139.

Zhou, J.F., Zadworny, D., Guemene, D., and Kuhnlein, U. (1996). Molecular cloning, tissue
distribution, and expression of the prolactin receptor during various reproductive states in
Meleagris gallopavo. *Biology of Reproduction* 55(5)**,** 1081-1090.

Ziegler, T.E. (2000). Hormones associated with non-maternal infant care: A review of mammalian and
avian studies. *Folia Primatologica* 71(1-2)**,** 6-21.

911