1

Prolactin and the regulation of parental care and helping behavior in

cooperatively breeding white-browed sparrow weaver societies

Lindsay A. Walker[1], Linda Tschirren[2], Jennifer E. York[1,3], Peter J. Sharp[4],

Simone L. Meddle[4], Andrew J. Young[1*]

[1] Centre for Ecology and Conservation, University of Exeter Cornwall Campus, Penryn, Cornwall TR10

9EZ, UK

[2] ZHAW School of Life Sciences and Facility Management, Institute of Natural Resource Sciences,

Grüental, 8820 Wädenswil

[3] Department of Zoology, University of Cambridge, Downing Street, Cambridge, CB2 3EJ, UK

[4] The Roslin Institute, The Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter

Bush, Midlothian EH25 9RG, Scotland, UK

[*] Corresponding author

A.J.Young@exeter.ac.uk

Centre for Ecology and Conservation

College of Life and Environmental Sciences

University of Exeter, Cornwall Campus

TR10 9EZ

UK

ABSTRACT

In many cooperatively breeding societies non-breeding individuals help to rear the offspring of breeders.

The physiological mechanisms that regulate such cooperative helping behavior are poorly understood,

but may have been co-opted, during the evolution of cooperative breeding, from pre-existing

mechanisms that regulated parental care. Key among these may be a role for prolactin. Here we

investigate whether natural variation in circulating prolactin levels predicts both parental and helper

contributions to nestling provisioning in cooperatively breeding white-browed sparrow weavers,

Plocepasser mahali. In sparrow weaver groups, a single dominant pair monopolize reproduction and

non-breeding subordinates help with nestling feeding. We show that: (i) among parents, dominant

females feed nestlings at higher rates, make longer provisioning visits and have higher prolactin levels

than dominant males; and (ii) among subordinates, engaged in cooperative helping behavior, those

within their natal groups feed nestlings at higher rates and have higher prolactin levels than immigrants.

Moreover, continuous variation in prolactin levels positively predicts nestling-provisioning rates and

mean provisioning visit durations when all bird classes are combined. These relationships are principally

driven by differences among bird classes in both circulating prolactin levels and provisioning traits; the

more limited within-class variation in prolactin and provisioning traits were not evidently correlated,

highlighting a likely role for additional mechanisms in the fine-scale regulation of care. Our findings

broadly support the hypothesis that parental care and cooperative helping behavior are regulated by a

common underlying mechanism and highlight the need for experimentation to now establish the

causality of any role for prolactin.

[250 words]

INTRODUCTION

In many cooperatively breeding societies, non-breeding helpers assist with the rearing of parents'

young, via cooperative contributions to diverse forms of care (e.g. incubation, babysitting and offspring

provisioning; Solomon and French 1997, Koenig and Dickinson 2004, 2016). The majority of research

on such 'helping behavior' has sought to explain its evolution, by identifying the effects of helping on

recipients and the means by which these yield fitness benefits to helpers (Cockburn 1998, Dickinson

and Hatchwell 2004, Koenig and Dickinson 2016). By contrast, our understanding of the proximate

physiological mechanisms that regulate the expression of cooperative behavior is less advanced

(Schoech et al. 2004, Soares et al. 2010, Sanderson et al. 2014, Dantzer et al. 2017), despite a surge

of interest in the origins of consistent individual differences in both cooperative behavior and endocrine

traits (Sanderson et al. 2015, English et al. 2010, Dantzer et al. 2019, Houslay et al. 2019). As

cooperatively breeding species commonly evolved from monogamous pair-breeding species in which

parental care was well developed (Cornwallis et al. 2010, Lukas and Clutton-Brock 2012), it seems likely

that the physiological mechanisms that regulate cooperative helping behavior among non-breeders

were co-opted from the pre-existing mechanisms that regulated parental care among breeders.

Attempts to identify the proximate mechanisms that regulate helping behavior may therefore be well

served by testing candidate mechanisms already identified for the regulation of parental care in non-

cooperative species (Ziegler 2000, Schoech et al. 2004, Carlson et al. 2006a, 2006b). One such

mechanism is the neuroendocrine pathway involving the anterior pituitary gland hormone prolactin

(Buntin 1996, Sharp et al. 1998, Ziegler 2000, Carlson et al. 2006a, 2006b, Angelier et al. 2016).

Numerous studies suggest that prolactin can play a causal role in the expression of parental care,

though its precise role is not clear and seems likely to vary across taxa (Buntin 1996, Sharp et al. 1998,

Angelier et al. 2016). In birds, prolactin is thought to play a causal role in the onset and maintenance of

parental care, but it is less clear whether variation in circulating prolactin levels is also involved in the

quantitative regulation of contributions to care once caring behavior has begun (Boos et al. 2007,

72 Angelier et al. 2016, Smiley and Adkins-Regan 2018). The transition from sexual activity to parenting is

73 typically associated with an increase in circulating prolactin levels, which peak during the care period

74 (Buntin 1996, Sharp et al. 1998, Angelier et al. 2016, e.g. Schoech et al. 1996, Badyaev and Duckworth

75 2005). Evidence that naturally low prolactin levels are commonly associated with breeding attempt

76 abandonment and/or failure (e.g. Chastel and Lormee 2002, Chastel et al. 2005), and that experimental

77 reductions in circulating prolactin levels during late incubation can disrupt post-natal care (e.g. Smiley

78 and Adkins-Regan 2018), suggest that these elevated prolactin levels are necessary for the onset and/or

79 maintenance of parental care. Indeed, experimental elevations of circulating prolactin suggest that

80 elevated prolactin levels can promote the onset of both incubation behavior (e.g. Sockman et al. 2000)

81 and nestling provisioning behavior (e.g. Badyaev and Duckworth 2005). Positive associations between

82 continuous variation in circulating prolactin levels and the rates at which parents provision their offspring

83 (e.g. Duckworth et al. 2003, Ouyang et al. 2011) highlight the possibility that prolactin levels also regulate

84 the *amount* of care that an actively caring parent provides to its offspring. However, this hypothesis has

85 yet to be experimentally tested, and such positive associations could arise instead via effects of

86 provisioning activity on a bird's circulating prolactin levels, as parental contact with offspring cues can

87 increase prolactin secretion (Hall 1987, Sharp et al. 1998). Causal links between prolactin and

88 provisioning rates may therefore exist in both directions. Indeed, such a feedback loop (in which

89 offspring cues stimulate prolactin secretion that in turn maintains and/or elevates the expression of

90 parental care) could conceivably both maintain parental care while offspring survive, and regulate its

91 expression according to offspring vigor and need.

93 A number of studies of cooperatively breeding species have now begun to investigate the relationships

94 between prolactin and care-giving behavior, both among parents and non-breeding helpers (Ziegler

95 2000, Schoech et al. 2004, Soares et al. 2010). Prolactin levels have been shown to rise in parents and

96 non-breeding helpers during the transition to incubation and nestling care in at least three species of

97 cooperatively breeding bird (Schoech et al. 1996, Brown and Vleck 1998, Khan et al. 2001, see also

Vleck et al. 1991). While few studies have investigated specifically whether variation in circulating

prolactin levels predicts variation in cooperative contributions to helping, studies of at least two

cooperative breeders have yielded compelling evidence in this regard. In Florida scrub jays

(*Aphelocoma coerulescens*), breeders fed offspring at higher rates than non-breeders and showed

higher circulating prolactin levels (Schoech et al. 1996, see also Vleck et al. 1991), and those non-

breeders that helped to feed offspring showed higher prolactin levels than those that did not (Schoech

et al. 1996). Indeed, continuous variation in circulating prolactin levels predicted continuous variation in

feeding contributions, both among all birds combined and specifically among non-breeders (Schoech et

al. 1996). Similarly, in cooperatively breeding meerkat (*Suricata suricatta*) societies, continuous variation

in the prolactin levels of helpers positively predicted their cooperative contributions to both babysitting

and pup-feeding (Carlson et al. 2006a, 2006b). In the pup-feeding study, prolactin levels only predicted

the pup-feeding rates of helpers in statistical models that did not allow for an independent positive effect

of circulating cortisol levels on the animal's pup-feeding rates (Carlson et al. 2006a). Experimental work

since highlights that this putative positive effect of cortisol on pup-feeding rates may not have been

causal, however, as glucocorticoid receptor blockade increased rather than decreased pup-feeding

rates among meerkat helpers (Dantzer et al. 2017). Such relationships between circulating prolactin

levels and helping behavior are not always apparent. For example, prolactin levels did not predict the

offspring provisioning rates of helpers in red-cockaded woodpecker (*Picoides borealis*) groups (Khan et

al. 2001), and the pituitary gland prolactin mRNA levels of a cooperatively breeding fish were not

evidently related to care-giving behavior (Bender et al. 2008); though the relevant sample sizes in both

studies were modest.

The likelihood that glucocorticoids play a role in mediating the state-dependence of cooperative helping

(Sanderson et al. 2014, Dantzer et al. 2017, see also Angelier and Chastel 2009, Angelier et al. 2016),

coupled with the expectation that other neuroendocrine pathways also modulate helping potentially

independent of circulating prolactin levels (Schoech et al. 1998, Ziegler 2000, Young et al. 2005, Carlson

et al. 2006a), highlights the potential complexity of seeking relationships between continuous variation

in prolactin levels and care. Notably too, given the potential for contact with offspring to raise circulating

prolactin levels (Sharp et al. 1998, Hall 1987), it is possible that the prolactin-helping associations

documented above arose not via causal effects of prolactin on care, but via either the reverse causal

relationship or in the absence of any causal link between the two. Nevertheless, the promising findings

to date highlight the need for further studies to investigate whether natural circulating levels of prolactin

predict variation in individual contributions to both parental care and helping behavior in cooperatively

breeding species, and ultimately the use of experimental manipulations of prolactin levels to test the

causality and nature of any relationships detected (Sockman et al. 2000, Carlson et al. 2003, Badyaev

and Duckworth 2005, Smiley and Adkins-Regan 2018).

Here we investigate whether natural variation in circulating prolactin levels positively predicts the

nestling provisioning behavior of both parents and non-breeding helpers in a wild cooperatively breeding

bird, the white-browed sparrow-weaver (*Plocepasser mahali*). White-browed sparrow weavers are rain-

dependent breeders that live in year-round territorial groups throughout the semi-arid regions of sub-

Saharan Africa (Lewis 1982, Wood et al. 2021). Within each social group, a single dominant male and

female completely monopolize within-group reproduction and up to 10 non-breeding subordinates of

both sexes help to feed their nestlings (Harrison et al. 2013a, 2013b, Capilla-Lasheras et al. 2021).

Subordinates are typically offspring from previous broods that have delayed dispersal from their natal

group (and so are helping to rear their parents' young), but subordinate immigrants of both sexes do

also occur (Harrison et al. 2013a, 2013b, Harrison et al. 2014). Subordinates contribute to several

cooperative activities year-round, including territorial defense, roost construction and anti-predator

vigilance (Lewis 1982, Walker et al. 2016, York et al. 2019), and during breeding periods they contribute

substantially to nestling provisioning (Cram et al. 2015a, Capilla-Lasheras et al. 2021). Helping behavior

by subordinates has a causal positive effect on the overall rate at which nestlings are fed (Capilla-

Lasheras et al. 2021), and appears to reduce environmentally-induced variance in the reproductive

success of the dominant pair (Capilla-Lasheras et al. 2021) and lighten the post-natal provisioning

workload of the dominant female (Capilla-Lasheras 2019). While the neuroendocrine correlates of white-

browed sparrow weaver reproduction, aggression and song production have been investigated (e.g.

Wingfield and Lewis 1993, Voigt et al. 2007, York et al. 2016), the regulation of parental and helper

contributions to offspring provisioning remains unexplored.

We test three predictions of the hypothesis that prolactin plays a role in regulating the expression of

both parental care (among dominants) and cooperative helping behavior (among non-breeding

subordinates) in cooperatively breeding societies. First, with regard to parental care, we predict that

differences between the nestling provisioning rates of dominant females and dominant males will be

mirrored by parallel differences in their mean circulating prolactin levels. Dominant females are expected

to provision nestlings at higher rates and to have higher prolactin levels than dominant males (as the

12-18% incidence of extra-group paternity in this population leaves dominant females more closely

related than dominant males, on average, to the offspring that they rear; Harrison et al. 2013a). Second,

with regard to helping behavior, we predict that differences between the nestling provisioning rates of

subordinates still residing within their natal group (hereafter 'natal subordinates') and immigrant

subordinates, will also be mirrored by parallel differences in their mean circulating prolactin levels. Natal

subordinates are expected to provision nestlings at higher rates than immigrants, as while the former

are typically rearing future generations of siblings born to their parents, the latter will typically be

unrelated to the nestlings in their group (Harrison et al. 2013a). Finally, we predict that continuous

variation in circulating prolactin levels will positively predict continuous variation in the provisioning rates

of birds, and that this relationship will be apparent (i) at the population level (when all four of the bird

classes above are combined), and (ii) within bird classes, having factored out the among-class

differences in prolactin levels and provisioning rates.

METHODS

General field methods

Data were collected in the context of a long-term research project that monitors ~40 cooperative groups

of white-browed sparrow weavers at Tswalu Kalahari Reserve, South Africa (27°160S, 22°250E). Data

were collected at a similar time in two separate breeding seasons (January to February 2013, and

January to March 2014). White-browed sparrow weavers in this population may breed at any time from

September through to May (the Southern summer), depending on the timing of unpredictable summer

rainfall (Capilla-Lasheras 2019, Wood et al. 2021). Each bird within our study population is fitted with a

metal ring and three color rings, providing a unique ring combination for identification in the field

(SAFRING license 1444). From around six months of age, males and females of the focal subspecies

(*Plocepasser mahali mahali*) can be distinguished by their bill color; males have a dark brown bill while

females have a paler grey-to-pink bill (Leitner et al. 2009). Dominance status and social group

compositions were determined via regular (at least twice weekly) group visits. Social dominance was

assigned based on the monitoring of key dominance-related behaviors: the dominant pair routinely

displace other group members and produce synchronized duet song, the dominant female is the sole

incubator, and the dominant male consistently produces dawn song during breeding periods (Harrison

et al. 2013a, Cram et al. 2015b, York et al. 2016). The dispersal status (natal or immigrant) of

subordinate birds was determined via the continuous monitoring of the study population since 2007.

Based on this information, four classes of birds were assigned: Dominant Females; Dominant Males;

Natal Subordinates and Immigrant Subordinates. Social group size was defined as the number of adult

(> 1 year of age) birds consistently seen foraging and roosting together at the time of the focal breeding

attempt. The breeding status of each group was determined by monitoring the contents of all woven

nest structures within each group's territory, at least every other day throughout the two study periods.

When one or more eggs were newly detected, the active nest was visited daily in the afternoon until no

new eggs were detected (the birds lay one egg per day in the morning, and typically lay clutches of 2

eggs (range 1-4); Harrison et al. 2013a). To determine hatch dates, daily monitoring of the active nest

resumed 14 days after the detection of the first egg (as incubation lasts 14-19 days; Harrison et al.

2013a). This method yielded accurate information on the day on which the first nestling in each clutch

hatched, which was termed 'Day 1' of the nestling provisioning period for the focal breeding attempt. All

protocols were approved by the Ethics Committees of the Universities of Exeter and Pretoria and

complied with regulations stipulated in The Association for the Study of Animal Behaviour (ASAB)

Guidelines for Use of Animals in Research.

Monitoring provisioning behavior

To identify individuals during the recording of nestling provisioning events, group members were

captured from their roost chambers during the incubation period (details below) and marked on the vent

with a unique dye-mark. The dominant female was left unmarked to minimize disturbance during

incubation, but could still be distinguished from other group members by being the only unmarked bird

within her group (only resident group members provision offspring). To record provisioning events, a

Panasonic SDR-S50 camcorder attached to a tripod (approximately 0.5 meters in height) was placed

on the ground beneath the entrance to the active nest two days before recording commenced (to allow

the birds to habituate to it). On the days of provisioning monitoring, the recordings were started between

06:15 and 07:54, with this start time being adjusted through the season to maintain an approximately

constant time offset from sunrise, and an expected video duration of approximately 3 hours. Provisioning

videos were collected in this way for all focal breeding attempts (n = 37 broods across 30 social groups)

on two mornings between Days 6 and 9 inclusive of the nestling provisioning period (typically for the

two consecutive mornings of Days 7 and 8; nestlings fledged from day 20). This approach yielded a

mean total duration of provisioning video of 6.08 hours (range 4.05 – 8.12 hours) per breeding attempt.

Video recordings were transcribed using VLC Media Player version 2.2, with the observer recording, for

each provisioning visit, the identity of the bird visiting the nest (determined via their distinct dye mark

and bill color, which reveals their sex) and the duration of time that they spent within the nest (the time

elapsed between passing in and out of the enclosed nest structure; hereafter 'Provisioning visit

duration'). Prior work on this study population using within-nest cameras has shown that all nest visits

during the nestling age window studied here entail the delivery of a single prey item to the brood, unless

the visiting bird is carrying a feather or grass in which case no food is delivered (Walker 2016). We

therefore excluded such feather- or grass-carrying nest visits from our provisioning visit records. From

the transcribed data for each focal brood we then calculated two provisioning trait values for each adult

group member: (i) 'Provisioning rate' (feeds / hr) was calculated as the total number of provisioning visits

that the bird conducted over the two monitored mornings divided by the total duration of video collected

over those two mornings, and (ii) 'Mean Visit Duration' (minutes) was calculated as the mean duration

of all provisioning visits conducted by the focal bird over the two monitored mornings.

Bird capture and blood sampling

To obtain a matched blood sample for prolactin measurement, we attempted to capture and blood

sample all adult (> 1 year old at the time of sampling) birds within the monitored brood's social group on

the evening of the second day of provisioning behavior recording. Birds were captured individually at

night from the woven roost chambers within their group's territory (Cram et al. 2015a) by flushing

individuals into a custom-made capture bag. All captures, leg ring fitting, dye-marking and blood

sampling were conducted by a single investigator. Birds were then immediately returned to a roost

chamber within their territory to pass the remainder of the night. Upon capture, a blood sample (*c.* 140

µL) was taken from the brachial vein of the bird using a 26-g needle and heparinized capillary tubes.

Captures occurred soon after dusk, once the birds were roosting in their woven chambers. Time of

capture was recorded (to allow us to fit the time lag from sunset to capture as a covariate predictor in

our prolactin analyses, in case of diel variation and/or effects of the time elapsed since roosting on a

bird's prolactin levels), along with the time lag between capture and the completion of blood sampling

(mean ± standard deviation [S.D.]: 3.12 ± 0.73 minutes [range 1.65 – 4.78 minutes]; to allow us to control

for potential mean effects of capture stress on prolactin levels in our statistical models). Blood samples

were immediately centrifuged in the field (12,000 g for 3 minutes; Haematospin 1400; Hawksley Medical

and Laboratory Equipment, Lancing, UK) and the plasma was drawn off and stored in a cryovial on ice

until it could be transferred to liquid nitrogen on return from the field (mean ± S.D. time lag from sample

collection to storage on liquid nitrogen = 148 ± 63 min). At the end of the field season, samples were

transferred to the UK on dry ice and then stored at -80 degrees Celsius until analysis for prolactin.

Prolactin Radioimmunoassay

The prolactin assay was carried out at the Roslin Institute (University of Edinburgh, Easter Bush,

Midlothian, Scotland, UK). Plasma prolactin levels were measured using a highly specific heterologous

micro-radioimmunoassay of donkey anti-rabbit serum to European starling (*Sturnus vulgaris*) prolactin

(Sharp antibody code 44/2). Prolactin was radiolabeled with iodine^{125} using chloramine-T. 168 (out of a

total of 208) samples were assayed in duplicate, and the remaining 40 samples were assayed as

singletons (not all samples assayed were for use in this study). All samples were measured in a single

assay, in which the intra-assay coefficient of variation for the duplicate samples was 3.31%.

Statistical methods

The above methods yielded a final data set of matched provisioning trait data (estimated from the focal

bird's average performance over two mornings of provisioning recordings; see above) and circulating

prolactin levels (when sampled on the evening of the second day of provisioning monitoring) for 70

different adult birds, each sampled once (for all traits), while feeding a total of 37 broods across 30 social

groups. For a small number of these birds, a second matched measure of the provisioning traits and

prolactin levels was also available from a subsequent breeding attempt. However, these few repeat

measures were not included in our final data set for analysis, as attempts to conduct mixed effects

models using bird ID to account for the presence of these repeated measures of individuals typically

failed to estimate non-zero variance for bird ID (quite possibly because too few repeated measures were

available). All analyses therefore utilized the data from only the first sampling event per bird, obviating

the need to include bird ID as a random effect within our mixed effects models (see below). While some

analyses utilized the whole data set (i.e. n = 70 adults birds each sampled once for all traits), others

used subsets of it (e.g. when focusing only on dominants or subordinates), and so the sample sizes for

each analysis are reported within the relevant results section and model output table. As mean

provisioning visit duration data were only available for birds that had a non-zero provisioning rate, the

sample sizes for mean visit duration analyses were sometimes smaller than those for provisioning rate.

All statistical models and visualizations were carried out in R (version 4.1.0; R Core Team). Mixed effects

modelling was conducted using the R package 'lme4' (Bates et al. 2015), and the importance of a given

combination of model predictors was assessed using an information-theoretic (IT) approach. Each

modelling exercise started with the specification of a 'global model' containing all predictors of interest.

This global model and all possible simpler models containing subsets of the global model's fixed effect

predictors (including the intercept-only model) were then fitted to the data using maximum likelihood

and ranked based on AICc (Akaike's Information Criterion corrected for small sample size; Burnham

and Anderson 2002). This 'all subsets' approach was appropriate as all simpler models were biologically

plausible. ΔAICc values (i.e. the difference in AICc between the focal model and the best supported 'top'

model) were then calculated for every model fitted. As lower AICc values are indicative of stronger

statistical support for a given model, the ΔAICc for the best-supported 'top' model = 0 and models that

attracted less statistical support had progressively more positive ΔAICc values. We gave consideration

to models within a ΔAICc value of six (Richards 2008, Richards et al. 2011) and subsequently reduced

this Δ6 'top model set' by applying the 'nesting rule' described in (Richards 2008). This rule aims to

avoid the retention of overly complex models that do not improve model fit, by discarding models that

are more complex versions of simpler (nested) models with weaker AIC support (Richards 2008, Arnold

2010). Adding variables with little or no explanatory power to a top model can weaken AIC by less than

6 points, leading to the retention within the Δ6 top model set of more complex versions of better

supported models, containing such uninformative variables (Arnold 2010). The nesting rule thus reduces

the chance of considering such overly complex models containing uninformative variables (Richards

2008). In the model selection tables presented within the main paper (Tables 1 & 2) each line presents

a model from the top model set (i.e. those models within ΔAICc = 6 of the best-supported model) after

the application of the nesting rule, while each of the fixed effect (predictor) columns presents the

estimated effect size for that fixed effect predictor within that model (or is blank if the focal fixed effect

was absent from that model). The corresponding full top model sets *prior* to the application of the nesting

rule are presented in the Appendices (Tables A1 & A2) in case of interest. The specific modelling

exercises conducted for each results section are described below.

1. Are differences in the nestling provisioning rates and nest visit durations of dominant females

and males (engaged in parental care) mirrored by differences in their circulating prolactin levels?

We used two separate mixed effects models with Gaussian error structure to model the causes of

variation in (i) the provisioning rates and (ii) the mean provisioning visit durations of dominant birds. The

two modeling exercises began with an identical global model structure. In addition to the primary

predictor of interest, 'bird class' (dominant female or dominant male), we fitted the following terms as

fixed effect predictors: brood size (the brood size being fed), adult group size (the number of adult group

members during the focal nestling provisioning period) and year (a two-level factor capturing the year in

which sampling occurred; 2013 or 2014). We fitted both social group ID and brood ID (the identity of the

brood being fed) as random effects, retaining them in the model structure regardless of the degree of

variance that they explained.

We then used a third mixed effects model with Gaussian error structure to model the causes of variation

in the circulating prolactin levels of dominant birds, starting with a global model structure containing the

same fixed and random effect predictors as the provisioning trait models just described, but with the

addition of two further fixed effects to account for potential methodological effects on prolactin

concentrations: (i) the time lag from sunset to the bird's capture for blood sampling (to allow for the

possibility of diel variation in prolactin levels and/or changes in prolactin levels once the birds entered

their roosts) and (ii) the time lag from capture to the completion of blood sampling (to allow for possible

effects of capture stress on circulating prolactin levels). While neither of these two time-lag variables

were present within the best-supported model that arose from model comparisons (suggesting no

appreciable effect of these variables on circulating prolactin levels; Table 1), we nevertheless verified

that our inferences regarding the effect of bird class on prolactin levels on the basis of the best-supported

model overall were also upheld by the best-supported model that contained both of these time lag terms

(see results text).

2. Are differences in the provisioning rates of natal and immigrant subordinates (engaged in

cooperative helping behavior) mirrored by differences in their circulating prolactin levels?

We then used two mixed effects models with Gaussian error structure to model the causes of variation

in the provisioning rates and prolactin levels of subordinate birds engaged in helping behavior. We fitted

the same set of fixed and random effect predictors to these models as were fitted to the corresponding

provisioning rate and prolactin level models conducted for dominant birds (see above) with two

exceptions: (i) here 'bird class' reflected whether the bird was a natal subordinate or an immigrant

subordinate, and (ii) here only brood ID was fitted as a random effect (social group ID was not, as all

subordinate birds from any given social group were sampled while feeding the same single brood,

leaving brood ID and social group ID with identical structure in this case). Again, we verified that our

inferences regarding the effect of bird class on prolactin levels on the basis of the best-supported model

overall were also upheld by the best-supported model that contained both of the methodological time

lag terms (see results text). We did not model the causes of variation in the mean provisioning visit

durations of subordinates as too few immigrant subordinates actually provisioned the focal broods,

leaving us with an insufficient sample size of measures of the provisioning visit durations of this bird

class.

3. Does continuous variation in prolactin levels predict continuous variation in provisioning rates?

To investigate whether natural variation in prolactin levels predicted continuous variation in the birds' nestling provisioning rates and mean provisioning visit durations at the population level (i.e. when all bird classes were combined) we conducted two mixed effect models (one for each provisioning trait response term), with circulating prolactin concentration as the sole fixed effect predictor (as we have not hypothesized specific mechanisms by which other variables might impact provisioning traits independent of prolactin levels) and social group ID and brood ID as random effects. Mean provisioning visit duration was logarithm transformed for analysis, to normalize model residuals. Inspection of the patterns of the mean prolactin levels and provisioning trait values of the different bird classes (i.e. dominant females, dominant males, natal subordinates and immigrant subordinates) suggested that any such continuous relationship between prolactin levels and provisioning trait values at the population level could be driven principally by the variation in these traits *among* the bird classes (Figures 3a & 3c). In order to then investigate whether the more limited variation in circulating prolactin levels *within* bird classes predicted the *within*-class variation in provisioning trait values, we first mean-centered each birds' prolactin level and provisioning trait values (the log transformed values in the case of mean provisioning visit duration) around the mean value of the focal trait for birds of their class (by subtracting from it the mean value of the focal trait for their bird class). We then conducted two mixed effects models (one for each mean-centered provisioning trait response term), with mean-centered circulating prolactin concentration as the sole fixed effect predictor and social group ID and clutch ID as random effects.

RESULTS

1. Are differences in the nestling provisioning rates and nest visit durations of dominant females

and males (engaged in parental care) mirrored by differences in their circulating prolactin levels?

Analyzing the provisioning behavior of dominant birds engaged in parental care (n = 46 dominants, 20

females and 26 males, feeding 31 broods at 28 social groups) revealed strong evidence that dominant

females feed offspring at higher rates than dominant males (Figure 1a; bird class was present in all

models down to ΔAICc = 42.36; Table 1) and that the mean provisioning visit durations of dominant

females are also longer than those of dominant males (Figure 1b; ΔAICc = 11.49; Table 1). There was

also evidence that brood size positively predicted the provisioning rates of dominant birds (ΔAICc =

1.92; Table 1), but not their mean visit durations (the best-supported visit duration model containing

brood size scored 1.42 AICc points below the top model and was rejected under the nesting rule; Table

1; Table A1). There was no compelling evidence that the provisioning rates or mean visit durations of

dominant birds were associated with either the year of study or adult group size (which were absent

from the best-supported model for both traits; Table 1).

Analyzing the circulating prolactin levels of dominant birds during the provisioning periods analyzed

above (again, n = 46 dominants, 20 females and 26 males, feeding 31 broods at 28 social groups)

revealed strong evidence that dominant females also have higher circulating prolactin levels than

dominant males (Figure 1c; bird class was present in all models down to ΔAICc = 19.93; Table 1). There

was no compelling evidence that the prolactin levels of dominants were associated with group size,

brood size, the time lag from sunset to capture or the time lag from capture to blood sampling (none of

which were present within the best-supported model; Table 1). Among the considered models that

included both of the methodological time lag terms (i.e. those models forced to account for any effects

of these terms, regardless of their importance), the best-supported model again contained an effect of

bird class (see Table A1; model number 22).

Figure 1. The (a) provisioning rates, (b) mean provisioning visit durations, and (c) circulating prolactin

concentrations of dominant females (Dom Fem) and dominant males (Dom Male) engaged in parental

care. Squares and error bars present the predicted means ± S.E. from the top model for the relevant

trait (Table 2) while controlling for the effects of the other variables present in that model. The points

show the raw data points.

Table 1. Modelling the Provisioning Behavior and Prolactin Levels of DOMINANTS

Model selection of the factors affecting the (i) nestling provisioning rates, (ii) mean provisioning visit

durations and (iii) prolactin levels of dominant females ('Dom Fem') and dominant males ('Dom Male')

engaged in parental care. The table shows all models within <6 ΔAICc of the top model, after applying

the model nesting rule (Richards et al., 2011; see methods). The grey lines indicate the response term

for the set of models beneath it. The numbers indicate coefficients (± standard errors; S.E.) and df is

the degrees of freedom. The 'Year' effect is the change from 2013 to 2014. Bird Class was present

within all models down to ΔAICc = 42.36 in the provisioning rate analysis, all models down to ΔAICc =

11.49 in the mean visit duration analysis, and all models down to ΔAICc = 19.93 in the prolactin analysis.

N = 46 dominant birds, 20 female and 26 male, feeding 31 broods at 28 social groups for the provisioning

rate and prolactin level analyses. The sample size for the mean provisioning visit duration analysis was

slightly smaller (n = 44 dominant birds), as 2 dominant males did not provision their brood, leaving us

without a measure of their mean visit duration. Table A1 presents these top model sets prior to

application of the model nesting rule.

Intercept	Bird Class (Dom Fem vs Dom Male)	Brood Size	Group Size	Year	df	AICc	ΔAICc
(i) Provisioning Rate (feeds / hr)							
-0.36 (0.87)	5.05 (0.58)	1.44 (0.46)			6	205.8	0.00
1.21 (0.50)	5.12 (0.59)			1.59 (0.59)	6	207.7	1.92
2.06 (0.42)	5.12 (0.61)				5	211.8	5.98
(ii) Mean Provisioning Visit Duration (minutes)							
0.58 (0.21)	1.28 (0.31)				5	139.7	0.00
(iii) Plasma Prolactin (ng / ml)							
1.98 (0.22)	1.62 (0.25)				5	149.6	0.00

2. Are differences in the provisioning rates of natal and immigrant subordinates (engaged in

helping behavior) mirrored by differences in their circulating prolactin levels?

Analyzing the provisioning behavior of subordinate birds engaged in cooperative helping behavior (n =

24 subordinates, 17 natal and 7 immigrant, feeding 16 broods at 16 social groups) revealed evidence

that subordinates within their natal groups feed offspring at higher rates than immigrant subordinates

(Figure 2a; bird class was present in all models down to ΔAICc = 10.64; Table 2). There was also

evidence that brood size positively predicted subordinate provisioning rates (ΔAICc = 2.13; Table 2).

There was no compelling evidence that helper provisioning rates were associated with either group size

or the year of study (neither term was present within the best-supported model; Table 2). No analysis of

the provisioning visit durations of subordinates was conducted as an insufficient number of subordinate

immigrants ever provisioned the broods (see Figure 2a).

Analyzing the circulating prolactin levels of subordinate birds during the provisioning periods analyzed

above (again, n = 24 subordinates, 17 natal and 7 immigrant, feeding 16 broods at 16 social groups)

revealed that natal subordinates also have higher prolactin levels than immigrant subordinates (Figure

2b; bird class was present in all models down to ΔAICc = 4.15; Table 2). There was also evidence that

subordinate prolactin levels were negatively associated with group size (ΔAICc = 4.15) and very weak

evidence that they were higher in the second year of study (ΔAICc = 0.02). There was no compelling

evidence that subordinate prolactin levels were associated with brood size or the time lags from sunset

to capture and from capture to blood sampling (none of which appeared in the best-supported model;

Table 2). Among the considered models that included both of the methodological time lag terms (i.e.

those models forced to account for any effects of these terms, regardless of their importance), the best-

supported model again contained an effect of bird class (see Table A2; model number 30).

Figure 2. The (a) provisioning rates, and (b) circulating prolactin levels of natal subordinates (Sub Natal) and immigrant subordinates (Sub Imm) engaged in cooperative helping behavior, feeding the broods of the dominant male and female. Y axis scales match those in Figure 1 to facilitate comparison. Squares and error bars (which do not always extend beyond the square) present the predicted means ± S.E. from the top model for the relevant trait (Table 2) while controlling for the effects of the other variables present in that model. The points show the raw data points. No analysis of provisioning visit durations was conducted as an insufficient number of subordinate immigrants ever provisioned the broods (see panel 2a).

Table 2. Modelling the Provisioning Behavior and Prolactin Levels of SUBORDINATES

Model selection of the factors affecting the (i) nestling provisioning rates, and (ii) prolactin levels of natal

subordinates ('Sub Natal') and immigrant subordinates ('Sub Imm') engaged in cooperative helping

behavior. The table shows all models within <6 ΔAIC of the top model, after applying the model nesting

rule (Richards et al., 2011; see methods). The grey lines indicate the response term for the set of models

beneath it. The numbers indicate coefficients (± standard errors; S.E.) and df is the degrees of freedom.

The 'Year' effect is the change from 2013 to 2014. Bird Class was present within all models down to

ΔAICc = 10.64 in the provisioning rate analysis. N = 24 subordinate birds, 17 natal and 7 immigrant,

feeding 16 broods at 16 social groups. Table A2 presents these top model sets prior to application of

the model nestling rule.

Intercept	Bird Class (Sub Natal vs Sub Imm)	Brood Size	Group Size	Year	df	AICc	ΔAICc
(i) **Provisioning Rate (feeds / hr)**							
-1.39 (0.68)	1.52 (0.34)	0.92 (0.37)			5	66.6	0.00
0.12 (0.34)	1.49 (0.37)				4	68.8	2.13
(ii) **Plasma Prolactin (ng/ml)**							
2.20 (0.54)	0.89 (0.28)		-0.41 (0.13)	0.56 (0.28)	6	57.13	0.00
2.82 (0.48)	0.91 (0.31)		-0.51 (0.12)		5	57.15	0.02
1.02 (0.17)				0.86 (0.32)	4	61.28	4.15
2.74 (0.56)			-0.34 (0.13)		4	61.42	4.29

3. Does continuous variation in prolactin levels predict continuous variation in provisioning rates?

Our analysis at the population level, including birds of all classes, revealed strong evidence that a bird's circulating prolactin level positively predicts both its provisioning rate (Figure 3a; effect size ± S.E. = 1.20 ± 0.23; the provisioning rate model containing the prolactin predictor scored 20.36 AICc points stronger than the intercept-only model; n = 70 birds feeding 37 broods at 30 social groups) and its mean provisioning visit duration (Figure 3c; effect size ± S.E. = 0.32 ± 0.076; the mean visit duration model containing the prolactin predictor scored 12.59 AICc points stronger than the intercept-only model; n = 59 birds feeding 36 broods at 30 social groups). Plotting out the mean prolactin levels and provisioning trait values of the different bird classes (Figure 3a & 3c), reveals that both of these population-level relationships between prolactin and provisioning traits are driven in large part by the among-bird-class differences in prolactin levels being mirrored by parallel among-bird-class differences in mean provisioning rate (Figure 3a) and mean provisioning visit duration (Figure 3c). Indeed, after mean-centering each bird's prolactin level and provisioning trait values around the focal trait's mean value for their bird class, we found no evidence that within-bird-class variation in prolactin levels predicted within-bird-class variation in either provisioning rate (Figure 3b; effect size ± S.E. = -0.15 ± 0.21; the model containing the mean-centered-prolactin predictor scored 1.87 AICc points weaker than an intercept-only model) or mean provisioning visit duration (Figure 3d; effect size ± S.E. = -0.049 ± 0.081; the model containing the mean-centered prolactin predictor scored 2.03 AICc points weaker than an intercept-only model).

Figure 3. At the population level, considering all bird classes together, natural variation in circulating prolactin levels predicts variation in both (**a**) provisioning rate and (**c**) mean provisioning visit duration. These relationships are driven principally by differences among the mean trait values of the different focal bird classes (presented ± S.E. by the squares and diamonds within panels **a** and **c**; see legend within panel **a**). Follow-up analyses revealed no evidence that variation in prolactin levels *within* these bird classes predicted within-bird-class variation in either (**b**) provisioning rate or (**d**) mean provisioning visit duration. In panels **a** and **c** the line and shaded ribbon present the predicted mean relationship and its standard error, while the shaded circular points within all panels present the raw data points.

DISCUSSION

This study investigated the hypothesis that prolactin plays a role in the regulation of nestling provisioning, both among dominant birds (engaged in parental care) and subordinate birds (that engage in cooperative helping behavior), in cooperatively breeding white-browed sparrow weaver societies. Among dominants engaged in parental care, we found that the dominant female (the mother) fed offspring at higher rates, made longer provisioning visits and had higher circulating prolactin levels than the dominant male (typically the father). Among subordinates, we found that natal subordinates helped to feed offspring at higher rates and had higher circulating prolactin levels than immigrant subordinates. Indeed, when all bird classes were combined, we found that continuous variation in the circulating prolactin levels of the birds predicted continuous variation in their provisioning rates and mean provisioning visit durations. These patterns appear to be driven principally by correlated differences among the four different bird classes in their prolactin levels and provisioning traits. We found no evidence that the more limited variation in circulating prolactin levels within the different bird classes predicted the more limited within-class variation in their provisioning traits. Together, these findings are broadly consistent with the hypothesis that parental care and cooperative helping behavior are regulated by a common underlying mechanism, and the hypothesis that prolactin plays a role in that pathway, but also highlight the need for experimental studies to now probe the causality and nature of any role for prolactin. Below, we discuss potential explanations for these findings, the different roles that prolactin could conceivably play in the regulation of parenting and cooperative helping in this species, and the wider implications of our findings for mechanistic and evolutionary research on cooperative behavior.

While our findings are broadly consistent with the hypothesis that parental care and helping behavior are regulated by a common mechanism in which prolactin plays a role, the lack of a relationship between within-class variation in prolactin levels and provisioning traits, coupled with the correlative nature of our findings, leave it important to consider the range of possible roles that prolactin could play in the regulation of provisioning behavior in this species. At least three main possibilities exist, which will now

require careful experimentation to tease apart. First, it is possible that circulating prolactin is one key

regulator of continuous variation in individual contributions to offspring provisioning, both among parents

and helpers. While most of our findings are consistent with this hypothesis, the absence of evident

relationships between within-class variation in prolactin levels and provisioning traits complicates this

view. However, the lack of *within*-class relationships could be attributable simply to a major source of

variation in both traits (*among*-class variation) having been factored out at this stage, leaving these

within-class analyses seeking more subtle prolactin-provisioning relationships (than those at the

population level) that could be obscured by a number of mechanisms. For example, difficulties with the

synchronous and accurate assessment of both prolactin levels and provisioning rates could have yielded

noise in the data set that precluded the detection of these more subtle prolactin-provisioning

relationships. While we sampled birds for prolactin on the evening following the morning provisioning-

monitoring session (a time lag comparable to, or shorter than, those of similar studies; e.g. Duckworth

et al. 2003, Ouyang et al. 2011), individuals may have differed in the way that their prolactin levels

changed during the day, leaving their evening prolactin levels only a modest proxy for those while

provisioning. The focal birds also varied in the timing of blood sampling, and while our analyses did not

detect any overall effects on prolactin levels of the time lags either from sunset to capture or from capture

to sampling, any individual variation in the circadian rhythm of prolactin secretion or in the prolactin

stress response (if this species shows one; Krause et al. 2015) could have further decoupled the

assessed prolactin levels from those during provisioning. Ultimately though, even if prolactin levels were

a key regulator of provisioning behavior, alternative mechanisms are also expected to impact

provisioning rates potentially independent of circulating prolactin levels, leaving the relationship between

natural variation in prolactin levels and provisioning behavior potentially modest in the first place

(Schoech et al. 1998, Angelier et al. 2016). Key among these could be (i) variation in other components

of a prolactin-mediated pathway (such as inter-individual and temporal variation in the density of

prolactin receptors; Zhou et al. 1996, Ohkubo et al. 1998), as well as (ii) mechanisms that may impact

provisioning behavior via prolactin-independent pathways (e.g. the effects of circulating testosterone;

Schoech et al. 1998). Moreover, even if prolactin levels determined provisioning 'motivation', the extent

to which variation in provisioning motivation was reflected in provisioning rates would depend upon the

prey capture skills of the focal bird and the environmental availability of prey. Indeed, all points

considered, it is arguably mechanistically naïve to expect particularly fine-grained associations between

the levels of a single hormone and behavior to be evident in natural populations even where a causal

link exists. To now robustly test the hypothesis that prolactin regulates continuous variation in the

magnitude of individual contributions to offspring provisioning, there is a need to experimentally elevate

the circulating prolactin levels of actively provisioning birds whose natural prolactin levels are not at the

upper end of the physiological range (dominant males and natal subordinates may serve this purpose

well; Figure 3a). This manipulation would allow one to test the key prediction that an increase in the

prolactin levels of an actively provisioning bird will cause it to increase its provisioning rate; a prediction

that to our knowledge has yet to best tested in either a parenting or helping context (the few experimental

elevations of endogenous prolactin secretion in a provisioning context to date have focussed on the

establishment of provisioning in non-provisioning birds rather than its quantitative variation within

actively provisioning birds; e.g. Badyaev and Duckworth 2005).

A second potential explanation for the balance of our findings is that prolactin could instead play a causal

role in the onset and maintenance of provisioning behavior among parents and helpers, without playing

a role in the quantitative regulation of contributions to provisioning among actively provisioning birds.

For example, a threshold level of prolactin may be required for the onset and/or maintenance of

provisioning behavior (Angelier et al. 2006, Boos et al. 2007). Under this scenario, the higher prolactin

levels of natal subordinates and dominant birds, relative to immigrant subordinates, could be causally

responsible for the former bird classes engaging in provisioning while the latter typically did not. This

could be the case without prolactin playing any causal role in regulating continuous variation in the

provisioning rates of actively provisioning birds; a scenario that could account for the lack of within-class

correlations between prolactin levels and provisioning behavior. The elevated prolactin levels of

dominant females (relative to dominant males and natal subordinates) could conceivably be a

downstream consequence of either a role for prolactin in incubation (Buntin 1996, Sharp et al. 1998,

Khan et al. 2001; as dominant females are the sole incubator in this species) and/or their differential

exposure to offspring cues during the nestling period (which can increase prolactin secretion; Hall 1987,

Sharp et al. 1998), given their markedly higher provisioning rates and mean visit durations than other

classes. The hypothesis that prolactin maintains provisioning behavior but does not quantitatively

regulate contributions to it could now be tested by (i) experimentally elevating the prolactin levels of

subordinate immigrants, to test the key prediction that this would cause these typically non-provisioning

birds to commence provisioning behavior (e.g. see Badyaev and Duckworth (2005) for a rare

demonstration of this transition in the context of parental nestling feeding), (ii) experimentally reducing

the prolactin levels of the actively-provisioning classes to test whether this eliminates provisioning

behavior (e.g. Smiley and Adkins-Regan 2018), and (iii) experimentally elevating the prolactin levels of

actively provisioning dominant males and/or natal subordinates (the manipulation proposed in the

previous paragraph), as doing so should *not* increase their provisioning rates if prolactin merely

maintains provisioning behavior without regulating contributions to it.

Given the correlative nature of our findings, it is also conceivable that prolactin plays no role in the onset,

maintenance or quantitative regulation of parenting and cooperative helping behavior. Under this

scenario, one might attribute the evident associations between prolactin and provisioning to a 'reverse

causal' relationship, in which provisioning interactions with offspring stimulate prolactin release (Hall

1987, Sharp et al. 1998). However, such a reverse causal argument alone cannot readily account for

our findings in their entirety, as within-class variation in provisioning rates and mean provisioning visit

durations were not evidently associated with circulating prolactin levels (though, again, the lack of such

an association could be attributable to challenges with accurately and simultaneously quantifying both

hormone and behavior; see above). When considering whether our findings could be attributable solely

to effects of provisioning behavior on prolactin levels (i.e. in the absence of any effect of prolactin on

provisioning), it is worth considering why selection would have left prolactin levels sensitive to offspring

interactions in the first place. Arguably the most plausible explanation is that such a mechanism plays

a role in a feedback loop in which a causal relationship exists in both directions: if prolactin did establish,

maintain and/or quantitatively regulate care, selection may have favored regulating prolactin secretion

according to offspring interactions in order to maintain care as long as offspring survive and/or regulate

care according to offspring viability or need (Hall 1987, Sharp et al. 1998, Angelier et al. 2016). As such,

where offspring cues do stimulate prolactin release, such a relationship might generally be expected to

occur alongside causal effects of prolactin on care. While the experiments outlined above would shed

light on the causality of the prolactin-provisioning associations detected here, wider investigations are

also needed to probe the role, if any, that such a feedback loop (with causal relationships in both

directions) may play in the maintenance and/or regulation of cooperative care.

Our analyses revealed evidence that the circulating prolactin levels of subordinates were lower in larger

social groups, but that the same relationship was not apparent among dominants. This finding is notable

in that it parallels the apparent effects of group size on survival in this study population: subordinate

birds are markedly less likely to survive in larger groups, but the same is not true for dominants

(O'Callaghan & Young *unpublished data*). This survival relationship may be driven by enhanced foraging

competition in larger groups (which subordinates may suffer from to a greater extent than dominants,

given their lower rank), as territory size does not increase proportionally with group size (Martin-Taylor

2018; likely leaving resources more strongly contested in larger groups). Accordingly, subordinates in

larger groups show significantly lower body condition (O'Callaghan & Young *unpublished data*); a

relationship that could conceivably account for their reduced prolactin levels in larger groups too.

Chronic exposure to nutritionally stressful conditions is often associated with reduced prolactin levels; a

relationship that could be due in part to negative effects of circulating corticosterone on prolactin release

(Delehanty et al. 1997, Criscuolo et al. 2005, Angelier and Chastel 2009, Riechert et al. 2014, Angelier

et al. 2016). Indeed, as the Kalahari experienced drought conditions during the two years of this study,

such negative effects of environmental stress on prolactin release might also explain why the plasma

prolactin concentrations detected here were lower than those generally reported in studies of other

passerines engaged in care (e.g. Vleck et al. 1991, Schoech et al. 1996, Khan et al. 2001, Duckworth

et al. 2003, Ouyang et al. 2011).

While experimental tests of causality are needed (see above), our findings are broadly consistent with

the hypothesis that pre-existing mechanisms that regulated parental care in ancestral bi-parental

species were co-opted for the regulation of cooperative helping behavior on the evolution of cooperative

breeding. The possibility that parenting and cooperative helping are indeed regulated by a common

mechanism has important evolutionary implications. Explanations for the evolution, maintenance and

optimization of cooperative behavior typically focus on the roles of the fitness benefits and costs of

cooperation *per se*. However, if cooperation and parenting *are* regulated by a common underlying

mechanism it is conceivable that this shared architecture for the regulation of care-giving is shaped as

much by the payoffs from its expression in a parental context as by the payoffs from its expression in a

cooperative helping context. While selection might independently optimize parental and helper caring

strategies (e.g. via the evolution of an entirely context-dependent caring strategy), it is conceivable that

mechanistic constraints may preclude their independent optimization. For example, genetic variants that

modified a bird's sensitivity to begging could conceivably impact both its parental and cooperative care,

yielding scope for intra-locus genetic conflict to constrain the independent optimization of both parental

and cooperative care (Pennell et al. 2018, see also the conceptual parallels with sexual conflict: Stewart

et al. 2010, Pennell and Morrow 2013). Where this is the case, attempts to understand the evolutionary

origins, maintenance and optimization of cooperative behavior may require attention to the extent to

which genetic correlations exist between parental and cooperative behavior. Notably, our findings

suggest that helping behavior in sparrow-weaver societies is not maintained by selection solely because

a genetic correlation with parenting has precluded the evolution of 'non-helping' (see Brown and Vleck

1998 for a similar debate), because a context-dependent helping strategy appears to have evolved.

Subordinates routinely help while within their natal groups but typically cease to do so as immigrants

(when their low relatedness to the brood would markedly reduce their indirect fitness payoff from helping;

Harrison et al. 2013a).

Conclusion

Our findings lend new support to the hypotheses that helping behavior in cooperatively breeding

societies has shared mechanistic underpinnings with parental care, and that prolactin plays a key role

in this pathway (see also Vleck et al., 1991; Schoech et al., 1996; Khan et al., 2001; Carlson et al.,

2006). Our findings and their complexity highlight the need for experimental studies to investigate both

the causality and nature of the relationship between prolactin and provisioning in this species, in both

parental and cooperative contexts. Our findings also highlight that attempts to understand the evolution

of cooperative helping may benefit from attention to the possibility of genetic constraints on the

independent optimization of cooperation and parenting. Our study has implications too for the growing

interest in the mechanistic origins of consistent individual differences in cooperative behavior

(Sanderson et al. 2015, Dantzer et al. 2019). Specifically, our findings highlight that such differences

could arise from consistent individual differences within the pathway by which prolactin acts (e.g. via

differences in prolactin secretion and/or reception; Ohkubo et al. 1998, Zhou et al. 1996).

ACKNOWLEDGMENTS

We thank all contributors to our long-term database, and Darren O'Connell for support with data

collection. We also thank E.O. and Son and all at Tswalu Kalahari Reserve for access to our field site

and exceptional logistical support (with particular gratitude to Cornelia le Roux and Dylan Smith), and

the Northern Cape Conservation Authority for granting permission to conduct our research. We are

grateful to D. L. Cram for assistance in training in bird capture and blood sampling, J.L. McDonald and

K.J. Walker for useful comments on an earlier draft, and to P. Capilla-Lasheras for statistical advice.

L.A.W. was funded by a NERC PhD studentship (NE/J500185/1). J.E.Y. was funded by the BBSRC

(BB/H022716/1). S.L.M. was funded by Roslin Institute Strategic Grant funding from the BBSRC

(BB/P013759/1). A.J.Y. was funded by BBSRC David Phillips and NERC Blue Skies Research

Fellowships (BB/H022716/1 and NE/E013481/1). We dedicate this manuscript to our co-author Peter

Sharp (P.J.S.), whose openness and interest in this project were key to its inception, and whose wider

contributions to avian endocrinology we celebrate. Any errors are our own, as he did not have a chance

to comment on the final manuscript.

REFERENCES

Angelier, F., C. Barbraud, H. Lormee, F. Prud'homme, and O. Chastel. 2006. Kidnapping of chicks in
emperor penguins: a hormonal by-product? Journal of Experimental Biology **209**:1413-1420.

Angelier, F., and O. Chastel. 2009. Stress, prolactin and parental investment in birds: A review.
General and Comparative Endocrinology **163**:142-148.

Angelier, F., J. C. Wingfield, S. Tartu, and O. Chastel. 2016. Does prolactin mediate parental and life-
history decisions in response to environmental conditions in birds? A review. Horm Behav
77:18-29.

Arnold, T. W. 2010. Uninformative parameters and model selection using Akaike's Information
Criterion. Journal of Wildlife Management **74**:1175-1178.

Badyaev, A., and R. A. Duckworth. 2005. Evolution of plasticity in hormonally-integrated parental
tactcis: An example with the house finch. Pages 375-386 *in* A. Dawson and P. J. Sharp,
editors. Functional Avian Endocrinology. Narosa Publishing House, New Delhi, India.

Bates, D., M. Machler, B. M. Bolker, and S. C. Walker. 2015. Fitting Linear Mixed-Effects Models
Using lme4. Journal of Statistical Software **67**:1-48.

Bender, N., M. Taborsky, and D. M. Power. 2008. The role of prolactin in the regulation of brood care
in the cooperatively breeding fish Neolamprologus pulcher. J Exp Zool A Ecol Genet Physiol
309:515-524.

Boos, M., C. Zimmer, A. Carriere, J. P. Robin, and O. Petit. 2007. Post-hatching parental care
behaviour and hormonal status in a precocial bird. Behavioural Processes **76**:206-214.

Brown, J. L., and C. M. Vleck. 1998. Prolactin and helping in birds: has natural selection strengthened
helping behavior? Behavioral Ecology **9**:541-545.

Buntin, J. D. 1996. Neural and hormonal control of parental behavior in birds. Advances in the Study
of Behavior **25**:161-213.

Burnham, K. P., and D. R. Anderson. 2002. Model Selection and Multimodel Inference. 2 edition.
Springer-Verlag New York, New York.
Capilla-Lasheras, P. 2019. The evolution of cooperation in an arid-zone bird: bet-hedging, plasticity
and constraints. University of Exeter, University of Exeter.
Capilla-Lasheras, P., E. M. Wood, X. Harrison, A. J. Wilson, and A. J. Young. 2021. Altruistic bet-
hedging and the evolution of cooperation in a Kalahari bird. Science Advances **7**:eabe8980.
Carlson, A. A., M. B. Manser, A. J. Young, A. F. Russell, N. R. Jordan, A. S. McNeilly, and T. Clutton-
Brock. 2006a. Cortisol levels are positively associated with pup-feeding rates in male
meerkats. Proceedings of the Royal Society of London Series B-Biological Sciences **273**:571-
577.
Carlson, A. A., L. Nicol, A. J. Young, A. F. Parlow, and A. S. McNeilly. 2003. Radioimmunoassay of
prolactin for the meerkat (Suricata suricatta), a cooperatively breeding carnivore. General and
Comparative Endocrinology **130**:148-156.
Carlson, A. A., A. F. Russell, A. J. Young, N. R. Jordan, A. S. McNeilly, A. F. Parlow, and T. Clutton-
Brock. 2006b. Elevated prolactin levels immediately precede decisions to babysit by male
meerkat helpers. Hormones and Behavior **50**:94-100.
Chastel, O., A. Lacroix, H. Weimerskirch, and G. W. Gabrielsen. 2005. Modulation of prolactin but not
corticosterone responses to stress in relation to parental effort in a long-lived bird. Hormones
and Behavior **47**:459-466.
Chastel, O., and H. Lormee. 2002. Patterns of prolactin secretion in relation to incubation failure in a
tropical seabird, the red-footed booby. Condor **104**:873-876.
Cockburn, A. 1998. Evolution of helping behavior in cooperatively breeding birds. Annual Review of
Ecology and Systematics **29**:141-177.
Cornwallis, C. K., S. A. West, K. E. Davis, and A. S. Griffin. 2010. Promiscuity and the evolutionary
transition to complex societies. Nature **466**:969-U991.
Cram, D. L., J. D. Blount, and A. J. Young. 2015a. The oxidative costs of reproduction are group-size
dependent in a wild cooperative breeder. Proceedings of the Royal Society B-Biological
Sciences **282**.
Cram, D. L., J. D. Blount, and A. J. Young. 2015b. Oxidative status and social dominance in a wild
cooperative breeder. Functional Ecology **29**:229-238.
Criscuolo, F., O. Chastel, F. Bertile, G. W. Gabrielsen, Y. Le Maho, and T. Raclot. 2005.
Corticosterone alone does not trigger a short term behavioural shift in incubating female
common eiders Somateria mollissima, but does modify long term reproductive success.
Journal of Avian Biology **36**:306-312.
Dantzer, B., C. Dubuc, I. B. Goncalves, D. L. Cram, N. C. Bennett, A. Ganswindt, M. Heistermann, C.
Duncan, D. Gaynor, and T. H. Clutton-Brock. 2019. The development of individual differences

in cooperative behaviour: maternal glucocorticoid hormones alter helping behaviour of
offspring in wild meerkats. Philos Trans R Soc Lond B Biol Sci **374**:20180117.
Dantzer, B., I. B. Goncalves, H. C. Spence-Jones, N. C. Bennett, M. Heistermann, A. Ganswindt, C.
Dubuc, D. Gaynor, M. B. Manser, and T. H. Clutton-Brock. 2017. The influence of stress
hormones and aggression on cooperative behaviour in subordinate meerkats. Proc Biol Sci
284.
Delehanty, D. J., L. W. Oring, A. J. Fivizzani, and M. E. ElHalawani. 1997. Circulating prolactin of
incubating male Wilson's Phalaropes corresponds to clutch size and environmental stress.
Condor **99**:397-405.
Dickinson, J. L., and B. J. Hatchwell. 2004. Fitness consequences of helping. Pages 48-66 *in* W. D.
Koenig and J. L. Dickinson, editors. Ecology and Evolution of Cooperative Breeding in Birds.
Cambridge University Press, Cambridge.
Duckworth, R. A., A. V. Badyaev, and A. F. Parlow. 2003. Elaborately ornamented males avoid costly
parental care in the house finch (Carpodacus mexicanus): a proximate perspective. Behavioral
Ecology and Sociobiology **55**:176-183.
English, S., S. Nakagawa, and T. H. Clutton-Brock. 2010. Consistent individual differences in
cooperative behaviour in meerkats (Suricata suricatta). Journal of Evolutionary Biology
23:1597-1604.
Hall, M. R. 1987. External stimuli affecting incubation behavior and prolactin secretion in the duck
(Anas platyrhynchos). Horm Behav **21**:269-287.
Harrison, X., J. York, D. Cram, M. Hares, and A. Young. 2013a. Complete reproductive skew within
white-browed sparrow weaver groups despite outbreeding opportunities for subordinates of
both sexes. Behavioral Ecology and Sociobiology **67**:1915-1929.
Harrison, X. A., J. E. York, D. L. Cram, and A. J. Young. 2013b. Extra-group mating increases
inbreeding risk in a cooperatively breeding bird. Molecular Ecology **22**:5700-5715.
Harrison, X. A., J. E. York, and A. J. Young. 2014. Population genetic structure and direct
observations reveal sex-reversed patterns of dispersal in a cooperative bird. Molecular Ecology
23:5740-5755.
Houslay, T. M., R. L. Earley, A. J. Young, and A. J. Wilson. 2019. Habituation and individual variation
in the endocrine stress response in the Trinidadian guppy (Poecilia reticulata). Gen Comp
Endocrinol **270**:113-122.
Khan, M. Z., F. M. A. McNabb, J. R. Walters, and P. J. Sharp. 2001. Patterns of testosterone and
prolactin concentrations and reproductive behavior of helpers and breeders in the
cooperatively breeding red-cockaded woodpecker (Picoides borealis). Hormones and Behavior
40:1-13.

Koenig, W. D., and J. L. Dickinson, editors. 2004. Ecology and Evolution of Cooperative Breeding in
Birds. Cambridge University Press, Cambridge.
Koenig, W. D., and J. L. Dickinson, editors. 2016. Cooperative breeding in vertebrates: studies of
ecology, evolution and behavior. Cambridge University Press.
Krause, J. S., S. L. Meddle, and J. C. Wingfield. 2015. The Effects of Acute Restraint Stress on
Plasma Levels of Prolactin and Corticosterone across Life-History Stages in a Short-Lived
Bird: Gambel's White-Crowned Sparrow (Zonotrichia leucophrys gambelii). Physiol Biochem
Zool **88**:589-598.
Leitner, S., P. Mundy, and C. Voigt. 2009. Morphometrics of White-browed Sparrow-Weavers
Plocepasser mahali in south-western Zimbabwe. Ostrich **80**:99-102.
Lewis, D. M. 1982. Cooperative Breeding in a Population of White-Browed Sparrow Weavers
Plocepasser-Mahali. Ibis **124**:511-522.
Lukas, D., and T. Clutton-Brock. 2012. Cooperative breeding and monogamy in mammalian societies.
Proceedings of the Royal Society B-Biological Sciences **279**:2151-2156.
Martin-Taylor, C. M. 2018. The impacts of out-group threats on within- and between-group behaviours
in the cooperatively breeding white-browed sparrow-weaver. University of Exeter.
Ohkubo, T., M. Tanaka, K. Nakashima, and P. J. Sharp. 1998. Relationship between prolactin
receptor mRNA in the anterior pituitary gland and hypothalamus and reproductive state in male
and female bantams (Gallus domesticus). General and Comparative Endocrinology **111**:167-
176.
Ouyang, J. Q., P. J. Sharp, A. Dawson, M. Quetting, and M. Hau. 2011. Hormone levels predict
individual differences in reproductive success in a passerine bird. Proceedings of the Royal
Society B-Biological Sciences **278**:2537-2545.
Pennell, T. M., L. Holman, E. H. Morrow, and J. Field. 2018. Building a new research framework for
social evolution: intralocus caste antagonism. Biological Reviews **93**:1251-1268.
Pennell, T. M., and E. H. Morrow. 2013. Two sexes, one genome: the evolutionary dynamics of
intralocus sexual conflict. Ecology and Evolution **3**:1819-1834.
Richards, S. A. 2008. Dealing with overdispersed count data in applied ecology. Journal of Applied
Ecology **45**:218-227.
Richards, S. A., M. J. Whittingham, and P. A. Stephens. 2011. Model selection and model averaging
in behavioural ecology: the utility of the IT-AIC framework. Behavioral Ecology and
Sociobiology **65**:77-89.
Riechert, J., P. H. Becker, and O. Chastel. 2014. Predicting reproductive success from hormone
concentrations in the common tern (Sterna hirundo) while considering food abundance.
Oecologia **176**:715-727.

Sanderson, J. L., I. Stott, A. J. Young, E. I. K. Vitikainen, S. J. Hodge, and M. A. Cant. 2015. The
856   origins of consistent individual differences in cooperation in wild banded mongooses, Mungos
857   mungo. Animal Behaviour **107**:193-200.

858   Sanderson, J. L., A. J. Young, S. J. Hodge, S. Kyabulima, S. L. Walker, and M. A. Cant. 2014.
859   Hormonal mediation of a carry-over effect in a wild cooperative mammal. Functional Ecology
860   **28**:1377-1386.

861   Schoech, S. J., E. D. Ketterson, V. Nolan, P. J. Sharp, and J. D. Buntin. 1998. The effect of
862   exogenous testosterone on parental behavior, plasma prolactin, and prolactin binding sites in
863   dark-eyed juncos. Hormones and Behavior **34**:1-10.

864   Schoech, S. J., R. L. Mumme, and J. C. Wingfield. 1996. Prolactin and helping behaviour in the
865   cooperatively breeding Florida scrub-jay, *Aphelocoma coerulescens*. Animal Behaviour
866   **52**:445-456.

867   Schoech, S. J., S. J. Reynolds, and R. K. Boughton. 2004. Endocrinology. Pages 128-141 *in* W. D.
868   Koenig and J. L. Dickinson, editors. Ecology and Evolution of Cooperative Breeding in Birds.
869   Cambridge University Press, Cambridge.

870   Sharp, P. J., A. Dawson, and R. W. Lea. 1998. Control of luteinizing hormone and prolactin secretion
871   in birds. Comparative Biochemistry and Physiology C-Pharmacology Toxicology &
872   Endocrinology **119**:275-282.

873   Smiley, K. O., and E. Adkins-Regan. 2018. Lowering prolactin reduces post-hatch parental care in
874   male and female zebra finches (Taeniopygia guttata). Horm Behav **98**:103-114.

875   Soares, M. C., R. Bshary, L. Fusani, W. Goymann, M. Hau, K. Hirschenhauser, and R. F. Oliveira.
876   2010. Hormonal mechanisms of cooperative behaviour. Philosophical Transactions of the
877   Royal Society B-Biological Sciences **365**:2737-2750.

878   Sockman, K. W., H. Schwabl, and P. J. Sharp. 2000. The role of prolactin in the regulation of clutch
879   size and onset of incubation behavior in the American kestrel. Hormones and Behavior **38**:168-
880   176.

881   Solomon, N. G., and J. A. French, editors. 1997. Cooperative Breeding in Mammals. Cambridge
882   University Press, Cambridge.

883   Stewart, A. D., A. Pischedda, and W. R. Rice. 2010. Resolving Intralocus Sexual Conflict: Genetic
884   Mechanisms and Time Frame. Journal of Heredity **101**:S94-S99.

885   Vleck, C. M., N. A. Mays, J. W. Dawson, and A. R. Goldsmith. 1991. Hormonal Correlates of Parental
886   and Helping-Behavior in Cooperatively Breeding Harris Hawks (Parabuteo-Unicinctus). Auk
887   **108**:638-648.

888   Voigt, C., S. Leitner, and M. Gahr. 2007. Socially induced brain differentiation in a cooperatively
889   breeding songbird. Proceedings of the Royal Society of London Series B-Biological Sciences
890   **274**:2645-2652.

Walker, L. A. 2016. The evolution and regulation of cooperation in the wild. University of Exeter.

Walker, L. A., J. E. York, and A. J. Young. 2016. Sexually selected sentinels? Evidence of a role for
intrasexual competition in sentinel behavior. Behavioral Ecology **27**:1461-1470.

Wingfield, J. C., and D. M. Lewis. 1993. Hormonal and Behavioral-Responses to Simulated Territorial
Intrusion in the Cooperatively Breeding White-Browed Sparrow Weaver, Plocepasser-Mahali.
Animal Behaviour **45**:1-11.

Wood, E. M., P. Capilla-Lasheras, D. L. Cram, L. A. Walker, J. E. York, A. Lange, P. B. Hamilton, C.
R. Tyler, and A. J. Young. 2021. Social dominance and rainfall predict telomere dynamics in a
cooperative arid-zone bird. Mol Ecol.

York, J. E., A. N. Radford, B. de Vries, T. G. Groothuis, and A. J. Young. 2016. Dominance-related
seasonal song production is unrelated to circulating testosterone in a subtropical songbird.
General and Comparative Endocrinology **233**:43-52.

York, J. E., K. J. Wells, and A. J. Young. 2019. Dominance-related contributions to collective territory
defence are adjusted according to the threat. Animal Behaviour **158**:25-34.

Young, A. J., A. A. Carlson, and T. H. Clutton-Brock. 2005. Trade-offs between extra-territorial
prospecting and helping in a cooperative mammal. Animal Behaviour **70**:829-837.

Zhou, J. F., D. Zadworny, D. Guemene, and U. Kuhnlein. 1996. Molecular cloning, tissue distribution,
and expression of the prolactin receptor during various reproductive states in Meleagris
gallopavo. Biology of Reproduction **55**:1081-1090.

Ziegler, T. E. 2000. Hormones associated with non-maternal infant care: A review of mammalian and
avian studies. Folia Primatologica **71**:6-21.

914   Table A1. Top model sets prior to applying the model nesting rule when modelling the

915   Provisioning Behavior and Prolactin Levels of DOMINANTS

Model selection of the factors affecting the (i) nestling provisioning rates, (ii) mean provisioning visit

durations and (iii) prolactin levels of dominant females ('Dom Fem') and dominant males ('Dom Male')

engaged in parental care. This table shows all models within <6 ΔAIC of the top model, *prior to* applying

the model nesting rule (Richards et al., 2011; see methods). Table 1 in the main paper presents the

subset of these models that were retained *after* applying the model nesting rule (which are flagged here

with 'YES' in the 'Retained' column). The grey lines indicate the response term for the set of models

beneath it. The presence of a continuous variable within a focal model is indicated by the presence of

its effect size estimate, while the presence of factorial variable is indicated with a '+'. The 'Year' effect

is the change from 2013 to 2014. 'Int' = Intercept. 'df' = degrees of freedom.

Model Number	Int	Bird Class (Sub Natal vs Sub Imm)	Brood Size	Group Size	Year	Sunset to Capture Lag	Capture to Bleed lag	df	AICc	ΔAICc	Retain
Provisioning Rate (feeds / hr)											
4	-0.36	+	1.44			N/A	N/A	6	205.78	0.00	YES
12	-0.29	+	1.08		+	N/A	N/A	7	206.06	0.28	
11	1.21	+			+	N/A	N/A	6	207.70	1.92	YES
8	-0.19	+	1.42	-0.05		N/A	N/A	7	208.54	2.76	
16	-0.46	+	1.08	0.05	+	N/A	N/A	8	208.97	3.19	
15	1.14	+		0.02	+	N/A	N/A	7	210.49	4.71	
3	2.06	+				N/A	N/A	5	211.76	5.98	YES
Mean Provisioning Visit Duration (minutes)											
3	0.58	+				N/A	N/A	5	139.66	0.00	YES
4	1.08	+	-0.30			N/A	N/A	6	141.08	1.42	
11	0.71	+			+	N/A	N/A	6	141.79	2.13	
7	0.37	+		0.07		N/A	N/A	6	142.11	2.45	
12	1.07	+	-0.26		+	N/A	N/A	7	143.81	4.15	
8	0.93	+	-0.28	0.04		N/A	N/A	7	143.84	4.18	

15	0.57	+		0.04	+	N/A	N/A	7	144.55	4.89	
Plasma Prolactin (ng/ml)											
5	1.98	+						5	149.62	0.00	YES
7	1.32	+	0.39					6	150.56	0.94	
13	1.31	+		0.23				6	150.68	1.07	
15	0.50	+	0.43	0.25				7	151.36	1.74	
37	1.77	+			+			6	151.41	1.79	
45	0.82	+		0.29	+			7	151.68	2.07	
6	2.51	+					-0.0028	6	151.79	2.17	
21	2.03	+				-0.0005		6	152.24	2.62	
14	1.81	+		0.22			-0.0025	7	153.06	3.44	
8	1.76	+	0.37				-0.0021	7	153.07	3.45	
39	1.32	+	0.33		+			7	153.17	3.55	
23	1.32	+	0.39			-0.0001		7	153.35	3.73	
29	1.29	+		0.23		0.0001		7	153.48	3.86	
47	0.47	+	0.30	0.28	+			8	153.76	4.14	
38	2.17	+			+		-0.0020	7	153.97	4.35	
53	1.86	+			+	-0.0013		7	153.99	4.37	
16	0.89	+	0.41	0.25			-0.0017	8	154.11	4.49	
31	0.42	+	0.44	0.26		0.0006		8	154.26	4.64	
61	0.90	+		0.29	+	-0.0009		8	154.53	4.91	
46	1.09	+		0.29	+		-0.0012	8	154.54	4.92	
22	2.54	+				-0.0003	-0.0027	7	154.57	4.95	

Table A2. Top model sets prior to applying the model nesting rule when modelling the

Provisioning Behavior and Prolactin Levels of SUBORDINATES

Model selection of the factors affecting the (i) nestling provisioning rates and (ii) prolactin levels of natal

subordinates ('Sub Nat') and immigrant subordinates ('Sub Imm') engaged in cooperative helping

behavior. This table shows all models within <6 ΔAIC of the top model, *prior to* applying the model

nesting rule (Richards et al., 2011; see methods). Table 2 in the main paper presents the subset of

these models that were retained *after* applying the model nesting rule (which are flagged here with 'YES'

in the 'Retained' column). The grey lines indicate the response term for the set of models beneath it.

The presence of a continuous variable within a focal model is indicated by the presence of its effect size

estimate, while the presence of factorial variable is indicated with a '+'. The 'Year' effect is the change

from 2013 to 2014. 'Int' = Intercept. 'df' = degrees of freedom.

Model Number	Int	Bird Class (Sub Nat vs Sub Imm)	Brood Size	Group Size	Year	Sunset to Capture Lag	Capture to Bleed lag	df	AICc	ΔAICc	Retain
Provisioning Rate (feeds / hr)											
4	-1.39	+	0.92			N/A	N/A	5	66.64	0.00	YES
12	-1.40	+	1.12		+	N/A	N/A	6	67.92	1.28	
3	0.12	+				N/A	N/A	4	68.77	2.13	YES
8	-0.65	+	0.86	-0.18		N/A	N/A	6	69.68	3.04	
16	-0.30	+	1.06	-0.27	+	N/A	N/A	7	70.56	3.93	
7	1.22	+		-0.31		N/A	N/A	5	70.73	4.09	
11	0.23	+			+	N/A	N/A	5	71.79	5.15	
Plasma Prolactin (ng / ml)											
45	2.20	+		-0.41	+			6	57.13	0.00	YES
13	2.82	+		-0.51				5	57.15	0.02	YES
29	2.47	+		-0.48		0.0038		6	59.10	1.97	
46	1.15	+		-0.35	+		0.0038	7	59.22	2.09	
15	2.62	+	0.08	-0.50				6	60.62	3.49	
14	2.83	+		-0.51			0.0000	6	60.76	3.62	
61	2.11	+		-0.40	+	0.0019		7	60.78	3.65	
47	2.27	+	-0.03	-0.41	+			7	61.17	4.04	
33	1.02				+			4	61.28	4.15	YES

9	2.74			-0.34				4	61.42	4.29	YES
25	2.22			-0.32		0.0057		5	61.62	4.49	
38	-0.86	+			+		0.0068	6	61.68	4.55	
41	2.09			-0.23	+			5	61.76	4.63	
37	0.64	+			+			5	62.00	4.87	
34	0.06				+		0.0046	5	62.57	5.43	
30	2.37	+		-0.48		0.0039	0.0004	7	63.12	5.99	

940