1

Tyr is Responsible for the _Cctq1a_ QTL and Links Developmental Environment to Central

Corneal Thickness Determination

Kacie J. Meyer[1§], Demelza R. Larson[1§], S. Scott Whitmore[2], Carly J. van der Heide[1,2], Adam

Hedberg-Buenz[1,3], Laura M. Dutca[3], Swanand Koli[4], Nicholas Pomernackas[1], Hannah E.

Mercer[1], Maurisa N. Mansaray[1], William J. Paradee[5], Kai Wang[6], K. Saidas Nair[4,7], Todd E.

Scheetz[2], and Michael G. Anderson[1,2,3]*

[1]Department of Molecular Physiology and Biophysics, University of Iowa, Iowa City, Iowa;

[2]Department of Ophthalmology and Visual Sciences, University of Iowa, Iowa City, Iowa;

[3]Center for the Prevention and Treatment of Visual Loss, Iowa City VA Health Care System,

Iowa City, Iowa; [4]Department of Ophthalmology, University of California, San Francisco,

California; [5]Genome Editing Core Facility, University of Iowa, Iowa City, Iowa; [6]Department of

Biostatistics, University of Iowa, Iowa City, Iowa; [7]Department of Anatomy, University of

California, San Francisco, California

[§]Equal contribution

*Corresponding author: Dr. Michael G. Anderson, Department of Molecular Physiology and

Biophysics, 3111 Medical Education and Research Facility, 375 Newton Road, Iowa City, IA

52242, (319) 335-7839 (telephone), (319) 335-7330 (FAX), michael-g-anderson@uiowa.edu

Abstract

Central corneal thickness is a quantitative trait with important associations to human health. In a phenotype-driven approach studying corneal thickness of congenic derivatives of C57BLKS/J and SJL/J mice, the critical region for a quantitative trait locus influencing corneal thickness, *Cctq1a*, was delimited to a 10-gene interval. Exome sequencing, RNAseq, and studying independent mutations eliminated multiple candidate genes and confirmed one. Though the causative gene, *Tyr*, has no obvious direct function in the transparent cornea, studies with multiple alleles on matched genetic backgrounds, both in isolation and genetic complementation crosses, confirmed allelism of *Tyr-Cctq1a*; albino mice lacking *Tyr* function had thin corneas. Albino mice also had increased axial length. Because albinism exposes eyes to increased light, the effect of dark-rearing was tested and found to rescue central corneal thickness. In sum, the results point to an epiphenomenon; developmental light exposure interacts with genotype as an important determinate of adult corneal thickness.

Introduction

Central corneal thickness has important associations with ocular disease, but its natural

determining factors remain predominately elusive. The cornea consists of three cellular layers

(an outermost epithelium, middle stroma, and innermost endothelium) separated by two thinner

basement membranes. The combined central thickness of these layers (central corneal

thickness, CCT) increases rapidly through infancy and early childhood, reaches adult values in

pre-teen ages, and remains relatively stable thereafter, with eventual modest age-related

thinning[1-3, 4]. For largely unknown reasons, average CCT can vary by dozens of microns

between ethnicities[4,5]. Thin CCT is associated with several corneal diseases, such as corneal

dystrophy, brittle cornea syndrome, keratoconus, and cornea plana; diseases of connective

tissue, such as Marfan syndrome, Ehlers-Danlos syndrome, Loeys-Dietz syndrome, and

osteogenesis imperfecta; and at least two diseases in which the nature of, and/or reason for, the

association is unclear, including myopia and primary open angle glaucoma[6].

CCT is a highly heritable trait[7,8], leading to many genetic studies. Variants influencing CCT have

been identified from familial studies of Mendelian syndromes and GWAS of various large

populations[9-12]. From these studies, some themes have begun to emerge. For example, some

of the same CCT loci are identified in both multigenic and Mendelian disease studies[10,13,14]. It

has also been common to identify variants related to collagen matrix integrity[10,12,14,15]. However,

it is also clear that much remains unknown. Among the known associations, most known SNPs

occur in non-coding regions, and the nearest genes typically have no obvious link to known

structural components of the cornea[15]. It is also clear that many important genes remain to be

discovered. Known SNPs give rise to a SNP-based heritability estimate of 42.5% and account

for only 14.2% of the CCT variance[9].

Here, an unique approach complementary to others is undertaken using inbred mouse strains to

identify quantitative trait loci (QTL) that influence CCT[16,17]. Similar to humans, there is natural

variation of CCT among inbred mouse strains[18]. Previous work using a quantitative approach

with intercrosses between two such strains (C57BLKS/J [KS] mice with thin corneas and SJL/J

[SJL] mice with thick corneas) identified the first CCT QTL, *Central corneal thickness QTL 1*

(*Cctq1*) on mouse chromosome 7[16]. Here, *Cctq1* was resolved into two closely linked regions,

Cctq1a and *Cctq1b*, which each influence CCT. Through multiple genetic approaches, a

mutation in the tyrosinase gene (*Tyr*) is identified as causative of the *Cctq1a* phenotype, which

appears to influence CCT via an epiphenomenon dependent on developmental light exposure.

Results

Cctq1 **contains two adjacent interacting QTL**

The original 95% Bayesian credible interval of *Cctq1* spanned a 38.3 cM region on chromosome

7 (34.1 cM – 72.4 cM)[16]. To reduce this interval, recombination mapping was used with 92

recombinant N4 intercross mice. However, the interval was originally recalcitrant to division—

more than one interval conferred the increased CCT phenotype. This suggested that there was

too much genetic heterogeneity for this trait at the N4 generation, potentially including the

presence of more than one CCT-regulating gene within or near *Cctq1*. To address these

possibilities, the original F2 dataset was subjected to additional evaluations while backcrossing

of the congenic mice to the N10 generation was continued.

The original analysis of (KS X SJL) F2 mice was based upon a significance threshold

determined empirically by stratified permutation testing with 1000 permutations[16,19], and did not

identify any loci that significantly interacted with *Cctq1*. Prompted by the recombination

mapping, the dataset was re-examined by performing a pairwise scan of the markers on

chromosome 7 using scantwo analysis with R/qtl[19,20]. Of the possible interactions, the markers

that produced the highest LOD scores were *D7Mit31* and *rs13479535* (Figure 1A; full LOD

score = 10.33, interactive LOD score = 5.05). *D7Mit31* lies within the 95% Bayes credible

interval of *Cctq1*; *rs13479535* is 2 cM distal to the end of the interval at 74.3 cM. The putative

linked loci were subsequently subjected to multiple regression analysis in which each locus and

the interaction component were sequentially dropped from the 2-QTL model. This analysis

indicated that both loci and their interaction had significant contributions to the model

(Supplementary Data 1). The QTL at *D7Mit31* was responsible for 31% of the phenotypic

variability while the QTL at *rs13479535* accounted for another 24% of the variability. These data

indicate that both loci are true QTL. *Cctq1* was thus resolved into two QTL, *Cctq1a* (95% Bayes

credible interval: *D7Mit318–D7Mit220*, spanning 49.0 cM, peak at *D7Mit31*), and *Cctq1b* (95%

Bayes credible interval: *D7Mit105–rs13479545*, spanning 74.3 cM, peak at *rs13479535*; Figure

1B).

To reduce genetic heterogeneity, N4 mice were further backcrossed onto the KS background to

the N10 generation. Because the congenic interval was relatively large, a panel of six markers

was used at each generation of backcrossing to keep the interval intact. At generation N10,

congenic mice were intercrossed, recombination within the interval was allowed, and mice with

all nine genetic combinations of *Cctq1a* and *Cctq1b* were phenotyped for CCT (Table 1).

Congenic control mice (KS.SJL-*Cctq1a^{KS},Cctq1b^{KS}*) had a CCT indistinguishable from inbred

KS mice (94.8± 2.4 µm vs. 94.5 ± 3.2 µm, respectively; one-way ANOVA with Tukey post-test;

Table 1). Congenic mice with SJL genotypes at *Cctq1a* (*i.e.*, KS.SJL-*Cctq1a^{SJL}*) had significantly

thinner corneas than inbred KS mice (87.9 ± 3.8 µm; *n* = 39; *p* < 0.001; Student's two-tailed *t*-

test) independent of the genotype at *Cctq1b* (*n* = 13 and *p* < 0.05 for each of three genotypes at

Cctq1b; one-way ANOVA with Tukey post-test; Table 1; Supplementary Data 2). The difference

in thickness mediated by *Cctq1a* is predominantly due to the thickness of the stroma (Δ 9.6 µm,

p < 0.001; Student's two-tailed *t*-test), though there is a small (Δ 1.2 µm) and marginally

significant (*p* = 0.022 Student's two-tailed *t*-test) decrease in thickness of the epithelium as well.

Congenic mice with KS genotypes at *Cctq1a* and SJL genotypes at *Cctq1b* (i.e., KS.SJL-

Cctq1a^{KS},Cctq1b^{SJL}) also had significantly thinner corneas (90.8 ± 2.6 µm; *n* = 13 mice; *p* < 0.05;

one way ANOVA with Tukey post-test) than inbred KS mice (Table 1, Supplementary Data 2).

No other genetic combinations caused significant changes in CCT compared to inbred KS

controls. As predicted from the analysis of the original F2 Dataset with R/qtl, these data with

congenic mice independently support that both *Cctq1a* and *Cctq1b* are true QTL capable of

altering the phenotypic variability of CCT on a uniform genetic background.

Mapping of *Cctq1a* using recombination mapping and sub-congenics

Because of its larger effect, initial efforts were focused on fine-scale mapping for *Cctq1a*. To

identify the gene underlying *Cctq1a*, *Cctq1a*-recombinant N10F2 mice, with KS genotypes at

Cctq1b, were used to narrow the critical region. From this recombination analysis, the gene

underlying *Cctq1a* was deduced to be between SNP markers *rs108403472* at 48.51 cM and

rs6247100 at 50.26 cM. Simultaneously, sub-congenic mice were created by continued

backcrossing of the KS.SJL-*Cctq1a^{HET};Cctq1b^{KS}* N10 mice. Eyes of N10 congenic mice were

overtly healthy, differing only in pigmentation between genotypes (Supplementary Data 2). At

N12, *Cctq1a* was physically reduced to a 15.9 cM region (KS.SJL-*Cctq1a(15.9cM)*) flanked by

D7mit347 and *D7mit321* and characterized by an association of SJL homozygosity with

decreased CCT (87.6 ± 2.3 µm vs. 76.8 ± 2.4 µm, *n* = 15 mice per genotype, *p* < 0.001,

Student's two-tailed *t*-test). At N15, *Cctq1a* was physically reduced to a 9.9 cM region (KS.SJL-

Cctq1a(9.9cM)) flanked by *rs3672782* and *D7mit321*; again characterized by an association of

SJL homozygosity with decreased CCT (91.6 ± 2.2 µm vs. 80.7 ± 2.9 µm, *n* = 7 mice per

genotype, *p* < 0.001, Student's two-tailed *t*-test). One N15F2 mouse harbored a recombination

event within the minimal sub-congenic interval. The phenotype of this mouse indicated the gene

underlying *Cctq1a* lies proximal to marker *rs13479393* (Figure 2). Using this recombinant

mouse in a progeny test, additional intercrossing to generation N15F7 confirmed the distal

breakpoint proximal to marker *rs13479393* at 49.65 cM, again characterized by an association

of SJL homozygosity with decreased CCT (85.7 ± 1.6 μm vs. 94.4 ± 3.3 μm, *n* = 8–10 mice per

genotype; $p < 0.001$, Student's two-tailed *t*-test; Figure 3A; Supplementary Data 3–5).

In sum, physical recombination mapping utilizing multiple generations of congenic and sub-

congenic mice conclusively indicated that the gene underlying *Cctq1a* lies on chromosome 7

between markers *rs108403472* at 48.51 cM and *rs13479393* at 49.32 cM, a 0.81 cM region

containing the entirety of eight RefSeq genes (*Vmn2r78*, *Vmn2r79*, *Nox4*, *Tyr*, *Grm5*, *Ctsc*,

Rab38, and *Tmem135*), the 3' portion of one gene (*Vmn2r77*), and the 5' portion of one gene

(*Fzd4*) (Figure 2).

Candidate identification and prioritization

To identify all the possible exonic variants within the *Cctq1a* critical region, whole exome

sequencing was conducted on KS and SJL inbred mice. In the entire exome, 15,261 missense,

frameshift, and splice-site mutations were found between KS and SJL (Supplementary Data 6).

There were six amino acid altering variants within the *Cctq1a* critical region, located within

Vmn2r79 (A223T, L243M, T257I, I265V), *Tyr* (C103S), and *Fzd4* (F27L). Vomeronasal receptor

genes, such as *Vmn2r79*, have an increased rate of coding sequence variants [21] and the four

altered residues between KS and SJL in *Vmn2r79* are poorly conserved. The amino acid

change in *Tyr* from cysteine to serine is the albinism-causing *Tyr^c* allele conferred by the albino

SJL strain. The *Fzd4* amino acid variant is within the signal sequence of the protein; KS mice

conferred the phenylalanine amino acid residue (the same residue as C57BL/6J mice) while

SJL mice conferred the mammalian-conserved leucine residue.

Transcriptional profiling was additionally used to prioritize candidates. Using previously

published microarray data comparing adult corneal RNA expression profiles in KS and SJL

mice[18], *Nox4*, *Ctsc*, *Rab38*, *Tmem135*, and *Fzd4* were all present in the adult cornea; there was

no evidence for adult corneal expression of *Vmn2r77*, *Vmn2r78*, *Vmn2r79*, *Tyr*, or *Grm5*. Of

those expressed, *Nox4, Ctsc, and Fzd4* showed differential expression between the two strains.

Nox4 and *Ctsc* were both down-regulated 2.5-fold in SJL, while *Fzd4* was up-regulated 1.6-fold

in SJL.

Additionally, RNAseq was performed on corneas of 3-week-old KS.SJL-*Cctq1a(15.9cM)*^{SJL}

N12F3 mice. Analysis was focused on three comparisons: 1) KS.SJL-*Cctq1a(15.9cM)*^{SJL} vs. KS

(experimental, identifying genes with altered corneal expression in the congenic interval), 2)

KS.SJL-*Cctq1a(15.9cM)*^{SJL} vs. KS.SJL-*Cctq1a(15.9cM)*^{KS} (experimental, also identifying genes

with altered corneal expression in the congenic interval), and 3) KS.SJL-*Cctq1a(15.9cM)*^{KS} vs.

KS (control, identifying genes in the background of the congenic strain with altered corneal

expression not associated with the congenic interval) (Supplementary Data 7). In each

comparison, genes were first filtered for those with a Q-value ≤ 0.001 and a FPKM ≥ 1 in at

least one of the strains. Gene lists were subsequently compared to one another, identifying 87

genes consistently altered in both experimental comparisons but not in the control comparison

(Supplementary Data 7). Among these 87, only one was localized to the *Cctq1a* critical region,

Ctsc, which was modestly (-0.8 $\log_2$ fold) but consistently and significantly ($p = 5 \times 10^{-5}$; $Q =$

0.0009) down regulated in comparing the SJL allele to the KS allele. Web Gestalt[22] was used for

over-representation analysis comparing the list of 87 differentially expressed genes to a

background list of all genes expressed in the cornea with a FPKM ≥ 1 in any strain. Results of

the analysis indicate a strong signal for several collagen-related categories (fibrillar collagen

trimer, abnormal cutaneous collagen fibril morphology, collagen biosynthesis and modifying

enzymes, collagen degradation, etc.), extracellular-matrix-related categories (extracellular

matrix component, ECM proteoglycans, degradation of the extracellular matrix, etc.) and ocular-

related categories (decreased corneal stroma thickness, abnormal corneal epithelium

morphology, abnormal cornea morphology, and abnormal eye morphology) (Supplementary

Data 8).

Functional tests of lead candidates

Based on candidate prioritization criteria, *Fzd4* and *Ctsc* were initially considered the top

candidates. To test the influence of *Fzd4* on CCT, we tested a strain with a targeted mutation of

Fzd4 (B6;129-*Fzd4^{tm1Nat}*/J) on a segregating B6 and 129 background[23]. *Fzd4^{tm1Nat}* homozygotes

had a chocolate coat color not expected from either background. This observation is meaningful

as the gene responsible for this phenotype, *Rab38*, is physically near *Fzd4* and within the *Cctq1*

critical region, i.e., the strain is likely a double mutant for *Fzd4* (genotype verified) and *Rab38*

(mutation unknown). However, there was no correlation between *Fzd4^{tm1Nat}* genotype and CCT

(*p* = 0.819; one-way ANOVA comparing all three genotypes among littermates; Figure 3B;

Supplementary Data 3–5). In genetic complementation crosses, *Fzd4^{tm1Nat}* mutation

complemented the congenic interval ([*Fzd4^{WT}*/*Cctq1a^{SJL}* F1] vs. [*Fzd4^{HET}*/*Cctq1a^{SJL}* F1]; *p* =

0.528; Student's two-tailed *t*-test; *n* = 4-6 per genotype; Figure 3C; Supplementary Data 3–5).

The complementation cross also highlighted the coat color phenotype associated with the

Fzd4^{tm1Nat} mutation, with *Fzd4^{WT}*/*Cctq1a^{SJL}* F1 mice (*Tyr^{HET}*) having a black coat color with

normally pigmented eyes and *Fzd4^{HET}*/*Cctq1a^{SJL}* F1 mice (*Tyr^{HET}*) having an unmistakable

lightened ("light chocolate") coat color and light brown irides instead of brown (Supplementary

Data 9). Therefore, *Fzd4* was ruled out as causative of the *Cctq1a* phenotype, and *Rab38*

further deprioritized as a candidate.

To test the influence of *Ctsc* on CCT, we imported a strain with a targeted mutation of *Ctsc*

(B6.Cg-*Ctsc*[tm1Ley]) on an N10 congenic B6 background[24]. Because the targeted mutation was

generated on an albino 129 background, and *Ctsc* is physically linked to *Tyr*, *Ctsc*[tm1Ley]

homozygotes are albino, i.e., the strain is a double mutant for *Ctsc* and *Tyr*. This is meaningful

because the SJL/J strain is albino and the *Tyr*[c] mutation is within the KS.SJL-*Cctq1a* congenic

interval. Homozygotes had a statistically significant decrease in CCT compared to littermate

controls ($p < 0.002$ for albino *Ctsc*[tm1Ley] vs. pigmented *Ctsc*[HET]; $p < 0.002$ for albino *Ctsc*[tm1Ley] vs.

pigmented *Ctsc*[WT]; one-way ANOVA with Tukey post-test comparing all three genotypes among

littermates; n=10–12 mice per genotype; Figure 3D; Supplementary Data 3–5). In genetic

complementation crosses, *Ctsc*[tm1Ley] failed to complement *Cctq1a*[SJL] ($p < 0.001$ for pigmented

[*Ctsc*[tm1Ley]/*Cctq1a*[KS] F1] vs. albino [*Ctsc*[tm1Ley]/*Cctq1a*[SJL] F1]; Student's two-tailed *t*-test; $n = 10–$

11 mice per genotype; Figure 3E; Supplementary Data 3–5). Therefore, *Ctsc or Tyr* were

determined to be causative of the *Cctq1a* phenotype.

To differentiate *Ctsc* and *Tyr* as the causative mutation, independent alleles on a pure B6

background were assessed. To test *Ctsc*, four new mutations predicted to result in null protein

were generated in B6 mice with CRISPR-Cas9 technology (Supplementary Data 10). To test

Tyr, a well-known spontaneous mutation that was commercially available, *Tyr*[c-2J], was

analyzed[25,26]. There was no association between *Ctsc*[KO] genotype and CCT ($p = 0.237$; one-

way ANOVA comparing all three genotypes among littermates; $n = 9–17$ mice per group; Figure

3F; Supplementary Data 3-5). In genetic complementation crosses with the congenic strain, the

Ctsc[tm1Mga] mutation (46bp-deletion in the coding sequence of exon 1 leading to no detectable

CTSC protein; Supplementary Data 10) complemented the congenic phenotype ($p = 0.696$;

Student's two-tailed *t*-test; $n = 8–12$ mice per group; Figure 3G; Supplementary Data 3–5). In

contrast, albino *Tyr*[c-2J] mice had decreased CCT relative to pigmented B6 ($p < 0.001$; Student's

two-tailed *t*-test; $n = 9–10$ mice per group; Figure 3H; Supplementary Data 3–5). In genetic

complementation crosses with the congenic strain, the Tyr^{c-2J} mutation failed to complement the

congenic phenotype ($p < 0.001$ for pigmented [$Cctq1a^{KS}/Tyr^{c-2J}$ F1] vs. albino [$Cctq1a^{SJL}/Tyr^{c-2J}$

F1]; Student's two-tailed t-test; n = 8 mice per genotype; Figure 3I; Supplementary Data 3–5).

Thus, only Tyr mutation was left as a feasible candidate for $Cctq1a$.

CRISPR-Cas9 technology was also used to generate new Tyr mutations on a C57BL/6J

background (Supplementary Data 11). One allele was selected for propagation, Tyr^{tm4Mga}, an

albinism-causing 4bp-deletion in the coding sequence of exon 1 that is predicted to cause a

frameshift leading to a premature stop codon and RNA-mediated decay; i.e., a presumed null

mutation. Tyr^{tm4Mga} mice had a significantly thinner cornea than littermate controls ($p < 0.002$ for

Tyr^{WT} vs. Tyr^{tm4Mga}; $p < 0.002$ for Tyr^{HET} vs. Tyr^{tm4Mga}; one-way ANOVA with Tukey post-test

comparing all three genotypes among littermates; n = 6–10 mice per genotype; Figure 4;

Supplementary Data 5 and 12). Thus, a presumed null allele of Tyr caused the same albinism

and thinning of CCT as found for the c and c-$2J$ alleles.

Mechanism of Tyr function on CCT

In considering the possible mechanism through which Tyr might influence CCT, three

hypotheses were tested:

The first candidate mechanism centered on the role of DOPA, which is a cofactor in the

oxidation of tyrosine by TYR, leading to melanin production[27], and a substrate for tyrosine

hydroxylase (TH), leading to dopamine synthesis. Dopamine is considered a key molecule in

ocular growth[28,29], and mice with a conditional knock-out of Th in the retina have previously

been shown to have decreased CCT[30]. Rationalizing that some DOPA may normally escape

from pigment producing cells to influence CCT via a TH-dependent mechanism, we tested

whether providing supplemental DOPA in the drinking water of albino mice would rescue the

decreased CCT associated with albinism. No statistically significant effect was observed (Figure

5; Supplementary Data 5), discounting the hypothesis that a DOPA deficit was rate-limiting for

CCT determination in *Tyr* mutant mice.

The second candidate mechanism revolves around the possibility of a gene-environment

interaction involving temperature and light. In this series of experiments, mice were reared in

environmental control chambers in combinations of different temperatures and light cycles.

Cohorts included pigmented B6, albino Tyr^{c-2J}, and the temperature sensitive himalayan

mutation (B6.Cg-Tyr^{c-h}/J)[31,32]. At ambient temperatures, Tyr^{c-h} homozygotes are only partially

pigmented on the coolest parts of the body (such as the ears, nose, tail, and eyes) and CCT is

intermediate between B6 and Tyr^{c-2J} mice (Supplementary Data 5, 13, and 14). For CCT of mice

raised at ambient temperature, genetic complementation tests again confirmed the influence of

Tyr-mediated albinism on CCT (p = 0.337 for [pigmented B6 x Tyr^{c-h} F1] vs. [pigmented B6 x

Tyr^{c-2J} F1]; $p < 0.002$ for [pigmented B6 x Tyr^{c-h} F1] vs. [albino Tyr^{c-h} x Tyr^{c-2J} F1]; and $p < 0.002$

for [pigmented B6 x Tyr^{c-2J} F1] vs. [albino Tyr^{c-h} x Tyr^{c-2J} F1]; one-way ANOVA with Tukey post-

test; n = 4–14 mice per group, Supplementary Data 5). At decreased temperatures Tyr^{c-h}

homozygotes can generate pigment more broadly, and at increased temperatures the albinism

is accentuated (Supplementary Data 13 and 14).

Regarding temperature, comparison of B6.Tyr^{c-h} mice, as well as B6 and B6.Tyr^{c-2J} controls,

raised at 10°C vs. 32°C in environmental control chambers with a standard light cycle, showed

that CCT followed pigment status; the thin CCT of hypopigmented Tyr^{c-h} mice reared at

increased temperature (i.e., lower TYR activity) was rescued by rearing the Tyr^{c-h} mice at

decreased temperature (i.e., higher TYR activity; Tyr^{c-h} at 32°C vs. Tyr^{c-h} at 10°C; Δ 10.9 μm; p

< 0.001; n = 13–15 mice per condition; one-way ANOVA with Sidak test; Figure 6;

Supplementary Data 5). In testing the effect of temperature on corneal thickness in controls,

there was a small (Δ 2.8 µm) and nominally significant (p = 0.03) decreased CCT in mice raised

at 10ºC compared to mice raised at 32ºC when considering all C57BL/6J and Tyr^{c-2J} mice

across the experiment (n = 16 mice vs. n = 19 mice, Type II ANOVA) and there was no

interaction between genotype and temperature. In sum, changing environmental temperature

changed CCT of the Tyr^{c-h} mice as predicted.

Regarding light, comparison of B6.Tyr^{c-h} mice, as well as B6 and B6.Tyr^{c-2J} controls, raised at

10°C vs. 32°C in environmental control chambers with dark-rearing of mice from conception to

10–15 weeks of age rescued the thin CCT phenotype associated with albinism (p < 0.001 for

[Tyr^{c-h} at 32°C standard light] vs. [Tyr^{c-h} at 32°C dark rear]; p < 0.03 for [Tyr^{c-2J} at 10°C standard

light vs. Tyr^{c-2J} at 10°C dark rear]; p < 0.001 for [Tyr^{c-2J} at 32°C standard light vs. Tyr^{c-2J} at 32°C

dark rear]; n = 5–16 per condition; one-way ANOVA with Sidak test; Figure 6; Supplementary

Data 5). Rearing pigmented B6 mice in constant light is known to lead to an increase in axial

length[33]. OCT examinations of independent cohorts of C57BL/6J and Tyr^{c-2J} mice for the

purpose of measuring axial length show that Tyr^{c-2J} mice have a 65.6 µm greater axial length on

average compared to C57BL/6J mice (3.453 mm vs. 3.388 mm, respectively; p < 0.001,

Student's two-tailed t-test; n = 10 male and 10 female mice per strain; Figure 7).

Discussion

Using a phenotype-driven quantitative genetic analysis of CCT, physical mapping led to

identification of a small critical region containing 10 genes, of which we ruled out three (*Fzd4*,

Rab38, *Ctsc*) and found via an analysis of a *Tyr* allelic series (*c*, *c-2J*, *c-h*) that *Tyr* is the

causative gene underlying the *Cctq1a* QTL. *Tyr* is by no means an unknown gene—it was in

fact one of the first known mammalian genes whose initial discovery predates the word

"gene"[27,34,35]. However, *Tyr* was a surprising gene to find linked to CCT. TYR is an oxidase

whose only known biological role relates to melanin synthesis[27,36]. Although there are small

numbers of pigmented cells in the corneal limbus, the cornea is by and large not only non-

pigmented, but transparent. Furthermore, previously published data indicate that *Tyr* is not even

expressed in the adult cornea[18], which is consistent with the current study which also found

near-zero expression in the cornea at 3 weeks of age (RNA-Seq max expression ~1.3 FPKM;

Supplementary Data 7). Thus, there is no molecular rationale for proposing that TYR has a

direct function in corneal cells. If not for the current experiments, there would also be sparse

biological rationales for proposing that TYR might influence corneal anatomy through any

mechanism. Regardless, the current agnostic QTL study led to the conclusion that *Tyr*

contributes to the primary genetic influence on CCT, at least in the context of KS x SJL hybrids.

Experiments using multiple alleles on matched genetic backgrounds, in isolation and genetic

complementation crosses, conclusively confirmed allelism of *Tyr-Cctq1a*; albino mice lacking

TYR function have thin corneas. Rationalizing that albinism would expose the developing eye to

increased light, one of the mechanistic experiments performed here compared the effect of

dark-rearing on the CCT of albino vs. pigmented C57BL/6J mice. The results showed that the

thin CCT phenotype of albino B6.*Tyr*^c-h mice raised at 32°C and B6.*Tyr*^c-2J mice was rescued by

dark-rearing. Thus, we are led to propose an epiphenomenon, whereby developmental light

exposure interacts with genotype as an important determinate of corneal thickness.

All current CCT measurements were done with mice 10–15 weeks of age, which is slightly past

the age at which the cornea of B6 mice reaches its final adult thickness (~P55)[37]. Transcriptomic

changes in the cornea related to *Tyr* genotype were detectable at 3 weeks of age. Thus, the

timeframe for when *Tyr* can impact the cornea is presumably during anterior chamber

development at some point preceding 3 weeks of age. Two mechanisms, which are both

conjectural, might feasibly contribute to this early acting phenomenon. 1) Corneal development

might be a component of refractive development. Emmetropization typically occurs in the first

months following eyelid opening, with impacts from both the amount of light and its focus on the

retina[33,38]. Thus, it is feasible that albinism could in effect cause blur (from light not being

absorbed by melanin and reflecting within the eye), which induces relative myopia and CCT

thinning with rearing in normal lights, but not in dark-rearing. 2) Corneal development might be

influenced by central or corneal circadian outputs. Circadian outputs arise from both central and

peripheral clocks, with the suprachiasmatic nucleus (SCN) being a central master pacemaker

that receives light signals from retinal ganglion cells and subsequently coordinates phasing to

peripheral tissues[39,40]. Notably, the SCN receives retinal input from the retinohypothalamic tract,

which is known to be expanded in albinos[41-43]. Thus, one possibility is that factors increasing

SCN output (such as albinism or cycling light conditions) might lead to decreased CCT and

those decreasing the signaling (such as dark rearing) might lead to increased CCT. Refractive

and circadian mechanisms could also be acting in an intertwined way[44]. Additional experiments

are needed to distinguish these, and possibly other, mechanisms relevant to our current

findings.

A leading candidate for contributing to the molecular mechanism causing thin CCT in albino

mice was DOPA, which is a cofactor for TYR[27,36], a substrate for TH leading to dopamine

(reviewed in [28]), and an endogenous ligand for the G-protein-coupled receptor GPR143[45] .

DOPA can modulate refractive development[27, 29, 40] and the circadian system[46], as well

development of multiple ocular tissues[47,48]. Notably, mice with a conditional knock-out of *Th* in

the retina have decreased CCT[30]. The current experiments with *Tyr* mutant mice were not able

to detect a role for DOPA in influencing CCT of these albino strains, though an important caveat

to point out is that only a single dosing schedule for DOPA supplementation was currently

tested.

It is unclear whether albinism or pigmentation influences CCT in humans, though our current

experiments suggest this is likely. In humans, loss of function mutations in *TYR* cause

oculocutaneous albinism type 1 (OCA 1)[36,49]. It's unclear from the literature if humans with OCA

1 have decreased CCT; it may be difficult to ascertain because of relatively small patient

populations and confounding variables such as eye rubbing[50].

The current study has several implications with respect to mouse genetics and mouse models of

disease: 1) The results highlight the potential for environmental influences on ocular

development, which have been quantitated here for CCT, but may extend to other tissues as

well. 2) The results indicate that albino mice should be tested for potentially being a naturally

occurring model of myopia. 3) Because CCT is a complex trait, there is little reason to suspect

that different inbred albino mouse strains would necessarily have the thinnest CCT in

comparison to other inbred strains that are pigmented[51], only that they would have thinner CCT

as albino mice compared to pigmented mice *within* an inbred strain. However, for any

experiment using such an albino strain, or cohorts in which an allele such as the common Tyr^c

allele is segregating, attention to the possibility of environmental influences is warranted. 4) *Tyr*

has been linked with many ophthalmic traits in mice[41-43,47,52-57]; in some instances, a

consideration of gene-environment interactions in the mechanism of these various models may

be warranted. And finally, 5) our study uncovers a genetic peculiarity. *Cctq1* was originally

reported as a single QTL on chromosome 7, detected in an F2 intercross of KS and SJL inbred

mice[16]. In studies of successive generations of congenic mice, N4F2 mice heterozygous for the

Cctq1 alleles showed the original differential phenotype (over-dominant, *increased* CCT

compared to littermate controls)[16], whereas in N10–N15 intercrosses, mice homozygous for the

SJL allele showed the differential phenotype (recessive, *decreased* CCT compared to littermate

controls). As the sub-congenics were intercrossed and analyzed for CCT, KS.SJL-

Cctq1a^{SJL};Cctq1b^{KS} mice consistently had thinner corneas than KS.SJL-*Cctq1a^{KS};Cctq1b^{KS}*

mice of the same generation, but there were also fluctuations in absolute value related to

generation and interval size. The likely explanation for these observations is that there was

more than one CCT-modifying gene in the original *Cctq1* interval, which was in fact found to be

the case by the resolution of *Cctq1* into *Cctq1a* and *Cctq1b*. This is consistent with the findings

of human GWAS, indicating that there are likely hundreds of CCT-influencing genes dispersed

throughout the genome, many of which will be physically close to one another. In mice, it is

known that as a locus is narrowed using congenics, genes can be segregated away from

nearby modifiers and the overall phenotype of the original QTL can be reduced, disappear, or

reverse its apparent effect[58-60]. The current data seem to exemplify this phenomenon.

In summary, our phenotype-driven genetic study of CCT identified *Tyr* as a significant regulator

of CCT in mice. The molecular findings of this study were unexpected. We propose that the

results can be explained by an epiphenomenon whereby a gene:environment interaction; i.e.,

Tyr-mediated albinism allowing increased exposure of the eye to light has an important

influence on corneal development.

Materials and Methods

Experimental animals

All animals were treated in accordance with the ARVO Statement for the Use of Animals in

Ophthalmic and Vision Research. The majority of mice were housed and bred at the University

of Iowa Research Animal Facility with approval for experimental protocols conferred by the

Institutional Animal Care and Use Committee of the University of Iowa. Two cohorts of mice, the

C57BL/6J (JAX Stock No. 000664) and B6.Cg-*Tyr*$^{c-2J}$/J (JAX Stock No. 000058) used for in vivo

axial length measurements (Figure 7) were purchased from The Jackson Laboratory at 10-

weeks-old and subsequently housed at the University of California San Francisco until 12-

weeks-old with approval for experimental protocols conferred by the Institutional Animal Care

and use Committee at the University of California San Francisco. Mouse strains used in this

study include: SJL/J (Stock No. 000686), C57BLKS/J (JAX Stock No. 000662), C57BL/6J (JAX

Stock No. 000664), B6.Cg-Tyr^{c-2J}/J (JAX Stock No. 000058), B6;129-$Fzd4^{tm1Nat}$/J (JAX Stock

No. 012823), B6.Cg-Tyr^{c-h}/J (JAX Stock No. 000104; Imported from Dr. Brian Brooks at the

NIH), B6.Cg-$Ctsc^{tm1Ley}$ (Imported from Dr. Christine Pham at Washington University), KS.Cg-

SJL^{Cctq1}, B6-$Ctsc^{tm1Mga}$, and B6-Tyr^{tm4Mga}. All experiments included male and female mice.

Chromosome 7 QTL analysis

434   The chromosome 7 quantitative trait locus analysis was performed with R/qtl, using the two-

435   dimensional genome-wide scan (scantwo). Significance thresholds were determined empirically

436   by permutation testing, using 1000 permutations. The validity of a multiple QTL model was

437   tested by performing a multiple regression analysis. Phenotypic variance was estimated and the

438   full model was statistically compared to reduced models in which one QTL was dropped.

Constructing congenic mice

A ~38.3cM genomic region (*i.e.*, *Cctq1*) spanning from *D7Mit318* (SSLP marker at 42.3 cM) to

rs13479545 (SNP marker at 81.2 cM) was transferred from SJL/J mice (abbreviated throughout

as SJL; thick cornea) onto the genetic background of C57BLKS/J mice (abbreviated throughout

as KS; thin cornea) by reiterative backcrossing. Mice carrying the SJL alleles within the region

(*i.e.*, KS.SJL-*Cctq1^{Het}*) were selected at each generation by using a panel of six markers that

were tested and found to be polymorphic between KS and SJL mice. At the N10 generation,

mice were intercrossed. At this point, *Cctq1* was treated as a digenic locus, renamed to *Cctq1a*

and *Cctq1b* (see Results and Figure 2). *Cctq1a* encompassed the region spanning from

D7Mit318 (42.3 cM) to *D7Mit220* (55.7 cM). *Cctq1b* spanned from *D7Mit105* (70.3 cM) to

rs13479545 (81.2 cM). All genotype combinations of *Cctq1a* and *Cctq1b* (9 possible

combinations; *i.e.*, homozygosity for KS alleles, heterozygous, or homozygosity for SJL alleles

at each locus) were analyzed for their effect on CCT.

Sub-congenic mice harboring reduced *Cctq1a* intervals (KS alleles at *Cctq1b*) were also

generated. At N12, *Cctq1a* was reduced to a 15.9 cM region spanning from *D7mit347* to

D7mit321. These N12 sub-congenic mice are referred to throughout as KS.SJL-

Cctq1a(15.9cM). At N15, *Cctq1a* was reduced to a 9.9 cM region spanning from *rs3672782* to

D7mit321. These sub-congenic N15 mice are referred to throughout as KS.SJL-*Cctq1a*(9.9cM).

Sub-congenic mice were intercrossed, and all genotypes were assessed for the CCT

phenotype. The KS.SJL-*Cctq1* line has been sperm cryopreserved.

CCT phenotyping

All measurements were recorded from adult mice. Mice were injected with a standard mixture of

ketamine/xylazine (intraperitoneal injection of 100 mg ketamine + 10 mg xylazine / kg body

weight). During induction of anesthesia, mice were provided supplemental indirect warmth by a

heating pad. Immediately following anesthesia, eyes were hydrated with balanced salt solution

(BSS) and corneal images were obtained with a Bioptigen optical coherence tomographer (SD-

OCT; Bioptigen, Inc., USA). A 12mm telecentric bore was used to image the anterior segment of

each eye. The bore was positioned such that the pupil of the eye was centered in the volume

intensity projection. Scan parameters were as follows: radial volume scans 2.0 mm in diameter,

1000 A-scans/B-scan, 100 B-scans/volume, 1 frame/B-scan, and 1 volume. Central corneal

thickness (CCT) was measured for each eye using vertical angle-locked B-scan calipers. Mice

were included in the analysis if the difference between the right and left eyes was less than 7

μm and if both eyes were free from opacity. The average CCT with standard deviation for each

genotype was statistically compared using Student's two-tailed *t*-test for comparison of two

cohorts or one-way ANOVA with a Tukey post-test for comparison of three or more cohorts.

Recombination mapping

For genetic mapping of the gene underlying *Cctq1a*, additional polymorphic markers were

identified and tiled into the region. Intercrosses of N10 mice were continued and mice with

informative recombination events were analyzed for the CCT phenotype. Based on the allelic

effects of the QTL on the CCT phenotype (see Results), the genomic boundaries of the QTL

(and hence, the region of the underlying gene) were deduced by comparing the phenotype of

the recombinant mice with the location of the recombination event within the critical interval. The

following is a complete list of all the polymorphic markers used for genotyping (listed in order

from centromeric to telomeric): *D7Mit318*, *rs13479346*, *rs13479362*, *D7Mit347*, *D7Mit62*,

rs6271685, *rs108403472*, *Cctq1a-STR5*, *D7Mit31*, *rs3672782*, *rs32438580*, *rs3663323*,

rs13479392, *rs13479393*, *rs6247100*, *rs13479395*, *D7Mit301*, *D7Mit321*, *D7Mit220*, *D7Mit238*,

D7Mit105, *rs13479535*, *rs13479536*, and *rs13479545*. Primer sequences are available upon

request.

Exome sequence analysis

High quality genomic DNA was harvested from KS and SJL spleen tissue using a Qiagen

DNeasy Blood and Tissue kit following the manufacturer's instructions; an RNA digestion step

was included. DNA samples were sent to BGI Americas for sequencing and passed their quality

control standards. Libraries were constructed with an Agilent SureSelect 50Mb Mouse Exome

Capture Kit and were sequenced with 50X coverage using an Illumina HiSeq2000. Standard

bioinformatics analysis was conducted in which the data were filtered (by removing adaptor

contamination and low-quality reads from raw reads), aligned, and SNPs were called and

annotated using a reference genome (GRCm38 build).

RNASeq analysis

N12F3 KS.SJL-*Cctq1a(15.9cM)*^{KS}, KS.SJL-*Cctq1a(15.9cM)*^{SJL}, and KS inbred mice were

euthanized at three weeks of age by cervical dislocation. Immediately upon death, mice were

enucleated, and the eyes were placed in RNase free dishes containing RNA*later* RNA

stabilization reagent. Corneas from mice were dissected in RNA*later* and pooled to make one

sample (6 corneas per sample); three samples were collected per genotype. Cornea samples

were either stored at -80°C in RNA*later* or processed immediately. Corneas were transferred

from RNA*later* to 0.7 mL of lysis/binding buffer from the *mir*Vana miRNA isolation kit (Ambion)

and homogenized for 1 minute using a tissue tearer. The homogenate was then passed through

a QIAshredder column (Qiagen) and the lysate was collected. For the remainder of the

procedure, the samples were processed using the *mir*Vana kit for total RNA according to the

manufacturer's instructions. The quality and concentration of the RNA was analyzed using a

NanoDrop 2000 and the Agilent Model 2100 Bioanalyzer. All samples had RNA integrity

numbers of 9.5 or greater, indicating high quality RNA with little degradation of the samples.

Samples were barcoded and stranded libraries were prepared by the Genomics Division of the

Iowa Institute of Human Genetics. The nine libraries were pooled together, split into two equal

parts, and run on two lanes of an Illumina HiSeq to obtain 100 base pair, paired-end sequence

reads.

Reads were mapped to the mm10 mouse genome build using Tophat2 (ver 2.0.11[61]). The '-r'

parameter was set to 135, and the '--no-coverage-search' option was used. Transcript

abundance was quantified using Cufflinks (ver. 2.1[62]) for RefSeq transcript models from the

Illumina iGenomes mm10 package. Ribosomal RNA and mitochondrial gene loci, obtained from

USCS Genome Table Viewer, were masked from the Cufflinks analysis, and the '--max-bundle-

frags' parameter was set to 20000000. Differential expression between genotypes was

performed using Cuffdiff (ver. 2.1.1[63]). Genes identified from each comparison were

subsequently filtered for only those with a q-value ≤ 0.001 and a mean FPKM (Fragments Per

Kilobase of transcript per Million mapped reads) value ≥ 1 in at least one of the strains.

Functional enrichment analysis was performed with WebGestalt [32, 33] with analysis

parameters detailed in Supplementary Data 8.

Constructing *Ctsc* null mice

B6-*Ctsc^{KO}* mice were generated by the Genome Editing Facility at The University of Iowa on a

pure C57BL/6J (JAX Stock No. 000664) background by targeting *Ctsc* Exon 1 with

CRISPR/Cas9 using guide sequence: CGTGCGCTCCGACACTCCTGCC. Founders were

crossed with C57BL/6J mice and offspring analyzed for germline transmission of *Ctsc*

mutations. From four founders, we observed transmission of four separate *Ctsc* mutations, all

predicted to be null based on Sanger sequencing results. Four *Ctsc* mutations were propagated

in separate intercross mouse lines and were validated as null mutations by Western Blot using

an antibody against CTSC (Catalog #AF1034; R&D Systems, Inc.; Minneapolis, MN;

Supplementary Data 10). One of these mutations, *Ctsc^{tm1Mga}*, is a 46 bp exon 1 coding

sequence deletion that was used for additional downstream analysis.

Constructing *Tyr* null mice

To generate mice with *Tyr* null mutations on a pure C57BL/6J (JAX Stock No. 000664)

background, the Genome Editing Facility at the University of Iowa targeted *Tyr* Exon 1 with

CRISPR/Cas9 using two guide sequences simultaneously: 1) CCATGGATGGGTGATGGGAG

and 2) TTCAAAGGGGTGGATGACCG. Founders were crossed with C57BL/6J mice and

offspring analyzed by Sanger sequencing for germline transmission of *Tyr* mutations. This

experiment generated more alleles than we could reasonably work with (15 unique mutations

identified via sequencing; Supplementary Data 12). For each unique mutation, we set up a

complementation cross with B6.*Tyr^{c-2J}* and screened progeny coat color for alleles conferring

novel function. F1 progeny were screened for 14 alleles; the mice harboring the remaining allele

did not produce F1 progeny. All 14 alleles produced a standard albino coat color in trans with

the *Tyr^{c-2J}* mutation, indicating failure to complement. Going forward, we chose to complete

additional studies for one allele, *Tyr^tm4Mga*, which is a 4 bp deletion in the coding sequence of

exon 1, causes a frameshift and leads to a premature stop codon predicted to cause RNA-

mediated decay (i.e., a presumed null mutation). Accordingly, homozygotes of this strain are

albino.

L-DOPA supplementation studies

Breeder cages of C57BL/6J, B6.Cg-*Tyr^c-2J*/J, and B6.Cg-*Tyr^c-h*/J were provided with water

bottles containing water only (control) or water with 200mg/L of L-DOPA, with 30mg/L of

benserazide to minimize the conversion of L-DOPA to dopamine in the peripheral nervous

system. Fresh water bottles were prepared and supplied every Monday, Wednesday, and

Friday. Litters were weaned into cages where they continued to receive the same treatment of

supplemented water until 13–16 weeks old.

Environmental chamber

Breeder cages of C57BL/6J, B6.Cg-*Tyr^c-2J*/J, and B6.Cg-*Tyr^c-h*/J were housed in rodent

environmental control chambers (Powers Scientific) either at 10°C or 32°C and with either a

standard light cycle (12 h on/12 h off) or dark-rearing. Resulting pups were born and group-

housed at the altered temperature and light cycle until adulthood, when CCT was analyzed by

OCT at 10–15 weeks old.

Axial length phenotyping

Envisu R4300 spectral-domain optical coherence tomography (SD-OCT, Leica/Bioptigen Inc.,

Research Triangle Park, NC, USA) was employed to measure the ocular axial length in adult

(12-week-old) mice[64]. Mice were anesthetized with ketamine/xylazine (100 mg/kg and 5mg/kg,

respectively; intraperitoneal) and their eyes dilated before placing the animal in a cylindrical

holder. The eye was hydrated with Genteal (Alcon, Fort Worth, TX, USA) and positioned in front

of the OCT light source. Correct alignment of the eye was achieved by placing the Purkinje

image in the center of the pupil. The images were acquired in rectangular volume and radial

volume scans. The axial length was calculated by measuring the distance from the anterior

corneal surface to the RPE/choroid interface for both the left and right eyes of a given mouse.

Measurements from the left and right eye of each mouse were averaged to give a single

measurement per animal. Measurements from all eyes were included in the analysis. To

minimize the possible effect of body weight on ocular size, we ensured that body weight of

littermates was within a narrow range in each of the comparative groups. The average axial

length with standard deviation for each genotype was statistically compared using Student's

two-tailed t-test.

Statistics

A multiple regression analysis, in which each locus and the interaction component were

sequentially dropped from the 2-QTL model, was used to analyze the presence of two

interacting loci on chromosome 7. Student's two-tailed t-test was used to evaluate the difference

between two independent genotypes for CCT (t-values and degrees of freedom for each

comparison are listed in Supplementary Data 5) and axial length (t-value = 5.415; degrees of

freedom = 38). A one-way ANOVA with a Tukey post-test was used to evaluate the difference

between three or more independent genotypes for CCT (f-values and degrees of freedom for

each comparison are listed in Supplementary Data 5). A one-way ANOVA with Sidak test was

used to evaluate four comparisons to determine the effect of two environmental conditions on

CCT in B6.cg-Tyr^{c-h} and B6.cg-Tyr^{c-2J} mice (f-value and degrees of freedom are listed in

Supplementary Data 5).

ACKNOWLEDGEMENTS

This work was supported in part by Merit Review Award (101RX001481) from the U.S.

Departments of Veterans Affairs RR&D Service to MGA and by internal faculty development

funds to MGA. DRL was supported in part by a NIH/NEI training grant, EY021436. AHB was

supported in part by a NIH/NIDDK training grant, DK112751. Equipment maintenance was

supported in part by the University of Iowa Multidisciplinary Investigations in Visual Science

service cores (a resource supported by a P30 grant from the NIH/NEI to the University of Iowa,

EY025580). We acknowledge use of the University of Iowa Genome Editing Core Facility (a

core resource supported in part by grants from the NIH and from the Roy J. and Lucille A.

Carver College of Medicine) directed by William Paradee, PhD, and we thank Norma Sinclair,

Patricia Yarolem and Joanne Schwarting for their technical expertise in generating transgenic

mice. The authors thank Karl Broman for analyzing chromosome 7 of our original F2 dataset

with us. The contents do not represent the views of the U.S. Department of Veterans Affairs or

the U.S. Government.

The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

K.J.M., D.R.L., and M.G.A. conceived and designed the experiments. K.J.M., D.R.L., C.J.V.,

A.H.B., L.M.D., N.P., H.E.M., M.N.A., and M.G.A. performed the experimental work. K.J.M.,

D.R.L., and M.G.A. performed experimental data analysis and interpreted the results with

contributions from all authors. S.S.W. analyzed the RNA-seq data supervised by T.E.S. W.J.P.

constructed the transgenic mice. S.K. performed the axial length experiment supervised by

K.S.N. K.W. verified and performed statistical analyses. K.J.M., D.R.L. and M.G.A. wrote the

manuscript with input from all authors. M.G.A. supervised the study and provided funding.

REFERENCES

1. Pediatric Eye Disease Investigator, G. *et al.* Central corneal thickness in children. *Arch Ophthalmol* **129**, 1132-1138, doi:10.1001/archophthalmol.2011.225 (2011).

2. Hashmani, N. *et al.* Effect of age, sex, and refractive errors on central corneal thickness measured by Oculus Pentacam((R)). *Clin Ophthalmol* **11**, 1233-1238, doi:10.2147/OPTH.S141313 (2017).

3. Tayyab, A., Masrur, A., Afzal, F., Iqbal, F. & Naseem, K. Central Corneal Thickness and its Relationship to Intra-Ocular and Epidmiological Determinants. *J Coll Physicians Surg Pak* **26**, 494-497, doi:2349 (2016).

4. Brandt, J. D., Beiser, J. A., Kass, M. A., Gordon, M. O. & Ocular Hypertension Treatment Study, G. Central Corneal Thickness in the Ocular Hypertension Treatment Study (OHTS). *Ophthalmology* **127**, S72-S81, doi:10.1016/j.ophtha.2020.01.028 (2020).

5. La Rosa, F. A., Gross, R. L. & Orengo-Nania, S. Central corneal thickness of Caucasians and African Americans in glaucomatous and nonglaucomatous populations. *Arch Ophthalmol* **119**, 23-27 (2001).

6. Swierkowska, J. & Gajecka, M. Genetic factors influencing the reduction of central corneal thickness in disorders affecting the eye. *Ophthalmic Genet* **38**, 501-510, doi:10.1080/13816810.2017.1313993 (2017).

7. Toh, T. *et al.* Central corneal thickness is highly heritable: the twin eye studies. *Investigative ophthalmology & visual science* **46**, 3718-3722, doi:10.1167/iovs.04-1497 (2005).

8. Zheng, Y. *et al.* Heritability of central corneal thickness in Chinese: the Guangzhou Twin Eye Study. *Investigative ophthalmology & visual science* **49**, 4303-4307, doi:10.1167/iovs.08-1934 (2008).

9. Choquet, H. *et al.* A multiethnic genome-wide analysis of 44,039 individuals identifies 41 new loci associated with central corneal thickness. *Commun Biol* **3**, 301, doi:10.1038/s42003-020-1037-7 (2020).

10. Iglesias, A. I. *et al.* Cross-ancestry genome-wide association analysis of corneal thickness strengthens link between complex and Mendelian eye diseases. *Nat Commun* **9**, 1864, doi:10.1038/s41467-018-03646-6 (2018).

11. Dhooge, T. *et al.* More than meets the eye: Expanding and reviewing the clinical and mutational spectrum of brittle cornea syndrome. *Hum Mutat*, doi:10.1002/humu.24199 (2021).

12. Lu, Y. *et al.* Genome-wide association analyses identify multiple loci associated with central corneal thickness and keratoconus. *Nature genetics* **45**, 155-163, doi:10.1038/ng.2506 (2013).

13. Lu, Y. *et al.* Common genetic variants near the Brittle Cornea Syndrome locus ZNF469 influence the blinding disease risk factor central corneal thickness. *PLoS Genet* **6**, e1000947, doi:10.1371/journal.pgen.1000947 (2010).

14. Dimasi, D. P. *et al.* Novel quantitative trait loci for central corneal thickness identified by candidate gene analysis of osteogenesis imperfecta genes. *Hum Genet* **127**, 33-44, doi:10.1007/s00439-009-0729-3 (2010).

15. Zhang, J., Wu, D., Dai, Y. & Xu, J. Functional relevance for central cornea thickness-associated genetic variants by using integrative analyses. *BioData Min* **11**, 19, doi:10.1186/s13040-018-0179-3 (2018).

16 Lively, G. D., Koehn, D., Hedberg-Buenz, A., Wang, K. & Anderson, M. G. Quantitative
trait loci associated with murine central corneal thickness. *Physiological genomics* **42**,
281-286, doi:10.1152/physiolgenomics.00140.2009 (2010).
17 Koehn, D. R., Meyer, K. J. & Anderson, M. G. Genetic Evidence for Differential
Regulation of Corneal Epithelial and Stromal Thickness. *Invest Ophthalmol Vis Sci* **56**,
5599-5607, doi:10.1167/iovs.15-17179 (2015).
18 Lively, G. D. *et al.* Genetic Dependence of Central Corneal Thickness among Inbred
Strains of Mice. *Investigative ophthalmology & visual science* **51**, 160-171,
doi:10.1167/iovs.09-3429 (2010).
19 Broman, K. W. & Sen, S. *A Guide to QTL Mapping with R/qtl.* (Springer
Science+Business Media, 2009).
20 Broman, K. W., Wu, H., Sen, S. & Churchill, G. A. R/qtl: QTL mapping in experimental
crosses. *Bioinformatics* **19**, 889-890 (2003).
21 Wynn, E. H., Sanchez-Andrade, G., Carss, K. J. & Logan, D. W. Genomic variation in
the vomeronasal receptor gene repertoires of inbred mice. *BMC Genomics* **13**, 415,
doi:10.1186/1471-2164-13-415 (2012).
22 Liao, Y., Wang, J., Jaehnig, E. J., Shi, Z. & Zhang, B. WebGestalt 2019: gene set
analysis toolkit with revamped UIs and APIs. *Nucleic acids research* **47**, W199-W205,
doi:10.1093/nar/gkz401 (2019).
23 Wang, Y., Huso, D., Cahill, H., Ryugo, D. & Nathans, J. Progressive cerebellar, auditory,
and esophageal dysfunction caused by targeted disruption of the frizzled-4 gene. *J
Neurosci* **21**, 4761-4771 (2001).
24 Pham, C. T. & Ley, T. J. Dipeptidyl peptidase I is required for the processing and
activation of granzymes A and B in vivo. *Proc Natl Acad Sci U S A* **96**, 8627-8632,
doi:10.1073/pnas.96.15.8627 (1999).
25 Le Fur, N., Kelsall, S. R. & Mintz, B. Base substitution at different alternative splice
donor sites of the tyrosinase gene in murine albinism. *Genomics* **37**, 245-248,
doi:10.1006/geno.1996.0551 (1996).
26 Green, E. L. Albino-2J (c<2J>). *Mouse News Lett* **49** (1973).
27 Beermann, F., Orlow, S. J. & Lamoreux, M. L. The Tyr (albino) locus of the laboratory
mouse. *Mamm Genome* **15**, 749-758, doi:10.1007/s00335-004-4002-8 (2004).
28 Zhou, X., Pardue, M. T., Iuvone, P. M. & Qu, J. Dopamine signaling and myopia
development: What are the key challenges. *Prog Retin Eye Res* **61**, 60-71,
doi:10.1016/j.preteyeres.2017.06.003 (2017).
29 Feldkaemper, M. & Schaeffel, F. An updated view on the role of dopamine in myopia.
Exp Eye Res **114**, 106-119, doi:10.1016/j.exer.2013.02.007 (2013).
30 Bergen, M. A. *et al.* Altered Refractive Development in Mice With Reduced Levels of
Retinal Dopamine. *Investigative ophthalmology & visual science* **57**, 4412-4419,
doi:10.1167/iovs.15-17784 (2016).
31 Kidson, S. & Fabian, B. Pigment synthesis in the Himalayan mouse. *J Exp Zool* **210**,
145-152, doi:10.1002/jez.1402100116 (1979).
32 Kwon, B. S., Halaban, R. & Chintamaneni, C. Molecular basis of mouse Himalayan
mutation. *Biochem Biophys Res Commun* **161**, 252-260, doi:10.1016/0006-
291x(89)91588-x (1989).
33 Tkatchenko, T. V. *et al.* Photopic visual input is necessary for emmetropization in mice.
Exp Eye Res **115**, 87-95, doi:10.1016/j.exer.2013.06.025 (2013).

34 Allen, G. M. The heredity of coat color in mice. *Proceedings of the American Academy*
of Arts and Sciences **40**, 61-163 (1904).
35 Castle, W. E. & Allen, G. M. The heredity of albinism. *Proceedings of the American*
Academy of Arts and Sciences **38**, 603-622 (1903).
36 Seruggia, D., Josa, S., Fernandez, A. & Montoliu, L. The structure and function of the
mouse tyrosinase locus. *Pigment Cell Melanoma Res*, doi:10.1111/pcmr.12942 (2020).
37 Hanlon, S. D., Patel, N. B. & Burns, A. R. Assessment of postnatal corneal development
in the C57BL/6 mouse using spectral domain optical coherence tomography and
microwave-assisted histology. *Exp Eye Res* **93**, 363-370, doi:10.1016/j.exer.2011.04.013
(2011).
38 Chakraborty, R., Ostrin, L. A., Benavente-Perez, A. & Verkicharla, P. K. Optical
mechanisms regulating emmetropisation and refractive errors: evidence from animal
models. *Clin Exp Optom* **103**, 55-67, doi:10.1111/cxo.12991 (2020).
39 Mohawk, J. A., Green, C. B. & Takahashi, J. S. Central and peripheral circadian clocks in
mammals. *Annu Rev Neurosci* **35**, 445-462, doi:10.1146/annurev-neuro-060909-153128
(2012).
40 Peirson, S. N., Brown, L. A., Pothecary, C. A., Benson, L. A. & Fisk, A. S. Light and the
laboratory mouse. *J Neurosci Methods* **300**, 26-36, doi:10.1016/j.jneumeth.2017.04.007
(2018).
41 Steininger, T. L., Rye, D. B., Gilliland, M. A., Wainer, B. H. & Benca, R. M. Differences
in the retinohypothalamic tract in albino Lewis versus brown Norway rat strains.
Neuroscience **54**, 11-14, doi:10.1016/0306-4522(93)90379-t (1993).
42 Miller, A. M., Chappell, R., Obermeyer, W. H. & Benca, R. M. Analysis of the
retinohypothalamic tract in congenic albino and pigmented rats. *Brain Res* **741**, 348-351,
doi:10.1016/s0006-8993(96)01050-5 (1996).
43 Fleming, M. D., Benca, R. M. & Behan, M. Retinal projections to the subcortical visual
system in congenic albino and pigmented rats. *Neuroscience* **143**, 895-904,
doi:10.1016/j.neuroscience.2006.08.016 (2006).
44 Chakraborty, R. *et al.* Circadian rhythms, refractive development, and myopia.
Ophthalmic Physiol Opt **38**, 217-245, doi:10.1111/opo.12453 (2018).
45 Lopez, V. M., Decatur, C. L., Stamer, W. D., Lynch, R. M. & McKay, B. S. L-DOPA is
an endogenous ligand for OA1. *PLoS Biol* **6**, e236, doi:10.1371/journal.pbio.0060236
(2008).
46 Korshunov, K. S., Blakemore, L. J. & Trombley, P. Q. Dopamine: A Modulator of
Circadian Rhythms in the Central Nervous System. *Front Cell Neurosci* **11**, 91,
doi:10.3389/fncel.2017.00091 (2017).
47 Libby, R. T. *et al.* Modification of ocular defects in mouse developmental glaucoma
models by tyrosinase. *Science* **299**, 1578-1581, doi:10.1126/science.1080095 (2003).
48 Lee, H., Scott, J., Griffiths, H., Self, J. E. & Lotery, A. Oral levodopa rescues retinal
morphology and visual function in a murine model of human albinism. *Pigment Cell*
Melanoma Res **32**, 657-671, doi:10.1111/pcmr.12782 (2019).
49 Kirkwood, B. J. Albinism and its implications with vision. *Insight* **34**, 13-16 (2009).
50 Rao, V. A., Swathi, P., Chaitra & Thappa, D. M. Bilateral keratoconus with
oculocutaneous albinism. *Indian J Dermatol Venereol Leprol* **74**, 407-409,
doi:10.4103/0378-6323.42929 (2008).

769 51 Dimasi, D. P. *et al.* Ethnic and mouse strain differences in central corneal thickness and
770   association with pigmentation phenotype. *PLoS One* **6**, e22103,
771   doi:10.1371/journal.pone.0022103 (2011).
772 52 Buttner, C. *et al.* Tyrosinase Is a Novel Endogenous Regulator of Developmental and
773   Inflammatory Lymphangiogenesis. *Am J Pathol* **189**, 440-448,
774   doi:10.1016/j.ajpath.2018.10.014 (2019).
775 53 Mason, C. & Slavi, N. Retinal Ganglion Cell Axon Wiring Establishing the Binocular
776   Circuit. *Annu Rev Vis Sci* **6**, 215-236, doi:10.1146/annurev-vision-091517-034306
777   (2020).
778 54 Savinova, O. V. *et al.* Intraocular pressure in genetically distinct mice: an update and
779   strain survey. *BMC Genet* **2**, 12, doi:10.1186/1471-2156-2-12 (2001).
780 55 Johnson, B. A., Cole, B. S., Geisert, E. E., Ikeda, S. & Ikeda, A. Tyrosinase is the
781   modifier of retinoschisis in mice. *Genetics* **186**, 1337-1344,
782   doi:10.1534/genetics.110.120840 (2010).
783 56 Anderson, M. G. *et al.* Genetic context determines susceptibility to intraocular pressure
784   elevation in a mouse pigmentary glaucoma. *BMC Biol* **4**, 20, doi:10.1186/1741-7007-4-
785   20 (2006).
786 57 Oh, J. *et al.* Genetic background-dependent role of Egr1 for eyelid development. *Proc*
787   *Natl Acad Sci U S A* **114**, E7131-E7139, doi:10.1073/pnas.1705848114 (2017).
788 58 Mott, R. & Flint, J. Dissecting quantitative traits in mice. *Annual review of genomics and*
789   *human genetics* **14**, 421-439, doi:10.1146/annurev-genom-091212-153419 (2013).
790 59 Solberg Woods, L. C. QTL mapping in outbred populations: successes and challenges.
791   *Physiological genomics* **46**, 81-90, doi:10.1152/physiolgenomics.00127.2013 (2014).
792 60 Meyer, K. J. & Anderson, M. G. Genetic modifiers as relevant biological variables of eye
793   disorders. *Hum Mol Genet* **26**, R58-R67, doi:10.1093/hmg/ddx180 (2017).
794 61 Kim, D. *et al.* TopHat2: accurate alignment of transcriptomes in the presence of
795   insertions, deletions and gene fusions. *Genome Biol* **14**, R36, doi:10.1186/gb-2013-14-4-
796   r36 (2013).
797 62 Trapnell, C. *et al.* Transcript assembly and quantification by RNA-Seq reveals
798   unannotated transcripts and isoform switching during cell differentiation. *Nat Biotechnol*
799   **28**, 511-515, doi:10.1038/nbt.1621 (2010).
800 63 Trapnell, C. *et al.* Differential analysis of gene regulation at transcript resolution with
801   RNA-seq. *Nat Biotechnol* **31**, 46-53, doi:10.1038/nbt.2450 (2013).
802 64 Paylakhi, S. *et al.* Muller glia-derived PRSS56 is required to sustain ocular axial growth
803   and prevent refractive error. *PLoS Genet* **14**, e1007244,
804   doi:10.1371/journal.pgen.1007244 (2018).
805

Figure 1. *Cctq1* contains two adjacent interacting QTL, *Cctq1a* and *Cctq1b*. a Heatmap of

the chromosome 7 pairwise scan. The *upper left triangle* displays the interactive LOD score

(LOD$_i$; left side of the heat map scale) and the *lower right triangle* displays the full LOD score

(LOD$_f$; right side of the heat map scale). Chromosome 7 positions (Mb) are based on NCBI

Build 33. **b** Genetic map of chromosome 7 showing the *Cctq1a* and *Cctq1b* loci. The area *boxed*

in red is the original *Cctq1* interval blown up to show the intervals of *Cctq1a* and *Cctq1b*, and a

subset of the polymorphic markers used for genotyping. The *arrows* indicate the peaks of the

two QTL.

Table 1. CCT phenotype results of adult N10F2 mice.

$Cctq1a$[a]	$Cctq1b$[a]	Avg. CCT ± StDev.[b]	# Mice
KS	KS	94.8 ± 2.4	13
KS	Het	93.3 ± 2.5	13
KS	SJL	90.8 ± 2.6	13
Het	KS	95.5 ± 2.9	13
Het	Het	96.2 ± 3.4	13
Het	SJL	97.0 ± 2.3	13
SJL	KS	88.1 ± 3.7*	13
SJL	Het	86.8 ± 4.2*	13
SJL	SJL	88.8 ± 3.6*	13

[a] KS, homozygous for KS alleles across the indicated QTL; Het, heterozygous for alleles across

the indicated QTL; SJL, homozygous for SJL alleles across the indicated QTL

[b] Asterisk (*), significantly different ($p < 0.01$) than inbred KS mice by one-way ANOVA with

Tukey post-test

Figure 2. Genetic mapping of *Cctq1a* on mouse chromosome 7 using intercrosses of

KS.SJL-*Cctq1a* congenic and sub-congenic mice. *Black boxes* represent the KS or HET

genotype associated with a thick cornea and *white boxes* represent the SJL genotype

associated with a thin cornea. The number of mice (*n*) with each haplotype is listed to the left of

each row, with progeny-tested mice denoted with an asterisk (*). The adult CCT phenotype

(*pheno*; measured by optical coherence tomography) relative to littermate controls and the

deduced genotype (*geno*) for each haplotype is listed to the right of each row as "Thick KS" or

"Thin SJL". The *vertical lines* across the chromosome represent markers that are polymorphic

between KS and SJL mice. *Black arrowheads* indicate partial genes.

Figure 3. Testing the influence of *Cctq1a* positional candidate genes on central corneal thickness (CCT). Each point on the graph represents the average CCT measured by optical coherence tomography from one adult mouse with age-matching across genotypes and *error bars* = mean ± standard deviation. **a** Plot comparing CCT for *Cctq1a^{KS}* vs. *Cctq1a^{SJL}* genotypes in KS.SJL-*Cctq1a* N15F7 sub-congenic littermate mice 12 to13 weeks old (Student's two-tailed *t*-test). **b** Plot comparing CCT between *Fzd4* genotypes for littermate mice 12 to 13 weeks old (one-way ANOVA with Tukey test). **c** Plot comparing the CCT for mice with a *Fzd4^{tm1Nat}* allele in trans with a *Cctq1a^{SJL}* allele to littermate controls having a *Fzd4^{WT}* allele in trans with a *Cctq1a^{SJL}* allele (Student's two-tailed *t*-test; 17 to 33 weeks old). **d** Plot comparing CCT between *Ctsc^{tm1Ley}* genotypes for 11-week-old littermate mice (one-way ANOVA with Tukey test). **e** Plot comparing CCT for mice with a *Ctsc^{tm1Ley}* allele in trans with a *Cctq1a^{SJL}* allele compared to littermate controls having a *Ctsc^{tm1Ley}* allele in trans with a *Cctq1a^{KS}* allele (Student's two-tailed *t*-test; 15 to18 weeks old). **f** Plot comparing CCT for mice harboring one of four *Ctsc^{KO}* alleles on a pure B6 background to littermate controls (one-way ANOVA with Tukey test; 10 to 12 weeks old). **g** Plot comparing CCT for mice with a *Ctsc^{tm1Mga}* allele, made on a pure B6 background, in trans with a *Cctq1a^{SJL}* allele to littermate controls having a *Ctsc^{WT}* allele in trans with a *Cctq1a^{SJL}* allele (Student's two-tailed *t*-test; 10 to 11 weeks old). **h** Plot comparing CCT for B6.*Tyr^{c-2J}* mice to B6 mice (Student's two-tailed *t*-test; 11 to 13 weeks old). **i** Plot comparing CCT for mice with a *Tyr^{c-2J}* allele in trans with a *Cctq1a^{SJL}* allele to littermate controls having a *Tyr^{c-2J}* allele in trans with a *Cctq1a^{KS}* allele (Student's two-tailed *t*-test; 10 to 12 weeks old).

Figure 4. The influence of a null *Tyr* mutation on central corneal thickness (CCT). Each

point on the graph represents the average CCT measured by optical coherence tomography

from one adult mouse, 13 to 17 weeks old, with age-matching across genotypes and *error bars*

= mean ± standard deviation. The plot compares CCT between mice homozygous for the

Tyr^tm4Mga allele on a pure B6 background to littermate controls with heterozygous (HET) or wild-

type (WT) alleles (one-way ANOVA with Tukey test).

863
864

Scatter plot with y-axis "CCT (µm)" ranging 75 to 115. Three groups: Tyr^WT (black), Tyr^HET (blue), Tyr^tm4Mga (red). Two comparison bars labeled p < 0.002.

1 **Figure 5. Testing the influence of DOPA on central corneal thickness (CCT).** Each point on the graph represents the average

2 CCT measured by optical coherence tomography from one adult mouse, 13 to 16 weeks old, with age-matching across genotypes

3 and *error bars* = mean ± standard deviation. Plots compare CCT between mice supplied with DOPA in the drinking water from

4 conception through 10-weeks-old to controls supplied with standard drinking water for the **a** B6, **b** B6.$Tyr^{c\text{-}h}$, and **c** B6.$Tyr^{c\text{-}2J}$ strains of

5 mice (Student's two-tailed *t*-test for each comparison).

Figure 6. The influence of temperature and light on central corneal thickness (CCT) for

the B6, B6.*Tyr^c-h*, and B6.*Tyr^c-2J* strains of mice. Each point on the graph represents the

average CCT measured by optical coherence tomography from one adult mouse, 10 to 15

weeks old, with age-matching across genotypes and *error bars* = mean ± standard deviation.

Cohorts were subjected to standard lighting or dark-rearing. Each grouping of *Tyr* genotypes

(WT B6 = *black*; *c-h* temperature-sensitive himalayan = *blue*; *c-2J* albino = *red*) has mice raised

at 10°C or 32°C (one-way ANOVA with Sidak test).

1   **Figure 7. The influence of *Tyr* on ocular axial length.** Each point on the graph represents the

2   average axial length measured by optical coherence tomography from one 12-week-old adult

3   mouse and *error bars* = mean ± standard deviation. The plot compares axial length between the

4   B6 and B6.*Tyr*^*c-2J* strains of mice, measured from the outer surface of the cornea to the retinal

5   pigment epithelium/choroid interface (Student's two-tailed *t*-test).