# Nonparametric Interrogation of Transcriptional Regulation in Single-Cell RNA and Chromatin Accessibility Multiomic Data

Yuchao Jiang[1,2,3,*], Yuriko Harigaya[4], Zhaojun Zhang[5], Hongpan Zhang[6,7], Chongzhi Zang[6,7,8], Nancy R Zhang[5,*]

[1]    Department of Biostatistics, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC 27599, USA.

[2]    Department of Genetics, School of Medicine, University of North Carolina, Chapel Hill, NC 27599, USA.

[3]    Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, NC 27599, USA.

[4]    Curriculum in Bioinformatics and Computational Biology, School of Medicine, University of North Carolina, Chapel Hill, NC 27599, USA.

[5]    Department of Statistics, The Wharton School, University of Pennsylvania, Philadelphia, PA 19104, USA.

[6]    Center for Public Health Genomics, University of Virginia, Charlottesville, VA 22908, USA.

[7]    Department of Biochemistry and Molecular Genetics, University of Virginia, Charlottesville, VA 22908, USA.

[8]    Department of Public Health Sciences, University of Virginia, Charlottesville, VA 22908, USA.

*    To whom correspondence should be addressed: yuchaoj@email.unc.edu, nzh@wharton.upenn.edu.

**Abstract**

Epigenetic control of gene expression is highly cell-type- and context-specific. Yet, despite its complexity, gene regulatory logic can be broken down into modular components consisting of a transcription factor (TF) activating or repressing the expression of a target gene through its binding to a *cis*-regulatory region. Recent advances in joint profiling of transcription and chromatin accessibility with single-cell resolution offer unprecedented opportunities to interrogate such regulatory logic. Here, we propose a nonparametric approach, TRIPOD, to detect and characterize three-way relationships between a TF, its target gene, and the accessibility of the TF's binding site, using single-cell RNA and ATAC multiomic data. We apply TRIPOD to interrogate cell-type-specific regulatory logic in peripheral blood mononuclear cells and contrast our results to detections from enhancer databases, *cis*-eQTL studies, ChIP-seq experiments, and TF knockdown/knockout studies. We then apply TRIPOD to mouse embryonic brain data during neurogenesis and gliogenesis and identified known and novel putative regulatory relationships, validated by ChIP-seq and PLAC-seq. Finally, we demonstrate TRIPOD on SHARE-seq data of differentiating mouse hair follicle cells and identify lineage-specific regulation supported by histone marks for gene activation and super-enhancer annotations.

**Keywords**: single-cell multiomics, transcriptional regulation, transcription factor, chromatin accessibility.

Context-specific regulation of gene transcription is central to cell identity and function in eukaryotes. Precision of transcriptional control is achieved through multitudes of transcription factors (TFs) that bind to the *cis*-regulatory regions of their target genes, dynamically modulating chromatin accessibility and recruiting transcription complexes in response to developmental and environmental cues[1]. Dissecting this regulatory logic is fundamental to our understanding of biological systems and our study of diseases. Over the past decades, molecular studies have elucidated the structure of TF complexes and provided mechanistic models into their function[2]. Methods based on high-throughput sequencing have enabled the genome-wide profiling of gene expression[3], TF binding[4], chromatin accessibility[5], and 3D genome structure[6]. TF knockdown/knockout studies have also identified, *en masse*, their species-, tissue-, and context-specific target genes[7]. Concurrently, novel statistical approaches have allowed for more precise identification and modeling of TF binding sites[8], and expression quantitative trait loci (eQTLs) databases now include associations that are tissue-specific[9] and will soon be cell-type specific[10]. Yet, despite this tremendous progress, our understanding of gene regulatory logic is still rudimentary. When a TF activates or represses the expression of a gene through binding to a regulatory element in *cis* to the gene, we call such a relationship a *regulatory trio*. Despite its complexity, gene regulatory logic can be broken down into modular components consisting of such peak-TF-gene trios. In this paper, we focus on the identification of regulatory trios using multiomic experiments that jointly profile gene expression and chromatin accessibility at single-cell resolution.

Single-cell RNA sequencing (scRNA-seq) and single-cell assay of transposase-accessible chromatin sequencing (scATAC-seq), performed separately, have already generated detailed cell-type-specific profiles of gene expression and chromatin accessibility. When the two modalities are not measured in the same cells, the cells can be aligned by computational methods[11], followed by association analyses of gene expression and peak accessibility. While these methods have been shown to align well-differentiated cell types correctly, they often fail for cell populations consisting of transient and closely similar cell states. Additionally, the alignment of cells between scRNA-seq and scATAC-seq necessarily assumes a peak-gene relationship which is usually learned from other datasets. Then, the post-alignment association analysis is plagued by logical

circularity, as it is difficult to disentangle new findings from prior assumptions that underlie
the initial cell alignment.

Single-cell multiomic experiments that sequence the RNA and ATAC from the
same cells directly enable joint modeling of a cell's RNA expression and chromatin state,
yet methods for the analysis of such data are still in their infancy. Almost all existing
methods for detecting and characterizing regulatory relationships between TF, regulatory
region, and target gene rely only on marginal relationships, i.e., associations between two
of the three entities without conditioning on the third. For example, Signac[12] and Ma *et*
*al.*[13] use marginal associations between peaks and genes to identify putative enhancer
regions, while Signac[12] and Seurat V4[14] link differentially expressed TFs to differentially
accessible motifs across cell types. Such pairwise marginal associations are sometimes
examined manually using low-dimensional embedding. One exception is PECA[15], which
uses a parametric model to characterize the joint four-way relationship between TF
expression, regulatory site accessibility, chromatin remodeler expression, and target
gene expression. Although PECA was designed to be applied to matched bulk
transcriptomic and epigenomic data, such joint modeling concepts could potentially be
very powerful for single-cell multiomic data. In this paper, we propose a nonparametric
approach as an alternative to PECA's parametric model, thus allowing for robustness and
computational scalability.

As we will show through examples, context-specific gene regulation, such as cell-
type-specific regulation, may be masked in marginal associations. For example,
associations between a TF and its target gene may be apparent only conditional on the
accessibility of its binding site. Or, associations between the accessibility of an enhancer
and its target gene may be apparent only after accounting for the expression of certain
transcription factors involved in, but not sufficient for, the remodeling of the enhancer
region. The identification and characterization of such context-specific relationships are
relevant, for example, in the interpretation of GWAS results, where marginal pairwise
associations between ATAC peaks and gene expression have had limited success in
linking disease-associated SNPs to genes[16].

We explore in this paper the use of higher-order models that interrogate conditional
and three-way interaction relationships for the identification of regulatory trios. First, as

proof of principle, we show that a simple model that integrates TF expression with *cis*-
peak accessibility significantly improves gene expression prediction, as compared to a
comparable model that utilizes peak accessibility alone. We present TRIPOD, a
computational framework for *t*ranscription *r*egulation *i*nterrogation through nonparametric
*p*artial association analysis of single-cell multi*o*mic sequencing *d*ata. TRIPOD detects two
types of trio relationships, which we call conditional level 1 and conditional level 2, through
robust nonparametric tests that are easy to diagnose. TRIPOD's nonparametric approach
for the identification of conditional associations avoids assumptions of linearity of
relationships and normality of errors, allowing for better adjustment for confounding. Thus,
given a multiome experiment that measures RNA expression and chromatin accessibility
for the same cells at single-cell resolution, TRIPOD outputs, for a list of transcription
factors, their putative regulatory gene targets and the *cis* regions where they putatively
bind to regulate each gene. This allows the prioritization of regulatory relationships for
downstream analyses. We also develop a novel influence measure that allows the
detection and visualization of cell states driving these regulatory relationships, applicable
to data consisting of discrete cell types as well as continuous cell trajectories.

We first apply TRIPOD to single-cell multiomic data of human peripheral blood
mononuclear cells (PBMCs) and compare the regulatory trios detected to relationships
detected through marginal associations. We show that the detections are coherent with
the vast amounts of existing knowledge from enhancer databases, bulk cell-type-specific
chromatin immunoprecipitation followed by sequencing (ChIP-seq) experiments, tissue-
specific TF knockdown/knockout studies, and *cis*-eQTL studies, but that conditional and
marginal models identify different sets of relationships. We next apply TRIPOD to the
interrogation of lineage-specific regulation in the developing mouse brain, where
relationships detected by TRIPOD are compared against those derived from existing
ChIP-seq and proximity ligation-assisted ChIP-seq (PLAC-seq) data. Here, TRIPOD
identifies known trio relationships, as well as putative novel regulatory crosstalk between
neuronal TFs and glial-lineage genes. We also apply TRIPOD to SHARE-seq data on
mouse hair follicle cell differentiation to illustrate trio detection and influence analysis in
data collected from different protocols. Through these analyses, we demonstrate how to

harness single-cell multiomic technologies in the study of gene regulation and how the

data from these technologies corroborate and complement existing data.

**Results**

**A simple interaction model between TF expression and peak accessibility improves**

**RNA prediction**. To motivate our methods, we start with a simple prediction-based

analysis, comparable to that done by existing methods[11]. We benchmarked against: (i)

Signac[12] and Cicero[17], which predict gene expression by the gene activity matrix derived

from the sum of the ATAC reads in gene bodies and promoter regions; (ii) MAESTRO[18],

which predicts gene expression using a regulatory potential model that sums ATAC reads

weighted based on existing gene annotations; and (iii) sci-CAR[19], which predicts gene

expression by a regularized regression on coverage of individual peaks nearby. We

compared the predictions derived from these methods to that of a regularized regression

model, where for predictors, peak accessibilities are replaced by products between peak

accessibilities and TF expressions. Only peaks within a certain range of the gene's

transcription start site (TSS) and only interactions between TFs and peaks containing

high-scoring binding motifs for the TFs are considered. We refer to this model as the

peak-TF LASSO model. Since this model is prediction-based, we do not expect the peak-

TF pairs selected by LASSO to necessarily have a causal regulatory relationship to the

gene. Comparing this model to (i)-(iii) allows us to assess whether the peak-TF interaction

terms are informative for gene expression. To avoid overfitting, we performed out-of-fold

prediction and adopted independent training and testing sets. See Methods for details.

We analyzed single-cell multiomic datasets from different human and mouse

tissues generated by different platforms – PBMC by 10X Genomics, embryonic mouse

brain by 10X Genomics, mouse skin by SHARE-seq[13], and adult mouse brain by SNARE-

seq[20]. Data summaries are included in Supplementary Table 1; reduced dimensions via

uniform manifold approximation and projection (UMAP)[21] are shown in Fig. 1a and

Supplementary Fig. 1, 2a. To mitigate the undesirable consequences of sparsity and

stochasticity in the single-cell data, we clustered cells to form metacells[14] and pooled

gene expression and chromatin accessibility measurements within each metacell.

Our results show that, across window sizes, the peak-TF LASSO model
significantly improves prediction accuracy across the transcriptome (Fig.1b), with
examples of specific genes shown in Fig. 1c. This improvement in prediction accuracy
holds true when an independent dataset is used for validation (Supplementary Fig. 3).
For the SNARE-seq data[20], sequencing depth is substantially shallower (Supplementary
Fig. 4), thus the improvement of the peak-TF LASSO model is diminished but still evident
(Supplementary Fig. 2b). This demonstrates that the product of TF expression and peak
accessibility significantly improves RNA prediction accuracy beyond simply using peak
accessibility, offering strong empirical evidence of three-way interaction relationships
between TF expression, peak accessibility, and target gene expression that can be
extracted from such multiomic experiments. However, we will not rely on coefficients from
the LASSO model to screen for such trios, as their significance is difficult to compute due
to the hazards of post-selection inference[22]. Additionally, accessibility of peaks and
expression of TF affecting the same gene are often highly correlated, in which case
LASSO tends to select the few with the highest associations and ignore the rest. In such
cases, we believe it is more desirable to report all trios.

**TRIPOD for the detection of peak-TF-gene trio regulatory relationships by single-**
**cell multiomic data.** We propose TRIPOD, a nonparametric method that screens single-
cell RNA and ATAC multiomic data for conditional associations and three-way
interactions between the expression of a TF $t$, the accessibility of a peak region
$p$ containing the TF's motif, and the expression of a putative target gene $g$ within a pre-
fixed distance of peak $p$ (Fig. 2a). Existing methods[12-14] screen for marginal associations
either between the TF and the peak or between the peak and the target gene. However,
three-way relationships may be complex: When a TF binds to a *cis*-regulatory region to
affect the expression of a gene, it can do so in multiple ways, leading to different patterns
in the data. The TF could be directly responsible for opening the chromatin of the
enhancer region, facilitating the binding of other TFs that recruit the RNA polymerase. In
such cases, expression of the TF is likely to be marginally correlated with the accessibility
of the enhancer region, but its correlation with the expression of the target gene may be
masked due to confounding of other involved TFs. Alternatively, the TF may not be

directly responsible for chromatin remodeling but may bind to already accessible
chromatin in recruiting other TFs or the RNA polymerase. In such cases, expression of
the TF may not be highly correlated with the accessibility of the enhancer region or with
the expression of the target gene. When marginal associations are masked, evidence for
binding of the TF at the peak in the regulation of a gene can be inferred from partial
associations: (i) with the peak open at a fixed accessibility, whether cells with higher TF
expression have higher gene expression; and (ii) with the TF expression fixed at a value
above a threshold, whether cells with higher peak accessibility have higher gene
expression. To identify such conditional associations without making linearity
assumptions on the marginal relationships, TRIPOD matches metacells by either their TF
expressions or peak accessibilities (Fig. 2b): for each matched metacell pair, the variable
being matched is controlled for, and differences between the pair in the other two
variables are computed. Then, across pairs, the nonparametric Spearman's test is used
to assess the association between the difference in target gene expression $\Delta Y_g$ and
difference in the unmatched variable (i.e., $\Delta Y_t$ if the cells were matched by $X_p$, or $\Delta X_p$ if
the cells were matched by $Y_t$). We call this the "conditional level 1 test."

For illustration, consider the metacell denoted by the black point in Fig. 2b: If we
were to match by peak accessibility, this metacell would be matched to the metacell
colored in red. We would then compute $\Delta Y_t$, the difference between TF $t$ expressions of
the matched pair. If we were to match by TF expression, the black dot would be matched
to the metacell in green, and we would compute $\Delta X_p$, the difference in peak $p$ accessibility
for this pair. In either case, we would compute $\Delta Y_g$, the difference in gene $g$ expressions
between the pair. We would then mask those metacell-pairs whose values, for the
variable being matched, are too low (i.e., those pairs where the TF is off or the peak is
closed). Then, $\Delta X_p$ or $\Delta Y_t$, together with $\Delta Y_g$, would be submitted for level 1 test. We call
such a triplet of TF, peak, and target gene a "regulatory trio."

Even stronger evidence for a regulatory trio could be claimed if the *degree* of
association between the pairwise differences depends on the matched variable. For
example, we would tend to believe that TF $t$ binds to peak $p$ to regulate gene $g$ if, in cells
with high expression of TF $t$, an increase in peak $p$ accessibility yields a much larger
increase in gene $g$ expression, as compared to in cells with low expression of TF $t$. One

could screen for such interactions by matching by either TF $t$ or peak $p$ accessibility.
TRIPOD screens for such interaction effects through a "conditional level 2 test", which
assesses the association between $\Delta Y_g$ and the product of the matched variable with the
difference in the unmatched variable, after taking partial residuals on the difference in the
unmatched variable. In summary, TRIPOD categorizes each identified trio relationship as
supported by marginal association, association between peak and gene conditioned on
TF expression, and/or association between TF and gene conditioned on peak
accessibility. The conditional relationships are further categorized to level 1 or level 2,
with level 2 indicative of a stronger relationship exhibiting multiplicative interaction effects
between TF expression and peak accessibility.

For significant trios, TRIPOD further carries out a sampling-based influence
analysis, where phenotypically contiguous sets of metacells are held out to measure their
influence on the estimated coefficients. The corresponding cell types/states that lead to
significant deviations from the null upon their removal have high influence scores, which
can be used to identify cell types/states that drive a regulatory relationship.

To highlight the differences between TRIPOD and existing methods based on
marginal associations, we show two canonical examples (Supplementary Fig. 5) where
the two approaches disagree. Fig. 2c outlines a significant trio detected by TRIPOD's
level 2 testing, yet the marginal peak-gene and TF-gene associations were insignificant.
It turns out that a subset of cells with high peak accessibility $\{X_p\}$ have close-to-zero TF
expressions $\{Y_t\}$, and, meanwhile, another subset of cells with high TF expressions $\{Y_t\}$
have close-to-zero peak accessibilities $\{X_p\}$. In these cells, either the peak is closed, or
the TF is not expressed, and this leads to the target gene not being expressed, which
masks the marginal associations. The high peak accessibility and TF expression in these
cells, which act through other regulatory trios, cancel out when we consider the interaction
$\{X_p \times Y_t\}$, leading to a significant interaction term detected by TRIPOD. Conversely, Fig.
2d outlines another trio, whose marginal associations were significant, yet TRIPOD did
not detect significant conditional associations from either level 1 or level 2 testing. In this
case, with almost constant TF expression, the large difference in peak accessibility leads
to a small difference in target gene expression. Meanwhile, the cells that drive the
significantly positive correlation between $\{Y_g\}$ and $\{Y_t\}$ have almost zero values for $\{X_p\}$.

Both observations suggest that this peak has little to do with the regulation of the target

gene *FGL2* by this specific TF MAFK. Notably, we do not claim that the significantly linked

peaks and TFs through marginal association are false positives, but rather this specific

trio is insignificant (i.e., the peak and TF may act through other TF and peak, respectively).

In summary, TRIPOD puts peak-TF-gene trios into one unified model, complementing

existing methods based on marginal associations and allowing for simultaneous

identification of all three factors and prioritization of a different set of regulatory

relationships.

**TRIPOD identifies three-way regulatory relationships in PBMCs with orthogonal**

**validations.** We first applied TRIPOD to identify regulatory trios in the 10k PBMC dataset.

Cell-type labels for this dataset were transferred from a recently released CITE-seq

reference of 162,000 PBMC cells measured with 228 antibodies[14]. After quality control,

we kept 7790 cells from 14 cell types pooled into 80 metacells, 103,755 peaks, 14,508

genes, and 342 TFs; the UMAP reduced dimensions are shown in Supplementary Fig.

1a. Distribution of the number of peaks 100kb/200kb upstream and downstream of the

TSS per gene, as well as distribution of the number of motifs per peak, are shown in

Supplementary Fig. 6.

As a proof of concept, we first illustrate two trios where the frameworks agree,

identified by level 1 conditional testing (regulation of *CCR7* by LEF1; Fig. 3a) and level 2

interaction testing (regulation of *GNLY* by TBX21; Fig. 3b). From the influence analyses,

TRIPOD identified B and T cells as the cell types where LEF1 regulates *CCR7*, and

natural killer (NK) cells as the cell types where TBX21 regulates *GNLY*. These cell type-

specific regulatory relationships are corroborated by motif's deviation scores using

chromVAR[23] (Fig. 3) and the enrichment of Tn5 integration events in the flanking regions

using DNA footprinting analyses[12] (Supplementary Fig. 7e). Unlike chromVar and DNA

footprinting analyses, which only give genome-wide average enrichments, TRIPOD

significantly enhances the resolution by identifying the specific *cis*-regulatory regions that

the TFs bind for the regulation of target genes.

Results from TRIPOD and marginal association tests overlap but, as expected,

exhibit substantial differences (Supplementary Fig. 8). The previous section showed

example trios where the two frameworks disagree. Additionally, results from TRIPOD's

matching scheme and those from random matching also overlap but exhibit substantial

differences, both on the global scale (Supplementary Fig. 9) and for each gene

(Supplementary Fig. 10). Notably, for the two counterexamples discussed in the previous

section, random matching could not identify the masked positive trio in Fig. 2c, yet it

retained significance for the negative trio shown in Fig. 2d, in a similar fashion to marginal

testing (Supplementary Fig. 11). Genome-wide $p$-value distributions from TRIPOD's two

levels of testing under the null with permuted peak accessibility and TF expression are

shown in Supplementary Fig. 12, indicating that TRIPOD's framework has good type I

error control. A master output of significant associations with Bonferroni correction is

shown in Supplementary Table 2, with scatterplots and pairwise correlations of genome-

wide *p*-values from different testing schemes shown in Supplementary Fig. 13.

To our best knowledge, no experimental technique can directly validate three-way

regulatory relationships at high resolution with high throughput. Therefore, we performed

validation and benchmarking by harnessing existing databases and orthogonal

sequencing experiments that interrogate each pairwise relationship among the three

factors (Table 1). The rationale is that true regulatory relationships should show

enrichment in all three marginal relationships. Fig. 4a illustrates the extensive validation

strategies that were undertaken.

First, to validate the *cis*-linkage between peak region and target gene, we used the

enhancer databases of blood and non-cancerous cells from FANTOM5[24] (from HACER[25]),

4DGenome[26] (from HACER[25]), and EnhancerAtlas 2.0[27], as well as *cis*-eQTLs in the

whole blood reported by the GTEx consortium[9]. We collapsed TRIPOD's trio calls into

peak-gene relationships and benchmarked against Signac's LinkPeaks[12] on single cells

and marginal association testing on metacells; for each target gene, we performed a

hypergeometric test for enrichment of the peak-gene linkages in the regulatory databases

and annotations (see Methods for details). For all four databases, TRIPOD's $p$-values for

enrichment are substantially significant (Fig. 4b). When stratified by the different levels of

testing, TRIPOD's level 1 and level 2 conditional testing returns more significant

enrichment compared to linkPeaks and marginal associations; the most significant

enrichment is from level 1 testing matching by TF expression, which is expected since

the "gold-standard" peak-gene relationship is directly captured by the level 1 testing
without TF interaction (Supplementary Fig. 14a). Additionally, the unique sets of trio
regulatory relationships identified by TRIPOD but not by random matching (which results
in only marginally associated linkages) have significant enrichment, demonstrating the
effectiveness of TRIPOD in identifying true trio relationships that complement existing
methods based on marginal association testing (Supplementary Fig. 14b).

Second, to validate the TF-gene edge in the TRIPOD-identified trios, we referred
to knockTF[7], a TF knockdown/knockout gene expression database, and hTFtarget[28], a
database of known TF regulatory targets. Specifically, in knockTF, we found seven TF
knockdown/knockout RNA-seq experiments in the peripheral blood category. For these
TFs, we identified significantly linked genes by marginal association and by TRIPOD and
found TRIPOD's results to have significantly higher precision and recall (Fig. 4c); the
improvement is robust to varying FDR thresholds (Supplementary Table 3). For hTFtarget,
we obtained, for each highly variable gene, its blood-specific TFs, and calculated the
gene-specific precision-recall rates – TRIPOD is more sensitive compared to marginal
association testing, although both suffered from inflated "false positives," which can also
be due to the low sensitivity in the *in silico* calls by hTFtarget (Fig. 4d). Precision and
recall rates with varying significance levels further confirm that TRIPOD has better
agreement with existing TF knockdown/knockout data, in comparison to marginal
association testing (Supplementary Fig. 15).

Third, to validate the TF-peak edge representing TF binding to peak regions, in
addition to the DNA footprinting analysis shown in Supplementary Fig. 7e, we
downloaded from the Cistrome portal[29] non-cancerous ChIP-seq data from sorted human
blood cells (B lymphocyte, T lymphocyte, and monocyte (Supplementary Table 4). The
peaks identified by TRIPOD had a substantially higher percentage of overlap with the
ChIP-seq peaks compared to the genome-wide baseline; TRIPOD's performance is better
than or on par with that from testing of marginal associations (Fig. 4e). Since ChIP-seq
peaks reflect only TF binding, without consideration for the gene target of regulation, it is
expected that it agrees well with marginal association test results, which are capturing
such a universal relationship.

In summary, existing databases and public data of different types from a wide

range of studies extensively support each of the three pairwise links in the trios reported

by TRIPOD, demonstrating its effectiveness in uncovering true regulatory relationships.

**TRIPOD identifies known and novel putative regulatory relationships during mouse**

**embryonic brain development.** We next applied TRIPOD to single-cell multiomic data

of 5k mouse embryonic brain cells at day 18 by 10X Genomics. The cell type labels were

transferred from an independent scRNA-seq reference[30] using SAVERCAT[31]. We kept

3,962 cells that had consistent transferred labels from seven major cell types: radial glia,

neuroblast, GABAergic neuron, glutamatergic neuron, glioblast, oligodendrocyte, and

Cajal−Retzius neuron (Supplementary Fig. 1b). We applied TRIPOD to 633 TFs, 1000

highly variable genes, and ATAC peaks 200kb up/downstream of the genes' TSSs.

On the genome-wide scale, the union of TRIPOD's level 1 and 2 tests gave a larger

number of unique peak-gene pairs and TF-gene pairs than LinkPeaks[12] and marginal

association testing, respectively (Supplementary Fig. 16a). To evaluate these results, we

first examined whether the peak-gene links were enriched in previously reported

enhancer-promoter chromatin contacts using PLAC-seq data of mouse fetal brain[32]

(Table 1, Supplementary Fig. 16b). We observed that the regulatory links detected by

both marginal association and TRIPOD showed significant enrichment in PLAC-seq

contacts (Supplementary Fig. 16b). Importantly, TRIPOD detected sets of peak-gene

pairs from trio relationships that were overlapping but distinct from the sets obtained by

marginal association, and a substantial fraction of the links identified by TRIPOD but not

by the marginal method were validated by PLAC-seq (Fig. 5a; Supplementary Fig. 16c).

This suggests that TRIPOD identifies real regulatory relationships that complement those

detected by existing methods.

We also note that the type of evidence that supports a regulatory relationship

matters when compared to other types of experimental data. For example, PLAC-seq

measures, for a fixed TF, the degree of promoter contacts in the TF-binding domains.

Conceptually, the closest analog to this measurement in our model is level 1 association,

conditioned on TF expression, between the motif-containing peak region and target gene

expression. Thus, it is not surprising that this level 1 test matching by TF gives the most

significant enrichment (Supplementary Fig. 16b, 17a, 18a). However, detection by
TRIPOD is pre-conditioned on the expression of the target gene at a high enough level,
which is irrelevant to the PLAC-seq data. Thus, not all detections made by PLAC-seq are
expected to be found by TRIPOD.

To validate the links between TFs and peaks, we used publicly available ChIP-seq
data for Olig2[33], Neurog2[34], Eomes[34], and Tbr1[35], TFs that play key roles in embryonic
brain development (Table 1). The Olig2 ChIP-seq data were generated in three types of
rat cells, oligodendrocyte precursor cells (OPC), immature oligodendrocytes (iOL), and
mature oligodendrocytes (mOL), while the Neurog2, Eomes, and Tbr1 ChIP-seq data
were generated in mouse embryonic cerebral cortices (see Methods for details). When
TF expression was matched, TF binding peaks identified by TRIPOD level 1 tests were
significantly enriched in the TF ChIP-seq peaks across all datasets except for the Olig2
ChIP-seq data of mature oligodendrocytes (mOL), which served as a negative control
and had a substantially lower degree of enrichment (Supplementary Fig. 17, 18).
Importantly, TRIPOD detected a substantial number of peak-TF pairs that were not
detected through marginal associations but validated by ChIP-seq (Fig. 5b).

The validations and global benchmarking demonstrate TRIPOD's effectiveness in
finding real regulatory relationships. Next, we focused on a set of TFs known to play
essential roles during mouse embryonic brain development. Specifically, we chose Pax6,
Neurog2, Eomes, Neurod1, and Tbr1, major TFs mediating glutamatergic neurogenesis[36],
and Olig2, Sox10, Nkx2-2, Sox9, Nfia, and Ascl1, which initiate and mediate gliogenesis[37];
the known regulatory cascades are shown in Fig. 5c. Here, the up and downstream TFs
in a link are used as the TF and the target gene in TRIPOD's analysis, respectively, and
we established a link if at least one of the TRIPOD tests returned a positive coefficient
estimate with FDR-adjusted *p*-values less than 0.01 for at least one trio involving the pair
of the TF and the target gene. TRIPOD's level 1 and level 2 testing successfully captured
five out of the seven known regulatory links (Fig. 5c, d, Supplementary Fig. 19, 20);
interestingly, TRIPOD's results also suggest substantial crosstalk between the two
cascades, where neurogenesis-specific TFs activate gliogenesis-specific TFs (Fig. 5c, d).
ChIP-seq data of Neurog2, Eomes, and Tbr1 supported four of the crosstalk links:
regulation of *Sox9* by Neurog2 and regulation of *Nfia* by Neurog2, Eomes, and Tbr1,

respectively (Supplementary Fig. 21). These crosstalk links that were validated by ChIP-

seq were also captured by conditional associations; two of them were captured by

marginal associations (Supplementary Fig. 19). Thus, we think it is highly plausible that

neurogenesis TFs activate gliogenesis genes at day 18 of embryonic mouse brain

development, which is exactly when the switch is being made from neurogenesis to

gliogenesis. To our best knowledge, these possible links between neurogenesis and

gliogenesis pathways have not been systematically explored and thus warrant future

investigation. Finally, for each of the neurogenesis and gliogenesis TFs, we performed a

gene ontology (GO) analysis of their significantly linked target genes using DAVID[38]; the

enriched terms were largely consistent with the regulatory functions of the TFs during

neurogenesis and gliogenesis (Fig. 5e). Specifically, the mouse embryonic brain cells are

collected during the transition phase between neurogenesis and gliogenesis, and the

enriched terms contain oligodendrocyte differentiation and regulation of neuron

differentiation, confirming TRIPOD's calling results. Other terms, such as regulation of

transcription and cell cycle, are enriched due to the transcriptional regulatory role of the

TFs.

So far, we have taken advantage of the cross-cell-type variation to identify the trio

regulatory relationships. To dissect cell-type-specific regulation, we next applied the

influence analysis framework (see Methods for details) to the significant trios involving

neurogenesis and gliogenesis TFs. For a given TF, the number of trios, for which a given

cell type was influential (FDR < 0.01), is summarized in Fig. 5f, with details for specific

example trios given in Supplementary Fig. 22. The analyses underpinned the cell types

in which the transcriptional regulation was active, and, reassuringly, the neurogenesis

and gliogenesis TFs have the most regulatory influence in neuroblasts and glioblasts,

respectively. Additionally, Ascl1 is active in GABAergic neurons in addition to neuroblasts

and glioblasts, consistent with its role as a GABAergic fate determinant[39]. Notably, the

highly influential cell types that lead to the significant trios involving several neurogenesis-

specific TFs include not only neuroblast but also glioblast, supporting our previous

findings on the crosstalk between the two cascades. Notably, these results are unlikely

due to the given TFs being overexpressed in the corresponding highly influential cell types,

since the influential cell types were not the same as the cell types where the TFs were

highly expressed (Fig. 5f, Supplementary Fig. 22, 23). Overall, TRIPOD allows fine
characterization of cell-type- and cell-state-specific functions of the TFs during
neurogenesis and gliogenesis.

Using this dataset, we further examined how varying window sizes and different
resolutions/constructions of metacells affect the model fitting results; this led to the
following observations. First, incorporating peaks 100kb/200kb up/downstream of genes'
TSSs leads to consistent and significant enrichment of validated gene-peak pairs by
PLAC-seq and peak-TF pairs by ChIP-seq, while narrowing the window size down to 50kb
decreased the degree of enrichment (Supplementary Fig. 17). Second, the validation
results were robust to changes in resolutions of the metacells (Supplementary Fig. 18),
since TRIPOD does not require the metacells to truly represent distinct and non-
overlapping segments of the transcriptome space.

**TRIPOD infers lineage-specific regulatory relationships in differentiating mouse**
**hair follicle cells.** As a last example, we applied TRIPOD to SHARE-seq[13] data
(Supplementary Fig. 1c) of mouse hair follicle cells, consisting of four broadly defined cell
types – transit-amplifying cells (TAC), inner root sheath (IRS), hair shaft, and medulla
cells – along a differentiation trajectory. The cell-type labels were curated based on
marker genes, TF motifs, and ATAC peaks from the original publication[13]; pseudotime
was inferred using Palantir[40] and overlaid on the cisTopic[41] reduced dimensions of the
ATAC domain. Cells were partitioned using both the pseudotime and the UMAP
coordinates to construct metacells (Fig. 6a). Due to the low RNA coverage
(Supplementary Fig. 4), we focused on 222 highly-expressed TFs, 794 highly expressed
genes reported to have more than ten linked *cis*-regulatory peaks[13], and peaks 100kb
up/downstream of the genes' TSSs.

For validation, we used H3K4me1 and H3K27ac ChIP-seq data from an isolated
mouse TAC population[42] (Table 1). H3K4me1 and H3K27ac are markers for poised and
active enhancers, respectively, and were used to benchmark TRIPOD's linked peaks
against previously reported domains of regulatory chromatin (DORCs)[13], as well as
randomly sampled peaks. The linked peaks by TRIPOD had higher scores for both
H3K4me1 and H3K27ac than DORCs, the latter identified through marginal associations

(Fig. 6b). To further validate the regulatory effects of the linked peaks, we obtained

previously characterized super-enhancers (SEs) in mouse TACs[42]. Target genes of the

381 SEs were assigned based on the gene's proximity to the SE, as well as the correlation

between loss of the SE and loss of the gene transcription[42]. TRIPOD was able to

successfully recapitulate the SE regions for the genes considered, with four examples

shown in Fig. 6c, where significantly linked peaks mostly resided in the SEs.

To demonstrate, Fig. 6d shows regulatory trios that are specific to the IRS lineage,

the hair shaft lineage, and the medulla lineage (Supplementary Fig. 24). These trios also

showed significant pairwise marginal associations (Fig. 6e), lending confidence that they

are real. The cell types where the regulation happens were identified by influence analysis,

for which the $p$-values were smoothed along the differentiation trajectory and overlaid on

the UMAP embedding (Fig. 6f). DNA footprinting analyses surveyed the enrichment of

Tn5 integration events surrounding the corresponding motif sites and showed cell-type-

specific enrichment (Fig. 6g), corroborating TRIPOD's results.

**Discussion**

We have considered the detection of regulatory trios, consisting of a TF binding to a

regulatory region to activate or repress the transcription of a nearby gene, using single-

cell RNA and ATAC multiomic sequencing data. The presented method, TRIPOD, is a

new nonparametric approach that goes beyond marginal relationships to detect

conditional associations and interactions on peak-TF-gene trios. We applied TRIPOD to

three single-cell multiomic datasets from different species and protocols with extensive

validations and benchmarks. We started our analyses with predicting gene expression

from both peak accessibility and TF expression. Supervised frameworks have been

proposed to predict gene expression from DNA accessibility[43], and vice versa[44], using

matched bulk transcriptomic and epigenomic sequencing data. Blatti *et al.*[45] showed that

joint analysis of DNA accessibility, gene expression, and TF motif binding specificity

allows reasonably good prediction of TF binding as measured by ChIP-seq. However,

none of these methods incorporate TF expression. By selecting peaks near the genes'

TSSs and TFs with high motif scores in the selected peaks, we constructed biologically

meaningful peak-TF pairs as predictors and showed that such a mechanistic model
significantly boosts the prediction accuracy of gene expression.

We next considered the detection and significance assessment for individual peak-
TF-gene trios, comprehensively comparing our detections with those made by tissue- and
cell-type-matched PLAC-seq and ChIP-seq experiments, by *cis*-eQTL and TF
knockdown/knockout studies, and by those recorded in the main enhancer databases.
The comparisons show that TRIPOD detections are substantially enriched for overlap
with all of these experiments, and in most cases, improve upon the overlap achieved by
existing methods. It is important to note that the recall rates in the comparisons to these
experiments should only be interpreted as relative metrics and not as absolute measures
of sensitivity. That is because each experiment measures a biological relationship that is
associated but different from what we aim to recover from TRIPOD. For example, ChIP-
seq aims to capture all locations where the TF binds, regardless of which gene it is
affecting, while TRIPOD aims to recover specific TF, enhancer, target gene trios.
KnockTF and hTFtarget, on the other hand, aims to identify all genes whose expressions
change when a TF is knocked out/down, which may not be genes that the TF directly
regulates through binding. An experiment that perhaps comes closest to measuring what
TRIPOD detects is PLAC-seq, which quantifies chromatin contacts anchored at genomic
regions bound by specific proteins. In addition to ChIP-seq, we used PLAC-seq data to
corroborate TRIPOD detections for the embryonic mouse brain data in Fig. 5a,
Supplementary Fig. 16b, 17a, 18a. Here, the overlap is also far from 100%, as TRIPOD
can only detect a PLAC-seq relationship if the expression of the target gene is high
enough. Also, PLAC-seq cannot detect TRIPOD relationships unless the *cis*-region in
question comes into direct contact with the promoter, which is not the only mechanism of
gene regulation. For example, TF binding may change the local chromatin conformation
as an insulator or may help recruit the binding of other TFs. Thus, it is expected that
TRIPOD only recovers a small fraction of the signals identified by these experiments. For
this reason, we choose to use the word "recall" rather than "sensitivity," as we are using
it as a metric of enrichment rather than as a measure of true positive rate.

Our current study is limited in several ways. A study in *Drosophila*[46] modeled motif
binding specificities and chromatin accessibilities in bulk RNA and ATAC sequencing data

to predict the cooperative binding of pairs of TFs, using *in vitro* protein-protein binding

experiments for validation. The detection of synergies between multiple TFs and peaks

on the genome-wide scale and in a cell-type-specific manner needs further investigation.

Additionally, while we have not differentiated between positive and negative regulation,

TRIPOD reports both types of relationships and categorizes them by sign. While we

describe the trios with a positive sign to be enhancers, it is not clear how to interpret the

trios with negative signs, the latter having lower overlap with other benchmarking datasets.

Transcription activation and repression have been active research areas in biology, with

a lot yet unknown[47]. TRIPOD's results provide potential targets for experimental follow-

up and detailed characterization.

TRIPOD uses cell matching as a nonparametric method of computing conditional

associations. One could, conceptually, match on more cell-level attributes in addition to

transcription factor expression or peak level accessibility. For example, to recover true

causal relationships, it seems tempting to match on more potential confounders, such as

cell type. However, one should be careful in matching by additional covariates such as

inferred cell type labels, as this could also reduce the signal. For example, condition-

specific regulation signals that are shared across multiple (but not all) cell types would be

much reduced if we were to match on cell type. For specificity, TRIPOD relies on the

careful curation of inputs to the regression (using only peaks that contain the TF motif

and are close to the target gene), rather than matching on all possible confounders.

Our analysis focused on three datasets where the RNA and ATAC modalities have

sufficient depths of coverage. For the SHARE-seq data, the sequencing depth for RNA is

very low, and thus we focused only on highly expressed genes and TFs (Fig. 6). For

SNARE-seq data, whose coverage in both modalities is even lower, we focused on

prediction models and not trio detection, where we saw only marginal improvement

beyond existing methods[20] (Supplementary Fig. 2b). For data where the coverage is even

lower, e.g., PAIRED-seq, cross-modality metacells could not be stably formed, making

such analyses impossible (Supplementary Table 1, Supplementary Fig. 4). With rapidly

increasing sequencing capacity and technological advancement, TRIPOD, applied to

more cells sequenced at higher depth, can uncover novel regulatory relationships at a

finer resolution. With increased data resolution and cell numbers, it would then be

meaningful to explore beyond the three-way relationships characterized by TRIPOD to
include higher-order models that can more realistically capture the complex regulatory
relationships between enhancers, modules consisting of multiple transcription factors,
and the transcription of the target gene.
549
**Methods**
**Data input and construction of metacells.** Denote $X_{ip}$ as the peak accessibility for peak
$p$ $(1 \leq p \leq P)$ in cell $i$ $(1 \leq i \leq N)$, $Y_{ig}$ as the gene expression for gene $g$ $(1 \leq g \leq G)$,
and $Y_{it}$ as the TF expression for TF $t$ $(1 \leq t \leq T)$. The TF expression matrix is a subset
of the gene expression matrix, and for single-cell multiomic data, the cell entries are
matched. To mitigate the effect of ATAC sparsity[48] and RNA expression stochasticity[49],
as a first step, TRIPOD performs cell-wise smoothing by pooling similar cells into
"metacells." This, by default, is performed using the weighted-nearest neighbor method
by Seurat V4[14] to jointly reduce dimension and identify cell clusters/states across different
modalities. In practice, the metacells can also be inferred using one modality – for
example, RNA may better separate the different cell types[30], and in other cases,
chromatin accessibility may prime cells for differentiation[13]. For data normalization, we
use sctransform[50] and TF-IDF[11] for scRNA-seq and scATAC-seq, respectively, followed
by dimension reduction and clustering[12]. To account for peaks overlapping with other
genes (Supplementary Fig. 6b), TRIPOD has the option to either remove the overlapped
peaks or to adjust the peak accessibilities by the expressions of the overlapped genes, in
a similar fashion to MAESTRO[18]. To reconstruct the RNA and ATAC features for the
metacells, we take the sum of the integer-valued ATAC and RNA read counts across cells
belonging to the metacells; library size is adjusted for both the RNA and ATAC domain
by dividing all counts by a metacell-specific size factor (total read counts divided by $10^6$).
For the analyses presented in the manuscript, position frequency matrices (PFM)
were by default obtained from the JASPAR database[51], and we used 633 and 107 pairs
of TFs and motifs annotated in human and mouse, respectively. TRIPOD provides an
option to use a more comprehensive set of motif annotations from the HOCOMOCO[52]
database. TRIPOD also allows for a binding motif to be shared across multiple TFs, as
well as user-defined and/or de novo motifs. We additionally examined the effects of

combining the accessibilities of ATAC peaks containing the TF binding sites within the

window centered at the gene's TSS and using the combined accessibility as input; we did

not observe an improvement in model performance (Supplementary Fig. 25).

**RNA prediction by TF expression and peak accessibility.** To predict RNA from ATAC,

Signac[12] and Cicero[17] take the sum of peak accessibilities in gene bodies and promoter

regions to construct a pseudo-gene activity matrix: $\hat{Y}_{ig} = \sum_{p \in E_g} X_{ip}$, where $E_g$ is the set of

peaks within gene bodies and upstream regions of TSSs. Instead of directly taking the

sum, MAESTRO[18] adopts a "regulatory potential" model by taking the weighted sum of

accessibilities across all nearby peaks: $\hat{Y}_{ig} = \sum_{p \in E_g} w_p^g X_{ip}$, with weights $\{w_p^g\}$ pre-

calculated based on existing gene annotations. Specifically, the method weighs peaks by

exponential decay from TSS, sums all peaks on the given gene exons as if they are on

the TSS, normalizes the sum by total exon lengths, and excludes the peaks from

promoters and exons of nearby genes. The strategy to take the unweighted/weighted sum

of accessibility as a proxy for expression has been adopted to align the RNA and ATAC

modalities when scRNA-seq and scATAC-seq are sequenced in parallel from the same

cell population but not the same cells[11]. For single-cell multiomic data, sci-CAR[19] performs

feature selection to identify *cis*-linked peaks via a LASSO regression: $Y_{ig} \sim \sum_{p \in E_g} \beta_p^g X_{ip}$,

where an L1 regularization is imposed on $\beta_p^g$. Compared to MAESTRO, which pre-fixes

the weights $\{w_p^g\}$, $\{\beta_p^g\}$ are estimated from the data by regressing RNA against matched

ATAC data. What we propose is a feature selection model involving both peak

accessibility and TF expression: $Y_{ig} \sim \sum_{p \in E_g} \sum_{t \in f_p} \beta_{pt}^g X_{ip} Y_{it}$, where $f_p$ contains the set of

TFs with high-scoring binding motifs in peak $p$ inferred from the JASPAR database[51].

**TRIPOD model and trio regulatory relationship.** For a given target gene $g$, a peak $p$

within a window centered at the gene's TSS, and a TF $t$ whose binding motif is high-

scoring in the peak, TRIPOD infers the relationship between a regulatory trio $(p, t, g)$.

TRIPOD focuses on one trio at a time and goes beyond the marginal associations to

characterize the function $Y_g = f(X_p, Y_t)$. In what follows, we first describe TRIPOD's

matching-based nonparametric approach and then describe a linear parametric approach,

followed by a discussion on the connections and contrasts between the two approaches.

For each cell $i$ whose TF expression is above a threshold $\delta$ (we only carry out

testing in cells that express the TF), we carry out a minimum distance pairwise cross-

match based on $\{Y_{it}|Y_{it} > \delta\}$. Let $\{(i_j, i_{j^*})\}$ be the optimal matching, after throwing away

those pairs that have $|Y_{i_j t} - Y_{i_{j^*} t}| > e$. For each pair $j$, $i_j$ and $i_{j^*}$ are two metacells with

matched TF expression, for which we now observe two, possibly different, values

$\{X_{i_j p}, X_{i_{j^*} p}\}$ for peak $p$, as well as two corresponding values $\{Y_{i_j g}, Y_{i_{j^*} g}\}$ for gene $g$. We

then compute the following auxiliary differentials within each pair:

$$\Delta X_{jp} = X_{i_j p} - X_{i_{j^*} p},$$

$$\Delta Y_{jg} = Y_{i_j g} - Y_{i_{j^*} g},$$

616 as well as

$$\bar{Y}_{jt} = (Y_{i_j t} + Y_{i_{j^*} t})/2.$$

For level 1 testing of conditional association, we estimate $\hat{r}_p^g = \rho(\Delta X_{jp}, \Delta Y_{jg})$, where $\rho$ is

Spearman correlation, and test $H_1 : r_p^g = 0$. For level 2 testing of interaction, we perform a

regression $\Delta Y_{jg} = \alpha \Delta X_{jp} + \gamma \bar{Y}_{jt} \times \Delta X_{jp}$, set $\hat{\gamma}_{pt}$ to be the least-squares solution for $\gamma$, and

test $H_2 : \gamma_{pt} = 0$. For visualization of the model fitting, we take the partial residuals of $\Delta Y_{jg}$

and $\bar{Y}_{jt} \times \Delta X_{jp}$ on $\Delta X_{jp}$, respectively. Note that even though TF expression is not included

in the model as a main term, it is controlled for (and not just in the linear sense) by the

matching. Similarly, we can also perform this procedure matching by peak accessibility.

As a summary, for level 1 testing of conditional association, we have:

$$\text{Match by } Y_t, \ \alpha = \rho(\Delta Y_g, \Delta X_p),$$

$$\text{Match by } X_p, \ \beta = \rho(\Delta Y_g, \Delta Y_t).$$

628 For level 2 testing of (TF expression)×(peak accessibility) interaction effects, we have:

$$\text{Match by } Y_t, \ \Delta Y_g = \alpha^* \Delta X_p + \gamma_1(\bar{Y}_t \times \Delta X_p),$$

$$\text{Match by } X_p, \ \Delta Y_g = \beta^* \Delta Y_t + \gamma_2(\bar{X}_p \times \Delta Y_t).$$

631 To test for the conditional associations and interactions, we can also apply a

632 parametric method, such as multiple linear regression:

$$Y_g = \mu + \alpha_L X_p + \beta_L Y_t,$$

634
$$Y_g = \mu + \alpha_L^* X_p + \beta_L^* Y_t + \gamma_L X_p Y_t.$$

See Supplementary Fig. 26 for linear testing results for trios shown in Fig. 3 and Fig. 6.
The estimated coefficients from the nonparametric and parametric methods are
correlated on the global scale (Supplementary Fig. 13, 27), and their interpretations are
similar: $\alpha$ and $\alpha_L$ estimate the change in gene expression per change in peak
accessibility, fixing TF expression; $\beta$ and $\beta_L$ estimate the change in gene expression per
change in TF expression, fixing peak accessibility; $\gamma_1$ and $\gamma_L$ measure how the change in
gene expression per change in peak accessibility at each fixed TF expression relies on
the TF expression; $\gamma_2$ and $\gamma_L$ measure how the change in gene expression per change in
TF expression at each fixed peak accessibility relies on the peak accessibility. However,
the underlying models and assumptions are different. Matching controls for not just the
linear variation in the matched variable, but also any nonlinear variation. This contrasts
with adding the variable as a covariate in the linear regression, where we simply remove
linear dependence. The main motivation for using the matching model above is our
reluctance to assume the simple linear relationship. Additionally, we use the rank-based
Spearman correlation, which will not be driven by outliers – a "bulk" association between
ranks is needed for significance. Thus, the nonparametric model of TRIPOD is more
stringent (Supplementary Fig. 28) and more robust to outliers.

**Identifying regulatory cell type(s) and cell state(s)**. For the significant trios detected
by TRIPOD, we next seek to identify the underlying regulatory cell type(s). Specifically,
we carry out a cell-type-specific influence analysis to identify cell types that are highly
influential in driving the significance of the trio. Traditional approaches (e.g., the Cook's
distance and the DFFITs) delete observations one at a time, refit the model on remaining
observations, and measure the difference in the predicted value from the full model and
that from when the point is left out. While they can be readily applied to detect "influential"
metacells one at a time (Supplementary Fig. 7a,b), these methods do not adjust for the
degree of freedom properly when deleting different numbers of metacells from different
cell types. That is, they do not account for the different numbers of observations that are
simultaneously deleted. Additionally, both methods adopt a thresholding approach to
determine significance, without returning $p$-values that are necessary for multiple testing

correction. We, therefore, develop a sampling-based approach to directly test for the

influence of multiple metacells and to return $p$-values (Supplementary Fig. 7c).

Here, we focus on the linear model for its ease of computation: $\hat{Y}_g = \hat{\mu} + \hat{\alpha}X_p +$

$\hat{\beta}Y_t + \hat{\gamma}X_pY_t.$ Given a set of observations $I = \{i: i\text{th metacell belongs to a cell type}\},$ we

remove these metacells, fit the regression model, and make predictions: $\hat{Y}_g^{(I)} = \hat{\mu}^{(I)} +$

$\hat{\alpha}^{(I)}X_p + \hat{\beta}^{(I)}Y_t + \hat{\gamma}^{(I)}X_pY_t.$ The test statistics are the difference in the fitted gene

expressions $\left|\hat{Y}_g - \hat{Y}_g^{(I)}\right|.$ We generate the null distribution via sampling. Specifically, within

each sampling iteration, we sample without replacement the same number of metacells,

denoted as a set of $I^*$, delete these observations, and refit the regression model on the

remaining observations: $\hat{Y}_g^{(I^*)} = \hat{\mu}^{(I^*)} + \hat{\alpha}^{(I^*)}X_p + \hat{\beta}^{(I^*)}Y_t + \hat{\gamma}^{(I^*)}X_pY_t.$ The $p$-value is

computed across $K$ sampling iterations as $p_{Y_g} = \sum_{I^*} 1\left(\sum|\hat{Y}_g - \hat{Y}_g^{(I)}| \geq \sum|\hat{Y}_g - \hat{Y}_g^{(I^*)}|\right)/K,$

where 1() is the indicator function. In addition to testing each cell type separately, the

framework can be extended to test for the influence of cell-type groups. For example, in

Fig. 3, we reconstruct the cell-type hierarchy using expression levels of highly variable

genes from the RNA domain and carry out the aforementioned testing scheme at each

split for its descendent cell types in the hierarchical structure.

For transient cell states, TRIPOD first identifies the neighbors of each metacell

along the trajectory and then carries out metacell-specific testing by simultaneously

removing each metacell and its neighbors using the framework described above. The

resulting $p$-values are, therefore, smoothed and can be visualized in the UMAP plot (Fig.

6f and Supplementary Fig. 22) to identify the underlying branches/segments that are key

in defining the significant regulatory trio. This approach can be directly applied to cells

with branching dynamics without the need to isolate cell subsets or to identify cell types.

**Validation resources and strategies.** Resources for validating the trio regulatory

relationships are summarized in Table 1. To validate the peak-gene relationships, we

referred to existing enhancer databases: FANTOM5[24] links enhancers and genes based

on enhancer RNA expression; 4DGenome[26] links enhancers and genes based on

physical interactions using chromatin-looping data including 3C, 4C, 5C, ChIA-PET, and

Hi-C; EnhancerAtlas 2.0[27] reports enhancers using 12 high-throughput experimental

methods including H3K4me1/H3K27ac ChIP-seq, Dnase-seq, ATAC-seq, and GRO-seq.

We only focused on blood and non-cancerous cells from these databases (Fig. 4b). A list

of *cis*-eQTLs within the whole blood mapped in European-American subjects was

downloaded from the GTEx consortium[9] (Fig. 4b). For the mouse embryonic brain dataset,

we additionally adopted H3K4me3-mediated PLAC-seq data[32], which reported enhancer-

promoter chromatin contacts mapped in mouse fetal forebrain (Fig. 5a, Supplementary

Fig. 16b, 17a, 18a). For the mouse skin dataset, we adopted TAC-specific ChIP-seq data

of H3K4me1 and H3K27ac[42], markers for poised and active enhancers, respectively (Fig.

6b); we also obtained previously reported super-enhancers in mouse TACs from *in vivo*

studies[42] (Fig. 6c). Genomic coordinates were lifted over from mm9 to mm10 when

necessary.

To validate the TF-gene relationships in the PBMC data, we utilized the knockTF[7]

and the hTFtarget[28] databases. knockTF interrogates the changes in gene expression

profiles in TF knockdown/knockout experiments to link the TFs to their target genes in a

tissue- or cell-type-specific manner. We downloaded 12 experiments, corresponding to

12 TFs (BCL11A, ELK1, GATA3, JUN, MAF, MYB, NFATC3, NFKB1, STAT3, STAT6,

TAL1, and ZNF148) in the peripheral blood category, and focused on seven TFs that

have at least one linked gene by any model benchmarked (Fig. 4c; Supplementary Table

3). hTFtarget computationally predicts TF-gene relationships using ChIP-seq data, and

we manually downloaded the TFs associated with each of the top 100 highly variable

genes in the blood tissue (Fig. 4d; Supplementary Fig. 15).

For peak enrichment analysis compared to the existing enhancers, *cis*-eQTLs, and

enhancer-promoter contacts, we carried out a hypergeometric test as follows. Let $k$ be

the number of significantly linked peaks, $q$ be the number of significantly linked peaks that

overlap with annotations (e.g., annotated enhancers), $m$ be the number of peaks that

overlap with the annotations, and $n$ be the number of peaks that do not overlap with

annotations. The $p$-value of enrichment is derived from the hypergeometric distribution

using the cumulative distribution function, coded as phyper(q, m, n, k, lower.tail=F) in R.

We used this hypothesis testing framework to validate and benchmark the reported peak-

gene links, with results shown in Fig. 4b.

To validate the peak-TF relationships, we downloaded non-cancerous cell-type-
specific ChIP-seq data of human blood (B lymphocyte, T lymphocyte, and monocyte) from
the Cistrome[29] portal for the PBMC data (Fig. 4e, Supplementary Table 4), and ChIP-seq
data of Olig2[33], Neurog2[34], Eomes[34], and Tbr1[35] for the mouse embryonic brain data. The
Olig2 ChIP-seq data were generated in three types of rat cells: data from oligodendrocyte
precursor cells (OPC) and immature oligodendrocytes (iOL) were used for validation,
while data from mature oligodendrocytes (mOL) serve as a negative control[33]. Genomic
coordinates were converted from rn4 to mm10. The Neurog2 and Eomes ChIP-seq data
were generated in mouse embryonic cerebral cortices at day 14.5[34]; the Tbr1 ChIP-seq
data was generated in the whole cortex dissected from embryos at day 15.5[35]. In addition,
DNA footprinting signatures were corrected for Tn5 sequence insertion bias and stratified
by cell types using the Signac package[12] and can be used to validate the identified
TFs/motifs in a cell-type-specific manner (Fig. 6g, Supplementary Fig. 7e).
Hypergeometric tests for peak enrichment in TF binding sites by ChIP-seq were carried
out (Supplementary Fig. 17b-d, 18b-d). The results presented in Fig. 4e were obtained in
several steps: (i) we obtained sets of trios, for which B cells, T cells, and monocytes were
significantly influential; (ii) we applied TRIPOD and took the union set of the significant
trios; and (iii) we took the intersection between the trios obtained by the two types of
analyses, collapsed the trios to TF-peak relationships, and computed the fraction of peaks
overlapping ChIP-seq peaks.

**Data availability**
This study analyzed existing and publicly available single-cell RNA and ATAC multiomic
data. 10X Genomics single-cell multiomic datasets of PBMC (10k and 3k) and mouse
embryonic brain were downloaded https://support.10xgenomics.com/single-cell-
multiome-atac-gex/datasets. SNARE-seq data of adult mouse brain and SHARE-seq
data of mouse skin are available from the Gene Expression Omnibus (GEO) database
with accession numbers GSE126074 and GSE140203. A detailed data summary is
provided in Supplementary Table 1. Validation resources based on existing databases
and high-throughput sequencing data are summarized in Table 1 and Supplementary
Table 4.

**Code availability**

TRIPOD is compiled as an open-source R package available at https://github.com/yharigaya/TRIPOD. Scripts used for analyses carried out in this paper are deposited in the GitHub repository.

**Acknowledgments**

This work was supported by the National Institutes of Health (NIH) grant R35 GM133712 (to C.Z.), R01 HG006137 (to N.R.Z), R01 GM125301 (to N.R.Z.), and R35 GM138342 (to Y.J.). The authors thank Dr. Sai Ma for support and guidance on the SHARE-seq data, Manas Tiwari for help on accessing the hTFtarget database, and Drs. Yun Li, Michael Love, Li Qian, and Jason Stein for helpful discussions and comments.

**Author contributions**

Y.J. and N.R.Z. initiated and envisioned the study. Y.J., Y.H., and N.R.Z. formulated the model and developed the algorithm. Y.J. led the analyses for gene expression prediction, PBMC, and mouse skin. Y.H. led the analyses for mouse embryonic brain. Z.Z. processed reference datasets and performed cell-type label transfer. H.Z. and C.Z. provided support on validation, offered consultation, and contributed to result interpretation. Y.J., Y.H., and N.R.Z. wrote the manuscript, which was read and approved by all authors.

**Competing Interests**

The authors declare no competing interests.

**Figure Legends**

**Fig. 1 | Interaction between TF expression and peak accessibility improves RNA prediction accuracy. a**, UMAP embedding of 10x Genomics PBMC (left), 10x Genomics embryonic mouse brain (center), and SHARE-seq mouse skin (right) cells from single-cell RNA and ATAC multiomic sequencing. Cell-type labels were transferred from existing single-cell references or curated based on marker genes, motifs, and peaks; metacells were constructed to mitigate sparsity and stochasticity. **b**, Genome-wide distributions of

Pearson correlations between observed and leave-one-out predicted RNA expression levels, with varying window sizes. Predictions are from gene activity, regulatory potential, peak LASSO regression, and peak-TF LASSO regression. **c**, Predicted and observed RNA expression levels for highly variable genes, *CCR7*, *Adamts6*, and *Ano7*, from the three datasets, respectively.

**Fig. 2 | TRIPOD infers peak-TF-gene trio regulatory relationships using single-cell multiomic data. a**, Data input and schematic on a peak-TF-gene trio. **b**, Overview of TRIPOD for inferring regulatory relationships. TRIPOD complements existing methods based on marginal associations by identifying conditional associations through matching by TF expression or peak accessibility. **c**, An example trio identified by TRIPOD, but not by the marginal associations due to the heterogeneity of cell-type-specific regulations. **d**, An example trio identified by the marginal associations, but not by TRIPOD. The peak and TF are significantly linked to the gene, yet they act through other TF and peak, and thus the regulatory trio is insignificant. The points represent metacells (left two panels) and pairs of matched metacells (right two panels). Genomic coordinates for the peaks are from hg38.

**Fig. 3 | Examples of trio regulatory relationships in PBMC single-cell multiomic dataset. a-b**, Example trios identified by TRIPOD. Violin plots show cell-type-specific distributions of gene expression, peak accessibility, and TF expression. Scatterplots show TRIPOD's level 1 and level 2 testing, respectively. Inner and outer circles around the points are color-coded based on the cell types of the matched metacells. Hierarchical clustering is performed on RNA expression levels of highly variable genes. Red/gray circles indicate whether removal of the corresponding branches of metacells significantly changes the model fitting; crosses indicate that removal of the groups of metacells resulted in inestimable coefficients. Genomic coordinates for the peaks are from hg38.

**Fig. 4 | TRIPOD identified trio regulatory relationships in PBMC single-cell multiomic dataset supported by extensive validations. a,** A schematic of validation strategies. Shown are external datasets and databases used to validate the links between

peak accessibility and target gene expression (peak-gene validation), those between
peak accessibility and TF expression (peak-TF validation), and those between TF
expression and target gene expression (TF-gene validation). **b**, Peak-gene validation
based on enhancer databases (FANTOM5, 4DGenome, and EnhancerAtlas) and tissue-
specific *cis*-eQTL data from the GTEx Consortium. Box plots show distributions of $p$-
values from gene-specific hypergeometric tests. **c**, TF-gene validation based on lists of
TF-gene pairs from the knockTF database. **d**, Precision and recall rates for TF-gene pairs
using ground truths from the hTFtarget database. **e**, Peak-TF validation based on eight
cell-type-specific TF ChIP-seq datasets (B lymphocytes, monocytes, and T lymphocytes).
Fractions of significantly linked peaks and all peaks that overlap with the ChIP-seq peaks
are shown.

**Fig. 5 | TRIPOD identified known and novel regulatory relationships during mouse**
**embryonic brain development. a**, Venn diagram of the number of peak-gene pairs
captured by PLAC-seq, the marginal model, and the union set of TRIPOD's level 1 and
level 2 testing matching TF expression and peak accessibility. **b**, The same as **a** but for
Peak-TF validation by ChIP-seq data for Olig2, Neurog2, Eomes, and Tbr1. **c**, A
schematic of well-characterized TF regulatory cascades during neurogenesis and
gliogenesis. **d**, Trio examples from known regulatory relationships, as well as from
crosstalks supported by ChIP-seq data, captured by TRIPOD. **e**, GO analysis of putative
target genes of the neurogenesis and gliogenesis TFs. The number of TRIPOD-identified
target genes in the GO categories is shown. The background heatmap shows negative
log $p$-values (FDR < 0.05) from hypergeometric tests examining enrichment of GO terms.
**f**, Bar plots showing the number of putative cell-type-specific trios mediated by the
neurogenesis- and gliogenesis-specific TFs.

**Fig. 6 | TRIPOD identified regulatory relationships in mouse hair follicles with**
**transient cell states. a**, UMAP embedding of hair follicle cells from the mouse skin data.
Cells are colored by cell types (TAC, IRS, hair shaft, and medulla) and pseudotime. **b**,
H3K4me1 and H3K27ac ChIP-seq scores for linked peaks identified by TRIPOD, DORCs
(regulatory domains identified by gene-peak correlations), and randomly sampled peaks.

**c**, TRIPOD's linked peaks for four representative genes were significantly enriched in

previously annotated super-enhancers in the mouse TAC population. **d**, Trios detected

by TRIPOD that were active in IRS (top), medulla (middle), and hair shaft (bottom),

respectively. **e**, Dot plots of gene expressions, peak accessibilities, and TF expressions

across different cell types. **f**, Influence analyses identified segments along the

differentiation trajectory where the regulation took effect. The colors in the UMAP

embedding correspond to the smoothed $p$-values from a sampling-based approach. **g**,

DNA footprinting assays showed cell-type-specific enrichments of Tn5 integration events.

The findings were consistent with those from the influence analyses.

**Table 1 | Resources for validating peak-TF-gene regulatory relationship.** While there

is no existing experimental approach to validate all three factors in a trio at high resolution

with high throughput, we resort to existing databases and orthogonal sequencing data to

validate peak-gene, peak-TF, and TF-gene pairs, completing the loop.

**References**

1.　Gasperini, M., Tome, J.M. & Shendure, J. Towards a comprehensive catalogue of validated and
target-linked human enhancers. *Nat Rev Genet* **21**, 292-310 (2020).
2.　Hobert, O. Gene regulation by transcription factors and microRNAs. *Science* **319**, 1785-1786
(2008).
3.　Mortazavi, A., Williams, B.A., McCue, K., Schaeffer, L. & Wold, B. Mapping and quantifying
mammalian transcriptomes by RNA-Seq. *Nat Methods* **5**, 621-628 (2008).
4.　Johnson, D.S., Mortazavi, A., Myers, R.M. & Wold, B. Genome-wide mapping of in vivo protein-
DNA interactions. *Science* **316**, 1497-1502 (2007).
5.　Buenrostro, J.D., Giresi, P.G., Zaba, L.C., Chang, H.Y. & Greenleaf, W.J. Transposition of native
chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins
and nucleosome position. *Nat Methods* **10**, 1213-1218 (2013).
6.　Lieberman-Aiden, E. et al. Comprehensive mapping of long-range interactions reveals folding
principles of the human genome. *Science* **326**, 289-293 (2009).
7.　Feng, C. et al. KnockTF: a comprehensive human gene expression profile database with
knockdown/knockout of transcription factors. *Nucleic Acids Res* **48**, D93-D100 (2020).
8.　Tompa, M. et al. Assessing computational tools for the discovery of transcription factor binding
sites. *Nat Biotechnol* **23**, 137-144 (2005).
9.　Consortium, G. Genetic effects on gene expression across human tissues. *Nature* **550**, 204-213
(2017).
10.　Kim-Hellmuth, S. et al. Cell type-specific genetic regulation of gene expression across human
tissues. *Science* **369** (2020).
11.　Stuart, T. et al. Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902 e1821
(2019).

12.    Stuart, T., Srivastava, A., Lareau, C. & Satija, R. Multimodal single-cell chromatin analysis with
889            Signac. *bioRxiv*, 2020.2011.2009.373613 (2020).
890    13.    Ma, S. et al. Chromatin Potential Identified by Shared Single-Cell Profiling of RNA and Chromatin.
891            *Cell* **183**, 1103-1116 e1120 (2020).
892    14.    Hao, Y. et al. Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587 e3529
893            (2021).
894    15.    Duren, Z., Chen, X., Jiang, R., Wang, Y. & Wong, W.H. Modeling gene regulation from paired
895            expression and chromatin accessibility data. *Proc Natl Acad Sci U S A* **114**, E4914-E4923 (2017).
896    16.    Li, B. & Ritchie, M.D. From GWAS to Gene: Transcriptome-Wide Association Studies and Other
897            Methods to Functionally Understand GWAS Discoveries. *Front Genet* **12**, 713230 (2021).
898    17.    Pliner, H.A. et al. Cicero Predicts cis-Regulatory DNA Interactions from Single-Cell Chromatin
899            Accessibility Data. *Mol Cell* **71**, 858-871 e858 (2018).
900    18.    Wang, C. et al. Integrative analyses of single-cell transcriptome and regulome using MAESTRO.
901            *Genome Biol* **21**, 198 (2020).
902    19.    Cao, J. et al. Joint profiling of chromatin accessibility and gene expression in thousands of single
903            cells. *Science* **361**, 1380-1385 (2018).
904    20.    Chen, S., Lake, B.B. & Zhang, K. High-throughput sequencing of the transcriptome and chromatin
905            accessibility in the same cell. *Nat Biotechnol* **37**, 1452-1457 (2019).
906    21.    Becht, E. et al. Dimensionality reduction for visualizing single-cell data using UMAP. *Nat
907            Biotechnol* (2018).
908    22.    Taylor, J. & Tibshirani, R.J. Statistical learning and selective inference. *Proc Natl Acad Sci U S A*
909            **112**, 7629-7634 (2015).
910    23.    Schep, A.N., Wu, B., Buenrostro, J.D. & Greenleaf, W.J. chromVAR: inferring transcription-factor-
911            associated accessibility from single-cell epigenomic data. *Nat Methods* **14**, 975-978 (2017).
912    24.    Andersson, R. et al. An atlas of active enhancers across human cell types and tissues. *Nature*
913            **507**, 455-461 (2014).
914    25.    Wang, J. et al. HACER: an atlas of human active enhancers to interpret regulatory variants.
915            *Nucleic Acids Res* **47**, D106-D112 (2019).
916    26.    Teng, L., He, B., Wang, J. & Tan, K. 4DGenome: a comprehensive database of chromatin
917            interactions. *Bioinformatics* **32**, 2727 (2016).
918    27.    Gao, T. & Qian, J. EnhancerAtlas 2.0: an updated resource with enhancer annotation in 586
919            tissue/cell types across nine species. *Nucleic Acids Res* **48**, D58-D64 (2020).
920    28.    Zhang, Q. et al. hTFtarget: A Comprehensive Database for Regulations of Human Transcription
921            Factors and Their Targets. *Genomics Proteomics Bioinformatics* **18**, 120-128 (2020).
922    29.    Mei, S. et al. Cistrome Data Browser: a data portal for ChIP-Seq and chromatin accessibility data
923            in human and mouse. *Nucleic Acids Res* **45**, D658-D662 (2017).
924    30.    La Manno, G. et al. Molecular architecture of the developing mouse brain. *Nature* **596**, 92-96
925            (2021).
926    31.    Huang, M., Zhang, Z. & Zhang, N.R. Dimension reduction and denoising of single-cell RNA
927            sequencing data in the presence of observed confounding variables. *bioRxiv*,
928            2020.2008.2003.234765 (2020).
929    32.    Zhu, C. et al. An ultra high-throughput method for single-cell joint analysis of open chromatin
930            and transcriptome. *Nat Struct Mol Biol* **26**, 1063-1070 (2019).
931    33.    Yu, Y. et al. Olig2 targets chromatin remodelers to enhancers to initiate oligodendrocyte
932            differentiation. *Cell* **152**, 248-261 (2013).
933    34.    Sessa, A. et al. The Tbr2 Molecular Network Controls Cortical Neuronal Differentiation Through
934            Complementary Genetic and Epigenetic Pathways. *Cereb Cortex* **27**, 5715 (2017).

35.    Notwell, J.H. et al. TBR1 regulates autism risk genes in the developing neocortex. *Genome Res*
**26**, 1013-1022 (2016).

36.    Mira, H. & Morante, J. Neurogenesis From Embryo to Adult - Lessons From Flies and Mice. *Front*
*Cell Dev Biol* **8**, 533 (2020).

37.    Emery, B. & Lu, Q.R. Transcriptional and Epigenetic Regulation of Oligodendrocyte Development
and Myelination in the Central Nervous System. *Cold Spring Harb Perspect Biol* **7**, a020461
(2015).

38.    Huang da, W., Sherman, B.T. & Lempicki, R.A. Systematic and integrative analysis of large gene
lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57 (2009).

39.    Achim, K., Salminen, M. & Partanen, J. Mechanisms regulating GABAergic neuron development.
*Cell Mol Life Sci* **71**, 1395-1415 (2014).

40.    Setty, M. et al. Characterization of cell fate probabilities in single-cell data with Palantir. *Nat*
*Biotechnol* **37**, 451-460 (2019).

41.    Bravo Gonzalez-Blas, C. et al. cisTopic: cis-regulatory topic modeling on single-cell ATAC-seq
data. *Nat Methods* **16**, 397-400 (2019).

42.    Adam, R.C. et al. Pioneer factors govern super-enhancer dynamics in stem cell plasticity and
lineage choice. *Nature* **521**, 366-370 (2015).

43.    Natarajan, A., Yardimci, G.G., Sheffield, N.C., Crawford, G.E. & Ohler, U. Predicting cell-type-
specific gene expression from regions of open chromatin. *Genome Res* **22**, 1711-1722 (2012).

44.    Zhou, W. et al. Genome-wide prediction of DNase I hypersensitivity using gene expression. *Nat*
*Commun* **8**, 1038 (2017).

45.    Blatti, C., Kazemian, M., Wolfe, S., Brodsky, M. & Sinha, S. Integrating motif, DNA accessibility
and gene expression data to build regulatory maps in an organism. *Nucleic Acids Res* **43**, 3998-
4012 (2015).

46.    Kazemian, M., Pham, H., Wolfe, S.A., Brodsky, M.H. & Sinha, S. Widespread evidence of
cooperative DNA binding by transcription factors in Drosophila development. *Nucleic Acids Res*
**41**, 8237-8252 (2013).

47.    Panigrahi, A. & O'Malley, B.W. Mechanisms of enhancer action: the known and the unknown.
*Genome Biol* **22**, 108 (2021).

48.    Urrutia, E., Chen, L., Zhou, H. & Jiang, Y. Destin: toolkit for single-cell analysis of chromatin
accessibility. *Bioinformatics* **35**, 3818-3820 (2019).

49.    Jiang, Y., Zhang, N.R. & Li, M. SCALE: modeling allele-specific gene expression by single-cell RNA
sequencing. *Genome Biol* **18**, 74 (2017).

50.    Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-seq data
using regularized negative binomial regression. *Genome Biol* **20**, 296 (2019).

51.    Fornes, O. et al. JASPAR 2020: update of the open-access database of transcription factor
binding profiles. *Nucleic Acids Res* **48**, D87-D92 (2020).

52.    Kulakovskiy, I.V. et al. HOCOMOCO: towards a complete collection of transcription factor
binding models for human and mouse via large-scale ChIP-Seq analysis. *Nucleic Acids Res* **46**,
D252-D259 (2018).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

| Validation | Database/Resource | Description | Organism | Tissue | PMID | GSE | GSM | URL |
|---|---|---|---|---|---|---|---|---|
| Peak-gene | FANTOM5 (HACER) | Tissue-specific enhancer database | Human | Blood cell lines | 24670763 | NA | NA | http://bioinfo.vanderbilt.edu/AE/HACER |
| | 4DGenome (HACER) | Tissue-specific enhancer database | Human | Blood cell lines | 25788621 | NA | NA | http://bioinfo.vanderbilt.edu/AE/HACER |
| | EnhancerAtlas2.0 | Tissue-specific enhancer database | Human | Blood cell lines | 31740966 | NA | NA | http://www.enhanceratlas.org |
| | GTEx | Tissue-specific cis-eQTLs | Human | Whole blood | 29022597 | NA | NA | https://gtexportal.org |
| | PLAC-seq | Enhancer-promoter interactions by PLAC-seq (rep 1) / Enhancer-promoter interactions by PLAC-seq (rep 2) | Mouse | Embryonic forebrain at day 16.5 | 31695190 | GSE130399 | GSM3819641 / GSM3819642 | https://www.ncbi.nlm.nih.gov/geo |
| | Super-enhancer | H3K4me1 and H3K27ac ChIP-seq | Mouse | Transit-amplifying cell (TAC) from mouse hair follicle | 25799994 | GSE61316 | GSM1502001 / GSM1502003 | https://www.ncbi.nlm.nih.gov/geo |
| Peak-TF | Cistrome | A data portal for ChIP-Seq and chromatin accessibility data in human and mouse | Human | B lymphocyte, T lymphocyte, monocyte | 27789702 | Supplementary Table S2 | | http://cistrome.org |
| | TF ChIP-seq | Olig2 ChIP-seq | Rat | Oligodendrocyte precursor cell (OPC) / Immature oligodendrocyte (iOL) / Mature oligodendrocyte (mOL) | 23332759 | GSE42454 | GSM1040156 / GSM1040157 / GSM1040158 | https://www.ncbi.nlm.nih.gov/geo |
| | | Neurog2 ChIP-seq / Eomes ChIP-seq | Mouse | Embryonic cerebral cortex at day 14.5 | 27600842 | GSE63620 | GSM1553880 / GSM1553879 | |
| | | Tbr1 ChIP-seq | | Embryonic whole cortex at day 15.5 | 27325115 | GSE71384 | GSM1833461 | |
| TF-gene | knockTF | A database of human gene expression profiles with knockdown/knockout of transcription factors | Human | Blood cells | 31598675 | NA | NA | http://www.licpathway.net/KnockTF |
| | hTFtarget | A database of human transcription factors and their targets | Human | Blood cells | 32858223 | NA | NA | https://bio.tools/hTFtarget |

Table 1