1

Deriving time-concordant event cascades from gene expression data: A case study for Drug-Induced Liver Injury (DILI)

Anika Liu[1,2,3*], Namshik Han[1,4], Jordi Munoz-Muriedas[2], Andreas Bender[3*]

[1] Milner Therapeutics Institute, University of Cambridge, Cambridge, United Kingdom

[2] Systems Modelling and Translational Biology, Data and Computational Sciences, GSK, London,

United Kingdom

[3] Centre for Molecular Informatics, Department of Chemistry, University of Cambridge, Cambridge,

United Kingdom

[4] Cambridge Centre for AI in Medicine, Department of Applied Mathematics and Theoretical

Physics, University of Cambridge, Cambridge, United Kingdom

16
17 *Corresponding author
18 E-mail: ab454@cam.ac.uk (AB), al862@cam.ac.uk (AL)
19

Abstract

Adverse event pathogenesis is often a complex process which compromises multiple events ranging from the molecular to the phenotypic level. In toxicology, Adverse Outcome Pathways (AOPs) aim to formalize this as temporal sequences of events, in which event relationships should be supported by causal evidence according to the tailored Bradford-Hill criteria. One of the criteria is whether events are consistently observed in a certain temporal order and, in this work, we study this time concordance using the concept of "first activation" as data-driven means to generate hypotheses on potentially causal mechanisms. As a case study, we analysed liver data from repeat-dose studies in rats from the TG-GATEs database which comprises measurements across eight timepoints, ranging from 3 hours to 4 weeks post-treatment. We identified time concordant gene expression-derived events preceding adverse histopathology, which serves as surrogate readout for Drug-Induced Liver Injury (DILI). We find known mechanisms in DILI to be time-concordant, and show further that significance, frequency and log fold change (logFC) of differential expression are metrics which can additionally prioritize events although not necessary to be mechanistically relevant. Moreover, we used the temporal order of transcription factor (TF) expression and regulon activity to identify transcriptionally regulated TFs and subsequently combined this with prior knowledge on functional interactions to derive detailed gene-regulatory mechanisms, such as reduced Hnf4a activity leading to decreased expression and activity of Cebpa. At the same time, also potentially novel events are identified such as Sox13 which is highly significantly time-concordant and shows sustained activation over time. Overall, we demonstrate how time-resolved transcriptomics can derive and support mechanistic hypotheses by quantifying time concordance and how this can be combined with prior causal knowledge, with the aim of both understanding mechanisms of toxicity, as well as potential applications to the AOP framework. We make our results available in the form of a Shiny app (https://github.com/anikaliu/DILICascades_App), which allows users to query events of interest in more detail.

Author Summary

Understanding mechanisms from systems-scale biological data is of great relevance in toxicology as well as drug discovery; however how to generate causal hypotheses instead of correlations is by no means clear. In this work, we study the conserved temporal order of events and present an automatable framework to quantify and characterize time concordance across a large set of time-series. We apply this concept to events derived from time-resolved gene expression and histopathology from the TG-GATEs *in vivo* liver data as a case study. We were able to recover known events involved in the pathogenesis of Drug-Induced Liver Injury (DILI), and identify potentially novel pathway and transcription factors (TFs) which precede adverse histopathology. As complementary sources of evidence for causality, we additionally show how time concordance and prior knowledge on plausible interactions between TFs can be combined to derive causal hypotheses on the TFs' mode of regulation and interaction partners. Overall, the results derived in our case study can serve as valuable hypothesis-free starting points for the development of Adverse Outcome Pathways for DILI, and demonstrate that our approach provides a novel angle to prioritize mechanistically relevant events.

Introduction

Adverse drug reactions are a major reason for compound failure in the clinical trials [1,2] and the significant cause for post-marketing withdrawals. To counter exposing patients to these risks, it is desired to identify adverse events earlier in the individual patient but also in the drug development process. Mechanistic understanding of how adverse event pathogenesis is crucial in this regard, i.e. to derive early safety biomarkers or *in vitro* assays. However, current understanding of toxicity is largely incomplete, in particular for complex phenotypes such as organ injury which can usually be

caused by a wide range of compounds perturbing the biological system at different points mediated

through multiple biological scales and entities [3,4].

Multiple interrelated concepts have been introduced to formalize mechanistic knowledge in the

context of toxicity including Adverse Outcome Pathways, AOPs [4,5]. These begin with a molecular

initiating event (MIE) which describes the first interaction of the compound with the system, e.g. a

target protein, which is then linked to the adverse event (AE) through a causal cascade of key

events (KEs) on different biological levels, like activation of cellular pathways or changes in the

tissue or organ. Thereby, the Organization for Economic Co-operation and Development [6]

published three criteria to evaluate causality between events within AOPs based on the original

Bradford Hill considerations in the context of epidemiological studies [7] and previous work on the

related Mode Of Action (MOA) concept [8]: i) Biological plausibility, ii) essentiality of key events and

iii) empirical support for key event relationships. Empirical support for a causal relation between

events E_{cause}, which could be a MIE or KE, and $E_{consequence}$, which could be a AE, is further separated

into time concordance (E_{cause} happens before $E_{consequence}$), dose concordance (E_{cause} happens at

lower dose than $E_{consequence}$) and incidence concordance (E_{cause} affects a larger population and is

hence more frequent than $E_{consequence}$).

Computational approaches can thereby support these predominantly expert- and knowledge-driven

mechanistic efforts by prioritizing mechanistically relevant events or by providing additional insight

on the relation between an event and a given phenotype. For instance, computationally predicted

AOPs (cpAOPs) prioritize plausible events and event relationships as starting points for expert-

driven AOP development by integrating functional and statistical associations between biological

entities on different levels [9–11]. In contrast, probabilistic quantitative AOPs (qAOPs) provide

additional insight on the predictivity of key event relationships (KERs) by aiming to predict the

adverse event from *in vitro* assays implementing the expert-curated AOP as scaffold [12,13]. To

support increasing implementation of *in vitro* methods, however, it needs to be better understood

which readouts describing molecular or cellular effects are indicative for systems-level adverse

effects in the first place.

Biological readouts such as transcriptomics are particularly suited to study such intermediate key

events as they provide broad insights into cellular changes, e.g. in contrast to target profiling, which

can then lead to the identification of predictive signatures and mechanistically relevant insights. This

is for example true in the context of DILI [14–18], which is a major cause for attrition in drug

development and accounts for around half of the cases of acute liver failure in the US and European

countries [19,20]. In this regard, in particular time-series data is interesting as it is able to trace the

dynamic effects throughout pathogenesis. Previous studies focussed on the time (and dose)

dependence of gene expression-derived events in the context of adverse findings [21,22], so the

changes of individual events across changes in time (and dose), and also aimed to predict later

adverse findings from fixed early timepoints [14,15]. From a mechanistic perspective, however,

neither activation at a certain timepoint nor a certain progression over time is mandatory, but only

time concordance, so activation of the key event before the downstream adverse effects.

In this study, we hence quantify time concordance across gene expression- derived cellular events

and adverse events based on histopathology across wide range of compounds. To do so, we

introduce the concept of "first activation" for mechanistic analysis, which focusses only on the

earliest timepoint an event can be reliably detected and then orders events within a time-series by

their timepoint of first activation (Fig 1A). In contrast to previous time concordance analyses in

AOPs which addressed a defined set of KER and known KE [23–25], this analysis derives statistical

evidence for temporal concordance across time-series and can do so for any combination of events

based on gene expression or histopathology. Although the confidence of these temporal orders per

time series is limited by the noisiness of gene expression data and the low time resolution, statistical

significance can be evaluated across time series and we only consider relations time-concordant if

the preceding event is significantly more frequently found before the later event than in time series

without (Fig 1B). Furthermore, this also allows us to separate out events which depict general

perturbation response but are unspecific, as well as rare events, which are predictive but only

observed for a small subset of compounds.

Fig 1: Quantifying time concordance based on first activation. (A) The event activation of the
events A-D and the later event is shown over time, as well as their timepoint of first
activation, at which the event first passes the defined activation criteria. If an event takes
place before a defined later event, which in our study is adverse histopathology, it is time-
concordant. Time concordance indicates that there is potentially a causal relation between
both events, and is distinct from time-dependence which is defined based on the correlation
to the later event or time. (B) Based on the frequency of an event before or at the same time
as the later event and its frequency in background time-series without the later event, a
confusion matrix and different time concordance metrics can be derived. (C) Time
concordance can both prioritize novel links and provide further evidence on potential
mechanistic links between events. Events are indicated as nodes and mechanistic links
between them as edges.

We demonstrate the utility of this concept in this work using liver gene expression and

histopathology data from repeat-dose studies in rats provided by the TG-GATEs database (S1 Fig).

This allows us to take advantage of previous data curation and work on the dataset itself, in

particular by Sutherland et al. [15] who provide an adverse classification of each compound-dose

combination and toxscores summarising histopathological findings in each condition. Furthermore,

Drug-Induced Liver Injury (DILI) is well understood in comparison to other organ-level toxicities and

we hence know which processes are expected to precede injury, including cell death, inflammation

and other adaptive stress responses [19]. The concept, however, is generally applicable beyond the

toxicity area and transcriptomics data and can be used to derive mechanistic event cascades from

time-series data of any kind as long as the first activation of events within the time-series can be

defined.

We first describe the time concordance for known processes, similar to mechanistic qAOPs, and

then prioritize predictive, time-concordant KE providing a strong data-driven, automatable starting

point for AOP development, aligning with the objective of cpAOPs (more detailed comparison in S1

Table). We then combine data-driven time concordance and prior knowledge on event relations

between transcription factors (TFs) and gene expression to generate hypotheses for causal gene-

regulatory mechanisms in DILI pathogenesis and to generally show how time concordance can

stratify and support other streams of causal evidence. Overall, we show that time-resolved gene

expression and histopathology data can be used to quantify time concordance across a large set of

compounds and events, which allows us to characterize known mechanistic links and to prioritize

new ones (Fig 1C).

Results and Discussion

In order to derive the time concordance between cellular events and later adverse histopathology,

we use the workflow outlined in Fig 2 with each step being also introduced in the subsequent

sections and details on their respective implementation in Methods. We first derived TF and

pathway activity across expression profiles from the same experiment and subsequently defined the

first up- or downregulation TFs or pathways as events. Furthermore, we obtained binary

histopathology labels describing the occurrence of each histopathological finding at different levels

of severity and frequency from the Toxscores provided by Sutherland et al. [15]. Subsequently, we

derive the earliest timepoint of each event, e.g. pathways or adverse histopathology, within each

time-series. As last step of the time concordance analysis, we then evaluate which gene

expression-derived events are significantly enriched before or at the time where adverse

histopathology is found, as well as additional time concordance metrics outlined in Table 1.

Fig 2: Workflow to quantify time concordance between preceding gene expression-derived
events and later adverse histopathology. First, events are derived from the gene expression
and histopathology data. Pathway and TF activity is inferred based on the expression of the
respective gene sets using GSVA[26] and binary histopathology labels are derived from the
continuous toxscores. Secondly, we derive the first activation of expression-based events as
well as of adverse histopathology. Lastly, we quantify the time concordance between
potential preceding events (PE) which are derived from gene expression and adverse
histopathology as potential later event (LE).

176   Adverse histopathological findings and their temporal relation

To define the earliest timepoint of adverse histopathology within each time-series, we used the

annotations of time series as adverse or non-adverse by Sutherland et al. [15], as well as the

toxscores, which summarise the severity and frequency for each histological finding and each

compound-dose-time combination as mean severity score and range from 0 (normal) to 4 (severe).

These toxscores were used to define three levels for each histological finding: "null" (toxscore > 0),

"low" (toxscore > 0.67) and "high" (toxscore >1.34). For example in case of a toxscore of 1, both

"null" and "low" are considered to be present. We then evaluated which histology groups were

frequently found in the adverse compound-dose combinations (observed in >10% of adverse time

series corresponding to at least 5 out of 40 cases) with at least 50% of findings being in adverse

time series (Fig 3A). All of the included histology groups are significantly enriched in adverse

conditions, however, these criteria were implemented to identify findings with a certain specificity

and frequency instead of allowing a trade-off between both. The histology groups which passed the

filtering are regarded as adverse histopathological findings and include hepatocellular single cell

necrosis and biliary hyperplasia at all toxscore thresholds. In contrast, only some of the three

toxscore thresholds were selected with the above criteria for all other findings, e.g. the two higher

toxscore cut-offs for hepatocellular necrosis and inflammation and only the "high" cut-off for

increased hepatocellular mitosis. In all cases, the lower toxscore level was also frequently observed

in non-adverse conditions and hence considered too unspecific. In contrast, only the two milder

levels of fibrosis were included in the selection, as severe fibrosis was observed rarely. While we

focus on the described definition of adverse histopathological findings in this study, the difficulty in

summarising a complex phenotype such as DILI into a binary classification, adverse or not adverse,

is well established [27,28] and is also demonstrated by the discrepancies between DILI

classifications from DILIst, DILIrank and those derived by Sutherland et al based on the TG-GATEs

data (S1). We are aware that also broader or more targeted phenotypes might be of interest, and

we hence provide a Shiny app where results for alternative definitions of adverse and non-adverse

histopathology groups can be explored.

For the adverse histopathology labels, the distribution of toxscores and first activation over time (Fig

3B) shows that some findings are predominantly found late, like fibrosis, while others are

predominantly found early, e.g. hepatocellular single cell necrosis. Next, out of all 360 time-series

with at least 6 measured timepoints, the 61 time-series in which any of the adverse histopathology

labels is found were identified, which covered 38 compounds (S2 Table). In those, the earliest

evidence of an adverse phenotype is used to approximate the timepoint of the primary adverse

phenotype. Across all timeseries with adverse histopathology, we find that hepatocellular single cell

necrosis is most frequently the primary adverse phenotype, while biliary hyperplasia at any severity

is in most cases a secondary effect (Fig 3C, S3 Fig).

Fig 3: Distribution and relation of histopathological findings across time series. A)
Histopathology labels are defined for each histopathological findings at 3 different toxscore
cut-offs, namely "null" (toxscore>0), "low" (toxscore>0.67) and "high" (toxscore>1.34. For
each label, the number of occurrences in the 40 adverse time series and the fraction of
adverse time series among all occurrences of the given histopathology label are shown.
Histopathological findings, out of which at least 50% and at least 5 of the occurrences were
found in adverse conditions timeseries were considered adverse B) Number of conditions
with histopathological findings at different timepoints, as well as the frequency of the
respective first activations C) Time of first activation across timeseries labelled as adverse

223 **or non-adverse. Each time series is annotated with the dose level in repeat-dose studies, as**
224 **well as whether or not the time series was considered adverse by Sutherland et al. [15].**

Known pathways in DILI preceding adverse histopathology

To identify cellular mechanisms in the early pathogenesis of DILI, we next studied time-concordant

cellular changes preceding later adverse histopathology (see Methods). This identified 911

pathway-level events (37.3%), and 108 TF-level events (33.6%) with significant enrichment (p-

value<0.05) before or at adverse histopathology. We next evaluated time concordance for a set of

ten known events in DILI (Fig 4 and S3 Table). Recycling of bile acids and salts was the most

significantly enriched geneset overall and hence also among the ones linked to known events. Also

down-regulation of the other bile acid gene sets was significantly enriched (p-value < 0.05) pointing

to an overall down-regulation of bile acid metabolism. While cell death was also only found to be up-

regulated, dysregulation in both directions was found to precede injury for all other key events (Fig

4). However, only for peroxisomal processes, namely peroxisomal protein import and beta-oxidation

of very long fatty acids, both directions were significantly enriched indicating that dysregulation in

either direction might be linked to injury. Overall, significantly enriched gene sets are found for all

ten represented known events in DILI (p-value < 0.05) indicating that our analysis is able to recover

known cellular events.

Fig 4: Enrichment of known events in DILI before adverse histopathology based on gene sets as well as individual gene members. The enrichment of first activation before or at adverse histopathology is shown for gene sets mapping to known key events in DILI, for which first activation was defined as first timepoint of differential GSVA-derived gene set activity. Furthermore, also enrichment of individual genes within these genesets is shown and was derived based on the first timepoints of differential expression. Aligning with the expected direction, a significant down-regulation of Liver X Receptor (LXR) signalling and bile acid-related pathways is observed, while all other gene sets were found to be more significantly up-regulated. Only for peroxisomal pathways, both directions were significantly enriched indicating that dysregulation in direction might be linked to adverse histopathology.

To gain insights on a more fine-grained level, we next analysed the enrichment of significantly and strongly (absolute log fold change > 1) dysregulated individual genes from the above gene set (S2 File). Among the ten most significantly enriched gene-level events, three are involved in known processes, namely the up-regulation of Acyl-CoA thioesterase 3 (*Acot3*) and Carnitine O-Acetyltransferase (*Crat*) which are involved in fatty acid beta oxidation [29,30]. Multiple genes among the ten most significantly enriched gene-level events are also involved in mitochondrial and peroxisomal processes except for *Gadd45a,* Growth Arrest And DNA Damage-Inducible Protein which has a known role in hepatic fibrosis [32], Neutral Cholesterol Ester Hydrolase 1 (*Nceh1*) which is involved in cholesterol metabolism in macrophages [33], Ras-Related Protein Rab-30

(*Rab30*) elevated in early liver regeneration [34], as well as the Serine/Threonine Protein Kinase

NIM1 (*Nim1k*).

For JNK signalling, we did not find any significantly enriched genes indicating that while the overall

process is changing none of the individual genes shows strong and frequent expression changes. In

contrast, the opposite was found for oxidative stress with the Jun Proto-Oncogene (*Jun*) being one

of the most significantly enriched gene-level events but lacking significant changes on the gene-set

level. This shows that both gene- and gene-set level analysis can provide complementary insights

into cellular changes preceding DILI, and that in some cases effects can be attributed in individual

genes which might give more detailed information about the cellular changes.

While significant enrichment before or at adverse histopathology can be regarded as a necessary

criterion for time concordance, the temporal event relationship can be further characterised based

on the observed behaviour across experimental conditions which may be useful to further prioritize

mechanistically relevant pathways in a hypothesis-free manner. Following the Bradford-Hill

considerations, we hypothesize that this might be the case for observed effect size, frequency and

specificity of event occurrence before adverse histopathology. Firstly, we investigated how strongly

pathways were dysregulated comparing the maximal absolute log fold changes (|logFCs|) before or

at adverse histopathology in each adverse time-series for significantly time-concordant events (Fig

5). High median maximal |logFCs| were overall found for mitochondrial and peroxisomal pathways

and the highest median maximal |logFC| among all significant events was found for mitochondrial

fatty acid oxidation of unsaturated fatty acids. At the same time, however, the high variance for

pathways with high median maximal |logFC| as well as the only moderately high |logFC|s observed

for other known pathways in DILI, such as programmed cell death. This indicates that a high

magnitude of |logFC| is not necessary to contribute to an adverse event, but at the same time can

be a useful property to further prioritize important pathways.

Fig 5: Observed max. |logFC| before adverse histopathology. For known processes in DILI which correspond to significantly enriched events before adverse histopathology, the max |logFC| before adverse histopathology is shown. In comparison to other known pathways and the overall background distribution, a high logFC is found for mitochondrial beta oxidation followed by peroxisomal beta oxidation and mitophagy.

We next analysed to what extent dysregulation in a pathway is predictive for a particular type of histopathology. To this end, we calculated across how many adverse time-series each pathway is observed, summarised by the true positive rate (TPR), and the positive predictive value (PPV) indicating whether presence of the key event is a confident indicator for the later adverse event (Fig 6). We focus on significantly enriched events only (p-value < 0.05) and find a trade-off with respect to the highest TPR and PPV (Fig 6; for distribution of all events see S4 Fig). This generally shows that either highly frequent events with lower specificity can be identified, e.g. increased mitophagy (TPR: 0.41, PPV: 0.72), or more specific events at the expense of lower relative frequency, e.g. bile acid recycling (TPR= 0.30, PPV:1).

Surprisingly, lower relative frequencies are particularly observed for stress response and signaling

pathways with only Liver X Receptor (LXR)-dependent gene expression linked to lipogenesis

reaching a TPR over 20%. One explanation for the lower observed frequencies is that these

pathways are predominantly and initially mediated through post-transcriptional alterations instead of

gene expression changes [35,36], making the expression of pathway members a weak proxy for

pathway activation in early pathogenesis and explaining the overall low frequencies. In fact, one

reason LXR-dependent changes might have achieved higher frequencies as they explicitly include

the downstream regulated genes unlike the other signalling and stress response pathways [37].

Due to the previously discussed complementarity of gene- and gene set-level analysis, we also

show TPR and PPV for individual genes with a focus on those which are involved in gene sets

mapping to known key events. The most significant genes, already highlighted in Fig 4, reveal a

high frequency for the up-regulation of the Acyl-CoA thioesterases *Acot2*, *Acot3* and *Acot4*, as well

as the for the Enoyl-CoA Hydratase 1 *Ech1* which aligns with the relatively high frequency of

pathway-level events linked to mitochondrial and peroxosomal processes. Furthermore, the most

frequent gene-level events with a PPV=1 are the up-regulation of Activating Transcription Factor 3

(*Atf3*) which was found to promotes hepatic fibrosis [38] and Enoyl-CoA Delta Isomerase 1 (*Eci1*).

Fig 6: True positive rate (TPR) and positive predictive value (PPV) before or at histopathology of genes and gene sets in known key events in DILI. Events related to the given known key event are shown in red or blue indicating an up- or downregulation, respectively. Genes with a p-value < 0.0001 involved in known key events in DILI are

additionally labelled. The background distribution of all significantly enriched genes or gene
sets is shown in grey (p-value < 0.05).

Known TFs in DILI preceding adverse histopathology

To gain insight into signalling and expression regulation preceding adverse histopathology, we next analysed transcription factors (TFs), which are involved in early perturbation response preceding downstream gene expression changes and also are likely to show strong signal in transcriptomics data given their direct link to gene expression. As known TFs in DILI, we thereby included TFs mediating the stress response and signalling pathways already introduced above, as well as nuclear receptors which take in important roles in liver physiology and malfunctions and can be, both, MIEs or KEs (mapping shown in Fig 7A). Consistent with the pathway-level results, an enriched up-regulation was found for Nuclear factor erythroid 2-related factor 2 (Nfe2l2) which is a key mediator of oxidative stress [18,39] as well as the Nf-κB subunits Rela and Nfkb1 indicating inflammation [40], while the Oxysterols Receptors LXRα (Nr1h3) and LXRβ (Nr1h2) which control lipid metabolism showed enriched down-regulation [41].

For ER stress, we included three TFs mapping to the three branches of unfolded protein response [42]: Activating transcription factor 4 (Atf4), Activating transcription factor 6 (Atf6) and X-box binding protein 1 (Xbp1). Atf4 up-regulation was found to be most significantly enriched, most frequent and also showing the largest logFC (Fig 7). This highlights its overall importance in mediating ER stress and is consistent with the known role for ATF4 in DILI [43]. While Atf4 is a member of the pro-apoptotic unfolded protein response branch, the ATF6 and XPB1-mediated branches tend to be cytoprotective [44]. In agreement with this, Atf6 was not significantly enriched, however, Xbp1 showed rare but significantly enriched down-regulation.

Transcription Factor AP-1 (Jun) which is one of downstream target TFs of c-Jun N-terminal kinase (JNK) signaling was not significantly enriched in either direction due its rare activation among adverse time series although JNK signaling up-regulation itself was significantly enriched with *Jun*

up-regulation being one of the most significantly enriched gene-level events. However, JNK

signaling is particularly known in acetaminophen-induced liver injury and in this context leads to

hepatocyte death through interactions with Sab on the mitochondrial outer membrane and not

through transcriptional regulation mediated by AP-1 [45,46]. As increased Jun activity is hence

known to be a consequence of JNK signaling but not a cause of injury, it would be plausible to see

enriched pathway activity but not in TF activity before adverse histopathology. Overall, we were

hence able to show significant enrichment of some of the known TFs in DILI before adverse

histopathology and can also biologically reason the absence of significance for others.

While none of the included TF-level events ranked as most significant or most strongly changing

before adverse histopathology as in the analysis of pathway-level events, the down-regulation of

Nr1h3, which is involved in lipid metabolism, was identified as most frequent event (Fig 7B)

indicating that the linked physiological changes are commonly but not specifically found before

injury. Similarly, the up-regulation of stress response, indicated by Nfe2l2 and Atf4, was found to be

frequent aligning with their role in adaptive stress response [47]. Overall, frequency might hence be

a useful metric to identify pre-adverse cellular events which precede injury but are not highly

specific.

Data-driven prioritization of cellular events taking place before adverse

368

histopathology

369

370 As many events were found to be significantly enriched before adverse histopathology, we next

371 aimed at identifying and characterizing events most supported by time concordance, and hence to

372 move closer to the eventual aim of constructing AOPs from data. In our analysis, some known

373 events in DILI ranked highest by enrichment p-value while others rank highest by max. |logFC|

374 before adverse histopathology. In contrast, known TFs in DILI were found as most frequent ones in

375 the dataset. We hence next looked into the top 10 TF- and pathway-level events identified using

376 max. |logFC|, the enrichment p-value, and the TPR before or at adverse histopathology. These are

377 shown in Fig 8 and S4 Table, while all time concordance metrics are summarised in S2 File. The

378 most significantly enriched pathway-level event is decreased bile acid and salt recycling and also

379 the down-regulation of multiple metabolic pathways, in particular targeting glycosaminoglycans, is

380 found among the most significant pathway-level events pointing towards reduced liver function.

381 Moreover, the most significantly enriched TF-level event was the down-regulation of Transcription

382 factor activating enhancer binding protein 4 (Tfap4) which shows emerging roles in cell fate

383 decisions [48], and is followed by Homeobox B13 (Hoxb13) for which expression has previously

384 been found to correlate with hepatic inflammation in hepatic fibrosis [49].

385 Among the up-regulated events, the most significant enrichment is found for cell cycle checkpoints

386 and DNA repair among the pathway-level events as well as E2F Transcription Factor 2 (E2f2),

387 which controls cellular proliferation and liver regeneration [50], and was found among the most

388 significant TF-level events. E2f2 up-regulation was also identified as 2nd most frequent TF event

after the down-regulation of Nr1h3 and among the top 10 most strongly changing TF events further

highlighting its strong time concordance. The most frequent genesets point to translation regulation

via Eukaryotic translation initiation factor 2A (EIF2a) including the upstream response mediated by

eIF-2-alpha kinase GCN2 and the downstream role in protein translation mediated through

interactions with tRNA. EIF2a is part of the same branch of Unfolded Protein Response (UPR) as

Atf4 and causes its preferential translation which, among others, mediates autophagy and

proapoptotic response [42,51] and is a known predictor of DILI [52]. Furthermore, increased folding

of actin by Chaperonin containing tailless complex polypeptide 1 (CCT) or tailless complex

polypeptide 1 ring complex (TRiC) is found frequently and with large effect size. It has been

previously linked to proteostasis and autophagy [53,54], but a role in DILI specifically is not yet

known. As most strongly dysregulated events, metabolic pathways are found pointing to increased

beta oxidation of fatty acids, as well as decreased cholesterol biosynthesis and tyrosine catabolism.

Also the most strongly down-regulated TFs point towards lipid metabolism, i.e. the Sterol Regulatory

Element Binding Transcription Factor 1, Srebf1, and the Sterol Regulatory Element Binding

Transcription Factor 2, Srebf2, as well as Nr1h3 which controls *Srebf1* expression. Overall, the

derived time concordant events, which take place between the beginning of treatment and onset of

adverse histopathology, hence include known and plausible events in liver injury which can be

further characterized based on their frequency, significance and logFC.

Fig 8: Highest ranking events by time concordance metrics. The ten transcription factor (TF)- and pathway-level events ranking highest by enrichment p-value, median max. |logFC| and true positive rate (TPR) before or at histopathology is shown.

Identifying mechanistic hypotheses combining known TF functions and time concordance

While both pathways and TFs constitute interpretable events in this study, further prior knowledge is available on how TFs can function on a molecular level allowing us to derive more detailed hypothesis. Firstly, TF activity can generally be modulated through changes in expression or in post-transcriptional regulation as consequence of cellular signaling or environmental changes. In case of transcriptional regulation, changes in mRNA levels should precede changes in TF activity estimated based on regulon expression and hence time concordance can be used to gain support for

transcriptional TF regulation. Being only interested in TF events with a potential mechanistic link to

liver injury, we studied how significantly concordant expression and activity for each TF are enriched

before adverse histopathology. The strongest evidence for a role in DILI pathogenesis is found for

18 TF events which show both significantly enriched TF expression and regulon activity, providing

complementary evidence of TF importance and hinting at transcriptional regulation (Fig 9A). While

this is not the case for the 17 TF events which only show significantly enriched TF activity but

insignificant enrichment of differential expression, including increased E2f2 activity, this can be

explained by post-transcriptional regulation potentially describing earlier response patterns which

are a direct consequence of upstream signaling. In contrast, 35 TF events with only significant gene

expression, such as increased Jun or Myc, might be already showing changes in expression but not

sufficiently large changes in activity yet. As this rather indicates a role in later pathogenesis and

expression is only regarded as supporting evidence, these TFs have not been included in the next

analysis steps.

Fig 9: Transcription Factor (TF) activity and expression before adverse histopathology. A) Significance of enrichment in adverse conditions for matched TF activity and expression-based events. Events only found on the expression or TF level are not included in the figure due to the inability to perform a statistical test for those. B) For significantly enriched TF activity-based events, the True Positive Rate (TPR) of observing TF activity before or at the time of adverse histopathology is shown, as well as the TPR for observing TF expression changes before TF activity in the time series where it precedes adverse histopathology.

To derive stronger mechanistic evidence for induction we next evaluated how frequently expression changes precede TF activity in the same adverse time-series and compare this against the overall frequency of TF event occurrences preceding adverse histopathology (Fig 9B). Among the events with significant enrichment of TF expression and activity, the most frequent evidence for induction was found for the down-regulation of CCAAT/enhancer-binding protein alpha (Cebpa). In humans, decreased expression of the homologous CEBPA is not only known across liver diseases, exogenously increased CEBPA expression has also been shown to reverse liver injury and is explored as therapeutic target in hepatocellular carcinoma [55]. The event with the 2nd highest relative frequency of expression preceding TF activity as well as the highest frequency of TF activity preceding injury is Atf4, for which expression of the homologous gene in humans is known to be induced as part of the ER stress response contributing to adverse liver phenotypes [56,57]. In contrast, it was found that for the Aryl Hydrogen Receptor (Ahr) and the Peroxisome Proliferator-activated Receptor Alpha (Ppara) changes in expression never preceded those in TF activity which aligns with their roles as nuclear receptors which are generally post-translationally activated via ligand binding [58,59]. As this provides counterevidence for transcriptional induction, these were not included as induced TF in the subsequent analysis.

After investigating the mode of regulation for individual TFs above, we next considered how these TFs are interlinked. To this end, we identified protein-protein interactions and, for induced TFs, TF-target gene interactions between significantly enriched TFs, which showed significant enrichment before adverse histopathology for both expression and regulon activity, as well as evidence of expression preceding TF activity within the same adverse time series. Results of this analysis are shown in Fig 10, and details on the observed absolute and relative frequencies, as well as the

source of the interaction are shown in S5 Table. One of the two identified interactions by highest

absolute frequency is Nr1h3 down-regulation resulting in reduced Srebf1 activity. Furthermore,

Srebf1 is also linked to upstream regulation by Nr1h2 which interacts with Peroxisome Proliferator-

Activated Receptor (Ppara) in both directions, and this cross-talk between Ppara and LXR

regulating Srebf1 expression has been explicitly studied in the context of fatty acid metabolism

regulation [60–62]. The 2^{nd} most frequently observed interaction is the down-regulation of

Transcription Factor 12 (Tcf12) inducing reduced activity of TEA Domain Transcription Factor 1

(Tead1). While Tead1 is indeed known to be involved in liver diseases and injury [63,64], the

interaction itself has not been reported before in the context of liver injury and the same applies also

for the other upstream Tead1 regulators identified. It should also be noted that for these interactions

first activation is only found at the same time but not in the time-concordant order providing weaker

evidence than, for example, the interaction between Nr1h3 and Srebf1. As additional larger TF

cluster, decreased activity of the Hepatocyte Nuclear Factor 1 (Hnf1a), Retinoic Acid Receptor

alpha (Rara) and Pancreatic And Duodenal Homeobox 1 (Pdx1) was found to lead to decreased

expression and activity of Hepatocyte Nuclear Factor 14 (Hnf4a) which is linked to reduced

expression and activity of CCAAT/enhancer-binding protein (Cebpa) through edges in both

directions. This cluster stands out due to the high confidence score of all interactions except the

edge between Pdx1 and Hnf4a indicating that there is strong support based on prior knowledge for

the involved interactions. Furthermore, it was previously found that artificially increased expression

of Hnf4a is able to reverse hepatic liver failure in rats, while also restoring expression of a highly

interconnected TF network including Hnf1a and Cebpa which supports the identified interactions

[65,66]. Two of the yet unknown TFs in DILI are Meis Homeobox 1 (Meis1) and Meis Homeobox 2

(Meis2) which are generally known in a developmental context [67,68]. However, their down-

regulation in early pathogenesis is supported by enriched TF activity, differential expression before

adverse histopathology as well as upstream regulators which are also enriched before adverse

histopathology.

Fig 10: Causal relationships between TFs supported by time concordance. For TFs which are significantly enriched before or at adverse histopathology, known causal relations are shown in which the upstream event is found before or at the downstream event in at least 20% of adverse cases. For induced TFs for which expression is found before regulon activity and significantly enriched, not only protein-protein interactions are considered but also upstream TF-target gene interactions annotated with DoRothEA[69] confidence scores (A: High confidence, C: Medium confidence).

Time-concordant events reflecting disease progression

While events do not have to be activated continuously to be causally involved in pathogenesis, events with consistent or increasing activation over time are particularly interesting as biomarkers as they can be experimentally measured without the chance of missing the timepoint of activation, and can potentially reflect disease progression beyond early pathogenesis. We therefore studied which TFs and pathways show time-dependent activation by testing for significant Spearman correlation between activation logFC and time in adverse time-series, and whether this overlaps with the previously derived time concordance (Fig 11). Overall, 118 pathways and 19 TFs were supported by both, significant time concordance and dependence, which represents 86.1% or 70.4% of the time concordant events, and 59.9% or 48.7% of the time-dependent events, respectively.

On the pathway level, multiple genesets pointed to a reduced level plasma lipoprotein particle assembly and remodelling which indicates changes in lipid distribution. This aligns with the known dyslipidaemia in chronic liver diseases, including decreasing serum values of LDL, HDL, total cholesterol, and triglycerides with increasing severity of disease, based on which previous studies suggested that routine monitoring of lipid profiles can improve the outcome for CLD patients [70]. Furthermore, a down-regulation of response to metal ions was found which could be related to

metallothioneins which protect against oxidative stress and are able to chelate heavy metals [71].

Both directions of dysregulation were previously observed in liver diseases: While a negative

correlation with disease progression was found in hepatocellular carcinoma [72], a positive

correlation was found in most other liver diseases including acetaminophen-induced liver injury [73].

This indicates that opposite directionality is more plausible based on current literature knowledge,

but cannot be fully clarified. The most time-concordant and -dependent TF event was down-

regulation of SRY-Box Transcription Factor 13 (Sox13) which is generally involved in cell fate [74]

and embryonal development [75]. As Sox13 does not yet have well understood functions on a more

detailed level, experimental validation of a potential role in DILI would be interesting. In contrast, the

next most significant time dependence is found for the hepatocyte nuclear factors Hnf1a and Hnf4a,

as well as Cebpa, which are known to negatively correlate with liver cirrhosis in rats [76,77]. Overall,

this shows that a mechanistic role for time-concordant and -dependent events is strongly supported

by the understanding of adverse liver phenotypes.

Fig 11: Combining time dependence and concordance to identify mechanistically supported biomarkers. A) The relation between time concordance, quantified by the enrichment p-value for event activation before adverse histopathology, and time dependence, quantified by the meta p-value for Spearman correlation between time and event activation across adverse conditions, is shown. B) For events with the most significant time-dependence, the distribution of correlation coefficients is shown providing further insight into the strength of correlation and consistency across adverse conditions.

While in general events with highly significant time dependence also showed highly significant time

concordance, some exceptions were found in which only one of both was highly significant. For

instance, the pathway with the 2nd most significant time-dependence (p-value < 10^{44}) is signaling via

advanced glycosylation end product receptor (RAGE) which contributes to inflammation and

oxidative stress generation and did not pass the significance threshold for time concordance (p-

value = 0.058). RAGE expression and activity, which are both induced by binding of RAGE ligands,

are thereby known to be up-regulated in various hepatic disorders resulting in a positive feedback

loop explaining increasing or sustained RAGE activation [78]. This indicates that, while RAGE

signaling is correlated with progression, there is no clear evidence for a role in early pathogenesis

preceding adverse histopathological changes. In contrast, SUMOylation of TFs, is time-concordant

(p-value = 0.002) but not -dependent (p-value = 0.48) indicating a mechanistic role in early

pathogenesis which is not sustained over time. This aligns with the finely regulated and pleiotropic

roles of SUMOylation in post-transcriptional regulation which have also been found to be involved in

the context of liver diseases [79].

Limitations of this study

In this study, we introduced a time-concordance based approach to derive mechanistic insight from

gene expression and histopathology data. We were able to recover known mechanisms in DILI as

well as able to propose novel and detailed mechanistic hypotheses. However, the present analysis

is based on a limited number time-series as well as only few timepoints within each time-series. This

does not only mean that rare events might be missed as they occur between measured timepoints

and that small effects might not be identified as significant, but also that there is potentially a bias

based on the tested compounds towards the represented modes of toxicity.

Furthermore, the analysis is limited by how confidently biological processes are inferred from the

data. This was for instance demonstrated by the differences between pathway and TF activation for

signalling and stress response pathways highlighting the discrepancy between protein activation

and gene expression. As only pathways induced through changes in gene expression or their

downstream expression footprints [69] can be confidently detected, this means that good estimates

of time concordance can predominantly be derived for intermediate or later key events while

preceding key events or molecular initiating events which are not mediated by transcriptional

regulation cannot be estimated based on the data. That being said, this is a limitation of gene

expression data in general and the time concordance approach would also be able to integrate
other data types describing events not covered yet.

Moreover, multiple choices were made to align our analysis to the AOP concept prioritizing
mechanisms supported by prior knowledge over purely data-driven hypothesis. First, detailed
insights might be lost by summarising results to the pathway level. While generally measurements
for individual genes can be noisy, this can be summarised in different ways e.g. based on similarity
in expression profiles [15]. In this study, however, we used curated gene sets due to their
interpretability and to derive modular events as defined in the AOP framework. Additionally, prior
knowledge was taken as ground truth, both in the gene set and interaction analysis, meaning that
only generally known pathways and interactions could be discovered. Like all methods based on
curated gene set and interactions, it was hence informed and biased by the current understanding
of biology. However, this prior biological knowledge contributes to the biological plausibility of the
derived events and relationships contributing to the weight of evidence of our findings in the context
of AOPs.

Lastly, it should be highlighted that time concordance is necessary for casual relations but not
sufficient to prove it. For instance, two events may be time-concordant because they are causally
linked to a shared preceding cause. To distinguish these effects, the additional Bradford-Hill
considerations can be helpful, but only prior knowledge has been considered in parts of this study.
In particular essentiality would provide strong evidence for causality, however, requires targeted
experiments and hence is unsuitable for hypothesis generation. In contrast, dose and incidence
concordance are generally feasible from a data-driven standpoint but were not pursued in this case
study due to the low number of doses and replicates.

Conclusion

In this study, we introduce "first activation" as concept to quantify the strength of temporal concordance between events across time series with the assumption that each activated event may have downstream effects irrespective of whether it is continuously or only transiently activated. With this approach, we study gene expression-based TF and pathway-level events found before adverse histopathology indicating liver injury in repeat-dose studies in rats from TG-GATEs as a case study. We find some known processes in DILI to be highly confident, e.g. bile acid recycling, while others are highly frequent but less specific including adaptive response pathways such as the eIF2α/ATF4 pathway [51].

Beyond quantifying time concordance for known and potentially novel events in DILI, we additionally show how time concordance can be combined with prior biological knowledge to generate hypothesis on potentially causal gene-regulatory cascades in DILI. Amongst others, this identifies LXRα down-regulation leading to decreased Srebf1 expression, an interaction known to regulate fatty acid synthesis in the liver [41], but also characterizes yet unknown TFs based on their time concordance, their mode of regulation (either transcriptional or post-transcriptional) and potential upstream regulators and downstream effectors. Two of the identified induced TFs are Meis1 and Meis 2 which is supported by significantly enriched decrease in expression and activity before adverse histopathology, as well as upstream regulators which also show significant enrichment of regulon activity and are found within the same time series. On top of time concordance, we also derive each event's time dependence and show that events mechanistically involved in early pathogenesis do not necessarily reflect disease progression and vice versa. However, for some events, e.g. Sox13, both properties are found and these may be useful biomarkers which reflect injury progression and already change preceding histopathological manifestation.

We believe that the described analysis can provide supporting evidence for mechanistic links

between events in line with the evolved Bradford-Hill considerations on time concordance and

biological plausibility and can hence e.g. support AOP development. Furthermore, the approach is

not limited to a particular adverse event and can instead quantify the interaction between any two

events represented in time series in a data-driven and automatable fashion. Consequentially, this

type of analysis could also be of interest to study the mechanism of action particular compound

classes or patterns of disease progression. We make the results of our analysis on the TG-GATEs

in vivo liver data publicly available in a Shiny app through which users can query the most time-

concordant events for more specific types of histopathology and study in detail in which time series

time concordance was observed or not observed (https://anikaliu.shinyapps.io/dili_cascades).

Methods

Open TG-GATES data processing

The TG-GATES gene expression data from studies in 6-week-old male Crl:CD Sprague-Dawley

(SD) rats with daily repeat-dosing (S1 Fig) was downloaded from the Life Science Data Archive

(DOI: 10.18908/lsdba.nbdc00954-01-000). The raw liver gene expression levels were background

corrected, $\log_2$ transformed, and quantile normalized with the rma function of the affy package per

treatment across all doses and timepoints [80]. Quality control was then performed using the

ArrayQualityMetrics package [81] and detected outliers with high distance to other experiments or

unusual signal distribution were removed (List of removed outliers summarised in S1 File). The

platform information for the Affymetrix Rat Genome 230 2.0 Array was derived from Gene

Expression Omnibus [82] (GEO accession: GPL1355) and was then used to summarise probe IDs

to rat gene symbols by median for all probes mapping uniquely to one gene symbol. Only the 360

compound-dose combinations with at least 6 measured timepoints after quality control were

included. Out of the these all eight timepoints were measured in most time series, while only six

timepoints were measured in two time series, and only seven timepoints in seven time series.

Definition of adverse histopathology

To characterize the extent of histological findings, we used the toxscores by Sutherland et al. [15] in

order to consider both severity and frequency of events in a single numerical output measure.

These are based on the lesion severity per animal which was first converted to a numerical scale

(normal = 0, minimal = 1, slight = 2, moderate = 3, marked or severe = 4) and then averaged across

all biological replicates as an aggregate measure for lesion frequency and severity. One

characteristic of this measure is that the overall distributions varied between different findings, e.g.

inflammation was more frequently annotated with low than with high toxscores while a more

balanced distribution of scores was observed for Hepatocellular single cell necrosis (S2 Fig).

To study which histological findings were enriched in adverse conditions, we first defined binary

histopathology labels describing the presence of histological findings with different extents in each

time-series. Based on the toxscore ranges used by Sutherland et al., three toxscore cut-offs are

implemented to describe each histopathological finding "Null" (toxscore > 0), "low" (toxscore > 0.67)

and "high" (toxscore > 1.34). We then studied which labels were over-represented in adverse time-

series. These were defined using the annotation of Sutherland et al., where pathologists classified

compound-dose combinations in the TG-GATEs database as adverse or non-adverse after 4 and

29 days of treatment. We used the 29 days classification to define 40 adverse time series and only

regarded time-series as non-adverse for compounds which were not classified as adverse at any

dose in the negative control, in order to account for the fact that some of the cellular changes of

interest might already take place at lower doses, although the resulting phenotype is not considered

adverse yet.

We defined findings as adverse histopathology if they are observed in at least 5 out of 40 adverse

time-series to remove rare histopathological findings, and additionally require that at least 50% of

findings are in adverse conditions to remove findings which are unspecific. All labels which were

identified with these criteria are significantly enriched among time-series labelled as adverse by

Sutherland et al. in comparison to those that were considered non-adverse using a one-sided

Fisher's Exact test (p-value > 0.0001), performed using the *fisher.test* function of the *stats* R

package [83]. However, this combination of additional criteria was chosen to exclude findings which

are rare or weakly associated.

While not all compounds in the TG-GATEs database are drugs and some mechanisms of toxicity

may not translate to humans, out of the 38 compounds represented in adverse time-series, 22 have

additionally been classified as hepatotoxic in DILIst [28] and 18 in DILIrank (vMost-DILI-Concern or

vLess-DILI-Concern) [27] (S2 Table). This overlap with compound-level DILI annotations by the

FDA shows that the compounds in this study partially represent known mechanisms of DILI in

humans, while also highlighting the fact that a clear classification is not possible.

Pathway and TF activity inference

The activity of pathways and TFs across all doses and timepoints of a treatment including vehicle

controls was derived based on the expression of its gene sets members using GSVA [26], which

computes a gene set enrichment by sample matrix from the gene expression by sample matrix. This

was performed using a Gaussian kernel requiring at least 5 genes per gene set, and overall

provides the basis for the subsequent pathway- and TF-centric steps. As prior knowledge, we used

pathway maps from Reactome [37] which were derived through MSigDB [86] and the msigdbr

package [87]. TF activity gene sets were derived from DoRothEA [88] and mapped from human to

rat gene symbols with biomaRt [89]. These gene sets describe known, functional TF-gene

interactions and are assigned a confidence level based on the strength of evidence of these

interactions. Thereby, only the 207 TFs with a high to medium confidence level of A-C were

included and the few TF-gene interactions with a negative mode of regulation were removed to

better infer TF directionality. To evaluate which pathway or TF is dysregulated, we computed the

differential activity in comparison to the vehicle control group, which was treated for the same

amount of time and as part of the same experiment, using the moderated t-statistic in limma [90].

We identify significantly dysregulated gene sets with a False Discovery Rate (FDR) < 0.05.

Temporal concordance of events

In this study, the order of events was derived based on each event's timepoint of first activation

within each time-series (Fig 1A). For pathways and TFs, we defined first activation as the earliest

time of measurement at which significant differential regulation was observed (FDR < 0.05) in each

direction, while an additional logFC cut-off has been implemented for individual genes. As first

evidence of adverse morphological changes in the liver, we used the first timepoint any of the

adverse histopathology label derived before were found.

We were then generally interested in potential preceding events **PE** which are first activated before

or at the same time as a potential later event or outcome **LE** and used multiple metrics to quantify

the degree of time concordance which can be related to the original work by Bradford Hill (Table 1).

Thereby, the key later event in this study was adverse histopathology but we used a more general

notation **LE**, as some of the following criteria to quantify time concordance are also applied in the

TF analysis, where gene expression-derived events are used as later event. First, we used the true

positive rate (TPR) which describes how frequently **PE** is observed before **LE** among all time-series

with **LE** and hence its consistency across compounds. Secondly, we use the maximal effect size of

PE observed before **LE**, summarised across time-series by median, to characterise the strength of

association. To evaluate the significance of the findings, we additionally defined a set of background

time-series unrelated to **LE** (Fig 1B). For adverse histopathology, these unrelated background time-

series were the 133 time-series without any observed histological changes. We then computed the

enrichment of **PE** before or at **LE** using the *fisher.test* function of the *stats* R package [83], first

estimating the odds ratio using the conditional maximum likelihood estimate and subsequently

testing the null hypothesis whether the odds ratio derived from a confusion matrix as described in

Fig 1 is equal to or smaller than 1.

and the positive predictive value (PPV) of **PE** for **LE**, which describes how likely **LE** is observed at

the same or a later time given the observation of **PE.**

Across all metrics, we only consider time-series in the statistics for which any event of the same

type as **PE**, e.g. TF or pathway, was observed at the included timepoints, so before or at **LE** or at

any timepoint in the background time-series. We do this to account for the fact that in some cases

no changes are found which may be a consequence of the fact that there isn't a measured timepoint

before **LE** or that at the available timepoints expression changes cannot be detected. We argue that

in these cases this should not be treated as evidence of absence of the given event, but rather as

absence of evidence.

Table 1: Metrics quantifying the time concordance between a potential preceding event *PE*
and potential later event *LE,* and their relation to the original Bradford Hill (BH)
considerations.

BH consideration	Metric	Formula	Description
Consistency	True positive rate (TPR)	p(**PE** →**LE**|**LE**)	Fraction of time series with event **PE** with specified temporal relation among time series with event **LE**
Specificity	Positive predictive value (PPV)	p(**PE**→ **LE**|**PE**)	Fraction of time series with event **LE** with specified temporal relation among time series with event **PE**
Temporality	p-value	One-sided Fisher's Exact test	Likelihood of observing event **PE** and **LE** with specified temporal relation with equal or higher frequency by chance assuming a hypergeometric distribution.
Strength	Effect size in time series with **LE**	Median (logFC)	Median logFC of **PE** observed in time-series with **LE** (in comparison to vehicle control)

Combining time concordance on TF-TF interactions

We used 3 sources of causal prior knowledge to derive mechanistic hypotheses linking TFs:

Protein-protein interaction between TFs derived from Omnipath through OmnipathR [91,92], TF-

target gene interactions from DoRothEA [88] and the link between gene expression and protein

levels following the central dogma of molecular biology. Using these interactions as backbone, we

then derived those additionally supported by time concordance. Thereby, the dysregulation of the

nodes was required to match the reported mode of regulation (edge sign) and the source node or

upstream event was required to be observed in at least 20% of cases before or at the same time as

the target node or downstream event. For induced TFs, significant enrichment of gene expression

(|logFC|>0.5) and TF activity before adverse histopathology was required, as well as evidence for

changes in expression preceding changes in the same direction in regulon activity within the same

time series.

Time dependence

In each adverse time-series, we tested for Spearman correlation between timepoint and event

activation logFC using the correlation R package [93], and include a logFC of 0 at timepoint 0 h

assuming that there are no differences in comparison to the control group before treatment. We

then identify pathways and TFs which only show significant Spearman correlation in one direction,

positive or negative. For those events, we apply the Fisher's combined probability test using the

metap R package [94] across all adverse time-series to evaluate whether overall significant

correlation between event activation and time is found.

Supporting information

S1 Table. Comparison of quantitative Adverse Outcome Pathway (qAOP) models. Comparison of the first activation concept and other qAOP models with respect to their potential roles in AOP development.

S2 Table. Compounds classified as adverse based on histopathology and concordance with previous annotations. For the annotations by Sutherland et al. [15], who classified each compound at each measured dose as adverse or non-adverse at day 4 and day 29, the adverse doses for each compound are listed. Furthermore, the binary classification as adverse (1) and non-adverse (0) from DILIst [84] are included as well as the vDILIConcern and Severity Class classifications from DILIRank [85] which describe evidence for liver side effects observed in humans derived from post-marketing data.

S3 Table. Time concordance metrics for Reactome pathway maps which map to known key events based on literature review.

S4 Table. Time concordance metrics for top 10 ranking events by True Positive Rate (TPR), significance and median max. |logFC|.

S5 Table. TF-TF relations supported by known relations and time concordance. For TF events which are significantly enriched before or at adverse histopathology, known interactions supported by time concordance are shown. With respect to the interaction, the absolute and relative frequency are shown for how often the source TF was observed "before" or "before or at" downstream TF activity. Additionally, the source of the interactions provided in Omnipath are shown for protein-protein interactions and the DoRothEA confidence level for TF-target gene interactions.

S1 Fig. Open TG-GATEs study design. 6-week-old male Crl:CD Sprague-Dawley (SD) rats were treated with a range of compounds using daily repeat-dosing. For each compound, four doses were used including a vehicle control, and samples were taken at 8 timepoints. For each combination of compound, timepoint and dose, histopathology was annotated and gene expression measured for 3 replicates.

S2 Fig. Distribution of toxscores across histopathological findings.

S3 Fig. Frequency of histopathological findings before and after first adverse histopathology. For adverse and non-adverse histopathological findings the frequency before or at first non-adverse histopathology is shown (left). For adverse findings, this indicates how frequently they were one of the first adverse histopathological findings given that they cannot occur before by definition. This identifies single-cell necrosis at any severity ("null"), as the most frequent finding, both in absolute and relative terms.

S4 Fig. Background distribution of temporal association metrics across pathway and Transcription Factor (TF) events. The dependency between different metrics is shown. A) Frequency of events by median max. |logFC| before or at histopathology and enrichment p-value. B) Frequency of events by true positive rate (TPR) and positive predictive value (PPV) before or at adverse histopathology. C) Direct relation between TPR, PPV and enrichment p-value. D) Direct relation between TPR, PPV and frequency in background time-series.

derive low p-values for events with high TPR and high PPV. Given a number of event
activations in non-adverse control time series, there is a monotonous relationship between
PPV and TPR with increasing PPV at lower event frequency in non-adverse control time
series.

S1 File. Removed outlier samples.

S2 File. Time concordance metrics for all TFs, pathways as well as genes using both a
minimal |logFC| of 0.5 and 1.

References

1.　Hwang TJ, Carpenter D, Lauffenburger JC, Wang B, Franklin JM, Kesselheim AS. Failure of

investigational drugs in late-stage clinical development and publication of trial results. JAMA

Internal Medicine. 2016;176: 1826–1833. doi:10.1001/jamainternmed.2016.6008

2.　Harrison RK. Phase II and phase III failures: 2013-2015. Nature Reviews Drug Discovery.

2016;15: 817–818. doi:10.1038/nrd.2016.184

3.　Bai JPF, Abernethy DR. Systems Pharmacology to Predict Drug Toxicity: Integration Across

Levels of Biological Organization *. 2012 [cited 5 Jun 2019]. doi:10.1146/annurev-pharmtox-

011112-140248

4.　Leist M, Ghallab A, Graepel R, Marchan R, Hassan R, Bennekou SH, et al. Adverse outcome

pathways: opportunities, limitations and open questions. Archives of Toxicology. 2017;91:

3477–3505. doi:10.1007/s00204-017-2045-3

5.　Ankley GT, Bennett RS, Erickson RJ, Hoff DJ, Hornung MW, Johnson RD, et al. Adverse

outcome pathways: A conceptual framework to support ecotoxicology research and risk

assessment. Environmental Toxicology and Chemistry Wiley Blackwell; Mar 1, 2010 pp. 730–

741. doi:10.1002/etc.34

6.　OECD (Organisation for Economic Co-operation and Development). USERS' HANDBOOK

SUPPLEMENT TO THE GUIDANCE DOCUMENT FOR DEVELOPING AND ASSESSING

804  AOPs. OECD Environment, Health and Safety Publications Series on Testing and
805  Assessment. 2018; 1–62.

806 7. Hill AB. The Environment and Disease: Association or Causation? Proc R Soc Med. 1965;
807  295–300.

808 8. Meek ME, Palermo CM, Bachman AN, North CM, Lewis RJ, Meek B, et al. Mode of action
809  human relevance (species concordance) framework: Evolution of the Bradford Hill
810  considerations and comparative analysis of weight of evidence. Journal of Applied Toxicology
811  John Wiley and Sons Ltd; 2014 pp. 595–606. doi:10.1002/jat.2984

812 9. Oki NO, Nelms MD, Bell SM, Mortensen HM, Edwards SW. Accelerating Adverse Outcome
813  Pathway Development Using Publicly Available Data Sources. Current Environmental Health
814  Reports. 2016;3: 53–63. doi:10.1007/s40572-016-0079-y

815 10. Oki NO, Edwards SW. An integrative data mining approach to identifying adverse outcome
816  pathway signatures. Toxicology. 2016;350–352: 49–61. doi:10.1016/J.TOX.2016.04.004

817 11. Bell SM, Angrish MM, Wood CE, Edwards SW. Integrating publicly available data to generate
818  computationally predicted adverse outcome pathways for fatty liver. Toxicological Sciences.
819  2016;150: 510–520. doi:10.1093/toxsci/kfw017

820 12. Burgoon LD, Angrish M, Garcia-Reyero N, Pollesch N, Zupanic A, Perkins E. Predicting the
821  Probability that a Chemical Causes Steatosis Using Adverse Outcome Pathway Bayesian
822  Networks (AOPBNs). Risk Analysis. 2020;40: 512–523. doi:10.1111/risa.13423

823 13. Jeong J, Garcia-Reyero N, Burgoon L, Perkins E, Park T, Kim C, et al. Development of
824  Adverse Outcome Pathway for PPARγAntagonism Leading to Pulmonary Fibrosis and
825  Chemical Selection for Its Validation: ToxCast Database and a Deep Learning Artificial

Neural Network Model-Based Approach. Chemical Research in Toxicology. 2019;32: 1212–

1222. doi:10.1021/acs.chemrestox.9b00040

14.   Kohonen P, Parkkinen JA, Willighagen EL, Ceder R, Wennerberg K, Kaski S, et al. A

transcriptomics data-driven gene space accurately predicts liver cytopathology and drug-

induced liver injury. Nature Communications. 2017;8. doi:10.1038/ncomms15932

15.   Sutherland JJ, Webster YW, Willy JA, Searfoss GH, Goldstein KM, Irizarry AR, et al.

Toxicogenomic module associations with pathogenesis: A network-based approach to

understanding drug toxicity. Pharmacogenomics Journal. 2018;18: 377–390.

doi:10.1038/tpj.2017.17

16.   Souza TM, Kleinjans JCS, Jennen DGJ. Dose and Time Dependencies in Stress Pathway

Responses during Chemical Exposure: Novel Insights from Gene Regulatory Networks.

Frontiers in Genetics. 2017;8: 142–142. doi:10.3389/fgene.2017.00142

17.   Rooney J, Hill T, Qin C, Sistare FD, Christopher Corton J. Adverse outcome pathway-driven

identification of rat liver tumorigens in short-term assays. Toxicology and Applied

Pharmacology. 2018;356: 99–113. doi:10.1016/j.taap.2018.07.023

18.   Rooney J, Oshida K, Vasani N, Vallanat B, Ryan N, Chorley BN, et al. Activation of Nrf2 in

the liver is associated with stress resistance mediated by suppression of the growth

hormone-regulated STAT5b transcription factor. PLoS ONE. 2018;13.

doi:10.1371/journal.pone.0200004

19.   Andrade RJ, Chalasani N, Björnsson ES, Suzuki A, Kullak-Ublick GA, Watkins PB, et al.

Drug-induced liver injury. Nature Reviews Disease Primers. 2019;5: 1–22.

doi:10.1038/s41572-019-0105-0

20.   Regev A. Drug-induced liver injury and drug development: Industry perspective. Seminars in

849   Liver Disease. 2014;34: 227–239. doi:10.1055/S-0034-1375962/ID/JR00749-45

850   21.   Aguayo-Orozco A, Bois FY, Brunak S, Taboureau O. Analysis of Time-Series Gene

851   Expression Data to Explore Mechanisms of Chemical-Induced Hepatic Steatosis Toxicity.

852   Frontiers in Genetics. 2018;9: 396. doi:10.3389/fgene.2018.00396

853   22.   Zhang JD, Berntenis N, Roth A, Ebeling M. Data mining reveals a network of early-response

854   genes as a consensus signature of drug-induced in vitro and in vivo toxicity.

855   Pharmacogenomics Journal. 2014;14: 208–216. doi:10.1038/tpj.2013.39

856   23.   Zgheib E, Gao W, Limonciel A, Aladjov H, Yang H, Tebby C, et al. Application of three

857   approaches for quantitative AOP development to renal toxicity. Computational Toxicology.

858   2019;11: 1–13. doi:10.1016/J.COMTOX.2019.02.001

859   24.   Hassan I, El-Masri H, Kosian PA, Ford J, Degitz SJ, Gilbert ME. Neurodevelopment and

860   Thyroid Hormone Synthesis Inhibition in the Rat: Quantitative Understanding Within the

861   Adverse Outcome Pathway Framework. Toxicological Sciences. 2017;160: 57–73.

862   doi:10.1093/TOXSCI/KFX163

863   25.   Spinu N, Cronin MTD, Enoch SJ, Madden JC, Worth AP. Quantitative adverse outcome

864   pathway (qAOP) models for toxicity prediction. Archives of Toxicology. Springer; 2020. pp.

865   1497–1510. doi:10.1007/s00204-020-02774-7

866   26.   Hänzelmann S, Castelo R, Guinney J. GSVA: gene set variation analysis for microarray and

867   RNA-Seq data. BMC Bioinformatics. 2013. Available: http://www.biomedcentral.com/1471-

868   2105/14/7http://www.bioconductor.org.Background

27.   Vall A, Sabnis Y, Shi J, Class R, Hochreiter S, Klambauer G. The Promise of AI for DILI

Prediction. Frontiers in Artificial Intelligence. 2021;4: 15.

doi:10.3389/FRAI.2021.638410/BIBTEX

28.   Liu A, Walter M, Wright P, Bartosik A, Dolciami D, Elbasir A, et al. Prediction and mechanistic

analysis of drug-induced liver injury (DILI) based on chemical structure. Biology Direct.

2021;16: 1–15. doi:10.1186/s13062-020-00285-0

29.   Tillander V, Alexson SEH, Cohen DE. Deactivating Fatty Acids: Acyl-CoA Thioesterase-

Mediated Control of Lipid Metabolism. Trends in Endocrinology and Metabolism. Elsevier

Inc.; 2017. pp. 473–484. doi:10.1016/j.tem.2017.03.001

30.   Violante S, Ijlst L, Ruiter J, Koster J, van Lenthe H, Duran M, et al. Substrate specificity of

human carnitine acetyltransferase: Implications for fatty acid and branched-chain amino acid

metabolism. Biochimica et Biophysica Acta - Molecular Basis of Disease. 2013;1832: 773–

779. doi:10.1016/j.bbadis.2013.02.012

31.   Priya Dharshini LC, Vishnupriya S, Sakthivel KM, Rasmi RR. Oxidative stress responsive

transcription factors in cellular signalling transduction mechanisms. Cellular Signalling.

2020;72: 109670. doi:10.1016/J.CELLSIG.2020.109670

32.   Hong L, Sun QF, Xu TY, Wu YH, Zhang H, Fu RQ, et al. New role and molecular mechanism

of Gadd45a in hepatic Fibrosis. World Journal of Gastroenterology. 2016;22: 2779–2788.

doi:10.3748/wjg.v22.i9.2779

33.   Okazaki H, Igarashi M, Nishi M, Sekiya M, Tajima M, Takase S, et al. Identification of neutral

cholesterol ester hydrolase, a key enzyme removing cholesterol from macrophages. J Biol

Chem. 2008;283: 33357–33364. doi:10.1074/JBC.M802686200

34.   Chiba M, Murata S, Myronovych A, Kohno K, Hiraiwa N, Nishibori M, et al. Elevation and
892   characteristics of Rab30 and S100a8/S100a9 expression in an early phase of liver
893   regeneration in the mouse. International Journal of Molecular Medicine. 2011;27: 567–574.
894   doi:10.3892/IJMM.2011.614/HTML

895   35.   Liu A, Trairatphisan P, Gjerga E, Didangelos A, Barratt J, Saez-Rodriguez J. From
896   expression footprints to causal pathways: contextualizing large signaling networks with
897   CARNIVAL. npj Systems Biology and Applications. 2019;5: 1–10. doi:10.1038/s41540-019-
898   0118-z

899   36.   Schubert M, Klinger B, Klünemann M, Sieber A, Uhlitz F, Sauer S, et al. Perturbation-
900   response genes reveal signaling footprints in cancer gene expression. Nature
901   Communications. 2018;9: 20. doi:10.1038/s41467-017-02391-6

902   37.   Fabregat A, Jupe S, Matthews L, Sidiropoulos K, Gillespie M, Garapati P, et al. The
903   Reactome Pathway Knowledgebase. Nucleic Acids Research. 2018;46: D649–D655.
904   doi:10.1093/nar/gkx1132

905   38.   Shi Z, Zhang K, Chen T, Zhang Y, Du X, Zhao Y, et al. Transcriptional factor ATF3 promotes
906   liver fibrosis via activating hepatic stellate cells. Cell Death & Disease 2020 11:12. 2020;11:
907   1–16. doi:10.1038/s41419-020-03271-6

908   39.   Copple IM, Goldring CE, Jenkins RE, Chia AJL, Randle LE, Hayes JD, et al. The hepatotoxic
909   metabolite of acetaminophen directly activates the Keap1-Nrf2 cell defense system.
910   Hepatology. 2008;48: 1292–1301. doi:10.1002/hep.22472

911   40.   Luedde T, Schwabe RF. NF-κB in the liver-linking injury, fibrosis and hepatocellular
912   carcinoma. Nature Reviews Gastroenterology and Hepatology. NIH Public Access; 2011. pp.
913   108–118. doi:10.1038/nrgastro.2010.213

914   41.   Schultz JR, Tu H, Luk A, Repa JJ, Medina JC, Li L, et al. Role of LXRs in control of

915   lipogenesis. Genes and Development. 2000;14: 2831–2838. doi:10.1101/gad.850400

916   42.   Hetz C, Zhang K, Kaufman RJ. Mechanisms, regulation and functions of the unfolded protein

917   response. Nature Reviews Molecular Cell Biology. Nature Research; 2020. pp. 421–438.

918   doi:10.1038/s41580-020-0250-z

919   43.   Wijaya LS, Trairatphisan P, Gabor A, Niemeijer M, Keet J, Alcalà Morera A, et al. Integration

920   of temporal single cell cellular stress response activity with logic-ODE modeling reveals

921   activation of ATF4-CHOP axis as a critical predictor of drug-induced liver injury. Biochemical

922   Pharmacology. 2021;190: 114591. doi:10.1016/j.bcp.2021.114591

923   44.   Fredriksson L, Wink S, Herpers B, Benedetti G, Hadi M, de Bont H, et al. Drug-induced

924   endoplasmic reticulum and oxidative stress responses independently sensitize toward TNFα-

925   mediated   hepatotoxicity.   Toxicological   Sciences.   2014;140:   144–159.

926   doi:10.1093/toxsci/kfu072

927   45.   Seki E, Brenner DA, Karin M. A liver full of JNK: Signaling in regulation of cell function and

928   disease pathogenesis, and clinical approaches. Gastroenterology. 2012;143: 307–320.

929   doi:10.1053/j.gastro.2012.06.004

930   46.   Win S, Than TA, Zhang J, Oo C, Min RWM, Kaplowitz N. New insights into the role and

931   mechanism of c-Jun-N-terminal kinase signaling in the pathobiology of liver diseases.

932   Hepatology. John Wiley and Sons Inc.; 2018. pp. 2013–2024. doi:10.1002/hep.29689

933   47.   Simmons SO, Fan C-Y, Ramabhadran R. Cellular Stress Response Pathway System as a

934   Sentinel Ensemble in Toxicological Screening. Toxicological Sciences. 2009;111: 202–225.

935   doi:10.1093/TOXSCI/KFP140

48. Wong MMK, Joyson SM, Hermeking H, Chiu SK. Transcription factor AP4 mediates cell fate

decisions: To divide, age, or die. Cancers. MDPI AG; 2021. pp. 1–15.

doi:10.3390/cancers13040676

49. Zuo L, Tan T, Wei C, Wang H, Tan L, Hao Y, et al. HOXB13 expression is correlated with

hepatic inflammatory activity of patients with hepatic fibrosis. J Mol Histol. 2020;51: 183–189.

doi:10.1007/S10735-020-09868-7

50. Delgado I, Fresnedo O, Iglesias A, Rueda Y, Syn WK, Zubiaga AM, et al. A role for

transcription factor E2F2 in hepatocyte proliferation and timely liver regeneration. American

Journal of Physiology - Gastrointestinal and Liver Physiology. 2011;301.

doi:10.1152/ajpgi.00481.2010

51. B'Chir W, Maurin AC, Carraro V, Averous J, Jousse C, Muranishi Y, et al. The eIF2α/ATF4

pathway is essential for stress-induced autophagy gene expression. Nucleic Acids Research.

2013;41: 7683–7699. doi:10.1093/nar/gkt563

52. Wijaya LS, Trairatphisan P, Gabor A, Niemeijer M, Keet J, Alcalà Morera A, et al. Integration

of temporal single cell cellular stress response activity with logic-ODE modeling reveals

activation of ATF4-CHOP axis as a critical predictor of drug-induced liver injury. Biochemical

Pharmacology. 2021;190: 114591. doi:10.1016/J.BCP.2021.114591

53. Pavel M, Imarisio S, Menzies FM, Jimenez-Sanchez M, Siddiqi FH, Wu X, et al. CCT

complex restricts neuropathogenic protein aggregation via autophagy. Nature

Communications. 2016;7: 1–18. doi:10.1038/ncomms13821

54. Grantham J. The Molecular Chaperone CCT/TRiC: An Essential Component of Proteostasis

and a Potential Modulator of Protein Aggregation. Frontiers in Genetics. Frontiers Media S.A.;

2020. doi:10.3389/fgene.2020.00172

55.   Reebye V, Huang KW, Lin V, Jarvis S, Cutilas P, Dorman S, et al. Gene activation of CEBPA

using saRNA: Preclinical studies of the first in human saRNA drug candidate for liver cancer.

Oncogene. 2018;37: 3216–3228. doi:10.1038/s41388-018-0126-2

56.   Fusakio ME, Willy JA, Wang Y, Mirek ET, Baghdadi RJTA, Adams CM, et al. Transcription

factor ATF4 directs basal and stress-induced gene expression in the unfolded protein

response and cholesterol metabolism in the liver. Molecular Biology of the Cell. 2016;27:

1536–1551. doi:10.1091/mbc.E16-01-0039

57.   Hao L, Zhong W, Dong H, Guo W, Sun X, Zhang W, et al. ATF4 activation promotes hepatic

mitochondrial dysfunction by repressing NRF1-TFAM signalling in alcoholic steatohepatitis.

Gut. 2020 [cited 6 May 2021]. doi:10.1136/gutjnl-2020-321548

58.   Larigot L, Juricek L, Dairou J, Coumoul X. AhR signaling pathways and regulatory functions.

Biochimie Open. 2018;7: 1–9. doi:10.1016/J.BIOPEN.2018.05.001

59.   Kersten S, Stienstra R. The role and regulation of the peroxisome proliferator activated

receptor   alpha   in   human   liver.   Biochimie.   2017;136:   75–84.

doi:10.1016/J.BIOCHI.2016.12.019

60.   Yoshikawa T, Ide T, Shimano H, Yahagi N, Amemiya-Kudo M, Matsuzaka T, et al. Cross-talk

between peroxisome proliferator-activated receptor (PPAR) α and liver X receptor (LXR) in

nutritional regulation of fatty acid metabolism. I. PPARS suppress sterol regulatory element

binding protein-1c promoter through inhibition of LXR signaling. Molecular Endocrinology.

2003;17: 1240–1254. doi:10.1210/ME.2002-0190

61.   Yoshikawa T, Shimano H, Amemiya-Kudo M, Yahagi N, Hasty AH, Matsuzaka T, et al.

Identification of Liver X Receptor-Retinoid X Receptor as an Activator of the Sterol

Regulatory Element-Binding Protein 1c Gene Promoter. Molecular and Cellular Biology.

2001;21: 2991–3000. doi:10.1128/mcb.21.9.2991-3000.2001

62. Boergesen M, Pedersen TA, Gross B, van Heeringen SJ, Hagenbeek D, Bindesboll C, et al.

Genome-Wide Profiling of Liver X Receptor, Retinoid X Receptor, and Peroxisome

Proliferator-Activated Receptor in Mouse Liver Reveals Extensive Sharing of Binding Sites.

Molecular and Cellular Biology. 2012;32: 852–867. doi:10.1128/mcb.06175-11

63. Kusumanchi P, Liang T, Zhang T, Ross RA, Han S, Chandler K, et al. Stress-Responsive

Gene FK506-Binding Protein 51 Mediates Alcohol-Induced Liver Injury Through the Hippo

Pathway and Chemokine (C-X-C Motif) Ligand 1 Signaling. Hepatology. 2021 [cited 6 Sep

2021]. doi:10.1002/HEP.31800

64. Manmadhan S, Ehmer U. Hippo Signaling in the Liver – A Long and Ever-Expanding Story.

Frontiers in Cell and Developmental Biology. 2019;7: 33. doi:10.3389/FCELL.2019.00033

65. Kyrmizi I, Hatzis P, Katrakili N, Tronche F, Gonzalez FJ, Talianidis I. Plasticity and expanding

complexity of the hepatic transcription factor network during liver development. Genes and

Development. 2006;20: 2293–2305. doi:10.1101/GAD.390906

66. Nishikawa T, Bell A, Brooks JM, Setoyama K, Melis M, Han B, et al. Resetting the

transcription factor network reverses terminal chronic hepatic failure. The Journal of Clinical

Investigation. 2015;125: 1533. doi:10.1172/JCI73137

67. Schulte D, Geerts D. MEIS transcription factors in development and disease. Development.

2019;146. doi:10.1242/DEV.174706

68. Berenguer M, Duester G. Role of Retinoic Acid Signaling, FGF Signaling and Meis Genes in

Control of Limb Development. Biomolecules. 2021;11: 1–11. doi:10.3390/BIOM11010080

69.   Garcia-Alonso L, Holland CH, Ibrahim MM, Turei D, Saez-Rodriguez J. Benchmark and

1004   integration of resources for the estimation of human transcription factor activities. Genome

1005   Research. 2019;29: 1363–1375. doi:10.1101/gr.240663.118

1006   70.   Farooque U, Lohano AK, Dahri Q, Arain N, Farukhuddin F, Khadke C, et al. The Pattern of

1007   Dyslipidemia   in   Chronic   Liver   Disease   Patients.   Cureus.   2021;13.

1008   doi:10.7759/CUREUS.13259

1009   71.   Oliva L, D'Incà R, Medici V, Sturniolo GC. Metallothioneins and liver diseases.

1010   Metallothioneins   in   Biochemistry   and   Pathology.   2008;   289–316.

1011   doi:10.1142/9789812778949_0014

1012   72.   Huang G-W, Yang L-Y. Metallothionein expression in hepatocellular carcinoma. World

1013   Journal of Gastroenterology. 2002;8: 650. doi:10.3748/WJG.V8.I4.650

1014   73.   Devisscher L, Campenhout S van, Lefere S, Raevens S, Tilleman L, Nieuwerburgh F van, et

1015   al. Metallothioneins alter macrophage phenotype and represent novel therapeutic targets for

1016   acetaminophen-induced   liver   injury.   Journal   of   Leukocyte   Biology.   2021;   1–11.

1017   doi:10.1002/JLB.3A0820-527R

1018   74.   Lefebvre V. The SoxD transcription factors – Sox5, Sox6, and Sox13 – are key cell fate

1019   modulators. The International Journal of Biochemistry & Cell Biology. 2010;42: 429–432.

1020   doi:10.1016/J.BIOCEL.2009.07.016

1021   75.   Wang Y, Ristevski S, Harley VR. SOX13 exhibits a distinct spatial and temporal expression

1022   pattern   during   chondrogenesis,   neurogenesis,   and   limb   development.   Journal   of

1023   Histochemistry and Cytochemistry. 2006;54: 1327–1333. doi:10.1369/jhc.6A6923.2006

1024   76.   Liu L, Yannam GR, Nishikawa T, Yamamoto T, Basma H, Ito R, et al. The microenvironment

1025   in hepatocyte regeneration and function in rats with advanced cirrhosis. Hepatology. 2012;55:

1026   1529–1539. doi:10.1002/HEP.24815

1027   77.   Guzman-Lepe J, Cervantes-Alvarez E, l'Hortet AC de, Wang Y, Mars WM, Oda Y, et al.

1028   Liver-enriched transcription factor expression relates to chronic hepatic failure in humans.

1029   Hepatology Communications. 2018;2: 582–594. doi:10.1002/HEP4.1172

1030   78.   Yamagishi S, Matsui T. Role of receptor for advanced glycation end products (RAGE) in liver

1031   disease. European Journal of Medical Research. 2015;20. doi:10.1186/S40001-015-0090-Z

1032   79.   Zeng M, Liu W, Hu Y, Fu N. Sumoylation in liver disease. Clinica Chimica Acta. Elsevier;

1033   2020. pp. 347–353. doi:10.1016/j.cca.2020.07.044

1034   80.   Gautier L, Cope L, Bolstad BM, Irizarry RA. Affy - Analysis of Affymetrix GeneChip data at the

1035   probe level. Bioinformatics. 2004;20: 307–315. doi:10.1093/bioinformatics/btg405

1036   81.   Kauffmann A, Gentleman R, Huber W. arrayQualityMetrics - A bioconductor package for

1037   quality   assessment   of   microarray   data.   Bioinformatics.   2009;25:   415–416.

1038   doi:10.1093/bioinformatics/btn647

1039   82.   Edgar R, Domrachev M, Lash AE. Gene Expression Omnibus: NCBI gene expression and

1040   hybridization   array   data   repository.   Nucleic   Acids   Research.   2002;30:   207–210.

1041   doi:10.1093/nar/30.1.207

1042   83.   R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria;

1043   2020. Available: http://www.r-project.org/

84.   Thakkar S, Li T, Liu Z, Wu L, Roberts R, Tong W. Drug-induced liver injury severity and
toxicity (DILIst): binary classification of 1279 drugs by human hepatotoxicity. Drug Discovery
Today. 2020;25: 201–208. doi:10.1016/J.DRUDIS.2019.09.022

85.   Chen M, Suzuki A, Thakkar S, Yu K, Hu C, Tong W. DILIrank: the largest reference drug list
ranked by the risk for developing drug-induced liver injury in humans. Drug Discovery Today.
2016;21: 648–653. doi:10.1016/j.drudis.2016.02.015

86.   Liberzon A, Subramanian A, Pinchback R, Thorvaldsdóttir H, Tamayo P, Mesirov JP.
Molecular signatures database (MSigDB) 3.0. Bioinformatics. 2011;27: 1739–1740.
doi:10.1093/bioinformatics/btr260

87.   Dolgalev I. msigdbr: MSigDB Gene Sets for Multiple Organisms in a Tidy Data Format. 2020.

88.   Garcia-Alonso L, Ibrahim MM, Turei D, Saez-Rodriguez J. Benchmark and integration of
resources for the estimation of human transcription factor activities. bioRxiv. 2018; 337915.
doi:10.1101/337915

89.   Smedley D, Haider S, Durinck S, Pandini L, Provero P, Allen J, et al. The BioMart community
portal: an innovative alternative to large, centralized data repositories. Nucleic Acids
Research. 2015;43: W589–W598. doi:10.1093/nar/gkv350

90.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. Limma powers differential
expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Research.
2015;43: e47. doi:10.1093/nar/gkv007

91.   Türei D, Korcsmáros T, Saez-Rodriguez J. OmniPath: Guidelines and gateway for literature-
curated signaling pathway resources. Nature Methods. 2016;13: 966–967.
doi:10.1038/nmeth.4077

92.   Türei D, Valdeolivas A, Gul L, Palacio-Escat N, Klein M, Ivanova O, et al. Integrated intra-
1067   and intercellular signaling knowledge for multicellular omics analysis. Molecular Systems
1068   Biology. 2021;17: e9923. doi:10.15252/msb.20209923

1069   93.   Makowski D, Ben-Shachar MS, Patil I, Lüdecke D. Methods and Algorithms for Correlation
1070   Analysis in R. Journal of Open Source Software. 2020;5: 2306. doi:10.21105/JOSS.02306

1071   94.   Dewey M. metap: meta-analysis of significance values. 2020.

1072