Tobramycin-induced secretion of *P. aeruginosa* 5′ tRNA-fMet halves suppresses lung

inflammation via AGO2 gene silencing

Zhongyou Li[1], Katja Koeppen[1], Alix Ashare[1,2], Deborah A. Hogan[1], Scott A. Gerber[3], and Bruce

A. Stanton[1*]

[1]Department of Microbiology and Immunology, Geisel School of Medicine at Dartmouth, Hanover,

NH, USA

[2]Pulmonary and Critical Care Medicine, Dartmouth-Hitchcock Medical Center, Lebanon, NH,

USA

[3]Norris Cotton Cancer Center, Geisel School of Medicine at Dartmouth, Lebanon, NH, USA

Corresponding author:

Bruce A. Stanton, Ph.D.

Department of Microbiology and Immunology

Geisel School of Medicine at Dartmouth

520 Remsen

Hanover NH 03755

Phone: 603-650-1775

Email: Bruce.A.Stanton@dartmouth.edu

Running title: Tobramycin suppresses inflammation via tRNA halves

Keywords: cystic fibrosis, tobramycin, outer membrane vesicles, tRNA halves, anti-

inflammatory

The authors have declared that no conflict of interest exists.

Abstract

Although inhaled tobramycin increases lung function in people with cystic fibrosis (pwCF), the

density of *P. aeruginosa* in the lungs is only modestly reduced by tobramycin; hence, the

mechanism whereby tobramycin improves lung function is unclear. Here, we demonstrate that

tobramycin increases the abundance of two 5′ tRNA-fMet halves in outer membrane vesicles

(OMVs) secreted by *P. aeruginosa* and that the 5′ tRNA-fMet halves reduce IL-8 secretion by

CF bronchial epithelial cells (CF-HBECs). In mouse lung, the 5′ tRNA-fMet halves attenuate KC

secretion and neutrophil recruitment. We also report that the 5′ tRNA-fMet halves suppress pro-

inflammatory network gene expression by an Argonaut 2 (AGO2)-mediated gene silencing

mechanism, thereby reducing IL-8 secretion in CF-HBECs. Moreover, tobramycin reduces the

IL-8 concentration and neutrophil content in bronchoalveolar lavage fluid of pwCF. Thus, we

conclude that tobramycin improves lung function in part by reducing chronic inflammation and

neutrophil-mediated lung damage in pwCF.

Introduction

Pseudomonas aeruginosa is an opportunistic pathogen that infects the lungs of

immunocompromised individuals, including those with chronic obstructive pulmonary disease

and cystic fibrosis (CF), and is an important cause of acute pneumonia (Williams et al., 2010;

Konstan et al., 2012; Lieberman and Lieberman, 2003; Novosad and Barker, 2013; Parker et

al., 2016). *P. aeruginosa* is one of the leading causes of nosocomial infections worldwide, and

ventilator-associated pneumonia mortality can be as high as 30% in some institutions (Williams

et al., 2010). *P. aeruginosa* contributes to 5–10% of the acute exacerbations in COPD, afflicting

24 million Americans, and is the 3rd-leading cause of death in the U.S. (Novosad and Barker,

2013; Sethi, 2010; Lieberman and Lieberman, 2003). *P. aeruginosa* also chronically colonizes

the lungs of ~70-80% of adults with CF, and its presence is strongly associated with reduced

forced expiratory volume (FEV_1) and a progressive loss of lung function (Stanton, 2017; Davies

and Martin, 2018; Cohen and Prince, 2012)

CF is a genetic disease caused by absent or aberrant function of the cystic fibrosis

transmembrane conductance regulator (CFTR), which leads to airway periciliary dehydration,

increased mucus viscosity, and decreased mucociliary clearance (Stanton, 2017; Stoltz et al.,

2015). Insufficient mucociliary clearance causes persistent bacterial infection, non-resolving

lung inflammation, and excessive neutrophil recruitment (Hauser et al., 2011; Lin and

Kazmierczak, 2017). Chronic neutrophilic airway inflammation damages the lungs by continuous

secretion of reactive oxygen species (ROS) and proteases, contributing to bronchiectasis and

progressive CF lung function loss (Roesch et al., 2018; Khan et al., 2019). *Pseudomonas*

aeruginosa is the most common pathogen identified in adult CF lungs (Hauser et al., 2011;

Cystic Fibrosis Foundation, 2020), and *P. aeruginosa* respiratory infection correlates with CF

lung disease severity and mortality (Emerson et al., 2002; Robinson et al., 2009).

Inhaled tobramycin is a commonly used antibiotic to suppress *P. aeruginosa* burden in CF and

to ameliorate lung function loss once chronic pulmonary colonization is established (Cystic

Fibrosis Foundation, 2020; Davies and Martin, 2018). The long-term use of inhaled tobramycin

significantly improves lung function and reduces mortality in people with CF (pwCF) (Sawicki et

al., 2012; Bowman, 2002). Inhaled tobramycin is administered in intermittent repeated cycles of

28 days on the drug and 28 days off. In a double-blind, placebo-controlled study, lung function

improved significantly after the first two weeks of treatment and correlated with a decrease of *P.*

aeruginosa colony-forming units (CFUs) in sputum by more than 158-fold (Ramsey et al., 1999).

Intriguingly, the magnitude of the reduction in bacterial CFUs was less than ten-fold in the third

cycle of therapy, although lung function improvement was maintained at a comparable level

(Ramsey et al., 1999). Furthermore, an open-label, follow-on trial with adolescent patients and

12 treatment cycles revealed that the reduction of *P. aeruginosa* CFUs in sputum only explained

11.7% of CF lung function improvement (Moss, 2002). Moreover, a more recent analysis of

sputum revealed that tobramycin has no significant effect on *P. aeruginosa* abundance (Nelson

et al., 2020). Together, these data suggest that tobramycin improves CF lung function by an

unknown mechanism in addition to its bactericidal activity. The goal of this study is to elucidate

this mechanism.

In the CF lungs, *P. aeruginosa* suppresses the host immune response by secreting outer

membrane vesicles (OMVs), which fuse with host cells and deliver virulence factors, DNA, small

RNAs (sRNAs), and transfer RNA (tRNA) fragments that mediate inter-kingdom host-pathogen

interaction (Li and Stanton, 2021; Kaparakis-Liaskos and Ferrero, 2015; Bomberger et al., 2009;

Coelho and Casadevall, 2019). OMVs are 50-300 nm lipopolysaccharide (LPS)-decorated

vesicles secreted by all gram-negative bacteria (Coelho and Casadevall, 2019; Jan, 2017).

Recently, we reported that *P. aeruginosa* secretes a 24-nucleotides (nt) long sRNA in OMVs,

which diffuse through the airway mucus layer and fuse with bronchial epithelial cells to transfer

the sRNA (Koeppen et al., 2016). The sRNA down-regulates the OMV-induced secretion of IL-8,

a potent neutrophil attractant, by airway epithelial cells, leading to attenuated recruitment of

neutrophils into mouse lungs (Koeppen et al., 2016).

This study aimed to test the hypothesis that tobramycin prevents the decline in lung function of

pwCF by increasing the level of anti-inflammatory sRNAs in OMVs secreted by *P. aeruginosa.*

Here, we demonstrate that tobramycin increases the abundance of two 5′ formyl-methionine

tRNA (tRNA-fMet) halves in OMVs, and that the 5′ tRNA-fMet halves are delivered into primary

CF-HBECs by OMVs. 5′ tRNA-fMet halves suppress IL-8 secretion by CF-HBECs and reduce

KC (a murine homolog of IL-8) levels and neutrophil recruitment in mouse lungs by an AGO2-

mediated post-transcriptional regulatory mechanism. This 5′ tRNA-fMet halves-mediated

reduction in lung neutrophils is predicted to mitigate lung damage. In pwCF, the IL-8

concentration and neutrophil content in bronchoalveolar lavage fluid (BALF) was significantly

reduced during the month of tobramycin administration compared to the month off tobramycin.

Taken together, these data reveal that the clinical benefit of tobramycin is due in part to an

increase in the secretion of 5′ tRNA-fMet halves in OMVs, leading to attenuation of IL-8 and

neutrophil-mediated CF lung damage.

Results

Tobramycin reduces the ability of OMVs secreted by *P. aeruginosa* to stimulate IL-8 secretion by CF-HBECs.

To test the hypothesis that tobramycin alters the virulence of OMVs secreted by *P. aeruginosa*, we designed an *in vitro* experiment depicted in Figure 1A. *P. aeruginosa* strain PA14 was grown in lysogeny broth (LB), and OMVs secreted by *P. aeruginosa* treated with vehicle (ctrl-OMVs) or tobramycin (Tobi-OMVs) were isolated as described in Methods. The concentration of tobramycin used (1 µg/mL) reduced growth by 33%, an amount similar to that observed in pwCF treated with tobramycin after three cycles of therapy (Ramsey et al., 1999) (Figure 1B). Tobramycin increased the secretion of OMVs by 38% compared to control (Figure 1C), which coincides with a previous report that antibiotics induce OMV production by *P. aeruginosa* (MacDonald and Kuehn, 2013).

To examine the effect of OMVs on the host immune response, polarized HBECs from CF donors (CF-HBECs) were grown in air-liquid interface (ALI) culture (Randell et al., 2011; Barnaby et al., 2019) and exposed to the same number of ctrl-OMVs or Tobi-OMVs for 6 hours, whereupon IL-8 secretion was measured. Tobi-OMVs induced 36% less IL-8 secretion than ctrl-OMVs (Figure 1D). Similar results were obtained even when cells were exposed to 40% more Tobi-OMVs (1.4X Tobi-OMVs) than ctrl-OMVs to reflect the finding that tobramycin increased OMV production (Figure 1D). We measured other cytokines secreted by CF-HBECs (EGF, GRO, IL17a, and IP10) in response to ctrl-OMVs and Tobi-OMVs, but of these additional cytokines only IP-10 was significantly reduced by Tobi-OMVs compared to ctrl-OMVs (Supplemental Figure 1).

Tobramycin increases the abundance of 5′ tRNA-fMet halves in OMVs, and the tRNA halves are transferred from OMVs to CF-HBECs

To test the hypothesis that tobramycin increases the abundance of anti-inflammatory sRNAs in

OMVs, we performed a small RNA-sequencing analysis to compare the sRNA content in ctrl-

OMVs and Tobi-OMVs. We identified 6145 unique sequences mapped to the PA14 genome,

and 1064 were differentially enriched in Tobi-OMVs. The sequence length ranged from 20 to 48

nucleotides; however, we excluded the 48-nt sequences from further analysis as they

represented RNA species longer than the read length.

We focused on the most abundant and differentially induced sRNAs in Tobi-OMVs (Figure 2A

and Table 1). We chose two 35-nt long sRNAs (#5 and #7 in Table 1) that were fragments of

two initiator tRNAs (tRNA-fMet1 and tRNA-fMet2 located at PA14_62790 and PA14_52320,

respectively) in PA14 for further analysis because they were bioinformatically predicted to

suppress IL-8 secretion by CF-HBECs. The sequence reads in Tobi-OMVs mapped to these

two loci had similar length distributions (Figure 2B and 2C; 80% of reads were 35-nt long),

suggesting the tRNA-fMet fragments were not products of random degradation. The similar

length distributions also imply common machinery for the biogenesis of these two tRNA

fragments. Both have low minimum free energy, suggesting stable secondary structures.

The two 35-nt long tRNA-fMet fragments are 5′ halves of tRNAs[fMet] (hereafter called 5′ tRNA-

fMet halves), which are products of cleavage in the anticodon loop (Figure 2D). Importantly, the

two 5′ tRNA-fMet halves have high sequence similarity with only one nucleotide difference,

suggesting similar sequence-based targeting functions. The high sequence similarity allowed us

to design qPCR primers to quantify both 5′ tRNA-fMet halves simultaneously. By qPCR, we

found that Tobi-OMVs secreted by PA14 and four clinical isolates (including two mucoid strains)

also contained significantly more 5′ tRNA-fMet halves than ctrl-OMVs (Figure 2E), indicating a

strain-independent phenotype that extends to clinically relevant strains. Moreover, we

reanalyzed our previously published small RNA-sequencing experiment (Koeppen et al., 2016),

in which we sought to detect PA14 sRNAs transferred into non-CF HBECs after ctrl-OMVs

exposure, and we were able to identify both 5′ tRNA-fMet halves in OMV-exposed cells but not

in the un-exposed group (Figure 2F).

Taken together, these observations demonstrate that the two 5′ tRNA-fMet halves are the most

abundant and most differentially induced sRNAs in Tobi-OMVs, have high sequence similarity,

and are delivered to airway epithelial cells by OMVs; thus, they are good candidates for further

investigation into their possible role in suppressing IL-8 secretion.

5′ tRNA-fMet halves reduce IL-8 secretion

To determine if 5′ tRNA-fMet halves reduce IL-8 secretion, we transformed PA14 with an

arabinose-inducible vector expressing 5′ tRNA-fMet1 half (tRNA1-OMVs) or an empty vector

control (V-OMVs). Small RNA-sequencing confirmed that the expression of 5′ tRNA-fMet1 half

in tRNA1-OMVs was significantly induced by 2.73 fold compared to V-OMVs (Supplemental

Figure 2). Primary CF-HBECs were exposed to V-OMVs or tRNA1-OMVs, and the secretion of

IL-8 was measured by ELISA. As predicted, tRNA1-OMVs induced less IL-8 secretion

compared to the same amount of V-OMVs (Figure 3A). To provide additional support for the

conclusion that 5′ tRNA-fMet halves reduce IL-8 secretion by CF-HBECs, we designed an

inhibitor, an RNA oligonucleotide with a complementary sequence to both 5′ tRNA-fMet halves.

CF-HBECs were transfected with the inhibitor or negative control inhibitor followed by exposure

to ctrl-OMVs or 1.4X Tobi-OMVs. As predicted, the inhibitor reduced the ability of 1.4X Tobi-

OMVs to suppress IL-8 secretion compared to ctrl-OMVs (Figure 3B).

Studies were conducted in mice to further support the conclusion that tobramycin reduces the

pro-inflammatory effect of OMVs by increasing the 5′ tRNA-fMet1 half content. Mice were

exposed to the same number of V-OMVs or tRNA1-OMVs by oropharyngeal aspiration for 5

hours, and BALF samples were harvested for analysis. The concentration of KC, a murine

functional homolog of IL-8 (Figure 3C), and neutrophil content (Figure 3D) were significantly

reduced in BALF obtained from mice exposed to tRNA1-OMVs compared to V-OMVs. Thus, 5′

tRNA-fMet1 half reduced the pro-inflammatory response of CF-HBECs *in vitro* and in a mouse

model of inflammation.

Inhaled tobramycin has an anti-inflammatory effect in *P. aeruginosa*-infected CF lungs

To determine if tobramycin reduces inflammation and neutrophil burden in CF lungs, we

performed a retrospective analysis to assess whether the administration of inhaled tobramycin

changes the inflammatory status in pwCF. BALF samples were collected from four CF patients

chronically infected with *P. aeruginosa* during the month of inhaled tobramycin (On Tobi) and

the month off tobramycin (Off Tobi). In BALF obtained On Tobi, average IL-8 levels were

reduced by 48.5% (Figure 3E), and the number of neutrophils was decreased by 25.9% (Figure

3F) compared to Off Tobi. This clinical observation is consistent with the *in vitro* and mouse

experiments, suggesting that OMVs secreted by tobramycin-exposed *P. aeruginosa* are less

pro-inflammatory than control OMVs.

5′ tRNA-fMet halves regulate gene expression by base-pairing with target genes in CF-

HBECs using an AGO2-mediated mechanism.

Although we and others have shown that prokaryotic sRNAs regulate eukaryotic gene

expression in a sequence-specific manner (Koeppen et al., 2016; Maute et al., 2013), the

mechanism is unknown. We therefore conducted experiments to determine if *P. aeruginosa* 5′

tRNA-fMet halves can utilize the eukaryotic Argonaute 2 (AGO2) dependent gene silencing

complex to suppress IL-8 secretion. We designed a three-step approach to identify the RNA

binding targets, followed by proteomic analysis to determine the effect of 5′ tRNA-fMet halves on

protein expression (Figure 4A). Ingenuity Pathway Analysis (IPA) (Krämer et al., 2014) was

performed at each step to identify significantly enriched and down-regulated pathways that are

relevant to CF and predicted to decrease IL-8 secretion (Table 2).

We first performed a miRanda microRNA (miRNA) target scan (Enright et al., 2003) to predict

the human binding targets of 5′ tRNA-fMet1 half. miRanda is an algorithm designed for RNA-

RNA binding prediction considering sequence complementarity and binding free energy. Given

the high sequence similarity between the two sRNAs, we used the sequence of 5′ tRNA-fMet1

half to scan the whole human transcriptome and adjusted the prediction for the gene expression

profile of polarized HBE cells to identify a list of 1518 predicted targets, accounting for 8.4% of

human coding genes. IPA identified several pro-inflammatory pathways in epithelial cells that

are predicted to be down-regulated by 5′ tRNA-fMet1 half, including integrin-linked kinase

signaling (Eucker et al., 2014; Ahmed et al., 2014; Gravelle et al., 2010), LPS-stimulated MAPK

Signaling (Koeppen et al., 2016), and HIF1a signaling (Cane et al., 2010) (Table 2).

The target gene prediction and pathway analysis encouraged us to map transcriptome-wide

interactions between 5′ tRNA-fMet1 half and target mRNAs mediated by AGO2. Although tRNA

fragments have been shown to regulate gene expression, to our knowledge, there is no direct

evidence that tRNA halves or any small noncoding RNA secreted by a prokaryotic organism can

suppress eukaryotic gene expression by interacting with the AGO2 gene silencing complex.

Thus, to determine if 5′ tRNA-fMet1 half interacts with eukaryotic mRNAs in the AGO2-

containing complex, we utilized the enhanced crosslinking and immunoprecipitation (eCLIP)

approach (Van Nostrand et al., 2016). Briefly, and as described in detail in methods, this

approach involved transfection of 5′ tRNA-fMet1 half into CF-HBECs, followed by ligation of 5′

tRNA-fMet1 half and other small RNAs to target mRNAs yielding sRNA-mRNA chimeric

fragments. Chimeric fragments immunoprecipitated with AGO2 were characterized with high-

throughput sequencing (chimeric eCLIP) (Figure 4B), which provided an unprecedented

resolution to identify sRNA-mRNA interactions. CF-HBECs were transfected with 5′ tRNA-fMet1

half or negative control siRNA (siNC) and subjected to AGO2 chimeric eCLIP analysis. This

analysis allowed us to profile transcriptome-wide AGO2 authentic binding sites mediated by 5′

tRNA-fMet1 half and other miRNAs with a stringent cutoff (IP vs. input cluster log2 fold

enrichment ≥ 3 and *P* value ≤ 0.001). Using these parameters, we identified 10947 AGO2

binding sites in 4454 genes. Within those authentic binding sites, we identified 629 chimeric

reads containing at least 18-nt long subsequences of 5′ tRNA-fMet1 half. Lengths of identified

subsequences ranged from 18-nt to the full size of 5′ tRNA-fMet1 half, and alignment positions

of subsequences were evenly distributed (Figure 4C). These findings suggest that the full length

5′ tRNA-fMet1 half was loaded into the AGO2 complex for gene targeting without being pre-

processed into a shorter sRNA, and the subsequences with different lengths identified in

chimeric reads were products of RNA fragmentation, a key step in the eCLIP sequencing library

preparation.

To deeply profile the target repertoire of 5′ tRNA-fMet1 half mediated by AGO2, we designed a

5′ tRNA-fMet1 half specific primer, which anneals to most of the identified subsequence (Figure

4C), for targeted sequencing (targeted chimeric eCLIP). The targeted chimeric eCLIP allowed

us to sequence 5′ tRNA-fMet1 half containing chimeric fragment at a much higher depth. We

identified that 5′ tRNA-fMet1 half targeted 5776 sites in 1945 genes, and those target sites were

not found in the negative control (siNC) transfected cells. Interestingly, although miRNA-AGO2

complexes usually target 3′ untranslated regions (3′ UTR), most 5′ tRNA-fMet1 half-AGO2 target

sites were in introns (Figure 4D). Furthermore, motif enrichment analysis in target sites revealed

that nucleotides 16-28 from the 5′-end of 5′ tRNA-fMet1 half (which do not include the only

distinct nucleotide between the two 5′ tRNA-fMet halves) could explain 77% of identified target

sites (Figure 4E). The fact that the most popular binding motif does not contain the unique

nucleotide differentiating the two 5′ tRNA-fMet halves suggests that they have many common

target genes. Moreover, we found that miRanda predicted the target genes identified by

chimeric eCLIP significantly better than expected by chance (Fisher′ s exact test, $P < 10^{-12}$),

suggesting that bioinformatic target prediction methods based on base-pairing can reliably

predict target genes. Also, IPA predicted that a similar set of pro-inflammatory and IL-8

induction pathways were inhibited by down-regulating these target genes (Table 2).

Lastly, to identify proteins whose abundances were changed by 5′ tRNA-fMet halves delivered

by OMVs, we utilized OMVs secreted by the 5′ tRNA-fMet1 half-overexpression and empty

vector clones (Supplemental Figure 2). Primary CF-HBECs from three donors were exposed to

V-OMVs or tRNA1-OMVs for 6 hours before being subjected to proteomic analysis. 8343

proteins were identified, and we selected the top 20% differentially expressed proteins by P

value, yielding 943 down-regulated proteins (Figure 4F). The statistically enriched and down-

regulated pathways identified by IPA overlapped with our previous analysis based on identified

target genes (Table 2). These down-regulated pathways included downstream signaling of

IL17A and IL-6, which are pro-inflammatory cytokines secreted by other cell types in CF lungs to

induce IL-8 secretion by CF-HBECs (McAllister et al., 2005; Courtney et al., 2004; Hsu et al.).

Considering the top 20% down-regulated proteins, IPA identified seven proteins that contributed

to IL-8 expression and predicted the decrease of IL-8 secretion (Figure 4G). Among the seven

significantly down-regulated proteins, MAPK10, IKBKG, and EP300 were the direct targets of 5′

tRNA-fMet1 half identified with our targeted chimeric eCLIP experiment, suggesting targeting of

a pro-inflammatory network involving MAPK and NFκB signaling.

In summary, our three-step approach demonstrated that 5′ tRNA-fMet1 halves transferred from

OMVs to CF-HBECs were loaded into the AGO2 complex to target specific genes via a base-

pairing mechanism, thus mediating the Tobi-OMVs induced reduction in IL-8 secretion.

Discussion

The goal of this study was to determine how tobramycin improves clinical outcomes in pwCF

without significantly reducing the abundance of *P. aeruginosa*. Our data reveal that tobramycin

increases the concentration of 5′ tRNA-fMet halves in OMVs secreted by *P. aeruginosa*, that the

OMVs deliver 5′ tRNA-fMet halves to CF-HBECs, and that the increased delivery of 5′ tRNA-

fMet halves to CF-HBECs suppresses IL-8 secretion by interacting with pro-inflammatory gene

transcripts, including MAPK10, IKBKG, and EP300, in an AGO2-mediated mechanism. Both *in*

vitro and *in vivo* experiments in mice are consistent with this conclusion. Moreover, our

retrospective analysis of pwCF on and off tobramycin is consistent with our data in mice that

tobramycin reduces IL-8 and the neutrophil content in BALF. The reduction in the neutrophil

content in BALF is predicted to mitigate lung damage in the CF lungs since CF neutrophils are

the source of significant lung damage in pwCF (Figure 5). To our knowledge, this is the first

report demonstrating that tRNA halves secreted by a prokaryotic organism suppress gene

expression in eukaryotic cells by an AGO2-mediated RNA silencing mechanism.

tRNA-derived fragments are a novel class of regulatory sRNAs in prokaryotes and eukaryotes

(Li and Stanton, 2021; Su et al., 2020). tRNAs are the most abundant RNA species by the

number of molecules, and fragments with different lengths have been reported in three domains

of life. miRNA-sized (~24-nt long) tRNA fragments in mammalian cells have garnered attention

as they have been found to associate with Argonaute (AGO) proteins to mediate gene silencing

by base-pairing with target mRNAs (Maute et al., 2013; Kumar et al., 2014; Kuscu et al., 2018).

Recently, a report showed that in *Bradyrhizobium japonicum*, a 21-nt tRNA fragment utilizes

host plant AGO1 to regulate host gene expression, a cross-kingdom symbiotic relationship

between bacteria and plants (Ren et al., 2019). However, tRNA halves have been thought not to

involve the AGO-miRNA-like mechanism since they are 32–50 nt in length. tRNA halves have

been shown to have both positive and negative effects on translation by regulating the formation

of ribosomes and the translation initiation complex (Ivanov et al., 2011; Gebetsberger et al.,

2012; Fricker et al., 2019; Mleczko et al., 2018). Here, we provide the first evidence that 35-nt 5′

tRNA-fMet halves from a bacterial pathogen are transferred into eukaryotic cells and loaded into

AGO2-containing protein complexes without pre-processing into shorter sRNAs to suppress

gene expression by base-pairing with target mRNAs. This finding not only provides a novel role

of tRNA halves in host-pathogen interactions but also suggests that siRNAs may be an effective

therapeutic approach to reduce lung damage caused by chronic inflammation and excessive

neutrophil recruitment.

The production of tRNA halves, also called tRNA-derived stress-induced RNAs (tiRNAs), by

cleavage in the anticodon loop of mature tRNAs is conserved across all life domains in

response to various stress conditions (Tao et al., 2020; Fricker et al., 2019; Thompson et al.,

2008); however, the molecular mechanism by which prokaryotes sense the stress and cleave

specific tRNAs, and the cell-autonomous effects of tRNA halves induced by stress remain

largely unknown. *Mycobacterium tuberculosis* maintains its persistence in host cells by cleaving

several tRNAs in half with endonucleases VapCs and MazF-mt9 to reduce the level of

translation (Winther et al., 2016; Schifano et al., 2016). *Escherichia coli* secretes colicin D, an

anticodon ribonuclease (ACNase), to cleave tRNA-Arg of competing *E. coli* strains in half by

recognizing the anticodon-loop sequence, and the cleaved tRNA-Arg blocks the ribosome A-site

to disrupt translation (Tomita et al., 2000; Ogawa et al., 2020). Similar mechanisms have been

identified in fungi to inhibit the cell growth of nonself competitors (Chakravarty et al., 2014; Lu et

al., 2005). Together, these reports and our findings suggest that *P. aeruginosa*, in response to

tobramycin exposure, up-regulates an unidentified ACNase that recognizes and cleaves the

anticodon loop of tRNA-fMets, which are essential initiator tRNAs, to slow down cell growth and

prevent cell death.

317 The interaction of *P. aeruginosa* 5′ tRNA-fMet halves with CF-HBEC mRNAs by base-pairing is

318 reminiscent of miRNA-mRNA interactions in mammalian cells. While canonical miRNAs use a

319 seed region, typically nucleotides 2-7 of miRNAs, to base-pair with the 3′ UTR of target mRNAs

320 (Chipman and Pasquinelli, 2019), we observed that tRNA-fMet halves used nucleotides 16-28 to

321 bind introns and coding sequences of target genes. This finding suggests that more studies are

322 needed to better understand the targeting rule of tRNA halves for more accurate target

323 prediction. Given the high sequence similarity of tRNAs in prokaryotes (Saks and Conery,

324 2007), precise target predictions would help generalize experimental findings to other fragments

325 or pathogens. For example, *Helicobacter pylori* secretes sR-2509025, a 31-nt 5′ tRNA-fMet

326 fragment, in OMVs that fuse with human gastric adenocarcinoma cells and sR-2509025

327 diminishes LPS-induced IL-8 secretion (Zhang et al., 2020). Due to the high sequence similarity

328 of tRNA-fMets from *P. aeruginosa* and *H. pylori* and the similar phenotype on regulating IL-8

329 secretion, sR-2509025 may interact with AGO2 in gastric epithelial cells to target the pro-

330 inflammatory network identified in this study.

331 Numerous reports have demonstrated that a myriad of signaling pathways, including NF-κB and

332 MAPK signaling pathways, induce IL-8 secretion (Lee et al., 2018; Li et al., 2002, 8; Jundi and

333 Greene, 2015). IKBKG, also known as NF-κB essential modulator (NEMO), is critical for NF-κB

334 pathway activation. Furthermore, EP300, also called P300, is a transcription co-factor required

335 for NF-κB-dependent IL-8 induction (Berghe et al., 1999; Huang et al., 2017). Moreover, a study

336 demonstrated that DNA damage leads to NF-κB activation followed by MAPK10-mediated IL-8

337 secretion (Biton and Ashkenazi, 2011). Indeed, the elevated DNA damage response correlates

338 with the non-resolving neutrophilic inflammation in the CF airways (Brown et al., 1995; Fischer

339 et al., 2013); hence, our findings revealing that 5′ tRNA-fMet halves-AGO2 complex decrease

340 IKBKG, EP300, and MAPK10 protein expression and thereby reduce IL-8 secretion and

341 neutrophil levels are consistent with the literature.

CFTR is a negative regulator of the pro-inflammatory response mediated by MAPK and NF-κB

signaling. Studies have shown that impaired CFTR leads to overactivation of NF-κB signaling

and enhanced secretion of IL-8 by epithelial cells (DiMango et al., 1998; Vij et al., 2009). Also,

CFTR down-regulates thermal injury-induced MAPK/NF-κB signaling, a pathway that leads to

IL-8 expression and pulmonary inflammation (Dong et al., 2015). Here, we demonstrate that 5′

tRNA-fMet halves target a pro-inflammatory network involving the MAPK and NF-κB signaling

pathways, which are intrinsically over-activated in CF, highlighting the importance of this

network in pulmonary inflammation.

There are a few limitations of our study. First, we performed a retrospective analysis of BALF

samples collected from pwCF on and off inhaled tobramycin; however, we could not collect

BALF in consecutive months on and off tobramycin in the same individuals because of the

invasive nature of the technique and IRB restrictions on research bronchoscopies at Dartmouth-

Hitchcock Medical Center. Nevertheless, after adjusting the number of days between collection

dates for each sample pair (range from 175 to 791 days; Supplemental Table 1), tobramycin-on

BALF had significantly lower IL-8 concentration and fewer neutrophil counts than tobramycin-off

BALF. Importantly, a similar observation was made for IL-8 in CF sputum samples collected in

consecutive months from pwCF on and off tobramycin (Husson et al., 2005). Because studies

have shown that IL-8 concentration in sputum is inversely correlated with pulmonary function

(Sagel et al., 2002, 2001), we conclude that inhaled tobramycin has an anti-inflammatory effect

in *P. aeruginosa*-infected CF lungs, resulting in improved lung function. Second, since there are

many other differences in the sRNA content and likely the virulence factor content of Tobi-OMVs

compared to ctrl-OMVs, we cannot rule out the possibility that other factors may contribute to

the difference in the immune response of CF-HBECs and mouse lungs to Tobi-OMVs versus

ctrl-OMVs. Nevertheless, since the inhibitor to 5′ tRNA-fMet halves transfected into CF-HBECs

blocked the Tobi-OMVs mediated reduction in IL-8 secretion compared to ctrl-OMVs, we

conclude that 5′ tRNA-fMet halves play a major role in suppressing IL-8 levels and neutrophil

recruitment. Third, tRNAs are known to have post-transcriptional modifications, which affect

RNA structure and RNA-protein interaction (Lorenz et al., 2017). Whether the modifications on

5′ tRNA-fMet halves modulate the anti-inflammatory effect requires additional study.

Highly effective CFTR modulator drugs have significantly improved outcomes in pwCF;

however, in a few recent studies, they have been shown to have either no effect or a modest

effect on the *P. aeruginosa* burden in the CF lungs (Davies and Martin, 2018; Yi et al., 2021).

Thus, new approaches are needed to reduce the bacterial load in the lungs of chronically

colonized pwCF. We propose that 5′ tRNA-fMet halves or similar miRNA-like molecules may be

utilized as a therapeutic strategy to reduce IL-8 and neutrophil content in the lungs of pwCF,

resulting in reduced lung damage and improved lung function.

Methods

***P. aeruginosa* cultures.** *P. aeruginosa* (strain PA14) and clinical isolates were grown in

lysogeny broth (LB, Thermo Fisher Scientific, Waltham, MA) liquid cultures at 37°C with shaking

at 225 rpm. Tobramycin (1 µg/mL), a concentration that reduces *P. aeruginosa* by an amount

similar to that observed clinically, or vehicle was added to the cultures. The clinical isolates, two

mucoid and two non-mucoid strains, have been characterized previously (Yu et al., 2012;

Moreau-Marquis et al., 2015). In some experiments, 5′ tRNA-fMet1 half (5′-

CGCGGGGTGGAGCAGTCTGGTAGCTCGTCGGGCTC-3′) was cloned into the arabinose-

inducible expression vector pMQ70 (Shanks et al., 2006) by cutting EcoRI and SmaI restriction

sites. GenScript (GenScript USA Inc., Piscataway, NJ, USA) performed the cloning procedure.

PA14 was transformed with the 5′ tRNA-fMet1 half expression vector or empty vector via

electroporation. *P. aeruginosa* strains with the arabinose-inducible vector and its derivatives

were grown in LB with 133 mM L-arabinose (2% w/v) and 300 µg/ml carbenicillin (both from

Sigma-Aldrich).

Growth kinetics of *P. aeruginosa*. *P. aeruginosa* overnight LB cultures were centrifuged,

washed, and resuspended in fresh LB before measuring the optical density at 600 nm (OD600)

to determine cell number. Bacteria were seeded at 1×10^5 cells per 100 µl LB with or without

tobramycin (1 µg/mL) in a transparent, flat bottom, 96-well plate covered with a lid. The plate

was cultured in a plate reader at 37°C for 24 h. The reader was programmed to measure the

OD600 every 10 minutes after shaking the plate for 5 seconds.

Outer membrane vesicle preparation and quantification. OMVs were isolated as described

by us previously (Koeppen et al., 2016; Bauman and Kuehn, 2006). Briefly, *P. aeruginosa*

overnight cultures were centrifuged for 1 h at 2800 g and 4°C to pellet the bacteria. The

supernatant was filtered twice through 0.45 µm PVDF membrane filters (Millipore, Billerica, MA,

USA) to remove bacteria and concentrated with 30K Amicon filters (Millipore, Billerica, MA,

USA) at 2800 g and 4°C to obtain ~200 µL concentrate. The concentrate was resuspended in

OMV buffer (20 mM HEPES, 500 mM NaCl, pH 7.4) and subjected to ultracentrifugation for 2 h

at 200,000 g and 4°C to pellet OMVs. OMV pellets were re-suspended in 60% OptiPrep Density

Gradient Medium (Sigma-Aldrich, Cat. # D1556) and layered with 40%, 35%, 30% and 20%

OptiPrep diluted in OMV buffer. OMVs in OptiPrep layers were centrifuged for 16 h at 100,000 g

and 4°C. 500 µl fractions were taken from the top of the gradient, with OMVs residing in

fractions 2 and 3, corresponding to 25% OptiPrep. The purified OMVs were quantified by

nanoparticle tracking analysis (NTA, NanoSight NS300, Malvern Panalytical Ltd, Malvern, UK)

before exposure of CF-HBECs or mice to OMVs.

CF-HBEC culture. De-identified primary human bronchial epithelial cells from four CF donors

(CF-HBECs, Phe508del homozygous) were obtained from Dr. Scott Randell (University of North

Carolina, Chapel Hill, NC, USA) and cultured as described previously (Fulcher and Randell,

2013; Koeppen et al., 2021). Briefly, cells were grown in BronchiaLife basal medium (Lifeline

Cell Technology, Frederick, MD, USA) supplemented with the BronchiaLife B/T LifeFactors Kit

(Lifeline) as well as 10,000 U/ml Penicillin and 10,000 µg/ml Streptomycin.

To polarize cells, CF-HBECs were seeded on polyester transwell permeable filters (#3405 for

24-mm transwell or #3801 for 12-mm Snapwell; Corning, Corning, NY) coated with 50 µg/ml

Collagen type IV (Sigma-Aldrich, St. Louis, MO). Air Liquid Interface (ALI) medium was added to

both apical and basolateral sides for cell growth. Once a confluent monolayer was obtained, the

apical medium was removed, and cells were cultured at an air-liquid interface and fed

basolaterally every other day with ALI media for 3-4 weeks before cells were fully polarized for

treatment (Randell et al., 2011).

Exposure of cells to OMVs. Polarized cells on 12-mm Snapwell filters were washed with PBS

to remove excess mucus, and 2 mL of serum-free ALI medium was added to the basolateral

side. 1.5×10^{10} purified OMVs or the same volume of Optiprep vehicle control in 200 µL serum-

free ALI medium were applied to the apical side of cells. $2.1×10^{10}$ Tobi-OMVs (1.4X Tobi-OMVs)

were also used. After a six-hour exposure, the basolateral medium was collected for cytokine

measurements.

Cytokine measurements. Cytokine secretion from primary CF-HBECs was measured with the

Human IL-8/CXCL8 DuoSet ELISA (#DY208, R&D Systems, Minneapolis, MN). Several

samples were also screened with MILLIPLEX MAP Human Cytokine/Chemokine 41-Plex

cytokine assay (Millipore). Cytokines in mouse BALF were analyzed with the Mouse CXCL1/KC

DuoSet ELISA (#DY453, R&D Systems, Minneapolis, MN).

RNA isolation and small RNA-seq analysis. PA14 was grown in T-broth (10 g tryptone and 5

g NaCl in 1 L H_2O) with or without tobramycin (1 μg/mL) to reduce small RNA reads from yeast

present in LB medium. The culture supernatants were processed as mentioned above to obtain

OMV pellets. The pellets were resuspended with OMV buffer and re-pelleted again by

centrifugation at 200,000 g for 2 h at 4°C and lysed with Qiazol followed by RNA isolation with

the miRNeasy kit (Qiagen) to obtain total RNA including the small RNA fraction. DNase-treated

total RNA was used to prepare cDNA libraries with the SMARTer smRNA-Seq Kit (Takara Bio,

Mountain View, CA). Libraries were sequenced as 50 bp single-end reads on an Illumina HiSeq

sequencer. The first three nucleotides of all reads and the adapter sequences were trimmed

using cutadapt (Martin, 2011) before sequence alignment.

To verify the overexpression of 5′ tRNA-fMet1 half, PA14 clones with the 5′ tRNA-fMet1 half

expression plasmid or the empty pMQ70 vector were grown in LB (with L-arabinose and

carbenicillin) for isolation of V-OMVs and tRNA1-OMVs. The OMV pellets were collected and

processed as described above to isolate RNA. The QIAseq miRNA Library Kit (Qiagen) was

used to prepare cDNA libraries, and 50 bp single-end sequencing was performed on an Illumina

MiniSeq system.

Reads were aligned to the PA14 reference genome using CLC Genomics Workbench (CLC-

Bio/Qiagen) with the following modifications from the standard parameters: a) the maximum

number of mismatches = zero to eliminate unspecific alignment and b) the maximum number of

hits for a read = 30 to capture all sRNAs aligned to the PA14 genome. Pileups of mapped reads

and frequency tables for each unique sequence were exported for normalization and further

analysis with the software package edgeR in the R environment (R Core Team, 2021; Robinson

et al., 2010).

Detection of 5′ tRNA-fMet halves by RT-PCR. The induction of 5′ tRNA-fMet halves by

tobramycin in OMVs of different *P. aeruginosa* strains was detected by custom Taqman Small

RNA Assay (#4398987, Thermo Fisher Scientific). According to the manufacturer′s instructions,

cDNA was synthesized with the TaqMan MicroRNA Reverse Transcription Kit (#4366596,

Thermo Fisher Scientific). PCR amplification and detection of 5′ tRNA-fMet halves were

performed using the TaqMan Universal PCR Master Mix (#4304437, Thermo Fisher Scientific)

as well as custom primers and probe design to target both 5′ tRNA-fMet halves specifically.

5′ tRNA-fMet1 half target prediction. The miRanda microRNA target scanning algorithm

(v3.3a) was used to predict human target genes of 5′ tRNA-fMet1 half (Enright et al., 2003). The

5′ tRNA-fMet1 half sequence was scanned against human RNA sequences (annotations from

GRCh38.p13 assembly) with a mimimum miRanda alignment score of 150 to generate a list of

predicted target genes and the corresponding interaction minimum free energies. To account for

the effect of gene expression on target prediction, for each predicted target the minimum free

energy was multiplied by the gene expression level (log2CPM) in polarized HBECs identified in

our previous publication (Goodale et al., 2017) to obtain an energy-expression score. 1518

genes (8.4% of all human genes) with energy-expression scores small than -200 were defined

as predicted targets for the Ingenuity Pathway Analysis (Krämer et al., 2014).

Transfection of CF-HBECs with 5′ tRNA-fMet1 half and chimeric eCLIP analysis

CF-HBECs were seeded on 15 cm dishes coated with PureCol Bovine Collagen Solution

(Advanced BioMatrix, Carlsbad, CA, USA) at 2.7×10^6 cells per dish. Three days after seeding

(at 80% confluence), cells were washed and fed with complete Lifeline medium with antibiotics

and transfected with 100 nM 5′ tRNA-fMet1 half (#10620310, Invitrogen custom siRNA, Thermo

Fisher Scientific) or 100 nM AllStars Negative Control siRNA (siNC) using HiPerFect

transfection reagent (both from Qiagen). One day after transfection, cells were washed and

covered with room temperature PBS before UV irradiation (254 nm, 400 mJ/cm^2). The irradiated

cells were partially digested with pre-warmed 37°C trypsin/EDTA followed by addition of cold

soybean trypsin inhibitor solution to round up cells before collection with scrapers and

centrifugation at 600 x g for 10 minutes at 4°C. Cell pellets were flash-frozen in liquid nitrogen

and shipped to Eclipse BioInnovations for chimeric eCLIP (Eclipse BioInnovations, San Diego,

CA).

The chimeric eCLIP experiment and initial data analysis were conducted by Eclipse

BioInnovations (Eclipse BioInnovations, San Diego, CA) as previously described (Van Nostrand

et al., 2016) with an additional ligation step to form chimeric RNA-RNA species before 3′ RNA

adapter ligation. In brief, cells were lysed and digested with RNase I. For each cell pellet, an

input and an immunoprecipitated sample using an anti-AGO2 antibody (Eclipse BioInnovations,

San Diego, CA) were generated for cDNA library preparation followed by paired-end 150 bp

sequencing on a NovaSeq platform. Non-chimeric reads were mapped to the human genome

(UCSC version GRCh38/hg38), AGO2 binding clusters were identified by CLIPper (Lovci et al.,

2013) in immunoprecipitated (IP) samples and normalized against the paired input sample to

define significant peaks (log2 fold change ≥ 3 of normalized reads and P value < 0.001

determined by Fisher's exact test). 5′ tRNA-fMet1 half-containing chimeric reads with at least 18

nt subsequences of 5′ tRNA-fMet1 half were identified, and the subsequences were trimmed

before mapping to the human genome.

For targeted chimeric eCLIP, a target-specific primer 5′ GGGTGGAGCAGTCTGGTA and a

sequencing adapter-specific primer were used to enrich 5′ tRNA-fMet1 half-containing cDNA

from the IP sample libraries before paired-end 150 bp sequencing on a NovaSeq platform. The

primer sequence was trimmed from the 5′ ends of reads, and the remainder of reads were

analyzed as non-chimeric reads as described above. The significant peak regions were

identified using the same cutoffs, and HOMER′s findMotifsGenome.pl program was used for

motif enrichment analysis (Heinz et al., 2010). The resulting list of target genes with significant

peaks in the transcripts was used as input for Ingenuity Pathway Analysis.

Proteomic analysis. Primary CF-HBECs were polarized on 24 mm transwell filters and washed

with PBS before treatment. 2 mL serum-free ALI medium was added to the basolateral side.

2.8×10^{10} purified V-OMVs or tRNA1-OMVs in 800 μL serum-free ALI medium were applied to

the apical side of cells. After a six-hour exposure, the cells were washed with PBS and

detached from the transwells with pre-warmed 37°C trypsin/EDTA. Cells were pelleted and

flash-frozen in liquid nitrogen for proteomic analysis.

The cell pellets were lysed in 8M urea/50mM Tris pH 8.1/100mM NaCl + protease inhibitors

(Roche) and quantified by BCA assay (Pierce), followed by trypsin digestion and desalting. 40

micrograms of peptides from each pellet were labeled with unique TMT reagent isobars; the

individual TMT-labeled samples were then combined and fractionated offline into 12 fractions by

PFP-RP-LC (Grassetti et al., 2017), followed by analysis on a UPLC-Orbitrap Fusion Lumos

tribrid instrument in SPS-MS3 mode (McAlister et al., 2014). The resulting tandem mass spectra

were data-searched using Comet; TMT reporter ion intensities were summed for each protein

and normalized for total intensity across all channels. Mean fold changes comparing tRNA1-

OMV-exposed cells with V-OMVs-exposed cells were calculated for each protein detected in all

samples. Proteins were ranked by paired t-test P value, and network analysis of the top 20%

proteins was performed with Ingenuity Pathway Analysis.

Transfection of CF-HBECs with 5′ tRNA-fMet halves inhibitor and OMV exposure. CF-

HBECs were seeded on PureCol-coated 12-well plates (Corning Inc.) at 50,000 cells per well.

Two days after seeding (~80% confluence), cells were washed and fed with the complete

Lifeline medium plus antibiotics and transfected with 50 nM custom mirVana miRNA inhibitor

(inhibitor sequence: 5′- GAGCCCGACGAGCUACCAGACUGCUCCA-3′, #4464086, Thermo

Fisher Scientific) or 50 nM mirVanna inhibitor negative control#1 (#4464077, Thermo Fisher

Scientific) using HiPerFect transfection reagent (Qiagen). 6 hours after transfection, cells were

exposed to Optiprep vehicle ctrl, ctrl-OMVs ($0.4×10^{10}$ per well), 1.4X Tobi-OMVs ($0.55×10^{10}$ per

well) for another 6 h, and the supernatants were collected for cytokine measurements.

Mouse exposure to OMVs

All animal experiments were approved by the Dartmouth Institutional Animal Care and Use

Committee (Protocol No. 00002026). 8–9 weeks old male and female C57BL/6J mice (The

Jackson Laboratory, Bar Harbor, ME, USA) were inoculated by oropharyngeal aspiration with

OMVs ($0.5×10^{10}$ OMVs per mouse) or vehicle following brief anesthesia with isoflurane. OMV

concentrations were adjusted with PBS to obtain 50 µl inoculation volume. 5 h after exposure,

mice were euthanized using isoflurane anesthesia, followed by cervical dislocation after

breathing stops. Mice trachea were surgically exposed, and a catheter tube was inserted into

the trachea and stabilized with sutures (#100–5000, Henry Schein Inc., Melville, NY, USA). The

catheter was prepared by fitting a 23 gauge needle (BD #305145, Becton, Dickinson and

Company, Franklin Lakes, NJ, USA) into transparent plastic tubing (BD #427411). BALF was

collected by pumping 1 ml of sterile PBS into the lungs and recovered with a syringe (BD

#309659). This process was repeated once to collect 2 mL of BALF.

Human subjects and bronchoscopy. All CF subjects were enrolled in a protocol approved by

the Dartmouth Hitchcock Institutional Review Board (Protocol No. 22781). CF subjects

(Phe508del homozygous) prescribed with an inhaled tobramycin regimen were enrolled if they

had an FEV1 > 50% predicted, and were not currently having an exacerbation. Following

informed consent, local anesthesia with nebulized lidocaine was administered to the posterior

pharynx. Under conscious sedation, a flexible fiberoptic bronchoscopy was performed

transorally. BALF was obtained from tertiary airways. After the bronchoscopy procedure, CF

subjects were monitored per institutional protocol until they were stable for discharge.

Quantification of neutrophils in BALF. Cells in BALF samples were pelleted and

resuspended in 100 µL RBC lysis buffer (Promega) for 1 min. After removing red blood cells, the

total number of cells in each BALF sample was counted, and concentrations were adjusted.

2×10^5 cells per sample were spun onto glass slides, air-dried, and stained with the Differential

Quik Stain Kit (Polysciences, Warrington, PA) according to the manufacturer′s protocol.

Neutrophils were counted under 100x magnification using a microscope. The neutrophil

concentration of BALF was calculated by accounting for the retrieved BALF volume and the

dilution factors used to adjust the cell concentration.

Statistics. Data were analyzed using the R software environment for statistical computing and

graphics version 4.1.0 (R Core Team, 2021) and Ingenuity Pathway Analysis (Krämer et al.,

2014). Statistical significance was calculated using a mixed effect linear model, Wilcoxon rank-

sum tests, paired t-tests, and likelihood ratio tests on gene-wise negative binomial generalized

linear models, as indicated in the figure legends. Data were visualized, and figures were created

using the R package ggplot2 (Wickham, 2016, 2).

Supplemental material. Fig. S1 contains data on the effect of Tobi-OMVs on cytokines

secreted by CF-HBEC. Fig. S2 provides validation that the 5′ tRNA-fMet1 half are over-

expressed in OMVs. Table S1 contains the human BALF sample collection dates.

574 **Data availability.** Small RNA-seq and chimera eCLIP data are available from the Gene

575 Expression Omnibus database (accession number GSE183895, GSE183897, and

576 GSE183898).

Author contributions

ZL, KK, AA, DAH, SAG, and BAS designed the research studies. ZL and KK conducted experiments, acquired data, and analyzed data. AA recruited human subjects, collected clinical samples and data for analysis. SAG performed the proteomic experiment. ZL prepared figures. ZL, KK, AA, DAH, SAG, and BAS wrote the manuscript. All authors contributed to the article and approved the submitted version.

Acknowledgments

This work was supported by the Cystic Fibrosis Foundation (STANTO19G0, STANTO20PO, STANTO19R0, and HOGAN19G0), the NIH (P30-DK117469, R01HL151385, P20-GM113132, S10OD016262), and NCCC Cancer Center Core Grants (5P30 CA023108-41, P30CA023108). We thank Dr. Fred W. Kolling for advice and support on the RNA-seq experiments.

Reference

Ahmed, A.U., S.T. Sarvestani, M.P. Gantier, B.R.G. Williams, and G.E. Hannigan. 2014.
Integrin-linked Kinase Modulates Lipopolysaccharide- and *Helicobacter pylori*-
induced Nuclear Factor κB-activated Tumor Necrosis Factor-α Production via
Regulation of p65 Serine 536 Phosphorylation. *Journal of Biological Chemistry*.
289:27776–27793. doi:10.1074/jbc.M114.574541.

Barnaby, R., K. Koeppen, and B.A. Stanton. 2019. Cyclodextrins reduce the ability of
Pseudomonas aeruginosa outer-membrane vesicles to reduce CFTR Cl−
secretion. *American Journal of Physiology-Lung Cellular and Molecular
Physiology*. 316:L206–L215. doi:10.1152/ajplung.00316.2018.

Bauman, S.J., and M.J. Kuehn. 2006. Purification of outer membrane vesicles from
Pseudomonas aeruginosa and their activation of an IL-8 response. *Microbes and
Infection*. 8:2400–2408. doi:10.1016/j.micinf.2006.05.001.

Berghe, W.V., K.D. Bosscher, E. Boone, S. Plaisance, and G. Haegeman. 1999. The
Nuclear Factor-κB Engages CBP/p300 and Histone Acetyltransferase Activity for
Transcriptional Activation of the Interleukin-6 Gene Promoter. *Journal of
Biological Chemistry*. 274:32091–32098. doi:10.1074/jbc.274.45.32091.

Biton, S., and A. Ashkenazi. 2011. NEMO and RIP1 Control Cell Fate in Response to
Extensive DNA Damage via TNF-α Feedforward Signaling. *Cell*. 145:92–103.
doi:10.1016/j.cell.2011.02.023.

Bomberger, J.M., D.P. MacEachran, B.A. Coutermarsh, S. Ye, G.A. O'Toole, and B.A.
Stanton. 2009. Long-Distance Delivery of Bacterial Virulence Factors by
Pseudomonas aeruginosa Outer Membrane Vesicles. *PLOS Pathogens*.
5:e1000382. doi:10.1371/journal.ppat.1000382.

Bowman, C.M. 2002. The long-term use of inhaled tobramycin in patients with cystic
fibrosis. *Journal of Cystic Fibrosis*. 1:194–198. doi:10.1016/S1569-
1993(02)00003-6.

Brown, R.K., A. McBurney, J. Lunec, and F.J. Kelly. 1995. Oxidative damage to DNA in
patients with cystic fibrosis. *Free Radical Biology and Medicine*. 18:801–806.
doi:10.1016/0891-5849(94)00172-G.

Cane, G., A. Ginouvès, S. Marchetti, R. Buscà, J. Pouysségur, E. Berra, P. Hofman,
and V. Vouret-Craviari. 2010. HIF-1α mediates the induction of IL-8 and VEGF
expression on infection with Afa/Dr diffusely adhering *E. coli* and promotes EMT-
like behaviour. *Cellular Microbiology*. 12:640–653. doi:10.1111/j.1462-
5822.2009.01422.x.

Chakravarty, A.K., P. Smith, R. Jalan, and S. Shuman. 2014. Structure, Mechanism,
and Specificity of a Eukaryal tRNA Restriction Enzyme Involved in Self-Nonself
Discrimination. *Cell Reports*. 7:339–347. doi:10.1016/j.celrep.2014.03.034.

Chipman, L.B., and A.E. Pasquinelli. 2019. miRNA Targeting: Growing beyond the
Seed. *Trends in Genetics*. 35:215–222. doi:10.1016/j.tig.2018.12.005.

Coelho, C., and A. Casadevall. 2019. Answers to naysayers regarding microbial
extracellular vesicles. *Biochemical Society Transactions*. 47:1005–1012.
doi:10.1042/BST20180252.

Cohen, T.S., and A. Prince. 2012. Cystic fibrosis: a mucosal immunodeficiency
syndrome. *Nat Med*. 18:509–519. doi:10.1038/nm.2715.

Courtney, J.M., M. Ennis, and J.S. Elborn. 2004. Cytokines and inflammatory mediators
in cystic fibrosis. *Journal of Cystic Fibrosis*. 3:223–231.
doi:10.1016/j.jcf.2004.06.006.

Cystic Fibrosis Foundation. 2020. Cystic Fibrosis Foundation Patient Registry 2019
Annual Data Report. *Cystic Fibrosis Foundation, Bethesda, MD.* 92.

Davies, J.C., and I. Martin. 2018. New anti-pseudomonal agents for cystic fibrosis- still
needed in the era of small molecule CFTR modulators? *Expert Opinion on*
Pharmacotherapy. 19:1327–1336. doi:10.1080/14656566.2018.1505864.

DiMango, E., A.J. Ratner, R. Bryan, S. Tabibi, and A. Prince. 1998. Activation of NF-
kappaB by adherent *Pseudomonas aeruginosa* in normal and cystic fibrosis
respiratory epithelial cells. *J. Clin. Invest.* 101:2598–2605. doi:10.1172/JCI2865.

Dong, Z.W., J. Chen, Y.C. Ruan, T. Zhou, Y. Chen, Y. Chen, L.L. Tsang, H.C. Chan,
and Y.Z. Peng. 2015. CFTR-regulated MAPK/NF-κB signaling in pulmonary
inflammation in thermal inhalation injury. *Sci Rep*. 5. doi:10.1038/srep15946.

Emerson, J., M. Rosenfeld, S. McNamara, B. Ramsey, and R.L. Gibson. 2002.
Pseudomonas aeruginosa and other predictors of mortality and morbidity in
young children with cystic fibrosis. *Pediatric Pulmonology*. 34:91–100.
doi:10.1002/ppul.10127.

Enright, A.J., B. John, U. Gaul, T. Tuschl, C. Sander, and D.S. Marks. 2003. MicroRNA
targets in *Drosophila*. *Genome Biology*. 5:R1. doi:10.1186/gb-2003-5-1-r1.

Eucker, T.P., D.R. Samuelson, M. Hunzicker-Dunn, and M.E. Konkel. 2014. The focal
complex of epithelial cells provides a signalling platform for interleukin-8
induction in response to bacterial pathogens. *Cellular Microbiology*. 16:1441–
1455. doi:https://doi.org/10.1111/cmi.12305.

Fischer, B.M., J.K. Wong, S. Degan, A.B. Kummarapurugu, S. Zheng, P. Haridass, and
J.A. Voynow. 2013. Increased expression of senescence markers in cystic
fibrosis airways. *American Journal of Physiology-Lung Cellular and Molecular*
Physiology. 304:L394–L400. doi:10.1152/ajplung.00091.2012.

Fricker, R., R. Brogli, H. Luidalepp, L. Wyss, M. Fasnacht, O. Joss, M. Zywicki, M.
Helm, A. Schneider, M. Cristodero, and N. Polacek. 2019. A tRNA half modulates
translation as stress response in *Trypanosoma brucei*. *Nature Communications*.
10:118. doi:10.1038/s41467-018-07949-6.

Fulcher, M.L., and S.H. Randell. 2013. Human Nasal and Tracheo-Bronchial
Respiratory Epithelial Cell Culture. *In* Epithelial Cell Culture Protocols: Second
Edition. S.H. Randell and M.L. Fulcher, editors. Humana Press, Totowa, NJ.
109–121.

Gebetsberger, J., M. Zywicki, A. Künzi, and N. Polacek. 2012. tRNA-Derived Fragments
Target the Ribosome and Function as Regulatory Non-Coding RNA in *Haloferax*
volcanii. *Archaea*. 2012:e260909. doi:https://doi.org/10.1155/2012/260909.

Goodale, B.C., E.J. Rayack, and B.A. Stanton. 2017. Arsenic alters transcriptional
responses to *Pseudomonas aeruginosa* infection and decreases antimicrobial
defense of human airway epithelial cells. *Toxicol Appl Pharmacol*. 331:154–163.
doi:10.1016/j.taap.2017.06.010.

Grassetti, A.V., R. Hards, and S.A. Gerber. 2017. Offline pentafluorophenyl (PFP)-RP
prefractionation as an alternative to high-pH RP for comprehensive LC-MS/MS
proteomics and phosphoproteomics. *Anal Bioanal Chem*. 409:4615–4625.
doi:10.1007/s00216-017-0407-6.

Gravelle, S., R. Barnes, N. Hawdon, L. Shewchuk, J. Eibl, J.S. Lam, and M. Ulanova.
2010. Up-regulation of integrin expression in lung adenocarcinoma cells caused
by bacterial infection: in vitro study. *Innate Immun*. 16:14–26.
doi:10.1177/1753425909106170.

Gruber, A.R., R. Lorenz, S.H. Bernhart, R. Neuböck, and I.L. Hofacker. 2008. The
Vienna RNA Websuite. *Nucleic Acids Research*. 36:W70–W74.
doi:10.1093/nar/gkn188.

Hauser, A.R., M. Jain, M. Bar-Meir, and S.A. McColley. 2011. Clinical Significance of
Microbial Infection and Adaptation in Cystic Fibrosis. *Clin Microbiol Rev*. 24:29–
70. doi:10.1128/CMR.00036-10.

Heinz, S., C. Benner, N. Spann, E. Bertolino, Y.C. Lin, P. Laslo, J.X. Cheng, C. Murre,
H. Singh, and C.K. Glass. 2010. Simple combinations of lineage-determining
transcription factors prime cis-regulatory elements required for macrophage and
B cell identities. *Mol Cell*. 38:576–589. doi:10.1016/j.molcel.2010.05.004.

Hsu, D., P. Taylor, D. Fletcher, R. van Heeckeren, J. Eastman, A. van Heeckeren, P.
Davis, J.F. Chmiel, E. Pearlman, and T.L. Bonfield. Interleukin-17
Pathophysiology and Therapeutic Intervention in Cystic Fibrosis Lung Infection
and Inflammation. *Infection and Immunity*. 84:2410–2421. doi:10.1128/IAI.00284-
16.

Huang, Z.-W., G.-S. Lien, C.-H. Lin, C.-P. Jiang, and B.-C. Chen. 2017. p300 and
C/EBPβ-regulated IKKβ expression are involved in thrombin-induced IL-8/CXCL8
expression in human lung epithelial cells. *Pharmacological Research*. 121:33–41.
doi:10.1016/j.phrs.2017.04.020.

Husson, M.O., N. Wizla-Derambure, D. Turck, P. Gosset, and B. Wallaert. 2005. Effect
of intermittent inhaled tobramycin on sputum cytokine profiles in cystic fibrosis.
Journal of Antimicrobial Chemotherapy. 56:247–249. doi:10.1093/jac/dki179.

Ivanov, P., M.M. Emara, J. Villen, S.P. Gygi, and P. Anderson. 2011. Angiogenin-
Induced tRNA Fragments Inhibit Translation Initiation. *Molecular Cell*. 43:613–
623. doi:10.1016/j.molcel.2011.06.022.

Jan, A.T. 2017. Outer Membrane Vesicles (OMVs) of Gram-negative Bacteria: A
Perspective Update. *Front. Microbiol.* 8. doi:10.3389/fmicb.2017.01053.

Jundi, K., and C.M. Greene. 2015. Transcription of Interleukin-8: How Altered
Regulation Can Affect Cystic Fibrosis Lung Disease. *Biomolecules*. 5:1386–
1398. doi:10.3390/biom5031386.

Kaparakis-Liaskos, M., and R.L. Ferrero. 2015. Immune modulation by bacterial outer
membrane vesicles. *Nature Reviews Immunology*. 15:375–387.
doi:10.1038/nri3837.

Khan, M.A., Z.S. Ali, N. Sweezey, H. Grasemann, and N. Palaniyar. 2019. Progression
of Cystic Fibrosis Lung Disease from Childhood to Adulthood: Neutrophils,
Neutrophil Extracellular Trap (NET) Formation, and NET Degradation. *Genes*.
10:183. doi:10.3390/genes10030183.

Koeppen, K., T.H. Hampton, M. Jarek, M. Scharfe, S.A. Gerber, D.W. Mielcarz, E.G.
Demers, E.L. Dolben, J.H. Hammond, D.A. Hogan, and B.A. Stanton. 2016. A
Novel Mechanism of Host-Pathogen Interaction through sRNA in Bacterial Outer
Membrane Vesicles. *PLOS Pathogens*. 12:e1005672.
doi:10.1371/journal.ppat.1005672.

Koeppen, K., A. Nymon, R. Barnaby, Z. Li, T.H. Hampton, A. Ashare, and B.A. Stanton.
2021. CF monocyte-derived macrophages have an attenuated response to
extracellular vesicles secreted by airway epithelial cells. *American Journal of
Physiology-Lung Cellular and Molecular Physiology*. 320:L530–L544.
doi:10.1152/ajplung.00621.2020.

Konstan, M.W., J.S. Wagener, D.R. VanDevanter, D.J. Pasta, A. Yegin, L. Rasouliyan,
and W.J. Morgan. 2012. Risk factors for rate of decline in FEV1 in adults with
cystic fibrosis. *Journal of Cystic Fibrosis*. 11:405–411.
doi:10.1016/j.jcf.2012.03.009.

Krämer, A., J. Green, J. Pollard, and S. Tugendreich. 2014. Causal analysis
approaches in Ingenuity Pathway Analysis. *Bioinformatics*. 30:523–530.
doi:10.1093/bioinformatics/btt703.

Kumar, P., J. Anaya, S.B. Mudunuri, and A. Dutta. 2014. Meta-analysis of tRNA derived
RNA fragments reveals that they are evolutionarily conserved and associate with
AGO proteins to recognize specific RNA targets. *BMC Biology*. 12:78.
doi:10.1186/s12915-014-0078-0.

Kuscu, C., P. Kumar, M. Kiran, Z. Su, A. Malik, and A. Dutta. 2018. tRNA fragments
(tRFs) guide Ago to regulate gene expression post-transcriptionally in a Dicer-
independent manner. *RNA*. 24:1093–1105. doi:10.1261/rna.066126.118.

Lee, J., Y.J. Yoon, J.H. Kim, N.T.H. Dinh, G. Go, S. Tae, K.-S. Park, H.T. Park, C. Lee,
T.-Y. Roh, D. Di Vizio, and Y.S. Gho. 2018. Outer Membrane Vesicles Derived
From *Escherichia coli* Regulate Neutrophil Migration by Induction of Endothelial
IL-8. *Front Microbiol*. 9:2268. doi:10.3389/fmicb.2018.02268.

Li, J., S. Kartha, S. Iasvovskaia, A. Tan, R.K. Bhat, J.M. Manaligod, K. Page, A.R.
Brasier, and M.B. Hershenson. 2002. Regulation of human airway epithelial cell
IL-8 expression by MAP kinases. *American Journal of Physiology-Lung Cellular
and Molecular Physiology*. 283:L690–L699. doi:10.1152/ajplung.00060.2002.

Li, Z., and B.A. Stanton. 2021. Transfer RNA-Derived Fragments, the Underappreciated
Regulatory Small RNAs in Microbial Pathogenesis. *Front. Microbiol.* 12.
doi:10.3389/fmicb.2021.687632.

Lieberman, D., and D. Lieberman. 2003. Pseudomonal Infections in Patients with
COPD. *Treat Respir Med*. 2:459–468. doi:10.1007/BF03256673.

Lin, C.K., and B.I. Kazmierczak. 2017. Inflammation: A Double-Edged Sword in the
Response to *Pseudomonas aeruginosa* Infection. *JIN*. 9:250–261.
doi:10.1159/000455857.

Lorenz, C., C.E. Lünse, and M. Mörl. 2017. tRNA Modifications: Impact on Structure
and Thermal Adaptation. *Biomolecules*. 7:35. doi:10.3390/biom7020035.

Lovci, M.T., D. Ghanem, H. Marr, J. Arnold, S. Gee, M. Parra, T.Y. Liang, T.J. Stark,
L.T. Gehman, S. Hoon, K.B. Massirer, G.A. Pratt, D.L. Black, J.W. Gray, J.G.
Conboy, and G.W. Yeo. 2013. Rbfox proteins regulate alternative mRNA splicing
through evolutionarily conserved RNA bridges. *Nat Struct Mol Biol*. 20:1434–
1442. doi:10.1038/nsmb.2699.

Lu, J., B. Huang, A. Esberg, M.J.O. Johansson, and A.S. Byström. 2005. The
Kluyveromyces lactis γ-toxin targets tRNA anticodons. *RNA*. 11:1648–1654.
doi:10.1261/rna.2172105.

MacDonald, I.A., and M.J. Kuehn. 2013. Stress-Induced Outer Membrane Vesicle
Production by *Pseudomonas aeruginosa*. *Journal of Bacteriology*. 195:2971–
2981. doi:10.1128/JB.02267-12.

Martin, M. 2011. Cutadapt removes adapter sequences from high-throughput
sequencing reads. *EMBnet.journal*. 17:10–12. doi:10.14806/ej.17.1.200.

Maute, R.L., C. Schneider, P. Sumazin, A. Holmes, A. Califano, K. Basso, and R. Dalla-
Favera. 2013. tRNA-derived microRNA modulates proliferation and the DNA
damage response and is down-regulated in B cell lymphoma. *PNAS*. 110:1404–
1409. doi:10.1073/pnas.1206761110.

McAlister, G.C., D.P. Nusinow, M.P. Jedrychowski, M. Wühr, E.L. Huttlin, B.K. Erickson,
R. Rad, W. Haas, and S.P. Gygi. 2014. MultiNotch MS3 Enables Accurate,
Sensitive, and Multiplexed Detection of Differential Expression across Cancer
Cell Line Proteomes. *Anal. Chem.* 86:7150–7158. doi:10.1021/ac502040v.

McAllister, F., A. Henry, J.L. Kreindler, P.J. Dubin, L. Ulrich, C. Steele, J.D. Finder, J.M.
Pilewski, B.M. Carreno, S.J. Goldman, J. Pirhonen, and J.K. Kolls. 2005. Role of
IL-17A, IL-17F, and the IL-17 Receptor in Regulating Growth-Related Oncogene-
α and Granulocyte Colony-Stimulating Factor in Bronchial Epithelium:
Implications for Airway Inflammation in Cystic Fibrosis. *J Immunol*. 175:404–412.
doi:10.4049/jimmunol.175.1.404.

Mleczko, A.M., P. Celichowski, and K. Bąkowska-Żywicka. 2018. Transfer RNA-derived
fragments target and regulate ribosome-associated aminoacyl-transfer RNA
synthetases. *Biochimica et Biophysica Acta (BBA) - Gene Regulatory
Mechanisms*. 1861:647–656. doi:10.1016/j.bbagrm.2018.06.001.

Moreau-Marquis, S., B. Coutermarsh, and B.A. Stanton. 2015. Combination of
hypothiocyanite and lactoferrin (ALX-109) enhances the ability of tobramycin and
aztreonam to eliminate *Pseudomonas aeruginosa* biofilms growing on cystic
fibrosis airway epithelial cells. *J Antimicrob Chemother*. 70:160–166.
doi:10.1093/jac/dku357.

Moss, R.B. 2002. Long-term Benefits of Inhaled Tobramycin in Adolescent Patients
With Cystic Fibrosis. *Chest*. 121:55–63. doi:10.1378/chest.121.1.55.

Nelson, M.T., D.J. Wolter, A. Eng, E.J. Weiss, A.T. Vo, M.J. Brittnacher, H.S. Hayden,
S. Ravishankar, G. Bautista, A. Ratjen, M. Blackledge, S. McNamara, L. Nay, C.
Majors, S.I. Miller, E. Borenstein, R.H. Simon, J.J. LiPuma, and L.R. Hoffman.
2020. Maintenance tobramycin primarily affects untargeted bacteria in the CF
sputum microbiome. *Thorax*. 75:780–790. doi:10.1136/thoraxjnl-2019-214187.

Novosad, S.A., and A.F. Barker. 2013. Chronic obstructive pulmonary disease and
bronchiectasis: *Current Opinion in Pulmonary Medicine*. 19:133–139.
doi:10.1097/MCP.0b013e32835d8312.

Ogawa, T., K. Takahashi, W. Ishida, T. Aono, M. Hidaka, T. Terada, and H. Masaki.
2020. Substrate recognition mechanism of tRNA-targeting ribonuclease, colicin
D, and an insight into tRNA cleavage-mediated translation impairment. *RNA*
Biology. 0:1–13. doi:10.1080/15476286.2020.1838782.

Parker, D., D. Ahn, T. Cohen, and A. Prince. 2016. Innate Immune Signaling Activated
by MDR Bacteria in the Airway. *Physiological Reviews*. 96:19–53.
doi:10.1152/physrev.00009.2015.

R Core Team. 2021. R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria.

Ramsey, B.W., M.S. Pepe, J.M. Quan, K.L. Otto, A.B. Montgomery, J. Williams-Warren,
M. Vasiljev-K, D. Borowitz, C.M. Bowman, B.C. Marshall, S. Marshall, and A.L.
Smith. 1999. Intermittent Administration of Inhaled Tobramycin in Patients with
Cystic Fibrosis. *N Engl J Med*. 340:23–30. doi:10.1056/NEJM199901073400104.

Randell, S.H., M.L. Fulcher, W. O'Neal, and J.C. Olsen. 2011. Primary Epithelial Cell
Models for Cystic Fibrosis Research. *In* Cystic Fibrosis: Diagnosis and Protocols,
Volume II: Methods and Resources to Understand Cystic Fibrosis. M.D. Amaral
and K. Kunzelmann, editors. Humana Press, Totowa, NJ. 285–310.

Ren, B., X. Wang, J. Duan, and J. Ma. 2019. Rhizobial tRNA-derived small RNAs are
signal molecules regulating plant nodulation. *Science*. 365:919–922.
doi:10.1126/science.aav8907.

Robinson, M.D., D.J. McCarthy, and G.K. Smyth. 2010. edgeR: a Bioconductor package
for differential expression analysis of digital gene expression data.
Bioinformatics. 26:139–140. doi:10.1093/bioinformatics/btp616.

Robinson, T.E., A.N. Leung, X. Chen, R.B. Moss, and M.J. Emond. 2009. Cystic fibrosis
HRCT scores correlate strongly with *Pseudomonas* infection. *Pediatric*
Pulmonology. 44:1107–1117. doi:10.1002/ppul.21107.

Roesch, E.A., D.P. Nichols, and J.F. Chmiel. 2018. Inflammation in cystic fibrosis: An
update. *Pediatric Pulmonology*. 53:S30–S50. doi:10.1002/ppul.24129.

Sagel, S.D., R. Kapsner, I. Osberg, M.K. Sontag, and F.J. Accurso. 2001. Airway
Inflammation in Children with Cystic Fibrosis and Healthy Children Assessed by
Sputum Induction. *Am J Respir Crit Care Med*. 164:1425–1431.
doi:10.1164/ajrccm.164.8.2104075.

Sagel, S.D., M.K. Sontag, J.S. Wagener, R.K. Kapsner, I. Osberg, and F.J. Accurso.
2002. Induced sputum inflammatory measures correlate with lung function in
children with cystic fibrosis. *The Journal of Pediatrics*. 141:811–817.
doi:10.1067/mpd.2002.129847.

Saks, M.E., and J.S. Conery. 2007. Anticodon-dependent conservation of bacterial
tRNA gene sequences. *RNA*. 13:651–660. doi:10.1261/rna.345907.

Sawicki, G.S., J.E. Signorovitch, J. Zhang, D. Latremouille-Viau, M. von Wartburg, E.Q.
Wu, and L. Shi. 2012. Reduced mortality in cystic fibrosis patients treated with
tobramycin inhalation solution. *Pediatr Pulmonol*. 47:44–52.
doi:10.1002/ppul.21521.

Schifano, J.M., J.W. Cruz, I.O. Vvedenskaya, R. Edifor, M. Ouyang, R.N. Husson, B.E.
Nickels, and N.A. Woychik. 2016. tRNA is a new target for cleavage by a MazF
toxin. *Nucleic Acids Research*. 44:1256–1270. doi:10.1093/nar/gkv1370.

Sethi, S. 2010. Infection as a comorbidity of COPD. *European Respiratory Journal*.
35:1209–1215. doi:10.1183/09031936.00081409.

Shanks, R.M.Q., N.C. Caiazza, S.M. Hinsa, C.M. Toutain, and G.A. O'Toole. 2006.
Saccharomyces cerevisiae -Based Molecular Tool Kit for Manipulation of Genes
from Gram-Negative Bacteria. *Appl Environ Microbiol*. 72:5027–5036.
doi:10.1128/AEM.00682-06.

Stanton, B.A. 2017. Effects of *Pseudomonas aeruginosa*on CFTR chloride secretion
and the host immune response. *Am J Physiol Cell Physiol*. 312:C357–C366.
doi:10.1152/ajpcell.00373.2016.

Stoltz, D.A., D.K. Meyerholz, and M.J. Welsh. 2015. Origins of Cystic Fibrosis Lung
Disease. *N Engl J Med*. 372:351–362. doi:10.1056/NEJMra1300109.

Su, Z., B. Wilson, P. Kumar, and A. Dutta. 2020. Noncanonical Roles of tRNAs: tRNA
Fragments and Beyond. *Annual Review of Genetics*. 54:47–69.
doi:10.1146/annurev-genet-022620-101840.

Tao, E.-W., W.Y. Cheng, W.-L. Li, J. Yu, and Q.-Y. Gao. 2020. tiRNAs: A novel class of
small noncoding RNAs that helps cells respond to stressors and plays roles in
cancer progression. *Journal of Cellular Physiology*. 235:683–690.
doi:10.1002/jcp.29057.

Thompson, D.M., C. Lu, P.J. Green, and R. Parker. 2008. tRNA cleavage is a
conserved response to oxidative stress in eukaryotes. *RNA*. 14:2095–2103.
doi:10.1261/rna.1232808.

Tomita, K., T. Ogawa, T. Uozumi, K. Watanabe, and H. Masaki. 2000. A cytotoxic
ribonuclease which specifically cleaves four isoaccepting arginine tRNAs at their
anticodon loops. *PNAS*. 97:8278–8283. doi:10.1073/pnas.140213797.

Van Nostrand, E.L., G.A. Pratt, A.A. Shishkin, C. Gelboin-Burkhart, M.Y. Fang, B.
Sundararaman, S.M. Blue, T.B. Nguyen, C. Surka, K. Elkins, R. Stanton, F. Rigo,
M. Guttman, and G.W. Yeo. 2016. Robust transcriptome-wide discovery of RNA

binding protein binding sites with enhanced CLIP (eCLIP). *Nat Methods*. 13:508–
514. doi:10.1038/nmeth.3810.

Vij, N., S. Mazur, and P.L. Zeitlin. 2009. CFTR Is a Negative Regulator of NFκB
Mediated Innate Immune Response. *PLOS ONE*. 4:e4664.
doi:10.1371/journal.pone.0004664.

Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New
York.

Williams, B.J., J. DehnbosᴛᴇL, and T.S. Blackwell. 2010. *Pseudomonas aeruginosa*:
Host defence in lung diseases. *Respirology*. 15:1037–1056. doi:10.1111/j.1440-
1843.2010.01819.x.

Winther, K., J.J. Tree, D. Tollervey, and K. Gerdes. 2016. VapCs of *Mycobacterium*
tuberculosis cleave RNAs essential for translation. *Nucleic Acids Research*.
44:9860–9871. doi:10.1093/nar/gkw781.

Yi, B., A.H. Dalpke, and S. Boutin. 2021. Changes in the Cystic Fibrosis Airway
Microbiome in Response to CFTR Modulator Therapy. *Front. Cell. Infect.*
Microbiol. 11. doi:10.3389/fcimb.2021.548613.

Yu, Q., E.F. Griffin, S. Moreau-Marquis, J.D. Schwartzman, B.A. Stanton, and G.A.
O'Toole. 2012. In vitro evaluation of tobramycin and aztreonam versus
*Pseudomonas aeruginosa*biofilms on cystic fibrosis-derived human airway
epithelial cells. *J Antimicrob Chemother*. 67:2673–2681. doi:10.1093/jac/dks296.

Zhang, H., Y. Zhang, Z. Song, R. Li, H. Ruan, Q. Liu, and X. Huang. 2020. sncRNAs
packaged by *Helicobacter pylori* outer membrane vesicles attenuate IL-8
secretion in human cells. *International Journal of Medical Microbiology*.
310:151356. doi:10.1016/j.ijmm.2019.151356.

Figures

Figure 1. Tobramycin reduces the ability of OMVs secreted by *P. aeruginosa* to stimulate

IL-8 secretion by CF-HBECs. (**A**) A schematic diagram shows the experimental design. (**B**)

Growth curve (in microtiter plates) of PA14 in LB alone (PA14) or in LB with tobramycin (1

μg/mL; PA14+Tobi). Lines represent the averages from three biological replicates, and error

bars indicate standard error of means (SEM). The presence of tobramycin inhibited PA14

growth by 33% between the 13 to 20 hour time points. (**C**) OMV concentration of purified ctrl-

OMVs and Tobi-OMVs (*n* = 12) measured with nanoparticle tracking analysis (Nanosight

NS300). The red line connects the mean concentration of the two groups and demonstrates a

38% increase in Tobi-OMVs concentration compared to ctrl-OMVs concentration. Data are

shown as the means ± SEM (**D**) Primary CF-HBECs from three donors (*n* = 3*)* were polarized in

ALI culture before being exposed to either the same number of ctrl-OMVs or Tobi-OMVs or 40%

more Tobi-OMVs (1.4X Tobi-OMVs) for 6 hours. The basolateral medium was collected to

measure IL-8. Lines connect experiments conducted with CF-HBECs from the same donor.

Horizontal red lines and red dots indicate means ± SEM. Paired t-tests (**C**); Linear mixed-effects

models with CF-HBEC donor as a random effect were used to calculate *P* values (**D**); *$P < 0.05$;

$P < 0.01$;*$P < 0.001$.

Figure. 2 Tobramycin increases the abundance of 5′ tRNA-fMet halves in OMVs, and the

tRNA halves are transferred into host cells. (**A**) MA plot comparing the small RNA

expression profile in Tobi-OMVs and ctrl-OMVs (*n* = 3 for each group). Each dot represents a

unique sequence read. The most abundant and most induced sRNAs by tobramycin treatment

are highlighted in red and listed in Table 1. (**B and C**) Length distribution of Tobi-OMVs sRNAs

mapped to gene locus PA14_62790 (**B**) and PA14_52320 (**C**). (**D**) Secondary cloverleaf

structure of tRNAs^{fMet} and cleavage site in the anticodon loop to generate 5′ tRNA-fMet halves.

The red line indicates the only different pair of nucleotides between the two tRNAs^{fMet}, and the

red dot represents the only nucleotide difference between the two 5′ tRNA-fMet halves. (**E**)

qPCR for 5′ tRNA-fMet halves in ctrl-OMVs and Tobi-OMVs purified from PA14 and four clinical

isolates (*n* = 5 strains), including two mucoid and two non-mucoid strains. The qPCR primers

and probe were designed to detect both 5′ tRNA-fMet halves. Horizontal lines indicate means ±

SEM. A paired t-test was used to establish significance. * *P* < 0.05. (**F**) Both 5′ tRNA-fMet

halves were detected in polarized primary HBE cells exposed to ctrl-OMVs but not in unexposed

cells using small RNA sequencing (from two donors; *n* = 2). Sequence reads in (F) are from our

previously published dataset (Koeppen et al., 2016).

Figure 3. 5′ tRNA-fMet halves reduce IL-8 secretion *in vitro* and *in vivo*. (A) Polarized CF-

HBECs (*n* = 4) exposed to tRNA1-OMV secreted less IL-8 compared to cells exposed to V-

OMV. (**B**) The Tobi-OMV effect of reducing IL-8 secretion was abolished by transfection of an

antisense RNA oligo inhibitor that anneals to both 5′ tRNA-fMet halves (5′ tRNA-fMet halves

inhibitor) but not by transfection of a negative control inhibitor (*n*= 4). Lines in panels A and B

connect data points using the cells from the same donor, a biological replicate. A linear mixed-

effects model with CF-HBEC donor as a random effect was used to calculate *P* values. (**C** and

D) BALF from mice exposed to V-OMVs or tRNA1-OMVs was collected to measure KC

concentration (**C**) and neutrophil number (**D**). Minimum-to-maximum whisker and box plots

showing the median and interquartile ranges. 9 to 10 mice were used per group, and Wilcoxon

rank-sum tests were used to test significance. (**E** and **F**) BALF samples collected from four CF

subjects (*n*= 4) during the 4-week administration of inhaled tobramycin (On Tobi) or not (Off

Tobi) with BALF IL-8 levels in panel (**E**) and BALF neutrophil content in panel (**F**). Lines connect

data points from the same subject, and the arrowheads indicate the sample collection order for

each CF subject. Linear mixed-effect models were used to account for donor-to-donor variability

and the number of days between collection dates for each sample pair (Supplemental Table 1).

Horizontal red lines and red dots indicate means ± SEM; N.S., not significant; *$P < 0.05$; **$P <$

0.001.

Figure 4. 5′ tRNA-fMet halves downregulate protein expression in a sequence-specific

manner mediated by AGO2. (**A**) Schematic representation of the three-step approach to

identify targets of 5′ tRNA-fMet halves leading to differential protein expression. (**B**) Diagram of

chimeric eCLIP to identify RNAs pulled down with AGO2. Samples were treated with RNase I

for RNA fragmentation followed by immunoprecipitation of AGO2-mRNA-sRNA complexes

before mRNA-sRNA ligation to generate chimeric reads. (**C**) Alignment of chimeric reads to the

5′ tRNA-fMet1 half sequence and the count distribution of each nucleotide (top density plot).

Chimeric reads were identified in tRNA-fMet1 half-transfected CF-HBECs (one donor; *n* = 1)

and contained at least 18-nt long 5′ tRNA-fMet1 half subsequences. The red lines indicate the

region where the primer performs targeted chimeric eCLIP anneals. (**D**) Distribution of target

site features identified with the targeted chimeric eCLIP. (**E**) Sequence logos of most significant

enriched target RNA motifs and the complementary sequence of 5′ tRNA-fMet halves. R

denotes a purine nucleotide (G/A). (**F**) Volcano plot of proteomic analysis of polarized CF-

HBECs (three donors; $n = 3$) treated with tRNA1-OMVs compared to cells treated with V-OMVs.

The top 20% differentially expressed proteins, determined by paired t-tests, are colored in blue.

Red dots with numbers represent down-regulated proteins corresponding to proteins numbered

in panel (G). (**G**) IPA identified a down-regulated pro-inflammatory network in the five

consensus pathways (table 2), leading to decreased IL-8 expression. mRNA transcripts

encoding MAPK10, IKBKG, and EP300 were identified as binding targets of tRNA-fMet1 half in

the targeted chimeric eCLIP experiment.

Figure 5. Graphical abstract indicating the anti-inflammatory effect of tobramycin

mediated by 5′ tRNA-fMet halves in *P. aeruginosa* OMVs. (A) *P. aeruginosa* colonizes the

CF lungs and secrets OMVs. OMVs diffuse through the mucus layer overlying bronchial

epithelial cells and induce IL-8 secretion, which recruits excessive neutrophils and causes lung

damage. (**B**) Tobramycin increases 5′ tRNA-fMet halves in OMVs secreted by *P. aeruginosa*. 5′

tRNA-fMet halves are delivered into host cells and loaded into the AGO2 protein complex to

down-regulate protein expression of MAPK10, IKBKG, and EP300, which suppresses OMV-

induced IL-8 secretion and neutrophil recruitment. A reduction in neutrophils in BALF is

predicted to improve lung function and decrease lung damage.

 Tables

 Table 1. Top 10 most abundant and most differentially induced sRNAs in Tobi-OMVs

 compared to ctrl-OMVs

#	PA14 locus	Gene product of the locus	Log2FC	Average Log2CPM	length	Minimum free energy (kcal/mol)[A]
1	Multiple	tRNA-Asp	4.95	10.93	23 (20, 23)[B]	-0.2
2	Multiple	16S rRNA	3.98	10.83	33 (30-39)[B]	-6.9
3	Multiple	23S rRNA	5.57	10.35	44	-14.6
4	Multiple	tRNA-Ala	3.19	10.13	34	-8
5	62790	tRNA-fMet1	4.82	9.88	35	-7.5
6	28740	tRNA-Pro	2.70	9.39	36	-9.7
7	52320	tRNA-fMet2	2.74	8.95	35	-9.3
8	61760	tRNA-Gln	6.23	8.88	20	-0.2
9	30720	tRNA-Cys	3.41	8.85	40	-4.6
10	multiple	5S rRNA	2.72	8.75	45	-3.8

[A]The minimum free energy for each sRNA was predicted using the RNAfold web server (Gruber
et al., 2008)

[B]Multiple reads of different lengths were mapped to the same locus, and the most abundant
read is listed in Table 1 and Figure 2A.

Table 2. The consensus of significantly enriched signaling pathways identified using

three approaches[A].

Canonical pathway	Gene target prediction		Gene target validation		Protein expression	
	miRanda (1518)[B]		chimeric eCLIP (1936)[B]		Proteomics (2168)[B]	
	P value	z-score[C]	*P* value	z-score[C]	*P* value	z-score[C]
Integrin-linked kinase (ILK) Signaling	0.0005	-3.26	0.0041	-0.53	1.1E-08	-2.27
LPS-stimulated MAPK Signaling	0.0087	-3.46	0.0005	-3.60	7.7E-05	-0.68
HIF1a Signaling	0.0022	-3	0.0072	-4.35	0.0015	-0.87
IL-17A Signaling in Airway Cells	0.0011	-2.53	0.0085	-1.66	0.0077	-0.30
IL-6 Signaling	0.0002	-4.02	0.0389	-3.46	0.0141	-0.94

999 [A] There are 38 consensus pathways. Only consensus pathways predicted to downregulate IL-8

1000 secretion in epithelial cells by IPA are listed.

1001 [B] Number in parentheses indicates the number of genes/proteins used to perform pathway

1002 enrichment analysis.

1003 [C] Negative z-scores indicate that the pathways are predicted to be down-regulated.

1004

1005

Supplemental Figure 1. OMV effects on EGF, GRO, IL17a, and IP10. CF-HBECs from three

donors (n = 3) were polarized in ALI culture before being exposed to either the same number of

ctrl-OMVs, Tobi-OMVs, or 40% more Tobi-OMVs (1.4X Tobi-OMVs). After six-hour exposure,

the basolateral medium was collected and diluted 4-fold for 41-plex measurements**.** Cytokines

other than IL-8, EGF, GRO, IL17a, and IP10 were below the detection limit. Lines connect

experiments conducted with CF-HBECs from the same donor. Horizontal red lines and red dots

indicate means ± SEM. Linear mixed-effects models with CF-HBEC donor as a random effect

were used to calculate P values; *P < 0.05.

1014

1015

1016 **Supplemental Figure 2. 5′ tRNA-fMet1 half is over-expressed in tRNA1-OMVs compared**

1017 **to V-OMVs.** Small RNAs in V-OMVs and tRNA1-OMVs harvested from LB culture with

1018 arabinose were subjected to small RNA sequencing to quantify 5′ tRNA-fMet1 half (n = 3). ****

1019 FDR < 0.0001.

1020

1021 **Supplemental Table 1. BALF sample collection dates**

Subject	TOBI	Sample collection date	# of Days apart[A]
A	Off	4/26/2017	0
A	On	3/28/2018	336
B	Off	3/7/2018	0
B	On	8/29/2018	175
C	Off	4/1/2015	791
C	On	1/30/2013	0
D	Off	6/14/2017	0
D	On	6/27/2018	378

1022 [A]For each pair of samples, the earlier collection date is marked as day 0.