Identification of microRNAs in the West Nile virus vector *Culex tarsalis*

Sultan Asad[1], Ahmed M. Mehdi[2], Sujit Pujari[1,3], Claudia Rüeckert[4,5], Gregory D. Ebel[5], and Jason L. Rasgon[1*]

1. Department of Entomology, The Center for Infectious Disease Dynamics, and the Huck Institutes of the Life Sciences. The Pennsylvania State University, University Park, PA, USA

2. The University of Queensland, Brisbane, Australia Diamantina Institute, Faculty of Medicine, The University of Queensland, Translational Research Institute, Brisbane, QLD, Australia

3. Department of Pharmacology Physiology and Neuroscience, University of South Carolina School of Medicine, Columbia, South Carolina, USA

4. Department of Biochemistry and Molecular Biology, University of Nevada Reno, Reno, NV, 89557, USA

5. Center for Vector-borne Infectious Diseases, Department of Microbiology, Immunology and Pathology, College of Veterinary Medicine and Biomedical Sciences, Colorado State University, Fort Collins, CO 80523, USA

*Corresponding Author: Jason L. Rasgon, phone number 814-863-3668, fax 814-865-3048, email 16 address jlr54@psu.edu

Abstract

Background: microRNAs (miRNAs) represent a group of small non-coding RNAs that are crucial gene regulators of important biological functions including development and pathogen defense in most living organisms. Presently, there is a lack of availability of information regarding the miRNAs in the mosquito *Culex tarsalis,* which is one of the most important vectors of West Nile virus (WNV) in the United States. We used small RNA sequencing data and *in vitr*o and *in vivo* experiments to identify and validate the presence of a repertoire of miRNAs in *Cx. tarsalis* mosquitoes.

Results: Using bioinformatic approaches we analyzed small RNA sequencing data from the *Cx. tarsalis* CT embryonic cell line to discover 86 miRNAs. Consistent with other mosquitoes such as *Aedes albopictus* and *Cx. quinquefasciatus*, mi-184 was found to be the most abundant miRNA in *Cx tarsalis*. We also identified an additional 20 novel miRNAs from the recently sequenced *Cx. tarsalis* genome, for a total of 106 miRNAs identified in this study. The presence of selected miRNAs was biologically validated in both cell line and adult *Cx. tarsalis* mosquitoes using RT-qPCR and sequencing.

Conclusions: *Cx. tarsalis* is an important vector of many medically important pathogens including WNV and Western Equine encephalitis. Here we report a detailed insight into the miRNA population in *Cx. tarsalis* mosquitoes. These results will open new avenues of research deciphering the role of miRNAs in different *Cx. tarsalis* biological events such as development, metabolism, immunity and pathogen infection.

Background

Recent reports of the involvement of miRNAs in insect immunity and host-pathogen interactions have highlighted the need for investigating miRNAs in mosquitoes that have significant impacts on global health. miRNAs are small non-coding RNAs ranging from 19-25 nucleotides in length which regulate post transcriptional gene expression [1, 2]. In the animal kingdom, miRNAs regulate gene expression by binding imperfectly to the mRNAs either at the 3'-untranslated regions (3'-UTR), 5'-untranslated regions (5'-UTR), or coding regions. Though the 2-8 nucleotides at the 5' end of the miRNA ("seed region") perfect complementarity to the target transcript is integral for miRNA function, the rest of the miRNA sequence can have mismatches and bulges [1, 3, 4]. miRNAs are transcribed inside the nucleus into a primary miRNA transcript which is processed into a pre-miRNA by Drosha and Pasha. The pre-miRNA is then exported to the cytoplasm with the help of exportin proteins. Once in cytoplasm, the pre-miRNA is processed by Dicer I into the ~22 nucleotide mature miRNA. Once maturation is completed the mature miRNA is loaded into the microRNA induced silencing complex (miRISC) where it becomes single-stranded. The single-stranded miRNA-miRISC complex then targets the mRNA with sequence similarity leading to target mRNA degradation or modulation of protein expression [5, 6]. miRNAs are critical for a variety of cellular processes including development [7], immunity [8] and pathogen response [9, 10], which make them an interesting choice to investigate in the host pathogen interactions of vector mosquitoes and the medically important pathogens they carry.

More than 3000 mosquito species are present worldwide, but many important mosquito species are underrepresented in miRbase (a database of all miRNAs reported from

different species) [11]. Recent advances in sequencing platforms and bioinformatic approaches have allowed us to successfully identify the miRNAs from species whose genomes are not yet properly sequenced, assembled and annotated.

Cx. tarsalis is a neglected yet important vector of many medically important viruses such as WNV [12-14], West Equine Encephalitis (WEEV) [15], St. Louis Encephalitis (SLEV) [16] and Cache Valley virus (CVV) [17] in North America. Despite many studies focusing on the ecology, feeding behavior and vector competence of this mosquito, there is little information regarding miRNAs in *Cx. tarsalis*. Here we have mined and analyzed publicly available small RNA sequencing data of the *Cx. tarsalis CT* cell line to identify 86 high confidence miRNAs that are present in *Cx tarsalis* mosquitoes. Ten randomly chosen miRNAs were confirmed both *in-vitro* and *in-vivo* in mosquitoes.

Results and Discussion

Data Used in this study

We accessed publicly available small RNA data generated from the *Cx. tarsalis* CT cell line and passed it through our in-house pipeline (Figure 1) to identify miRNAs, using CLC genomic work bench 20. SRA read files described in Figure 2A were directly downloaded into CLC genomics workbench. All reads were trimmed by removing the RA3 adapter (TGGAATTCTCGGGTGCCAAGGG) sequences from the 3' end of raw sequencing reads. Any reads with low quality scores or that did not have an adapter sequence were excluded. After removal of the RA3 adapter sequences we saw a cluster of RNA sequences ranging from 21-24 nucleotides with the highest peak at 22

nucleotides which is the most abundant size of mosquito miRNAs (Figure 2B). In order to further examine the profile of the trimmed RNA reads, we used the web-based server sRNA toolbox [18] to determine that approximately 70 percent of the reads were mappable to miRbase (Figure 2C).

Discovery of miRNAs in *Cx tarsalis*

miRNAs have been demonstrated to play a pivotal role in insect development [7, 19], reproduction [20, 21], metabolism and longevity [22, 23], insecticide resistance [24, 25], immunity, and host-pathogen interactions [26, 27], and have been shown to govern processes relevant to public health in various mosquito species [28-30]. Using CLC genomic workbench 20 we annotated and mapped trimmed small RNA reads from 3 biological replicates from the *Cx. tarsalis* CT cell line to mature miRNAs of both *Ae. aegypti* and *Cx. quinquefasciatus* using default parameters. miRNAs discovered were considered to be true hits only if they were present in at least two out of three biological replicates. Our analysis found 60 and 84 mature miRNAs in *Cx. tarsalis* using *Cx. quinquefasciatus* (Figure 3A) and *Ae. aegypti* (Figure 3B) mature miRNAs as reference, respectively, with a 58 mature miRNA overlap using Jvenn online server (figure 3C, Supplementary table 1) [31]. Finally, we successfully identified a total of 86 high confidence miRNAs in *Cx. tarsalis* mosquitoes (Table 1).

Majority of *Cx. tarsalis* miRNAs arise from the 3' arm and sequence variations occur mostly at 3' arm of mature miRNAs

Mature miRNAs are formed as a result of a Dicer-mediated processing of precursor miRNA (pre-miRNA) having two arms/strands named 5p and 3p, leading to selective loading and retention of either 3p or 5p in the miRISC complex to become functionally active, while the other strand is cleaved out of the complex and degraded [5]. This strand selection has been demonstrated to be influenced by the thermodynamic instability of the duplex, 5' end starting nucleotides, and miRNA duplex length [32-34]. In our results we were able to detect mature miRNA that arose from 5p, 3p and both 5p and 3p of pre-miRNAs. Our data showed that 49 out of 86 miRNAs identified in this study were derived from the 3' arm of the pre-miRNA which indicates that in the *Cx. tarsalis* CT cell line, there may be a preference toward 3p arm processing of miRNAs (Figure 4A). However, this finding needs further experimental validation.

In each of the data libraries used in this study, adapter trimmed small RNA reads mapped to the reference sequences of mature miRNAs from miRbase showed variability. These variants with slight difference in length and sequence are commonly known as isomiRs. These isomiRs are usually produced due to imperfect processing/cleavage by Drosha or Dicer [35-37]. IsomiRs were long considered as an artifact, but recent advances in deep sequencing and computational algorithms have identified isomiRs in many species [38-40], including in *Ae. aegypti* [41]. IsomiRs are classified into three main categories: 5' isomiRs, 3' isomiRs, and polymorphic isomiRs with 5' and 3' isomiRs subclassified according to the nucleotide's insertions and deletions [42]. To find out and quantify the miRNA isomiRs, we used CLC Genomic workbench 20 to quantify the isomiRs based on nucleotide additions, insertions and substitutions. We focused on insertions and deletions of nucleotides at 3' or 5' of mature

miRNAs and limited our discovery to 2 nucleotide addition or deletion at the either end. IsomiRs that were present in all three biological replicates were used for analysis (Supplementary table 2). Our results show most modifications occurring at the 3' end of the mature miRNAs, in particularly 2 nucleotide insertion followed by the 1 nucleotide insertion (Figure 4B). Our results agree with previous studies that highlighted majority of sequence variations leading to isomiRs formation are at the 3' end of mature miRNAs [43]. Although the 5' variations are rare, they have more evolutionary and functional importance, as most of the 5' variations lead to a shift in the seed region that might affect regulatory ability of the miRNA [44]. Expanded analysis revealed that cta-miR-184 showed a total of 240 different types of isomiRs, followed by cta-miR-2940-3p that showed 161 different types of isomiRs. Although both cta-miR-184 and cta-miR-2940-3p showed the highest amount of isomiRs formation, there was a striking difference in the modification pattern as the former showed the majority of sequence variations at the 3' end (Figure 4C) while the later showed the highest amount of sequence variations at the 5' end (Figure 4D). These results warrant further investigation into the role of isomiRs in mosquito physiology and host pathogen interactions, for example in the case of WNV infections.

Validation of *Cx. tarsalis* miRNAs

We randomly selected 10 miRNAs: cta-miR-317-3p, cta-miR-7, cta-miR-999, cta-miR-71-3p, cta-miR-33, cta-miR-2940-5p, cta-miR-998, cta-miR-92b-3p, cta-miR-2951-5p and cta-miR-2945 for validation both *in vitro* (CT cell line) and *in-vivo* in 7 day old sugar fed female *Cx. tarsalis* mosquitoes (Figure 5A, B). All qPCR amplicons were cloned into

pJET and mature miRNA sequences confirmed through Sanger sequencing (Supplementary Figure 1).

Prediction and validation of novel miRNAs from the *Cx. tarsalis* genome

During the preparation of this manuscript, the *Cx. tarsalis* genome was published [45], which we used to identify novel miRNAs (present in all 3 replicates) that could be mapped to the *Cx. tarsalis* genome. Our results identified 20 novel miRNAs in *Cx. tarsalis* mosquitoes (Table 2). RNAfold (rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi) was used to determine the secondary structure of the predicted pre-miRNA sequences (Figure 6). Primers were designed for 12 randomly selected novel miRNAs and validated through RT-qPCR in both the CT cell line and *Cx. tarsalis* mosquitoes. Our results confirm that all 12 novel selected miRNAs were expressed in CT cells (Figure 7A), but in mosquitoes we were only able to detect 8 out of 12 (Figure 7B), likely reflecting physiological differences between cultured cells and live mosquitoes.

Conclusions

Our study has identified a total of 106 miRNAs, including 86 already reported in other mosquitoes, and 20 novel miRNAs that are present in the *Cx. tarsalis* genome. These results will lay the foundation for functional studies and open new avenues for research in *Cx. tarsalis* biology and pathogen transmission.

Methods

Small RNA libraries and data analysis

To identify *Cx. tarsalis* miRNAs we have used publicly available small RNA data of *Cx. tarsalis* CT cell line [46] from National Center for Biotechnology Information Sequence Read archive. A total of 3 libraries were downloaded. All libraries were generated using the TruSeq Small RNA Sample Prep Kit (Illumina) and sequenced on the Illumina HiSeq 2000 platform. The data were analyzed using CLC genome workbench 20. Libraries were trimmed of adapter sequences and the trimmed libraries were mapped to known miRNAs and miRNAs star strands of *Ae. aegypti* and *Cx. quinquefasciatus*. For novel miRNA prediction the miRDeep2 pipeline was used [47].

Mosquitoes and cell lines

Cx. tarsalis mosquitoes (YOLO strain) were reared at 26 ^{o}C with 40-50% relative humidity and a 16:8 hr light and dark cycle. The mosquitoes were provided with 10% sucrose solution *ad libitum*. The *Cx. tarsalis* cell line CT (generously provided by Dr Aaron Brault, CDC) was maintained at 28 ^{o}C with 0.5% CO_2 in Schneider medium supplemented with 10% of fetal bovine serum (GIBCO) and 1% antibiotics (GIBCO). Cells were passaged once per week.

RNA extractions and miRNA RT-qPCR

Qiazol was used to extract total RNA from both cells and mosquitoes following the manufacturer's suggested protocol. For mosquito RNA samples, 5 *Cx. tarsalis* sugar fed females (7 days old) were pooled per biological replicate. RNA was quantified using a Nanodrop and 2ug of total RNA was used to synthesize miRNA cDNA using the miScript RT kit following manufacturer's suggested protocol. cDNA was 10 times diluted and approximately 18ng of cDNA was used to validate miRNAs identified in this study using the miScript qPCR kit according to manufacturer's suggested protocol. All qPCR reactions were performed with 3 biological replicates and 2-3 technical replicates. U6 RNA was used as internal control for RT-qPCR. CT values were imported into Qiagen qPCR data analysis excel template to get normalized expression values.

Cloning and sequencing of miRNA amplicons

qPCR amplicons were seperated using 1% agarose gel electrophoresis. Single amplicon bands were cut using a sterile surgical blade and DNA was extracted with the Zymo Gel extraction kit according to manufacturer's suggested protocol. Eluted amplicons were directly cloned in to pJET1.2/Blunt vector according to manufacturer's suggested protocol. Positive clones were sent to GENEWIZ for Sanger sequencing. Sequencing data was analyzed with CLC genomic workbench 20.

Figure Legends

Figure 1. Schematic diagram of miRNA discovery pipeline Diagram showing use of an integrated approach of bioinformatics and wet laboratory experiments to identify *Cx. tarsalis* miRNAs.

Figure 2. Small RNA sequencing data A*) Culex tarsalis* small RNA data used in this study. B) Read length distribution after the adapter removal showed enrichment of reads around 21-24 nucleotides confirming the successful removal of adapter sequences. C) RNA profile of reads after adapter removal showing around 70 percent of read mapped to miRbase.

Figure 3. Identification of *Cx tarsalis* miRNAs A) Venn diagram of miRNAs identified across 3 biological replicates. B) Venn diagram showing common miRNAs identified within 3 biological replicates. C) Overlap of miRNAs identified with *Ae. aegypti* and *Cx. quinquefasciatus* annotation. D) 10 most abundant miRNAs in *Cx. tarsalis* RNA samples, each bar represents average normalized expression value of individual miRNA, error bars represent the standard deviation, and the dots shows the values of individual biological replicates.

Figure 4*. Cx. tarsalis* miRNA mostly arise from 3p arm having most of sequence variation at 3'end of miRNA A) Number of miRNAs processed from either 3p or 5p

arm of pre-miRNA. B) Overall sequence variations at 3' or 5' of mature miRNA sequences. C) Sequence variation found in cta-miR-184 showing most of the variations at the 3' end of the mature miRNA. D) Sequence variation found in cta-miR-2940-3p showing most of the variations at the 5' end of the mature miRNA.

Figure 5. Validation of *Cx. tarsalis* miRNAs. Primer extension-based confirmation of 10 randomly selected miRNAs with RT-qPCR using 3 biological replicates. U6 was used as reference for relative quantification. Graphs showing relative expression of cta-miRNAs/U6 in A) CT cell line B) *Cx. tarsalis* female mosquitoes. Error bars represent standard deviation and dots represent individual biological replicate values. Each biological replicate is made up of 5 *Cx. tarsalis* sugar fed 7 days old females.

Figure 6. Prediction of secondary structure of novel miRNAs. Stem-loop structures of predicted novel pre-miRNAs. Mature novel miRNAs are highlighted in red while their corresponding strand are highlighted in blue/magenta.

Figure 7. Validation of novel *Cx tarsalis* miRNAs. RT-qPCR based amplification of novel miRNAs in A) CT cells and B) *Cx. tarsalis* female mosquitoes. Error bars represent standard deviation and dots represent individual biological replicate values. Each biological replicate is made up of 5 sugar fed *Cx. tarsalis* 7 day old females.

Acknowledgements

This work was supported by NIH grants R01AI15025, R01AI128201, and R01AI116636 to JLR. We are thankful to Dr Aaron C. Brault from CDC for generously providing the CT cell line.

References

1. Bartel DP: **MicroRNAs: target recognition and regulatory functions**. *Cell* 2009, **136**(2):215-233.

2. Bartel DP, Chen CZ: **Micromanagers of gene expression: the potentially widespread influence of metazoan microRNAs**. *Nat Rev Genet* 2004, **5**(5):396-400.

3. Rigoutsos I: **New tricks for animal microRNAS: targeting of amino acid coding regions at conserved and nonconserved sites**. *Cancer Res* 2009, **69**(8):3245-3248.

4. Schnall-Levin M, Zhao Y, Perrimon N, Berger B: **Conserved microRNA targeting in *Drosophila* is as widespread in coding regions as in 3'UTRs**. *Proc Natl Acad Sci U S A* 2010, **107**(36):15751-15756.

5. Asgari S: **Chapter Two - microRNAs as Regulators of Insect Host–Pathogen Interactions and Immunity**. In: *Advances in Insect Physiology.* Edited by Smagghe G, vol. 55: Academic Press; 2018: 19-45.

6. Flynt AS, Greimann JC, Chung WJ, Lima CD, Lai EC: **MicroRNA biogenesis via splicing and exosome-mediated trimming in *Drosophila***. *Mol Cell* 2010, **38**(6):900-907.

7. Alvarez-Garcia I, Miska EA: **MicroRNA functions in animal development and human disease**. *Development* 2005, **132**(21):4653-4662.

8. Momen-Heravi F, Bala S: **miRNA regulation of innate immunity**. *J Leukoc Biol* 2018.

9.   Miesen P, Ivens A, Buck AH, van Rij RP: **Small RNA Profiling in Dengue Virus 2-Infected *Aedes* Mosquito Cells Reveals Viral piRNAs and Novel Host miRNAs**. *PLoS Negl Trop Dis* 2016, **10**(2):e0004452.

10.   Saldana MA, Etebari K, Hart CE, Widen SG, Wood TG, Thangamani S, Asgari S, Hughes GL: **Zika virus alters the microRNA expression profile and elicits an RNAi response in *Aedes aegypti* mosquitoes**. *PLoS Negl Trop Dis* 2017, **11**(7):e0005760.

11.   Kozomara A, Birgaoanu M, Griffiths-Jones S: **miRBase: from microRNA sequences to function**. *Nucleic Acids Res* 2019, **47**(D1):D155-D162.

12.   Goddard LB, Roth AE, Reisen WK, Scott TW: **Vector competence of California mosquitoes for West Nile virus**. *Emerg Infect Dis* 2002, **8**(12):1385-1391.

13.   Goldberg TL, Anderson TK, Hamer GL: **West Nile virus may have hitched a ride across the Western United States on *Culex tarsalis* mosquitoes**. *Mol Ecol* 2010, **19**(8):1518-1519.

14.   Venkatesan M, Broman KW, Sellers M, Rasgon JL: **An initial linkage map of the West Nile Virus vector *Culex tarsalis***. *Insect Mol Biol* 2009, **18**(4):453-463.

15.   Barnett HC: **The transmission of Western equine encephalitis virus by the mosquito *Culex tarsalis Coq***. *Am J Trop Med Hyg* 1956, **5**(1):86-98.

16.   Hammon WM, Reeves WC: **Laboratory Transmission of St. Louis Encephalitis Virus by Three Genera of Mosquitoes**. *J Exp Med* 1943, **78**(4):241-253.

17. Ayers VB, Huang YS, Lyons AC, Park SL, Higgs S, Dunlop JI, Kohl A, Alto BW, Unlu I, Blitvich BJ *et al*: **Culex tarsalis is a competent vector species for Cache Valley virus**. *Parasit Vector*s 2018, **11**(1):519.

18. Aparicio-Puerta E, Lebrón R, Rueda A, Gómez-Martín C, Giannoukakos S, Jaspez D, Medina JM, Zubkovic A, Jurak I, Fromm B, Marchal JA, Oliver J, & Hackenberg M: **sRNAbench and sRNAtoolbox 2019: intuitive fast small RNA profiling and differential expression**. *Nucleic Acids Research* 2019, **47**(W1), W530–W535.

19. Zhang Y, Zhou X, Ge X, Jiang J, Li M, Jia S, Yang X, Kan Y, Miao X, Zhao G *et al*: **Insect-Specific microRNA Involved in the Development of the Silkworm Bombyx mori**. *PLoS One* 2009, **4**(3):e4677.

20. Song J, Li W, Zhao H, Gao L, Fan Y, Zhou S: **The microRNAs let-7 and miR-278 regulate insect metamorphosis and oogenesis by targeting the juvenile hormone early-response gene Krüppel-homolog 1**. *Development* 2018, **145**(24):dev170670.

21. Zhang Q, Dou W, Pan D, Chen EH, Niu JZ, Smagghe G, Wang JJ: **Genome-Wide Analysis of MicroRNAs in Relation to Pupariation in Oriental Fruit Fly**. *Front Physiol* 2019, **10**:301.

22. Ling L, Kokoza VA, Zhang C, Aksoy E, Raikhel AS: **MicroRNA-277 targets insulin-like peptides 7 and 8 to control lipid metabolism and reproduction in Aedes aegypti mosquitoes**. *Proceedings of the National Academy of Sciences* 2017, **114**(38):E8017-E8024.

23. Ma K-S, Li F, Liu Y, Liang P-Z, Chen X-W, Gao X-W: **Identification of microRNAs and their response to the stress of plant allelochemicals in *Aphis gossypii* (Hemiptera: Aphididae)**. *BMC Molecular Biology* 2017, **18**(1):5.

24. Li X, Guo L, Zhou X, Gao X, Liang P: **miRNAs regulated overexpression of ryanodine receptor is involved in chlorantraniliprole resistance in *Plutella xylostella* (L.)**. *Scientific Reports* 2015, **5**(1):14095.

25. Etebari K, Afrad MH, Tang B, Silva R, Furlong MJ, Asgari S: **Involvement of microRNA miR-2b-3p in regulation of metabolic resistance to insecticides in *Plutella xylostella***. *Insect Mol Biol* 2018, **27**(4):478-491.

26. Saldaña MA, Etebari K, Hart CE, Widen SG, Wood TG, Thangamani S, Asgari S, Hughes GL: **Zika virus alters the microRNA expression profile and elicits an RNAi response in *Aedes aegypti* mosquitoes**. *PLOS Neglected Tropical Diseases* 2017, **11**(7):e0005760.

27. Wong RR, Abd-Aziz N, Affendi S, Poh CL: **Role of microRNAs in antiviral responses to dengue infection**. *Journal of Biomedical Science* 2020, **27**(1):4.

28. Dubey SK, Shrinet J, Sunil S: ***Aedes aegypti* microRNA, miR-2944b-5p interacts with 3'UTR of chikungunya virus and cellular target vps-13 to regulate viral replication**. *PLOS Neglected Tropical Diseases* 2019, **13**(6):e0007429.

29. Yan H, Zhou Y, Liu Y, Deng Y, Chen X: **miR-252 of the Asian tiger mosquito *Aedes albopictus* regulates dengue virus replication by suppressing the expression of the dengue virus envelope protein**. *J Med Virol* 2014, **86**(8):1428-1436.

30.   Slonchak A, Hussain M, Torres S, Asgari S, Khromykh AA: **Expression of Mosquito MicroRNA Aae-miR-2940-5p Is Downregulated in Response to West Nile Virus Infection To Restrict Viral Replication**. *Journal of Virology* 2014, **88**(15):8457-8467.

31.   Bardou P, Mariette J, Escudié F, Djemiel C, Klopp C: **jvenn: an interactive Venn diagram viewer**. *BMC Bioinformatics* 2014, **15**(1):293.

32.   Czech B, Hannon GJ: **Small RNA sorting: matchmaking for Argonautes**. *Nat Rev Genet* 2011, **12**(1):19-31.

33.   Han BW, Hung JH, Weng Z, Zamore PD, Ameres SL: **The 3'-to-5' exoribonuclease Nibbler shapes the 3' ends of microRNAs bound to Drosophila Argonaute1**. *Curr Biol* 2011, **21**(22):1878-1887.

34.   Lucas K, Raikhel AS: **Insect microRNAs: biogenesis, expression profiling and biological functions**. *Insect Biochem Mol Biol* 2013, **43**(1):24-38.

35.   Kuchenbauer F, Morin RD, Argiropoulos B, Petriv OI, Griffith M, Heuser M, Yung E, Piper J, Delaney A, Prabhu AL *et al*: **In-depth characterization of the microRNA transcriptome in a leukemia progression model**. *Genome Res* 2008, **18**(11):1787-1797.

36.   Starega-Roslan J, Krol J, Koscianska E, Kozlowski P, Szlachcic WJ, Sobczak K, Krzyzosiak WJ: **Structural basis of microRNA length variety**. *Nucleic Acids Res* 2011, **39**(1):257-268.

37.   Wu H, Ye C, Ramirez D, Manjunath N: **Alternative processing of primary microRNA transcripts by Drosha generates 5' end variation of mature microRNA**. *PLoS One* 2009, **4**(10):e7566.

38. Wyman SK, Knouf EC, Parkin RK, Fritz BR, Lin DW, Dennis LM, Krouse MA, Webster PJ, Tewari M: **Post-transcriptional generation of miRNA variants by multiple nucleotidyl transferases contributes to miRNA transcriptome complexity**. *Genome Res* 2011, **21**(9):1450-1461.

39. Lee LW, Zhang S, Etheridge A, Ma L, Martin D, Galas D, Wang K: **Complexity of the microRNA repertoire revealed by next-generation sequencing**. *RNA* 2010, **16**(11):2170-2180.

40. Amsel D, Vilcinskas A, Billion A: **Evaluation of high-throughput isomiR identification tools: illuminating the early isomiRome of *Tribolium castaneum***. *BMC Bioinformatics* 2017, **18**(1):359.

41. Etebari K, Osei-Amo S, Blomberg SP, Asgari S: **Dengue virus infection alters post-transcriptional modification of microRNAs in the mosquito vector *Aedes aegypti***. *Sci Rep* 2015, **5**:15968.

42. Neilsen CT, Goodall GJ, Bracken CP: **IsomiRs--the overlooked repertoire in the dynamic microRNAome**. *Trends Genet* 2012, **28**(11):544-549.

43. Newman MA, Mani V, Hammond SM: **Deep sequencing of microRNA precursors reveals extensive 3' end modification**. *RNA* 2011, **17**(10):1795-1803.

44. Tan GC, Chan E, Molnar A, Sarkar R, Alexieva D, Isa IM, Robinson S, Zhang S, Ellis P, Langford CF *et al*: **5' isomiR variation is of functional and evolutionary importance**. *Nucleic Acids Res* 2014, **42**(14):9424-9435.

45. Main BJ, Marcantonio M, Rasgon JL, Brown CT, Barker CM: **Whole genome assembly of *Culex tarsalis***. *G3 Genes|Genomes|Genetics* 2021, **11**(2):jkaa063.

46.   Rückert C, Prasad AN, Garcia-Luna SM, Robison A, Grubaugh ND, Weger-Lucarelli J, Ebel GD: **Small RNA responses of *Culex* mosquitoes and cell lines during acute and persistent virus infection.** Insect Biochem Mol Biol 2019, 109:13-23.

46. Friedlander MR, Mackowiak SD, Li N, Chen W, Rajewsky N: **miRDeep2 accurately identifies known and hundreds of novel microRNA genes in seven animal clades**. *Nucleic Acids Res* 2012, **40**(1):37-52.

Table 1. 86 miRNAs identified by homology to *Ae. aegypti* and *Cx. quinquefasciatus.*

S. no	Name	Sequence	Length	*Ae. aegypti*	*Cx. quinquefasciatus*
1	cta-bantam-3p	UGAGAUCAUUUUGAAAGCUGA	21	Y	Y
2	cta-bantam-5p	CCGGUUUUCAUUUUCGAUCUGAC	23	Y	Y
3	cta-let-7-5p	UGAGGUAGUUGGUUGUAUAGU	21	Y	Y
4	cta-miR-10-5p	ACCCUGUAGAUCCGAAUUUGUU	22	Y	Y
5	cta-miR-100-5p	AACCCGUAGAUCCGAACUUGUG	22	Y	Y
6	cta-miR-11-3p	CAUCACAGUCUGAGUUCUUGCU	22	Y	Y
7	cta-miR-11-5p	CGAGAACUCCGGCUGUGACC	20	Y	Y
8	cta-miR-12-3p	UGAGUAUUACAUCAGGUACUGGU	23	Y	Y
9	cta-miR-13-3p	UAUCACAGCCAUUUUGACGAGU	22	Y	Y
10	cta-miR-14-3p	UCAGUCUUUUUCUCUCUCCUAU	22	Y	Y
11	cta-miR-184-3p	UGGACGGAGAACUGAUAAGGGC	22	Y	Y
12	cta-miR-1889-3p	CACGUUACAGAUGGGGUUUCC	22	Y	Y
13	cta-miR-1889-5p	UAAUCUCAAAUUGUAACAGUGG	22	Y	Y
14	cta-miR-1890-3p	UGAAAUCUUUGAUUAGGUCUGG	22	Y	Y
15	cta-miR-190-5p	AGAUAUGUUUGAUAUUCUUGGUUG	24	Y	Y
16	cta-miR-210-3p	CUUGUGCGUGUGACAACGGCUAU	23	Y	Y
17	cta-miR-2525-5p	CUAAGUACUAGUGCCGCAGGAG	22	Y	Y
18	cta-miR-263a-5p	AAUGGCACUGGAAGAAUUCACGG	23	Y	Y
19	cta-miR-275-3p	UCAGGUACCUGAAGUAGCGC	20	Y	Y
20	cta-miR-276-3p	UAGGAACUUCAUACCGUGCUCU	22	Y	Y
21	cta-miR-276-5p	AGCGAGGUAUAGAGUUCCU	19	Y	Y
22	cta-miR-277-3p	UAAAUGCACUAUCUGGUACGAC	22	Y	Y
23	cta-miR-278-3p	UCGGUGGGACUUUCGUCCGUUU	22	Y	Y
24	cta-miR-279-3p	UGACUAGGAUCCACACUCAUUAA	22	Y	Y
25	cta-miR-281-3p	UGUCAUGGAAUUGCUCUCUUU	21	Y	Y
26	cta-miR-281-5p	AAGAGAGCUAUCCGUCGACAGU	22	Y	Y
27	cta-miR-283-5p	CAAUAUCAGCUGGUAAUUCUGGG	23	Y	Y
28	cta-miR-285-3p	UAGCACCAUUCGAAAUCAGUAC	22	Y	Y
29	cta-miR-2941-3p	UAGUACGGCUAGAACUCCACGG	22	Y	Y
30	cta-miR-2a-3p	UAUCACAGCCAGCUUUGAAGAGC	23	Y	Y
31	cta-miR-305-3p	CGGCACAUGUUGGAGUACAC	20	Y	Y
32	cta-miR-305-5p	AUUGUACUUCAUCAGGUGCUCU	22	Y	Y
33	cta-miR-306-5p	UCAGGUACUGAGUGACUCUCAG	22	Y	Y
34	cta-miR-308-3p	AAUCACAGGAGUAUACUG	18	Y	Y
35	cta-miR-31-5p	UGGCAAGAUGUUGGCAUAGCUGA	23	Y	Y
36	cta-miR-315-5p	UUUUGAUUGUUGCUCAGAAAGC	22	Y	Y
37	cta-miR-317-3p	UGAACACAGCUGGUGGUAUCU	21	Y	Y
38	cta-miR-33-5p	GUGCAUUGUAGUUGCAUUGCA	21	Y	Y
39	cta-miR-375-3p	UUUGUUCGUUUGGCUCGAGUUA	22	Y	Y
40	cta-miR-7-5p	UGGAAGACUAGUGAUUUUGUUGU	23	Y	Y
41	cta-miR-71-3p	UCUCACUACCUUGUCUUUCAUG	22	Y	Y
42	cta-miR-71-5p	AGAAAGACAUGGGUAGUGAGAU	22	Y	Y
43	cta-miR-79-5p	GCUUUGGCGCUUUAGCUGUAUGA	23	Y	Y
44	cta-miR-8-3p	UAAUACUGUCAGGUAAAGAUGU	22	Y	Y
45	cta-miR-8-5p	CAUCUUACCGGGCAGCAUUAGA	22	Y	Y
46	cta-miR-87-3p	GUGAGCAAAUUUCAGGUGUGU	22	Y	Y
47	cta-miR-92a-3p	AUUGCACUUGUCCCGGCCU	19	Y	Y
48	cta-miR-957-3p	UGAAACCGUCCAAAACUGAGGC	22	Y	Y
49	cta-miR-965-3p	UAAGCGUAUAGCUUUUCCCAUU	22	Y	Y
50	cta-miR-970-3p	UCAUAAGACACACGCGGCUAU	21	Y	Y
51	cta-miR-980-3p	UAGCUGCCUAGUGAAGGGC	19	Y	Y
52	cta-miR-988-3p	CCCUUGUUGCAAACCUCACGC	21	Y	Y
53	cta-miR-988-5p	UGUGUGCUUUUGUGACAACGAGA	22	Y	Y
54	cta-miR-996-3p	UGACUAGAUUACAUGCUCGU	20	Y	Y
55	cta-miR-998-3p	UAGCACCAUGAGAUUCAGC	19	Y	Y
56	cta-miR-999-3p	UGUUAACUGUAAGACUGUGUCU	22	Y	Y
57	cta-miR-9c-3p	UAAAGCUUUAGUACCAGAGGUC	22	Y	Y
58	cta-miR-9c-5p	UCUUUGGUAUUCUAGCUGUAGA	22	Y	Y
59	cta-miR-2952-3p	AGAGCUCAGCACGCAGGGGUGGC	23		Y
60	cta-miR-2951-5p	UAGUACGGCCAUGACUGAGGGC	22		Y
61	cta-miR-11893-3p	UUCCUGCACUUAUACGCUUACCU	22	Y	
62	cta-miR-11899-3p	UAUGACCGAUUUAAUAUAUGGCU	23	Y	
63	cta-miR-12-5p	UGAGUAUUACAUCAGGUACUGGU	23	Y	
64	cta-miR-13-5p	CAGUACUUAUGUUAUGCUCUCU	22	Y	
65	cta-miR-275-5p	CGCGCUAAGCAGGAACCGAGAC	22	Y	
66	cta-miR-277-5p	CGUGUCAGAAGUGCAUUUACA	21	Y	
67	cta-miR-278-5p	ACGGACGAUAGUCUUCAGCGGCC	23	Y	
68	cta-miR-282-3p	ACAUAGCCUGACAGAGGUUAGG	22	Y	
69	cta-miR-286a-3p	UGACUAGACCGAACACUCGCGUCCU	25	Y	
70	cta-miR-286b-3p	UGACUAGACCGAACACUCGUAUCCC	25	Y	
71	cta-miR-2940-3p	GUCGACAGGGAGAUAAAUCACU	22	Y	
72	cta-miR-2940-5p	UGGUUUAUCUAAUCUGUCGAGGC	23	Y	
73	cta-miR-2944a-5p	GAAGGAACUUCUGCUGUGAUCUGA	24	Y	
74	cta-miR-2944b-3p	UAUCACAGCAGUAGUUACCUGA	22	Y	
75	cta-miR-2945-3p	UGACUAGAGGCAGACUCGUUUA	22	Y	
76	cta-miR-2945-5p	AGCGGGUCCGUUUCUAGUGUCAUG	24	Y	
77	cta-miR-2b-3p	UAUCACAGCCAGCUUUGAUGAGCU	24	Y	
78	cta-miR-2c-3p	UAUCACAGCCAGCUUUGAUGAGC	23	Y	
79	cta-miR-307-3p	CACAACCUCCUUGAGUGAGCGA	22	Y	
80	cta-miR-308-5p	CGCGGUAUAUUCUUGUGGCUUG	22	Y	
81	cta-miR-34-5p	UGGCAGUGUGGUUAGCUGGUUG	22	Y	
82	cta-miR-79-3p	UAAAGCUAGAUUACCAAAGCAU	22	Y	
83	cta-miR-92a-5p	CGGUACGGACAGGGGCAACAUU	22	Y	
84	cta-miR-92b-3p	CGGUACGGACAGGGGCAACAUU	22	Y	
85	cta-miR-92b-5p	AAUUGCACUUGUCCCGGCCUGC	22	Y	
86	cta-miR-9b-5p	UCUUUGGUUGAUUUUAGCUGUAUGC	24	Y	

Table 2. Novel miRNAs identified from the *Cx. tarsalis* genome.

S. no	Name	Sequence	Length	Predicted Consensus Precursor Sequence
1	cta-novel-1	UUAACAAUAUUUGUGUGACCUG	22	UUAACAAUAUUUGUGUGACCUGUUGUUAAUACGAUAUGAAGGUCACACGAAUAUUGCUAAGU
2	cta-novel-2	UGACUAGAGGCAGACUCGUUU	21	AGCGGGUCCGUUUCUAGUGUCAUGUGCUACGUUUGAAAGUCAUGACUAGAGGCAGACUCGUUU
3	cta-novel-3	UUUUAGACCGGUUUUGAACAC	21	UGUUCAAAACCGGUCUAAAACGUGAUUUUUCCGUUUUAGACCGGUUUUGAACAC
4	cta-novel-4	AGAUAGGACCUUUUGAAAAAGU	22	UUUUUCAAAAGAUCCCAUCUGCUAUGGGUUUUCUAUGCAGAUAGGACCUUUUGAAAAAGU
5	cta-novel-5	AUGGCGCUGUAAAAAAGCUUUUU	23	AAAGCUUUACACAACGCCAUCUAUCAGAAAAACUUAAUGACAAUCAGAUGGCGCUGUAAAAAAGCUUUUU
6	cta-novel-6	UAUGCUAUCUUGGGACAUGUCU	22	UAUGCUAUCUUGGGACAUGUCUGCUGUAAAAAGAUAGGACGUGUCACAAGAUAGCACACA
7	cta-novel-7	CAUGAUAUCUUGUGACAUGUCU	22	CAUGAUAUCUUGUGACAUGUCUGCUGUAAAAAGAUAGGACGUGUCACAAGAUAGCACA
8	cta-novel-8	AGCCGGACAGCCUCGACCUGGA	22	AGCCGGACAGCCUCGACCUGGACUACGACAUGUGGCAGGGCCAGUUCGAGUUUGUCGGGCCGGC
9	cta-novel-9	GAAGGAACUUCUGCUGUGAUCU	22	GAAGGAACUUCUGCUGUGAUCUGAGAUGUGUUCAUAUCACAGUAGUUGUACUUUAA
10	cta-novel-10	UGACUAGACCGAACACUCGUAUC	23	GGCGAUUGUCGGCUUGGUCGCUGUCUUUACCCAAGGCACAGUUGCACUUCAGUGACUAGACCGAACACUCGUAUC
11	cta-novel-11	UAUGACCGAAUUAAUGUAUGGCU	23	CCAUAUAUGAUUUUGGUCAUAUAUUGAAUUAUGAUAUUUUAUAUGACCGAAUUAAUGUAUGGCU
12	cta-novel-12	UUCCUGACUUAUAUACGCUUACCCCU	24	GUAAGUAGAUAAUUCAGAAAGGCGGCAACGUUUCUUCCUGACUUAUAUACGCUUACCCCU
13	cta-novel-13	UGUAACCCGUCAUGAACUGUCA	22	UGUAACCCGUCAUGAACUGUCAAAUUGGUGAAGCAGUUUGACAGUUCUUGUUGGGUUACAUU
14	cta-novel-14	CAAACCGGAAGUUGUAAACUGU	22	AGUUCAUCAUUUCCGGUUUGAUAGAGGAAUCAAACCGGAAGUUGUAAACUGU
15	cta-novel-15	AUCCCUGCCUUCGGAUGCCUA	21	AAAGAGGUAGGGAUAACUCGCCAUCCCUGCCUUCGGAUGCCUA
16	cta-novel-16	UAUGCUAUCGUGUGACAUGUCU	22	UAUGCUAUCGUGUGACAUGUCUGCUGUAAAAAAAUAGGACGUGUCACAAGAUAGCACUCA
17	cta-novel-17	UUUCGUUCCGGACACAAAAAUC	22	UUUCGUUCCGGACACAAAAUCAGGAAUCGGCCUGAGGAUUUUUGUUUCCGGAAUGUCCACAGAAC
18	cta-novel-18	UCUACCGAUCGAUCUUCAUAGC	22	CGUGAAGAUCUUCCGGUGCAGACAGGCUCGUUGAGGCCACUCAAAGUCACCAGCGUGUCUACCGAUCGAUCUUCAUAGC
19	cta-novel-19	AACAAAGCCGAUCGGAAAAGU	21	UUUUUCGUCGGCAGUGGCCAGUGUGAGAUACGCGCAACAAAGCCGAUCGGAAAAGU
20	cta-novel-20	GUAGAAUCGCGUCAAAUCGACUGA	24	AGUCGAGUGAUGCAAUUCUUUCCGCAAACAACUUUUGCCCACUUUCGCGUAGAAUCGCGUCAAAUCGACUGA

Figure 1

Download SRA files form NCBI in CLC Genome work bench 20.04

Identification and removal of adapter sequence

Discard sequences with less **than** 5 reads

Discard reads **without** Adapter sequence

Annotate Trimmed reads with miRbase 22.1 with *Culex quinquefasciatus mature* miRNAs as reference

Annotate Trimmed reads with miRbase 22.1 with *Aedes aegypti* mature miRNAs as reference

Export the mature miRNA list in Excel

miRNA considered for discovery when identified in at least in two out of three biological replicates

Pooled list of miRNAs discovered after miRbase annotation with both *Culex quinquefasciatus* and Aedes aegypti mature miRNAs

Validation of randomly selected miRNAs with qPCR

Figure 2

A

SRA ID	Sample	Raw Reads	Number of Bases in millions
SRR8936281	Mock infected CT cells 2dpi rep1	8808220	669.4
SRR8936288	Mock infected CT cells 2dpi rep2	6848904	520.4
SRR8936289	Mock infected CT cells 2dpi rep3	7797739	197.7

B

C

Figure 3

A **Culex quinquefasciatus**

B **Aedes aegypti**

C **Overlap between miRNAs identified with *Aedes* and *Culex* as reference**

Aedes aegypti   *Culex quinquefasciatus*

D

Figure 4

Figure 5

A

CT cell line

B

***Cx. tarsalis* female mosquitoes**

Figure 6

cta-novel-miR-1
cta-novel-miR-2
cta-novel-miR-3
cta-novel-miR-4
cta-novel-miR-5
cta-novel-miR-6
cta-novel-miR-7
cta-novel-miR-8
cta-novel-miR-9
cta-novel-miR-10
cta-novel-miR-11
cta-novel-miR-12
cta-novel-miR-13
cta-novel-miR-14
cta-novel-miR-15
cta-novel-miR-16
cta-novel-miR-17
cta-novel-miR-18
cta-novel-miR-19
cta-novel-miR-20

Figure 7

A

CT cell line

B

***Cx. tarsalis* female mosquitoes**