Negative regulation of IMD contributes to disease tolerance during systemic bacterial infection in *Drosophila*

Arun Prakash*, Katy M. Monteith, Pedro F. Vale*

Institute of Evolutionary Biology

School of Biological Sciences

University of Edinburgh

Edinburgh EH9 3FL

*Correspondence

arun.prakash@ed.ac.uk

pedro.vale@ed.ac.uk

Keywords: Disease tolerance, Immune deficiency (IMD) pathway, Immunopathology, Negative immune regulators, Pathogen clearance

Abstract

Disease tolerance is an infection phenotype where hosts show relatively high health despite harbouring elevated pathogen loads. Compared to the mechanisms of immune clearance our knowledge of the mechanisms underlying increased tolerance remains incomplete. Variation in the ability to reduce immunopathology may explain why some hosts can tolerate higher pathogen burdens with reduced pathology. Negative immune regulation would therefore appear to be a clear candidate for a mechanism underlying disease tolerance but this has not been tested directly for bacterial infections. Here, we examined how the negative regulation of the immune deficiency (IMD) pathway affects disease tolerance in *Drosophila melanogaster* when infected with the gram-negative bacterial pathogen *Pseudomonas entomophila*. We find that UAS^RNAi-mediated reduced expression of the negative regulators of IMD (*pirk* & *caudal*) severely reduced the ability to tolerate infection in both males and females across a wide range of infectious doses. While flies unable to regulate the IMD response exhibited higher expression of antimicrobial peptides and lower bacterial loads as expected, this was not accompanied by a proportional reduction in mortality. Instead, tolerance (measured as fly survival relative to its microbe load) was drastically reduced, likely due to the combination of increased immunopathology and cytotoxicity of elevated AMP expression. Our results therefore highlight that in addition to regulating an efficient pathogen clearance response, negative regulators of IMD also contribute to disease tolerance.

Introduction

To survive infection hosts must not only identify and eliminate the pathogen but must also reduce infection-induced tissue damage. Tissue damage may arise either directly from pathogen growth and secreted virulence factors, or as a negative side-effect of the immune response in the form of immunopathology (Graham, Allen, and Read 2005; Sears et al. 2011; Badinloo et al. 2018). For example, the upregulation of antimicrobial peptides (AMPs) after pathogenic infection (Badinloo et al. 2018; De Gregorio et al. 2002; Zerofsky et al. 2005), the accumulation of toxic/damage molecules such as reactive oxygen species (*ROS*) (Han and Ulevitch 2005; Schneider 2007; Lambeth 2007), and the phenoloxidase (*PO)* response (a fast-acting immune effector in insects) are all known to cause tissue damage (Sadd and Siva-Jothy 2006; Cerenius, Lee, and Söderhäll 2008; Khan, Agashe, and Rolff 2017). Overall, the fine balance between clearing pathogens and avoiding tissue damage requires tightly regulated induction and resolution of immune responses (Sears et al. 2011; Soares, Gozzelino, and Weis 2014).

Understanding the distinct pathogen- or host-derived causes of tissue damage is important not only from a therapeutic perspective (Vale, Fenton, and Brown 2014; Vale et al. 2016; Louie et al. 2016), it also offers important mechanistic explanations for the widely variable infection phenotypes observed between individuals (Hill 1998; Råberg, Graham, and Read 2009; Kutzer and Armitage 2016). The field of host-pathogen evolutionary ecology typically describes individuals with low pathogen loads and relatively high health during infection as 'resistant' because they are able to clear - or otherwise quantitatively reduce - the establishment and growth of pathogens within the host (Hill 1998; Råberg, Graham, and Read 2009; Lazzaro, Sceurman, and Clark 2004; Malo and Skamene 1994). Other infection phenotypes are more difficult to explain as the consequence of immune clearance, such as the observation that some individuals exhibit low levels of pathology (measured as either mortality or physiological performance), despite harbouring relatively high pathogen loads (Råberg, Graham, and Read 2009; Kutzer and Armitage 2016; Raberg, Sim, and Read 2007; Ayres and Schneider 2012). Instead, tolerating higher pathogen loads is more likely to be under the control of mechanisms that prevent, limit or repair tissue damage (Soares, Gozzelino, and Weis 2014; Vale, Fenton, and Brown 2014; Soares, Teixeira, and Moita 2017; R. Martins et al. 2019). However, compared to the detailed knowledge of immune clearance mechanisms we associate with resistance, our knowledge of the specific mechanisms underlying increased tolerance remains limited (Kutzer and Armitage 2016). Previous work in *Drosophila* has employed GWAS & transcriptomics to uncover several candidate genes associated with disease

tolerance, including many that do not have an obvious immune related function such as *CrebA*

(Troha et al. 2018), *grainyhead and debris buster* (Howick and Lazzaro 2014), *dFOXO* (Dionne et al.

2006; Lissner and Schneider 2018). Curiously, genes related to canonical immune pathways are

conspicuously absent from these genome-wide screens of tolerance, even though we would expect

the regulation of inflammation to be a key mechanism underlying disease tolerance (Sears et al.

2011; Vale, Fenton, and Brown 2014; Ayres and Schneider 2012; Merkling et al. 2015).

In the present work, we take a more targeted approach to specifically investigate if the negative

regulation of immune responses has any measurable effect on the phenotype of disease tolerance.

We focused on the immune deficiency (IMD) pathway, one the best described immune signalling

pathways in *Drosophila* (Leulier et al. 2003; Bruno Lemaitre and Hoffmann 2007; Myllymäki,

Valanne, and Rämet 2014). Following infection, DAP-type peptidoglycans from gram-negative

bacteria are recognised by the pathogen-sensing receptors *PGRPs (LC & LE)* (Vodovar et al. 2005;

Dieppois et al. 2015). These receptors then recruit the adaptor molecule IMD and activate the

intracellular signalling cascade that leads to the activation of a *NF-κB relish* (see **Fig**. 1) also

reviewed in (Myllymäki, Valanne, and Rämet 2014; Kuraishi, Hori, and Kurata 2013). Upon

activation, *relish* translocates to the nucleus and induces transcription of a set of IMD-responsive

AMPs. Minutes after pathogen recognition, a cocktail of AMPs such as *Diptericins, Attacins &*

Cecropins is secreted by the fat body (Bruno Lemaitre and Hoffmann 2007; B. Lemaitre, Reichhart,

and Hoffmann 1997). *Relish* also induces the transcription of negative regulators of IMD (*pirk,*

caudal, PGRPs-LB, LC, LE,) which ensures an appropriate level of immune response, while

avoiding immunopathology (Lee and Ferrandon 2011; Kleino and Silverman 2014). *Pirk*, for

example, interferes with the interaction of *PGRP-LC & -LE* with the molecule IMD, and limits

the activation of the IMD pathway (see **Fig**. 1) (Aggarwal et al. 2008; Kleino et al. 2008; Basbous

et al. 2011). Similarly, the homeobox protein *caudal* downregulates the expression of AMPs in flies

(Ryu et al. 2008). Taken together, the negative regulation of IMD-pathway ensures an appropriate

level of immune response and enables individuals to reduce immune activity in response to

reductions in bacterial numbers (Lee and Ferrandon 2011; Kleino et al. 2008; Paredes et al. 2011).

We measured disease tolerance during systemic infection with the gram-negative bacterium

Pseudomonas entomophila using *Drosophila* lines where individual components of the IMD-pathway

were knocked down using UAS^{RNAi} including (i) the transcription factor- *relish*, its negative

regulators (ii) *pirk* and (iii) *caudal*, and (iv) a major IMD-responsive AMP *Diptericin-B*, because of

substantial evidence that *Diptericins* are very effective during defense against Gram-negative

bacteria (Bruno Lemaitre and Hoffmann 2007; Wicker et al. 1990). Given the widespread evidence

of sex differences in immunity and response to infection, reviewed in (Klein and Flanagan 2016;

Belmonte et al. 2020), we also specifically tested whether female and male flies differ in their ability

to tolerate bacterial infection.

Figure 1: A simplified version of *Drosophila* IMD (immune deficiency) pathway activated by the
DAP type pedptidoglycan following the gram-negative bacterial infection (*Pseudomonas entomophila*).
These receptors then recruit the adaptor molecule Imd and activate the intracellular signalling
cascade that leads ↘   to the activation of a NF-κB *relish*. Upon activation, relish translocates ↘
to the nucleus and induces transcription of a set of cocktail of IMD-responsive AMPs such as
Diptericins, Attacins & Cecropins, minutes after pathogen recognition. The negative regulators ⊢
of IMD such as *pirk, caudal, PGRPs-LB* ensure an appropriate level of immune response, while
avoiding immunopathology

Materials and methods

<u>Fly strains and husbandry</u>: We used the following fly lines: UAS-*relish*[RNAi] (*Bloomington stock#*

33661), UAS-*caudal*[RNAi] (*Bloomington stock# 57546*), UAS-*pirk*[RNAi] (*Bloomington stock# 67011*) & UAS-

*Diptericin*B[RNAi] (*Bloomington stock# 28975*) also see Supplementary Information **Table** S1 for details

about fly genotypes. All fly lines were obtained from the Bloomington *Drosophila* stock centre at

the Indiana University. Each of the transgenic fly line contains a homologous hairpin under the
control of the Upstream Activating Sequence (UAS^{RNAi}) to a Gal4 temperature-sensitive
transcriptional activator (*Bloomington stock# 67065*). This fly line drives the Gal4 expression in the
entire body of fly and contains a temperature sensitive Gal80, which inhibits Gal4 at the permissive
temperature (25℃). All fly stocks were reared in plastic vials (12 ml) on a standard Lewis medium
at 25±℃ with a 12 hour light: dark on 14 days cycle (by transferring flies into new food vials). All
the crosses were made at 25℃ and ~3 days after eclosion, offspring were transferred to agar vials
(~12 flies per vial) at the non-permissive temperature (29℃) for two days to facilitate silencing of
the candidate genes until experimental day.

<u>Bacterial culture preparation</u>: The gram-negative bacterium *Pseudomonas entomophila* has a broad
host range, infecting insects, nematodes, plants, and higher vertebrates (Dieppois et al. 2015). In
flies, *P. entomophila* infection results in pathology to the gut epithelium during systemic and oral
infections, eventually causing death (Vodovar et al. 2005; Dieppois et al. 2015; Troha and Buchon
2019). To obtain pure cultures, frozen bacterial stock cultures (-80℃) were streaked onto fresh LB
agar plates and single colonies were inoculated into 20 ml LB broth and incubated overnight at
37°C with shaking at 120 rpm (revolutions per minute). The overnight cultures were diluted 1: 100
into 500 ml of fresh LB broth and incubated again at 37°C with shaking at 120 rpm. At the mid-
log phase (optical density $OD_{600} = {\sim}0.75$), we harvested the bacterial cells by centrifugation at
5000 rpm for 15 min at 4°C and re-suspended the bacterial pellet in 5% sucrose solution. The final
inoculum was adjusted to $OD_{600} = 1.0$, and this was the starting bacterial inoculum used for all
flies for systemic infection.

<u>Figure 2</u>: Design of experiments to assay **(1)** gene expression of the major IMD-regulated AMPs
(2) internal bacterial load and **(3)** survival following systemic bacterial infection with *Pseudomonas*
entomophila to test impact of the negative regulation of IMD pathway on phenotype of disease
tolerance **(4)**.

Gene expression: We analysed the antimicrobial peptides (AMPs) gene expression data to examine

the effects of knocking down negative regulators of IMD-pathway on the fly's immune response.

We quantified the expression of major IMD-responsive AMPs namely, *diptericin (Dpt), attacin-C*

AttC), cecropin-A1 (CecA1), by quantitative real-time polymerase chain reaction (qRT-PCR). We

randomly selected a subset of IMD-pathway mutants & functional flies both males & females at

8h post-infection for RNA extraction, we included 15 flies i.e., 5-7 replicates of 3 flies pooled

together for each treatment (Knockdown & functional) and sexes. We first homogenised whole

flies (3 in group) using sterile micro-pestles into 1.5µl microcentrifuge tubes containing 80µL of

TRIzol reagent (Invitrogen, Life Technologies). The tubes containing the fly homogenate were

kept frozen at -70°C until RNA extraction. We performed mRNA extractions using the standard

phenol-chloroform method and included a DNase treatment (Ambion, Life Technologies). We

confirmed the purity of eluted samples using a Nanodrop 1000 Spectrophotometer (version 3.8.1)

before proceeding with RT (reverse transcription). The cDNA was synthesized from 2µL of the

eluted RNA using M-MLV reverse transcriptase (Promega) and random hexamer primers, and

then diluted 1:1 in nuclease free water. We then performed quantitative RT-PCR (qRT-PCR) on

an Applied Biosystems StepOnePlus machine using Fast SYBR Green Master Mix (Invitrogen

Ltd.) using a 10 µL reaction containing 1.5 L of 1:1 diluted cDNA, 5 µL of Fast SYBR Green

Master Mix a 3.5 µL of a primer stock containing both forward and reverse primer at 1 µM

suspended in nuclease free water (final reaction concentration of each primer being 0.35 µM). For

each cDNA sample, we performed two technical replicates for each set of primers and the average

threshold cycle (Ct) was used for the analysis. We obtained the AMP primers from Sigma-Aldrich

Ltd (see supplementary **Table** S2 for information about primers). We optimised the annealing

temperature (T_a) and the efficiency (Eff) of the *Dpt* primer pair was calculated by 10-fold serial

dilution of a target template (each dilution was assayed in duplicate); *Dpt*: T_a= 59 °C, Eff= 102%;

AttC: T_a= 60 °C, Eff= 94%; *CecA1*: T_a= 59.5 °C, Eff= 109%.

Fly survival following bacterial infection: To test the impact of *Pseudomonas entomophila* infection on fly

survival, we challenged experimental flies to 4 different bacterial concentrations (infection doses)

ranging between (I) ~650 cells (II) ~1400 cells (III) ~2000 cells (IV) ~2900 cells/µl of *P.*

entomophila culture, obtained by serial dilution. Flies (n=10-18 fly vials /sex/infection dose/fly line)

were infected systemically by intra-thoracic pricking (Khalil et al. 2015) with a needle immersed in

bacterial suspension under minimal CO_2 anaesthesia. The effect of injury caused by pricking was

controlled by including mock infections performed using a needle dipped in sterile 5% sucrose
solution. We observed the mortality for 10 days post-infection every 3 hours (±0.5h) for the first
2 days (between 6 am and 11 pm) and every 12 hours for the following 7 days.

Quantification of bacterial load following bacterial infection: During the survival assay, flies were randomly
sampled (10-18 live fly vials/sex/infection dose/fly line) and the bacterial load was quantified at
~8 hours following infection. Initially, flies (group of 3) were transferred to a 1.5 ml
microcentrifuge tubes and surface sterilized by adding 70% ethanol for ~30 secs to ensure we
counted only colony forming units (CFUs) inside the flies. Ethanol was then removed by briefly
washing flies twice with sterile distilled water. To ensure that this method was efficient in surface
cleaning flies, we plated 4 µl of the second wash on LB agar (from random flies) ensuring that the
flies are free from surface bacteria (i.e., no viable CFUs). We then placed in 1.5 ml Eppendorf
tubes and added 100 µl of LB broth to each tube and the flies were thoroughly homogenized using
a motorized pestle. We immediately plated this homogenate in serial dilutions ranging from 100 to
10^{-6} using 1xPBS (phosphate buffer) on LB agar plates and incubated at 30°C for ~18 hours after
which viable CFUs were counted manually (Siva-Jothy et al. 2018).

Statistical analysis

1. *Gene expression*: We analysed the gene expression data by first calculating the ΔCT value
(Livak and Schmittgen 2001) for the expression of a gene of interest relative to the
reference house-keeping gene rp49 (Gupta and Vale 2017). We then used analysis of
variance (ANOVA) model using the R package 'ggplot2' (Wickham 2016; Bunn and
Korpela, 2019) to test for the effects of the experimental treatment on the ΔCT values of
each gene i.e., whether the AMPs expression differed significantly between knockdown
and fully functional flies of each IMD-pathway fly line.

2. *Bacterial load*: We then quantified differences in bacterial load following bacterial infection
by analysing the bacterial measurement ($\log_{10}$) at ~8h post-infection after *P. entomophila*
infections. We found that residuals of bacterial load data were normally distributed when
tested with Shapiro-Wilks test in 'R' and hence we used ANOVA and analysed the bacterial

load data by fitting 'knockdown' (functional flies & IMD-pathway knockdown flies) and

'sex' as categorical fixed-effects, while infection dose (4 concentrations of *P. entomophila*)

and 'vials' (replicates) as a random-effects for each of the fly lines (functional flies & IMD-

pathway knockdown flies).

3. *Survival post-infection*: We analysed the survival data after *P. entomophila* bacterial infection for

all the IMD-pathway mutants by a mixed effects Cox model using the R package 'coxme'

(Therneau 2015) for both males and females respectively. In each case, we specified the

model as: survival ~ knockdown *(functional & knockdown)* x sex *(male & female)* x dose *(4 concentrations of P. entomophila)*

(1|vial *(n replicates)*), with 'knockdown' and 'sex' as fixed effects, and vials as a random effect

for all the fly lines (IMD-pathway mutants). We fit separate models for each fly lines during

P. entomophila infections, since they were assayed on different days. Next, to understand

how different knockdown flies (i.e., having fully functional vs. knockdown of IMD-

pathway) differ in survival to bacterial infections in detail, we also analysed survival data

for *P. entomophila* infections for each of the IMD-pathway mutant/fly lines separately using

Cox Proportional Hazard survival analysis (Package 'survival'). We estimated the impact

of harbouring IMD-pathway knockdowns as the hazard ratio of IMD-knockdowns vs.

fully functional fly lines. A hazard ratio significantly greater than one indicates higher risk

of mortality in the IMD-mutants relative to fully functional individuals; hence, a significant

impact on fly's survival.

4. *Measuring disease tolerance*: Finally, to understand how the negative immune regulation

contribute to disease tolerance during bacterial infections we plotted tolerance by

calculating (median lifespan/mean bacterial load) for male & female flies (i.e., flies with

functional & knockdown IMD pathway regulators) after bacterial exposure with 4 different

infection doses. Tolerance is usually measured as a reaction-norm or slope which could be

either linear or non-linear, where the pathogen burden is plotted on the x-axis and health

or fitness is plotted on the y-axis (Raberg, Sim, and Read 2007; Ayres and Schneider 2012;

Medzhitov, Schneider, and Soares 2012; Louie et al. 2016; Gupta and Vale 2017; Oliveira,

Bahia, and Vale 2020). This standard reaction norm approach that measures tolerance as

the rate of health decrease for increasing microbe loads is only statistically sound when

mortality and microbe loads measurements are taken from the same individual or from the

same group of replicate individuals. Given the destructive sampling required to measure
microbe loads in flies we chose instead to use another commonly used metric of tolerance
using the average mortality of flies of a given line relative to the average bacterial load of
that line, see for example (Adelman et al. 2013; Knutie et al. 2017; Cornet et al. 2014). This
is sometimes referred to as 'point' tolerance and is intuitive to interpret: relative to a given
microbe load, does one line die more (less tolerant) or less (more tolerant) (Little et al.
2010).

Results & discussion:

1. <u>Loss of IMD negative regulation leads to increased AMP expression and decreased</u>
<u>bacterial load</u>

We first began by quantifying major IMD-regulated AMP expression (*AttC, CecA1 & Dpt*)
following infection with *P. entomophila* in both male & female flies (**Fig. 3A, Table-S3)**. As
expected, *relish* knockdown flies had lower AMP expression while loss of negative regulators *caudal*
& *pirk* resulted in higher AMP expression (**Fig. 3A, Table-S3)**. Next, we wanted to test whether
AMP expression correlated with the bacterial replication in flies. Consistent with their AMP
expression, both male and female flies with reduced negative regulation (*caudal* & *pirk*) exhibited a
reduction in bacterial numbers while flies lacking *relish* exhibited higher levels of bacterial numbers
(compare **Fig. 3A** gene expression **& Fig. 3B** bacterial load data), clearly indicating the expected
importance of AMPs in defence against this bacterial pathogen (**Fig. 3B, Table-S4)**.

2. Flies with reduced expression of *pirk* or *caudal* show increased mortality despite having lower bacterial loads

In flies, systemic infection with *P. entomophila* results in severe pathology, eventually causing death (N. E. Martins et al. 2013). Induction of AMPs can occur as early as 2 hours after infection (B. Lemaitre, Reichhart, and Hoffmann 1997). *Relish* knockdown flies are known to be short-lived and also exhibit increased bacterial loads after infection. For instance, even a single *E. cloacae* cell

infection is sufficient to kill the fly (Hedengren et al. 1999). Given our observation that knocking
down negative regulators of IMD resulted in an increase in AMP expression, we would expect
mortality in these flies to be lower than controls if variation in mortality was mainly explained by
pathogen clearance. However, while mortality was higher in *relish* and *DiptericinB* knockdowns as
expected, we also observed higher mortality when knocking down the expression of both negative
regulators, *pirk* and *caudal* (**Fig. 4, Table-S5)**, despite the increased expression of AMPs and their
relatively lower bacterial load **(Fig. 3)**. This increased mortality of flies lacking negative regulators
of IMD such as *pirk* and *caudal* has been observed previously (Kleino et al. 2008; Ryu et al. 2008;
Morris et al. 2016) and suggests that mechanisms other than pathogen clearance actively contribute
to the variation in mortality during systemic bacterial infection.

Figure 4: **(A)** Survival curves for male and female *UAS^RNAi* flies (n=16-24 fly
vial/treatment/sex/fly line) (a) *UAS-Relish^RNAi* (b) *UAS-Caudal^RNAi* (c) *UAS-Pirk^RNAi* (d) *UAS-*
DptB^RNAi after systemic infection of *P. entomophila* (dose ~2000 cells/μl). **(B)** hazard ratio
calculated from the survival curves. Asterisks in panel B indicate increased susceptibility of
knockdown flies compared to functional flies (p<0.05).

3.　Knocking down of *caudal* or *pirk* expression in flies results in reduced disease tolerance

By affecting the mortality of flies independently of their microbe loads, negative regulators of IMD
would fit the functional definition of a tolerance mechanisms (Kutzer and Armitage 2016; Ayres
and Schneider 2012; Lissner and Schneider 2018). To quantify the tolerance response of each fly

line more precisely, we measured the tolerance phenotype (measured as fly survival relative to its bacterial load) of male & female flies differing in functional components of IMD-pathway (*relish, caudal, pirk & DptB*). Overall, we found that flies without the negative IMD regulators *caudal* and *pirk* showed reduced tolerance to systemic *P. entomophila* infection compared to flies with fully intact immune system (**Fig. 5, Table-S6**). Flies without the transcription factor *relish* or the AMP *DptB* also showed reduced tolerance, which may reflect feedbacks between increased pathogen growth and decreased ability to limit damage when microbe loads get too high (that is, the per-pathogen damage becomes higher in flies unable to clear infection in the first place) (Råberg 2014).

Figure 5: Disease tolerance in IMD-pathway mutants and fully functional flies. The relationship between host (fly) survival and bacterial load (single infection dose) analysing using point tolerance at ~8 hours post *P. entomophila* infection for male and female flies. Each data represents 'point tolerance', the ratio estimates calculated from median lifespan and average bacterial load for IMD-pathway knockdown and flies with fully intact IMD functioning. Asterisks inside the panel indicates significant difference in disease tolerance between knockdown flies and flies with fully intact IMD functioning (p<0.05).

To further characterise the tolerance response of each IMD fly line across a wide range of pathogen exposure, we infected the IMD-pathway mutants with different doses of *P. entomophila* (ranging from ~650 bacterial cells infection dose to leading up to ~2900 cells). When IMD mutant flies were exposed to very high bacterial infection, all fly lines showed reduced tolerance (**Fig. 6, Table-S6**). We found similar effect for both females and males suggesting that overall tolerance effects

both sexes similarly when exposed to systemic *P. entomophila* bacteria. Altogether, our data show that the negative regulators of IMD pathway play a major role in antibacterial protection and mutant flies without negative regulators of IMD are less tolerant to infection.

Figure 6: Disease tolerance in IMD-pathway mutants and fully functional flies. The relationship between host (fly) survival and bacterial load of 4 infection doses [i.e., (I) ~650 cells (II) ~1400 cells (III) ~2000 cells (IV) ~2900 cells] of *P. entomophila*, analysing using point tolerance at ~8 hours following *P. entomophila* infection for **(A)** male and **(B)** female flies. Each data represents 'point tolerance', the ratio estimates calculated from median lifespan and average bacterial load for IMD-pathway knockdown and flies with fully functional IMD. Asterisks inside the panel indicates significant difference in disease tolerance between knockdown flies and with fully intact IMD functioning (p<0.05).

One likely cause of flies lacking negative IMD regulation still succumb to death despite reducing pathogen burden is immunopathology due to continued immune activation. There is a precedent for the effects of negative regulators on disease tolerance. For example*, G9a*, a negative regulator of Jak/Stat, was identified as being important in tolerating *Drosophila C Virus* infections by reducing immunopathology during the antiviral immunity (Merkling et al. 2015). During bacterial infections, *PGRP-LB* functions as a negative regulator of the *Imd* pathway by regulating the IMD-responsive AMP *Diptericin* (Zaidman-Rémy et al. 2006). More recently it has been shown that *PGRP-LB* mutant flies experience sex-differences in survival despite undetectable differences in their bacterial loads (Vincent and Dionne 2021). Though this study did not directly measure disease tolerance (i.e., survival relative to microbe load), its findings are consistent with the phenotype of

disease tolerance. Here, we have explicitly analysed the tolerance response in the IMD mutants

and found a clear effect of its negative regulators on disease tolerance, though this was comparable

in both males and females. How these negative regulators contribute to disease tolerance is likely

linked to their role in reducing immunopathology. In the case of *caudal* and *pirk*, there is a possible

source of tissue damage through the cytotoxic effects of AMPs, because we observed higher AMP

expression in these mutants. AMPs are known to be harmful to both prokaryotic and eukaryotic

cells and tissues (Song et al. 2005; Paredes-Gamero et al. 2012) and overexpression of AMPs in

flies leads to cytotoxicity, causing neurodegeneration and age-related mortality (Badinloo et al.

2018; Cao et al. 2013). This would support the idea that elevated levels of several AMPs are

sufficient to cause cytotoxicity to cells/tissues, contributing to immunopathological consequences.

Concluding remarks

In the present work, we used a highly virulent fly pathogen (*P. entomophila*) and flies lacking various

components of IMD-signalling. Our data show that flies with reduced expression of the negative

regulators of IMD *pirk* & *caudal* experienced reduced tolerance despite having higher AMP

expression and reducing the bacterial numbers. We hypothesise that this is likely due to

immunopathology arising through the accumulation of cytotoxicity (AMPs and ROS). Our results

highlight that the negative regulators *caudal* & *pirk* are important not only in regulating pathogen

clearance by the IMD pathway, but also play a role in disease tolerance (**Fig. 7**). Uncovering the

role of these regulators in disease tolerance phenotype opens up the possibility of using them as

therapeutic targets where the aim is to enhance host disease tolerance (R. Martins et al. 2019; Vale

et al. 2016; Vale, Fenton, and Brown 2014; Schneider and Ayres 2008). Future work may also

investigate if natural genetic variation in disease tolerance can be explained by variation in negative

immune regulators.

Figure 7: Summary highlighting that in addition to regulating an efficient pathogen clearance response, negative regulators of IMD also contribute to the phenotype of disease tolerance. Flies with reduced expression of the negative regulators *pirk* & *caudal* show reduced tolerance (overall health state) despite having higher AMP expression and reducing the bacterial numbers. Flies with reduced *relish* & *DptB* expression also exhibited reduced tolerance and survival mostly due to pathogen-induced mortality because of lower AMPs to target the bacterial replication.

Acknowledgements:

We thank Jennifer C. Regan, Amy B. Pedersen, Daniel H. Nussey and Ashworth fly group members for helpful discussion, and Fergal M. Waldron for help with *Drosophila* genetics. We thank Angela Reid and Lucinda Rowe for help with media preparation. Figure-1 was created using BioRender.com.

Author contribution:

A.P & P.F.V designed the experiments; A.P carried out experiments; A.P & K.M.M performed qPCR experiments; A.P & P.F.V analysed the data and wrote the manuscript.

Funding:

We acknowledge funding support from a Branco Weiss fellowship and a Chancellor's Fellowship

to P.F.V; a Darwin Trust PhD studentship to A.P. We thank School of Biological Sciences,

University of Edinburgh. United Kingdom.

Competing interests:

None.

References:

Adelman, James S., Laila Kirkpatrick, Jessica L. Grodio, and Dana M. Hawley. 2013. "House
Finch Populations Differ in Early Inflammatory Signaling and Pathogen Tolerance at
the Peak of *Mycoplasma Gallisepticum* Infection." *The American Naturalist* 181 (5):
674–89. https://doi.org/10.1086/670024.
Aggarwal, Kamna, Florentina Rus, Christie Vriesema-Magnuson, Deniz Ertürk-Hasdemir,
Nicholas Paquette, and Neal Silverman. 2008. "Rudra Interrupts Receptor Signaling
Complexes to Negatively Regulate the IMD Pathway." Edited by David S. Schneider.
PLoS Pathogens 4 (8): e1000120. https://doi.org/10.1371/journal.ppat.1000120.
Ayres, Janelle S, and David S Schneider. 2012. "Tolerance of Infections," 26.
Badinloo, Marziyeh, Elizabeth Nguyen, Winston Suh, Faisal Alzahrani, Jovelyn Castellanos,
Vladimir I. Klichko, William C. Orr, and Svetlana N. Radyuk. 2018. "Overexpression
of Antimicrobial Peptides Contributes to Aging through Cytotoxic Effects in
Drosophila Tissues." *Archives of Insect Biochemistry and Physiology* 98 (4): e21464.
https://doi.org/10.1002/arch.21464.
Basbous, Nada, Franck Coste, Philippe Leone, Renaud Vincentelli, Julien Royet, Christine
Kellenberger, and Alain Roussel. 2011. "The *Drosophila* Peptidoglycan-recognition
Protein LF Interacts with Peptidoglycan-recognition Protein LC to Downregulate the
Imd Pathway." *EMBO Reports* 12 (4): 327–33.
https://doi.org/10.1038/embor.2011.19.
Belmonte, Rebecca L., Mary-Kate Corbally, David F. Duneau, and Jennifer C. Regan. 2020.
"Sexual Dimorphisms in Innate Immunity and Responses to Infection in Drosophila
Melanogaster." *Frontiers in Immunology* 10 (January): 3075.
https://doi.org/10.3389/fimmu.2019.03075.
Bunn, Andy, and Mikko Korpela. 2019. "An Introduction to DplR," July, 16.
http://r.meteo.uni.wroc.pl/web/packages/dplR/vignettes/intro-dplR.pdf.
Cao, Yang, Stanislava Chtarbanova, Andrew J. Petersen, and Barry Ganetzky. 2013. "Dnr1
Mutations Cause Neurodegeneration in Drosophila by Activating the Innate Immune
Response in the Brain." *Proceedings of the National Academy of Sciences* 110 (19):
E1752–60. https://doi.org/10.1073/pnas.1306220110.
Cerenius, Lage, Bok Luel Lee, and Kenneth Söderhäll. 2008. "The ProPO-System: Pros and
Cons for Its Role in Invertebrate Immunity." *Trends in Immunology* 29 (6): 263–71.
https://doi.org/10.1016/j.it.2008.02.009.

Cornet, Stéphane, Coraline Bichet, Stephen Larcombe, Bruno Faivre, and Gabriele Sorci. 2014. "Impact of Host Nutritional Status on Infection Dynamics and Parasite Virulence in a Bird-Malaria System." *Journal of Animal Ecology* 83 (1): 256–65. https://doi.org/10.1111/1365-2656.12113.

De Gregorio, Ennio, Paul T.Spellman, Phoebe Tzou, Gerald M. Rubin, and Bruno Lemaitre. 2002. "The Toll and Imd Pathways Are the Major Regulators of the Immune Response in Drosophila." *The EMBO Journal* 21 (11): 2568–79. https://doi.org/10.1093/emboj/21.11.2568.

Dieppois, Guennaelle, Onya Opota, Jorge Lalucat, and Bruno Lemaitre. 2015. "Pseudomonas Entomophila: A Versatile Bacterium with Entomopathogenic Properties." In *Pseudomonas*, edited by Juan-Luis Ramos, Johanna B. Goldberg, and Alain Filloux, 25–49. Dordrecht: Springer Netherlands. https://doi.org/10.1007/978-94-017-9555-5_2.

Dionne, Marc S., Linh N. Pham, Mimi Shirasu-Hiza, and David S. Schneider. 2006. "Akt and Foxo Dysregulation Contribute to Infection-Induced Wasting in Drosophila." *Current Biology* 16 (20): 1977–85. https://doi.org/10.1016/j.cub.2006.08.052.

Graham, Andrea L., Judith E. Allen, and Andrew F. Read. 2005. "Evolutionary Causes and Consequences of Immunopathology." *Annual Review of Ecology, Evolution, and Systematics* 36 (1): 373–97. https://doi.org/10.1146/annurev.ecolsys.36.102003.152622.

Gupta, Vanika, and Pedro F. Vale. 2017. "Nonlinear Disease Tolerance Curves Reveal Distinct Components of Host Responses to Viral Infection." *Royal Society Open Science* 4 (7): 170342. https://doi.org/10.1098/rsos.170342.

Han, Jiahuai, and Richard J Ulevitch. 2005. "Limiting Inflammatory Responses during Activation of Innate Immunity." *Nature Immunology* 6 (12): 1198–1205. https://doi.org/10.1038/ni1274.

Hedengren, Marika, BengtÅsling, Mitchell S Dushay, Istvan Ando, Sophia Ekengren, Margareta Wihlborg, and Dan Hultmark. 1999. "Relish, a Central Factor in the Control of Humoral but Not Cellular Immunity in Drosophila." *Molecular Cell* 4 (5): 827–37. https://doi.org/10.1016/S1097-2765(00)80392-5.

Hill, Adrian V. S. 1998. "The Immunogenetics of Human Infectious Diseases." *Annual Review of Immunology* 16 (1): 593–617. https://doi.org/10.1146/annurev.immunol.16.1.593.

Howick, Virginia M., and Brian P. Lazzaro. 2014. "Genotype and Diet Shape Resistance and Tolerance across Distinct Phases of Bacterial Infection." *BMC Evolutionary Biology* 14 (1): 56. https://doi.org/10.1186/1471-2148-14-56.

Khalil, Sarah, Eliana Jacobson, Moria C. Chambers, and Brian P. Lazzaro. 2015. "Systemic Bacterial Infection and Immune Defense Phenotypes in Drosophila Melanogaster." *Journal of Visualized Experiments : JoVE*, no. 99 (May): 52613. https://doi.org/10.3791/52613.

Khan, Imroze, Deepa Agashe, and Jens Rolff. 2017. "Early-Life Inflammation, Immune Response and Ageing." *Proceedings of the Royal Society B: Biological Sciences* 284 (1850): 20170125. https://doi.org/10.1098/rspb.2017.0125.

Klein, Sabra L., and Katie L. Flanagan. 2016. "Sex Differences in Immune Responses." *Nature Reviews Immunology* 16 (10): 626–38. https://doi.org/10.1038/nri.2016.90.

Kleino, Anni, Henna Myllymäki, Jenni Kallio, Leena-Maija Vanha-aho, Kaisa Oksanen, Johanna Ulvila, Dan Hultmark, Susanna Valanne, and Mika Rämet. 2008. "Pirk Is a Negative Regulator of the *Drosophila* Imd Pathway." *The Journal of Immunology* 180 (8): 5413–22. https://doi.org/10.4049/jimmunol.180.8.5413.

Kleino, Anni, and Neal Silverman. 2014. "The Drosophila IMD Pathway in the Activation of the Humoral Immune Response." *Developmental & Comparative Immunology* 42 (1): 25–35. https://doi.org/10.1016/j.dci.2013.05.014.

Knutie, Sarah A., Christina L. Wilkinson, Qiu Chang Wu, C. Nicole Ortega, and Jason R. Rohr. 2017. "Host Resistance and Tolerance of Parasitic Gut Worms Depend on Resource Availability." *Oecologia* 183 (4): 1031–40. https://doi.org/10.1007/s00442-017-3822-7.

Kuraishi, Takayuki, Aki Hori, and Shoichiro Kurata. 2013. "Host-Microbe Interactions in the Gut of Drosophila Melanogaster." *Frontiers in Physiology* 4. https://doi.org/10.3389/fphys.2013.00375.

Kutzer, Megan A.M., and Sophie A.O. Armitage. 2016. "Maximising Fitness in the Face of Parasites: A Review of Host Tolerance." *Zoology* 119 (4): 281–89. https://doi.org/10.1016/j.zool.2016.05.011.

Lambeth, J. David. 2007. "Nox Enzymes, ROS, and Chronic Disease: An Example of Antagonistic Pleiotropy." *Free Radical Biology and Medicine* 43 (3): 332–47. https://doi.org/10.1016/j.freeradbiomed.2007.03.027.

Lazzaro, Brian P., Bonnielin K. Sceurman, and Andrew G. Clark. 2004. "Genetic Basis of Natural Variation in D. Melanogaster Antibacterial Immunity." *Science* 303 (5665): 1873–76. https://doi.org/10.1126/science.1092447.

Lee, Kwang-Zin, and Dominique Ferrandon. 2011. "Negative Regulation of Immune Responses on the Fly: Immune Responses on the Fly." *The EMBO Journal* 30 (6): 988–90. https://doi.org/10.1038/emboj.2011.47.

Lemaitre, B., J.-M. Reichhart, and J. A. Hoffmann. 1997. "Drosophila Host Defense: Differential Induction of Antimicrobial Peptide Genes after Infection by Various Classes of Microorganisms." *Proceedings of the National Academy of Sciences* 94 (26): 14614–19. https://doi.org/10.1073/pnas.94.26.14614.

Lemaitre, Bruno, and Jules Hoffmann. 2007. "The Host Defense of *Drosophila Melanogaster*." *Annual Review of Immunology* 25 (1): 697–743. https://doi.org/10.1146/annurev.immunol.25.022106.141615.

Leulier, François, Claudine Parquet, Sebastien Pili-Floury, Ji-Hwan Ryu, Martine Caroff, Won-Jae Lee, Dominique Mengin-Lecreulx, and Bruno Lemaitre. 2003. "The Drosophila Immune System Detects Bacteria through Specific Peptidoglycan Recognition." *Nature Immunology* 4 (5): 478–84. https://doi.org/10.1038/ni922.

Lissner, Michelle M, and David S Schneider. 2018. "The Physiological Basis of Disease Tolerance in Insects." *Current Opinion in Insect Science*, Global change biology * Molecular physiology, 29 (October): 133–36. https://doi.org/10.1016/j.cois.2018.09.004.

Little, Tom J., David M. Shuker, Nick Colegrave, Troy Day, and Andrea L. Graham. 2010. "The Coevolution of Virulence: Tolerance in Perspective." Edited by Marianne Manchester. *PLoS Pathogens* 6 (9): e1001006. https://doi.org/10.1371/journal.ppat.1001006.

Livak, Kenneth J., and Thomas D. Schmittgen. 2001. "Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2−ΔΔCT Method." *Methods* 25 (4): 402–8. https://doi.org/10.1006/meth.2001.1262.

Loch, Gerrit, Ingo Zinke, Tetsushi Mori, Pilar Carrera, Jonas Schroer, Haruko Takeyama, and Michael Hoch. 2017. "Antimicrobial Peptides Extend Lifespan in Drosophila." Edited by Gregg Roman. *PLOS ONE* 12 (5): e0176689. https://doi.org/10.1371/journal.pone.0176689.

Louie, Alexander, Kyung Han Song, Alejandra Hotson, Ann Thomas Tate, and David S. Schneider. 2016. "How Many Parameters Does It Take to Describe Disease

Tolerance?" Edited by Andy P. Dobson. *PLOS Biology* 14 (4): e1002435. https://doi.org/10.1371/journal.pbio.1002435.

Malo, Danielle, and Emil Skamene. 1994. "Genetic Control of Host Resistance to Infection." *Trends in Genetics* 10 (10): 365–71. https://doi.org/10.1016/0168-9525(94)90133-3.

Martins, Nelson E., Vitor G. Faria, Luis Teixeira, Sara Magalhães, and Élio Sucena. 2013. "Host Adaptation Is Contingent upon the Infection Route Taken by Pathogens." *PLOS Pathogens* 9 (9): e1003601. https://doi.org/10.1371/journal.ppat.1003601.

Martins, Rui, Ana Rita Carlos, Faouzi Braza, Jessica A. Thompson, Patricia Bastos-Amador, Susana Ramos, and Miguel P. Soares. 2019. "Disease Tolerance as an Inherent Component of Immunity." *Annual Review of Immunology* 37 (1): 405–37. https://doi.org/10.1146/annurev-immunol-042718-041739.

Medzhitov, Ruslan, David S. Schneider, and Miguel P. Soares. 2012. "Disease Tolerance as a Defense Strategy." *Science* 335 (6071): 936–41. https://doi.org/10.1126/science.1214935.

Merkling, Sarah H., Alfred W. Bronkhorst, Jamie M. Kramer, Gijs J. Overheul, Annette Schenck, and Ronald P. Van Rij. 2015. "The Epigenetic Regulator G9a Mediates Tolerance to RNA Virus Infection in Drosophila." *PLOS Pathogens* 11 (4): e1004692. https://doi.org/10.1371/journal.ppat.1004692.

Morris, Otto, Xi Liu, Celia Domingues, Christopher Runchel, Andrea Chai, Shaherin Basith, Tencho Tenev, et al. 2016. "Signal Integration by the IκB Protein Pickle Shapes Drosophila Innate Host Defense." *Cell Host & Microbe* 20 (3): 283–95. https://doi.org/10.1016/j.chom.2016.08.003.

Myllymäki, Henna, Susanna Valanne, and Mika Rämet. 2014. "The Drosophila Imd Signaling Pathway." *The Journal of Immunology* 192 (8): 3455–62. https://doi.org/10.4049/jimmunol.1303309.

Oliveira, José Henrique, Ana Cristina Bahia, and Pedro F. Vale. 2020. "How Are Arbovirus Vectors Able to Tolerate Infection?" *Developmental & Comparative Immunology* 103 (February): 103514. https://doi.org/10.1016/j.dci.2019.103514.

Paredes, Juan C., David P. Welchman, Mickaël Poidevin, and Bruno Lemaitre. 2011. "Negative Regulation by Amidase PGRPs Shapes the Drosophila Antibacterial Response and Protects the Fly from Innocuous Infection." *Immunity* 35 (5): 770–79. https://doi.org/10.1016/j.immuni.2011.09.018.

Paredes-Gamero, Edgar J., Marta N. C. Martins, Fábio A. M. Cappabianco, Jaime S. Ide, and Antonio Miranda. 2012. "Characterization of Dual Effects Induced by Antimicrobial Peptides: Regulated Cell Death or Membrane Disruption." *Biochimica et Biophysica Acta (BBA) - General Subjects* 1820 (7): 1062–72. https://doi.org/10.1016/j.bbagen.2012.02.015.

Raberg, L., D. Sim, and A. F. Read. 2007. "Disentangling Genetic Variation for Resistance and Tolerance to Infectious Diseases in Animals." *Science* 318 (5851): 812–14. https://doi.org/10.1126/science.1148526.

Råberg, Lars. 2014. "How to Live with the Enemy: Understanding Tolerance to Parasites." *PLOS Biology* 12 (11): e1001989. https://doi.org/10.1371/journal.pbio.1001989.

Råberg, Lars, Andrea L Graham, and Andrew F Read. 2009. "Decomposing Health: Tolerance and Resistance to Parasites in Animals." *Philosophical Transactions of the Royal Society B: Biological Sciences* 364 (1513): 37–49. https://doi.org/10.1098/rstb.2008.0184.

Ryu, J.-H., S.-H. Kim, H.-Y. Lee, J. Y. Bai, Y.-D. Nam, J.-W. Bae, D. G. Lee, S. C. Shin, E.-M. Ha, and W.-J. Lee. 2008. "Innate Immune Homeostasis by the Homeobox Gene Caudal and Commensal-Gut Mutualism in Drosophila." *Science* 319 (5864): 777–82. https://doi.org/10.1126/science.1149357.

Sadd, Ben M, and Michael T Siva-Jothy. 2006. "Self-Harm Caused by an Insect's Innate Immunity." *Proceedings of the Royal Society B: Biological Sciences* 273 (1600): 2571–74. https://doi.org/10.1098/rspb.2006.3574.

Schneider, David S. 2007. "How and Why Does a Fly Turn Its Immune System Off?" *PLoS Biology* 5 (9): e247. https://doi.org/10.1371/journal.pbio.0050247.

Schneider, David S., and Janelle S. Ayres. 2008. "Two Ways to Survive Infection: What Resistance and Tolerance Can Teach Us about Treating Infectious Diseases." *Nature Reviews. Immunology* 8 (11): 889–95. https://doi.org/10.1038/nri2432.

Sears, Brittany F., Jason R. Rohr, Judith E. Allen, and Lynn B. Martin. 2011. "The Economy of Inflammation: When Is Less More?" *Trends in Parasitology* 27 (9): 382–87. https://doi.org/10.1016/j.pt.2011.05.004.

Siva-Jothy, Jonathon A., Arun Prakash, Radhakrishnan B. Vasanthakrishnan, Katy M. Monteith, and Pedro F. Vale. 2018. "Oral Bacterial Infection and Shedding in Drosophila Melanogaster." *Journal of Visualized Experiments*, no. 135 (May): 57676. https://doi.org/10.3791/57676.

Soares, Miguel P., Raffaella Gozzelino, and Sebastian Weis. 2014. "Tissue Damage Control in Disease Tolerance." *Trends in Immunology* 35 (10): 483–94. https://doi.org/10.1016/j.it.2014.08.001.

Soares, Miguel P., Luis Teixeira, and Luis F. Moita. 2017. "Disease Tolerance and Immunity in Host Protection against Infection." *Nature Reviews Immunology* 17 (2): 83–96. https://doi.org/10.1038/nri.2016.136.

Song, Yun Mi, Yoonkyung Park, Shin Saeng Lim, Sung-Tae Yang, Eun-Rhan Woo, Il-Seon Park, Jung Sup Lee, et al. 2005. "Cell Selectivity and Mechanism of Action of Antimicrobial Model Peptides Containing Peptoid Residues." *Biochemistry* 44 (36): 12094–106. https://doi.org/10.1021/bi050765p.

Therneau, Terry. 2015. "Mixed Effects Cox Models." In *Mixed Effects Cox Models*. CRAN repository. https://repo.bppt.go.id/cran/web/packages/coxme/vignettes/coxme.pdf.

Troha, Katia, and Nicolas Buchon. 2019. "Methods for the Study of Innate Immunity in Drosophila Melanogaster." *WIREs Developmental Biology* 8 (5): e344. https://doi.org/10.1002/wdev.344.

Troha, Katia, Joo Hyun Im, Jonathan Revah, Brian P. Lazzaro, and Nicolas Buchon. 2018. "Comparative Transcriptomics Reveals CrebA as a Novel Regulator of Infection Tolerance in D. Melanogaster." *PLOS Pathogens* 14 (2): e1006847. https://doi.org/10.1371/journal.ppat.1006847.

Vale, Pedro F., Andy Fenton, and Sam P. Brown. 2014. "Limiting Damage during Infection: Lessons from Infection Tolerance for Novel Therapeutics." *PLoS Biology* 12 (1): e1001769. https://doi.org/10.1371/journal.pbio.1001769.

Vale, Pedro F., Luke McNally, Andrea Doeschl-Wilson, Kayla C. King, Roman Popat, Maria R. Domingo-Sananes, Judith E. Allen, Miguel P. Soares, and Rolf Kümmerli. 2016. "Beyond Killing: Can We Find New Ways to Manage Infection?" *Evolution, Medicine, and Public Health* 2016 (1): 148–57. https://doi.org/10.1093/emph/eow012.

Vincent, Crystal M., and Marc S. Dionne. 2021. "Disparate Regulation of IMD Signaling Drives Sex Differences in Infection Pathology in *Drosophila Melanogaster*." *Proceedings of the National Academy of Sciences* 118 (32): e2026554118. https://doi.org/10.1073/pnas.2026554118.

Vodovar, N., M. Vinals, P. Liehl, A. Basset, J. Degrouard, P. Spellman, F. Boccard, and B. Lemaitre. 2005. "Drosophila Host Defense after Oral Infection by an Entomopathogenic Pseudomonas Species." *Proceedings of the National Academy of Sciences* 102 (32): 11414–19. https://doi.org/10.1073/pnas.0502240102.

Wicker, C., J. M. Reichhart, D. Hoffmann, D. Hultmark, C. Samakovlis, and J. A. Hoffmann.
1990. "Insect Immunity. Characterization of a Drosophila CDNA Encoding a Novel
Member of the Diptericin Family of Immune Peptides." *Journal of Biological*
Chemistry 265 (36): 22493–98. https://doi.org/10.1016/S0021-9258(18)45732-8.
Wickham, Hadley. 2016. "R for Data Science: Import, Tidy, Transform, Visualize, and
Model Data." O'Reilly Media, Inc. https://mran.microsoft.com/snapshot/2015-02-
16/web/packages/ggplot2/ggplot2.pdf.
Zaidman-Rémy, Anna, Mireille Hervé, Mickael Poidevin, Sébastien Pili-Floury, Min-Sung
Kim, Didier Blanot, Byung-Ha Oh, Ryu Ueda, Dominique Mengin-Lecreulx, and
Bruno Lemaitre. 2006. "The Drosophila Amidase PGRP-LB Modulates the Immune
Response to Bacterial Infection." *Immunity* 24 (4): 463–73.
https://doi.org/10.1016/j.immuni.2006.02.012.
Zerofsky, Melissa, Ephat Harel, Neal Silverman, and Marc Tatar. 2005. "Aging of the Innate
Immune Response in Drosophila Melanogaster." *Aging Cell* 4 (2): 103–8.
https://doi.org/10.1111/j.1474-9728.2005.00147.x.

Supplementary information

Figure S1. Survival curves of female flies of different IMD-pathway mutants (n=16-24 flies/treatment/sex/fly line) **(a)** *UAS-Relish*[RNAi] **(b)** *UAS-Caudal*[RNAi] **(c)** *UAS-Pirk*[RNAi] **(d)** *UAS-DptB*[RNAi] during 4 concentrations of systemic *Pseudomonas entomophila* infection ranging between (1) ~650 cells (2) ~1400 cells (3) ~2000 cells (4) ~2900 cells/µl of *P. entomophila* culture. Since all 4 infection concentrations were performed simultaneously for each of the fly line, we used a single mock-controls (SI) across all 4 different infection doses.

Figure S2. Survival curves of male flies of different IMD-pathway mutants (n=16-24
flies/treatment/sex/fly line) **(a)** *UAS-Relish*[RNAi] **(b)** *UAS-Caudal*[RNAi] **(c)** *UAS-Pirk*[RNAi] **(d)** *UAS-*
DptB[RNAi] during 4 concentrations of systemic *Pseudomonas entomophila* infection ranging between (1)
~650 cells (2) ~1400 cells (3) ~2000 cells (4) ~2900 cells/μl of *P. entomophila* culture. Since all 4
infection concentrations were performed simultaneously for each of the fly line, we used a single
mock-controls (SI) across all 4 different infection doses.

Figure S3. The estimated hazard ratios calculated from survival curves (**Fig. S1 & S2**) for flies with fully functional immune system & IMD-pathway mutants for both Sexes. The different IMD-pathway mutants include **(a)** *UAS-Relish*[RNAi] **(b)** *UAS-Caudal*[RNAi] **(c)** *UAS-Pirk*[RNAi] **(d)** *UAS-DptB*[RNAi]. We used 4 different concentrations of systemic *P. entomophila* infection. A greater hazard ratio >1 shown as horizontal dashed grey lines indicates higher susceptibility of IMD-mutant flies than the functional flies to *P. entomophila* infection.

672 **Figure S4.** Bacterial loads at ~24 hours after different concentrations of systemic *P. entomophila*
673 infection for **(A)** female and **(B)** male flies for IMD-pathway mutants **(i)** *UAS-Relish*[RNAi] **(ii)**
674 *UAS-Caudal*[RNAi] **(iii)** *UAS-Pirk*[RNAi] **(iv)** *UAS-DptB*[RNAi].

675
676

677 **Table S1:** Fly strains used in this study
678

Fly line	Bloomington stock #	Genotype
UAS-*relish*RNAi	33661	*y[1]sc[*]v[1]; P{y[+t7.7]v[+t1.8]=TRiP.HMS00070}attP2*
UAS-*caudal*RNAi	57546	*y[1]sc[*]v[1];P{y[+t7.7] v[+t1.8]=TRiP.HMC04863}attP40),*
UAS-*pirk*RNAi	67011	*y[1]sc[*]v[1];P{y[+t7.7] v[+t1.8]=TRiP.HMS05477}attP40*
UAS-*Diptericin*BRNAi	28975	*y[1]v[1];P{y[+t7.7] v[+t1.8]=TRiP.HM05186}attP2*
(UASRNAi) ubiquitous driver	67065	*w*;P{UAS-3xFLAG.dCas9.VPR}attP40, P{tubP-GAL80ts}10; P{tubP-GAL4}LL7/TM6B, Tb1*

679
680
681 **Table S2**: Primers used in this study
682

	Obtained from	AMPs	Sequence
1.	(Troha et al. 2018)	*Dpt*_Forward:	5' GACGCCACGAGATTGGACTG 3'
		*Dpt*_Rverese	5' CCCACTTTCCAGCTCGGTTC 3'
		*AttC*_Forward:	5' TGCCCGATTGGACCTAAGC 3'
		*AttC*_Reverse:	5' GCGTATGGGTTTTGGTCAGTTC 3'
2.	(Loch et al. 2017)	*CecA1*_Forward	5' TCTTCGTTTTCGTCGCTCTCA 3',
		*CecA1*_Reverse	5' ATTCCCAGTCCCTGGATT GTG 3'.
3.	(Gupta and Vale 2017)	*Rp49*_Forward:	5' ATGCTAAGCTGTCGCACAAATG 3'
		*Rp49*_Reverse:	5' GTTCGATCCGTAACCGATGT 3'

Genotype	Source	df	Sum Sq	F value	p
UAS-Relish^RNAi	Knockdown *(knockdown & functional flies)*	1	1160.3	148.38	<0.001
	AMPs *(major IMD-responsive AMPs)*	2	11.2	0.717	0.49
	Sex	1	0.4	0.051	0.82
	Knockdown x AMPs	2	49.1	3.139	0.05
	Knockdown x Sex	1	0.3	0.033	0.85
	AMPs x Sex	2	1.4	0.087	0.91
	Knockdown x AMPs x Sex	2	11.7	0.748	0.47
UAS-Caudal^RNAi	Knockdown	1	460.4	159.2	<0.001
	AMPs	2	10.6	1.835	0.17
	Sex	1	23.5	8.137	<0.001
	Knockdown x AMPs	2	82.3	14.24	<0.001
	Knockdown x Sex	1	1.1	0.394	0.53
	AMPs x Sex	2	4.4	0.770	0.46
	Knockdown x AMPs x Sex	2	25.2	4.365	<0.001
UAS-Pirk^RNAi	Knockdown	1	538.3	338.9	<0.001
	AMPs	2	34.2	10.77	<0.001
	Sex	1	0.0	0.000	0.99
	Knockdown x AMPs	2	47.1	14.82	<0.001
	Knockdown x Sex	1	8.5	5.360	<0.001
	AMPs x Sex	2	35.6	11.22	<0.001
	Knockdown x AMPs x Sex	2	7.2	2.255	0.11

Genotype	Source	df	Sum Sq	F value	p
UAS-Relish^RNAi	Knockdown	3	351.07	45.4374	<0.001
	Sex	1	39.32	15.2654	<0.001
	Dose	3	518.53	67.1097	<0.001
	Knockdown x Sex	1	15.53	6.0297	<0.001
	Knockdown x dose	3	14.43	1.8681	0.13
	Sex x dose	3	1.64	0.2120	0.88
	Knockdown x Sex x dose	1	3.72	1.4450	0.23
	Random effect	*Std err*			
	Vials	*1.605*			
UAS-Caudal^RNAi	Knockdown	3	157.03	20.6574	<0.001
	Sex	1	115.72	45.6679	<0.001
	Dose	3	528.70	69.5504	<0.001
	Knockdown x Sex	1	0.38	0.1497	0.69
	Knockdown x dose	3	27.01	3.5529	0.014
	Sex x dose	3	3.55	0.4665	0.70
	Knockdown x Sex x dose	1	1.10	0.4360	0.50
	Random effect	*Std err*			
	vials	*1.592*			
UAS-Pirk^RNAi	Knockdown	3	164.02	18.3271	<0.001
	Sex	1	98.90	33.1544	<0.001
	Dose	3	407.46	45.5296	<0.001
	Knockdown x Sex	1	2.57	0.8601	0.35

	Source				p
	Knockdown x dose	3	3.17	0.3541	0.78
	Sex x dose	3	2.17	0.2428	0.86
	Knockdown x Sex x dose	1	0.06	0.0215	0.88
	Random effect	*Std err*			
	vials	*1.727*			
UAS-DptB^RNAi	Knockdown	3	128.94	13.2207	<0.001
	Sex	1	87.43	26.8915	<0.001
	Dose	3	615.60	63.1179	<0.001
	Knockdown x Sex	1	0.44	0.1354	0.71
	Knockdown x dose	3	27.20	2.7885	0.04
	Sex x dose	3	7.48	0.7674	0.51
	Knockdown x Sex x dose	1	1.43	0.4411	0.50
	Random effect	*Std err*			
	vials	*1.803*			

Table S5: Summary of mixed effects Cox model, fitting the model to estimate response to *P. entomophila* infection in flies with fully intact immune system and flies with deficiencies in IMD-pathway. We used data from the individuals of ~3-day adult males and females infected with 4 concentration (doses) of *P. entomophila* for each fly lines (IMD-pathway mutants) and specified the model as: survival ~ Knockdown * Sex * dose (1|vial), with 'Knockdown' and 'Sex' as fixed effects, and 'vials' as a random effect. The table shows model output (ANOVA) for survival post-infection for flies with fully function immune system and defects in IMD functioning.

Genotype	*Source*	*Loglik*	*Chisq*	*df*	*p*
UAS-Relish^RNAi	Knockdown	1344.5	33.429	1	<0.001
	Sex	1344.5	0.0003	1	0.98
	Dose	1273.7	141.55	4	<0.001
	Knockdown x Sex	1273.5	0.4192	1	0.51
	Knockdown x dose	1270.0	7.0930	4	0.13
	Sex x dose	1269.5	0.9557	4	0.91
	Knockdown x Sex x dose	1266.8	5.4275	4	0.24
	Random effect	*Std dev*			
	Vials	*0.244*			
UAS-Caudal^RNAi	Knockdown	1337.6	28.624	1	<0.001
	Sex	1337.5	0.0156	1	0.9
	Dose	1285.3	104.43	4	<0.001
	Knockdown x Sex	1285.3	0.0005	1	0.98
	Knockdown x dose	1283.6	3.4040	4	0.49
	Sex x dose	1280.6	6.0031	4	0.19
	Knockdown x Sex x dose	1279.2	2.8850	4	0.57
	Random effect	*Std dev*			
	vials	*0.02*			
UAS-Pirk^RNAi	Knockdown	1412.1	21.126	1	<0.001
	Sex	1412.1	0.0001	1	0.99
	Dose	1357.4	109.30	4	<0.001
	Knockdown x Sex	1356.6	1.6275	1	0.20
	Knockdown x dose	1355.3	2.5330	4	0.63
	Sex x dose	1353.9	2.9391	4	0.56
	Knockdown x Sex x dose	1352.3	3.2265	4	0.52
	Random effect	*Std dev*			
	Vials	*0.14*			
UAS-DptB^RNAi	Knockdown	1386.1	29.146	1	<0.001
	Sex	1386.0	0.2848	1	0.59
	Dose	1331.1	109.81	4	<0.001
	Knockdown x Sex	1331.0	0.1187	1	0.73
	Knockdown x dose	1330.8	0.3249	4	0.98
	Sex x dose	1329.1	3.5419	4	0.47
	Knockdown x Sex x dose	1328.0	2.1198	4	0.71
	Random effect	*Std dev*			

	Vials			0.15	

Table S6: 'Point tolerance' for all doses - Summary of ANCOVA. To assess differences bacterial loads following infection with 4 concentrations (dose) of *P. entomophila*, we analysed the average bacterial load, measured at ~8 hours following infection using ANCOVA and fitted 'Knockdown' and 'Sex' as categorical fixed effects and 'bacterial load' as continuous co-variate for each fly line.

Genotype	Source	df	F ratio	p
UAS-Relish^{RNAi}	Knockdown	1	17.584	0.003
	Sex	1	2.2509	0.17
	Dose	3	5.2387	0.02
	Bacterial load	1	0.0728	0.79
	Knockdown * Bac. load	1	13.549	0.006
UAS-Caudal^{RNAi}	Knockdown	1	9.1734	0.01
	Sex	1	0.5894	0.46
	Dose	3	3.7512	0.059
	Bacterial load	1	0.0386	0.84
	Knockdown * Bac. load	1	5.9732	0.04
UAS-Pirk^{RNAi}	Knockdown	1	25.071	0.001
	Sex	1	0.9570	0.35
	Dose	3	2.8131	0.10
	Bacterial load	1	8.6071	0.01
	Knockdown * Bac. load	1	0.1030	0.75
UAS-DptB^{RNAi}	Knockdown	1	25.486	0.001
	Sex	1	0.0266	0.87
	Dose	3	10.069	0.004
	Bacterial load	1	4.2575	0.07
	Knockdown * Bac. load	1	17.282	0.003