1
Negative immune regulation contributes to disease tolerance in Drosophila

Arun Prakash*, Katy M. Monteith, Pedro F. Vale*

Institute of Evolutionary Biology

School of Biological Sciences

University of Edinburgh

Edinburgh EH9 3FL

*Correspondence

arun.prakash@ed.ac.uk

pedro.vale@ed.ac.uk

Keywords: Disease tolerance, Immune deficiency (IMD) pathway, Immunopathology, Negative
immune regulators, Pathogen clearance

Abstract

Disease tolerance is an infection phenotype where hosts show relatively high health despite
harbouring elevated pathogen loads. Variation in the ability to reduce immunopathology may explain
why some hosts can tolerate higher pathogen burdens with reduced pathology. Negative immune
regulation would therefore appear to be a clear candidate for a mechanism underlying disease
tolerance. Here, we examined how the negative regulation of the immune deficiency (IMD) pathway
affects disease tolerance in *Drosophila melanogaster* when infected with four doses of the gram-
negative bacterial pathogen *Pseudomonas entomophila*. We find that while flies unable to regulate
the IMD response exhibited higher expression of antimicrobial peptides and lower bacterial loads as
expected, this was not accompanied by a proportional reduction in mortality. Instead, UAS-RNAi
knockdown of negative regulators of IMD (*pirk* and *caudal*) substantially increased the per-pathogen-
mortality in both males and females across all tested infectious doses. Our results therefore highlight
that in addition to regulating an efficient pathogen clearance response, negative regulators of IMD
also contribute to disease tolerance.

Keywords: Immune deficiency (IMD) pathway, immune effectors, disease tolerance,
immunopathology, negative immune regulators

Introduction

To survive infection hosts must achieve a fine balance between clearing pathogens and avoiding immune-induced pathology by tightly regulating the induction and resolution of immune responses [1–3]. Understanding these distinct pathogen- or host-derived causes of tissue damage is important not only from a therapeutic perspective [4–6], but also offers important mechanistic explanations for the variable infection phenotypes that are often observed between individuals [7–9]. Individuals with low pathogen loads and relatively high health during infection are commonly described as 'resistant' because they are able to clear - or otherwise quantitatively reduce - the establishment and growth of pathogens within the host [7,8,10,11]. Other infection phenotypes are more difficult to explain as the consequence of resistance, such as the observation that some individuals exhibit low levels of pathology (measured as either mortality or physiological performance), despite harbouring relatively high pathogen loads [8,9,12,13].

Instead, tolerating higher pathogen loads is more likely to result from mechanisms that prevent, limit or repair tissue damage [14–17]. Previous work in *Drosophila* uncovered several candidate genes associated with disease tolerance, including many that do not have an obvious immune related function such as *CrebA* [18], *grainyhead and debris buster* [19], *dFOXO* [20,21]. Curiously, genes related to canonical immune pathways are conspicuously absent from these genome-wide screens of tolerance, even though inflammation is often an underlying cause of damage during infection, and we might therefore expect the regulation of inflammation to be a key mechanism underlying disease tolerance [2,8,13,22].

Other work, however, suggests that minimization of inflammation may be key to understanding tolerance. For example, pro-inflammatory cytokines tend to be associated with decreased tolerance in populations of house finches (*Haemorhous mexicanus*) infected with a bacterial pathogen *Mycoplasma gallisepticum,* which exhibit reduced cytokine expression compared with a less tolerant population [23]. Lower levels of circulating pro-inflammatory cytokines are also associated with increased tolerance of malaria [24]. In mice, treating *Mycobacterium tuberculosis* infection with the anti-inflammatory drug Ibuprofen resulted in marked improvements in infection tolerance [25], while negative regulation of inflammation-like responses permit greater tolerance of Drosophila C virus infection in fruit flies [22,26,27]. Negative regulation of immune responses would therefore appear to be a promising candidate for a general a mechanism that promotes disease tolerance [16,24,28,29].

In the present work, we investigate if the negative regulation of immune responses has any measurable effect on the phenotype of disease tolerance during systemic bacterial infection in Drosophila. We focused on the regulation of the immune deficiency (IMD) pathway (Fig. S1), one the best described immune signalling pathways in *Drosophila* [30–32]. Following infection with gram-negative bacteria, DAP-type (Diaminopimelic acid) peptidoglycans are recognised by pathogen-sensing receptors *PGRPs (LC* and *LE)* [33,34]. These receptors then recruit the adaptor molecule

IMD and activate an intracellular signalling cascade that leads to the activation and translocation to
the nucleus of a *NF-κB relish,* inducing the transcription of a set of IMD-responsive antimicrobial
peptides (AMPs)[32]. *Relish* also induces the transcription of negative regulators of IMD (*pirk*,
caudal, PGRPs-LB, LC, LE,) which ensures an appropriate level of immune response, while avoiding
immunopathology [35,36]. *Pirk*, for example, interferes with the interaction of *PGRP-LC* and *-LE* with
the molecule IMD, and limits the activation of the IMD pathway [37–39]. Similarly, the homeobox
protein *caudal* downregulates the expression of AMPs in flies [40] (Fig. S1). Taken together, the
negative regulation of IMD-pathway ensures an appropriate level of immune response and enables
individuals to reduce immune activity in response to reductions in bacterial numbers [35,38,41].

We tested whether disrupting optimal regulation of the IMD pathway can impact the ability of
Drosophila to tolerate bacterial infection. We measured disease tolerance during systemic infection
with the gram-negative bacterium *Pseudomonas entomophila* using *Drosophila* lines where the
expression of specific components of the IMD-pathway were knocked down using UAS-RNAi,
including the transcription factor- *relish*, its negative regulators *pirk* and *caudal*, and a major IMD-
responsive AMP *Diptericin-B*.

Materials and methods

Experimental methods. To test the role of IMD negative regulation on systemic bacterial infection,
we used fly lines where the UAS/Gal4 system was used for ubiquitous downregulation of target
genes with RNA interference (RNAi): UAS-RNAi-*relish*, UAS-RNAi-*caudal* UAS-RNAi-*pirk* and UAS-
RNAi-*diptericinB* (see Supplementary Methods and Table S1 for details about fly genotypes). We
infected these flies with four inoculation doses of the gram-negative bacterium *Pseudomonas*
entomophila [34] OD_{600} = 0.1, 0.05, 0.01 and 0.005*,* delivering approximately I) ~650 cells (II) ~1400
cells (III) ~2000 cells (IV) ~2900 cells, respectively). Following infection, we quantified the
expression of three IMD-responsive AMPs namely, diptericin (Dpt), attacin-C AttC), cecropin-A1
(CecA1), by quantitative real-time polymerase chain reaction (qRT-PCR) and quantified the effects
of downregulating each gene of interest on the mortality of each fly and the internal bacterial loads
achieved during the infection. A detailed description of fly rearing conditions and bacterial culture
protocols, infection protocols, and the measurement of survival and bacterial loads is given in
Supplementary Methods

Statistical analysis

All data and code are available at. https://doi.org/10.5281/zenodo.5564519.

Gene expression: We analysed the gene expression data by first calculating the ΔCT value [42] for
the expression of a gene of interest relative to the reference house-keeping gene rp49 [26]. We
analysed, using ANOVA, the effects of the experimental treatment on the ΔCT values, indicating the
difference in AMPs expression between knockdown and fully functional flies of each IMD-pathway fly
line. We used the R package 'ggplot2' for graphics and data visualization (Wickham 2016; Bunn and
Korpela, 2019).

Bacterial load: We quantified differences in bacterial load following bacterial infection by analysing
the bacterial measurement ($\log_{10}$) 8 hours following *P. entomophila* infection. We found that
residuals of bacterial load data were normally distributed when tested with Shapiro-Wilks test and
hence we used ANOVA and analysed the bacterial load data by fitting 'knockdown' (functional flies
and IMD-pathway knockdown flies), 'sex' and 'infection dose' (4 different infectious doses of *P.*
entomophila) as categorical fixed-effects, while 'vials' (replicates) as a random-effects for each of the
fly lines (functional flies and IMD-pathway knockdown flies).

1. *Survival post-infection*: We analysed the survival data after *P. entomophila* bacterial infection for all
the IMD-pathway mutants using a mixed-effects Cox model using the R package 'coxme' [45] for
both males and females respectively. In each case, we specified the model as: survival ~
knockdown *(functional and knockdown)* x sex *(male and female)* x dose *(4 concentrations of P. entomophila)* + (1|vial *(n replicates)*),
with 'knockdown' and 'sex' as fixed effects, and vials as a random effect for all the fly lines (IMD-
pathway mutants). We fit separate models for each knockdown fly line and functional (wild type) fly
line, since they were assayed independently on different days. We estimated the impact of
harbouring IMD-pathway knockdowns as the hazard ratio of IMD-knockdowns vs. fully functional fly
lines. A hazard ratio significantly greater than one indicates higher risk of mortality in the IMD-
mutants relative to fully functional individuals; hence, a significant impact on fly's survival.

Measuring disease tolerance: We calculated tolerance as fly survival relative to its bacterial load
(lifespan/mean bacterial load, also known as the per pathogen mortality (hours survival^{-1}/ CFU/fly),
akin to the often used per-pathogen-pathogenicity. This is metric of tolerance is intuitive to interpret:
relative to a given microbe load, does one line die more (less tolerant) or less (more tolerant) [46–
49]. We analysed the per-pathogen-mortality (PPM) as PPM ~ knockdown *(functional and knockdown)* x sex
(male and female) x dose *(4 concentrations of P. entomophila)*.

Results

Relish knockdown flies had lower expression of IMD-regulated AMP expression (*AttC, CecA1* and *Dpt*) and higher internal loads of *P. entomophila*, as expected. Loss of negative regulators *caudal* and *pirk* resulted in higher AMP expression and lower bacterial loads (**Fig. 1, Table**-*S3-4).* These results were consistent at three other infection doses (**Fig**.*S2* for bacterial load across different infection doses).

Given our observation that knocking down negative regulators of IMD resulted in an increase in AMP expression and lower microbe loads, we would expect mortality in these flies to be lower than controls if variation in mortality was mainly explained by pathogen clearance. However, while mortality was higher in *relish* and *DiptericinB* knockdowns as expected under reduced resistance, we also observed higher mortality when knocking down the expression of both negative regulators, *pirk* and *caudal* (*Fig. S3-S5, Table S5 for survival curves and hazard ratios across different infection doses*). This occurred despite the increased expression of AMPs and their relatively lower bacterial load, which is suggestive of reduced tolerance.

To quantify the tolerance response of each fly line more precisely, we measured the per-pathogen-mortality (measured as fly survival relative to its bacterial load) of male and female RNAi-lines of the IMD-pathway (*relish, caudal, pirk* and *DptB*), across four infectious doses (ranging from ~650 bacterial cells infection dose to leading up to ~2900 cells). Overall, we found that flies with UAS-RNAi-mediated knockdown in expression any of the genes of interest showed an increase in the per-pathogen-mortality, for all tested infection doses, and without substantial differences between sexes (Fig. 2, Table 1). In flies with reduced *caudal* expression, we detected significant sex-by-knockdown interaction, driven by smaller effects of UAS-RNAi of *caudal* in males at higher doses (Table 1).

Discussion

Compared to the mechanisms of immune clearance our knowledge of the mechanisms underlying increased tolerance remains incomplete. We predicted that knocking down the expression of negative immune regulators would reduce the ability to flies to tolerate infection with *P. entomophilia*, reflected as an increase in the per-pathogen-mortality. While this was observed, we also observed higher per-pathogen mortality in *relish* and *dptB* knockdown flies. However, the underlying cause of this increase is likely to differ between lines (Figure S6). Flies with reduced expression of *relish* or *DptB* showed higher per-pathogen mortality, but internal loads in these flies were at least an order of magnitude higher than controls because of disrupted AMP production (Fig 1). A higher per-pathogen-mortality in this case may therefore reflect feedbacks between increased pathogen growth and decreased ability to limit damage when microbe loads get too high (that is, the per-pathogen mortality becomes higher in flies unable to control pathogen loads in the first place).

By contrast, UAS-RNAi of the negative regulators *caudal* and *pirk* caused higher per-pathogen
mortality despite these flies being able to achieve lower pathogen loads compared to functional
control flies (Fig. 1). This is indicative of reduced tolerance, as flies experienced greater mortality
despite harbouring fewer pathogens. Our results therefore highlight that the negative regulators
caudal and *pirk* are important not only in regulating pathogen clearance by the IMD pathway, but by
affecting the mortality of flies independently of their microbe loads, negative regulators of IMD would
fit the functional definition of a tolerance mechanisms [9,13,21].

This increased mortality of flies lacking negative regulators of IMD such as *pirk* and *caudal* has been
observed previously [38,40,50] and highlights that mechanisms other than pathogen clearance
actively contribute to the variation in mortality during systemic bacterial infection. One likely cause of
flies lacking negative IMD regulation still succumb to death despite reducing pathogen burden is
immunopathology due to continued immune activation. In a similar example of negative immune
regulation affecting disease tolerance, *G9a*, a negative regulator of Jak/Stat, was identified as being
important in tolerating *Drosophila C Virus* infections by reducing immunopathology during the
antiviral immunity [22]. During bacterial infections, *PGRP-LB* functions as a negative regulator of the
IMD pathway by regulating the IMD-responsive AMP *Diptericin* [51], and it has been shown that
PGRP-LB mutant flies experience sex-differences in survival despite undetectable differences in
their bacterial loads [52], which is consistent with an effect on tolerance. We note that in the current
experiment we did not detect substantial sex differences in the per-pathogen-mortality.

Recognising the role of negative immune regulators in disease tolerance phenotype opens up the
possibility of using them as therapeutic targets to improve host health [4,5,14,53]. Future work may
also investigate if natural genetic variation in disease tolerance can be explained by variation in
negative immune regulators [27], offering a more complete mechanistic understanding of host
heterogeneity in infection outcomes.

Acknowledgments

We thank Angela Reid, Lucinda Rowe, James King and Alison Fulton for help with media
preparation. We acknowledge funding and support from the Branco Weiss fellowship and a
Chancellor's Fellowship to PFV; a Darwin Trust PhD studentship to AP from the School of Biological
Sciences, The University of Edinburgh. For the purpose of open access, the author has applied a
Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising
from this submission.

References

1. Graham AL, Allen JE, Read AF. 2005 Evolutionary Causes and Consequences of Immunopathology. *Annu. Rev. Ecol. Evol. Syst.* **36**, 373–397. (doi:10.1146/annurev.ecolsys.36.102003.152622)

2. Sears BF, Rohr JR, Allen JE, Martin LB. 2011 The economy of inflammation: when is less more? *Trends Parasitol.* **27**, 382–387. (doi:10.1016/j.pt.2011.05.004)

3. Badinloo M, Nguyen E, Suh W, Alzahrani F, Castellanos J, Klichko VI, Orr WC, Radyuk SN. 2018 Overexpression of antimicrobial peptides contributes to aging through cytotoxic effects in *Drosophila* tissues. *Arch. Insect Biochem. Physiol.* **98**, e21464. (doi:10.1002/arch.21464)

4. Vale PF, Fenton A, Brown SP. 2014 Limiting Damage during Infection: Lessons from Infection Tolerance for Novel Therapeutics. *PLoS Biol.* **12**, e1001769. (doi:10.1371/journal.pbio.1001769)

5. Vale PF, McNally L, Doeschl-Wilson A, King KC, Popat R, Domingo-Sananes MR, Allen JE, Soares MP, Kümmerli R. 2016 Beyond killing: Can we find new ways to manage infection? *Evol. Med. Public Health* **2016**, 148–157. (doi:10.1093/emph/eow012)

6. Louie A, Song KH, Hotson A, Thomas Tate A, Schneider DS. 2016 How Many Parameters Does It Take to Describe Disease Tolerance? *PLOS Biol.* **14**, e1002435. (doi:10.1371/journal.pbio.1002435)

7. Hill AVS. 1998 The Immunogenetics of Human Infectious Diseases. *Annu. Rev. Immunol.* **16**, 593–617. (doi:10.1146/annurev.immunol.16.1.593)

8. Råberg L, Graham AL, Read AF. 2009 Decomposing health: tolerance and resistance to parasites in animals. *Philos. Trans. R. Soc. B Biol. Sci.* **364**, 37–49. (doi:10.1098/rstb.2008.0184)

9. Kutzer MAM, Armitage SAO. 2016 Maximising fitness in the face of parasites: a review of host tolerance. *Zoology* **119**, 281–289. (doi:10.1016/j.zool.2016.05.011)

10. Lazzaro BP, Sceurman BK, Clark AG. 2004 Genetic Basis of Natural Variation in D. melanogaster Antibacterial Immunity. *Science* **303**, 1873–1876. (doi:10.1126/science.1092447)

11. Malo D, Skamene E. 1994 Genetic control of host resistance to infection. *Trends Genet.* **10**, 365–371. (doi:10.1016/0168-9525(94)90133-3)

12. Raberg L, Sim D, Read AF. 2007 Disentangling Genetic Variation for Resistance and Tolerance to Infectious Diseases in Animals. *Science* **318**, 812–814. (doi:10.1126/science.1148526)

13. Ayres JS, Schneider DS. 2012 Tolerance of Infections. , 26.

14. Martins R, Carlos AR, Braza F, Thompson JA, Bastos-Amador P, Ramos S, Soares MP. 2019 Disease Tolerance as an Inherent Component of Immunity. *Annu. Rev. Immunol.* **37**, 405–437. (doi:10.1146/annurev-immunol-042718-041739)

15. Prakash A, Monteith KM, Vale PF. 2022 Mechanisms of damage prevention, signalling, and repair impact disease tolerance. *Proc. R. Soc. B* , 2021.10.03.462916. (doi:10.1101/2021.10.03.462916)

16. Soares MP, Teixeira L, Moita LF. 2017 Disease tolerance and immunity in host protection against infection. *Nat. Rev. Immunol.* **17**, 83–96. (doi:10.1038/nri.2016.136)

17. Soares MP, Gozzelino R, Weis S. 2014 Tissue damage control in disease tolerance. *Trends*
Immunol. **35**, 483–494. (doi:10.1016/j.it.2014.08.001)

18. Troha K, Im JH, Revah J, Lazzaro BP, Buchon N. 2018 Comparative transcriptomics reveals
CrebA as a novel regulator of infection tolerance in D. melanogaster. *PLOS Pathog.* **14**,
e1006847. (doi:10.1371/journal.ppat.1006847)

19. Howick VM, Lazzaro BP. 2014 Genotype and diet shape resistance and tolerance across
distinct phases of bacterial infection. *BMC Evol. Biol.* **14**, 56. (doi:10.1186/1471-2148-14-56)

20. Dionne MS, Pham LN, Shirasu-Hiza M, Schneider DS. 2006 Akt and foxo Dysregulation
Contribute to Infection-Induced Wasting in Drosophila. *Curr. Biol.* **16**, 1977–1985.
(doi:10.1016/j.cub.2006.08.052)

21. Lissner MM, Schneider DS. 2018 The physiological basis of disease tolerance in insects. *Curr.*
Opin. Insect Sci. **29**, 133–136. (doi:10.1016/j.cois.2018.09.004)

22. Merkling SH, Bronkhorst AW, Kramer JM, Overheul GJ, Schenck A, Rij RPV. 2015 The
Epigenetic Regulator G9a Mediates Tolerance to RNA Virus Infection in Drosophila. *PLOS*
Pathog. **11**, e1004692. (doi:10.1371/journal.ppat.1004692)

23. Adelman JS, Kirkpatrick L, Grodio JL, Hawley DM. 2013 House finch populations differ in early
inflammatory signaling and pathogen tolerance at the peak of Mycoplasma gallisepticum
infection. *Am. Nat.* **181**, 674–89. (doi:10.1086/670024)

24. Nahrendorf W, Ivens A, Spence PJ. 2021 Inducible mechanisms of disease tolerance provide an
alternative strategy of acquired immunity to malaria. *eLife* **10**, e63838. (doi:10.7554/eLife.63838)

25. Vilaplana C, Marzo E, Tapia G, Diaz J, Garcia V, Cardona PJ. 2013 Ibuprofen therapy resulted
in significantly decreased tissue bacillary loads and increased survival in a new murne
experimental model of active tuberculosis. *J. Infect. Dis.* **208**, 199–202.

26. Gupta V, Vale PF. 2017 Nonlinear disease tolerance curves reveal distinct components of host
responses to viral infection. *R. Soc. Open Sci.* **4**, 170342. (doi:10.1098/rsos.170342)

27. Kutzer M, Gupta V, Neophytou K, Doublet V, Monteith KM, Vale PF. 2023 Intraspecific genetic
variation in host vigor, viral load, and disease tolerance during Drosophila C Virus infection.
Open Biol. , 2022.06.09.495537. (doi:10.1101/2022.06.09.495537)

28. Prakash A, Monteith KM, Vale PF. 2021 Negative regulation of IMD contributes to disease
tolerance during systemic bacterial infection in Drosophila. *bioRxiv* , 2021.09.23.461574.
(doi:10.1101/2021.09.23.461574)

29. Prakash A, Bonnet M, Monteith KM, Vale PF. 2021 The Jak/Stat pathway mediates disease
tolerance during systemic bacterial infection in Drosophila. *bioRxiv* , 2021.09.23.461578.
(doi:10.1101/2021.09.23.461578)

30. Leulier F, Parquet C, Pili-Floury S, Ryu J-H, Caroff M, Lee W-J, Mengin-Lecreulx D, Lemaitre B.
2003 The Drosophila immune system detects bacteria through specific peptidoglycan
recognition. *Nat. Immunol.* **4**, 478–484. (doi:10.1038/ni922)

31. Lemaitre B, Hoffmann J. 2007 The Host Defense of *Drosophila melanogaster. Annu. Rev.*
Immunol. **25**, 697–743. (doi:10.1146/annurev.immunol.25.022106.141615)

32. Myllymäki H, Valanne S, Rämet M. 2014 The Drosophila Imd Signaling Pathway. *J. Immunol.*
192, 3455–3462. (doi:10.4049/jimmunol.1303309)

33. Vodovar N, Vinals M, Liehl P, Basset A, Degrouard J, Spellman P, Boccard F, Lemaitre B. 2005
Drosophila host defense after oral infection by an entomopathogenic Pseudomonas species.
Proc. Natl. Acad. Sci. **102**, 11414–11419. (doi:10.1073/pnas.0502240102)

34. Dieppois G, Opota O, Lalucat J, Lemaitre B. 2015 Pseudomonas entomophila: A Versatile
Bacterium with Entomopathogenic Properties. In *Pseudomonas* (eds J-L Ramos, JB Goldberg,
A Filloux), pp. 25–49. Dordrecht: Springer Netherlands. (doi:10.1007/978-94-017-9555-5_2)

35. Lee K-Z, Ferrandon D. 2011 Negative regulation of immune responses on the fly: Immune
responses on the fly. *EMBO J.* **30**, 988–990. (doi:10.1038/emboj.2011.47)

36. Kleino A, Silverman N. 2014 The Drosophila IMD pathway in the activation of the humoral
immune response. *Dev. Comp. Immunol.* **42**, 25–35. (doi:10.1016/j.dci.2013.05.014)

37. Aggarwal K, Rus F, Vriesema-Magnuson C, Ertürk-Hasdemir D, Paquette N, Silverman N. 2008
Rudra Interrupts Receptor Signaling Complexes to Negatively Regulate the IMD Pathway. *PLoS*
Pathog. **4**, e1000120. (doi:10.1371/journal.ppat.1000120)

38. Kleino A, Myllymäki H, Kallio J, Vanha-aho L-M, Oksanen K, Ulvila J, Hultmark D, Valanne S,
Rämet M. 2008 Pirk Is a Negative Regulator of the *Drosophila* Imd Pathway. *J. Immunol.* **180**,
5413–5422. (doi:10.4049/jimmunol.180.8.5413)

39. Basbous N, Coste F, Leone P, Vincentelli R, Royet J, Kellenberger C, Roussel A. 2011 The
Drosophila peptidoglycan⊥recognition protein LF interacts with peptidoglycan⊥recognition
protein LC to downregulate the Imd pathway. *EMBO Rep.* **12**, 327–333.
(doi:10.1038/embor.2011.19)

40. Ryu J-H *et al.* 2008 Innate Immune Homeostasis by the Homeobox Gene Caudal and
Commensal-Gut Mutualism in Drosophila. *Science* **319**, 777–782.
(doi:10.1126/science.1149357)

41. Paredes JC, Welchman DP, Poidevin M, Lemaitre B. 2011 Negative Regulation by Amidase
PGRPs Shapes the Drosophila Antibacterial Response and Protects the Fly from Innocuous
Infection. *Immunity* **35**, 770–779. (doi:10.1016/j.immuni.2011.09.018)

42. Livak KJ, Schmittgen TD. 2001 Analysis of Relative Gene Expression Data Using Real-Time
Quantitative PCR and the 2−ΔΔCT Method. *Methods* **25**, 402–408.
(doi:10.1006/meth.2001.1262)

43. Wickham H. 2016 R for data science: import, tidy, transform, visualize, and model data.

44. Bunn A, Korpela M. 2019 An Introduction to dplR. , 16.

45. Therneau T. 2015 Mixed Effects Cox Models. In *Mixed Effects Cox Models*, CRAN repository.

46. Adelman JS, Kirkpatrick L, Grodio JL, Hawley DM. 2013 House Finch Populations Differ in Early
Inflammatory Signaling and Pathogen Tolerance at the Peak of *Mycoplasma gallisepticum*
Infection. *Am. Nat.* **181**, 674–689. (doi:10.1086/670024)

47. Bertels F *et al.* 2018 Dissecting HIV Virulence: Heritability of Setpoint Viral Load, CD4+ T-Cell
Decline, and Per-Parasite Pathogenicity. *Mol. Biol. Evol.* **35**, 27–37.
(doi:10.1093/molbev/msx246)

48. Hidalgo BA, Silva LM, Franz M, Regoes RR, Armitage SAO. 2022 Decomposing virulence to
understand bacterial clearance in persistent infections. *Nat. Commun.* **13**, 5023.
(doi:10.1038/s41467-022-32118-1)

49. Knutie SA, Wilkinson CL, Wu QC, Ortega CN, Rohr JR. 2017 Host resistance and tolerance of
parasitic gut worms depend on resource availability. *Oecologia* **183**, 1031–1040.
(doi:10.1007/s00442-017-3822-7)

50. Morris O *et al.* 2016 Signal Integration by the IκB Protein Pickle Shapes Drosophila Innate Host
Defense. *Cell Host Microbe* **20**, 283–295. (doi:10.1016/j.chom.2016.08.003)

51. Zaidman-Rémy A *et al.* 2006 The Drosophila amidase PGRP-LB modulates the immune
response to bacterial infection. *Immunity* **24**, 463–473. (doi:10.1016/j.immuni.2006.02.012)

52. Vincent CM, Dionne MS. 2021 Disparate regulation of IMD signaling drives sex differences in
infection pathology in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci.* **118**, e2026554118.
(doi:10.1073/pnas.2026554118)

53. Lopez BS. 2022 Can Infectious Disease Control Be Achieved without Antibiotics by Exploiting
Mechanisms of Disease Tolerance? *ImmunoHorizons* **6**, 730–740.
(doi:10.4049/immunohorizons.2200043)

Caudal	Df	F value	p-value
UAS-RNAi	1	58.83	2.1×10^{-13}
Sex	1	1.09	0.30
Dose	3	4.77	2.9×10^{-03}
UAS-RNAi $\times$ Sex	1	7.36	0.01
UAS-RNAi $\times$ Dose	3	0.71	0.55
Sex $\times$ Dose	3	2.96	0.03
DptB			
UAS-RNAi	1	53.22	2.3×10^{-12}
Sex	1	0.01	0.91
Dose	3	8.52	1.8×10^{-05}
UAS-RNAi $\times$ Sex	1	0.16	0.69
UAS-RNAi $\times$ Dose	3	0.64	0.59
Sex $\times$ Dose	3	2.25	0.08
Pirk			
UAS-RNAi	1	36.00	5.2×10^{-09}
Sex	1	0.36	0.55
Dose	3	6.46	2.9×10^{-04}
UAS-RNAi $\times$ Sex	1	0.69	0.41
UAS-RNAi $\times$ Dose	3	0.97	0.41
Sex $\times$ Dose	3	0.68	0.57
Relish			
UAS-RNAi	1	46.90	3.8×10^{-11}
Sex	1	3.01	0.08
Dose	3	1.77	0.15
UAS-RNAi $\times$ Sex	1	0.17	0.68
UAS-RNAi $\times$ Dose	3	0.33	0.80
Sex $\times$ Dose	3	1.01	0.39

343 **Table 1.** ANOVA testing the effects on the per-pathogen-mortality of RNAi knockdown of the
344 expression of each gene of interest, fly sex and infection dose of *P. entomophila*. Details of statistical
345 analysis are given in the methods section.

Figure 1: **(A)** Quantification of major IMD-responsive AMPs by qRT-PCR (n=15-21 flies/sex/treatment) for IMD-pathway mutants and functional flies (males and females) around 8 hours following *P. entomophila* systemic infection (0.05 OD or ~2000 bacterial cells/µl). Each data point represents pooled data of 3 flies for each infection treatment, fly line and sex **(B)** internal bacterial load after 8 hours following infection (n=16-24 flies/sex/treatment). Asterisks inside the panels indicate significant differences in AMP expression (that is, impact of knocking down on overall AMPs measured) and bacterial load between knockdown flies and flies with fully intact IMD functioning (p<0.05). The error bars represent standard error.

Figure 2: Disease tolerance in IMD-pathway mutants and fully functional flies. The relationship between host (fly) survival and bacterial load of 4 infection doses [that is, (I) ~650 cells (II) ~1400 cells (III) ~2000 cells (IV) ~2900 cells] of *P. entomophila*, analysing using point tolerance 8 hours following *P. entomophila* infection for **(A)** male and **(B)** female flies. Each data represents 'point tolerance', the ratio estimates calculated from median lifespan and average bacterial load for IMD-pathway knockdown and flies with fully functional IMD. Asterisks inside the panel indicates significant difference in disease tolerance between knockdown flies and with fully intact IMD functioning ($p < 0.05$) across different infection doses. The error bars represent standard error.

365