1 # *Duox* and *Jak/Stat* signalling influence disease tolerance in
2 # Drosophila during *Pseudomonas entomophila* infection
3

Arun Prakash[1,2]*, Katy M. Monteith[1], Mickael Bonnet[3],Pedro F. Vale[1]*

[1]Institute of Ecology and Evolution, School of Biological Sciences, University of Edinburgh, UK

[2]Current address: Department of Biological Sciences, Vanderbilt University, USA

[3]UFR De Biologie, Campus Universitaire Des Cezeaux, France

* Correspondance

arunpadmaprakash@gmail.com

pedro.vale@ed.ac.uk

14
15

16 **Author Contributions:** AP and PFV conceived and designed the experiments. AP, MB, KMM,
17 performed the experiments. AP analysed the data with help from PFV. AP and PFV wrote the
18 paper with additional inputs from MB & KMM.
19
20 **Competing Interest Statement:** The authors declare no competing interests.
21
22
Keywords: Antibacterial immunity, Damage signalling, Disease tolerance, immune regulation,
reactive oxygen species*, Drosophila melanogaster*

Abstract

Disease tolerance describes an infected host's ability to maintain health independently of the
ability to clear microbe loads. The Jak/Stat pathway plays a pivotal role in humoral innate
immunity by detecting tissue damage and triggering cellular renewal, making it a candidate
tolerance mechanism. Here, we find that in *Drosophila melanogaster* infected with *Pseudomonas*
entomophila disrupting *ROS*-producing *dual oxidase (duox)* or the negative regulator of Jak/Stat
Socs36E, render male flies less tolerant. Another negative regulator of Jak/Stat*, G9a* - which has
previously been associated with variable tolerance of viral infections – did not affect the rate of
mortality with increasing microbe loads compared to flies with functional *G9a*, suggesting it does
not affect tolerance of bacterial infection as in viral infection. Our findings highlight that ROS
production and Jak/Stat signalling influence the ability of flies to tolerate bacterial infection sex-
specifically and may therefore contribute to sexually dimorphic infection outcomes in *Drosophila*.

1. Introduction

When organisms experience infection, they face two major challenges to return to a healthy state. The first challenge is to identify and clear the source of the infection. Individuals capable of dealing with the first challenge exhibit low microbe loads because their immune clearance mechanisms are very effective, and are typically labelled 'resistant'(Boon et al., 2009; Ganz and Ebert, 2010; Lazzaro et al., 2006; Wang et al., 2017). The mechanisms underlying host resistance have been well characterized empirically and often involve the detection of pathogen-derived molecular patterns such as peptidoglycans, and triggering signalling cascades including the immune deficiency (IMD) and Toll pathways, resulting in the downstream expression of antimicrobial peptides (AMPs) that directly kill pathogens (Kleino and Silverman, 2014; Myllymäki et al., 2014; Myllymäki and Rämet, 2014; Palmer et al., 2018; Valanne et al., 2011).

While crucial, pathogen clearance alone will not result in a healthy host, because after pathogen elimination what is left is the tissue damage caused by pathogen growth and as a side-effect of immunopathology. The second challenge to return to healthy state is therefore to repair and regenerate damaged tissues (Martins et al., 2019; Medzhitov et al., 2012; Prakash et al., 2022; Schneider and Ayres, 2008; Soares et al., 2017, 2014). Effective mechanisms of damage signalling and repair may explain why some individuals are tolerant of infection, and are able to experience relatively high health even if their pathogen loads remain high or are not completely cleared (Martins et al., 2019; Soares et al., 2014).

Compared to well-described pathogen clearance mechanisms, we are only beginning to unravel the mechanistic basis of disease tolerance (Martins et al., 2019; Medzhitov et al., 2012; Prakash et al., 2022; Soares et al., 2017, 2014). Likely candidate mechanisms underlying effective tolerance of infection include those that regulate inflammation to reduce immunopathology (Adelman et al., 2013; Cornet et al., 2014; Prakash et al., 2021; Sears et al., 2011); detoxification of host or pathogen derived metabolites (Ferreira et al., 2011; Soares et al., 2017; Vale et al., 2014); or tissue protection and regeneration (Jamieson et al., 2013; Prakash et al., 2022; Soares et al., 2017, 2014). However, the few disease tolerance candidate genes arising from genome-wide association or transcriptomic studies - such as *ghd* (*grainyhead*)*, dsb (debris buster), crebA (cyclic response element binding protein) and, dfoxo* (*forkhead box, sub-group O*) - do not appear to be directly associated with classical immune pathways (Dionne et al., 2006; Howick and Lazzaro, 2014; Lissner and Schneider, 2018; Troha et al., 2018).

Here we take advantage of the detailed knowledge of *Drosophila* immunity to investigate the role of damage signalling plays in disease tolerance during systemic bacterial infection. In response to mechanical injury, oxidative stress, and infection, the Jak/Stat pathway is activated by cytokine-like ligands of the unpaired family namely *upd-1, upd-2* and *upd-3* (Agaisse et al., 2003; Chakrabarti et al., 2016; Dostert et al., 2005; Ekengren et al., 2001; Ekengren and Hultmark, 2001; Gilbert et al., 2005; Harrison et al., 1998). *Upd-3* is produced during damage caused by reactive oxygen species (*ROS*), which in turn are produced by *dual oxidase* (*duox*) (Babior, 1995; Klebanoff, 1974; Lee and Kim, 2014). The extracellular binding of *upd-3* to *Domeless* (*dome*), leads to the phosphorylation of *Hopscotch* (*hop*). This then leads to the phosphorylation of *Stat92E*, and its translocation to the nucleus (Myllymäki and Rämet, 2014). In the nucleus, in addition to the production of factors that are necessary for repairing cellar damage, *Stat92E* also induces the expression of *Socs36E*, a negative regulator of *Hopscotch* (Kiu and Nicholson, 2012). Recent work has also highlighted the role of the histone H3 lysine 9

methyltransferase (also called *G9a*) in negatively regulating the expression of the Jak/Stat
pathway during infection (Merkling et al., 2015).

Focusing on its role in immunity, there is substantial evidence that Jak/Stat signalling
plays a key role in wound healing, gut immunity, and downstream AMP production (Chakrabarti
et al., 2016; Kemp et al., 2013; Lamiable and Imler, 2014; Tafesh-Edwards and Eleftherianos,
2020). For instance, during enteric bacterial infection in *Drosophila*, the Jak/Stat pathway
contributes to intestinal immunity by regulating intestinal stem cell (ISC) proliferation and
epithelial cell renewal via epidermal growth factor (*EGFR)* signalling (Buchon et al., 2010;
Chakrabarti et al., 2016; Ohlstein and Spradling, 2006). The absence of epithelial renewal leads
to a loss of structural integrity and increased susceptibility to bacterial infections (Buchon et al.,
2009). In cellular immunity, Jak/Stat signalling is central to the production, differentiation and
maintenance of blood cells in insects (Banerjee et al., 2019; Meister and Lagueux, 2003). The
Jak/Stat pathway is also important in humoral immunity to viral infection (Dostert et al., 2005),
where a loss of regulation of Jak/Stat by the epigenetic negative regulators *G9a* results in
reduced tolerance of *Drosophila C virus* infections due to increased immunopathology (Merkling
et al., 2015). This specific result motivated us to question whether the effects of *G9a*-mediated
Jak/Stat regulation on tolerance were specific to viral infection, or if the regulation of Jak/Stat
also affects disease tolerance during bacterial infection.

We investigated the tolerance response of *Drosophila* during septic infection with the
bacterial pathogen *P. entomophila*, using transgenic flies lacking various components of Jak/Stat
signalling and regulation. Further motivated by the widespread observation of sexually dimorphic
immunity *reviewed in* (Belmonte et al., 2020; Klein and Flanagan, 2016) and particularly that the
effects of *G9a* on tolerance of *DCV* infection are more pronounced in female flies (Gupta and
Vale, 2017; Merkling et al., 2015), we also focused on assessing sex differences in how Jak/Stat
signalling affects tolerance of *P.entomophila* infection.

110
111
2. Materials and methods
2.1   Fly strains and maintenance
We used several *D*. melanogaster transgenic lines with TE mobilization using a P-element
construct and subsequent loss-of-function for *Duox* - P{SUPor-P}Duox^{KG07745} (Hurd et al., 2015) ,
Domeless - P{SUPor-P}^{KG08434}*, Hopscotch* - P{SUPor-P}hop^{KG01990}(Bellen et al., 2004), *Socs36E*
- P{EPgy2}Socs36E^{EY06665} (Monahan and Starz-Gaiano, 2013). All lines were on the *yw*
background (Eleftherianos et al., 2014) which served as a control genotype (detailed information
is presented in **Fig** S1 and S2 and Table S1). We also used *G9a* mutant flies (that is, *G9a*$^{-/-}$, also
known as *G9a*^{DD2} generated previously by mobilization of the P-element *KG01242* located in the
5' UTR of the gene(Kramer et al., 2011)) and control *G9a*$^{+/+}$ (Merkling et al., 2015). We
maintained all the fly lines in a 12ml plastic vials on a standard cornmeal diet *see* (Siva-Jothy et
al., 2018), at 25°C (±2°C). We used 3-5-day-old adult flies for all our experiments (see below).
First, we housed 2 males and 5 females for egg laying (48 hours) in a vial containing fresh food.
We then removed the adults and the vials containing the eggs were kept in 25°C incubator for 14
days, or until pupation. We placed the newly eclosed individuals (males and females separately)
in fresh food vials until the experimental day (3 days).

128
129 ### 2.2 Bacterial culture preparation
130 We used *P. entomophila* cultured overnight in Luria broth (LB) at 37°C under constant agitation
131 that is, 120 revolutions per minute (rpm). *P. entomophila* is a gram-negative bacterium naturally

found in soil and aquatic environments, known to be highly pathogenetic for *D. melanogaster* (Dieppois et al., 2015; Vodovar et al., 2005). Upon reaching 0.75 OD_{600} we pelleted the culture by centrifuging during 5 minutes at 5000rpm at 4°C, and then removed the supernatant. We resuspended the bacteria in 1xPBS (phosphate buffer saline) and prepared the final infection inoculum of OD_{600} of 0.05 for all our infection assays.

2.3 Systemic infection assay

We used a split-vial experimental design (see *Fig. S3*), where, after infection, each vial containing 25 flies (of each sex and fly line combination) were divided into 2 vials for measuring **(A)** survival following infection (n= 15 vials of 15-17 flies/vial/infection treatment/sex/fly line) and **(B)** internal bacterial load (n= 15 vials of 8-10 flies/vial/infection treatment/sex/fly line). With this split-vial design we were able to use replicate-matched data for both survival and bacterial load to estimate disease tolerance for each fly line (that is, for each replicate group, mean fly survival with respect to mean internal bacterial load). We infected 3-5-day old male and female adult flies using a 0.14mm insect minutein needles bent at 90° angle to avoid damaging the internal tissues by dipping in *P. entomophila* bacterial inoculum of OD_{600} of 0.05, resulting in 50-70 bacterial cells/fly. For mock controls we substituted bacterial solution with sterile 1xPBS. After stabbed the flies in the sternopleural region of the thorax (Khalil et al., 2015). We then placed males and females separately onto fresh food vials and incubated at 25°C. We scored the flies (both infected and control) every 2-3 hours for the first 48-hours following infection, then 2-3 times each day for the next 6 days (150 hours).

2.4 Measuring bacterial load

To quantify internal bacterial load after 24-hours following systemic *P. entomophila* infection first, we thoroughly washed each fly with 70% ethanol for 30 sec to surface sterilize and then rinsed twice with autoclaved distilled water. We plated the second wash on LB agar plates and confirmed that the surface bacteria were successfully removed after sterilization. We then transferred individual fly onto 1.5ml micro centrifuge tubes and homogenized using a motorized pestle for approximately 30-60 seconds in 100μl LB broth (n=30 fly homogenates/sex/infection treatment/ fly line). We performed serial dilution of each fly homogenate up to 10^{-6} fold and added 4μL aliquot on a LB agar plate. We incubated the plate overnight for 18h at 30°C and counted the resultant bacterial colonies manually (Siva-Jothy et al., 2018). We note that mock-infected control fly homogenates did not produce any colonies on LB agar plates.

2.5 Statistical analyses

2.5.1 Survival: We analysed the survival data with a Cox mixed effects model using the R package 'coxme' (*Therneau 2015*) for different treatment groups (*P. entomophila* systemic infection and mock controls) across males and females. We specified the model as: survival ~ fly line * treatment * sex * (1|vials/block), with 'fly line', 'treatment' and 'sex' and their interactions as fixed effects, and 'vials' nested within a 'block' as a random effect.

2.5.2 Bacterial load: We found that the bacterial load data were not normally distributed (tested with Shapiro–Wilks's test for normality). We therefore used a non-parametric one-way ANOVA Kruskal-Wallis test to test the effects of each fly line and sex on internal bacterial load**.**

2.5.3 Measuring disease tolerance: We analysed disease tolerance as the linear relationship between fly survival against bacterial load (Ayres and Schneider, 2012; Louie et al., 2016; Oliveira et al., 2020; Raberg et al., 2007). To this end, we employed ANCOVA by fitting 'fly line' and 'sex' as categorical fixed effects and 'bacterial load' as a continuous covariate, and their interactions as fixed effects. Since we were interested in identifying how each transgenic line

180 differed from the control line, we compared the estimates of the model slope using pairwise
181 comparison (f-test; *yw* vs. different transgenic lines) to test the extent to which each transgenic
182 line significantly differed from the control in tolerating bacterial infections.
183
184 **3. Results and Discussion**
185
3.1. Following systemic bacterial infection, disruption of *Duox* or different components of
Jak/Stat pathway result in variable survival outcomes

Overall, we found that disruption of *Duox* or the Jak/Stat pathway (either by disrupting the
positive regulators *upd3* and *domeless*, or overactivation by disrupting the negative regulator
socs36E) affected fly survival during bacterial *P. entomophila* infections (**Fig.** *1A* and *B,* **Table** *1*
and *SI-2*). Both male and female flies lacking *duox* (*ROS* producing *dual oxidase*) were more
susceptible to *P. entomophila* infections compared to the control line (*yw*) (**Fig.** *1A* and *B,* **Table**
1 and *SI-2*). However, other transgenic lines showed slightly improved survival relative to the
functional control line. These included male and female flies lacking the transmembrane receptor
domeless, and males lacking the negative regulator *Soc36E* (see hazard ratio in **Fig.** *1B,* **Table** *1*
and *SI-2*).

197
198**1. 3.2 Control *yw* and Duox / Jak/Stat transgenic deletion lines exhibit similar bacterial loads**

We investigated whether the variation we observed between transgenic lines in mortality
could be explained by differences in their bacterial load. Given that most mortality occurred just
after 24 hours for most of our fly genotypes (**Fig.** *1A*) we quantified bacterial load at 24 hours
following infection. Both control and transgenic lines exhibited similar levels of bacterial load 24
hours following infection with *P. entomophila* (**Fig.** *1C,* **Table** *SI-3*). Therefore, despite no
substantial difference in microbe loads at 24-hours post infection, transgenic lines showed
variable survival. This would fit the functional definition of disease tolerance as for the same
bacterial load some lines appear to be more tolerant (survive longer, such as *domeless*) while
others are less tolerant (e.g., *duox*).

Figure 1. (A) Survival curves for control *yw* flies and flies lacking Jak/Stat pathway components for females and males exposed to systemic *P. entomophila* of infection dose OD_{600}=0.05 (n= 15 vials with 15-17 flies each vial/fly line/treatment/sex/infection dose). ['*' indicates that the Jak/Stat transgenic lines are significantly different from *yw* flies]. **(B)** Estimated hazard ratios calculated from the survival curves for males and female flies (*yw* and with flies lacking components of Jak/Stat signalling and *duox*). A greater hazard ratio (>1) indicates higher susceptibility of Jak/Stat mutants than control while (<1) indicates transgenic lines have better survival than control flies to systemic bacterial infection (p=<0.05). **(C)** Bacterial load (mean log_{10}) measured 24 hours following infection (n= 15 vials with 8-10 flies each vial/fly line/treatment/sex/infection dose). [significantly different fly lines are connected by different letters using Tukey's HSD as a post hoc analysis of pairwise comparisons].

221 2. 3.3 Disrupted expression of *Duox* or *Jak-Stat* signalling leads to differences in disease tolerance phenotypes

While the results above are indicative of variable tolerance depending on the Jak/Stat disruption, we carried out a formal analysis of disease tolerance using the slope of the linear

reaction norm between fly survival and microbe load, where each data point is the matched
survival / CFU data for one replicate vial (see methods and *Fig. S3* for description of split-vial
design). Here, the differences in tolerance between Jak/Stat deletion and the control fly line are
indicated by a significant interaction between the bacterial load and the fly line for survival, which
reflects the overall rate at which fly health (survival) changes with bacterial load between fly lines.
Overall, we found that the transgenic lines showed differences in disease tolerance phenotypes
compared to control in both males and females, and this effect was driven mainly the Duox-
deficient lines, which showed a much steeper decline in survival with increasing *P. entomophila*
bacterial loads (***Fig.** 2A and 2B, **Tables 2 and** 3*). Given the role of *duox* in producing *ROS*, one
possible explanation for decreased tolerance in the *duox* transgenic line is flies require
intracellular *ROS* (oxidative burst) such as H_2O_2 (hydrogen peroxide) for the activation of cellular
reponses during wounding and injury, in addition to Toll and Jak/Stat activation (Chakrabarti and
Visweswariah, 2020). In other work, wild type ($w^{Dahomey}$) males showed higher levels of *duox*
expression and *ROS* following *Ecc* (*Erwinia catovora*) infection (Regan et al., 2016), which may
suggest that loss of function of *duox* might impact males more than females, as observed in this
experiment (Fig. 2B).

An unexpected observation was that flies lacking *domeless* showed slightly increased
survival relative to the *yw* control (Fig 1) (and a trend for increased tolerance, though not
statistically significant, Fig 2). Given the role of *domeless* as an activator of Jak-Stat signalling ,
this might suggest that Jak/Stat activation may be costly to flies. While immune deployment and
regulation is highly energy demanding across most species (McKean et al., 2008; Nystrand and
Dowling, 2020; Schwenke et al., 2016; Vale et al., 2015), the physiological costs of specific
individual immune components and pathways remains understudied and an open question for
future research.

***Figure 2.* (A).** The relationship between fly survival (measured as mean lifespan) and mean
bacterial load (as mean CFUs - Colony Forming Units) analysed using linear models for female
and male flies. Each point shows data for median lifespan and mean CFUs of 15 vials (with each
vial containing 25 flies/sex/fly line combination after 24 hours post systemic bacterial exposure.
The data shown here are for the infection doses (OD_{600}=0.05). **(B).** Represents estimates of
negative slope of the linear reaction norm extracted from the linear models. [Maroon asterisks '*'
on the lower side of the panel B indicates that transgenic lines are significantly different from
control *yw*, analysed using the F-test pairwise comparisons of estimates of the linear reaction
norm for both males and females separately (see Table-3)]. Grey asterisks '*' on the upper side
of the panel B indicates sex differences within the fly line that is, males and females significantly
differ in tolerance to systemic bacterial *P. entomophila* infection.

261
262

3. 3.4 Disruption of *G9a* does not affect tolerance of *P.entomophila*

The negative regulator of Jak/Stat, *G9a*, was previously identified as being important for
tolerating *Drosophila C Virus* (DCV) infections (Merkling et al., 2015). Subsequent work exploring
sex differences in this response found that *G9a^{+/+}* (control) females had higher tolerance than
G9a^{−/−} females, when measured across a range of viral DCV doses (Gupta and Vale, 2017). We
wanted to test whether the loss of function of *G9a* also affects fly survival and disease tolerance
in response to bacterial infections. Overall, we found that loss of *G9a* makes both males and
females more susceptible to *P. entomophila* infections, (***Fig. 3A*** for survival and ***Fig. 3B*** for

hazard ratio, **Table 4** and **Table SI-4**). To test if this increased mortality in *G9a^{-/-}* flies was
associated with higher bacterial replication we measured bacterial load following 24 hours *P.*
entomophila systemic infection. We found that *G9a^{-/-}* females exhibited higher bacterial load than
G9a^{+/+} (control) flies, while males showed similar bacterial load as *G9a^{+/+}* flies (**Fig. 3C, Table** *SI-*
5). However, the overall ability to tolerate *P. entomophila* bacterial infections (that is, measured
as *G9a* fly's survival relative to its bacterial load) remained similar across both males and
females *G9a* flies that is, both *G9a^{-/-}* and *G9a^{+/+}* controls (**Fig. 3D, Table 5**, and **Table 6** for
comparison between estimates of tolerance slope). Thus, despite the previously identified role of
this negative regulator of Jak/Stat in tolerating viral infections by reducing immunopathology
(Gupta and Vale, 2017; Merkling et al., 2015), *G9a* does not appear to affect bacterial disease
tolerance in either sex.

Figure 3. (A) Survival curves for control *G9a^{+/+}* flies and *G9a^{-/-}* flies lacking *G9a* the epigenetic
regulator of Jak/Stat for female and male flies exposed to systemic *P. entomophila* of infection
dose $OD_{600}=0.05$ [n=15 vials with 15-17 flies in each vial/fly line/treatment/sex]. **(B)** Estimated
hazard ratios calculated from the survival curves for males and female flies (control y*w* and flies
without *G9a*). A greater hazard ratio (>1) indicates higher susceptibility of *G9a^{-/-}* to bacterial
infection relative to control flies. ['*' indicates that the *G9a^{-/-}* flies are significantly different from
G9a^{+/+} flies]. **(C)** Bacterial load (mean $\log_{10}$) measured 24 hours following infection (n=15 vials
with 8-10 flies in each vial/fly line/ treatment and sex combination). [Significantly different fly lines

are connected by different letters using Tukey's HSD as a post hoc analysis of pairwise
comparisons]. **(D)** Linear tolerance to *P. entomophila* infection – the relationship between *G9a* fly
survival (measured as mean lifespan) and bacterial load (as mean CFUs - Colony Forming Units)
analysed using linear models for female and male flies (both $G9a^{-/-}$ and $G9a^{+/+}$).

295
296 4. Concluding remarks

Tissue damage signalling and repair mechanisms such as Jak/Stat are important from a
therapeutic perspective because they have the potential to boost host tolerance by minimising
disease severity (Soares et al., 2014; Vale et al., 2016). Our data show that loss of Jak/Stat
pathway components reduces overall survival following *P.entomophila* infection and that this is
not caused by impaired pathogen clearance but due to lower disease tolerance. These
observations have parallels in human infection. For instance, dysregulation of cytokines and
interferons (JAK signalling - Tyrosinekinase2) result in immunodeficiency while defective STAT
increases the risk of autoimmunity (O'Shea et al., 2014, 2013). Drugs that inhibit JAK have been
shown to be effective in treating several autoimmune diseases by targeting cytokine-dependent
pathways, while STAT inhibitors have been promising candidates in the context of cancer
(Miklossy et al., 2013; Pérez-Jeldres et al., 2019; Salas et al., 2020). It may therefore be possible
to repurpose these existing drugs to improve host tolerance of infection. In summary, our work
highlights that Jak/Stat directly impacts the ability to tolerate bacterial infection and that this
response differs between males and females. Jak/Stat mediated disease tolerance may be a
potential source of sexually dimorphic response to infection in *Drosophila*.

312
313 5. Acknowledgments
314

We thank Ashworth fly group members for helpful discussion, and Angela Reid, Lucinda Rowe,
James King and Alison Fulton for help with media preparation. We acknowledge funding and
support from the Branco Weiss fellowship and a Chancellor's Fellowship to PFV; a Darwin Trust
PhD studentship to AP from the School of Biological Sciences, The University of Edinburgh. For
the purpose of open access, the author has applied a Creative Commons Attribution (CC BY)
licence to any Author Accepted Manuscript version arising from this submission.

322 6. References

Adelman, J.S., Kirkpatrick, L., Grodio, J.L., Hawley, D.M., 2013. House Finch Populations Differ
in Early Inflammatory Signaling and Pathogen Tolerance at the Peak of *Mycoplasma*
gallisepticum Infection. Am. Nat. 181, 674–689. https://doi.org/10.1086/670024
Agaisse, H., Petersen, U.-M., Boutros, M., Mathey-Prevot, B., Perrimon, N., 2003. Signaling Role
of Hemocytes in Drosophila JAK/STAT-Dependent Response to Septic Injury. Dev. Cell
5, 441–450. https://doi.org/10.1016/S1534-5807(03)00244-2
Ayres, J.S., Schneider, D.S., 2012. Tolerance of Infections 26.
Babior, B.M., 1995. The respiratory burst oxidase. Curr. Opin. Hematol. 55–60.
Banerjee, U., Girard, J.R., Goins, L.M., Spratford, C.M., 2019. Drosophila as a Genetic Model for
Hematopoiesis. Genetics 211, 367–417. https://doi.org/10.1534/genetics.118.300223
Bellen, H.J., Levis, R.W., Liao, G., He, Y., Carlson, J.W., Tsang, G., Evans-Holm, M., Hiesinger,
P.R., Schulze, K.L., Rubin, G.M., Hoskins, R.A., Spradling, A.C., 2004. The BDGP Gene
Disruption Project: Single Transposon Insertions Associated With 40% of Drosophila
Genes. Genetics 167, 761–781. https://doi.org/10.1534/genetics.104.026427

Belmonte, R.L., Corbally, M.-K., Duneau, D.F., Regan, J.C., 2020. Sexual Dimorphisms in Innate
Immunity and Responses to Infection in Drosophila melanogaster. Front. Immunol. 10,
3075. https://doi.org/10.3389/fimmu.2019.03075

Boon, A.C.M., deBeauchamp, J., Hollmann, A., Luke, J., Kotb, M., Rowe, S., Finkelstein, D., Lu,
L., Williams, R.W., Webby, R.J., 2009. Host Genetic Variation Affects Resistance to
Infection with a Highly Pathogenic H5N1 Influenza A Virus in Mice. J. Virol. 83, 10.
https://doi.org/10.1128/JVI.00514-09

Buchon, N., Broderick, N.A., Kuraishi, T., Lemaitre, B., 2010. DrosophilaEGFR pathway
coordinates stem cell proliferation and gut remodeling following infection. BMC Biol. 8,
152. https://doi.org/10.1186/1741-7007-8-152

Buchon, N., Broderick, N.A., Poidevin, M., Pradervand, S., Lemaitre, B., 2009. Drosophila
Intestinal Response to Bacterial Infection: Activation of Host Defense and Stem Cell
Proliferation. Cell Host Microbe 5, 200–211. https://doi.org/10.1016/j.chom.2009.01.003

Chakrabarti, S., Dudzic, J.P., Li, X., Collas, E.J., Boquete, J.-P., Lemaitre, B., 2016. Remote
Control of Intestinal Stem Cell Activity by Haemocytes in Drosophila. PLOS Genet. 12,
e1006089. https://doi.org/10.1371/journal.pgen.1006089

Chakrabarti, S., Visweswariah, S.S., 2020. Intramacrophage ROS Primes the Innate Immune
System via JAK/STAT and Toll Activation. Cell Rep. 33, 108368.
https://doi.org/10.1016/j.celrep.2020.108368

Cornet, S., Bichet, C., Larcombe, S., Faivre, B., Sorci, G., 2014. Impact of host nutritional status
on infection dynamics and parasite virulence in a bird-malaria system. J. Anim. Ecol. 83,
256–265. https://doi.org/10.1111/1365-2656.12113

Dieppois, G., Opota, O., Lalucat, J., Lemaitre, B., 2015. Pseudomonas entomophila: A Versatile
Bacterium with Entomopathogenic Properties, in: Ramos, J.-L., Goldberg, J.B., Filloux, A.
(Eds.), Pseudomonas. Springer Netherlands, Dordrecht, pp. 25–49.
https://doi.org/10.1007/978-94-017-9555-5_2

Dionne, M.S., Pham, L.N., Shirasu-Hiza, M., Schneider, D.S., 2006. Akt and foxo Dysregulation
Contribute to Infection-Induced Wasting in Drosophila. Curr. Biol. 16, 1977–1985.
https://doi.org/10.1016/j.cub.2006.08.052

Dostert, C., Jouanguy, E., Irving, P., Troxler, L., Galiana-Arnoux, D., Hetru, C., Hoffmann, J.A.,
Imler, J.-L., 2005. The Jak-STAT signaling pathway is required but not sufficient for the
antiviral response of drosophila. Nat. Immunol. 6, 946–953.
https://doi.org/10.1038/ni1237

Ekengren, S., Hultmark, D., 2001. A Family of Turandot-Related Genes in the Humoral Stress
Response of Drosophila. Biochem. Biophys. Res. Commun. 284, 998–1003.
https://doi.org/10.1006/bbrc.2001.5067

Ekengren, S., Tryselius, Y., Dushay, M.S., Liu, G., Steiner, H., Hultmark, D., 2001. A humoral
stress response in Drosophila. Curr. Biol. 11, 714–718. https://doi.org/10.1016/S0960-
9822(01)00203-2

Eleftherianos, I., More, K., Spivack, S., Paulin, E., Khojandi, A., Shukla, S., 2014. Nitric Oxide
Levels Regulate the Immune Response of Drosophila melanogaster Reference
Laboratory Strains to Bacterial Infections. Infect. Immun. 82, 4169–4181.
https://doi.org/10.1128/IAI.02318-14

Ferreira, A., Marguti, I., Bechmann, I., Jeney, V., Chora, Â., Palha, N.R., Rebelo, S., Henri, A.,
Beuzard, Y., Soares, M.P., 2011. Sickle Hemoglobin Confers Tolerance to Plasmodium
Infection. Cell 145, 398–409. https://doi.org/10.1016/j.cell.2011.03.049

Ganz, H.H., Ebert, D., 2010. Benefits of host genetic diversity for resistance to infection depend
on parasite diversity. Ecology 91, 1263–1268. https://doi.org/10.1890/09-1243.1

Gilbert, M.M., Weaver, B.K., Gergen, J.P., Reich, N.C., 2005. A novel functional activator of the
Drosophila JAK/STAT pathway, unpaired2, is revealed by an in vivo reporter of pathway
activation. Mech. Dev. 122, 939–948. https://doi.org/10.1016/j.mod.2005.03.004

Gupta, V., Vale, P.F., 2017. Nonlinear disease tolerance curves reveal distinct components of
host responses to viral infection. R. Soc. Open Sci. 4, 170342.
https://doi.org/10.1098/rsos.170342

Harrison, D.A., McCoon, P.E., Binari, R., Gilman, M., Perrimon, N., 1998. Drosophila unpaired
encodes a secreted protein that activates the JAK signaling pathway. Genes Dev. 12,
3252–3263. https://doi.org/10.1101/gad.12.20.3252

Howick, V.M., Lazzaro, B.P., 2014. Genotype and diet shape resistance and tolerance across
distinct phases of bacterial infection. BMC Evol. Biol. 14, 56.
https://doi.org/10.1186/1471-2148-14-56
Hurd, T.R., Liang, F.-X., Lehmann, R., 2015. Curly Encodes Dual Oxidase, Which Acts with
Heme Peroxidase Curly Su to Shape the Adult Drosophila Wing: e1005625. PLoS Genet.
11. https://doi.org/10.1371/journal.pgen.1005625
Jamieson, A.M., Pasman, L., Yu, S., Gamradt, P., Homer, R.J., Decker, T., Medzhitov, R., 2013.
Role of Tissue Protection in Lethal Respiratory Viral-Bacterial Coinfection. Science 340,
1230–1234. https://doi.org/10.1126/science.1233632
Kemp, C., Mueller, S., Goto, A., Barbier, V., Paro, S., Bonnay, F., Dostert, C., Troxler, L., Hetru,
C., Meignin, C., Pfeffer, S., Hoffmann, J.A., Imler, J.-L., 2013. Broad RNA Interference–
Mediated Antiviral Immunity and Virus-Specific Inducible Responses in Drosophila. J.
Immunol. 190, 650–658. https://doi.org/10.4049/jimmunol.1102486
Khalil, S., Jacobson, E., Chambers, M.C., Lazzaro, B.P., 2015. Systemic Bacterial Infection and
Immune Defense Phenotypes in Drosophila Melanogaster. J. Vis. Exp. JoVE 52613.
https://doi.org/10.3791/52613
Kiu, H., Nicholson, S.E., 2012. Biology and significance of the JAK/STAT signalling pathways.
Growth Factors 30, 88–106. https://doi.org/10.3109/08977194.2012.660936
Klebanoff, S.J., 1974. Role of the Superoxide Anion in the Myeloperoxidase-mediated
Antimicrobial System. J. Biol. Chem. 249 No. 12, 3724–3728.
Klein, S.L., Flanagan, K.L., 2016. Sex differences in immune responses. Nat. Rev. Immunol. 16,
626–638. https://doi.org/10.1038/nri.2016.90
Kleino, A., Silverman, N., 2014. The Drosophila IMD pathway in the activation of the humoral
immune response. Dev. Comp. Immunol. 42, 25–35.
https://doi.org/10.1016/j.dci.2013.05.014
Kramer, J.M., Kochinke, K., Oortveld, M.A.W., Marks, H., Kramer, D., de Jong, E.K., Asztalos, Z.,
Westwood, J.T., Stunnenberg, H.G., Sokolowski, M.B., Keleman, K., Zhou, H., van
Bokhoven, H., Schenck, A., 2011. Epigenetic regulation of learning and memory by
Drosophila EHMT/G9a. PLoS Biol. 9, e1000569.
https://doi.org/10.1371/journal.pbio.1000569
Lamiable, O., Imler, J.-L., 2014. Induced antiviral innate immunity in Drosophila. Curr. Opin.
Microbiol., Host–microbe interactions: fungi/parasites/viruses 20, 62–68.
https://doi.org/10.1016/j.mib.2014.05.006
Lazzaro, B.P., Sackton, T.B., Clark, A.G., 2006. Genetic Variation in Drosophila melanogaster
Resistance to Infection: A Comparison Across Bacteria. Genetics 174, 1539–1554.
https://doi.org/10.1534/genetics.105.054593
Lee, W.-J., Kim, S.-H., 2014. Role of DUOX in gut inflammation: lessons from Drosophila model
of gut-microbiota interactions. Front. Cell. Infect. Microbiol. 3, 116.
https://doi.org/10.3389/fcimb.2013.00116
Lissner, M.M., Schneider, D.S., 2018. The physiological basis of disease tolerance in insects.
Curr. Opin. Insect Sci., Global change biology * Molecular physiology 29, 133–136.
https://doi.org/10.1016/j.cois.2018.09.004
Louie, A., Song, K.H., Hotson, A., Thomas Tate, A., Schneider, D.S., 2016. How Many
Parameters Does It Take to Describe Disease Tolerance? PLOS Biol. 14, e1002435.
https://doi.org/10.1371/journal.pbio.1002435
Martins, R., Carlos, A.R., Braza, F., Thompson, J.A., Bastos-Amador, P., Ramos, S., Soares,
M.P., 2019. Disease Tolerance as an Inherent Component of Immunity. Annu. Rev.
Immunol. 37, 405–437. https://doi.org/10.1146/annurev-immunol-042718-041739
McKean, K.A., Yourth, C.P., Lazzaro, B.P., Clark, A.G., 2008. The evolutionary costs of
immunological maintenance and deployment. BMC Evol. Biol. 8, 76.
https://doi.org/10.1186/1471-2148-8-76
Medzhitov, R., Schneider, D.S., Soares, M.P., 2012. Disease Tolerance as a Defense Strategy.
Science 335, 936–941. https://doi.org/10.1126/science.1214935
Meister, M., Lagueux, M., 2003. Drosophila blood cells. Cell. Microbiol. 5, 573–580.
https://doi.org/10.1046/j.1462-5822.2003.00302.x

Merkling, S.H., Bronkhorst, A.W., Kramer, J.M., Overheul, G.J., Schenck, A., Rij, R.P.V., 2015.
The Epigenetic Regulator G9a Mediates Tolerance to RNA Virus Infection in Drosophila.
PLOS Pathog. 11, e1004692. https://doi.org/10.1371/journal.ppat.1004692
Miklossy, G., Hilliard, T.S., Turkson, J., 2013. Therapeutic modulators of STAT signalling for
human diseases. Nat. Rev. Drug Discov. 12, 611–629. https://doi.org/10.1038/nrd4088
Monahan, A.J., Starz-Gaiano, M., 2013. Socs36E attenuates STAT signaling to optimize motile
cell specification in the Drosophila ovary. Dev. Biol. 379, 152–166.
https://doi.org/10.1016/j.ydbio.2013.03.022
Myllymäki, H., Rämet, M., 2014. JAK/STAT Pathway in Drosophila Immunity. Scand. J. Immunol.
79, 377–385. https://doi.org/10.1111/sji.12170
Myllymäki, H., Valanne, S., Rämet, M., 2014. The Drosophila Imd Signaling Pathway. J.
Immunol. 192, 3455–3462. https://doi.org/10.4049/jimmunol.1303309
Nystrand, M., Dowling, D.K., 2020. Effects of immune challenge on expression of life-history and
immune trait expression in sexually reproducing metazoans—a meta-analysis. BMC Biol.
18, 135. https://doi.org/10.1186/s12915-020-00856-7
Ohlstein, B., Spradling, A., 2006. The adult Drosophila posterior midgut is maintained by
pluripotent stem cells. Nature 439, 470–474. https://doi.org/10.1038/nature04333
Oliveira, J.H., Bahia, A.C., Vale, P.F., 2020. How are arbovirus vectors able to tolerate infection?
Dev. Comp. Immunol. 103, 103514. https://doi.org/10.1016/j.dci.2019.103514
O'Shea, J.J., Holland, S.M., Staudt, L.M., 2013. JAKs and STATs in Immunity,
Immunodeficiency, and Cancer. N. Engl. J. Med. 368, 161–170.
https://doi.org/10.1056/NEJMra1202117
O'Shea, J.J., Schwartz, D.M., Villarino, A.V., Gadina, M., McInnes, I.B., Laurence, A., 2014. The
JAK-STAT Pathway: Impact on Human Disease and Therapeutic Intervention 21.
Palmer, W.H., Varghese, F.S., Van Rij, R.P., 2018. Natural Variation in Resistance to Virus
Infection in Dipteran Insects. Viruses 10, 118. https://doi.org/10.3390/v10030118
Pérez-Jeldres, T., Tyler, C.J., Boyer, J.D., Karuppuchamy, T., Yarur, A., Giles, D.A., Yeasmin,
S., Lundborg, L., Sandborn, W.J., Patel, D.R., Rivera-Nieves, J., 2019. Targeting
Cytokine Signaling and Lymphocyte Traffic via Small Molecules in Inflammatory Bowel
Disease: JAK Inhibitors and S1PR Agonists. https://doi.org/10.3389/fphar.2019.00212
Prakash, A., Monteith, K.M., Vale, P.F., 2022. Mechanisms of damage prevention, signalling, and
repair impact disease tolerance. Proc. R. Soc. B 2021.10.03.462916.
https://doi.org/10.1101/2021.10.03.462916
Prakash, A., Monteith, K.M., Vale, P.F., 2021. Negative regulation of IMD contributes to disease
tolerance during systemic bacterial infection in Drosophila. bioRxiv 2021.09.23.461574.
https://doi.org/10.1101/2021.09.23.461574
Raberg, L., Sim, D., Read, A.F., 2007. Disentangling Genetic Variation for Resistance and
Tolerance to Infectious Diseases in Animals. Science 318, 812–814.
https://doi.org/10.1126/science.1148526
Regan, J.C., Khericha, M., Dobson, A.J., Bolukbasi, E., Rattanavirotkul, N., Partridge, L., 2016.
Sex difference in pathology of the ageing gut mediates the greater response of female
lifespan to dietary restriction. eLife 5, e10956. https://doi.org/10.7554/eLife.10956
Salas, A., Hernandez-Rocha, C., Duijvestein, M., Faubion, W., McGovern, D., Vermeire, S.,
Vetrano, S., Vande Casteele, N., 2020. JAK–STAT pathway targeting for the treatment of
inflammatory bowel disease. Nat. Rev. Gastroenterol. Hepatol. 17, 323–337.
https://doi.org/10.1038/s41575-020-0273-0
Schneider, D.S., Ayres, J.S., 2008. Two ways to survive infection: what resistance and tolerance
can teach us about treating infectious diseases. Nat. Rev. Immunol. 8, 889–895.
https://doi.org/10.1038/nri2432
Schwenke, R.A., Lazzaro, B.P., Wolfner, M.F., 2016. Reproduction–Immunity Trade-Offs in
Insects. Annu. Rev. Entomol. 61, 239–256. https://doi.org/10.1146/annurev-ento-010715-
023924
Sears, B.F., Rohr, J.R., Allen, J.E., Martin, L.B., 2011. The economy of inflammation: when is
less more? Trends Parasitol. 27, 382–387. https://doi.org/10.1016/j.pt.2011.05.004
Siva-Jothy, J.A., Prakash, A., Vasanthakrishnan, R.B., Monteith, K.M., Vale, P.F., 2018. Oral
Bacterial Infection and Shedding in Drosophila melanogaster. J. Vis. Exp. 57676.
https://doi.org/10.3791/57676

Soares, M.P., Gozzelino, R., Weis, S., 2014. Tissue damage control in disease tolerance. Trends
Immunol. 35, 483–494. https://doi.org/10.1016/j.it.2014.08.001
Soares, M.P., Teixeira, L., Moita, L.F., 2017. Disease tolerance and immunity in host protection
against infection. Nat. Rev. Immunol. 17, 83–96. https://doi.org/10.1038/nri.2016.136
Tafesh-Edwards, G., Eleftherianos, I., 2020. JNK signaling in Drosophila immunity and
homeostasis. Immunol. Lett. 226, 7–11. https://doi.org/10.1016/j.imlet.2020.06.017
Troha, K., Im, J.H., Revah, J., Lazzaro, B.P., Buchon, N., 2018. Comparative transcriptomics
reveals CrebA as a novel regulator of infection tolerance in D. melanogaster. PLOS
Pathog. 14, e1006847. https://doi.org/10.1371/journal.ppat.1006847
Valanne, S., Wang, J.-H., Rämet, M., 2011. The Drosophila Toll Signaling Pathway. J. Immunol.
186, 649–656. https://doi.org/10.4049/jimmunol.1002302
Vale, P.F., Fenton, A., Brown, S.P., 2014. Limiting Damage during Infection: Lessons from
Infection Tolerance for Novel Therapeutics. PLoS Biol. 12, e1001769.
https://doi.org/10.1371/journal.pbio.1001769
Vale, P.F., Lafforgue, G., Gatchitch, F., Gardan, R., Moineau, S., Gandon, S., 2015. Costs of
CRISPR-Cas-mediated resistance in Streptococcus thermophilus. Proc. Biol. Sci. 282,
20151270. https://doi.org/10.1098/rspb.2015.1270
Vale, P.F., McNally, L., Doeschl-Wilson, A., King, K.C., Popat, R., Domingo-Sananes, M.R.,
Allen, J.E., Soares, M.P., Kümmerli, R., 2016. Beyond killing: Can we find new ways to
manage infection? Evol. Med. Public Health 2016, 148–157.
https://doi.org/10.1093/emph/eow012
Vodovar, N., Vinals, M., Liehl, P., Basset, A., Degrouard, J., Spellman, P., Boccard, F., Lemaitre,
B., 2005. Drosophila host defense after oral infection by an entomopathogenic
Pseudomonas species. Proc. Natl. Acad. Sci. 102, 11414–11419.
https://doi.org/10.1073/pnas.0502240102
Wang, J.B., Lu, H.-L., Leger, R.J.S., 2017. The genetic basis for variation in resistance to
infection in the Drosophila melanogaster genetic reference panel. PLOS Pathog. 13,
e1006260. https://doi.org/10.1371/journal.ppat.1006260
534
535

536 **Tables**

537 *Table 1*: Summary of estimated hazard ratio from the cox proportional model. A greater hazard
538 ratio estimates (>1) indicates that Jak/Stat mutant flies are more susceptible to *P. entomophila*
539 infection than *yw* control flies while lower ratio (<1) indicates that transgenic lines have better
540 survival than *yw* control.
541

Fly line	*sex*	*estimate*	*P*	*lower 95%*	*upper 95%*
Domeless	Female	0.462	<0.001	0.391	0.548
	Male	0.383	<0.001	0.322	0.457
Duox	Female	2.017	<0.001	1.712	2.384
	Male	1.707	<0.001	1.455	2.009
Hopscotch	Female	0.830	0.03	0.701	0.986
	Male	0.694	<0.001	0.585	0.824
Socs36e	Female	0.990	0.91	0.843	1.167
	Male	0.795	0.006	0.676	0.937

545   **Table 2.** Summary of ANCOVA. To assess differences in infection tolerance (fly survival with
546   increasing bacterial burden) following systemic *P. entomophila* infection with $OD_{600}=0.05$
547   infection dose, 24 hours following infection. We analysed ANCOVA and fitted 'sex' as categorical
548   fixed effects, 'mean bacterial load (log_{10})' as a continuous covariate and their interactions as
549   fixed effects for the transgenic lines.
550

Fly line	Source	DF	Sum of Sq.	F ratio	P
Female	Fly line	4	24817.1	15.27	**<0.001**
	Bac. load	1	4482.9	11.03	**0.0012**
	Fly line X bac. load	4	7642.8	4.7	**0.0015**
Male	Fly line	4	16964.8	9.22	**<0.001**
	Bac. load	1	6122.6	13.32	**0.0004**
	Fly line X bac. load	4	5737.5	3.12	**0.017**

551
552
553
554

555   **Table 3***: Summary of F-test pairwise comparisons of estimates of linear slopes (from the linear
556   model) transgenic lines compared to the *yw* control.
557

sex	line	SSE	ddf	slope diff	std err	F ratio	p
Female	*Duox*	10018.68	30	-13.15	3.09	0.53	0.47
	Domeless	16075.94	45	1.83	2.25	8.38	**0.0058**
	Hopscotch	11897.99	34	-5.99	2.64	1.39	0.24
	Socs36E	27135.53	47	-3.77	2.28	1.48	0.22
Male	*Duox*	17512.77	34	-11.25	4.47	5.25	**0.028**
	Domeless	27106.63	49	-0.46	2.67	0.19	0.65
	Hopscotch	15019.55	36	-3.2	2.8	0.01	0.91
	Socs36E	22285.12	47	-6.17	2.75	0.54	0.46

558
559
560
561   **Table 4**: Summary of estimated hazard ratio from the cox proportional model. A greater hazard
562   ratio (>1) indicates that *G9a*[-/-] flies are more susceptible to *P. entomophila* infection than control
563   (*G9a*[+/+]) flies.
564

sex	Fly line	estimate	p	Std err
Female	G9a-/-	2.2	<0.001	0.75
Male	G9a-/-	1.41	<0.001	0.45

565
566
567

568 **Table 5.** Summary of ANCOVA. To assess differences in infection tolerance (fly survival with
569 increasing bacterial burden) following systemic *P. entomophila* infection with OD_{600}=0.05
570 infection dose, 24 hours following infection. We analysed ANCOVA and fitted 'sex' as categorical
571 fixed effects, 'mean bacterial load (log_{10})' as a continuous covariate and their interactions as
572 fixed effects for each of the fly lines (*G9a*).

	Fly line	*Source*	*DF*	*Sum of Sq.*	*F ratio*	*p*
G9a	*Female*	Fly line	1	6802.2	20.21	**<0.001**
		Bac. load	1	53.19	0.158	0.69
		Fly line X bac. load	1	1.610	0.004	0.94
	Male	Fly line	1	3042.5	8.685	**0.005**
		Bac. load	1	533.13	1.521	0.22
		Fly line X bac. load	1	166.21	0.474	0.49

574
575

576 **Table 6**: Summary of F-test pairwise comparisons of estimates of the linear slopes (linear
577 reaction norm) for *G9a $^{-/-}$* relative to *G9a $^{+/+}$* control fly lines.

Sex	*Fly line*	*Fly line*	*F Ratio*	*p*
Female	*G9a $^{-/-}$*	*G9a $^{+/+}$*	0.005	0.94
Male	*G9a $^{-/-}$*	*G9a $^{+/+}$*	0.474	0.49