1 # The *Jak/Stat* pathway mediates disease tolerance during systemic
2 # bacterial infection in *Drosophila*
3

Arun Prakash[1]*, Mickael Bonnet[2], Katy M. Monteith[1], Pedro F. Vale[1]*

[1]Institute of Evolutionary Biology

School of Biological Sciences, University of Edinburgh, UK

[2]UFR De Biologie,

Campus Universitaire Des Cezeaux, France

* Correspondance

arun.prakash@ed.ac.uk

pedro.vale@ed.ac.uk

Keywords: Antibacterial immunity, Damage signalling mechanisms, Disease tolerance, Negative
immune regulators

Abstract

Disease tolerance describes a host's ability to maintain health independently of the ability to clear

microbe loads. However, we currently know little about the mechanisms that underlie disease

tolerance or how known mechanisms of tissue damage signalling and repair may contribute to

variation in tolerance. The Jak/Stat pathway plays a pivotal role in *Drosophila* humoral innate

immunity, signalling tissue damage and triggering cellular renewal, making it a potential mechanism

underlying the disease tolerance phenotype. Here, we show that disrupting the Jak/Stat pathway

in *Drosophila melanogaster* alters disease tolerance during *Pseudomonas entomophila* systemic infection.

Overall, flies with disrupted Jak/Stat show variation in survival that is not explained by variation

in pathogen loads. For instance, mutations disrupting the function of *ROS*-producing *dual oxidase*

(duox) or the negative regulator of Jak/Stat, *Socs36E* render males less tolerant to systemic bacterial

infection but not females. We also investigated whether the negative regulator of Jak/Stat, *G9a* -

which has previously been associated with tolerance of viral infections – is also implicated in

tolerance of bacterial infection. While female flies lacking *G9a* showed higher mortality and

reduced bacterial clearance, disease tolerance did not differ between *G9a* mutants and the wildtype.

This suggests that *G9a* does not affect tolerance during systemic bacterial infection as it appears

to do with viral infection. Overall, our findings highlight that Jak/Stat signalling mediates disease

tolerance during systemic bacterial infection and that this response differs between males and

females. Our work therefore suggests that differences in Jak/Stat mediated disease tolerance may

be a potential source of sexually dimorphic response to infection in *Drosophila.*

Introduction

When organisms acquire infection, they face two major challenges in order to return to a healthy state. The first challenge is to identify and clear the source of the infection. While crucial, pathogen clearance alone will not result in a healthy host, because after pathogen elimination, what is left is the tissue damage caused by pathogen growth and as a side-effect of immunopathology. The second challenge to return to healthy state is therefore to minimise damage even before it occurs, and to repair and regenerate tissue after it has been damaged during infection (Martins et al. 2019; McCarville and Ayres 2018; Oliveira, Bahia, and Vale 2020; Soares, Teixeira, and Moita 2017; Vale, Fenton, and Brown 2014).

Individuals capable of dealing with the first challenge are typically labelled 'resistant', and exhibit low microbe loads because their immune clearance mechanisms are very effective (Boon et al. 2009; Ganz and Ebert 2010; Lazzaro, Sackton, and Clark 2006; Wang, Lu, and Leger 2017). The mechanisms underlying host resistance have been well characterized empirically and often involve detecting pathogen-derived molecular patterns such as peptidoglycans and triggering signalling cascades that result in the expression of antimicrobial peptides (AMPs) which directly kill the replicating pathogen (Kleino and Silverman 2014; Henna Myllymäki, Valanne, and Rämet 2014; H. Myllymäki and Rämet 2014; Palmer, Varghese, and Van Rij 2018; Valanne, Wang, and Rämet 2011). These responses have been described in great detail in the fruit fly *Drosophila*, and include the immune deficiency (Imd), Toll & Jak/Stat pathways (Bang 2019; Ferrandon et al. 2007; Hoffmann 2003; Hultmark 2003; Lemaitre and Hoffmann 2007).

Individuals with very effective mechanisms of damage limitation and repair, however, may not necessarily have a strong capacity for pathogen clearance, but will be successful in preventing or repairing tissue damage (Martins et al. 2019; Medzhitov, Schneider, and Soares 2012; Schneider and Ayres 2008; Soares, Teixeira, and Moita 2017; Soares, Gozzelino, and Weis 2014). Effective mechanisms of damage signalling and repair may explain why some individuals are tolerant of infection, and are able to experience relatively high health even if their pathogen loads remain high or are not completely cleared (Martins et al. 2019; Soares, Gozzelino, and Weis 2014). Compared to well-described pathogen clearance mechanisms, we are only beginning to unravel the mechanistic basis of disease tolerance. Likely candidate mechanisms include those that regulate inflammation to reduce immunopathology (Adelman et al. 2013; Cornet et al. 2014; Sears et al.

2011); detoxification of host or pathogen derived metabolites (Ferreira et al. 2011; Soares, Teixeira,
and Moita 2017; Vale, Fenton, and Brown 2014); or tissue protection and regeneration (Jamieson
et al. 2013; Soares, Teixeira, and Moita 2017; Soares, Gozzelino, and Weis 2014). However, the
few disease tolerance candidate genes arising from genome-wide association or transcriptomic
studies - such as *ghd (grainyhead), dsb (debris buster), crebA (cyclic response element binding protein) and, dfoxo*
(*forkhead box, sub-group O*) - do not appear to be associated with these functions, or with classical
immune pathways (Dionne et al. 2006; Howick and Lazzaro 2014; Lissner and Schneider 2018;
Troha et al. 2018). This may indicate that disease tolerance is a complex phenotype and that
dissecting its mechanistic basis may be more successful through a more focused examination of
the effect of specific candidate signalling pathways.

Here we take advantage of the detailed knowledge of *Drosophila* infection and immunity to
investigate whether damage signalling plays a role in disease tolerance during systemic bacterial
infection. We focus on the Janus kinase/signal transducer and activator of transcription (Jak/Stat)
pathway which is known to have important roles in signalling damage and tissue renewal. The
Jak/Stat pathway was originally shown to control multiple biological processes including
development, homeostasis, metabolism, and immunity in insects and mammals (Brivanlou and
Darnell 2002; Harrison 2012; H. Myllymäki and Rämet 2014; Stark and Darnell 2012; Zhou and
Agaisse 2012). In response to different types of stresses such as mechanical injury, oxidative stress,
and infection the Jak/Stat pathway is activated by cytokine-like ligands of the unpaired family
namely *upd-1, upd-2* and *upd-3* (Agaisse et al. 2003; Chakrabarti et al. 2016; Dostert et al. 2005;
Ekengren et al. 2001; Ekengren and Hultmark 2001; Gilbert et al. 2005; Harrison et al. 1998). *Upd-*
3 is produced during damage caused by reactive oxygen species (*ROS*), which in turn are produced
by *dual oxidase (duox)* (Klebanoff 1974; Babior 1995; Lee and Kim 2014) (see **Fig**. *1*). The
extracellular binding of *upd-3* to *Domeless (dome)*, leads to the phosphorylation of *Hopscotch (hop)*.
This then leads to the phosphorylation of *Stat92E*, and its translocation to the nucleus (H.
Myllymäki and Rämet 2014). In the nucleus, in addition to the production of factors that are
necessary for repairing cellar damage, *Stat92E* also induces the expression of *Socs36E*, a negative
regulator of *Hopscotch* (Kiu and Nicholson 2012). Recent work has also highlighted the role of the
histone H3 lysine 9 methyltransferase (also called *G9a*) in negatively regulating the expression of
the Jak/Stat pathway during infection (Merkling et al. 2015).

Focusing on its role in immunity, there is substantial evidence that Jak/Stat signalling plays a key
role in wound healing, gut immunity, and downstream AMP production (Chakrabarti et al. 2016;
Kemp et al. 2013; Lamiable and Imler 2014; Tafesh-Edwards and Eleftherianos 2020). For
instance, during enteric bacterial infection in *Drosophila*, the Jak/Stat pathway contributes to
intestinal immunity by regulating intestinal stem cell (ISC) proliferation and epithelial cell renewal
via epidermal growth factor (*EGFR) signalling* (Buchon et al. 2010; Chakrabarti et al. 2016; Ohlstein
and Spradling 2006). The absence of epithelial renewal leads to a loss of structural integrity and
increased susceptibility to bacterial infections (Buchon et al. 2009). In cellular immunity, Jak/Stat
signalling is central to the production, differentiation and maintenance of blood cells in insects
(Banerjee et al. 2019; Meister and Lagueux 2003).

Jak/Stat is also important in humoral immunity to viral infection (Dostert et al. 2005), where a loss
of regulation of Jak/Stat by the epigenetic negative regulators *G9a* results in reduced tolerance of
Drosophila C virus infections due to increased immunopathology (Merkling et al. 2015). This specific
result led us to question whether the effects of *G9a*-mediated Jak/Stat regulation on tolerance
were specific to viral infection, or if the regulation of Jak/Stat also affects disease tolerance during
bacterial infection. Here we investigated the tolerance response of *Drosophila* during septic
infection with the bacterial pathogen *P. entomophila*, using deletion mutant flies lacking various
components of Jak/Stat signalling and regulation. Further motivated by the widespread
observation of sexually dimorphic immunity *reviewed in* (Belmonte et al. 2020; Klein and Flanagan
2016) and particularly that the effects of *G9a* on tolerance of *DCV* infection are more pronounced
in female flies (Gupta and Vale 2017; Merkling et al. 2015), we also focused on assessing how
Jak/Stat affects males and females separately.

Abbreviations:
ISC - Intestinal stem cells
DUOX - Dual-oxidases
Dom - Domeless
Hops - Hopscotch
Upd3 - Unpaired 3
ROS - Reactive oxygen species
Socs36E - Suppressor of cytokine signaling at 36E
Stat92E - Signal-transducer & activator of transcription protein 92E

Figure 1. A simplified version of *Drosophila* Jak/Stat pathway known to have important roles in signalling damage and tissue renewal. In response to infection, the Jak/Stat pathway is activated by cytokine-like ligands of the unpaired family namely *upd-3*, produced during damage caused by reactive oxygen species (*ROS*), which in turn are produced by *dual oxidase* (*duox*). The extracellular binding of *upd-3* to *Domeless* (*dome*), leads to the phosphorylation of *Hopscotch* (*hop*). This then leads to the phosphorylation of *Stat92E*, and its translocation to the nucleus. In the nucleus, *Stat92E* induces the expression of *Socs36E*, a negative regulator of *Hopscotch*. Recent work has also highlighted the role of the *histone H3 lysine 9 methyltransferase* (also called *G9a*) in negatively regulating the expression of the Jak/Stat pathway during infection.

Materials and methods

<u>*Fly strains and maintenance:*</u> we used several *D. melanogaster* strains with loss-of-function mutations for the *Jak/Stat* pathway. All the fly strains were on the same genetic background as that of *yw (189)* including *Domeless (Bloomington stock# 14726)*, *Hopscotch (15097)*, *Socs36E (16744)*, *Duox (16468)*. We also used *G9a* mutant flies (i.e., $G9a^{-/-}$) and control ($G9a^{+/+}$), also known as $G9a^{DD2}$ see (Merkling et al. 2015) for details about *G9a* mutants. Previous work has shown that *yw* flies have lower basal levels of nitric oxide and are more susceptible to infection compared to other lab wild types such as w^{1118}, *Oregon R, Canton-S* (Eleftherianos et al. 2003). However, we found that males and females of both *yw* and another wildtype of *outcrossed* flies showed comparable survival following *P. entomophila* systemic infection and exhibited similar levels of internal bacterial loads when measured ~24 hours following infection (see supplementary *Fig. S1A & B*, *Table S1* for

survival; *Fig. S1C,* for bacterial load). We maintained all the fly lines in a 12ml plastic vials on a

standard cornmeal diet or Lewis's medium *see* (Siva-Jothy et al. 2018), at 25°C (±2°C). We used

~3-day old adult flies for all our experiments (see below). First, we housed 2 males and 5 females

for egg laying (48 hours) onto a vial containing fresh food. We then removed the adults and the

vials containing the eggs were kept in 25°C incubator for ~14 days, or until pupation. We placed

the newly eclosed individuals (males and females separately) in fresh food vials until the

experimental day (~3 days).

Bacterial culture preparation: We used *P. entomophila* cultured overnight in Luria broth (LB) at 37°C

under constant agitation i.e., 120 revolutions per minute (rpm). *P. entomophila* is a gram-negative

bacterium naturally found in soil and aquatic environments, known to be highly pathogenetic for

D. melanogaster (Dieppois et al. 2015; Vodovar et al. 2005)). Upon reaching ~0.75 OD_{600} we pelleted

the culture using a 4°C cold centrifuge during 5 minutes at 5000rpm and then removed the

supernatant. We resuspended the bacteria in 1X phosphate buffer saline (PBS) and prepared the

final infection dose of OD_{600} of 0.05 for all our infection assays.

Systemic infection assay: We used a split-vial experimental design (see **Fig. 2**), wherein after infection

each vial containing about 25 flies (of each treatment, sex and fly line combination) were divided

into 2 vials for measuring **(A)**. survival following infection (~15 flies/combination) and **(B)**.

internal bacterial load (see **Fig. 2II**). With this split-vial design we were able to use replicate-

matched data for both survival and bacterial load in order to estimate disease tolerance for each

fly line (i.e., fly survival with respect to internal bacterial load). We infected ~3-day old male and

female adult flies using a 0.15mm insect minutein needles bent at ~90° angle to avoid damaging

the internal tissues by dipping in *P. entomophila* bacterial solution. For mock controls we substituted

bacterial solution with sterile 1X PBS. After stabbed the flies in the sternopleural region of the

thorax (Khalil et al. 2015). We then placed males and females separately onto fresh food vials and

incubated at 25°C. We scored the flies (both infected and control) every 2-3 hours for the first 48

hours following infection, then 2-3 times each day for the next 6 days.

Measuring bacterial load: To quantify internal bacterial load after *P. entomophila* infection first, we

thoroughly washed each fly with 70% ethanol for ~30 sec to surface sterilize and then rinsed twice

with autoclaved distilled water. We plated the second wash on LB agar plates to confirm whether

the surface bacteria were successfully removed after sterilization. We then transferred individual

fly onto 1.5ml micro centrifuge tubes and homogenized using a motorized pestle for approximately

30-60 sec in 100μl LB broth (n=30 homogenates/sex/infection treatment/ fly line). We

performed serial dilution of each fly homogenate up to 10^{-6} fold and added 4μL aliquot on a LB

agar plate. We incubated the plate overnight for ~18h at 30°C and counted the resultant bacterial

colonies manually (Siva-Jothy et al. 2018). We note that mock-infected control fly homogenates

did not produce any colonies on LB agar plates.

Statistics.

1.   *Survival post oral bacterial infection*: We analysed the survival data with a Cox mixed effects model using the R package 'coxme' (*Therneau 2015*) for different treatment groups (*P. entomophila* systemic infection and mock controls) across males and females. We specified the model as: survival ~ fly line * treatment * sex * (1|vials/block), with 'fly line', 'treatment' and 'sex' and their interactions as fixed effects, and 'vials' nested within a 'block' as a random effect for wild type *yw* and *Jak/Stat* mutant fly lines.

2.   *Bacterial load*: We found that the bacterial load data were not normally distributed (tested with Shapiro–Wilks's test for normality). Hence, we first log transformed the data and then confirmed that the transformed residuals were still not normally distributed. Hence, we used a non-parametric one-way ANOVA Kruskal-Wallis test to test the effects of each fly lines (wild type *yw* and *Jak/Stat* mutants) on males and females following bacterial infection**.**

3.   *Measuring infection tolerance*: Finally, to understand how the Jak/Stat signalling contributes to the phenotype disease we analysed tolerance to bacterial infection by linear reaction norm (i.e., measuring the linear relationship between fly survival against bacterial load) (Raberg, Sim, and Read 2007; Ayres and Schneider 2012; Louie et al. 2016; Oliveira, Bahia, and Vale 2020). We analysed using ANCOVA for each of the fly lines (wild type *yw* and *Jak/Stat* mutants) by fitting 'fly line' and 'sex' as categorical fixed effects and 'bacterial load' as a continuous covariate and their interactions as fixed effects. Because we were interested in identifying how the Jak/Stat mutants differ from wildtype flies during disease tolerance - we compared the estimates of the slope using pairwise comparison (f-test; *w^{1118}* vs. different *Jak/Stat* mutants) to find out what extent these Jak/Stat mutants significantly differed from *yw* wild type in tolerating bacterial infections.

Figure 2: **(I)** Design of experiments to assay **(1)** survival and **(2)** internal bacterial load following systemic bacterial infection with *Pseudomonas entomophila* to test the role of the Jak/Stat pathway on **(3)** disease tolerance. **(II)** A split-vial experimental design, wherein after infection each vial containing about 25 flies of each treatment, sex and fly line combination were divided into 2 vials for measuring **(A)**. survival following infection (~15 flies/combination) and **(B)**. internal bacterial load (~10 flies/combination).

Results & discussion

1. **Following systemic bacterial infection, loss of different components of Jak/Stat pathway result in variable survival outcomes**

Overall, we found that disruption of Jak/Stat signalling affected fly survival during bacterial *P. entomophila* infections (**Fig.** *3A and B,* **Table** *1 & S2*). Both male and female flies lacking *duox* (*ROS* producing *dual oxidase*) were more susceptible to *P. entomophila* infections compared to the wildtype (*yw*) (**Fig.** *3A and B,* **Table** *1 & S2*). However, other mutants showed improved survival relative to wild types. These included male and female flies lacking the transmembrane receptor *domeless*, males lacking the negative regulator *Soc36E* and both sexes lacking *hopscotch* (see hazard ratio in **Fig.** *3B,* **Table** *1 & S2*).

2. **Both wildtype (*yw*) and Jak/Stat mutant flies exhibit similar bacterial loads**

Next, we tested whether the variation we observed between mutants in mortality could be explained by differences in their bacterial load. Given that most mortality occurred just after 24 hours for most of our fly genotypes (**Fig.** *3A*) we quantified bacterial load at 24 hours following infection. Both wild type and Jak/Stat mutants exhibited similar levels of bacterial load around 24 hours following infection with *P. entomophila* (**Fig.** *3C,* **Table** *S3*). Therefore, despite no substantial difference in microbe loads at 24 hours post infection, Jak/Stat mutants showed variable survival. This would fit the functional definition of disease tolerance as for the same bacterial load some lines appear to be more tolerant (survive longer, such as *domeless*) while others are less tolerant (e.g., *duox*).

Figure 3. (A) Survival curves for wildtype *yw* flies and flies lacking Jak/Stat pathway components for females and males exposed to systemic *P. entomophila* of infection dose OD$_{600}$=0.05 (n=20-30 flies/vial (~15 vials)/fly line/treatment/sex/infection dose). ['*' indicates that the *Jak/Stat* mutants are significantly different from *yw* flies]. **(B)** Estimated hazard ratios calculated from the survival curves for males and female flies (y*w* & with flies lacking components of Jak/Stat). A greater hazard ratio (>1) indicates higher susceptibility of Jak/Stat mutants than wild type while (<1) indicates Jak/Stat mutants have better survival than wild type *yw* flies to systemic bacterial infection (p=<0.05). **(C)** Bacterial load (mean ln10) measured around 24 hours following infection. [significantly different fly lines are connected with different alphabets using Tukey's HSD as a post hoc analysis of pairwise comparisons].

Fly line	sex	estimate	P	lower 95%	upper 95%
Domeless	Female	0.462	<0.001	0.391	0.548
	Male	0.383	<0.001	0.322	0.457

Duox	Female	2.017	<0.001	1.712	2.384
	Male	1.707	<0.001	1.455	2.009
Hopscotch	Female	0.830	0.03	0.701	0.986
	Male	0.694	<0.001	0.585	0.824
Socs36e	Female	0.990	0.91	0.843	1.167
	Male	0.795	0.006	0.676	0.937

Table 1: Summary of estimated hazard ratio from the cox proportional model. A greater hazard ratio (>1) indicates that Jak/Stat mutant flies are more susceptible to *P. entomophila* infection than *yw* wildtype flies while lower ratio (<1) indicates that Jak/Stat mutants have better survival than wildtype.

3. **During systemic bacterial infection, loss of Jak-Stat signalling leads to sex differences in disease tolerance phenotypes**

While the results above are indicative of variable tolerance depending on the Jak/Stat mutation, we carried out a formal analysis of disease tolerance using the slope of the linear reaction norm between fly survival and microbe load, where each data point is the matched survival / CFU data for one replicate vial (see methods for description of split-vial design and ***Fig. 2***). Here, the differences in tolerance between Jak/Stat mutant and *yw* fly lines are indicated by a significant interaction between the bacterial load and the fly line for survival, which reflects the overall rate at which fly health (survival) changes with bacterial load (tolerance) between fly lines. Overall, we found that the Jak/Stat mutants showed differences in disease tolerance phenotypes compared to wild type *yw* in both males and females (***Fig. 4, Table 2***). Females lacking *duox* or *Socs36E* showed higher tolerance than males to *P. entomophila* (***Fig. 4B and 4A, Table 3***). Given the role of *duox* in producing *ROS*, one possible explanation for increased tolerance in *duox* mutants is that *ROS*-derived damage is lower. However, flies also require intracellular *ROS* (oxidative burst) such as H_2O_2 (hydrogen peroxide) for the activation of macrophages during wounding and injury, in addition to Toll and Jak/Stat activation (Chakrabarti and Visweswariah 2020). It is possible that these effects are more pronounced in males which would explain our observation that female *duox* mutants have comparatively higher tolerance. In other work, wild type ($w^{Dahomey}$) males showed higher levels of *duox* expression and *ROS* following *Ecc* (*Erwinia catovora*) infection (Regan et al. 2016), which may suggest that loss of function of *duox* might impact males more than females.

Regarding the sex difference in disease tolerance in flies lacking *Socs36E,* it could be due to sex differences in regulation of Jak/Stat, and the resulting differences in immunopathology. For example, female flies lacking the negative epigenetic regulator *G9a*, were found to be more tolerant

than males during viral infections (Gupta and Vale 2017). Similar examples also exist from other innate immune pathways where, disrupting the negative regulator of IMD, *PGRP-LB* (*peptidoglycan receptor-LB*), affected survival to a greater extent in females following *E. coli* infection, suggesting the sex-specific role of some of these regulators (Vincent and Dionne 2021). The combination of these observations might explain why flies lacking *Socs36E* showed sex differences in disease tolerance, particularly in light of the crosstalk between Jak/Stat and IMD pathways (Kemp et al. 2013; Bang 2019; Dostert et al. 2005). An unexpected result is that we found that female flies lacking *domeless* showed increased survival (and tolerance) relative to the *yw* wildtype (***Fig. 3*** - survival *and **Fig**. 4* - tolerance), suggesting that Jak/Stat activation may in fact be costly to flies. Immune deployment and regulation is highly energy demanding (McKean et al. 2008; Schwenke, Lazzaro, and Wolfner 2016; Nystrand and Dowling 2020). The physiological cost of specific individual immune components and pathways remains an open question for future research.

(A).

Figure 4. **(A).** The relationship between fly survival (measured as median lifespan) and bacterial load (as mean CFUs - Colony Forming Units) analysed using linear models for female and male flies (*yw* & Jak/Stat mutants). Each point shows data for median lifespan and mean CFUs of 15 vials (with each vial containing ~25 flies/sex/fly line combination after 24 hours post systemic bacterial exposure. The data shown here are for the infection doses (OD_{600}=0.05). **(B).** Represents

estimates of negative slope of the linear reaction norm extracted from the linear models. [Asterix '*' on the lower side of the panel B indicates that Jak/Stat mutants are significantly different from wild type *yw*, analysed using the F-test pairwise comparisons of estimates of the linear reaction norm for both males and females separately (see Table-3)]. Asterix '*' on the upper side of the panel B indicates sex differences within the fly line i.e., males and females significantly differ in tolerance to systemic bacterial *P. entomophila* infection.

Fly line		Source	DF	Sum of Sq.	F ratio	P
Jak/Stat	*Female*	Fly line	1	100928.9	21.6	<0.001
		Bac. load	1	4273.8	3.66	0.06
		Fly line X bac. load	1	22262.4	4.77	0.001
	Male	Fly line	1	48485.2	5.72	0.0003
		Bac. load	1	23206.2	10.9	0.001
		Fly line X bac. load	1	26181.0	3.09	0.01

Table 2. Summary of ANCOVA. To assess differences in infection tolerance (fly survival with increasing bacterial burden) following systemic *P. entomophila* infection with OD_{600}=0.05 infection dose, around 24 hours following infection. We analysed ANCOVA and fitted 'sex' as categorical fixed effects, 'average bacterial load (ln_{10})' as a continuous covariate and their interactions as fixed effects for the Jak/Stat mutants.

Sex	Fly line	Difference (slope)	Std err.	F Ratio	p
Female	*duox vs. yw*	-18.11	7.32	2.13	0.15
	domeless vs. yw	28.04	6.28	6.32	0.01
	hopscotch vs. yw	7.87	6.76	0.05	0.81
	socs36E vs. yw	0.87	6.21	2.92	0.09
Male	*duox vs. yw*	7.20	8.64	12.51	0.001
	domeless vs. yw	34.74	6.80	0.52	0.47
	hopscotch vs. yw	23.30	7.52	1.65	0.20
	socs36E vs. yw	11.67	6.94	6.23	0.01

Table 3: Summary of F-test pairwise comparisons of estimates of linear slopes (from the linear model) Jak/Stat-mutants compared to the *yw* wildtype.

4. **Loss of *G9a* does not affect bacterial disease tolerance**

The epigenetic negative regulator of Jak/Stat *G9a* was previously identified as being important for tolerating *Drosophila C Virus* (DCV) infections (Merkling et al. 2015). Subsequent work exploring sex differences in this response found that $G9a^{+/+}$ (control) females had higher tolerance than $G9a^{-/-}$ females, when measured across a range of viral DCV doses (Gupta and Vale 2017). We wanted to test whether the loss of function of *G9a* also affects fly survival and disease tolerance in response to bacterial infections. Overall, we found that loss of *G9a* makes both males and females more susceptible to *P. entomophila* infections, (**Fig. 5A** for survival and **Fig. 5B** for hazard

ratio, **Table 4** and **Table** S4). To test if this increased mortality in *G9a*[-/-] flies was associated with higher bacterial replication we measured bacterial load following ~24 hours *P. entomophila* systemic infection. We found that *G9a*[-/-] females exhibited higher bacterial load than *G9a*[+/+] (control) flies, while males showed similar bacterial load as *G9a*[+/+] flies (**Fig. 5C, Table** S5). However, the overall ability to tolerate *P. entomophila* bacterial infections (i.e., measured as *G9a* fly's survival relative to its bacterial load) remained similar across both males and females *G9a* flies i.e., both *G9a*[-/-] and *G9a*[+/+] controls (see **Fig. 5D, Table 5,** and **Table 6** for comparison between estimates of tolerance slope). Thus, despite the previously identified role of this negative regulator of Jak/Stat in tolerating viral infections by reducing immunopathology (Gupta and Vale 2017; Merkling et al. 2015), *G9a* does not appear to affect bacterial disease tolerance in either sex.

Figure 5. (A) Survival curves for wildtype *yw* flies and flies lacking *G9a* the epigenetic regulator of Jak/Stat for female and male flies exposed to systemic *P. entomophila* of infection dose $OD_{600}=0.05$

sex	flyline	estimate	p	Std err
Female	G9a-/-	2.2	<0.001	0.75
Male	G9a-/-	1.41	<0.001	0.45

353 ***Table 4***: Summary of estimated hazard ratio from the cox proportional model. A greater hazard
354 ratio (>1) indicates that *G9a$^{-/-}$* mutant flies are more susceptible to *P. entomophila* infection than
355 control (*G9a$^{+/+}$*) flies.

	Fly line	Source	DF	Sum of Sq.	F ratio	P
G9a	**Female**	Fly line	1	12373.5	8.028	0.007
		Bac. load	1	414.4	0.268	0.60
		Fly line X bac. load	1	1469.8	0.953	0.33
	Male	Fly line	1	394.8	0.692	0.41
		Bac. load	1	343.4	0.602	0.44
		Fly line X bac. load	1	8.480	0.014	0.90

357 ***Table 5***. Summary of ANCOVA. To assess differences in infection tolerance (fly survival with
358 increasing bacterial burden) following systemic *P. entomophila* infection with OD_{600}=0.05 infection
359 dose, around ~24 hours following infection. We analysed ANCOVA and fitted 'sex' as categorical
360 fixed effects, 'average bacterial load (ln_{10})' as a continuous covariate and their interactions as fixed
361 effects for each of the fly lines (*G9a*).

Sex	Fly line	Fly line	F Ratio	p
Female	*G9a$^{-/-}$*	*G9a$^{+/+}$*	0.95	0.33
Male	*G9a$^{-/-}$*	*G9a$^{+/+}$*	0.01	0.90

363 ***Table 6***: Summary of F-test pairwise comparisons of estimates of the linear slopes (linear reaction
364 norm) for *G9a$^{-/-}$* relative to *G9a$^{+/+}$* wildtype fly lines.

5. Concluding remarks

367 Tissue damage signalling and repair mechanisms such as Jak/Stat are important from a therapeutic

368 perspective because they have the potential to boost host tolerance by minimising disease severity

369 (Soares, Gozzelino, and Weis 2014; Vale et al. 2016). Our data show that loss of Jak/Stat pathway

370 components following bacterial infection disrupts disease tolerance and overall survival. These

371 observations have parallels in human infection. For instance, dysregulation of cytokines and

interferons (JAK signalling - Tyrosinekinase2) result in immunodeficiency while defective STAT increases the risk of autoimmunity (O'Shea, Holland, and Staudt 2013; O'Shea et al. 2014). Drugs that inhibit JAK have been shown to be effective in treating several autoimmune diseases by targeting cytokine-dependent pathways, while STAT inhibitors have been promising candidates in the context of cancer (Salas et al. 2020; Pérez-Jeldres et al. 2019; Miklossy, Hilliard, and Turkson 2013). It may therefore be possible to repurpose these existing drugs to improve host tolerance of infection. In summary, our work highlights that Jak/Stat signalling mediates disease tolerance during systemic bacterial infection and that this response differs between males and females. Therefore, we suggest that differences in Jak/Stat mediated disease tolerance may be a potential source of sexually dimorphic response to infection in *Drosophila*.

Acknowledgements:

We thank Ronald P. van Rij and David Finnegan for sharing the fly lines, Vincent Doublet and Emily Robertshaw for laboratory help. We are grateful to Darren Obbard, Jennifer C. Regan, Luke Holman, Amy B. Pedersen, Daniel H. Nussey, Tiina S. Salminen and the Ashworth fly group members for helpful discussion. Finally, we thank Angela Reid, Alison Fulton, Lucinda Rowe and James King for help with media preparation. Figure-1 was created using BioRender.com.

Authors contribution:

A.P and P.F.V conceived the study and designed the experiments; A.P carried out the experimental work with assistance from M.B. and K.M.M; A.P and M.B. acquired the data; A.P analysed the data and drafted the manuscript; A.P & P.F.V wrote the manuscript.

Funding:

We acknowledge funding support to P.F.V from the Branco Weiss fellowship and Chancellor's Fellowship; A.P was supported by the Darwin Trust PhD studentship. We thank School of Biological Sciences, University of Edinburgh. United Kingdom.

Competing interests:

None.

References:

Adelman, James S., Laila Kirkpatrick, Jessica L. Grodio, and Dana M. Hawley. 2013.
"House Finch Populations Differ in Early Inflammatory Signaling and Pathogen
Tolerance at the Peak of *Mycoplasma Gallisepticum* Infection." *The American*
Naturalist 181 (5): 674–89. https://doi.org/10.1086/670024.
Agaisse, Hervé, Ulla-Maja Petersen, Michael Boutros, Bernard Mathey-Prevot, and Norbert
Perrimon. 2003. "Signaling Role of Hemocytes in Drosophila JAK/STAT-Dependent
Response to Septic Injury." *Developmental Cell* 5 (3): 441–50.
https://doi.org/10.1016/S1534-5807(03)00244-2.
Ayres, Janelle S, and David S Schneider. 2012. "Tolerance of Infections," 26.
Babior, Bernard M. 1995. "The Respiratory Burst Oxidase." *Current Opinion in Hematology*,
no. 2: 55–60.
file:///Users/arunprakash/Downloads/The_respiratory_burst_oxidase.8.pdf.
Banerjee, Utpal, Juliet R Girard, Lauren M Goins, and Carrie M Spratford. 2019. "Drosophila
as a Genetic Model for Hematopoiesis." *Genetics* 211 (2): 367–417.
https://doi.org/10.1534/genetics.118.300223.
Bang, In Seok. 2019. "JAK/STAT Signaling in Insect Innate Immunity." *Entomological*
Research 49 (8): 339–53. https://doi.org/10.1111/1748-5967.12384.
Belmonte, Rebecca L., Mary-Kate Corbally, David F. Duneau, and Jennifer C. Regan. 2020.
"Sexual Dimorphisms in Innate Immunity and Responses to Infection in Drosophila
Melanogaster." *Frontiers in Immunology* 10 (January): 3075.
https://doi.org/10.3389/fimmu.2019.03075.
Boon, Adrianus C. M., Jennifer deBeauchamp, Anna Hollmann, Jennifer Luke, Malak Kotb,
Sarah Rowe, David Finkelstein, Lu Lu, Robert W. Williams, and Richard J. Webby.
2009. "Host Genetic Variation Affects Resistance to Infection with a Highly
Pathogenic H5N1 Influenza A Virus in Mice." *Journal of Virology* 83 (20): 10.
https://doi.org/10.1128/JVI.00514-09.
Brivanlou, Ali H., and James E. Darnell. 2002. "Signal Transduction and the Control of Gene
Expression." *Science* 295 (5556): 813–18. https://doi.org/10.1126/science.1066355.
Buchon, Nicolas, Nichole A. Broderick, Takayuki Kuraishi, and Bruno Lemaitre. 2010.
"DrosophilaEGFR Pathway Coordinates Stem Cell Proliferation and Gut Remodeling
Following Infection." *BMC Biology* 8 (1): 152. https://doi.org/10.1186/1741-7007-8-
152.
Buchon, Nicolas, Nichole A. Broderick, Mickael Poidevin, Sylvain Pradervand, and Bruno
Lemaitre. 2009. "Drosophila Intestinal Response to Bacterial Infection: Activation of
Host Defense and Stem Cell Proliferation." *Cell Host & Microbe* 5 (2): 200–211.
https://doi.org/10.1016/j.chom.2009.01.003.
Chakrabarti, Sveta, Jan Paul Dudzic, Xiaoxue Li, Esther Jeanne Collas, Jean-Phillipe
Boquete, and Bruno Lemaitre. 2016. "Remote Control of Intestinal Stem Cell Activity
by Haemocytes in Drosophila." Edited by Utpal Banerjee. *PLOS Genetics* 12 (5):
e1006089. https://doi.org/10.1371/journal.pgen.1006089.

Chakrabarti, Sveta, and Sandhya S. Visweswariah. 2020. "Intramacrophage ROS Primes the Innate Immune System via JAK/STAT and Toll Activation." *Cell Reports* 33 (6): 108368. https://doi.org/10.1016/j.celrep.2020.108368.

Cornet, Stéphane, Coraline Bichet, Stephen Larcombe, Bruno Faivre, and Gabriele Sorci. 2014. "Impact of Host Nutritional Status on Infection Dynamics and Parasite Virulence in a Bird-Malaria System." *Journal of Animal Ecology* 83 (1): 256–65. https://doi.org/10.1111/1365-2656.12113.

Dieppois, Guennaelle, Onya Opota, Jorge Lalucat, and Bruno Lemaitre. 2015. "Pseudomonas Entomophila: A Versatile Bacterium with Entomopathogenic Properties." In *Pseudomonas*, edited by Juan-Luis Ramos, Johanna B. Goldberg, and Alain Filloux, 25–49. Dordrecht: Springer Netherlands. https://doi.org/10.1007/978-94-017-9555-5_2.

Dionne, Marc S., Linh N. Pham, Mimi Shirasu-Hiza, and David S. Schneider. 2006. "Akt and Foxo Dysregulation Contribute to Infection-Induced Wasting in Drosophila." *Current Biology* 16 (20): 1977–85. https://doi.org/10.1016/j.cub.2006.08.052.

Dostert, Catherine, Emmanuelle Jouanguy, Phil Irving, Laurent Troxler, Delphine Galiana-Arnoux, Charles Hetru, Jules A. Hoffmann, and Jean-Luc Imler. 2005. "The Jak-STAT Signaling Pathway Is Required but Not Sufficient for the Antiviral Response of Drosophila." *Nature Immunology* 6 (9): 946–53. https://doi.org/10.1038/ni1237.

Ekengren, Sophia, and Dan Hultmark. 2001. "A Family of Turandot-Related Genes in the Humoral Stress Response of Drosophila." *Biochemical and Biophysical Research Communications* 284 (4): 998–1003. https://doi.org/10.1006/bbrc.2001.5067.

Ekengren, Sophia, Ylva Tryselius, Mitchell S. Dushay, Gang Liu, Håkan Steiner, and Dan Hultmark. 2001. "A Humoral Stress Response in Drosophila." *Current Biology* 11 (9): 714–18. https://doi.org/10.1016/S0960-9822(01)00203-2.

Eleftherianos, Ioannis, Kareen More, Stephanie Spivack, Ethan Paulin, Arman Khojandi, and Sajala Shukla. 2003. "Nitric Oxide Levels Regulate the Immune Response of Drosophila Melanogaster Reference Laboratory Strains to Bacterial Infections." https://doi.org/10.1128/IAI.02318-14 •.

Ferrandon, Dominique, Jean-Luc Imler, Charles Hetru, and Jules A. Hoffmann. 2007. "The Drosophila Systemic Immune Response: Sensing and Signalling during Bacterial and Fungal Infections." *Nature Reviews Immunology* 7 (11): 862–74. https://doi.org/10.1038/nri2194.

Ferreira, Ana, Ivo Marguti, Ingo Bechmann, Viktória Jeney, Ângelo Chora, Nuno R. Palha, Sofia Rebelo, Annie Henri, Yves Beuzard, and Miguel P. Soares. 2011. "Sickle Hemoglobin Confers Tolerance to Plasmodium Infection." *Cell* 145 (3): 398–409. https://doi.org/10.1016/j.cell.2011.03.049.

Ganz, Holly H., and Dieter Ebert. 2010. "Benefits of Host Genetic Diversity for Resistance to Infection Depend on Parasite Diversity." *Ecology* 91 (5): 1263–68. https://doi.org/10.1890/09-1243.1.

Gilbert, M. Melissa, Brian K. Weaver, J. Peter Gergen, and Nancy C. Reich. 2005. "A Novel Functional Activator of the Drosophila JAK/STAT Pathway, Unpaired2, Is Revealed by an in Vivo Reporter of Pathway Activation." *Mechanisms of Development* 122 (7): 939–48. https://doi.org/10.1016/j.mod.2005.03.004.

Gupta, Vanika, and Pedro F. Vale. 2017. "Nonlinear Disease Tolerance Curves Reveal Distinct Components of Host Responses to Viral Infection." *Royal Society Open Science* 4 (7): 170342. https://doi.org/10.1098/rsos.170342.

Harrison, Douglas A. 2012. "The JAK/STAT Pathway." *Cold Spring Harbor Perspectives in Biology* 4 (3): a011205. https://doi.org/10.1101/cshperspect.a011205.

Harrison, Douglas A., Patricia E. McCoon, Richard Binari, Michael Gilman, and Norbert Perrimon. 1998. "Drosophila Unpaired Encodes a Secreted Protein That Activates the JAK Signaling Pathway." *Genes & Development* 12 (20): 3252–63. https://doi.org/10.1101/gad.12.20.3252.

Hoffmann, Jules A. 2003. "The Immune Response of Drosophila." *Nature* 426 (6962): 33–38. https://doi.org/10.1038/nature02021.

Howick, Virginia M., and Brian P. Lazzaro. 2014. "Genotype and Diet Shape Resistance and Tolerance across Distinct Phases of Bacterial Infection." *BMC Evolutionary Biology* 14 (1): 56. https://doi.org/10.1186/1471-2148-14-56.

Hultmark, Dan. 2003. "Drosophila Immunity: Paths and Patterns." *Current Opinion in Immunology* 15 (1): 12–19. https://doi.org/10.1016/S0952-7915(02)00005-5.

Jamieson, Amanda M., Lesley Pasman, Shuang Yu, Pia Gamradt, Robert J. Homer, Thomas Decker, and Ruslan Medzhitov. 2013. "Role of Tissue Protection in Lethal Respiratory Viral-Bacterial Coinfection." *Science* 340 (6137): 1230–34. https://doi.org/10.1126/science.1233632.

Kemp, Cordula, Stefanie Mueller, Akira Goto, Vincent Barbier, Simona Paro, François Bonnay, Catherine Dostert, et al. 2013. "Broad RNA Interference–Mediated Antiviral Immunity and Virus-Specific Inducible Responses in Drosophila." *The Journal of Immunology* 190 (2): 650–58. https://doi.org/10.4049/jimmunol.1102486.

Khalil, Sarah, Eliana Jacobson, Moria C. Chambers, and Brian P. Lazzaro. 2015. "Systemic Bacterial Infection and Immune Defense Phenotypes in Drosophila Melanogaster." *Journal of Visualized Experiments : JoVE*, no. 99 (May): 52613. https://doi.org/10.3791/52613.

Kiu, Hiu, and Sandra E. Nicholson. 2012. "Biology and Significance of the JAK/STAT Signalling Pathways." *Growth Factors* 30 (2): 88–106. https://doi.org/10.3109/08977194.2012.660936.

Klebanoff, Seymour J. 1974. "Role of the Superoxide Anion in the Myeloperoxidase-Mediated Antimicrobial System." *The Journal of Biological Chemistry* 249 No. 12 (June 25): 3724–28.

Klein, Sabra L., and Katie L. Flanagan. 2016. "Sex Differences in Immune Responses." *Nature Reviews Immunology* 16 (10): 626–38. https://doi.org/10.1038/nri.2016.90.

Kleino, Anni, and Neal Silverman. 2014. "The Drosophila IMD Pathway in the Activation of the Humoral Immune Response." *Developmental & Comparative Immunology* 42 (1): 25–35. https://doi.org/10.1016/j.dci.2013.05.014.

Lamiable, Olivier, and Jean-Luc Imler. 2014. "Induced Antiviral Innate Immunity in Drosophila." *Current Opinion in Microbiology*, Host–microbe interactions: fungi/parasites/viruses, 20 (August): 62–68. https://doi.org/10.1016/j.mib.2014.05.006.

Lazzaro, Brian P, Timothy B Sackton, and Andrew G Clark. 2006. "Genetic Variation in Drosophila Melanogaster Resistance to Infection: A Comparison Across Bacteria." *Genetics* 174 (3): 1539–54. https://doi.org/10.1534/genetics.105.054593.

Lee, Won-Jae, and Sung-Hee Kim. 2014. "Role of DUOX in Gut Inflammation: Lessons from Drosophila Model of Gut-Microbiota Interactions." *Frontiers in Cellular and Infection Microbiology* 3: 116. https://doi.org/10.3389/fcimb.2013.00116.

Lemaitre, Bruno, and Jules Hoffmann. 2007. "The Host Defense of *Drosophila Melanogaster*." *Annual Review of Immunology* 25 (1): 697–743. https://doi.org/10.1146/annurev.immunol.25.022106.141615.

Lissner, Michelle M, and David S Schneider. 2018. "The Physiological Basis of Disease Tolerance in Insects." *Current Opinion in Insect Science*, Global change biology *

Molecular physiology, 29 (October): 133–36.
https://doi.org/10.1016/j.cois.2018.09.004.

Louie, Alexander, Kyung Han Song, Alejandra Hotson, Ann Thomas Tate, and David S.
Schneider. 2016. "How Many Parameters Does It Take to Describe Disease
Tolerance?" Edited by Andy P. Dobson. *PLOS Biology* 14 (4): e1002435.
https://doi.org/10.1371/journal.pbio.1002435.

Martins, Rui, Ana Rita Carlos, Faouzi Braza, Jessica A. Thompson, Patricia Bastos-Amador,
Susana Ramos, and Miguel P. Soares. 2019. "Disease Tolerance as an Inherent
Component of Immunity." *Annual Review of Immunology* 37 (1): 405–37.
https://doi.org/10.1146/annurev-immunol-042718-041739.

McCarville, JL, and JS Ayres. 2018. "Disease Tolerance: Concept and Mechanisms."
Current Opinion in Immunology, Innate immunity 2018, 50 (February): 88–93.
https://doi.org/10.1016/j.coi.2017.12.003.

McKean, Kurt A., Christopher P. Yourth, Brian P. Lazzaro, and Andrew G. Clark. 2008.
"The Evolutionary Costs of Immunological Maintenance and Deployment." *BMC
Evolutionary Biology* 8 (1): 76. https://doi.org/10.1186/1471-2148-8-76.

Medzhitov, Ruslan, David S. Schneider, and Miguel P. Soares. 2012. "Disease Tolerance as a
Defense Strategy." *Science* 335 (6071): 936–41.
https://doi.org/10.1126/science.1214935.

Meister, Marie, and Marie Lagueux. 2003. "Drosophila Blood Cells." *Cellular Microbiology*
5 (9): 573–80. https://doi.org/10.1046/j.1462-5822.2003.00302.x.

Merkling, Sarah H., Alfred W. Bronkhorst, Jamie M. Kramer, Gijs J. Overheul, Annette
Schenck, and Ronald P. Van Rij. 2015. "The Epigenetic Regulator G9a Mediates
Tolerance to RNA Virus Infection in Drosophila." *PLOS Pathogens* 11 (4):
e1004692. https://doi.org/10.1371/journal.ppat.1004692.

Miklossy, Gabriella, Tyvette S. Hilliard, and James Turkson. 2013. "Therapeutic Modulators
of STAT Signalling for Human Diseases." *Nature Reviews Drug Discovery* 12 (8):
611–29. https://doi.org/10.1038/nrd4088.

Myllymäki, H., and M. Rämet. 2014. "JAK/STAT Pathway in Drosophila Immunity."
Scandinavian Journal of Immunology 79 (6): 377–85.
https://doi.org/10.1111/sji.12170.

Myllymäki, Henna, Susanna Valanne, and Mika Rämet. 2014. "The Drosophila Imd
Signaling Pathway." *The Journal of Immunology* 192 (8): 3455–62.
https://doi.org/10.4049/jimmunol.1303309.

Nystrand, M., and D. K. Dowling. 2020. "Effects of Immune Challenge on Expression of
Life-History and Immune Trait Expression in Sexually Reproducing Metazoans—a
Meta-Analysis." *BMC Biology* 18 (1): 135. https://doi.org/10.1186/s12915-020-
00856-7.

Ohlstein, Benjamin, and Allan Spradling. 2006. "The Adult Drosophila Posterior Midgut Is
Maintained by Pluripotent Stem Cells." *Nature* 439 (7075): 470–74.
https://doi.org/10.1038/nature04333.

Oliveira, José Henrique, Ana Cristina Bahia, and Pedro F. Vale. 2020. "How Are Arbovirus
Vectors Able to Tolerate Infection?" *Developmental & Comparative Immunology* 103
(February): 103514. https://doi.org/10.1016/j.dci.2019.103514.

O'Shea, John J., Steven M. Holland, and Louis M. Staudt. 2013. "JAKs and STATs in
Immunity, Immunodeficiency, and Cancer." *New England Journal of Medicine* 368
(2): 161–70. https://doi.org/10.1056/NEJMra1202117.

O'Shea, John J, Daniella M Schwartz, Alejandro V Villarino, Massimo Gadina, Iain B
McInnes, and Arian Laurence. 2014. "The JAK-STAT Pathway: Impact on Human
Disease and Therapeutic Intervention," 21.

Palmer, William H., Finny S. Varghese, and Ronald P. Van Rij. 2018. "Natural Variation in
 Resistance to Virus Infection in Dipteran Insects." *Viruses* 10 (3): 118.
 https://doi.org/10.3390/v10030118.

Pérez-Jeldres, Tamara, Christopher J. Tyler, Joshua D. Boyer, Thangaraj Karuppuchamy,
 Andres Yarur, Daniel A. Giles, Shaila Yeasmin, et al. 2019. "Targeting Cytokine
 Signaling and Lymphocyte Traffic via Small Molecules in Inflammatory Bowel
 Disease: JAK Inhibitors and S1PR Agonists."
 https://doi.org/10.3389/fphar.2019.00212.

Raberg, L., D. Sim, and A. F. Read. 2007. "Disentangling Genetic Variation for Resistance
 and Tolerance to Infectious Diseases in Animals." *Science* 318 (5851): 812–14.
 https://doi.org/10.1126/science.1148526.

Regan, Jennifer C, Mobina Khericha, Adam J Dobson, Ekin Bolukbasi, Nattaphong
 Rattanavirotkul, and Linda Partridge. 2016. "Sex Difference in Pathology of the
 Ageing Gut Mediates the Greater Response of Female Lifespan to Dietary
 Restriction." *ELife* 5 (February): e10956. https://doi.org/10.7554/eLife.10956.

Salas, Azucena, Cristian Hernandez-Rocha, Marjolijn Duijvestein, William Faubion, Dermot
 McGovern, Severine Vermeire, Stefania Vetrano, and Niels Vande Casteele. 2020.
 "JAK–STAT Pathway Targeting for the Treatment of Inflammatory Bowel Disease."
 Nature Reviews Gastroenterology & Hepatology 17 (6): 323–37.
 https://doi.org/10.1038/s41575-020-0273-0.

Schneider, David S., and Janelle S. Ayres. 2008. "Two Ways to Survive Infection: What
 Resistance and Tolerance Can Teach Us about Treating Infectious Diseases." *Nature
 Reviews Immunology* 8 (11): 889–95. https://doi.org/10.1038/nri2432.

Schwenke, Robin A., Brian P. Lazzaro, and Mariana F. Wolfner. 2016. "Reproduction–
 Immunity Trade-Offs in Insects." *Annual Review of Entomology* 61 (1): 239–56.
 https://doi.org/10.1146/annurev-ento-010715-023924.

Sears, Brittany F., Jason R. Rohr, Judith E. Allen, and Lynn B. Martin. 2011. "The Economy
 of Inflammation: When Is Less More?" *Trends in Parasitology* 27 (9): 382–87.
 https://doi.org/10.1016/j.pt.2011.05.004.

Siva-Jothy, Jonathon A., Arun Prakash, Radhakrishnan B. Vasanthakrishnan, Katy M.
 Monteith, and Pedro F. Vale. 2018. "Oral Bacterial Infection and Shedding in
 Drosophila Melanogaster." *Journal of Visualized Experiments*, no. 135 (May): 57676.
 https://doi.org/10.3791/57676.

Soares, Miguel P., Raffaella Gozzelino, and Sebastian Weis. 2014. "Tissue Damage Control
 in Disease Tolerance." *Trends in Immunology* 35 (10): 483–94.
 https://doi.org/10.1016/j.it.2014.08.001.

Soares, Miguel P., Luis Teixeira, and Luis F. Moita. 2017. "Disease Tolerance and Immunity
 in Host Protection against Infection." *Nature Reviews Immunology* 17 (2): 83–96.
 https://doi.org/10.1038/nri.2016.136.

Stark, George R., and James E. Darnell. 2012. "The JAK-STAT Pathway at Twenty."
 Immunity 36 (4): 503–14. https://doi.org/10.1016/j.immuni.2012.03.013.

Tafesh-Edwards, Ghada, and Ioannis Eleftherianos. 2020. "JNK Signaling in Drosophila
 Immunity and Homeostasis." *Immunology Letters* 226 (October): 7–11.
 https://doi.org/10.1016/j.imlet.2020.06.017.

Troha, Katia, Joo Hyun Im, Jonathan Revah, Brian P. Lazzaro, and Nicolas Buchon. 2018.
 "Comparative Transcriptomics Reveals CrebA as a Novel Regulator of Infection
 Tolerance in D. Melanogaster." *PLOS Pathogens* 14 (2): e1006847.
 https://doi.org/10.1371/journal.ppat.1006847.

Valanne, Susanna, Jing-Huan Wang, and Mika Rämet. 2011. "The Drosophila Toll Signaling
Pathway." *The Journal of Immunology* 186 (2): 649–56.
https://doi.org/10.4049/jimmunol.1002302.
Vale, Pedro F., Andy Fenton, and Sam P. Brown. 2014. "Limiting Damage during Infection:
Lessons from Infection Tolerance for Novel Therapeutics." *PLoS Biology* 12 (1):
e1001769. https://doi.org/10.1371/journal.pbio.1001769.
Vale, Pedro F., Luke McNally, Andrea Doeschl-Wilson, Kayla C. King, Roman Popat, Maria
R. Domingo-Sananes, Judith E. Allen, Miguel P. Soares, and Rolf Kümmerli. 2016.
"Beyond Killing: Can We Find New Ways to Manage Infection?" *Evolution,*
Medicine, and Public Health 2016 (1): 148–57.
https://doi.org/10.1093/emph/eow012.
Vincent, Crystal M., and Marc S. Dionne. 2021. "Disparate Regulation of IMD Signaling
Drives Sex Differences in Infection Pathology in *Drosophila Melanogaster*."
Proceedings of the National Academy of Sciences 118 (32): e2026554118.
https://doi.org/10.1073/pnas.2026554118.
Vodovar, N., M. Vinals, P. Liehl, A. Basset, J. Degrouard, P. Spellman, F. Boccard, and B.
Lemaitre. 2005. "Drosophila Host Defense after Oral Infection by an
Entomopathogenic Pseudomonas Species." *Proceedings of the National Academy of*
Sciences 102 (32): 11414–19. https://doi.org/10.1073/pnas.0502240102.
Wang, Jonathan B., Hsiao-Ling Lu, and Raymond J. St Leger. 2017. "The Genetic Basis for
Variation in Resistance to Infection in the Drosophila Melanogaster Genetic
Reference Panel." *PLOS Pathogens* 13 (3): e1006260.
https://doi.org/10.1371/journal.ppat.1006260.
Zhou, Feng, and Hervé Agaisse. 2012. "JAK/STAT Signaling and Invertebrate Immune
Responses." In *Jak-Stat Signaling : From Basics to Disease*, edited by Thomas
Decker and Mathias Müller, 133–51. Vienna: Springer. https://doi.org/10.1007/978-3-
7091-0891-8_9.
667
668

        Supplementary Information

***Figure S1.* (A)** Survival curves for males and females of wildtype *yw* and *Outcross* population
exposed to systemic *P. entomophila* of infection dose OD_{600}=0.05 (n=20-30 flies/vial (~15 vials)/fly
line/treatment/sex/infection dose). **(B)** Estimated hazard ratios calculated from the survival
curves for males and female flies (*Outcross* and *yw* flies). A greater hazard ratio (>1) indicates higher
susceptibility to bacterial infection. ['n.s' in the panel indicates that *Outcross* flies are not significantly
different from *yw* flies]. **(C)** Internal bacterial load (as mean CFUs- colony forming units) measured
around 24 hours post systemic *P. entomophila* infection. [significantly different fly lines are
connected with different alphabets using Tukey's HSD as a post hoc analysis of pairwise
comparisons].

Both wildtype (*yw*) & *Outcrossed* flies show similar response to systemic *P. entomophila*
infection: All our *Jak/Stat*-pathway mutants were in *yw* genetic background. Previous work has
shown that *yw* flies have lower basal levels of nitric oxide and are more susceptible to infection
compared to other lab wild types such as *w^1118, Oregon R, Canton-S* (*Eleftherianos et al., 2014*). Before
addressing whether the *Jak/Stat* pathway is involved in disease tolerance of bacterial infection, we
wanted to better understand how wild type (*yw*) flies survive *P. entomophila* infections relative to
other wild-type strains. We used an *Outcrossed* fly population as another control for comparison,
which was originally created using 100 pairwise crosses of 113 *DRGP* lines and maintained as an
outcrossed population for over 19 generations (see *Savola et al., 2021* for generation of *DGRP*
Outcrossed population). We found that males and females of both wild type *yw* and the *Outcross*ed
flies showed comparable survival following *P. entomophila* systemic infection and exhibited similar
levels of internal bacterial loads when measured ~24h post-infection (***Fig.* *S1A & S1B, Table *S1***
for survival; ***Fig.* *S1C,** for bacterial load). Since both the wildtype (*yw*) and *Outcross* fly lines showed
similar infection responses to *P. entomophila*, all remaining experiments only used *yw* as the wildtype
control line.

Table S1. Summary of mixed effects Cox model, for wildtype *yw* & flies without *Jak/Stat* pathway components during systemic *P. entomophila* infection in. We used data from the individuals of 3-day adult males and females infected with OD_{600} = 0.05 infection dose of *P. entomophila* for each fly lines (*Jak/Stat* -pathway mutants) and specified the model as: survival ~ treatment x sex x fly line (1|vial), with 'treatment', 'sex' & 'fly line' as fixed effects, and 'vials' as a random effect. The table shows model output (ANOVA) for survival post-infection for flies with fully function immune system and lacking *Jak/Stat* - pathway.

	Source	Log lik.	Chi sq.	df	p
Wildtype vs. outcross	Treatment (SI vs I)	-5956.3	500.18	1	<0.001
	Sex	-5953.8	5.0844	1	0.38
	Fly line	-5945.8	15.929	1	0.54
	Treatment x sex	-5945.7	0.3030	1	0.58
	Treatment x fly line	-5945.7	0.0077	1	0.93
	Sex x fly line	-5945.3	0.7848	1	0.37
	Treatment x sex x fly line	-5942.8	4.9490	1	0.02
	Random effect	*Std dev*			
	vial	*0.61*			

Table S2. Summary of mixed effects Cox model, for each fly line, wildtype *yw* & flies without *Jak/Stat* pathway mutants. We used data from the individuals of 3-day adult males and females infected with OD_{600} = 0.05 infection dose of *P. entomophila* for each fly lines (*Jak/Stat* -pathway mutants) and specified the model as: survival ~ treatment x sex x fly line (1|vial), with 'treatment', 'sex' & 'fly line' as fixed effects, and 'vials' as a random effect. The table shows model output (ANOVA) for survival post-infection for flies with fully function immune system (*yw*) and flies lacking *Jak/Stat* – pathway components.

	Source	Log lik.	Chi sq.	df	p
Jak/Stat mutants	Treatment	-30378	1042.2	1	<0.001
	Sex	-30357	42.39	1	<0.001
	Fly line	-29746	1222.9	4	<0.001
	Treatment x sex	-29746	0.050	1	0.82
	Treatment x fly line	-29744	2.732	4	0.60
	Sex x flyline	-29736	15.57	4	0.003
	Treatment x fly line x sex	-29734	4.506	4	0.34
	Random effect	*Std dev*			
	vial/block	*0.49*			

716 ***Table S3.*** Summary of ln$_{10}$ transformed bacterial load data around 24 hours following
717 OD$_{600}$=0.05 *P. entomophila* systemic infection for wildtype (*yw)* flies & *Jak/Stat* pathway mutants,
718 analysed using non-parametric one-way ANOVA (K-W test) by fitting 'fly line' (*yw* and Jak/Stat
719 mutants) as categorical fixed-effects for males and females separately.
720

K-W	Sex	Source	DF	Chi Sq.	P
Jak/Stat	Female	Fly line	4	9.1704	0.057
mutants	Male	Fly line	4	9.4565	0.0506

Table S4. Summary of mixed effects Cox model, for G9a *Jak/Stat* pathway components during
systemic *P. entomophila* infection in. We used data from the individuals of 3-day adult males and
females infected with OD$_{600}$ = 0.05 infection dose of *P. entomophila* for each fly lines (*G9a*) and
specified the model as: survival ~ treatment x sex x fly line (1|vial), with 'treatment', 'sex' & 'fly
line' as fixed effects, and 'vials' as a random effect. The table shows model output (ANOVA) for
survival post-infection for flies with fully function immune system and lacking *G9a*.

	Source	Log lik.	Chi sq.	df	p
G9a	Treatment	-8485.2	310.1	1	<0.001
	Sex	-8481.2	8.07	1	0.004
	Fly line	-8436.6	89.1	1	<0.001
	Treatment x sex	-8436.5	0.22	1	0.63
	Treatment x fly line	-8433.9	5.13	1	0.002
	Sex x fly line	-8426.3	15.2	1	<0.001
	Treatment x sex x fly line	-8426.3	0.00	1	1.0
	Random effect	*Std dev*			
	vial/block	*0.25*			

729
730
731
732
733 ***Table S5.*** Summary of log$_{10}$ transformed bacterial load data around 24 hours following OD$_{600}$=0.05
734 *P. entomophila* systemic infection for *G9a* mutants, analysed using non-parametric one-way
735 ANOVA (K-W test) by fitting 'Fly line' as categorical fixed-effects for males and females
736 separately.

Fly line	Sex	S	Z	p
G9a	Female	641.5	3.065	0.002
	Male	548.5	1.479	0.139

737
738