Local adaptation to novel habitats and climatic changes in a wild tomato species via selective sweeps

Kai Wei *, Gustavo A Silva-Arias *+, Aurélien Tellier

Professorship for Population Genetics, Department of Life Science Systems, School of Life Sciences, Technical University of Munich, Liesel-Beckmann Strasse 2, 85354 Freising, Germany

* these authors contributed equally

+Corresponding author: Gustavo A Silva-Arias: gustavo.silva@tum.de

Word count

        Main body: 6165
        Introduction: 1172
        Materials and Methods: 1233
        Results: 2879
        Discussion: 881

5 figures (all in colour)

2 tables

3 supporting information files

Summary

● Positive Darwinian selection is the driving force underpinning local adaptation,
and leaves footprints of selective sweeps at the underlying major genes.
Quantifying the timing of selection and revealing the genetic bases of local
adaptation in plants species occurring in steep and varying environmental
gradients is crucial to predict a species' ability to respond to climate changes
and explore new niches.

● We use whole genome sequence data from six populations across three
different habitats of the wild tomato species *Solanum chilense*, to 1) infer the
past demographic history, and 2) search for genes under strong positive
selection. We then correlate current and past climatic projections with the
demographic history, allele frequencies in space and time, the age of
selection events, and changes in habitat suitability.

● We find evidence for several selective sweeps targeting regulatory networks
involved in root hair development in low altitude, and response to photoperiod
and vernalization in high altitude populations. These sweeps occur in a
concerted fashion in a given regulatory gene network at particular periods of
substantial climatic change.

● We decipher the genetic bases and the timing of local adaptation during plant
colonization of semi-arid habitats using a unique combination of genome
scans for selection and modelling of past climatic data.

Keywords:

Local adaptation, selective sweeps, age of sweeps, gene network, *Solanum chilense*, climate change

Introduction

Positive Darwinian selection is a key process underlying local adaptation, the process by which populations of a species evolve in response to geographically variable selection (Kawecki & Ebert, 2004). Spatial local adaptation results from the spatial heterogeneity of habitats promoting different selective pressures across populations, especially after species range expansion (Kawecki & Ebert, 2004; Savolainen *et al.*, 2013; Tiffin & Ross-Ibarra, 2014). Temporal local adaptation results from abiotic/biotic conditions slowly varying over time (over hundreds up to thousands of years) at a given location and differently across populations (Polechová *et al.*, 2009). In order to predict the future adaptive potential of plants in the face of climatic change, it is desirable to assess 1) how many genes and gene networks underpin spatial and temporal local adaptation, 2) what is the timing (time period) of local adaptation on a given trait, and 3) if selection at a given gene can be linked to past changes in climatic conditions.

Population genetics theory investigates the characteristics and polymorphism footprints of both types of polygenic local adaptation: spatial adaptation to the new local fixed optimum of the colonized habitat (Chevin *et al.*, 2010), and adaptation to track the moving environmental optimum (Polechová *et al.*, 2009; Matuszewski *et al.*, 2014; Jain & Stephan, 2017a). Under large enough population sizes, both models predict that more significant steps of local adaptation occur first at sites with strong selective coefficients, possibly generating selective sweeps (Chevin *et al.*, 2010; Matuszewski *et al.*, 2014; Jain & Stephan, 2017a). Selective sweeps are polymorphism patterns (footprints) in the genome due to the rapid (tens to hundreds of generations) fixation of advantageous alleles and the associated hitchhiking effect (Smith & Haigh, 1974; Kim & Stephan, 2002). Population genomics studies often assume implicitly, as an approximation, that a gene under strong selection (selective sweep) in one population but not in another is indicative of potential involvement in

local adaptation (see discussion in Tiffin and Ross-Ibarra 2014). A rigorous

demonstration requires validating the fitness effect of that gene in common garden

experiments (Kawecki & Ebert, 2004). Thus, complete studies to uncover the genetic

bases of spatial adaptation have been conducted in animals such as humans or

invertebrates (*Caenorhabditis elegans, Drosophila melanogaster*) but also in few

amenable plant species (*Arabidopsis* spp., *Arabis* sp., *Capsella* spp., *Helianthus* spp.

Senecio lautus, *Mimulus* spp.) in which the molecular bases for adaptation can be

studied in exquisite details using Genome Wide Association studies and functional

studies (review in *e.g.* Savolainen *et al.*, 2013; Josephs *et al.*, 2017; Fagny &

Austerlitz, 2021). Numerous population genomic studies of spatial local adaptation

are also conducted in non-model plant species, albeit with lower resolution regarding

the underlying molecular bases (Pannell & Fields, 2014; Bragg *et al.*, 2015).

Not surprisingly, we note that most spatial local adaptation is due to recent

selective sweeps estimated to be less than approximately 25,000 years old (*e.g.*

(Wright *et al.*, 2015; Schrider & Kern, 2017; Zou *et al.*, 2017; Garud *et al.*, 2021), and

often associated with either strong bottlenecks during colonization events, recent

(re)colonization processes, or selection on standing genetic variation. While recent

sweeps underpin spatial local adaptation, we note in the literature the absence of

reported older sweeps underpinning adaptation to past climatic changes (local

temporal adaptation). This is explained by several methodological and theoretical

limitations to detect loci under strong positive selection (Lotterhos & Whitlock, 2015;

Hoban *et al.*, 2016). First, there is a confounding effect of the past demographic

history of the populations (Stephan, 2016) especially under strong bottlenecks due to

the colonization of new habitats (François *et al.*, 2016; Hoban *et al.*, 2016). Indeed,

under strong demographic bottlenecks, it is notoriously difficult to carefully control the

statistical rate of false-positive sweeps and sweep signatures pre-dating the

bottleneck are erased (Stephan, 2016; Hoban *et al.*, 2016; Schrider & Kern, 2017;

Garud *et al.*, 2021). Second, strong selection at few genes is only the tip of the

iceberg, as adaptation likely occurs for polygenic traits with selection acting over

many genes. Different number of major genes with selective sweeps signatures can

be observed depending on the distribution of selection coefficients, the efficiency of

selection (a function of effective population size and recombination rate), the

architecture of the traits, place of genes in gene networks/pathways, and gene

pleiotropy (Jain & Stephan, 2017b; Barghi *et al.*, 2020). Third, theory predicts that

strong selective sweeps can be observed and detected reliably only for ca. $0.1Ne$

generations after selection, where Ne is the effective population size (Kim & Stephan,

2002). As a result of these three factors, in plant species undergoing habitat range

expansion with strong bottleneck during colonization and life-history traits decreasing

effective population size (*e.g.* selfing), we expect 1) only few sweeps to occur under

polygenic selection, and 2) only recent sweeps to be detected using polymorphism

data. In order to potentially recover and infer older selective events of local temporal

and spatial adaptation, the study of alternative outcrossing plant species with larger

effective population sizes during colonization events is required.

Solanum chilense is an outcrossing wild tomato species found in southern Peru

and northern Chile in mesic to very arid habitats (Nakazato *et al.*, 2010), and its

ancestral range is likely in marginal desert habitat of the coast and mid-altitude 'pre-

cordillera' region (800 - 2000 m altitude) of southern Peru. *S. chilense* colonized

independently two different southern, but arid, isolated regions around the Atacama

desert at different time points (Böndel *et al.*, 2015; Stam *et al.*, 2019b): an early

divergence (older than 50,000 years ago [ya]) with the colonization of coastal

habitats (in Lomas formations), and a more recent lineage divergence (less than

20,000 ya) restricted to highland altitudes (above 2,400m) of the Andean plateau.

Signatures of natural selection (positive or balancing) at genes involved in stress

adaptation were found when scanning few candidate genes for biotic and abiotic

stress response (Xia *et al.*, 2010; Fischer *et al.*, 2011; Böndel *et al.*, 2015; Nosenko

et al., 2016; Böndel *et al.*, 2018; Stam *et al.*, 2019b). In the present study, we obtain

full genome sequence data for 30 diploid and highly heterozygous plants from six

populations representing the three main habitats of the species (Fig. 1a, defined in

Böndel et al. 2015): the central group (area of origin at low to high altitude), south-

coastal group, and south-highland group. The south-highland group strongly differs

from the central group for its current climatic conditions, while the south-coastal

appears as only marginally different from the environment prevailing in the central

group (Fig. 1b). Our aims are first to infer accurately the past demographic history of

the species colonization and to reconstruct recent dynamics of the species'

distribution range in response to climatic history. Second, we conduct genome scans

for selective sweeps and assign functions and gene network topology to these

candidate genes. Third, we link climatic and genetic data at candidate genes using

genotype-environment association analysis to highlight the relevance of key gene

regulatory networks (pathways) for local adaptation, and discuss the history of spatial

and temporal local adaptation in *S. chilense*.

Materials and Methods

Sample collection, sequencing and bioinformatics

Plants were grown from individual seeds obtained from the Tomato Genetics

Resource Center (TGRC, University of California, Davis, USA). We sampled five

diploid plants from accessions LA1963, LA3111, LA2931, LA4330, LA2932, LA4107

(Fig. 1a; Table S1) grown in standard glasshouse conditions. Genomic DNA was

extracted using the DNA extraction kit from Quiagen and sequenced on an Illumina

HiSeq 2500 with standard library size of 300 bp (Eurofins Genomics, Germany). The

30 *S. chilense* whole-genome sequencing data are available on ENA in BioProject

PRJEB47577.

We performed quality control of the raw reads and trimmed calls with

insufficient quality or adapter contamination. The clean reads were mapped to the

Solanum pennellii reference genome (Bolger *et al.*, 2014) available from Solanaceae

Genomics Network using the Burrows-Wheeler Alignment tool (v0.7.16) with default

settings (Li & Durbin, 2009) and sorted with Samtools (v1.5) (Wysoker *et al.*, 2009).

The raw alignments were then processed to add read groups, mark duplicates and fix

mates. Variant calling was performed using the HaplotypeCaller tool of GATK

(McKenna *et al.*, 2010) (Auwera et al. 2013)with default parameters for each sample.

Individual genomic variant files were then combined into a variant matrix with the

GenotypeGVCFs tool and annotated based on the gene annotation of the *S.pennellii*

reference (details in the Supplementary SI text 1).

Population genetics analyses and inference of demographic history

For all population genetics analyses we used *S. pennellii* population LA716 as

outgroup. We built a maximum likelihood (ML) phylogenetic tree and performed a

principal component analysis (PCA) and the inference of population structure with

ADMIXTURE (Alexander *et al.*, 2009). Population genetics statistics, nucleotide

diversity (π), Tajima's D and F_{ST}, were calculated with VCFtools (v1.15) (Danecek *et*

al., 2011) for each population (or pairs of populations) over 100 kb sliding non-

overlapping windows. The linkage disequilibrium (LD) levels were calculated per

population as the genotype correlation coefficient (r^2) between two loci using

VCFtools with a maximum distance of 1,000 kb.

The demographic inference was conducted using the Multiple Sequentially

Markovian Coalescent method (MSMC2) with phased VCF files and 40 hidden states

(Malaspinas *et al.*, 2016). The cross-coalescence analysis was performed for each

pairwise comparison of genomes between pairs of populations to estimate the

population separation history and the migration rate with MSMC-IM (Wang *et al.*,

2020). The phased data were generated with SHAPEIT (v2.17) (Delaneau *et al.*,

2012). We assumed generation time of 5 years (uncertainty interval 3-7) and a

mutation rate per generation of 1×10^{-8} (uncertainty interval 5.1×10^{-9} – 2.5×10^{-8},

based on Roselius *et al.*, 2005), accounting for ambiguity in these estimates.

Modelling present and past species distribution

We reconstructed and visualized the environmental space occupied by *S. chilense*

extracting the environmental conditions at the current occurrence points and

summarize them by PCA (Fig. 1b) (Legendre & Legendre, 2012). The environmental

data include 63 climatic layers obtained from three public databases WorldClim2

(Fick & Hijmans, 2017), ENVIREM (Title & Bemmels, 2018), and the Consultative

Group on International Agricultural Research (CGIAR) (Trabucco & Zomer, 2018)

(Dataset S5). The PCA was performed by the *prcomp* function in R (R Core Team,

2020).

We then performed an ensemble modelling framework (Araujo & New, 2007)

using the BIOMOD package (Thuiller *et al.*, 2009, 2014) in R, including all known

localities covering the entire geographic range of *S. chilense* using eight modelling

algorithms, five cross-validation replicates, and ten pseudo-absence sampling sets,

therefore completing a total of 400 models. Consensus niche models were obtained

using a TSS-weighted average method to account for the predictive power of each

fitted model. Models with low predictive power (TSS < 0.7) were discarded. All fitted

suitability models were then projected to infer the distribution suitable habitat of *S.*

chilense under current climatic conditions and during the Last Glacial Maximum

(LGM; ~21 Kya).

Genome-wide selection scans and statistical power

We identified selective sweeps using biallelic SNPs by SweeD (Pavlidis *et al.*, 2013)

and OmegaPlus (Alachiotis *et al.*, 2012). The CLR statistics in SweeD were

calculated with default parameters with 10 kb intervals. OmegaPlus statistics (ω)

were computed at 10 kb intervals. We specified a minimum window of 10 kb and a

maximum window of 100 kb to be used for computing LD values between SNPs,

respectively. Outlier CLR and ω statistics indicative of a selective sweep are defined

by comparison to the genome-wide distribution values and the thresholds are defined

by coalescent simulations of the inferred demographic history. We used the

coalescent simulator SCRM (Staab *et al.*, 2015) to generate 10,000 neutral datasets

of 10 Mb based on the demographic history of each population and assuming a

varying recombination rate every 100 kb within each 10 Mb simulated block

(recombination rate varied between $0.1*\theta$ and $10*\theta$). Using the genomic coordinates,

we then extracted the overlap regions between the two methods, which are regarded

as high confident selective sweep regions. As independent confirmation of the sweep

regions, we used McSwan (Tournebize *et al.*, 2019) to detect sweeps and estimate

their age. McSwan was run with the same parameters as SweeD.

To evaluate the sensitivity of our sweeps detection pipeline, we simulated

1,000 selective sweeps per selective coefficient value (*s*=0.01, 0.1, 0.5, 1) for each of

the six populations under the inferred demographic model (with a random sweep age

distribution). We used the *generate_pseudoobs* function based on MSMS simulator

implemented in the McSwan R package (Ewing & Hermisson, 2010; Tournebize *et*

al., 2019). We then ran SweeD, OmegaPlus and McSwan on all simulated data sets

using the same parameters and thresholds defined above to quantify the percentage

of sweeps detected per population (and age of the detected sweeps).

GO enrichment analysis and gene networks

Due to the lack of a complete gene function annotation database, we performed a

BLASTX against the NCBI database of non-redundant proteins (nr) screened for

green plants (e-value cutoff was 10^{-6}) and used Blast2GO to assign GO terms for

each gene identified in the genome scan analysis (Conesa *et al.*, 2005; Conesa &

Götz, 2008) as well as performed a blast to the *A. thaliana* dataset TAIR10

separately to remove redundant terms (Berardini *et al.*, 2015). The false discovery

rates (FDR) were calculated to estimate the extent to which genes were enriched in

given GO categories (significance cutoff of P-values < 0.05). For each of the genes

enriched in specific biological processes, we retrieved the interacting gene

neighbours using GeneMANIA (Warde-Farley *et al.*, 2010). We generated aggregate

interaction networks in GeneMANIA, based on physical interactions, predicted and

co-expression. Finally, we performed hierarchical clustering and manually optimized

the weighted value cutoff for displaying the gene network.

Redundancy analysis (RDA)

We tested for genotype-environment association (GEA) using RDA (Capblancq &

Forester, 2021) using the *rda* function from the vegan package implemented in R

(Oksanen *et al.*, 2015), modelling genotypes as a function of the same climatic

predictor variables used for the niche reconstruction analyses, and producing

constrained axes and representative predictors. The significance of RDA constrained

axes was assessed using the *anova.cca* function and significant axes were then

used to identify candidate loci ($P < 0.001$). Candidate loci were identified using 2.5

standard deviation as cut-off (two-tailed p-value = 0.012). In order to measure the

rate of false-positive associations due to the demographic history, we also performed

the same RDA analysis using 1) a set of 1,000 randomly chosen SNPs from non-

sweep regions, and 2) polymorphism data from the neutral simulations used to

calibrate the sweeD and OmegaPlus thresholds.

Results

Past colonization events and climatic variations in *Solanum chilense*

We sequence whole genomes of 30 heterozygous plants from *S. chilense* from six

populations (LA3111, LA1963, LA2931, LA2932, LA4107, LA4330) (Fig. 1a; Table

S1) and used the reference genome assembly of *S. pennellii*. All 30 *S. chilense*

individuals show high-quality sequence and mapping scores with more than 97% of

mapping paired reads, individual genome coverage ranging between 16 to 24 reads

per base, and >70% genome coverage per sample (Dataset S1). After SNP calling

and stringent filtering, a total of 34,109,217 SNPs are identified across all 30 samples

(Table S2) for a genome size estimated approximately to 914Mb (Stam *et al.*, 2019a).

Phylogenetic analysis, principal component analysis (PCA) and population genetics

statistics (Fig. 1c, S1, S2; Table S3, S4) support the population structure into three

genetic groups, confirming the results in (Böndel *et al.*, 2015): a central group

(LA1963, LA3111, LA2931), the south-highland group (LA4330), and the south-coast

group (LA2932 and LA4107). The two south-coast populations constitute

independent groups (best K=4; Fig. 1d, S1b-d). Only the individuals of the population

LA2931 (the southernmost of the central group) display small admixed ancestry

coefficients (< 5%) with the south-highland group (LA4330, Fig. 1d). There is no

significant correlation between genetic (pairwise Nei's distance) and geographical

distance (Pearson test, r = 0.35, P = 0.20; Fig. S1e).

As we confirm that *S. chilense* independently colonized the coastal and

highland southern habitats from a lowland area located north of the central group

region (Böndel *et al.*, 2015; Stam *et al.*, 2019b), we further refine our estimates of the

historical changes in effective population size (N_e, Fig. 2a, S3), divergence and

potential post-divergence gene flow (Fig. S4), and finally construct a consensus

demographic model (Fig. 2b; Dataset S2; see Fig. S3 for confidence intervals related

to mutation rate and generation time uncertainty). These estimates are compared in

Fig. 2b with the reconstructed past climatic variation highlighting five marine isotope

stages (MIS) climatic periods (Lisiecki & Raymo, 2005; Ritter *et al.*, 2019). The two

south-coast populations found in Lomas habitats (LA2932 and LA4107) show early

divergence consistent with the admixture analysis (during the Last Inter-Glacial

period, MIS5) likely from the lowland area of the central group (LA1963). The

colonization of the highland likely occurred later, first in the central group region

(LA3111, LA2931) between the last interglacial and Last Glacial Maximum (LGM)

periods (ca. 75-130 kya, MIS3-4) and then with further colonization of southern

highlands (from 30 kya, MIS1-2, LA4330). All populations show a moderate effective

size reduction matching with the estimated time of the LGM characterized as a cold

and dry period and supported by a contraction of the suitable habitats to a narrow

strip in lower altitudes, and a subsequent expansion thereafter (Fig. 2a,c). Indeed,

the local habitat at the current location of LA2931 and LA4330 was likely unsuitable

for the establishment of southern highland populations until 15 kya (after the LGM,

i.e. during MIS1-2; Fig. 2c). The lower genetic diversity of the south populations (and

estimated *Ne*) is thus due to a mild colonization bottleneck during the southward

expansion (Fig. 2a,c, S2; Table S3). Both south-coast populations show consistent

signals of long-term history of colonization, subsequent isolation with negligible gene

flow, and possible local specialization to sparsely suitable Lomas habitats along the

coast (Fig. 2b,c, S3).

The divergence between the central group populations (during MIS3-4) occurs

in parallel to the colonization of the coastal habitat (Fig. 2b, S4), but before the

colonization of the south-highland (LA4330). Moreover, strong post-divergence gene-

flow and low differentiation are found in the central group, especially among the pairs

LA1963-LA3111 and LA3111-LA2931 (Fig. S2c, S4), consistent with their

geographical and/or environmental proximity (Fig. 1a,b) and the range contraction

during the LGM (MIS2 in Fig. 2). The colonization of high-altitude regions in the

central group is thus accompanied by high levels of gene flow despite these

populations ranging across a large altitudinal gradient (2500m of altitude difference

between LA1963 and LA3111 or LA2931). The divergence history results in the

south-coast and south-highland populations to be fairly isolated from one another (as

separated by the Atacama desert) leading to the suggestion of an incipient speciation

process (Fig. 2b, S4) (Raduski & Igić, 2021).

Selective sweeps underpin local adaptation

In total, we find 2,921 candidate sweep regions with SweeD (mean size 212,858 bp

+/- 3,938) and 13,106 with OmegaPlus (mean size 59,618 bp +/- 521) across all six

populations (Table 1), yielding a total of 520 overlapping regions (mean size 41,082

bp +/- 1,618). These regions contain 799 protein-coding candidate genes assumed to

likely be under positive selection (Fig. S5; Dataset S3). In LA4107, we find 61

candidate genes and about 100 candidate genes are detected in each of the other

four populations (Table 1). The largest number of candidate genes (354) is found in

LA4330 (Table 1), likely because the population has been established recently (Fig.

2a,b), and its habitat is ostensible different from the rest of the species range (Fig.

1b).

We present here two arguments supporting the fact that our cutoff values are

well designed on the basis of the population demography to reasonably discriminate

between demography and selection signals (as shown in Huber *et al.*, 2014). First,

comparison of genome-wide genetic diversity statistics (π and Tajima's D) between

the observed data and the simulations show that our demographic scenario captures

well the genomic diversity patterns in all populations (Fig. S2). Second, we estimate

by simulations the accuracy (statistical power) to detect sweeps due to strong

selection under our demographic model to be between 64 and 88% across

populations and for each method (Table 2). LA4330 exhibits even relatively high

statistical power compared to the other populations (Table 2). This demonstrates that

thresholds for sweep detection are conservative and allow minimizing the rate of

false positives, at the small cost of not detecting all selective sweeps, especially if the

selection coefficients are too small (Table 2, S5). Further, only a few candidate genes

are shared among different populations, with the central and south-highland

populations sharing a small number of candidate genes, while almost none are

shared between the two south-coastal populations (Fig. S5c). This lack of common

candidate genes among populations is likely due to 1) the high effective population

sizes (Fig. 2a) generating new variants across many genes which are then

differentially picked up by selection across different populations, and 2) the relatively

old inter-population divergence and timing of local adaptation.

An overview of population genetics statistics shows that our candidate regions

exhibit typical characteristics (lower nucleotide diversity, higher LD, more negative

Tajima's D and higher pairwise F_{ST} values) of positively selected regions when

compared to the genome-wide statistics (see SI Text 2; Fig. S6; Table S3, S4).

Furthermore, we find an overlap between our candidate genes under selective sweep

and genes exhibiting signals of positive selection in previous studies in *S. chilense*

which are based on few chosen genes, different plants, different populations and

different sample sizes than ours. Among our candidate genes, we indeed find three

genes (*JERF*3, *TPP* and *CT*189) involved in abiotic stress tolerance such as salt,

drought or cold (Böndel *et al.*, 2015) as well as three NLRs (nucleotide binding

leucine rich repeat, SOLCI006592800, SOLCI001535800, SOLCI005342400)

possibly linked to resistance to pathogens (Stam *et al.*, 2019b). We also find that two

of the seven most up-regulated genes under cold conditions in a transcriptomic study

of *S. chilense* (Nosenko *et al.*, 2016) do appear in our selection scan in high altitude

populations: *CBF*3 (Solyc03g026270) in LA2931, and *CBF*1 (Solyc03g026280) in

LA4330. These results indicate that our genome-wide selective sweep scan

generalizes the previous selection studies in *S. chilense* and supports the functional

relevance of our candidate genes.

Gene regulatory networks underlying local adaptation in *S. chilense*

A Gene Ontology (GO) enrichment analysis of the 799 candidate genes reveals significantly different and common biological processes across populations. We find common GO categories in all populations, including basic cell metabolism, immune response, specific organ development, and response to external stimuli (Fig. S7). Most interesting, however, are four GO categories specifically found in several populations and linked to local differences in habitats: (i) root hair cell differentiation functions are enriched in 15 candidate genes, only in the three coastal populations (LA1963, LA2932 and LA4107); (ii) response to circadian rhythm, photoperiodicity and flowering time are enriched in 12 candidate genes in two high-altitude (LA3111 and LA4330) and a south-coast (LA2932) populations; (iii) vernalization response is enriched in eight candidate genes in the three high-altitude populations (LA2931, LA3111, LA4330), and (iv) protein lipidation is enriched in seven candidate genes in the south-highland population (LA4330). Based on the wealth of available data in cultivated tomato, *S. pennellii* and *A. thaliana*, we further study the gene regulatory networks to which the candidate genes belong.

For adaptation to high-altitude conditions, 15 candidate genes are interconnected in a flowering gene network, which is itself subdivided into two sub-networks related to flowering, photoperiod and vernalization control pathways (Fig. 3a; Dataset S4). Photoperiod responsive genes can sense changes in sunlight and affect the circadian rhythm to regulate plant flowering (Johansson & Staiger, 2015; Song *et al.*, 2015), while vernalization genes regulate the flowering and germination through long-term low temperature (Guo *et al.*, 2018; Xu & Chong, 2018; Iida & Mähönen, 2020). These two sub-networks are connected through several key genes, some of which appear as candidate genes entailing local adaptation in our populations: FL FLOWERING LOCUS C (FLC or AGL25), FLOWERING LOCUS T

(FT) and AGAMOUS-LIKE genes (AGL) (Fig. 3a,b). These key genes are essential

regulators acting on the flowering regulation pathway (Michaels & Amasino, 1999;

Sheldon *et al.*, 2000; Turck *et al.*, 2008; Putterill & Varkonyi-Gasic, 2016).

Remarkably, some candidate genes in the recently diverged south-highland

population (LA4330) aggregate into an independent network involved in circadian

rhythm regulation, connected to the photoperiod network by JUMONJI DOMAIN

CONTAINING 5 (JMJD5) also a candidate gene in LA3111 (Fig. 3a). In the central-

highland population (LA3111), several other candidate genes of the photoperiod

network also regulate circadian rhythm and flowering time. The three high-altitude

populations (LA3111, LA2931, and LA4330) have candidate genes of the

AGAMOUS-LIKE (AGL) gene family in the vernalization network (Fig. 3a). We also

note that the network of protein lipidation genes appears to be related to the

synthesis of fatty acids in the south-highland population (Fig. 3d; Dataset S4). We

speculate here that this latter adaptation may be related to adaptation to lowest-

temperature stress of LA4330 (Dataset S5) (Maksimov *et al.*, 2017; Jiang *et al.*,

2018). Adaptation to high altitude involves the regulation of the flowering, including

photoperiod and vernalization pathways, but through different genes in different

populations, while cold stress and its consequence (adaptation in lipidation pathway)

may be relevant for adaptation to the highest altitudes (LA4330).

Regarding adaptation to coastal conditions, we find 11 candidate genes related

to root development and cellular homeostasis functions clustered in a single network

(Fig. 3c; Dataset S4). We speculate that the drought and water shortage typical of

the coastal conditions (Dataset S5) would promote the differentiation and extension

of plant roots (Xiong *et al.*, 2013; Li *et al.*, 2017). The cell WALL ASSOCIATED

LINASE 4 (WAK4), a candidate gene identified LA4107, acts as a linker of signal

from the cell wall to the plasma membrane and thus serve a vital role in lateral root

development (He *et al.*, 1999; Lally *et al.*, 2001). It is also possible that the soft soil at

the seaside is also beneficial to the extension of root hairs to increase the water

absorption area (White & Kirkegaard, 2010; Bengough *et al.*, 2016). In addition to

root development, we also find genes involved in cell homeostasis (Fig. 3c; Dataset

S4), which would be critical for the coastal drought and salinity conditions to maintain

the stability of the intracellular environment in the coastal habitats (Forni *et al.*, 2017;

Zhao *et al.*, 2020).

Candidate genes show genotype-environment associations

Our candidate loci are hypothesized to be responsible for adaptation to local climatic

conditions, so we test for genotype-environment association (GEA) using redundancy

analysis (RDA). We perform first a "present day" RDA using 144,713 SNPs from all

candidate regions and 63 climatic variables representing current (present) conditions

for temperature, precipitation, solar radiation, and wind (Dataset S5). We find that the

two-first RDA axes are significant (ANOVA's $P < 0.001$) and retain most (38% and

21%) of the putative adaptive genetic variance identified in the genome scans in all

populations (Fig. 4a). Tables S6 and S7 summarize outlier SNPs in different RDA

models and their correlation with climatic variables. In concordance with the PCAs of

both climatic and genomic variation (Fig. 1b,c), the two main RDA axes cluster the

individuals into three groups corresponding to the main geographical regions (central,

south-highland, and south-coastal) supporting that those axes synthesize the

principal selective pressures for local spatial adaptation along with the species

distribution (Fig. 4a; Table S7). RDA1 represents the differentiation of the two south-

coast populations in correlation with higher precipitation of the coldest quarter (Bio19)

and annual variation of solar radiation (CV_R) and RDA2 summarizes a climatic

gradient differentiating the south-highland population mainly driven by annual

potential evapotranspiration (annualPET) and temperature annual range (Bio7)

(Table S7).

Further RDA analyses based on gene variants of the GO categories circadian

rhythm-photoperiodism, vernalization, root-hair differentiation, and protein lipidation

highlight combinations of climatic variables and genetic variants related to local

spatial adaptation (Fig. S8a,c,e,g). These analyses show that the two main RDA axes

explain 40% of the variation. Climatic variables representing temperature variability

through the year such as temperature seasonality (Bio4) and temperature annual

range (Bio7) are consistently correlated with adaptive variation of the south-highland

population (Fig. 4a, S8). A total of 68 SNPs within candidate genes of the population

LA4330 are strongly associated with these two variables (Bio4, Bio7) in three of the

RDA based on the GO categories (circadian rhythm-photoperiodism, vernalization,

and protein lipidation; Dataset S6). The RDA based on the root-hair differentiation

GO category exhibits a strong differentiation between lowland and highland

populations based on atmosphere water vapour availability variables (ann_Vmin,

ann_AET; Fig. S8e). Note that the RDA testing for false positives implemented on

1,000 random SNPs from non-sweep regions and neutral simulations produced no

significant RDA axes and correlations with any climatic variable.

To assess the occurrence of selection in *S. chilense* as a response to past

climatic changes, we implement an "LGM" RDA using 37 climate variables projected

to the Last Glacial Maximum conditions (Fig. 4b, S8b,d,f,h; Dataset S5). This LGM

RDA analysis aims to uncover additional genomic variation selected in response to

temporal climatic changes and underlying the niche expansion towards the south-

highland region (Fig. 1b). The LGM RDA analyses capture a smaller proportion of the

genetic variability in the first two constrained axes (30%) compared to that using the

current climatic variables. About 30% outlier SNPs are identified in genomic regions

correlated with past climatic variables and not with the current variables (Table S6,

S7). For example, the central populations LA3111 and LA2931 are separated in the

past RDA of vernalization genes using LGM climatic variables indicating that warmer

climate after LGM may drive gene flow among central populations as seen in the

current RDA (Fig. 2b, S4, S8c,d). The past RDA of LGM climatic variables unveils

that high-altitude populations, especially LA4330, have SNPs correlating with

temperature (*i.e.* annual mean minimum temperature; ann_mTmin, and temperature

annual range; Bio7) whereas coastal populations SNPs do correlate with precipitation

and potential evapotranspiration of the coldest and driest seasons (Fig. 4b). We

advise caution in interpreting the LGM RDA results as these are based on climatic

values from locations that likely had little or no population occurrence in the past,

especially those in highland areas (but only mild bottlenecks could suggest local

persistence, Fig. 2a,c). We suggest that this analysis is nevertheless useful for

identifying alleles that arose in response to sudden changes in adaptive climatic

optima during glacial-interglacial transitions, especially in the highland populations .

Age of selective sweeps and timing of selection

We finally estimate the age, that is the time of the appearance of the selected allele,

of 112 selective sweep regions chosen to conservatively overlap between the three

detection methods (McSwan, SweeD and OmegaPlus, Table 1). These regions

contain 175 genes and exhibit a mean sweep age of ca. 28,000 years. The ages of

sweeps range from as early as 65 kya up to 2.5 kya (Table 1, Fig. 5). The highland

populations exhibit more recent sweeps (2.5 - 35 kya) than those at the coastal

populations, consistent with the recent (re)colonization of higher altitudes (Fig. 5).

The south-coastal populations exhibit older and large distributions of sweep age

consistent with older events of colonization (2.5 - 65 kya). Regarding the key gene

networks of relevance for local adaptation highlighted above (root hair, protein

lipidation, vernalization and photoperiod), each of them exhibits a narrow range of

sweep age values across several populations (Fig. 5). The averages of sweep ages

observed (Table 1) are perfectly in line with the estimates obtained from the sweep

simulations under our demographic model (Table 2), demonstrating that our statistical

power is adequate to estimate sweep ages under the demographic model and that

old sweeps in the highland populations cannot be recovered (even if they occurred)

in contrast to the coastal populations.

Discussion

After detecting selective sweeps at candidate genes, we draw correlations between

allele frequencies and climatic data, age of seeps and species distribution range

shifts driven by climate changes. We demonstrate that the distribution of the ages of

selective sweeps at few key gene networks are restricted in time and can be

associated with local spatial adaptation due to the colonization of new habitats , or

climatic changes responsible for local temporal adaptations in marginal populations.

To our knowledge, this is the first study to attempt to dissect the complex spatial and

temporal selective processes involved during and after the colonization of new highly

stressful environments such as arid and hyper-arid regions in plants.

Taking our demographic and selection results altogether, we formulate the

following scenario for the highland colonization. During the past colder climate

phases (LGM-MIS2 at 30-15kya), the suitable areas of the species likely decreased

at high altitude (Fig. 2c). We speculate that the populations were already established

at high altitude before the LGM (MIS 3-5; Fig. 2) likely in the northern part of the

range (from the location of LA3111 up to that of LA2931), before a contraction of the

species range occurred towards lower altitudes during the LGM, and the subsequent

colonization of new southern locations concluded ~15kya (post-LGM, LA4330). The

highland populations (LA3111, LA2931, LA4330) show adaptation by a burst, *i.e.*

over a short time, of numerous selective sweeps across several gene networks (Fig.

5) but also show older selective events pre-dating the LGM (MIS3-4). Interestingly,

the population LA4330 exhibits selective sweeps in the vernalization and photoperiod

which pre-date its establishment. These selective events likely occurred in the

northern part of the range (LA3111, LA2931) during MIS2-4 acting as pre-adaptation

requisite for colonizing the more divergent and extreme environments of the south-

highlands (LA4330; Fig. 1b).

The *S. chilense* lineage likely originates from coastal up to 'pre-cordillera' (800 -

2000 m altitude) habitats in southern Peru, explaining the early divergence and

southward colonization process, accompanied by habitat fragmentation and

contraction, which yields two highly isolated populations on the coast (Fig. 2b,c;

LA2932 and LA 4107). The coastal colonization process seems to involve fewer

sweeps than the adaptation to higher altitudes, for example a burst of selective

sweeps in genes related to root anatomical traits during the LGM-MIS2 period (Fig.

5). We speculate here that these sweeps are due to temporal adaptation to changes

in the habitat after colonization. However, some of the adaptive genomic signals in

the coastal populations could be blurred due to the older divergence time and

stronger drift (due to habitat fragmentation along the coast), or be

incomplete/partial/soft sweeps (with small selection coefficients) which we do not

detect (*e.g.* Garud *et al.*, 2021).

We find between 60 and 350 selective sweeps per population, but contrary to

our naïve expectations and previous findings in the literature, sweeps show a large

distribution of ages, especially in the south-coastal populations. We suggest that

several sweeps do occur concomitantly in a given gene pathway/network at a given

time period, either to promote local spatial adaptation or in response to a moving

environmental optimum (our climatic periods, temporal adaptation) as predicted

under the polygenic model of adaptation (Polechová *et al.*, 2009; Chevin *et al.*, 2010;

Matuszewski *et al.*, 2014; Jain & Stephan, 2017a). Selective sweeps at genes with

large selection coefficients can be observed because the populations of *S. chilense*

exhibit large effective sizes (Fig. 2; Böndel *et al.*, 2015), especially when compared to

the small above ground abundance (census size) reported in these semi-arid habitats

(Tellier *et al.*, 2011). *S. chilense* is outcrossing and exhibits persistent seed banking.

Both factors contribute to generate large effective population sizes by 1) decreasing

linkage disequilibrium and the effect of linked selection, 2) buffering the negative

impact of colonization bottlenecks, and 3) enhance the recovery post-colonization

(Fig. 2; Tellier *et al.*, 2011; Živković & Tellier, 2018). Therefore, the detection of old

sweeps dating up to 65 kya for the coastal populations and up to 35 kya for the

highland populations (pre-dating the recent post-LGM colonization; Fig. 2) is made

possible and stretching beyond the theoretical limit of $0.1N_e$ computed without the

effect of seed banking (Kim & Stephan, 2002).

As a word of caution, we focus on four main GO categories, which can be

reliably associated with physiological traits likely underlying adaptation: root hair

differentiation, vernalization, photoperiod, and protein lipidation. Pinpointing the

regulatory or non-coding SNPs under selection was not possible with our sample

sizes and functional information on many candidate genes in Figure 5 and S6 is still

lacking to provide a complete picture. Furthermore, we focus here on selective

sweeps resulting from strong positive selection as we cannot assess in our data the

occurrence of weaker positive (polygenic) selection or signatures of soft or

incomplete sweeps which would be favoured by the presence of seed banking

(Živković & Tellier, 2018). Yet, we are confident that our candidate genes under

selection are functionally relevant, as demonstrated by the overlap with previous

studies (Böndel *et al.*, 2015; Nosenko *et al.*, 2016; Stam *et al.*, 2019b). Our selection

scans are not exhaustive and future work requires larger sample sizes as well as

original material from *S. chilense* populations from the field to reveal the extent of

positive and balancing selection in this species.

Supplementary information

Available as a single pdf with additional figures, tables and fully detailed methods.

Acknowledgments

AT acknowledges funding from DFG (Deutscheforschungsgemeinschaft) Grant Number: 317616126 (TE809/7-1); KW was funded by the Chinese Scholarship Council, and GSA funded by the Technical University of Munich. We thank the Tomato Genetics Resource Center (TGRC) of the University of California, Davis for generously providing us with the seeds of the accession included in this study. We thank Daniela Scheikl for assistance with plant work and Christine Würmser for the Illumina sequencing.

Author contributions

Conceptualization, GSA and AT; Software, Formal analysis and Investigation, KW and GSA, Writing KW, GSA and AT; Funding Acquisition, AT; Supervision, GSA and AT.

Data availability

The raw pair-end sequencing genomic data used in all the analyses can be accessed at the European Nucleotide Archive (ENA) project accession PRJEB47577.

All code used in this study and other previously published genomic data is available at the sources referenced. The developed scripts and extra data formats are found on our gitlab: https://gitlab.lrz.de/population_genetics/wild-tomato-genomics.

 Figure legends

Fig. 1 Geographic distribution and population structure of *S. chilense.* (a) Map with

distribution of all *S. chilense* populations by the TGRC, the six *S. chilense*

populations in this study (black circles), and the four population groups (colorful

circles). (b) The PCA of all populations using 63 current climatic variables (Dataset

S5). Population structure using SNP data based on (c) PCA and (d) Admixture

analysis (optimal K value is 4; Fig S1b).

Fig. 2 Demographic history and species distribution model of *S. chilense* for current
and Last Glacial Maximum (LGM) climate conditions (a) The estimation of historical
patterns of effective population size (N_e) for 10 pairwise genome comparisons per
population using the MSMC model. (b) Interpreted demographic scenario for the six
samples populations of *S. chilense* including likely estimations of effective population
size, divergence times and gene-flow. The width of the boxes represents the relative
effective population size, arrows represent the migration between population pairs.
Grey background boxes indicate five Marine isotope stages (MIS) climatic periods.
(c) Overlay of the reconstruction of the distribution model for *S. chilense* using
current climatic variables (red) and LGM past climatic variables (blue). Darker color
of the gradient indicates higher suitable habitat for a given climatic period.

Fig. 3 Interaction genetic networks of candidate genes. (a) The network of flowering

regulation involved two sub-networks, photoperiod and vernalization pathways, for

regulation of flowering. (b) The schematic diagram of flowering regulation involved

photoperiod and vernalization is adapted from (Xu & Chong, 2018). "┤"indicates

repressive effects on gene expression; "→"indicates promotive effects on gene

expression. (c) The network of root development and cell homeostasis. (d) The

networks of protein lipidation. Connections represent gene interactions based on

physical   interactions,   informatics   predictions   and   co-expression   analyses.

Connection thickness is proportional to weighted value of the connected genes. The

black lines connected two sub-networks, genes under selection were connected by

solid lines and other genes were connected by dashed line. Node colors correspond

to genes were detected the different populations in genome scans. Gray circles, not

detected in genome scan, but present in *S. chilense*; gray squares: not present in *S.*

chilense.

618 Fig. 4 Redundancy analysis (RDA) ordination bi-plots between the climatic variables,

619 populations and the genetic variants in all candidate sweeps. (a) RDA using current

620 climatic variables. (b) RDA using LGM climatic variables. Arrows indicated the

621 direction and magnitude of variables correlated with the populations. The

622 abbreviations of climatic variables are shown in Dataset S5.

629   Fig. 5 Distribution of sweep age across the five MIS climatic periods.

633   Table 1. The summary of genome scans and estimation of sweep age.

Population	Genome Scans				Sweeps age			
	N_{SweeD}	$N_{OmegaPlus}$	$N_{overlaps1}$	N_{genes1}	N_{McSwan}	$N_{overlaps2}$	N_{genes2}	Age_{mean} (kyr)
LA1963	385	2,474	98[b]	86	267	16	14	38 ± 16
LA2931	517	2,268	109	125	355	24	28	20 ± 10
LA2932	374	1,717	46	101	302	15	29	36 ± 15
LA3111	663	2,307	105	107	377	22	22	23 ± 11
LA4107	203	2,047	37	61	194	11	13	34 ± 10
LA4330	779	2,293	125	354	438	36	71	17 ± 8

N_{SweeD}, number of outlier regions from SweeD; $N_{OmegaPlus}$, number of outlier regions from OmegaPlus;

$N_{overlaps1}$, number of overlapping regions between SweeD and OmegaPlus; N_{genes1}, number of candidate

genes in overlaps1, and all candidate genes show in Data S1C; N_{McSwan}, number of outlier regions from

McSwan; $N_{overlaps2}$, number of overlapping regions between McSwan and overlaps1; N_{genes2}, number of

genes in overlaps2; Age_{mean}, mean age ± standard deviation of overlaps2.

Table 2. The estimation of sensitivity to detect selective sweeps using our pipeline with
different selection strengths.

Population	s=0.5			s=1			Mean age ± sd (kya)
	SweeD	OmegaPlus	McSwan	SweeD	OmegaPlus	McSwan	
LA1963	22.4	31.2	29.8	74.2	81.6	76.4	36 ± 14
LA2931	27.3	35.1	30.6	76.3	84.1	77.2	24 ± 12
LA2932	19.7	27.4	29.4	68.5	78.3	75.3	35 ± 16
LA3111	26.8	33.5	30.2	74.1	80.5	78.4	27 ± 12
LA4107	17.4	21.7	28.7	63.6	74.8	71.7	33 ± 14
LA4330	30.1	37.9	34.6	81.7	88.3	79.8	19 ± 10

644 The validation of our pipeline to detect selective sweeps by simulating 1,000 data sets of selective
645 sweeps with four different selective strengths (s=0.01, 0.1, 0.5, 1; see also Table S5) in six populations.
646 The numbers denote the percentage of the detected simulated sweeps using different methods. The
647 mean age represents average age of simulated sweeps detected by McSwan.
648
649

References

Alachiotis N, Stamatakis A, Pavlidis P. **2012**. OmegaPlus: a scalable tool for rapid detection of selective sweeps in whole-genome datasets. *Bioinformatics* **28**: 2274–2275.

Alexander DH, Novembre J, Lange K. **2009**. Fast model-based estimation of ancestry in unrelated individuals. *Genome research* **19**: 1655–1664.

Araujo M, New M. **2007**. Ensemble forecasting of species distributions. *Trends in Ecology & Evolution* **22**: 42–47.

Barghi N, Hermisson J, Schlötterer C. **2020**. Polygenic adaptation: A unifying framework to understand positive selection. *Nature Reviews Genetics* **21**: 769–781.

Bengough AG, Loades K, McKenzie BM. **2016**. Root hairs aid soil penetration by anchoring the root surface to pore walls. *Journal of Experimental Botany* **67**: 1071–1078.

Berardini TZ, Reiser L, Li D, Mezheritsky Y, Muller R, Strait E, Huala E. **2015**. The Arabidopsis information resource: making and mining the "gold standard" annotated reference plant genome. *genesis* **53**: 474–485.

Bolger A, Scossa F, Bolger ME, Lanz C, Maumus F, Tohge T, Quesneville H, Alseekh S, Sørensen I, Lichtenstein G, *et al.* 2014. The genome of the stress-tolerant wild tomato species *Solanum pennellii*. *Nature Genetics* **46**: 1034–1038.

Böndel KB, Lainer H, Nosenko T, Mboup M, Tellier A, Stephan W. **2015**. North–south colonization associated with local adaptation of the wild tomato species *Solanum chilense*. *Molecular Biology and Evolution* **32**: 2932–2943.

Böndel KB, Nosenko T, Stephan W. **2018**. Signatures of natural selection in abiotic stress-responsive genes of *Solanum chilense*. *Royal Society Open Science* **5**: 171198.

Bragg JG, Supple MA, Andrew RL, Borevitz JO. **2015**. Genomic variation across landscapes: insights and applications. *New Phytologist* **207**: 953–967.

Capblancq T, Forester BR. **2021**. Redundancy analysis: A Swiss Army Knife for landscape genomics. *Methods in Ecology and Evolution* **12**: 2298–2309.

Chevin L-M, Martin G, Lenormand T. **2010**. Fisher's model and the genomics of adaptation: Restricted pleiotropy, heterogenous mutation, and parallel evolution. *Evolution* **64**: 3213–3231.

Conesa A, Götz S. **2008**. Blast2GO: a comprehensive suite for functional analysis in
plant genomics. *International journal of plant genomics* **2008**.

Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M. **2005**.
Blast2GO: a universal tool for annotation, visualization and analysis in functional
genomics research. *Bioinformatics* **21**: 3674–3676.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker
RE, Lunter G, Marth GT, Sherry ST. **2011**. The variant call format and VCFtools.
Bioinformatics **27**: 2156–2158.

Delaneau O, Marchini J, Zagury J-F. **2012**. A linear complexity phasing method for
thousands of genomes. *Nature methods* **9**: 179–181.

Ewing G, Hermisson J. **2010**. MSMS: a coalescent simulation program including
recombination, demographic structure and selection at a single locus. *Bioinformatics*
26: 2064–2065.

Fagny M, Austerlitz F. **2021**. Polygenic adaptation: integrating population genetics
and gene regulatory networks. *Trends in Genetics* **37**: 631–638.

Fick SE, Hijmans RJ. **2017**. WorldClim 2: New 1-km spatial resolution climate
surfaces for global land areas. *International Journal of Climatology* **37**: 4302–4315.

Fischer I, Camus-Kulandaivelu L, Allal F, Stephan W. **2011**. Adaptation to drought
in two wild tomato species: the evolution of the *Asr* gene family. *New Phytologist* **190**:
1032–1044.

Forni C, Duca D, Glick BR. **2017**. Mechanisms of plant response to salt and drought
stress and their alteration by rhizobacteria. *Plant and Soil* **410**: 335–356.

François O, Martins H, Caye K, Schoville SD. **2016**. Controlling false discoveries in
genome scans for selection. *Molecular Ecology* **25**: 454–469.

Garud NR, Messer PW, Petrov DA. **2021**. Detection of hard and soft selective
sweeps from *Drosophila melanogaster* population genomic data. *PLOS Genetics* **17**:
e1009373.

Guo X, Liu D, Chong K. **2018**. Cold signaling in plants: Insights into mechanisms
and regulation. *Journal of integrative plant biology* **60**: 745–756.

He Z-H, Cheeseman I, He D, Kohorn BD. **1999**. A cluster of five cell wall-associated
receptor kinase genes, Wak1–5, are expressed in specific organs of *Arabidopsis*.
Plant molecular biology **39**: 1189–1196.

Hoban S, Kelley JL, Lotterhos KE, Antolin MF, Bradburd G, Lowry DB, Poss ML, Reed LK, Storfer A, Whitlock MC. **2016**. Finding the Genomic Basis of Local Adaptation: Pitfalls, Practical Solutions, and Future Directions. *The American Naturalist* **188**: 379–397.

Huber CD, Nordborg M, Hermisson J, Hellmann I. **2014**. Keeping it local: Evidence for positive selection in swedish *Arabidopsis thaliana*. *Molecular Biology and Evolution* **31**: 3026–3039.

Iida H, Mähönen AP. **2020**. Growth-mediated sensing of long-term cold in plants. *Nature* **583**: 690–691.

Jain K, Stephan W. **2017a**. Rapid adaptation of a polygenic trait after a sudden environmental shift. *Genetics* **206**: 389–406.

Jain K, Stephan W. **2017b**. Modes of rapid polygenic adaptation. *Molecular Biology and Evolution* **34**: 3169–3175.

Jiang H, Zhang X, Chen X, Aramsangtienchai P, Tong Z, Lin H. **2018**. Protein lipidation: occurrence, mechanisms, biological functions, and enabling technologies. *Chemical reviews* **118**: 919–988.

Johansson M, Staiger D. **2015**. Time to flower: interplay between photoperiod and the circadian clock. *Journal of experimental botany* **66**: 719–730.

Josephs EB, Stinchcombe JR, Wright SI. **2017**. What can genome-wide association studies tell us about the evolutionary forces maintaining genetic variation for quantitative traits? *New Phytologist* **214**: 21–33.

Kawecki TJ, Ebert D. **2004**. Conceptual issues in local adaptation. *Ecology Letters* **7**: 1225–1241.

Kim Y, Stephan W. **2002**. Detecting a local signature of genetic hitchhiking along a recombining chromosome. *Genetics* **160**: 765.

Lally D, Ingmire P, Tong H-Y, He Z-H. **2001**. Antisense expression of a cell wall–associated protein kinase, WAK4, inhibits cell elongation and alters morphology. *The Plant Cell* **13**: 1317–1332.

Legendre P, Legendre L. **2012**. *Numerical ecology*. Amsterdam: Elsevier.

Li X, Cai W, Liu Y, Li H, Fu L, Liu Z, Xu L, Liu H, Xu T, Xiong Y. **2017**. Differential TOR activation and cell proliferation in *Arabidopsis* root and shoot apexes. *Proceedings of the National Academy of Sciences* **114**: 2765–2770.

Li H, Durbin R. **2009**. Fast and accurate short read alignment with Burrows–Wheeler
transform. *bioinformatics* **25**: 1754–1760.

Lisiecki LE, Raymo ME. **2005**. A Pliocene-Pleistocene stack of 57 globally
distributed benthic $\delta^{18}O$ records. *Paleoceanography* **20**: PA1003.

Lotterhos KE, Whitlock MC. **2015**. The relative power of genome scans to detect
local adaptation depends on sampling design and statistical method. *Molecular*
Ecology **24**: 1031–1046.

Maksimov EG, Mironov KS, Trofimova MS, Nechaeva NL, Todorenko DA,
Klementiev KE, Tsoraev G V, Tyutyaev E V, Zorina AA, Feduraev P V. **2017**.
Membrane fluidity controls redox-regulated cold stress responses in cyanobacteria.
Photosynthesis research **133**: 215–223.

Malaspinas A-S, Westaway MC, Muller C, Sousa VC, Lao O, Alves I, Bergström
A, Athanasiadis G, Cheng JY, Crawford JE, *et al.* **2016**. A genomic history of
Aboriginal Australia. *Nature* **538**: 207–214.

Matuszewski S, Hermisson J, Kopp M. **2014**. Fisher's geometric model with a
moving optimum. *Evolution* **68**: 2571–2588.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A,
Garimella K, Altshuler D, Gabriel S, Daly M, *et al.* **2010**. The Genome Analysis
Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing
data. *Genome Research* **20**: 1297–1303.

Michaels SD, Amasino RM. **1999**. FLOWERING LOCUS C encodes a novel MADS
domain protein that acts as a repressor of flowering. *The Plant Cell* **11**: 949–956.

Nakazato T, Warren DL, Moyle LC. **2010**. Ecological and geographic modes of
species divergence in wild tomatoes. *American Journal of Botany* **97**: 680–693.

Nosenko T, Böndel KB, Kumpfmüller G, Stephan W. **2016**. Adaptation to low
temperatures in the wild tomato species *Solanum chilense*. *Molecular Ecology* **25**:
2853–2869.

Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB,
Simpson GL, Solymos P, Stevens MHH, Wagner H. **2015**. *VEGAN: Community*
ecology package. R package version 2.3-0.

Pannell JR, Fields PD. **2014**. Evolution in subdivided plant populations: concepts,
recent advances and future directions. *New Phytologist* **201**: 417–432.

Pavlidis P, Živković D, Stamatakis A, Alachiotis N. **2013**. SweeD: Likelihood-

Based Detection of Selective Sweeps in Thousands of Genomes. *Molecular Biology and Evolution* **30**: 2224–2234.

Polechová J, Barton N, Marion G. **2009**. Species' range: Adaptation in space and time. *The American Naturalist* **174**: E186–E204.

Putterill J, Varkonyi-Gasic E. **2016**. FT and florigen long-distance flowering control in plants. *Current Opinion in Plant Biology* **33**: 77–82.

R Core Team. **2020**. *R: A language and environment for statistical computing*. Vienna, Austria. http://www.R-project.org/: R Foundation for Statistical Computing.

Raduski AR, Igić B. **2021**. Biosystematic studies on the status of *Solanum chilense*. *American Journal of Botany* **108**: 520–537.

Ritter B, Wennrich V, Medialdea A, Brill D, King G, Schneiderwind S, Niemann K, Fernández-Galego E, Diederich J, Rolf C, *et al.* 2019. Climatic fluctuations in the hyperarid core of the Atacama Desert during the past 215 ka. *Scientific Reports* **9**: 5270.

Roselius K, Stephan W, Städler T. **2005**. The relationship of nucleotide polymorphism, recombination rate and selection in wild Tomato species. *Genetics* **171**: 753–763.

Savolainen O, Lascoux M, Merilä J. **2013**. Ecological genomics of local adaptation. *Nature Reviews Genetics* **14**: 807–820.

Schrider DR, Kern AD. **2017**. Soft sweeps are the dominant mode of adaptation in the human genome. *Molecular Biology and Evolution* **34**: 1863–1877.

Sheldon CC, Rouse DT, Finnegan EJ, Peacock WJ, Dennis ES. **2000**. The molecular basis of vernalization: the central role of FLOWERING LOCUS C (FLC). *Proceedings of the National Academy of Sciences* **97**: 3753–3758.

Smith JM, Haigh J. **1974**. The hitch-hiking effect of a favourable gene. *Genetical Research* **23**: 23–35.

Song YH, Shim JS, Kinmonth-Schultz HA, Imaizumi T. **2015**. Photoperiodic flowering: time measurement mechanisms in leaves. *Annual review of plant biology* **66**: 441–464.

Staab PR, Zhu S, Metzler D, Lunter G. **2015**. scrm: Efficiently simulating long sequences using the approximated coalescent with recombination. *Bioinformatics* **31**: 1680–1682.

Stam R, Nosenko T, Hörger AC, Stephan W, Seidel M, Kuhn JMM, Haberer G, Tellier A. **2019a**. The *de novo* reference genome and transcriptome assemblies of the wild tomato species *Solanum chilense* highlights birth and death of NLR genes between tomato species. *G3: Genes, Genomes, Genetics* **9**: 3933–3941.

Stam R, Silva-Arias GA, Tellier A. **2019b**. Subsets of NLR genes show differential signatures of adaptation during colonization of new habitats. *New Phytologist* **224**: 367–379.

Stephan W. **2016**. Signatures of positive selection: From selective sweeps at individual loci to subtle allele frequency changes in polygenic adaptation. *Molecular Ecology* **25**: 79–88.

Tellier A, Laurent SJY, Lainer H, Pavlidis P, Stephan W. **2011**. Inference of seed bank parameters in two wild tomato species using ecological and genetic data. *Proceedings of the National Academy of Sciences* **108**: 17052–17057.

Thuiller W, Georges D, Engler R. **2014**. *BIOMOD2: Ensemble platform for species distribution modeling. R package   version 3.1-62/r677*.

Thuiller W, Lafourcade B, Engler R, Araújo MB. **2009**. BIOMOD - a platform for ensemble forecasting of species distributions. *Ecography* **32**: 369–373.

Tiffin P, Ross-Ibarra J. **2014**. Advances and limits of using population genetics to understand local adaptation. *Trends in Ecology & Evolution* **29**: 673–680.

Title PO, Bemmels JB. **2018**. ENVIREM: An expanded set of bioclimatic and topographic variables increases flexibility and improves performance of ecological niche modeling. *Ecography* **41**: 291–307.

Tournebize R, Poncet V, Jakobsson M, Vigouroux Y, Manel S. **2019**. McSwan: A joint site frequency spectrum method to detect and date selective sweeps across multiple population genomes. *Molecular Ecology Resources* **19**: 283–295.

Trabucco A, Zomer RJ. **2018**. Global aridity index and potential evapotranspiration (ET0) climate database v2. *CGIAR Consort Spat Inf*.

Turck F, Fornara F, Coupland G. **2008**. Regulation and identity of Florigen: FLOWERING LOCUS T moves center stage. *Annual Review of Plant Biology* **59**: 573–594.

Wang K, Mathieson I, O'Connell J, Schiffels S. **2020**. Tracking human population structure through time from whole genome sequences. *PLoS genetics* **16**: e1008552.

Warde-Farley D, Donaldson SL, Comes O, Zuberi K, Badrawi R, Chao P, Franz

M, Grouios C, Kazi F, Lopes CT. **2010**. The GeneMANIA prediction server:
biological network integration for gene prioritization and predicting gene function.
Nucleic acids research **38**: W214–W220.

White RG, Kirkegaard JA. **2010**. The distribution and abundance of wheat roots in a
dense, structured subsoil–implications for water uptake. *Plant, cell & environment* **33**:
133–148.

**Wright KM, Hellsten U, Xu C, Jeong AL, Sreedasyam A, Chapman JA, Schmutz
J, Coop G, Rokhsar DS, Willis JH**. **2015**. Adaptation to heavy-metal contaminated
environments proceeds via selection on pre-existing genetic variation. *bioRxiv*
029900.

Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. **2009**.
The Sequence alignment/map (SAM) format and SAMtools. *Bioinformatics* **25**: 2078–
2079.

Xia H, Camus-Kulandaivelu L, Stephan W, Tellier A, Zhang Z. **2010**. Nucleotide
diversity patterns of local adaptation at drought-related candidate genes in wild
tomatoes. *Molecular Ecology* **19**: 4144–4154.

Xiong Y, McCormack M, Li L, Hall Q, Xiang C, Sheen J. **2013**. Glucose–TOR
signalling reprograms the transcriptome and activates meristems. *Nature* **496**: 181–
186.

Xu S, Chong K. **2018**. Remembering winter through vernalisation. *Nature Plants* **4**:
997–1009.

Zhao C, Zhang H, Song C, Zhu J-K, Shabala S. **2020**. Mechanisms of plant
responses and adaptation to soil salinity. *The innovation* **1**: 100017.

Živković D, Tellier A. **2018**. All but sleeping? Consequences of soil seed banks on
neutral and selective diversity in plant species. In: Morris RJ, ed. Mathematical
Modelling in Plant Biology. Cham: Springer International Publishing, 195–212.

**Zou Y-P, Hou X-H, Wu Q, Chen J-F, Li Z-W, Han T-S, Niu X-M, Yang L, Xu Y-C,
Zhang J, *et al.* 2017**. Adaptation of *Arabidopsis thaliana* to the Yangtze River basin.
Genome Biology **18**: 239.