Local and temporal adaptation to climatic change in a wild tomato species via

selective sweeps

Kai Wei *, Gustavo A Silva-Arias *, Aurélien Tellier

Professorship for Population Genetics, Department of Life Science Systems, School

of Life Sciences, Technical University of Munich, Liesel-Beckmann Strasse 2, 85354

Freising, Germany

* these authors contributed equally

Corresponding author:

Aurélien Tellier tellier@wzw.tum.de

Keywords:

Local adaptation, selective sweeps, age of sweeps, last glacial maximum, gene

network

Summary

Positive Darwinian selection is the driving force underpinning local (in space) and temporal (in time) adaptation, and leaves footprints of selective sweeps at the underlying major genes. These two adaptive processes are classically considered independently, so that most genomic selection scans uncover only recent sweeps underpinning spatial adaptation. However, understanding if these adaptive processes are intermingled and share common genetic bases is crucial to predict a species' evolutionary potential, for example in response to changing environmental conditions. We use whole genome data from six populations across three different habitats of the wild tomato species *Solanum chilense*, to 1) infer the past demographic history, and 2) search for genes under strong positive selection. We then correlate the demographic history, allele frequencies in space and time, the age of selection events and the reconstructed historical ecological distribution of the species over five main climatic periods spanning 150,000 years. We find evidence for several selective sweeps targeting regulatory networks involved in root hair development in low altitude, and response to photoperiod and vernalization in high altitude populations. These sweeps occur in a concerted fashion in a given regulatory gene network and only at particular time periods, thereby underpinning temporal local adaptation. These genes under positive selection provide subsequently the basis for spatial local adaptation to novel habitats when new ecological niches become available. Our results reveal the importance of jointly studying spatial and temporal adaptations in species during habitat colonization.

Introduction

Positive Darwinian selection is a key process driving macro and micro-evolution and underlying local adaptation when species expand their range and colonize novel habitats. Local adaptation encompasses thus the spatial component of adaptation to a new habitat, but also a temporal component when climatic/abiotic conditions vary in time after colonization[1]. The latter is understood here at the scale of hundreds up to thousands of years during which climatic conditions are changing due to glacial/inter-glacial periods or medium to long-term changes in abiotic and biotic environmental conditions (the so-called moving environmental optimum). Population genetics theory investigates the characteristics and polymorphism footprints of local adaptation while separating the two processes: spatial adaptation to a local optimum based on Fisher's geometric model of adaptation[2] and temporal adaptation to a moving optimum[1,3,4]. Under both models and large enough population sizes, it is typically expected that bigger steps of adaptation occur first at sites with strong selective coefficients, possibly generating selective sweeps[2–4]. Selective sweeps are polymorphism patterns (footprints) in the genome due to the rapid fixation of advantageous alleles and the associated hitch-hiking effect[5,6]. However, there is so far no complete theory investigating the more realistic scenario combining both local spatial adaptation to newly colonized habitats and further temporal adaptation to temporally variable climatic conditions and the expected patterns of selection.

We identify therefore a number of unanswered theoretical and empirical questions regarding the joint effect of spatial and temporal local adaptation. How many genes underlie adaptation to new habitats and/or changing habitat in time? How many genes underpinning temporal or spatial adaptation show strong selection (selective sweeps)? Are the same genes involved in both adaptive processes? Do these genes belong to one or few gene pathways/regulatory networks? What is the timing of adaptation and can selection at a given gene be linked to past changes in climatic conditions? This lack of knowledge regarding the underpinnings of local

adaptation represents a major bottleneck to predict/assess the species potential for

adaptability and resilience in the face of the current global climatic change. Indeed,

despite numerous studies highlighting genes or even SNPs underlying spatial

climatic adaptation[7,8], future predictions of the strength of selection and change in

allele frequencies in response to the predicted climate change trends remain fairly

speculative.

To date, numerous studies uncovering the genetic bases of spatial adaptation,

have been conducted in few model organisms such as humans, invertebrates

(*Caenorhabditis elegans, Drosophila melanogaster*) or plants (*Arabidopsis* spp.,

Arabis sp., *Capsella* spp.). Using genome scan methods based on the selective

sweep/hitchhiking model[5,6], Genome Wide Association studies (GWAs) and

functional studies, the crucial role of changes in genes/proteins (coding regions)

and/or gene expression and the relevant gene network/pathways has been

investigated (review in [9,10]). Interestingly, most model study systems exhibit spatial

local adaptation due to recent selective sweeps being dated less than 25,000 years

old[11–13] which are associated with strong bottlenecks during colonization events (*e.g.*

colonization of Europe in humans and *Drosophila*) or due to recent recolonization

after glacial period, bottlenecks in glacial refugia and selfing (in *Arabidopsis thaliana*,

Capsella sp. and *Arabis* sp.). The observation of recent sweeps, and lack of older

selection events, is explained by the lack of power to detect loci under strong positive

local and temporal selection due to three factors. First, there is a confounding effect

of past demographic history of the populations[14] especially under strong bottlenecks

often characterizing the colonization of new habitats[15]. Disentangling the two types of

signatures, neutral demography and selection, is possible by inferring the past

demography over the whole genome and search for outlier portions of the genome

under selection. However, under strong demographic bottlenecks, it is difficult to

carefully control the statistical rate of false positive sweeps and sweep signatures

pre-dating the bottleneck are blurred[12–14]. Second, strong selection at few genes is

only the tip of the iceberg, as adaptation likely occurs for traits which are polygenic

with selection acting over many genes. Different number of major genes with

selective sweeps signatures can be observed depending on the distribution of

selection coefficients, the efficiency of selection (a function of effective population

size and recombination rate), the architecture of the traits, place of genes in gene

networks/pathways, and gene pleiotropy[16,17]. While theoretical predictions exist, a

general understanding of the genetic architecture of local adaptation is still missing

(but see [16]). Third, under temporal and polygenic selection model, strong selection

can only occur if population sizes are large enough[3,14]. Furthermore, theory predicts

that selective sweeps can be observed and detected reliably only for ca. $0.1 Ne$

generations after selection, where Ne is the effective population size[5]. As a result, in

species undergoing habitat range expansion with strong bottleneck during

colonization and life history traits decreasing effective population size (*e.g.* selfing),

we expect 1) only few sweeps to occur under polygenic selection, and 2) only recent

sweeps to be detected using polymorphism data. Therefore, in order to reveal the

underpinnings of long-term spatial and temporal local adaptation due to strong

selection (selective sweeps), the study of alternative species with larger effective

population sizes during colonization events is required.

Solanum chilense is an outcrossing wild tomato species found in southern Peru

and northern Chile in mesic to very arid habitats[18]. The ancestral species range is

found around the Peruvian and Chilean border mainly along the (mid-altitude) 'pre-

cordillera' region (800 - 2000 m altitude) characterized as marginal desert habitat. *S.*

chilense colonized independently two different southern, but arid, isolated regions

around the Atacama desert at different time points[19,20]: an early divergence (older

than 50,000 ya) with the colonization of coastal habitats (in Lomas formations), and a

more recent lineage divergence (less than 20,000 ya) restricted to highland altitudes

(above 2,400m) of the Andean plateau. Signatures of natural selection (positive or

balancing) at few genes involved in abiotic stress (drought, cold, salt) adaptation

were found when scanning few candidate genes for biotic and abiotic stresses[19–24]. In

the present study, we obtain full genome sequence data for 30 diploid and highly

heterozygous plants from six populations (five plants per population) representing the

three main habitats of the species (Figure 1A): the central group (area of origin at low

to high altitude), south-coastal group, and south-highland group. These habitats differ

in their current climatic conditions: the south-highland group is different from the

central group, while the south-coastal appears as only marginally different from the

environment prevailing in the central group (Figure 1B). Our aims are first to infer

accurately the past demographic history of the species colonization as well as to

reconstruct the past climatic history of the species' range. Second, we conduct

genome scans for genes under selection in these populations and assign functions

and gene network position to these candidate genes. We then link climatic and

genetic data at candidate genes to highlight the relevance of few key gene regulatory

networks (pathways) for local spatial adaptation. Finally, we connect past species

niche reconstruction and age estimates of the selective sweeps in order to reveal the

history of the temporal local adaptation to ever changing climatic conditions.

Results

Past demography of *S. chilense* is influenced by colonization events and
climatic variations

We sequence whole genomes of 30 heterozygous plants from *Solanum chilense*

from six populations (LA3111, LA1963, LA2931, LA2932, LA4107, LA4330) (Figure

1A, Table S1). All reads are aligned to the reference genome assembly of *S. pennellii*

and all 30 *S. chilense* individuals show high-quality sequence and mapping scores

with more than 97% of mapping paired reads, individual genome coverage ranging

between 16 to 24 reads per base, and >70% genome coverage per sample (Data

S1A). After SNP calling and stringent filtering, a total of 34,109,217 SNPs are

identified across all samples (Table S2) for a genome size estimated approximately

to 914Mb[25]. Phylogenetic and principal component analysis (PCA) analyses support

population structuring into three genetic groups (Figure 1C; Figure S1C; confirming

the results in [19]): a central group (LA1963, LA3111, LA2931), the south-highland

group (LA4330), and the south-coast group (LA2932 and LA4107). Moreover,

population structure analyses support that the two south-coast populations constitute

independent groups (best K=4; Figure 1D; Figure S1C-E). Only the individuals of the

population LA2931 (the southernmost of the central group) display small admixed

ancestry coefficients (< 5%) with the south-highland group (Figure 1D). There is no

significant correlation between genetic (pairwise Nei's distance) and geographical

distance (Pearson test, r = 0.35, P = 0.20; Figure S1A-B).

As indicated by nucleotide diversity and differentiation statistics (Figure S2;

Table S3 and S4), we confirm the previous results obtained using a reduced number

of genetic markers: *S. chilense* has colonized independently the coastal and highland

southern-habitats, and the species area of origin is likely the mid- altitude of the

central region[19,20]. Using multiple sequentially Markovian coalescent analyses, we

estimate historical changes in effective population size (N_e, Figure 2A), divergence

and potential post-divergence gene flow (Figure S2), and construct a consensus

demographic model (Figure 2B; Data S1B). These estimates are compared with the

reconstructed past climatic variation highlighting five Marine isotope stages (MIS)

climatic periods[26,27]. The two south-coast populations found in Lomas habitats show

early divergence consistent with the admixture analysis (during the Last Inter-Glacial

period, MIS5) likely from the lowland area of the central group (LA1963). The

colonization of the highland likely occurred later, first in the central group region

(LA3111, LA2931) between the last inter-glacial and Last Glacial Maximum (LGM)

periods (ca. 75-130 kya, MIS3-4) and then with further colonization of southern

highlands (from 30 kya, MIS1-2, LA4330). All populations show a reduction of

diversity matching with the estimated time of the LGM characterized as a cold and

dry period and supported by a contraction of the suitable habitats to a narrow strip in

lower altitudes, and a subsequent expansion thereafter (Figure 2C). Indeed, the local

habitat at the current location of LA2931 and LA4330 was likely unsuitable for the

establishment of the most southern highland population until 15 kya (after the LGM,

i.e. during MIS1-2). The lower genetic diversity of the south populations (and

estimated *Ne*) is thus due to the mild bottleneck processes during the southward

expansion (Figure 2A; Figure S2; Table S3). Both south-coast populations show

consistent signals of long-term history of colonization, subsequent isolation with

negligible gene flow, and possible local specialization to sparsely suitable Lomas

habitats along the coast (Figure 2B and 2C, Figure S2).

The divergence between the central group populations (during MIS3-4) occurs

after the colonization of the coastal habitat (Figure S3), but before the colonization of

the south-highland (LA4330). Moreover, strong post-divergence gene-flow and low

differentiation are found in the central group, especially among the pairs LA1963-

LA3111 and LA3111-LA2931 (Figure S2), consistent with their geographical and

environmental proximity (Figure 1B) and the range contraction during the LGM (MIS2

in Figure 2). The colonization of high-altitude regions in the central group is thus

accompanied by high levels of gene flow despite these populations ranging across a

large altitudinal gradient (2500m of altitude difference between LA1963 and LA3111

or LA2931). The divergence history results in the south-coast and south-highland

populations to be fairly isolated from one another (as separated by the Atacama

desert) leading to the suggestion of an incipient speciation process (Figure 2B;

Figure S2)[28]. We want to assess if and how many selective sweeps underpin such

adaptation to novel habitats (colonization of the coastal and highlands) and to

temporally variable climatic conditions (past climatic conditions).

Selective sweeps underpin local adaptation

To detect candidate genes under positive selection we scan the genomes for
selective sweep signals using both an SFS-based method (site-frequency spectrum
based) SweeD, and an LD-based method (linkage disequilibrium based) OmegaPlus.
To disentangle outlier regions putatively under selection from the neutral expected
distribution, we set conservative threshold values for the SweeD and OmegaPlus
statistics obtained from coalescent simulations with varying rates of recombination
based on the results of the demographic inference (Table 1; Figure S3). The
overlapping genomic regions under selection between the two methods are then
considered as the candidate regions under selection. In total, we find 2,921 regions
with SweeD (mean size 212,858bp +/- 3,938) and 13,106 with OmegaPlus (mean
size 59,618 bp +/- 521) across all six populations, yielding a total of 520 overlapping
regions (mean size 41,082 bp +/- 1,618). Within those regions we find a total of 799
protein-coding candidate genes detected as being likely under positive selection
(Data S1C). Among them, the largest number of candidate genes (354) is detected in
LA4330 (Table 1), likely because the population 1) has been established recently
(Figure 2B), 2) has a lower population size and thus higher amount of linkage
disequilibrium so that selective sweeps are easier to detect[29] (Figure S4C), and 3) its
habitat is very different from the rest of the species range (Figure 1B). In LA4107, we
find 61 candidate genes and about 100 candidate genes were detected in each of the
other four populations (Table 1). However, only a few candidate genes are shared
among different populations, with the central and south-highland populations sharing
a small number of candidate genes, while almost none are shared between the two
south-coastal populations (Figure S3C). This lack of common candidate genes
among populations is likely due to 1) the high nucleotide diversity and high effective
population sizes generating new variants across many genes which are then
differentially picked up by selection across different populations, and 2) the relatively
old inter-population divergence and timing of local adaptation.

Our candidate regions exhibit typical characteristics of positively selected

regions when compared to the genome-wide statistics, and do not show any peculiar

pattern of recombination (or lack thereof) which could invalidate our scan results (see

SI Text 2; Figure S4A-D; Table S3 and S4). Furthermore, we find an overlap between

our candidate genes under selective sweep and genes exhibiting signals of positive

selection in previous studies in *S. chilense*. It is noteworthy that these previous

studies were based on few chosen genes, different plants, different populations and

different sample sizes than ours. Among our candidate genes, we indeed find three

genes (*JERF*3, *TPP* and *CT*189) involved in abiotic stress tolerance such as salt,

drought or cold[19] as well as three NLRs (nucleotide binding leucine rich repeat,

SOLCI006592800, SOLCI001535800, SOLCI005342400) possibly linked to

resistance to pathogens[20]. We also find that two of the seven most up-regulated

genes under cold conditions in a transcriptomic study of *S. chilense*[21] do appear in

our selection scan in high altitude populations: *CBF*3 (Solyc03g026270) in LA2931,

and *CBF*1 (Solyc03g026280) in LA4330. These overlapping results indicate that our

genome-wide selective sweep scan generalizes the previous studies of selection in *S.*

chilense and support the functional relevance of our candidate genes.

We use the function integrated in the program McSwan to detect sweeps and

compute their age, that is the time of the appearance of the selected allele (Figure 3).

Conservatively, we select only sweeps detected with McSwan, SweeD and

OmegaPlus (Table 1). A total of 112 sweep regions (mean size 22,171 bp +/- 3,240)

overlap with our candidate regions from SweeD and OmegaPlus, containing 175

genes and with a mean sweep age of ca. 61,000 years. The ages of sweeps range

from as early as 150 kya to 5 kya, and only the highland populations exhibit sweeps

more recent than 15 kya, consistent with recent (re)colonization of higher altitudes

(Figure 3). Contrary to previous observations in other species (see introduction), we

find remarkably old sweep ages. We explain this result by the large effective

population sizes (per population in Figure 2, but also species wide[19,30]), mild

bottlenecks during colonization (Figure 2A) and the presence of strong seed banks[30].

Persistent seed banks indeed decrease the effect of genetic drift[31] and allow to retain

for longer periods of time the footprints of selection[32]. We then further investigate the

function of our candidate genes.

Gene regulatory networks underlying local adaptation in *S. chilense*

A Gene Ontology (GO) enrichment analysis of the 799 candidate genes reveals

significantly different and common biological processes across populations. We find

common GO categories to all populations such as basic cell metabolism, immune

response, specific organ development and response to external stimuli (Figure S5).

Most interesting, however, are four GO categories specifically found in several

populations and link to local differences in habitats: (i) root hair cell differentiation

functions are enriched in 15 candidate genes, only in the three coastal populations

(LA1963, LA2932 and LA4107); (ii) response to circadian rhythm, photoperiodicity

and flowering time are enriched in 12 candidate genes in two high-altitude (LA3111

and LA4330) and a south-coast (LA2932) populations; (iii) vernalization response is

enriched in eight candidate genes in the three high-altitude populations (LA2931,

LA3111, LA4330), and (iv) protein lipidation is enriched in seven candidate genes in

the south-highland population (LA4330). Based on the wealth of available data in

cultivated tomato and *S. pennellii* (with a well annotated genome) as well as gene

ontology in *A. thaliana*, we further study the gene regulatory networks to which the

candidate genes belong.

For adaptation to high-altitude conditions, 15 candidate genes are found

interconnected in a flowering gene network, which is itself sub-divided into two sub-

networks related to flowering, photoperiod and vernalization control pathways (Figure

4A; Data S1D). Photoperiod responsive genes can sense changes in sunlight and

affect the circadian rhythm to regulate plant flowering[33,34], while vernalization genes

regulate the flowering and germination through long-term low temperature [35–37].

These two sub-networks are connected through several key genes, some of which

appearing as candidate genes entailing local adaptation in our populations: FL

FLOWERING LOCUS C (FLC or AGL25), FLOWERING LOCUS T (FT) and

AGAMOUS-LIKE genes (AGL) (Figure 4A and 4B). These key genes are essential

regulators acting on the flowering regulation pathway[38–41]. Remarkably, some

candidate genes in the recently diverged south-highland population (LA4330)

aggregate into an independent network involved in circadian rhythm regulation,

connected to the photoperiod network by JUMONJI DOMAIN CONTAINING 5

(JMJD5) also a candidate gene in LA3111 (Figure 4A). In the central-highland

population (LA3111) several other candidate genes of the photoperiod network also

regulate circadian rhythm and flowering time. Interestingly, the three high-altitude

populations (LA3111, LA2931, and LA4330) present candidate genes of the

AGAMOUS-LIKE (AGL) gene family in the vernalization network (Figure 4A). We

also note that the network of protein lipidation genes appears to be related to the

synthesis of fatty acids in the south-highland population (Figure 4D; Data S1D). We

speculate here that this latter adaptation may be related to adaptation to lowest-

temperature stress of LA4330 (Data S1E)[42,43]. Adaptation to high altitude involves the

regulation of the flowering, including photoperiod and vernalization pathways, but

through different genes in different populations, while cold stress and its

consequence (adaptation in lipidation pathway) may be relevant for adaptation to the

highest altitudes.

Regarding adaptation to coastal conditions, we find 11 candidate genes related

to root development and cellular homeostasis functions clustered in a single network

(Figure 4C; Data S1D). We speculate that the drought and water shortage typical of

the coastal conditions (Data S1E) would promote the differentiation and extension of

plant roots[44,45]. The cell WALL ASSOCIATED LINASE 4 (WAK4), a candidate gene

identified LA4107, is shown to act as linker of signal from the cell wall to plasma

membrane and thus serve a vital role in lateral root development[46,47]. It is also

possible that the soft soil at the seaside is also beneficial to the extension of root

hairs to increase the water absorption area[48,49]. In addition to root development, we

find also genes involved in cell homeostasis (Figure 4C; Data S1D), which would be

critical for the coastal drought and salinity conditions to maintain stability of

intracellular environment in the coastal habitats[50,51].

Candidate genes show genotype-environment associations to local climatic

conditions

Our candidate loci are hypothesized to be responsible for adaptation to local climatic

conditions, so we test for genotype-environment association (GEA) using redundancy

analysis (RDA). RDA tests for correlation between allele frequencies and climatic

variables and can be used 1) to search more finely for specific selective variants

within the genomic sweep regions, and 2) reveal selective gradients defined by

multiple environmental variables[52,53]. We here perform first a "present day" RDA

using 144,713 SNPs from all candidate regions and 63 climatic variables

representing current (present) conditions for temperature, precipitation, solar

radiation, and wind (Data S1E). We find that the two-first RDA axes are significant

(ANOVA's $p < 0.001$) and retain most (38% and 21%) of the putative adaptive

genetic variance identified in the genome scans in all populations (Figure 5A). Table

S5 and S6 summarize outlier SNPs in different RDA models and their correlation with

climatic variables. In concordance to the PCAs of both climatic and genomic variation

(Figures 1B and 1C), the two main RDA axes cluster the individuals into three groups

corresponding to the main geographical regions (central, south-highland, and south-

coastal) supporting that those axes synthetize the principal selective pressures for

spatial local adaptation along the species distribution (Figure 5A). RDA1 represents

the differentiation of the two south-coast populations in correlation with higher

precipitation of coldest quarter (Bio19) and annual variation of solar radiation (CV_R).

There are 480 and 501 SNPs strongly associated with these two climatic variables,

respectively (Table S6). RDA2 summarizes a climatic gradient differentiating the

south-highland population manly driven by annual potential evapotranspiration

(annualPET) and temperature annual range (Bio7) with 1,184 and 372 strongly

associated SNPs, respectively (Table S6).

Further RDA analyses based on gene variants of the GO categories circadian

rhythm-photoperiodism, vernalization, root-hair differentiation, and protein lipidation

highlight combinations of climatic variables and genetic variants related to local

spatial adaptation (Figure S6A, 6C, 6E and 6G). These analyses show that about 40%

of the variation is explained by the two main RDA axes. Climatic variables

representing temperature variability through the year such as temperature

seasonality (Bio4) and temperature annual range (Bio7) are consistently correlated

with adaptive variation of the south-highland population (Figure 5A; Figure S6). A

total of 68 SNPs within candidate genes of the population LA4330 are strongly

associated to these two variables (Bio4, Bio7) in three of the RDA based on the GO

categories (circadian rhythm-photoperiodism, vernalization, and protein lipidation;

Data S1F). The RDA based on the root-hair differentiation GO category shows a

strong differentiation between lowland and highland populations, with stronger

correlation for the southern populations (Figure S6E). This differentiation correlates

with atmosphere water vapor availability variables such as annual minimum water

vapor pressure (ann_Vmin) and annual actual evapotranspiration (ann_AET)

variables for the coastal populations.

Considering that the sweep age estimations indicate that a fraction of the

adaptive variation currently observed in *S. chilense* may have originated from the

influence of past climate change processes (Figure 3), we implement an "LGM" RDA

using 37 climate variables projected to the Last Glacial Maximum conditions (Figure

5B; Figure S6B, 6D, 6F and 6H; Data S1E). This analysis implies the assumption that

the species did not shift its geographical range following the late Pleistocene climatic

oscillations, but rather underwent a shift in its niche while retaining large portions of

its distribution[54,55]. This process is especially relevant for the highland regions where

past distribution projections show no suitable habitat, while population persistence

being likely since the demographic reconstruction indicates 1) only mild population

size reductions (Figure 2C), and 2) pre-LGM divergence of the highland populations

(LA3111 and LA2931 during MIS3-4 climatic periods, Figure 2B). Therefore, we

expect that this analysis can uncover additional genomic variation selected in

response to temporal climatic changes and underlying the niche expansion towards

highly differentiated conditions in the south-highland region (Figure 1B). RDA

analyses using LGM climatic variables capture a smaller proportion of the genetic

variability in the first two constrained axes (30%) compared to that using the current

climatic variables. We find that some outlier/candidate SNPs appear not to show any

correlation with the past climatic variables, but about 30% outlier SNPs are identified

in additional genomic regions correlated with past climatic variables and not with the

current variables (Table S5 and S6). The analyses also emphasize the differentiation

of central, south-highland, and south-coast populations. Indeed, we find a different

RDA for vernalization candidates using the two sets of (present and past) climatic

variables. For example, the central populations LA3111 and LA2931 are separated in

the past RDA of vernalization using LGM climatic variables indicating warmer climate

after LGM may drive gene flow among central populations as seen in the current

RDA (Figure 2B; Figure S6C and S6D; Figure S3). The past RDA of LGM climatic

variables unveils that high-altitude populations, especially LA4330, have SNPs

correlating with temperature (*i.e.* annual mean minimum temperature; ann_mTmin,

and temperature annual range; Bio7) whereas coastal populations SNPs do correlate

with precipitation and potential evapotranspiration of the coldest and driest seasons

(Figure 5B).

Discussion

By correlating selective sweeps at candidate genes to current and past climatic data,

we demonstrate that local spatial adaptation to novel habitats and temporal

adaptation to changing conditions are intermingled. Nevertheless, our results show

that it is possible to separate the underpinnings of the spatial and temporal

adaptation in a species with large effective population size using niche modeling and

past climatic reconstruction. As a word of caution, we note that our discussion

focuses on four main GO categories which can be reliably associated with

physiological traits likely underlying adaptation (though pinpointing the regulatory or

coding SNPs under selection was not possible with our sample sizes), while

functional information on many candidate genes in Figures 3 and S5 is still lacking to

provide a complete picture. Yet, we are confident that our candidate genes under

selection are functionally relevant, as demonstrated by the overlap with previous

studies focusing on few genes involved in abiotic and biotic stress tolerance and

transcriptomic data under cold conditions[19–21].

For each of the five past climatic periods depicted in Figures 2 and 3, we

show different temporally restricted bursts of sweeps for either the south-highland or

south-coastal groups of populations. When overlaying specifically the age of the four

gene network categories of candidate genes (photoperiod, vernalization, protein

lipidation, and root hair differentiation), we find that the sweeps within a given

category/gene network are found at the time of a given particular climatic period

(Figure 3). This demonstrates that selective sweeps occur following temporal

adaptation in response to variable climatic conditions, as suggested under the

adaptation to a moving optimum theory[1]. We observe that during the colder climate

phases (MIS2 at 30-15kya), the suitable areas of the species decreased to a narrow

belt along lower altitude areas (Figure 2C). The persistence of populations at higher

altitude would have been possible by niche expansion driven by selection at genes

involved in the vernalization gene networks (Figure 3). We also observe a correlation

between the age of the sweeps and our "LGM" RDA analysis based on the past

climatic data. Therefore, we suggest the photoperiod and protein lipidation

adaptations to have occurred before the LGM (Figure 3) and to be instrumental pre-

adaptations favoring subsequently the colonization of new habitats in the southern

highlands (after the LGM). During this most recent colonization event, sweeps

occurred at other genes, including the complete photoperiod gene sub-network in

LA4330 (Figure 4A). The mentioned pre-adaptation sweeps could have arisen in the

ancestral populations from the highland-central region and by gene flow have

facilitated the recent expansion of the species' geographic range[56]. We do not detect

these sweeps in the central-highland populations probably because these haplotypes

can be broken by the combined effect of recombination (more efficient in central

populations with higher N_e) and gene flow. In addition, adaptations in the photoperiod

and protein lipidation gene networks in the highland populations are adaptations

older than the LGM, presumably due to earlier colonization of higher altitude habitats

under the warmer climate of the period MIS3-4 (Figure 3). Therefore, we suggest that

temporal adaptation does constrain or facilitate spatial local adaptation at later time

steps. The respective contributions of spatial and temporal can be disentangled by

comparing the significant genes (and SNPs) found in the past and present RDA

analyses.

The *S. chilense* lineage likely originated from coastal habitats in southern Peru.

This explains the early divergence and southward colonization process,

accompanied by habitat fragmentation and contraction, which yields two highly

isolated populations (Figure 2C). The coastal colonization seems to involve fewer

sweeps than the adaptation to higher altitudes, for example a burst of selective

sweeps in genes related to root anatomical traits during the LGM period (Figures 3,

4C, S6E-F). However, some of the adaptive genomic signals in the coastal

populations could be blurred due to the older timing or stronger drift (due to habitat

fragmentation along the coast), or be incomplete or partial sweeps which we do not

detect with our current scans.

We tentatively suggest some answers to the questions highlighted in the

introduction on the basis of our empirical results. We find between 60 and 350

selective sweeps per population, but contrary to our naïve expectations, sweeps

have a large variance in age. Indeed, our initial assumption was that most sweeps in

the south populations would be recent and occurring during only the recent spatial

local adaptation to a new habitat. On the contrary, the resulting picture of local

adaptation we observe is that of a complex process involving several discrete

adaptive steps, each due to temporally restricted burst of selective sweeps within a

given gene network. This adaptive process can be traced back using a combination

of sweep age estimates, RDA analyses and past and present climatic models. We

suggest that several sweeps do occur concomitantly in a given gene

pathway/network at a given time period, possibly as a temporal adaptation to a

moving environmental optimum (our climatic periods). Thereafter, these sweeps in a

given gene (sub-)network may lay the foundations, in other words, provide

exaptation[57,58], for colonizing new habitats at future climatic periods when new

ecological niches/habitats become available. The enabled future spatial adaptation

seems to involve selection at genes in different gene networks (possibly improving

upon and fine-tuning the previous phenotypic adaptation). Note that we focus here on

selective sweeps resulting from strong positive selection as we cannot assess in our

data the occurrence of weaker positive or polygenic selection. It appears that few

gene networks seem to be under strong selection at any climatic period, as predicted

under polygenic selection model of adaptation to a moving optimum[1,3,4,16]. We

speculate further, with the aim to broaden the current theoretical framework, that

strong selection underlying temporal adaptation to a moving optimum would occur at

one (or few) phenotypic trait, while local spatial adaptation to a new optimum requires

strong selection at several traits.

To our knowledge, this is the first study to attempt to disentangle the complex

spatial and temporal processes involved in colonization to new highly stressful

environments such as hyper-dry high-mountain regions. In the wild tomato *Solanum*

chilense, the recruitment of multiple genes belonging to gene networks responsible

for regulating the circadian cycle and flowering time is evident. This divergent selection process arises in response to recent climate change processes. The detectability of this polygenic pattern of selection is favored by the recent divergence of the populations involved, persistence of high genetic variability due to mild bottleneck during the expansion process and presence of seed banks, as well as the use of a combination of genome scans and availability of past climatic data. We demonstrate the power to use past climatic reconstruction with scans for selection to detect the underpinnings of local and temporal adaptation, when old selective sweeps can be recovered. Specifically, adaptation in space and time occurs over short time windows during which few genes of a given gene network are strongly selected upon, consistent with a polygenic model of adaptation to a moving optimum.

Acknowledgments

AT acknowledges funding from DFG (Deutscheforschungsgemeinschaft) Grant Number: 317616126 (TE809/7-1); KW was funded by the Chinese Scholarship Council, and GSA funded by the Technical University of Munich. We thank the Tomato Genetics Resource Center (TGRC) of the University of California, Davis for generously providing us with the seeds of the accession included in this study. We thank Daniela Scheikl for assistance with plant work and Christine Würmser for the Illumina sequencing.

Authors   contributions

Conceptualization, GSA and AT; Software, Formal analysis and Investigation, KW and GSA, Writing KW, GSA and AT; Funding Acquisition, AT; Supervision, GSA and AT.

Declaration   of   interests

The authors declare no competing interests

Resource availability

Lead contact

Further information on materials, datasets, and protocols should be directed to and

will be fulfilled by the Lead Contact, Aurélien Tellier (tellier@wzw.tum.de).

Data and code availability

The raw pair-end sequencing genomic data used in all the analyses can be accessed

at the European Nucleotide Archive (ENA) project accession PRJEB47577.

All code used in this study and other previously published genomic data is available

at the sources referenced in the key resource table.

Figure legends

Figure 1. Geographic distribution and population structure of *S. chilense.* (A) Map
with distribution of all *S. chilense* populations by the TGRC, the six *S. chilense*
populations in this study (black circles), and the four population groups (colorful
circles). (B) The PCA of all populations using 63 current climatic variables (Data S1E).
Population structure using SNP data based on (C) PCA and (D) Admixture analysis
(optimal K value is 4, Figure S1D).

529
530

Figure 2. Demographic history and species distribution model of *S. chilense* for
current and Last Glacial Maximum (LGM) climate conditions (A) The estimation of
historical patterns of effective population size (N_e) for 10 pairwise genome
comparisons per population using the MSMC model. (B) Interpreted demographic
scenario for the six samples populations of *S. chilense* including likely estimations of
effective population size, divergence times and gene-flow. The width of the boxes
represents the relative effective population size, arrows represent the migration
between population pairs. Grey background boxes indicate five Marine isotope
stages (MIS) climatic periods. (C) Overlay of the reconstruction of the distribution
model for *S. chilense* using current climatic variables (red) and LGM past climatic
variables (blue). Darker color of the gradient indicates higher suitable habitat for a
given climatic period.

Figure 3. Distribution of sweep age across the five MIS climatic periods.

Figure 4. Interaction genetic networks of candidate genes. (A) The network of
flowering regulation involved two sub-networks, photoperiod and vernalization
pathways, for regulation of flowering. (B) The schematic diagram of flowering
regulation involved photoperiod and vernalization is adapted from [36]. "⊣" indicates
repressive effects on gene expression; "→" indicates promotive effects on gene
expression. (C) The network of root development and cell homeostasis. (D) The
networks of protein lipidation. Connections represent gene interactions based on
physical interactions, informatics predictions and co-expression analyses.
Connection thickness is proportional to weighted value of the connected genes. The
black lines connected two sub-networks, genes under selection were connected by
solid lines and other genes were connected by dashed line. Node colors correspond
to genes were detected the different populations in genome scans. Gray circles, not
detected in genome scan, but present in *S. chilense*; gray squares: not present in *S.*
chilense.

Figure 5. Redundancy analysis (RDA) ordination bi-plots between the climatic
variables, populations and the genetic variants in all candidate sweeps. (A) RDA
using current climatic variables. (B) RDA using LGM climatic variables. Arrows
indicated the direction and magnitude of variables correlated with the populations.
The abbreviations of climatic variables are shown in Data S1E.
567

Table 1. Summary of genome scan results and estimation of sweep age.

Populations	Genome Scans				Sweeps age			
	N_{SweeD}	$N_{OmegaPlus}$	$N_{overlaps1}$	N_{genes1}	N_{McSwan}	$N_{overlaps2}$	N_{genes2}	Age_{mean} (kyr)
						16		
LA1963	385	2,474	98	86	267	39	14	71±55
LA2931	517	2,268	109	125	355	19	26	62±41
LA2932	374	1,717	46	101	302	15	29	65±48
LA3111	663	2,307	105	107	377	22	22	60±51
LA4107	203	2,047	37	61	194	10	10	63±40
LA4330	779	2,293	125	354	438	39	74	41±30

N_{SweeD}, number of outlier regions from SweeD; $N_{OmegaPlus}$, number of outlier regions from OmegaPlus; $N_{overlaps1}$, number of overlapping regions between SweeD and OmegaPlus; N_{genes1}, number of candidate genes in overlaps1, and all candidate genes show in Data S1C; N_{McSwan}, number of outlier regions from McSwan; $N_{overlaps2}$, number of overlapping regions between McSwan and overlaps1; N_{genes2}, number of genes in overlaps2; Age_{mean}, mean age of overlaps2.

STAR Methods

Sample collection, sequencing and bioinformatics

Plants were grown from individual seeds obtained from the Tomato Genetics

Resource Center (TGRC, University of California, Davis, USA;

http://tgrc.ucdavis.edu). We sampled five plants from accessions LA1963, LA3111,

LA2931, LA4330, LA2932, LA4107; Table S1) which were grown in individual pots in

standard glasshouse conditions at the Dürnast plant research facilities of the TUM

School of Life Sciences. Genomic DNA was extracted using the DNA extraction kit

from Quiagen and sequenced on an Illumina HiSeq 2500 with standard library size of

300 bp (Eurofins Genomics, Germany). The 30 *S. chilense* whole genome

sequencing data are available on ENA in BioProject PRJEB47577.

Quality control of the raw reads was performed with FastQC (v0.11.6)[59]. Base

calls with insufficient quality or showing adapter contamination were removed using

Trimmomatic (v0.36) using default parameters[60]. The high-quality clean reads were

mapped to the *Solanum pennellii* reference genome[61] available from Solanaceae

Genomics Network using Burrows-Wheeler Alignment tool (v0.7.16) with default

settings [62] and afterward sorted with Samtools (v1.5) [63]. The raw alignments were

then processed to add read groups, mark duplicates and fix mates using GATK (v3.3)

suite[64] and PicardTools (v2.10.6)[65]. Variant calling was performed using the

HaplotypeCaller tool of GATK with default parameters for each sample and individual

genomic variant files were then combined into a variant matrix with the

GenotypeGVCFs tool. Finally, after filtering, the variants were annotated using the

snpEff software (v4.3) with default settings[66]. For each SNP, the annotation were

identified based on the gene annotation of the *S.pennellii* reference. For details on

commands and parameters used see Supplementary SI text 1.

 Population genetics analyses

For all population genetics analyses we use *S. pennellii* population LA716 as

outgroup. The phylogenetic tree was constructed by the maximum likelihood (ML)

method using SNPhylo pipeline (v20160204) based on all SNPs[67]. The reliability of

each branch was evaluated by bootstrapping with 1,000 replicates and the tree was

visualized with iTOL (v5.0)[68]. A principal component analysis (PCA) was performed

for the whole SNP dataset using GCTA (v1.91.4) [69]. The inference of population

structure was performed using the program ADMIXTURE (v1.3.0)[70]. Nine scenarios

(ranging from K = 2 to K = 10) were assessed for genetic clustering.

Population genetics statistics, nucleotide diversity (π), Tajima's D and F_{ST},

were calculated with VCFtools (v1.15)[71] for each population (or pairs of populations)

over 100 kb sliding non-overlapping windows. Watterson's estimator of theta (θ_w) and

Fay and Wu's H were calculated using ANGSD (v0.921)[72]. The linkage disequilibrium

(LD) levels were calculated per population as the genotype correlation coefficient (r^2)

between two loci (within the same chromosome) using VCFtools with a maximum

distance of 1,000 kb. LD decay was then assessed for all pairs of SNPs within 1,000

kb distance using PopLDdecay (v3.27)[73].

Inference of demographic history

The demographic analysis was conducted using the Multiple Sequentially Markovian

Coalescent method (MSMC2) with phased VCF files and 40 hidden states[74]. The

cross-coalescence analysis was performed for each pairwise comparison of

genomes between pairs of populations to estimate the population separation history

and the migration rate with MSMC-IM[75]. The phased data were generated with

SHAPEIT (v2.17)[76]. We assumed generation time of 5 years and a rate of 1×10^{-8}

mutations per generation to scale time in the result plots.

Modeling present and past species distribution

We reconstructed and visualized the environmental space occupied by *S. chilense*

extracting the environmental conditions at the occurrence points and summarize

them by PCA[77]. The environmental data include 63 climatic layers obtained from

three public databases WorldClim2[78], ENVIREM[79], and the Consultative Group on

International Agricultural Research (CGIAR)[80] (Data S1E). The PCA was performed

by the *prcomp* function in R (v4.0.3)[81].

We then performed an ensemble modeling framework[82] using the BIOMOD

package[83,84] in R including all known localities (from TGRC database) covering the

full extant geographic range of *S. chilense* using eight modeling algorithms, five

cross-validation replicates, and ten pseudo-absence sampling sets, therefore

completing a total of 400 models. The environmental predictors include five variables

selected from those used in the climatic PCA. Variable selection was implemented by

keeping low correlated predictors. We assessed the predictive performance of the

models through the AUC and TSS statistics. Consensus niche models were obtained

using a TSS-weighted average method to account for the predictive power of each

fitted model. Models with low predictive power (TSS < 0.7) were discarded. All fitted

suitability models were then projected to infer the distribution of the *S. chilense*

suitable habitat under current and past conditions during the Last Glacial Maximum

(LGM; ~21 Kya).

Genome-wide selection scans

We identified selective sweeps using biallelic SNPs by SweeD (v3.3.1)[85] and

OmegaPlus (v2.3.0)[86]. The CLR statistics in SweeD were calculated with default

parameters except for using a resolution of 10 kb intervals within each chromosome.

OmegaPlus statistics (ω) were computed at 10 kb intervals. We specified a minimum

window of 10 kb and a maximum window of 100 kb to be used for computing LD

values between SNPs, respectively. Outlier CLR and ω statistics indicative of a

selective sweep are defined by comparison to the genome-wide distribution values

and the thresholds are defined by coalescent simulations of the the inferred

demographic history. We used the coalescent simulator SCRM (v1.7.3)[87] to generate

10,000 neutral datasets of 10 Mb based on the demographic history of each

population. To reduce false positives, we set up simulations with varying

recombination rate every 100 kb within each 10 Mb simulated block. We then

extracted the overlap regions of the two methods using the genomic coordinates and

those regions were regarded as high confident selective sweep regions. The age of

sweeps was estimated using McSwan[88] (see the supplementary methods).

686   GO enrichment analysis and gene networks

Due to the lack of a complete gene function annotation database, we performed a

BLASTX against the NCBI database of non-redundant proteins (nr) screened for

green plants (e-value cutoff was 10^{-6}) and used Blast2GO to assign GO terms for

each gene identified in the genome scan analysis[89,90] as well as performed a blast to

the *A. thaliana* dataset TAIR10 separately to remove redundant terms[91]. The false

discovery rates (FDR) were calculated to estimate the extent to which genes were

enriched in given GO categories. P-values smaller than 0.05 were used as cutoff for

a significant level of correlation. For each of the genes enriched in specific biological

processes, we retrieved the interacting gene neighbors using GeneMANIA (v3.5.2)[92].

We generated aggregate interaction networks in GeneMANIA, based on physical

interactions, predicted and co-expression. Finally, we performed hierarchical

clustering and manually optimized the weighted value cutoff for displaying the gene

network. The functions of networks were extracted form GeneMANIA outputs with

FDR cutoff 0.05 (Data S1D).

Redundancy analysis (RDA) analysis

The RDA was then performed using the *rda* function from the vegan package implemented in R[93], modeling genotypes as a function of the same climatic predictor variables used for the niche reconstruction analyses, and producing constrained axes and representative predictors. The significance of RDA constrained axes was assessed using the *anova.cca* function and significant axes were then used to identify candidate loci (P < 0.001). Candidate loci were identified using 2.5 standard deviation as cutoff (two-tailed p-value = 0.012).

Supplementary   information

Available as a single pdf with additional figures, tables and methods.

References

1. Polechová, J., Barton, N., and Marion, G. (2009). Species' range: Adaptation in space and time. The American Naturalist *174*, E186–E204.

2. Chevin, L.-M., Martin, G., and Lenormand, T. (2010). Fisher's model and the genomics of adaptation: Restricted pleiotropy, heterogenous mutation, and parallel evolution. Evolution *64*, 3213–3231.

3. Jain, K., and Stephan, W. (2017). Rapid adaptation of a polygenic trait after a sudden environmental shift. Genetics *206*, 389–406.

4. Matuszewski, S., Hermisson, J., and Kopp, M. (2014). Fisher's geometric model with a moving optimum. Evolution *68*, 2571–2588.

5. Kim, Y., and Stephan, W. (2002). Detecting a local signature of genetic hitchhiking along a recombining chromosome. Genetics *160*, 765.

6. Smith, J.M., and Haigh, J. (1974). The hitch-hiking effect of a favourable gene. Genet. Res. *23*, 23–35.

7. Exposito-Alonso, M., Burbano, H.A., Bossdorf, O., Nielsen, R., and Weigel, D. (2019). Natural selection on the Arabidopsis thaliana genome in present and future climates. Nature *573*, 126–129.

8. Aguirre-Liguori, J.A., Ramírez-Barahona, S., and Gaut, B.S. (2021). The

evolutionary genomics of species' responses to climate change. Nat Ecol
Evol, 1–11.

9. Savolainen, O., Lascoux, M., and Merilä, J. (2013). Ecological genomics
of local adaptation. Nature Reviews Genetics *14*, 807–820.

10. Fagny, M., and Austerlitz, F. (2021). Polygenic adaptation: Integrating
population genetics and gene regulatory networks. Trends in Genetics *37*,
631–638.

11. Zou, Y.-P., Hou, X.-H., Wu, Q., Chen, J.-F., Li, Z.-W., Han, T.-S., Niu, X.-M.,
Yang, L., Xu, Y.-C., Zhang, J., et al. (2017). Adaptation of *Arabidopsis
thaliana* to the Yangtze River basin. Genome Biology *18*, 239.

12. Schrider, D.R., and Kern, A.D. (2017). Soft sweeps are the dominant
mode of adaptation in the human genome. Molecular Biology and
Evolution *34*, 1863–1877.

13. Garud, N.R., Messer, P.W., and Petrov, D.A. (2021). Detection of hard and
soft selective sweeps from *Drosophila melanogaster* population genomic
data. PLOS Genetics *17*, e1009373.

14. Stephan, W. (2016). Signatures of positive selection: From selective
sweeps at individual loci to subtle allele frequency changes in polygenic
adaptation. Molecular Ecology *25*, 79–88.

15. Li, H., and Stephan, W. (2006). Inferring the demographic history and rate
of adaptive substitution in *Drosophila*. PLOS Genetics *2*, e166.

16. Jain, K., and Stephan, W. (2017). Modes of rapid polygenic adaptation.
Molecular Biology and Evolution *34*, 3169–3175.

17. Barghi, N., Hermisson, J., and Schlötterer, C. (2020). Polygenic adaptation:
A unifying framework to understand positive selection. Nat Rev Genet *21*,
769–781.

18. Nakazato, T., Warren, D.L., and Moyle, L.C. (2010). Ecological and
geographic modes of species divergence in wild tomatoes. American
Journal of Botany *97*, 680–693.

19. Böndel, K.B., Lainer, H., Nosenko, T., Mboup, M., Tellier, A., and Stephan,
W. (2015). North–south colonization associated with local adaptation of
the wild tomato species *Solanum chilense*. Molecular Biology and
Evolution *32*, 2932–2943.

20. Stam, R., Silva-Arias, G.A., and Tellier, A. (2019). Subsets of NLR genes

show differential signatures of adaptation during colonization of new habitats. New Phytologist *224*, 367–379.

21. Nosenko, T., Böndel, K.B., Kumpfmüller, G., and Stephan, W. (2016). Adaptation to low temperatures in the wild tomato species *Solanum chilense*. Molecular Ecology *25*, 2853–2869.

22. Xia, H., Camus-Kulandaivelu, L., Stephan, W., Tellier, A., and Zhang, Z. (2010). Nucleotide diversity patterns of local adaptation at drought-related candidate genes in wild tomatoes. Molecular Ecology *19*, 4144–4154.

23. Fischer, I., Camus-Kulandaivelu, L., Allal, F., and Stephan, W. (2011). Adaptation to drought in two wild tomato species: the evolution of the *Asr* gene family. New Phytologist *190*, 1032–1044.

24. Böndel, K.B., Nosenko, T., and Stephan, W. (2018). Signatures of natural selection in abiotic stress-responsive genes of *Solanum chilense*. Royal Society Open Science *5*, 171198.

25. Stam, R., Nosenko, T., Hörger, A.C., Stephan, W., Seidel, M., Kuhn, J.M.M., Haberer, G., and Tellier, A. (2019). The *de novo* reference genome and transcriptome assemblies of the wild tomato species *Solanum chilense* highlights birth and death of NLR genes between tomato species. G3 *9*, 3933–3941.

26. Lisiecki, L.E., and Raymo, M.E. (2005). A Pliocene-Pleistocene stack of 57 globally distributed benthic δ18O records. Paleoceanography *20*.

27. Ritter, B., Wennrich, V., Medialdea, A., Brill, D., King, G., Schneiderwind, S., Niemann, K., Fernández-Galego, E., Diederich, J., Rolf, C., et al. (2019). Climatic fluctuations in the hyperarid core of the Atacama Desert during the past 215 ka. Sci Rep *9*, 5270.

28. Raduski, A.R., and Igić, B. (2021). Biosystematic studies on the status of *Solanum chilense*. American Journal of Botany *108*, 520–537.

29. Koropoulis, A., Alachiotis, N., and Pavlidis, P. (2020). Detecting positive selection in populations using genetic data. In (Humana, New York, NY), pp. 87–123.

30. Tellier, A., Laurent, S.J.Y., Lainer, H., Pavlidis, P., and Stephan, W. (2011). Inference of seed bank parameters in two wild tomato species using ecological and genetic data. Proceedings of the National Academy of Sciences *108*, 17052–17057.

31. Tellier, A. (2019). Persistent seed banking as eco-evolutionary determinant
of plant nucleotide diversity: novel population genetics insights. New
Phytologist *221*, 725–730.

32. Živković, D., and Tellier, A. (2018). All but sleeping? Consequences of soil
seed banks on neutral and selective diversity in plant species. In
Mathematical Modelling in Plant Biology, R. J. Morris, ed. (Springer
International Publishing), pp. 195–212.

33. Song, Y.H., Shim, J.S., Kinmonth-Schultz, H.A., and Imaizumi, T. (2015).
Photoperiodic flowering: time measurement mechanisms in leaves. Annual
review of plant biology *66*, 441–464.

34. Johansson, M., and Staiger, D. (2015). Time to flower: interplay between
photoperiod and the circadian clock. Journal of experimental botany *66*,
719–730.

35. Guo, X., Liu, D., and Chong, K. (2018). Cold signaling in plants: Insights
into mechanisms and regulation. Journal of integrative plant biology *60*,
745–756.

36. Xu, S., and Chong, K. (2018). Remembering winter through vernalisation.
Nature Plants *4*, 997–1009.

37. Iida, H., and Mähönen, A.P. (2020). Growth-mediated sensing of long-term
cold in plants. Nature *583*, 690–691.

38. Michaels, S.D., and Amasino, R.M. (1999). FLOWERING LOCUS C
encodes a novel MADS domain protein that acts as a repressor of
flowering. The Plant Cell *11*, 949–956.

39. Sheldon, C.C., Rouse, D.T., Finnegan, E.J., Peacock, W.J., and Dennis,
E.S. (2000). The molecular basis of vernalization: the central role of
FLOWERING LOCUS C (FLC). Proceedings of the National Academy of
Sciences *97*, 3753–3758.

40. Turck, F., Fornara, F., and Coupland, G. (2008). Regulation and identity of
Florigen: FLOWERING LOCUS T moves center stage. Annual Review of
Plant Biology *59*, 573–594.

41. Putterill, J., and Varkonyi-Gasic, E. (2016). FT and florigen long-distance
flowering control in plants. Current Opinion in Plant Biology *33*, 77–82.

42. Jiang, H., Zhang, X., Chen, X., Aramsangtienchai, P., Tong, Z., and Lin, H.
(2018). Protein lipidation: occurrence, mechanisms, biological functions,

and enabling technologies. Chemical reviews *118*, 919–988.

43. Maksimov, E.G., Mironov, K.S., Trofimova, M.S., Nechaeva, N.L.,
Todorenko, D.A., Klementiev, K.E., Tsoraev, G. V, Tyutyaev, E. V, Zorina,
A.A., and Feduraev, P. V (2017). Membrane fluidity controls redox-
regulated cold stress responses in cyanobacteria. Photosynthesis
research *133*, 215–223.

44. Li, X., Cai, W., Liu, Y., Li, H., Fu, L., Liu, Z., Xu, L., Liu, H., Xu, T., and
Xiong, Y. (2017). Differential TOR activation and cell proliferation in
Arabidopsis root and shoot apexes. PNAS *114*, 2765–2770.

45. Xiong, Y., McCormack, M., Li, L., Hall, Q., Xiang, C., and Sheen, J. (2013).
Glucose–TOR signalling reprograms the transcriptome and activates
meristems. Nature *496*, 181–186.

46. He, Z.-H., Cheeseman, I., He, D., and Kohorn, B.D. (1999). A cluster of
five cell wall-associated receptor kinase genes, Wak1–5, are expressed in
specific organs of *Arabidopsis*. Plant molecular biology *39*, 1189–1196.

47. Lally, D., Ingmire, P., Tong, H.-Y., and He, Z.-H. (2001). Antisense
expression of a cell wall–associated protein kinase, WAK4, inhibits cell
elongation and alters morphology. The Plant Cell *13*, 1317–1332.

48. White, R.G., and Kirkegaard, J.A. (2010). The distribution and abundance
of wheat roots in a dense, structured subsoil–implications for water uptake.
Plant, cell & environment *33*, 133–148.

49. Bengough, A.G., Loades, K., and McKenzie, B.M. (2016). Root hairs aid
soil penetration by anchoring the root surface to pore walls. Journal of
Experimental Botany *67*, 1071–1078.

50. Forni, C., Duca, D., and Glick, B.R. (2017). Mechanisms of plant response
to salt and drought stress and their alteration by rhizobacteria. Plant and
Soil *410*, 335–356.

51. Zhao, C., Zhang, H., Song, C., Zhu, J.-K., and Shabala, S. (2020).
Mechanisms of plant responses and adaptation to soil salinity. The
innovation *1*, 100017.

52. Capblancq, T., Luu, K., Blum, M.G.B., and Bazin, E. (2018). Evaluation of
redundancy analysis to identify signatures of local adaptation. Molecular
Ecology Resources *18*, 1223–1233.

53. Forester, B.R., Lasky, J.R., Wagner, H.H., and Urban, D.L. (2018).

Comparing methods for detecting multilocus adaptation with multivariate genotype–environment associations. Molecular Ecology *27*, 2215–2233.

54. Jezkova, T., Olah-Hemmings, V., and Riddle, B.R. (2011). Niche shifting in response to warming climate after the Last Glacial Maximum: Inference from genetic data and niche assessments in the chisel-toothed kangaroo rat (*Dipodomys microps*). Global Change Biology *17*, 3486–3502.

55. Sillero, N., Reis, M., Vieira, C.P., Vieira, J., and Morales-Hojas, R. (2014). Niche evolution and thermal adaptation in the temperate species *Drosophila americana*. Journal of Evolutionary Biology *27*, 1549–1561.

56. Kottler, E.J., Dickman, E.E., Sexton, J.P., Emery, N.C., and Franks, S.J. (2021). Draining the swamping hypothesis: Little evidence that gene flow reduces fitness at range edges. Trends in Ecology & Evolution *36*, 533–544.

57. Gould, S.J., and Vrba, E.S. (1982). Exaptation—a missing term in the science of form. Paleobiology *8*, 4–15.

58. Larson, G., Stephens, P.A., Tehrani, J.J., and Layton, R.H. (2013). Exapting exaptation. Trends in Ecology & Evolution *28*, 497–498.

59. Andrews, C. (2010). FastQC: A quality control tool for high throughput sequence data. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

60. Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence data. Bioinformatics *30*, 2114–2120.

61. Bolger, A., Scossa, F., Bolger, M.E., Lanz, C., Maumus, F., Tohge, T., Quesneville, H., Alseekh, S., Sørensen, I., Lichtenstein, G., et al. (2014). The genome of the stress-tolerant wild tomato species *Solanum pennellii*. Nature Genetics *46*, 1034–1038.

62. Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler transform. bioinformatics *25*, 1754–1760.

63. Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and Durbin, R. (2009). The Sequence alignment/map (SAM) format and SAMtools. Bioinformatics *25*, 2078–2079.

64. Auwera, G.A.V. der, Carneiro, M.O., Hartl, C., Poplin, R., Angel, G. del, Levy-Moonshine, A., Jordan, T., Shakir, K., Roazen, D., Thibault, J., et al. (2013). From fastq data to high-confidence variant calls: The Genome

Analysis Toolkit best practices pipeline. Current Protocols in Bioinformatics
43, 11.10.1-11.10.33.

65. Li, H. (2011). A statistical framework for SNP calling, mutation discovery,
association mapping and population genetical parameter estimation from
sequencing data. Bioinformatics *27*, 2987–2993.

66. Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., Land,
S.J., Lu, X., and Ruden, D.M. (2012). A program for annotating and
predicting the effects of single nucleotide polymorphisms, SnpEff. Fly *6*,
80–92.

67. Lee, T.-H., Guo, H., Wang, X., Kim, C., and Paterson, A.H. (2014).
SNPhylo: A pipeline to construct a phylogenetic tree from huge SNP data.
BMC Genomics *15*, 162.

68. Letunic, I., and Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: An
online tool for phylogenetic tree display and annotation. Nucleic Acids
Research *49*, W293–W296.

69. Yang, J., Lee, S.H., Goddard, M.E., and Visscher, P.M. (2011). GCTA: a
tool for genome-wide complex trait analysis. The American Journal of
Human Genetics *88*, 76–82.

70. Alexander, D.H., Novembre, J., and Lange, K. (2009). Fast model-based
estimation of ancestry in unrelated individuals. Genome research *19*,
1655–1664.

71. Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo,
M.A., Handsaker, R.E., Lunter, G., Marth, G.T., and Sherry, S.T. (2011).
The variant call format and VCFtools. Bioinformatics *27*, 2156–2158.

72. Korneliussen, T.S., Albrechtsen, A., and Nielsen, R. (2014). ANGSD:
analysis of next generation sequencing data. BMC bioinformatics *15*, 1–13.

73. Zhang, C., Dong, S.S., Xu, J.Y., He, W.M., and Yang, T.L. (2019).
PopLDdecay: A fast and effective tool for linkage disequilibrium decay
analysis based on variant call format files. Bioinformatics *35*.

74. Schiffels, S., and Durbin, R. (2014). Inferring human population size and
separation history from multiple genome sequences. Nature genetics *46*,
919–925.

75. Wang, K., Mathieson, I., O'Connell, J., and Schiffels, S. (2020). Tracking
human population structure through time from whole genome sequences.

PLoS genetics *16*, e1008552.

76. Delaneau, O., Marchini, J., and Zagury, J.-F. (2012). A linear complexity phasing method for thousands of genomes. Nature methods *9*, 179–181.

77. Legendre, P., and Legendre, L. (2012). Numerical ecology Third English edition. (Elsevier).

78. Fick, S.E., and Hijmans, R.J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas. International Journal of Climatology *37*, 4302–4315.

79. Title, P.O., and Bemmels, J.B. (2018). ENVIREM: An expanded set of bioclimatic and topographic variables increases flexibility and improves performance of ecological niche modeling. Ecography *41*, 291–307.

80. Trabucco, A., and Zomer, R.J. (2018). Global aridity index and potential evapotranspiration (ET0) climate database v2. CGIAR Consort Spat Inf.

81. R Core Team (2020). R: A language and environment for statistical computing (R Foundation for Statistical Computing).

82. Araujo, M., and New, M. (2007). Ensemble forecasting of species distributions. Trends in Ecology & Evolution *22*, 42–47.

83. Thuiller, W., Lafourcade, B., Engler, R., and Araújo, M.B. (2009). BIOMOD - a platform for ensemble forecasting of species distributions. Ecography *32*, 369–373.

84. Thuiller, W., Georges, D., and Engler, R. (2014). BIOMOD2: Ensemble platform for species distribution modeling. R package   version 3.1-62/r677.

85. Pavlidis, P., Živković, D., Stamatakis, A., and Alachiotis, N. (2013). SweeD: Likelihood-Based Detection of Selective Sweeps in Thousands of Genomes. Molecular Biology and Evolution *30*, 2224–2234.

86. Alachiotis, N., Stamatakis, A., and Pavlidis, P. (2012). OmegaPlus: a scalable tool for rapid detection of selective sweeps in whole-genome datasets. Bioinformatics *28*, 2274–2275.

87. Staab, P.R., Zhu, S., Metzler, D., and Lunter, G. (2015). scrm: Efficiently simulating long sequences using the approximated coalescent with recombination. Bioinformatics *31*, 1680–1682.

88. Tournebize, R., Poncet, V., Jakobsson, M., Vigouroux, Y., and Manel, S. (2019). McSwan: A joint site frequency spectrum method to detect and

date selective sweeps across multiple population genomes. Molecular
Ecology Resources *19*, 283–295.

89.Conesa, A., Götz, S., García-Gómez, J.M., Terol, J., Talón, M., and Robles,
M. (2005). Blast2GO: a universal tool for annotation, visualization and
analysis in functional genomics research. Bioinformatics *21*, 3674–3676.

90. Conesa, A., and Götz, S. (2008). Blast2GO: a comprehensive suite for
functional analysis in plant genomics. International journal of plant
genomics *2008*.

91. Berardini, T.Z., Reiser, L., Li, D., Mezheritsky, Y., Muller, R., Strait, E., and
Huala, E. (2015). The Arabidopsis information resource: making and
mining the "gold standard" annotated reference plant genome. genesis *53*,
474–485.

92. Warde-Farley, D., Donaldson, S.L., Comes, O., Zuberi, K., Badrawi, R.,
Chao, P., Franz, M., Grouios, C., Kazi, F., and Lopes, C.T. (2010). The
GeneMANIA prediction server: biological network integration for gene
prioritization and predicting gene function. Nucleic acids research *38*,
W214–W220.

93.Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'Hara,
R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., and Wagner, H.
(2015). VEGAN: Community ecology package. R package version 2.3-0.

990