SyNPL: Synthetic Notch pluripotent cell lines to monitor and manipulate cell
interactions *in vitro* and *in vivo*.

Running title: Engineering cell interactions

Mattias Malaguti[1,‡], Rosa Portero Migueles[1], Jennifer Annoh[1], Daina Sadurska[1,2], Guillaume Blin[1],
Sally Lowell[1,‡]

1. Centre for Regenerative Medicine, Institute for Stem Cell Research, School of Biological
Sciences, University of Edinburgh, 5 Little France Drive, Edinburgh EH16 4UU, UK
2. Current address: Sir William Dunn School of Pathology, University of Oxford, Oxford, United
Kingdom

‡ Authors for correspondence: mattias.malaguti@ed.ac.uk sally.lowell@ed.ac.uk

Keywords: Cell interaction; SynNotch; pluripotent stem cells; cell engineering, patterning.

ABSTRACT

Cell-cell interactions govern differentiation and cell competition in pluripotent cells during early
development, but the investigation of such processes is hindered by a lack of efficient analysis tools.
Here we introduce SyNPL: clonal pluripotent stem cell lines which employ optimised Synthetic Notch
(SynNotch) technology to report cell-cell interactions between engineered "sender" and "receiver"
cells in cultured pluripotent cells and chimaeric mouse embryos. A modular design makes it
straightforward to adapt the system for programming differentiation decisions non-cell-
autonomously in receiver cells in response to direct contact with sender cells. We demonstrate the
utility of this system by enforcing neuronal differentiation at the boundary between two cell
populations. In summary, we provide a new tool which could be used to identify cell interactions and
to profile changes in gene or protein expression that result from direct cell-cell contact with defined
cell populations in culture and in early embryos, and which can be adapted to generate synthetic
patterning of cell fate decisions.

INTRODUCTION

During embryogenesis, pluripotent cells undergo a series of cell fate decisions that are controlled by
interactions between epiblast cells, their early differentiated derivatives, and the surrounding
extraembryonic tissues (Arnold and Robertson, 2009; Nowotschin and Hadjantonakis, 2010; Rossant
and Tam, 2009). The transcriptional changes that accompany exit from pluripotency and
differentiation into specific cell types have been extensively characterised, and the long-range signals
that control these changes are now well understood (De Los Angeles et al., 2015; Kinoshita and Smith,
2018; Pera and Tam, 2010; Posfai et al., 2021; Tam and Loebel, 2007). Less is known about how early
developmental decisions are influenced by direct interactions of cells with their neighbours. It is well
established that cell-cell interactions play a key role in development (Dias et al., 2014; Gurdon, 1987;
Johnson and Ziomek, 1983; Schultz, 1985), but until recently there has been a paucity of molecular
and technological tools available to study these processes in detail in relevant settings (Nishida-Aoki
and Gujral, 2019; Yang et al., 2021).

Quantitative image analysis can be used to identify and infer the effect of neighbours on the properties
of cells of interest in fixed samples (Blin et al., 2018; Fischer et al., 2020; Forsyth et al., 2021; Toth et
al., 2018). We have recently developed a software suite for automated neighbour identification during
live imaging (Blin et al., 2019), which provides researchers with a further dimension to study the
effects on cell-cell interactions on cell fate decisions. Whilst live image analysis provides high-
resolution visual information, this approach is labour-intensive and only leads to neighbour
identification *a posteriori*.

The nascent field of Synthetic Developmental Biology (Davies, 2017; Ebrahimkhani and Ebisuya, 2019;
Ho and Morsut, 2021; Santorelli et al., 2019; Schlissel and Li, 2020) seeks to understand the
mechanisms of patterning and cell differentiation through the engineering of genetic circuits (Cachat
et al., 2016; Matsuda et al., 2015; Sekine et al., 2018). By re-engineering the Notch/Delta signalling
cascade, Lim and colleagues generated a synthetic circuit capable of reporting and manipulating cell-
cell interactions in real time (Morsut et al., 2016). In this system, a "sender" cell presenting an
extracellular membrane-bound antigen of interest is recognised by a "receiver" cell expressing a
chimaeric Synthetic Notch (SynNotch) receptor, composed of an extracellular antigen-recognition
domain, the Notch1 core transmembrane domain containing proteolytic cleavage sites, and an
intracellular synthetic effector domain (Fig. 1). Thanks to the modularity of the SynNotch circuitry, it
is therefore possible to interrogate and manipulate the effects of interactions between cell types of
interest.

SynNotch technology has been used for monitoring cell-cell interactions, generating synthetic
patterns, generating synthetic morphogen gradients, inducing contact-mediated gene editing, and
generating custom antigen receptor T-cells (Cho et al., 2018; Choe et al., 2021; He et al., 2017; Huang
et al., 2020; Roybal et al., 2016; Sgodda et al., 2020; Toda et al., 2018; Toda et al., 2020; Wang et al.,
2021). SynNotch technology has been established in *Drosophila* (He et al., 2017) as well as in
immortalised cell lines and differentiated cell types, but its potential in the study of mammalian
developmental events remains largely untapped.

Mouse embryonic stem cells (ESCs) can be differentiated into any cell type of interest *in vitro*, can give
rise to chimaeric embryos and can be used to establish transgenic mouse lines (Bradley et al., 1984;
Evans and Kaufman, 1981; Martin, 1981). Adapting the SynNotch system for use in mouse ESCs would
therefore permit monitoring and manipulation of cell-cell interactions in a developmental context
both *in vivo* and *in vitro*. The original system designed by (Morsut et al., 2016) used lentiviral
transduction of immortalised and primary cell lines, where transgene expression was driven from the
retroviral SFFV promoter. Lentiviral transduction can lead to multiple copy transgene integration in
mouse ESCs (Pfeifer et al., 2002), and the SFFV promoter is prone to silencing in mouse pluripotent
cells and their derivatives (Herbst et al., 2012; Pfaff et al., 2013; Wu et al., 2011), making this system
suboptimal for mouse ESCs.

In this study, we made a number of adaptations to the original SynNotch system (Morsut et al., 2016)
in order to establish clonal modular SynNotch pluripotent cell lines (SyNPL). We characterised the
properties of this SyNPL system by monitoring interactions between eGFP-expressing sender cells and
mCherry-inducible receiver cells *in vitro*, then showed that this system can report interactions
between neighbouring cells *in vivo* in chimaeric mouse embryos, that it can be used for synthetic
patterning, and that its flexible and modular design can be exploited to conveniently manipulate cell-
cell interactions and drive contact-mediated synthetic cell fate engineering.

RESULTS

Design of SyNPL ESCs

We set out to adapt the SynNotch system, which was previously established through viral transduction of immortalised mouse L929 fibroblasts and K562 erythroleukaemic cells (Morsut et al., 2016), for use in mouse ESCs. In this system, sender cells are labelled with membrane-tethered extracellular eGFP (Fig. 1A). Receiver cells constitutively express a SynNotch receptor composed of an anti-GFP nanobody (LaG17) (Fridy et al., 2014), the mouse Notch1 minimal transmembrane core (Uniprot: Q01705, residues 1427-1752), and a tetracycline transactivator (tTA) (Gossen and Bujard, 1992), and contain a tetracycline response element (TRE) promoter capable of driving *mCherry* expression in response to tTA binding (Fig. 1B). Interaction of eGFP on sender cells with the anti-GFP nanobody on receiver cells leads to cleavage of the Notch1 core, releasing the tTA which can translocate to the nucleus, bind to the TRE promoter and drive *mCherry* expression (Fig. 1C). In addition, we constitutively labelled receiver cells with a tagBFP-3xNLS construct (Fig. 1B,C) to conveniently identify them by fluorescence microscopy and flow cytometry even in the absence of a contact-dependent mCherry signal.

Our aims when adapting the SynNotch system were to generate ESC lines with low cell-cell variability, robust and sustained transgene expression, and a modular design to allow convenient transgene exchange. In order to avoid cell-cell variability we elected to generate clonal cell ESC lines with stable genomic integration of the SynNotch system components, delivering transgenes by electroporation rather than lentiviral transduction to reduce the risk of multi-copy integration (Boggs et al., 1986; Charrier et al., 2011; Pfeifer et al., 2002; Smithies et al., 1985). We sought to ensure uniform levels of transgene expression by screening clonal lines and/or targeting transgenes to specific genomic locations, and by replacing the silencing-prone SFFV retroviral promoter used by (Morsut et al., 2016) with CAG (Niwa et al., 1991) or mouse *Pgk1* (McBurney et al., 1991) promoters, which have been extensively characterised in mouse ESCs (Chen et al., 2011; Hong et al., 2007). Finally, we set out to introduce modularity to our system by generating a "landing platform" master cell line to allow recombination-mediated cassette exchange (RMCE) of transgenes of interest.

Generation of extracellular membrane-tethered eGFP-expressing sender ESCs

We first generated clonal sender cell lines expressing membrane-tethered extracellular eGFP. The CAG and mouse *Pgk1* promoters are both silencing-resistant promoters commonly used to drive ubiquitous transgene expression in ESCs. We set out to determine which of these promoters can generate viable sender cells with strong and uniform expression of membrane eGFP. We also wished to explore whether eGFP molecules with HA and Myc protein tags can retain "sender" function in pluripotent cells.

We electroporated mouse ESCs with four sender constructs, containing either CAG or mouse *Pgk1* promoters driving expression of either untagged or HA- and Myc-tagged eGFP fused to a membrane-tethering domain (Fig. 2A-D). We isolated and expanded 64 clonal lines derived from stable genomic integration of the four constructs, and screened them by flow cytometry, analysing median eGFP intensity (Fig. 2E,F), percentage of eGFP-positive cells (Fig. 2G,H) and eGFP distribution (Figs. S1, S2). Both CAG and *Pgk1* promoters are capable of driving high uniform expression of eGFP, and as expected, there is considerable variability in eGFP expression between clonal lines.

We selected one untagged eGFP sender clone (CmGP1) and one HA- and Myc-tagged eGFP sender clone (CHmGMP19), exhibiting high, uniform and similar levels eGFP expression (Fig. 2I) for further analysis. For both clones, the pattern of eGFP expression appeared to be consistent with membrane localisation, and, in the case of CHmGMP19, eGFP co-localised with Myc (Fig. 2J,K).

Generation of a safe harbour site landing pad master ESC line

To facilitate convenient and repeated modification of the genome, we generated a clonal ESC line
carrying a "landing pad" targeted to the *Rosa26* locus, a safe harbour site in the mouse genome
(Friedrich and Soriano, 1991). This landing pad contains a splice acceptor, the *Neo* (G418/geneticin
resistance) gene, and a CAG promoter driving expression of *mKate2-3xNLS*, which encodes a red
fluorescent protein with no evident phenotypic effect in mouse embryos (Malaguti et al., 2013;
Shcherbo et al., 2009). This entire cassette is flanked by two *attP50* sites, which allows for φC31
integrase-mediated recombination with cassettes flanked by two *attB53* sites (Huang et al., 2009;
Tosti et al., 2018) (Fig. S3A). After confirming insertion at the correct genomic locus (Fig. S3B), we
verified that all cells express high and uniform levels of mKate2-3xNLS (Fig. S3C,D). We named this cell
line EM35.

An "all-in-one" design fails to generate fully functional mCherry inducible receiver cells

We first set out to investigate whether it would be possible to target all transcriptional units required
for receiver cell activity to the *Rosa26* landing pad in EM35 ESCs, and whether this would lead to the
generation of functional receiver ESCs. The design and characterisation of the resulting cell lines is
explained in detail in Supplementary Methods and Figs. S4-S9. Briefly, we found that mCherry could,
as expected, be induced by subpopulations of tagBFP-positive receiver cells in response to interaction
with eGFP-positive sender cells, but that this particular receiver cell design was hampered by variable
levels of tagBFP, variable inducibility of mCherry, and low levels of the SynNotch receptor. We
conclude that the SynNotch receptor construct and *TRE-mCherry* cassette can function as expected in
ESCs, but that further modifications to the design are required to obtain a reliable contact-reporting
system.

A multi-step design produces fully functional mCherry-inducible receiver ESCs

We hypothesised that two independent events may be affecting mCherry inducibility during our initial
attempts to generate "all-in-one-locus" receiver cells (Figs. S4-S9). Firstly, we speculated that *mCherry*
and *tagBFP* transgenes may have been lost from the genome due to mitotic recombination (Stern,
1936) or due to errors in replication at similar DNA sequences in close proximity (*Pgk1* promoters,
bGHpA signals). Secondly, we suspected that the expression levels of the SynNotch receptor were not
sufficient for efficient mCherry induction (Fig, S9).

We planned to circumvent the issue of potential loss of DNA by physically separating the three
transcriptional units through random genomic integration of the SynNotch receptor cassette and of
the *tagBFP-3xNLS* cassette, and by removing identical DNA sequences. To attempt to increase levels
of SynNotch receptor and to obtain uniform levels of tagBFP-3xNLS, we added an internal ribosome
entry site (IRES) followed by *Ble* (zeocin resistance gene) downstream of the SynNotch receptor
sequence, and an IRES followed by *Hph* (hygromycin B resistance gene) downstream of the *tagBFP-*
3xNLS sequence (Fig. 3A).

In order to generate these cells, we first randomly integrated the SynNotch receptor construct into
the genome of EM35 landing pad ESCs (Fig. S10A). We screened 87 clones for Myc expression, and
selected two clones (35SRZ9, 35SRZ86) with high, uniform Myc expression (Fig. S10B). The levels of
Myc in these clones were higher than those observed in receiver cells generated with an all-in-one
design (clones SNCB+4 and SNCB-6) and higher than those observed in Myc-tagged sender cells
(CHmGMP19) (Fig. S10C). We then randomly integrated the *tagBFP-3xNLS* transgene into the genome
of these two clones (Fig. S10D). We screened a total of 36 clones and selected one for each parental

line with high, uniform expression of tagBFP-3xNLS (PSNB-A clone 10, PSNB-B clone 3) (Fig. S10E,F).
We renamed these lines PSNB (Parental SynNotch tagBFP) clones A and B respectively.

Next, we performed RMCE at the *Rosa26* landing pad in PSNB cells to replace the *mKate2* transgene
with one of three constructs: the *TRE-mCherry* cassette present in the SCNB construct, a *tetO-mCherry*
cassette with more tTA binding sequences elements in the inducible promoter (to test whether this
led to improved mCherry induction), or an empty vector cassette to generate tagBFP-positive mKate2-
and mCherry-negative control cell lines (Fig. S11A). We verified that integration of the empty vector
cassette led to loss of mKate2 expression, and used these control cell lines to confirm that tagBFP
signal was able to unambiguously identify receiver cells (Figs. 3B,C, S11B).

We then set out to test whether the new receiver cell lines containing inducible mCherry cassettes
expressed mCherry in response to co-culture with sender cells. We screened 27 clones for tagBFP and
mCherry expression by culturing them in the presence or absence of sender cells for 24 hours (Fig.
S11C-F). We observed that all genetically identical clones behaved very similarly, suggesting we were
not experiencing silencing or loss of DNA. Clones containing the larger *tetO-mCherry* cassette
exhibited high levels of mCherry leakiness in the absence of sender cells. Co-culture with sender cells
led to significant mCherry induction, but the distributions in the presence and absence of sender cells
overlapped significantly (Fig. S11D,F). Clones containing the smaller *TRE-mCherry* cassette exhibited
mCherry leakiness in the absence of sender cells, however co-culture with sender cells led to an
increase in mCherry expression to levels which displayed minimal overlap with those seen in cells
cultured in the absence of sender cells (Fig. S11C,E). We selected three clones with minimal leakiness
and high inducibility for downstream analysis (PSNBA-TRE1, PSNBB-TRE10, PSNBB-TRE9). We renamed
these cells STC (for SynNotch-TRE-mCherry) clones A1, B1 and B2 respectively. We verified that these
receiver lines could induce mCherry following cell-cell interaction with sender cells by co-culturing
sender and receiver cells together for 24 hours. We observed specific induction of mCherry by tagBFP-
positive receiver cells in contact with eGFP-positive sender cells (Fig. 3D). We used flow cytometry to
confirm that mCherry is robustly induced in the majority of tagBFP-positive receiver cells following co-
culture with 9-fold excess of CmGP1 sender cells at confluence (Fig 3B,C,E,F).

These observations suggest that physical separation of the three transcriptional units in the genome
of receiver cells, coupled to the use of internal ribosome entry sites and selectable markers within the
units, can lead to the generation of receiver ESC lines which exhibit clear and specific induction of
mCherry upon interaction with eGFP-expressing sender cells.

Extracellular membrane-tethered eGFP is required for contact-mediated transgene induction in
receiver cells

It would be useful to make use of existing GFP fluorescent reporter ESCs (for example, cell-state
reporters or signalling reporters) to act as sender cells, in order to test how particular cell states may
influence direct neighbours. However, many such cell lines make use of non-membrane-tethered GFP,
which is unlikely to interact with the anti-GFP nanobody on STC receiver cells. We therefore wished
to test whether membrane tethering of eGFP to the extracellular space was absolutely necessary for
effective neighbour-labelling.

We profiled mCherry levels in STC receiver cells by flow cytometry following 24 hours of co-culture
with "sender" cells carrying different transgenes (Fig. 3E-H). We cultured STC receiver cells alone (Fig.
3E), in the presence of CmGP1 sender cells (Fig. 3F), in the presence of a control cell line expressing
untagged intracellular eGFP (E14GIP1) (Fig. 3G), or in the presence of CmGP1 sender cells containing
an extra untagged intracellular eGFP transgene (CmGP1GH1) (Fig. 3H). We found that E14GIP1 cells,

which do not express membrane-tethered eGFP, did not induce mCherry above baseline levels in in
co-cultured STC receiver cells (Fig. 3E-G). Furthermore, we observed that mCherry is induced to similar
levels following co-culture with either CmGP1 or CmGP1GH1 sender cells (Fig. 3F,H), suggesting that
the additional untagged eGFP transgene in CmGP1GH1 cells does not interfere with mCherry induction
by the extracellular eGFP transgene. We conclude that cells containing intracellular GFP would not be
capable of functioning as sender cells unless supplemented with extracellular membrane tethered
eGFP.

Increasing sender:receiver cell ratios leads to increased transgene induction in receiver cells

We next set out to establish how differing sender:receiver cell ratios could affect the efficiency of
neighbour-labelling. We co-cultured STC receiver cells with different proportions of sender cells for 24
hours (Fig. S12). We observed that as few as 20% of sender cells were sufficient to induce mCherry in
approximately half of STC receiver cells, and that 90% sender cells could induce mCherry in over 90%
of STC receiver cells (Fig. S12A,B). mCherry fluorescence follows a bimodal distribution in receiver cells
exposed to "non-saturating" numbers of sender cells, and a unimodal distribution in receiver cells
exposed to "saturating" numbers of sender cells (Fig. S12C-H). This suggests that STC receiver cells
which have come into contact with sender cells can uniformly induce high levels of mCherry
expression, as observed by immunofluorescence when co-culturing cells at a 9:1 sender:receiver cell
ratio (Fig. S12I-K).

HA- and Myc-tags on eGFP do not affect sender cell function

An epitope tag fused to eGFP could be helpful for unequivocally identifying and isolating sender cells,
but carries the risk of interfering with sender function. Having generated sender cells with HA- and
Myc-tagged (CHmGMP19) and untagged (CmGP1) eGFP molecules, we sought to understand whether
tagged and untagged eGFP differed in their ability to induce transgene expression in receiver cells. We
therefore compared the ability of each sender line to induce mCherry expression in the previously
described experiment, where we cultured STC receiver cells with CmGP1 or CHmGMP19 sender cells
at varying sender:receiver cell ratios (Fig. S12). Both CmGP1 and CHmGMP19 sender cell lines can
drive mCherry induction to similar extents (Fig. S12A-H), suggesting that the LaG17 nanobody on the
SynNotch receptor can recognise both untagged and HA- and Myc-tagged eGFP, and that these tags
do not compromise sender function.

Kinetics of contact-dependent transgene induction in receiver cells

We next set out to ask how long sender cells need to be in contact with receiver cells in order to trigger
mCherry labelling, and to characterise the kinetics of mCherry induction. We filmed STC receiver cells
co-cultured with CmGP1GH1 sender cells for 24 hours (Fig. 4A, Movie 1). We first observed mCherry
induction 5-6 hours after initial sender-receiver contact (Fig. 4A, yellow arrowheads), with a rapid
increase in mCherry levels until the 16-hour timepoint, and a slower increase thereafter. We observed
STC receiver cells which did not make contact with sender cells and remained mCherry-negative (Fig.
4A, magenta arrowheads), and an STC receiver cell that made contact with a sender cell 2 hours before
the cells were fixed at the 24-hour timepoint for immunofluorescence, and which remained mCherry-
negative (Fig. 4A, cyan arrowheads).

We then quantified the kinetics of mCherry induction. We co-cultured 10% STC receiver cells with 90%
CmGP1 sender cells, and analysed mCherry expression by flow cytometry over the course of 72 hours
(Fig. 4B). mCherry is first induced at low levels at the 5-hour timepoint, and its expression keeps
increasing until the 48-hour timepoint (Figs. 4C,D, S13A,B).

mCherry protein maturation will introduce a time-lag between initiation of mCherry transcription and
the detection of mCherry fluorescence. For example, when analysing further timelapse experiments
(Fig. 4E, Movie 2), we noticed an example of an STC receiver cell which remained in contact with a
sender cell for 8 hours, lost contact for 12 hours following a cell division, but kept increasing its levels
of mCherry in the meantime (Fig. 4E, white arrowheads). Relying on direct detection of mCherry (Fig
4A-D) is therefore likely to overestimate the minimum duration of cell contact required for mCherry
induction.

We designed an experimental strategy to overcome this problem. We co-cultured 10% STC receiver
cells with 90% CmGP1 sender cells for various time points between 0 and 24 hours, then added
doxycycline to the culture medium for a further 16 hours (Fig. 4F). Doxycycline prevents tTA from
binding to *TRE* sequences (Gossen and Bujard, 1992), hence we expect doxycycline administration to
halt *mCherry* transcription in receiver cells while still allowing time for mCherry protein to mature.
This means that any mCherry signal observed after doxycycline administration should be ascribable to
cell-contact-dependent transcription that took place during the initial period of co-culture in
doxycycline-free medium. In this experimental setting, we observed low but detectable induction of
mCherry when cells had experienced only 2 hours of doxycycline-free co-culture (Figs. 4G,H, S13C,D).

These data collectively suggest that 2 hours of sender-receiver contact may be sufficient for induction
of low levels of mCherry, and that mCherry levels will keep increasing in receiver cells for a period of
time following the loss of sender-receiver contact. This neighbour-labelling system can therefore
identify receiver cells that have had relatively brief interactions with sender cells or which have
recently lost contact with sender cells.

Kinetics of contact-dependent mCherry perdurance in receiver cells

We next sought to establish how long mCherry signal persisted following loss of tTA-mediated
mCherry transcription. We co-cultured 10% STC receiver cells with 90% CmGP1GH1 cells for 24 hours,
then added doxycycline to the culture medium (to block the activity of tTA and halt mCherry
transcription) and filmed the cells over 48 hours (Fig. S14A,B, Movies 3,4). We observed little change
in mCherry fluorescence for the initial 10-12 hours, then a gradual decrease in signal until its
extinguishment around 38-40 hours after doxycycline addition (Fig. S14A,B). In order to quantify this
process, we co-cultured 10% STC receiver cells with 90% CmGP1 sender cells for 24 hours, then added
doxycycline to the culture medium and analysed mCherry fluorescence by flow cytometry at various
timepoints (Fig. S14C). No reduction of mCherry signal was observed for the initial 8 hours following
doxycycline administration, with median mCherry expression decreasing by approximately half by 16
hours, and mCherry signal returning to background levels within 48 hours (Fig. S14D-F).

Taken together, these results suggest that induction of mCherry occurs more rapidly than loss of
mCherry signal, presumably due to the high stability of this fluorescent protein, further confirming the
utility of this system for identifying recent as well as current cell-cell interactions.

The SyNPL SynNotch cell-cell interaction reporter is functional in early mouse embryos

Having comprehensively characterised the SynNotch system in clonal mouse ESC lines, we set out to
test whether the system could also function *in vivo* in early mouse embryos. We aggregated wild-type
morulae with CmGP1GH1 sender cells and/or STC receiver cells, and cultured these to the blastocyst
stage (Fig. 5). As expected, all chimaeric blastocysts (55/55) containing both sender and STC receiver
cells induced expression of mCherry, whereas no wild-type blastocysts nor blastocysts containing only
sender cells displayed mCherry expression (Fig. 5A,B). 12 out of 13 chimaeras containing STC receiver
cells alone did not express readily detectable levels of mCherry (Fig. 5B), in line with the low proportion

of mCherry-high cells observed *in vitro* in STC receiver cells cultured alone. All three STC clonal lines
reliably induced mCherry within chimaeric embryos that also contained CmGP1GH1 sender cells (Fig.
5B, Fig. S15). These results suggest that the SyNPL neighbour-labelling system is functional, efficient
and reliable *in vivo*.

Spatial confinement of sender and receiver cells leads to synthetic patterning

SynNotch technology has been successfully employed to generate synthetic patterns. Strategies to
achieve this include co-culturing cells in a low sender:receiver cell ratio in order to create 2-
dimensional activated receiver cell rings surrounding a clone of sender cells (Morsut et al., 2016),
creating self-organising cell aggregates through contact-mediated induction of adhesion molecules
(Toda et al., 2018), and recreating morphogen gradients through the use of anchor proteins to capture
diffusible receiver cell-activating signal (Toda et al., 2020).

We set out to test whether our clonal ESC lines could be used to generate a synthetic pattern: a stripe
of transgene expression at the region of contact between sender and receiver cells. To this end, we
plated CmGP1 sender and STC receiver cells in separate chambers of a removable multi-chamber cell
culture insert and allowed them to reach confluence. We then removed the insert, allowing cells to
grow in the space between chambers, make contact and induce transgene expression in receiver cells
(Fig. 6A). We assessed expression of mCherry 24 hours after initial sender:receiver contact, and
observed a distinct stripe of mCherry expression at the sender:receiver border (Fig. 6B). This result
suggest that SynNotch technology can be successfully employed in mouse ESCs to generate synthetic
patterns of gene expression.

Harnessing modularity of SyNPL SynNotch ESCs to synthetically alter cell fate

One advantage of introducing modularity to our SyNPL SynNotch system design is that we can
generate clonal receiver cell lines with inducible expression of any gene of interest. Having
characterised the system in mCherry-inducible STC receiver cells, we wished to test whether we could
alter receiver cell fate following interaction with sender cells by generating Neurog1-inducible receiver
cells.

The transcription factor Neurog1 (Neurogenin 1) is necessary for neural development in mice, where
it drives neuronal differentiation of progenitor cells (Cau et al., 2002; Ma et al., 1998; Yuan and Hassan,
2014). Ectopic expression of Neurog1 has been shown to lead to neuronal differentiation both in
ectoderm and, surprisingly, in mesoderm (Cai et al., 2000; Ma et al., 1996; Perez et al., 1999). This
ability of Neurog1 to induce differentiation in non-permissive conditions was exploited to drive
neuronal differentiation of mouse ESCs in pluripotent culture conditions (Velkey and O'Shea, 2013).
We therefore set out to ask whether contact-mediated induction of a *Neurog1* transgene could induce
neuronal differentiation specifically in receiver cells that come into contact with eGFP-expressing
sender cells.

Firstly, we generated STN (SynNotch TRE-Neurog1) receiver cells by performing RMCE at the *Rosa26*
landing pad in PSNB cell lines to replace the *mKate2-3xNLS* transgene with a *TRE-3xFlag-Neurog1*
cassette (Figs. 6C, S16A). We then tested whether we could drive contact-mediated induction of the
Neurog1 transgene by co-culturing STN receiver cells with CmGP1 sender cells for 48 hours, the
timepoint at which we observed maximum mCherry induction in STC receiver cells (Figs. 4C,D, S13A,B).
We confirmed this resulted in robust induction of 3xFlag-Neurog1 in STN receiver cells compared to
STN receiver cells cultured alone (Fig. S16B), which suggests that PSNB landing pad cell lines can be
used to generate receiver cells harbouring inducible transgenes other than mCherry.

We then set out to test whether we could induce contact-mediated neuronal differentiation of
receiver cells in pluripotent culture conditions, and whether we could engineer differentiation to occur
in a synthetic pattern. To this end, we repeated the synthetic stripe patterning experiment described
above (Fig. 6A,B), using STN receiver cells in place of STC receiver cells. We assessed the expression of
the neuronal marker Tubb3 96 hours after initial sender:receiver contact, and observed evident
induction of Tubb3 and acquisition of neuronal morphology by STN receiver cells at the
sender:receiver border (Fig. 6D).

We conclude that the interaction between eGFP-expressing sender cells and STN receiver cells can
lead to contact-mediated *Neurog1* induction and neuronal differentiation of receiver cells in non-
permissive culture conditions. This demonstrates that the SyNPL system can be readily used to
generate clonal ESC lines for contact-mediated induction of transgenes of interest, and that these cell
lines can in turn be used to manipulate cell-cell interactions in order to program synthetic cell fate
decisions in response to contact with a particular cell population at desired locations in space.

DISCUSSION

Engineering SynNotch machinery (Morsut et al., 2016) into pluripotent cells opens up many
opportunities for understanding how direct cell-cell interactions between neighbouring cells can
control differentiation decisions, mediate cell competition (Sancho et al., 2013) and orchestrate
morphogenesis (Gorfinkiel and Martinez Arias, 2021) as cells differentiate in 2D or 3D culture. Mouse
ES cells have the unique ability to contribute to chimaeric embryos, meaning that appropriately
engineered cell lines can also be used to understand and control cell-cell interactions during early
embryonic development. There are however particular challenges associated with engineering
existing SynNotch technologies into pluripotent cells. Here we describe how we overcame these
challenges to generate the SyNPL system: a set of clonal SynNotch "sender" and "receiver" mouse ES
cells engineered with optimised and modular SynNotch technology. We demonstrate the utility of the
SyNPL system for monitoring cell-cell interactions both in culture and in early mouse embryos, and
show that we can use this system to engineer contact-dependent cell fate decisions at the boundary
between two populations of pluripotent cells.

Properties of sender cells

We generated sender cell lines expressing high and uniform levels of extracellular membrane-tethered
eGFP. This transgenic construct was previously used for SynNotch sender cells (Morsut et al., 2016;
Sgodda et al., 2020), and comprises eGFP fused to an N-terminal mouse IgGK signal sequence and a C-
terminal human PDGFRB transmembrane domain. The addition of HA and Myc tags at the N- and C-
termini of eGFP did not affect the ability of sender cells to induce mCherry expression in STC receiver
cells (Fig. S12), suggesting that the antigenic region of eGFP recognised by the LaG17 anti-GFP
nanobody component of the SynNotch receptor is not masked by the HA and Myc protein tags.
Furthermore, the LaG17 anti-GFP nanobody can also bind to *Aequorea victoria* YFP, CFP and BFP, and
Aequorea macrodactyla CFP (Fridy et al., 2014), raising the interesting possibility that transgene
induction in our receiver cells could be induced by sender cells expressing different membrane-
tethered fluorophores.

Mouse ESCs and mouse lines labelled with lipid anchor-tethered GFP constructs have previously been
described (Kondoh et al., 1999; Nowotschin et al., 2009; Rhee et al., 2006; Shioi et al., 2011). It would
be interesting to test if these lines could function as SynNotch sender cells; this would require
extracellular GFP localisation and generation of sufficient tensile force upon receptor interaction

(Morsut et al., 2016). GPI anchors GFP to the outer leaflet of the plasma membrane (Rhee et al., 2006;
Sevcsik et al., 2015); GFP-GPI-labelled cells should therefore be capable of acting as sender cells. This
does indeed appear to be the case in *Drosophila* (He et al., 2017).

"All-in-one" locus receiver cells display suboptimal functionality

Our initial attempts at generating "all-in-one" *Rosa26*-targeted receiver cells (termed SNCB+ and
SNCB- cells) were unsuccessful. The large variation in tagBFP expression and low proportion of
mCherry-inducible ESCs we saw within all of our purportedly genetically identical clonal lines suggests
that either widespread transgene silencing or loss of DNA occurred at the *Rosa26* safe harbour locus
in pluripotent cells. Furthermore, the observation that rederivation of clonal lines following
fluorescence-activated cell sorting of single SNCB+ and SNCB- cells led to re-establishment of the initial
heterogeneous distributions of fluorophore expression (Figs. S7, S8) suggests that the all-in-one design
is not optimal for use in pluripotent cells. We were able to overcome these problems by switching to
a random-integration strategy, but it is possible that inclusion of IRES-antibiotic resistance cassettes
and/or insulator sequences may provide an alternative route towards generating a reliable system
without sacrificing the all-in-one-locus approach.

Landing pad ESCs provide system modularity

Targeting of a landing pad to the *Rosa26* safe harbour locus has been previously characterised as an
efficient strategy for the rapid generation of multiple cell lines through RMCE (Seibler et al., 2005;
Tchorz et al., 2012; Tosti et al., 2018). We found this to be true for our cell lines harbouring a *Rosa26-*
attP50-Neo-mKate2-attP50 landing pad, which we used to generate several different cell lines with
ease. The advantage of such a system is the ability to target different transgenes to the same genomic
locus in the same parental cell line. Our parental PSNB lines harbouring the *Rosa26* landing pad
allowed us to initially test the functionality of SynNotch in ESCs with an inducible *mCherry* transgene
in STC receiver cells, prior to generating genetically equivalent STN receiver cells with a *Neurog1*
transgene in place of *mCherry*. This modular design therefore makes it possible to readily switch
between using SyNPL for monitoring and profiling the consequences of defined cell-cell interactions
(based on contact-dependent mCherry expression) and using SyNPL for engineering contact-
dependent cell behaviours (based on contact-dependent expression of any cell behaviour-
determinant).

Describing the properties of the SyNPL system

We characterised various aspects of the SyNPL system that will help inform the experimental design
for users of these cells.

Previous studies have co-cultured senders and receiver cells at different ratios (ranging from 1:50 to
5:1), and for varying times (ranging from 10 minutes to several days) (Cho et al., 2018; Choe et al.,
2021; He et al., 2017; Huang et al., 2020; Luo et al., 2019; Matsunaga et al., 2020; Morsut et al., 2016;
Roybal et al., 2016; Sgodda et al., 2020; Srivastava et al., 2019; Toda et al., 2018; Toda et al., 2020;
Wang et al., 2021; Yang et al., 2020). We analysed transgene induction in STC receiver cells at 11
different sender:receiver cell ratios (ranging from 1:19 to 9:1), and observed that higher proportions
of sender cells in culture result in a higher proportion of receiver cells inducing mCherry. This is in line
with the results obtained by (Sgodda et al., 2020) when comparing three different sender:receiver cell
ratios, and with the observations of (Morsut et al., 2016), who exposed receiver cells to varying
concentrations of sender ligand. By finely varying the concentrations of sender ligand, Lim and
colleagues described the transgene induction response as sigmoidal (Morsut et al., 2016), which was

not evident in our data. It is however possible that by testing lower sender:receiver cell ratios this
might hold true in our SynNotch system too.

When analysing transgene expression within single samples, we found that mCherry distribution
follows a bimodal on/off response, indicative of the presence of receiver cells which do not interact
with sender cells at low sender:receiver cell ratios. This bimodal pattern of transgene induction is also
evident in data from Cantz and colleagues (Sgodda et al., 2020). The ability of individual sender cells
to induce mCherry induction in STC receiver cells (as seen in Movies 1 and 2) implies that this system
can be effectively employed to study the effect of interactions between receiver cells and individual
and/or rare sender cells in relevant model systems.

We performed a high-resolution study of the kinetics of mCherry induction and downregulation in STC
receiver cells. We observed low levels of mCherry induction in STC receiver cells following 2 hours of
co-culture with sender cells, provided we allowed time for subsequent protein maturation, and
observed maximum mCherry induction following 48 hours of co-culture. Previous studies making use
of lentiviral-delivered transgenes suggest that 10 minutes may be sufficient for transgene activation
in HEK293 receiver cells, and that 1 hour may be sufficient for transgene induction in L929 receiver
cells, as long as protein maturation time is allowed (Morsut et al., 2016; Sgodda et al., 2020). This is
significantly faster than what we observed in this study, and may be ascribable to lentiviral
transduction leading to higher levels of SynNotch receptor and/or integration of multiple transgene
copies compared to our clonal mouse ESC lines.

In our system, mCherry downregulation did not commence for at least 8 hours following simulated
loss of sender:receiver cell contact, with full loss of signal occurring after more than 40 hours. This is
in line with previous observations in L929 receiver cells, where inducible GFP transgene expression
was lost between 24 and 50 hours after sender cells were removed from culture (Morsut et al., 2016).
The kinetics of mCherry induction and downregulation suggest that this SynNotch system is suited for
the study of cell-cell interactions with a temporal range of hours rather than minutes, and that
"memory" of such interactions will persist for a few days. Should this persistence of mCherry signal
prove inconvenient for the study of particular processes, the PSNB landing pad parental cell lines can
be used to readily generate cell interaction reporter receiver cells harbouring destabilised inducible
transgenes with short half-lives.

Exploring the roles of cell-cell interactions *in vivo* and *in vitro*

We demonstrated that our clonal mouse ESC lines can be used *in vivo* in chimaeric embryos. The ability
to conveniently switch between *in vitro* and *in vivo* experimentation was a key reason for us to
establish SynNotch technology in mouse ESCs. Both the receiver lines we generated and the parental
PSNB landing pad cell lines offer the power and flexibility to address questions we have so far been
unable to answer in *in vitro* and *in vivo* settings. For example, the system could be employed in cell
competition studies: STC receiver cells could be used to identify and isolate the direct neighbours of
eGFP-tagged "loser" cells, and profiled to study what changes are induced upon interaction with loser
cells in order to bring about their elimination. Receiver cells could also be engineered to express
candidate fitness-altering transgenes in response to interaction with eGFP-tagged wild-type sender
cells, as successfully demonstrated in *Drosophila* by (He et al., 2017).

The establishment of this system in mouse ESCs also allows to monitor and manipulate the effects of
cell-cell interactions in specific cell types obtained through directed differentiation. We also
demonstrated that our cell lines can be used to generate synthetic patterns of gene expression,
resulting in spatially-defined programming of cell fate. The combination of directed differentiation of

ESCs, spatial confinement of sender and receiver cells, and contact-mediated cell fate engineering
provides many possibilities for the study of cell-cell interactions in any developmental process of
interest.

Concluding remarks

Cell-cell interactions are a shared feature of the development of all multicellular organisms. Whilst the
particulars of these interactions vary greatly amongst eukaryotic supergroups, it is clear that they play
an essential role in development (Armingol et al., 2021). The Synthetic Biology field has recently
developed several applications to monitor cell communication, such as SynNotch (Morsut et al., 2016)
and derivative systems (Zhu et al., 2021), direct transfer of fluorophores to neighbouring cells
(Ombrato et al., 2019; Tang et al., 2020), reconstitution of a fluorophore following interaction
between different cell types carrying non-fluorescent fluorophore fragments (Kinoshita et al., 2020).
We have here demonstrated how SynNotch technology can be used to monitor and manipulate cell-
cell interactions in mouse ESCs and in mouse embryos. We hope that the SyNPL system will prove a
useful and adaptable tool for the mouse Stem Cell and Developmental Biology community.

MATERIALS AND METHODS

Reagents

REAGENT	SOURCE	IDENTIFIER	CONCENTRATION
Antibodies and nuclear counterstain			
Mouse monoclonal anti-Flag	Sigma-Aldrich	Cat# F3165; RRID: AB_259529	1:1000
Chicken polyclonal anti-GFP	Abcam	Cat# ab13970; RRID: AB_300798	1:1000
Goat polyclonal anti-Myc	Abcam	Cat# ab9132; RRID: AB_307033	1:1000
Mouse monoclonal anti-Myc AlexaFluor 488 conjugate	Santa Cruz	Cat# sc-40 AF488; RRID: AB_2892598	1:200
Rat monoclonal anti-mCherry	Invitrogen	Cat# M11217; RRID: AB_2536611	1:1000
Mouse monoclonal anti-Nup107 (NPC)	Abcam	Cat# ab24609; RRID: AB_448181	1:1000
Rabbit polyclonal anti-tRFP (tagBFP)	Evrogen	Cat# AB233; RRID: AB_2571743	1:1000
Mouse monoclonal anti-Tubb3	Biolegend	Cat# 801201; RRID: AB_2313773	1:1000
DRAQ7	Abcam	Cat# ab109202	1µM (IF); 300nM (flow cytometry)
Chemicals and solutions			

2-Mercaptoethanol	Gibco	Cat# 31350010	100nM
Accutase	Sigma-Aldrich	A6964	
Chorulon (hCG)	Intervet	Cat# CH-475-1	100 IU/ml
DMEM, high glucose, no glutamine, no phenol red	Gibco	Cat# 31053028	
Donkey serum	Sigma-Aldrich	Cat# D9663	3% v/v
Doxycycline hyclate	Sigma-Aldrich	Cat# D9891	1µg/ml
Dulbecco's phosphate buffered saline (PBS)	Sigma-Aldrich	Cat# D8537	
Fibronectin from bovine plasma solution	Sigma-Aldrich	Cat# F1141	
Foetal calf serum	Life Technologies	Cat# 10270106	10% v/v
Formaldehyde 37-41%	Fisher Scientific	Cat# F/1501/PB08	4% w/v
Gelatin	Sigma-Aldrich	Cat# G1890	0.1% w/v
Geneticin (G418)	Gibco	Cat# 11811031	200µg/ml
Glasgow Minimum Essential Medium	Sigma-Aldrich	Cat# G5154	
Hygromycin B	Gibco	Cat# 10687010	200µg/ml
L-Glutamine	Thermo Scientific	Cat# 25030024	2mM
Lipofectamine 3000 transfection reagent	Life Technologies	Cat# L3000008	
M2 medium	Sigma-Aldrich	Cat# M7167	
MEM Non-essential amino acids solution (100X)	Gibco	Cat# 11140050	1X
Penicillin/streptomycin	Gibco	Cat# 15140122	100 U/ml
PMSG	ProSpec	Cat# HOR-272	100 IU/ml
Prolong Gold Antifade Mountant	Thermo Fisher	Cat# P36930	
Puromycin dihydrochloride	Sigma-Aldrich	Cat# P8833	2µg/ml
Sodium pyruvate	Gibco	Cat# 11360039	1 mM
Triton X-100	Sigma-Aldrich	Cat# X100	0.1% v/v
Trypsin EDTA 0.25%	Gibco	Cat# 25200072	0.05% w/v
Tyrode's solution, acidic	Sigma-Aldrich	Cat# T1788	
Zeocin	Gibco	Cat# R25001	100µg/ml
Cell culture substrates and inserts			
3 well silicone culture insert	Ibidi	Cat# 80369	
µ-Slide 8 Well	Ibidi	Cat# 80826	
Glass coverslip, 24mm diameter	VWR	Cat# 631-1583	
Plastic flasks and plates for routine cell culture	Corning	Various	
DNA constructs			
pHR_EGFPligand	(Morsut et al., 2016)	Addgene 79129	
pHR_SFFV_LaG17_synNotch_TetRVP64	(Morsut et al., 2016)	Addgene 79128	
pDisplay-eGFP-TM	(Han et al., 2004)		

pHR_TRE-mCherry-PGK-tagBFP-WPRE	Elise Cachat, University of Edinburgh		
pRosa26-DEST-1lox	(Tosti et al., 2018)		
CAG-φC31 integrase	(Tosti et al., 2018)		
pENTR-2xAttP50	(Tosti et al., 2018)		
pENTR-2xAttB53	(Tosti et al., 2018)		
pPyCAG-eGFP-TM-IRES-Pac	This study		
pPyPGK-eGFP-TM-IRES-Pac	This study		
pPyCAG-HA-eGFP-Myc-TM-IRES-Pac	This study		
pPyPGK-HA-eGFP-Myc-TM-IRES-Pac	This study		
pPyPGK-Myc-LaG17-Notch1TM-tTA-IRES-Ble	This study		
pPyCAG-tagBFP-3xNLS-IRES-Hph	This study		
pENTR-attB53-Pac-bGHpA-attB53	This study		
pENTR-attB53-Pac-bGHpA-(TRE-mCherry-SV40pA)-attB53	This study		
pENTR-attB53-Pac-bGHpA-(tetO-mCherry-rBGpA)-attB53	This study		
pENTR-attB53_SNCB+_attB53	This study		
pENTR-attB53_SNCB-_attB53	This study		
pENTR-attB53-Pac-bGHpA-(TRE-3xFlag-Neurog1-SV40pA)-attB53	This study		
Cell lines			
E14Ju09 mouse ESCs (129/Ola, male)	(Hamilton and Brickman, 2014)		
CmGP1 sender mouse ESCs (129/Ola, male)	This study		
CHmGMP19 sender mouse ESCs (129/Ola, male)	This study		
CmGP1GH1 sender mouse ESCs (129/Ola, male)	This study		
E14GIP1 mouse ESCs (129/Ola, male)	This study		
EM35 mouse ESCs (129/Ola, male)	This study		
SNCB+ receiver mouse ESCs (129/Ola, male)	This study		
SNCB- receiver mouse ESCs (129/Ola, male)	This study		
35SRZ mouse ESCs (129/Ola, male)	This study		
PSNB mouse ESCs (129/Ola, male)			
PSNB-E mouse ESCs (129/Ola, male)	This study		
PSNB-tetO receiver mouse ESCs (129/Ola, male)	This study		
STC receiver mouse ESCs (129/Ola, male)	This study		

| STN receiver mouse ESCs (129/Ola, male) | This study | | |

Animal care and use

496 Animal experiments were performed under the UK Home Office project license PEEC9E359, approved
497 by the Animal Welfare and Ethical Review Panel of the University of Edinburgh and within the
498 conditions of the Animals (Scientific Procedures) Act 1986.

Chimaera generation

C57BL/6 female mice (Charles River) were superovulated (100 IU/ml PMSG and 100 IU/ml hCG
intraperitoneal injections 48 hours apart) and crossed with wild-type stud male mice. Pregnant mice
were culled at embryonic day 2.5 (E2.5) by cervical dislocation, ovaries with oviducts were dissected
and collected in pre-warmed M2 medium. Oviducts were flushed using PBS and a 20-gauge needle
attached to a 1ml syringe and filled with PB1 (Whittingham, 1974). E2.5 embryos were collected and
washed in PB1, their zona pellucida was removed using acidic Tyrode's solution, and they were
transferred to a plate with incisions where two clumps of approximately 8 sender and 8 receiver cells
were added to each embryo. Embryos were then incubated at 37°C in 5% CO_2 for 48 hours prior to
fixation.

Mouse ESC culture

Mouse embryonic stem cells were routinely maintained on gelatinised culture vessels (Corning) at
37°C and 5% CO_2 in Glasgow Minimum Essential Medium (GMEM) supplemented with 10% foetal calf
serum (FCS), 100U/ml LIF (produced in-house), 100nM 2-mercaptoethanol, 1X non-essential amino
acids, 2mM L-Glutamine), 1mM Sodium Pyruvate (medium referred to as "ES cell culture medium" or
"LIF+FCS"). The medium was supplemented with 200μg/ml G418, 2μg/ml puromycin, 200μg/ml
hygromycin B and/or 100μg/ml zeocin as appropriate.
For live imaging, GMEM was replaced with phenol red-free Dulbecco's Modified Eagle Medium
(DMEM), with all other components of the culture medium used at identical concentrations.

DNA constructs

pHR_SFFV_LaG17_synNotch_TetRVP64 (Addgene plasmid #79128) (Morsut et al., 2016) and
pHR_EGFPligand (Addgene plasmid #79129) (Morsut et al., 2016) were kind gifts from Dr Wendell Lim
and Dr Leonardo Morsut. pDisplay-GFP-TM (Han et al., 2004) was a kind gift of Dr Luis Ángel Fernández
(CNB-CSIC). pHR_TRE-mCherry-PGK-tagBFP-WPRE was a kind gift of Dr Elise Cachat (The University of
Edinburgh). pRosa26-DEST-1lox, CAG-φC31 integrase, pENTR-2xAttP50 and pENTR-2xAttB53 (Tosti et
al., 2018) constructs were kind gifts of Dr Keisuke Kaji (The University of Edinburgh).

Untagged transmembrane eGFP constructs were generated by digesting pHR_EGFPligand with
XhoI+NotI, and ligating the *IgGK signal-eGFP-PDGFRB TMD* cassette into XhoI+NotI-digested *pPyCAG-*
IRES-Pac (Malaguti et al., 2019) or *pPyPGK-IRES-Pac* (Rao et al., 2020) vector backbones. HA- and Myc-
tagged eGFP constructs were generated by PCR-amplifying an *IgGK signal-HA-eGFP-Myc-PDGFRB TMD*
cassette flanked by PspXI and NotI sites from pDisplay-GFP-TM, digesting the amplicon with
PspXI+NotI, and ligating the insert into XhoI+NotI-digested *pPyCAG-IRES-Pac* or *pPyPGK-IRES-Pac*
vector backbones.

The *pPyPGK-CD8a signal-Myc-LaG17-Notch1 minimal transmembrane core-tTA-IRES-Ble* SynNotch
receptor construct was generated by PCR-amplifying a *CD8a signal-Myc-LaG17-Notch1 minimal*

transmembrane   core-tTA   cassette   flanked   by   XhoI   and   Bsu36I   sites   from
pHR_SFFV_LaG17_synNotch_TetRVP64, digesting the amplicon with XhoI+Bsu36I, and ligating the
insert into a XhoI+Bsu36I-digested *pPyPGK-IRES-Ble* vector backbone. The mouse Notch1 minimal
transmembrane core consists of residues 1427-1752 (Uniprot: Q01705).

The *pPyCAG-tagBFP-3xNLS-IRES-Hph* construct was generated by PCR-amplifying a *tagBFP* cassette
flanked by XhoI and KasI sites from pHR_TRE-mCherry-PGK-tagBFP-WPRE, digesting the amplicon with
XhoI+KasI, and ligating the insert into a XhoI+NotI-digested *pPyCAG-IRES-Hph* backbone (Malaguti et
al., 2019) alongside oligonucleotides annealed to generate a *3xNLS* fragment with KasI and NotI
overhangs (Malaguti et al., 2013).

The *Rosa26* landing pad targeting vector was generated by Gateway Cloning (Invitrogen) of an *attL1-*
attP50-Neo-SV40pA-(CAG-mKate2-3xNLS-bGHpA)-attP50-attL2 cassette into the pRosa26-DEST-1lox
targeting vector. Its final structure is as follows: *Rosa26 5'HA-splice acceptor-loxP-attP50-Neo-*
SV40pA-(CAG-mKate2-3xNLS-bGHpA)-attP50-Rosa26 3'HA-PGK-DTA-bGHpA. Sequence in brackets is
on -strand.

A *pENTR-attB53-Pac-bGHpA-attB53* "empty vector" construct for RMCE at the *Rosa26* locus was
generated by adding a *Pac-bGHpA* cassette followed by an EcoRV restriction site to pENTR-2xAttB53
by Gibson assembly.
The TRE-mCherry RMCE construct used to generated STC receiver cells from PSNB landing pad lines
was generated by PCR-amplifying a *TRE-mCherry-SV40pA* cassette flanked by EcoRV-AscI and EcoRV-
BamHI sites from pHR_TRE-mCherry-PGK-tagBFP-WPRE, digesting the amplicon with EcoRV, ligating
the insert into EcoRV-digested *pENTR-attB53-Pac-bGHpA-attB53* backbone, and screening for
insertion on the -strand. The tetO-mCherry RMCE construct used to generate PSNB-tetO cells from
PNSB landing pad lines was generated by PCR-amplifying a *tetO-mCherry-rBGpA* cassette flanked by
MluI and BamHI sites, digesting the amplicon with MluI+BamHI and ligating the insert into
AscI+BamHI-digested *pENTR-attB53-Pac-bGHpA-(TRE-mCherry-SV40pA)*.
The TRE-3xFlag-Neurog1 RMCE construct used to generate STN receiver cells from PSNB landing pad
lines was generated by PCR-amplifying a *3xFlag-Neurog1* cassette flanked by NdeI and MfeI sites from
wild-type mouse cDNA, digesting the amplicon with NdeI+MfeI and ligating the insert into NdeI+MfeI-
digested *pENTR-attB53-Pac-bGHpA-(TRE-mCherry-SV40pA)* backbone (in which *mCherry* is flanked by
NfeI and MfeI sites).

SNCB+ and SNCB- constructs were generated in two steps. First, the base *pENTR-attB53-Pac-bGHpA-*
attB53 RMCE construct was linearised with EcoRV, and ligated with a HincII-*PGK-CD8a signal-Myc-*
LaG17-Notch1 minimal transmembrane core-tTA-HindIII fragment (digested from the SynNotch
receptor construct described above) and a HindIII-*bGHpA*-EcoRV fragment, and clones were screened
for insertion of the SynNotch receptor on the +strand. Next, the resulting construct was linearised
with EcoRV, and ligated with an EcoRV-*TRE-mCherry-SV40pA-PGK-tagBFP*-PacI fragment (digested
from pHR_TRE-mCherry-PGK-tagBFP-WPRE) and a PacI-*bGHpA*-EcoRV fragment. Correct assembly on
the + and - strands generated the SNCB+ and SNCB- constructs respectively.

Abbreviations: TMD: transmembrane domain; IRES: internal ribosome entry site; 5'/3' HA: 5'/3'
homology arm; SV40pA: SV40 polyadenylation signal sequence; bGHpA: bovine growth hormone
polyadenylation signal sequence; rBGpA: rabbit beta-globin polyadenylation signal sequence; NLS:
nuclear localisation signal; PGK: mouse *Pgk1* promoter; Pac: puromycin N-acetyltransferase; Hph:
hygromycin B phosphotransferase; Ble: bleomycin resistance gene (confers zeocin resistance); tTA:
tetracycline transactivator.

Transfections

For electroporations, 10^7 ESCs were electroporated with 100µg DNA using a BioRad GenePulser set to 800V/3µF. For nucleofections, 5×10^5 ESCs were nucleofected with 5µg DNA with the Lonza P3 Primary Cell Nucleofector Unit and kit, using program CG-104, and following manufacturer instructions. For lipofections, 10^5 ESCs were lipofected with 3µg DNA mixed with 3µl Lipofectamine 3000 and 6µl P3000 solution, following manufacturer instructions. For φC31-mediated RMCE, equal masses of RMCE constructs and CAG-φC31 vector were transfected.

Clonal ESC lines were generated by transfecting constructs of interest into ESCs, then plating cells at low density onto gelatinised 9cm dishes in the absence of selection. Selective medium was added 48 hours post-transfection and replaced every other day. After 7-10 days, clones were manually picked, dissociated, and replated into gelatinised 96-well culture plates. Clones were transferred to gelatinised vessels with larger culture areas when confluent, screened as appropriate, expanded and cryopreserved.

Cell lines

E14Ju09 ESCs are a 129/Ola male wild-type clonal line derived from E14tg2a (Hamilton and Brickman, 2014; Hooper et al., 1987).

Sender cells were generated by electroporating E14Ju09 ESCs with one of four constructs: *pPyCAG-IgGK signal-eGFP-PDGFRB TMD-IRES-Pac* (CmGP cells), *pPyPGK-IgGK signal-eGFP-PDGFRB TMD-IRES-Pac* (PmGP cells), *pPyCAG-IgGK signal-HA-eGFP-Myc-PDGFRB TMD-IRES-Pac* (CHmGMP cells), *pPyPGK-IgGK signal-HA-eGFP-Myc-PDGFRB TMD-IRES-Pac* (PHmGMP cells). Simplified versions of the constructs are displayed in Fig. 2. CmGP1GH1 sender cells were generated by lipofecting CmGP1 sender cells with a *pPyCAG-eGFP-IRES-Hph* construct. E14GIP1 "cytoplasmic sender" cells were generated by lipofecting E14Ju09 ESCs with a *pPyCAG-eGFP-IRES-Pac* construct.

EM35 landing pad cells were generated by electroporating E14Ju09 ESCs with the *Rosa26* landing pad targeting vector described above. Correct targeting was verified by genomic DNA PCR with the following primers: Forward: GGCGGACTGGCGGGACTA, Reverse: GGGACAGGATAAGTATGACATCATCAAGG. Primer locations and expected band sizes are displayed in Fig. S3A. This PCR strategy was modified from that described by (Mort et al., 2014) to suit the different sequence of our *Rosa26* targeting vector.

SNCB+ and SNCB- receiver cells were generated by electroporating EM35 ESCs with the constructs depicted in Figs. S4A, S6A, and a *CAG-φC31 integrase* construct to mediate RMCE.

35SRZ landing pad cells were generated by electroporating EM35 landing pad cells with *pPyPGK-CD8a signal-Myc-LaG17-Notch1 minimal transmembrane core-tTA-IRES-Ble*.

PSNB landing pad cell lines were generated by nucleofecting 35SRZ ESCs with a *pPyCAG-tagBFP-3xNLS-IRES-Hph* construct. PSNB-A cells were derived from 35SRZ clone 9 (PSNB-A clone 10 renamed PSNB clone A), PSNB-B cells were derived from 35SRZ clone 86 (PNSB-B clone 3 renamed PSNB clone B).

STC receiver cells were generated by nucleofecting PSNB ESCs with *CAG-φC31 integrase* and the following RMCE construct: *attB53-Pac-bGHpA-(TRE-mCherry-SV40pA)-attB53*. Sequence in brackets is on -strand. STC clone A1 was derived from PSNB clone A, STC clones B1 and B2 were derived from PSNB clone B.

PSNB-tetO cells were generated by nucleofecting PSNB ESCs with *CAG-φC31 integrase* and the following RMCE construct: *attB53-Pac-bGHpA-(tetO-mCherry-SV40pA)-attB53*. Sequence in brackets is on -strand.

PNSB-E cells were generated by nucleofecting PSNB ESCs with *CAG-φC31 integrase* and the following RMCE construct: *attB53-Pac-bGHpA-attB53*.

623 STN receiver cells were generated by nucleofecting PSNB clone A ESCs with *CAG-φC31 integrase* and
624 the following RMCE construct: *attB53-Pac-bGHpA-(TRE-3xFlag-Neurog1-SV40pA)-attB53*. Sequence in
625 brackets is on -strand.

Co-culture experiments

Sender and receiver cells were detached from culture vessels with accutase, quenched in ESC culture
medium, pelleted by spinning at 300g for 3 minutes, resuspended in ESC culture medium
supplemented with 2µg/ml puromycin and counted. Cells were plated at ratios described in figure
legends, and at empirically determined optimal densities.
For flow cytometry experiments, cells were plated onto 12-well plates coated with 7.5µg/ml
fibronectin, at the following densities: experiments carried out in the absence of doxycyline: 1h-8h:
$4x10^5$ cells/well; 16h: $2.4x10^5$ cells/well; 24h: $1.6x10^5$ cells/well; 48h: $8x10^4$ cells/well; 72h: $4x10^4$
cells/well. mCherry induction with 16h doxycycline (Fig. 4F-H): 0-4h: $2.4x10^5$ cells/well; 5-8h: $1.6x10^5$
cells/well; 24h: $8x10^4$ cells/well. mCherry downregulation experiments (Fig. S14): 0h-8h: $1.6x10^5$
cells/well; 16h: $1.2x10^5$ cells/well; 24h: $8x10^4$ cells/well; 48h: $4x10^4$ cells/well.
For immunofluorescence experiments, cells were plated on flamed 24mm glass coverslips housed in
a 6-well plate coated with 7.5µg/ml fibronectin.
For live imaging experiments, cells were plated onto an 8-well imaging slide coated with 7.5µg/ml
fibronectin, at the following densities: mCherry induction experiments: $3x10^4$ cells/well; 0-mCherry
downregulation experiments: 0-24h: $2x10^4$ cells/well; 24-48h: 10^4 cells/well. ESC culture medium was
supplemented with 200µg/ml hygromycin B and 1X penicillin/streptomycin.
To test induction of 3xFlag-Neurog1 in STN receiver cells, CmGP1 sender cells and STN receiver cells
were plated at a 9:1 ratio at a concentration of $2x10^5$ cells/well onto a 24mm flamed glass coverslip
housed in a 6-well plate coated with 7.5µg/ml fibronectin, then fixed and stained after 48 hours.

Synthetic patterning experiments

A 24mm glass coverslip housed in a 6-well plate was coated with 7.5µg/ml fibronectin, then allowed
to air dry. When fully dry, forceps were used to place a culture insert 3-well silicon chamber on top of
the coverslip; downward force was carefully exerted to secure it in place. $4x10^4$ cells were plated
overnight in 70µl culture medium in each of the three wells. Sender cells were plated in the central
well, and receiver cells were plated in the outside wells. 2ml culture medium were added outside of
the 3-well insert in order to prevent evaporation. The next day, the 2ml culture medium outside of
the 3-well insert were aspirated, and the 70µl in each of the three wells were carefully removed in
order not to dislodge the 3-well insert. Each well was quickly washed with 70µl PBS to get rid of any
remaining cells in suspension. Forceps were used to detach the 3-well insert from the glass coverslip,
and 2.5ml culture medium were added to the well. Culture medium was replaced daily. Growth of
cells into the gaps between wells were monitored daily; following contact between sender and
receiver cells, cells were kept in culture for a further 24 hours (STC receivers + CmGP1 senders) or a
further 96 hours (STN receivers + CmGP1 senders) prior to fixation and immunofluorescence.

Flow cytometry

Cells were washed in PBS, then detached from culture vessels with accutase. They were resuspended
in ice-cold PBS+10% FCS, pelleted by spinning at 300g for 3 minutes, resuspended in ice-cold PBS+10%
FCS+300nM DRAQ7 and placed on ice before analysing on a BD LSRFortessa flow cytometer. Forward
and side-scatter width and amplitude were used to identify single cells in suspension, dead cells were
excluded by gating on DRAQ7-negative cells, and tagBFP, GFP and mCherry/mKate2 expression were
then analysed using V 450/50-A, B 530/30-A, Y/G 610/20-A laser/filters combinations respectively.

Immunofluorescence

Cells were plated on flamed glass coverslips coated with 7.5μg/ml fibronectin and cultured as indicated in figure legends. Cells were washed with PBS, fixed in 4% formaldehyde in PBS for 20 minutes at room temperature, then washed three times in PBS for a total of 15 minutes. Cells were blocked overnight at 4°C in blocking solution (PBS+3% donkey serum+0.1% Triton X-100). Primary antibodies diluted in blocking solution were added for 3 hours at room temperature, the coverslips were washed 3 times in PBS for a total of 30 minutes, secondary antibodies diluted in blocking solution were added for 1 hour at room temperature, the coverslips were washed 3 times in PBS for a total of 30 minutes. The coverslips were then mounted onto glass slides in Prolong Gold mounting medium. For synthetic patterning experiments and blastocyst stainings, antibodies were incubated overnight at 4°C or 37°C respectively to improve penetration. Blastocysts were imaged in PBS in an imaging chamber, and scoring of mCherry-HI cells was performed manually using chimaeras containing both sender and receiver cells as a reference. All imaging was performed on a Leica SP8 confocal microscope with a 40X immersion lens unless otherwise indicated.

Live imaging

Cells to be imaged were allowed to adhere at room temperature for 15 minutes after plating, after which they were placed in a 37°C 5% CO_2 humidified chamber and imaged with a widefield Nikon Ti-E microscope, 20X lens, and Hamamatsu camera. Images were taken at 10 minute intervals for 24 hours, and xy coordinates were saved. Following live imaging, cells were fixed and stained for fluorophore expression, and imaged at the previously saved xy coordinates.

ACKNOWLEDGEMENTS

We thank Elise Cachat and Keisuke Kaji for advice and reagents; Luis Ángel Fernández for the pDisplay-eGFP-TM plasmid; Dónal O'Carroll and Pedro Moreira for assistance with generating chimaeric mice; Eve Moutaux for cloning initial versions of the RMCE constructs; Alexandre Veiga and Matthew French for performing preliminary experiments with the original SynNotch constructs. We thank Matthieu Vermeren for assistance with timelapse imaging, Fiona Rossi and Claire Cryer for flow cytometry support, and Theresa O'Connor, Helen Henderson and Marilyn Thomson for cell culture support. We are grateful to Val Wilson, Leonardo Morsut and members of the Lowell and Wilson labs for helpful discussions.

COMPETING INTERESTS

The authors declare no competing interests.

FUNDING

This work was funded by a Wellcome Trust Senior Fellowship to SL (WT103789AIA) and a Wellcome Trust Sir Henry Wellcome Fellowship to GB (WT100133).

REFERENCES

Armingol, E., Officer, A., Harismendy, O. and Lewis, N. E. (2021). Deciphering cell–cell interactions and communication from gene expression. *Nature Reviews Genetics* **22**, 71–88.

Arnold, S. J. and Robertson, E. J. (2009). Making a commitment: cell lineage allocation and axis patterning in the early mouse embryo. *Nature Reviews Molecular Cell Biology* **10**, 91–103.

Blin, G., Wisniewski, D., Picart, C., Thery, M., Puceat, M. and Lowell, S. (2018). Geometrical confinement controls the asymmetric patterning of Brachyury in cultures of pluripotent cells. *Development*.

Blin, G., Sadurska, D., Migueles, R. P., Chen, N., Watson, J. A. and Lowell, S. (2019). Nessys: A new set of tools for the automated detection of nuclei within intact tissues and dense 3D cultures. *PLOS Biol.* **17**, e3000388.

Boggs, S. S., Gregg, R. G., Borenstein, N. and Smithies, O. (1986). Efficient transformation and frequent single-site, single-copy insertion of DNA can be obtained in mouse erythroleukemia cells transformed by electroporation. *Experimental Hematology* **14**, 988–994.

Bradley, A., Evans, M., Kaufman, M. H. and Robertson, E. (1984). Formation of germ-line chimaeras from embryo-derived teratocarcinoma cell lines. *Nature* **309**, 255–256.

Cachat, E., Liu, W., Martin, K. C., Yuan, X., Yin, H., Hohenstein, P. and Davies, J. A. (2016). 2- and 3-dimensional synthetic large-scale de novo patterning by mammalian cells through phase separation. *Sci. Rep.* **6**, 20664.

Cai, L., Morrow, E. M. and Cepko, C. L. (2000). Misexpression of basic helix-loop-helix genes in the murine cerebral cortex affects cell fate choices and neuronal survival. *Development* **127**, 3021–3030.

Cau, E., Casarosa, S. and Guillemot, F. (2002). Mash1 and Ngn1 control distinct steps of determination and differentiation in the olfactory sensory neuron lineage. *Development* **129**, 1871–1880.

Charrier, S., Ferrand, M., Zerbato, M., Précigout, G., Viornery, A., Bucher-Laurent, S., Benkhelifa-Ziyyat, S., Merten, O. W., Perea, J. and Galy, A. (2011). Quantification of lentiviral vector copy numbers in individual hematopoietic colony-forming cells shows vector dose-dependent effects on the frequency and level of transduction. *Gene Therapy* **18**, 479–487.

Chen, C., Krohn, J., Bhattacharya, S. and Davies, B. (2011). A Comparison of Exogenous Promoter Activity at the ROSA26 Locus Using a PhiC31 Integrase Mediated Cassette Exchange Approach in Mouse ES Cells. *PLOS ONE* **6**, e23376.

Cho, J. H., Okuma, A., Al-Rubaye, D., Intisar, E., Junghans, R. P. and Wong, W. W. (2018). Engineering Axl specific CAR and SynNotch receptor for cancer therapy. *Scientific Reports* **8**, 3846.

Choe, J. H., Watchmaker, P. B., Simic, M. S., Gilbert, R. D., Li, A. W., Krasnow, N. A., Downey, K. M., Yu, W., Carrera, D. A., Celli, A., et al. (2021). SynNotch-CAR T cells overcome challenges of specificity, heterogeneity, and persistence in treating glioblastoma. *Sci Transl Med* **13**,.

Davies, J. (2017). Using synthetic biology to explore principles of development. *Development* **144**,
1146–1158.

De Los Angeles, A., Ferrari, F., Xi, R., Fujiwara, Y., Benvenisty, N., Deng, H., Hochedlinger, K.,
Jaenisch, R., Lee, S., Leitch, H. G., et al. (2015). Hallmarks of pluripotency. *Nature* **525**, 469–
478.

Dias, A. S., Almeida, I. de, Belmonte, J. M., Glazier, J. A. and Stern, C. D. (2014). Somites Without a
Clock. *Science* **343**, 791–795.

Ebrahimkhani, M. R. and Ebisuya, M. (2019). Synthetic developmental biology: build and control
multicellular systems. *Curr. Opin. Chem. Biol.* **52**, 9–15.

Evans, M. J. and Kaufman, M. H. (1981). Establishment in culture of pluripotential cells from mouse
embryos. *Nature* **292**, 154–156.

Fischer, S. C., Corujo-Simon, E., Lilao-Garzon, J., Stelzer, E. H. K. and Muñoz-Descalzo, S. (2020). The
transition from local to global patterns governs the differentiation of mouse blastocysts.
PLOS ONE **15**, e0233030.

Forsyth, J. E., Al-Anbaki, A. H., Fuente, R. de la, Modare, N., Perez-Cortes, D., Rivera, I., Kelly, R. S.,
Cotter, S. and Plusa, B. (2021). IVEN: A quantitative tool to describe 3D cell position and
neighbourhood reveals architectural changes in FGF4-treated preimplantation embryos.
PLOS Biology **19**, e3001345.

Fridy, P. C., Li, Y., Keegan, S., Thompson, M. K., Nudelman, I., Scheid, J. F., Oeffinger, M.,
Nussenzweig, M. C., Fenyö, D., Chait, B. T., et al. (2014). A robust pipeline for rapid
production of versatile nanobody repertoires. *Nature Methods* **11**, 1253–1260.

Friedrich, G. and Soriano, P. (1991). Promoter traps in embryonic stem cells: a genetic screen to
identify and mutate developmental genes in mice. *Gene Dev* **5**, 1513–1523.

Gorfinkiel, N. and Martinez Arias, A. (2021). The cell in the age of the genomic revolution: Cell
Regulatory Networks. *Cells & Development* 203720.

Gossen, M. and Bujard, H. (1992). Tight control of gene expression in mammalian cells by
tetracycline-responsive promoters. *Proceedings of the National Academy of Sciences* **89**,
5547–5551.

Gurdon, J. B. (1987). Embryonic induction — molecular prospects. *Development* **99**, 285–306.

Hamilton, W. B. and Brickman, J. M. (2014). Erk signaling suppresses embryonic stem cell self-
renewal to specify endoderm. *Cell Reports* **9**, 2056–2070.

Han, H.-J., Park, S.-G., Kim, S.-H., Hwang, S.-Y., Han, J., Traicoff, J., Kho, W.-G. and Chung, J.-Y.
(2004). Epidermal growth factor-like motifs 1 and 2 of Plasmodium vivax merozoite surface
protein 1 are critical domains in erythrocyte invasion. *Biochemical and Biophysical Research*
Communications **320**, 563–570.

He, L., Huang, J. and Perrimon, N. (2017). Development of an optimized synthetic Notch receptor as
an in vivo cell–cell contact sensor. *Proc. Natl. Acad. Sci.* **114**, 5467–5472.

Herbst, F., Ball, C. R., Tuorto, F., Nowrouzi, A., Wang, W., Zavidij, O., Dieter, S. M., Fessler, S.,
Hoeven, F. van der, Kloz, U., et al. (2012). Extensive Methylation of Promoter Sequences
Silences Lentiviral Transgene Expression During Stem Cell Differentiation In Vivo. *Molecular*
Therapy **20**, 1014–1021.

Ho, C. and Morsut, L. (2021). Novel synthetic biology approaches for developmental systems. *Stem*
Cell Reports **16**, 1051–1064.

Hong, S., Hwang, D.-Y., Yoon, S., Isacson, O., Ramezani, A., Hawley, R. G. and Kim, K.-S. (2007).
Functional Analysis of Various Promoters in Lentiviral Vectors at Different Stages of In Vitro
Differentiation of Mouse Embryonic Stem Cells. *Molecular Therapy* **15**, 1630–1639.

Hooper, M., Hardy, K., Handyside, A., Hunter, S. and Monk, M. (1987). HPRT-deficient (Lesch–
Nyhan) mouse embryos derived from germline colonization by cultured cells. *Nature* **326**,
292–295.

Huang, J., Zhou, W., Dong, W., Watson, A. M. and Hong, Y. (2009). Directed, efficient, and versatile
modifications of the Drosophila genome by genomic engineering. *Proceedings of the*
National Academy of Sciences **106**, 8284–8289.

Huang, H., Zhang, X., Lv, J., Yang, H., Wang, X., Ma, S., Shao, R., Peng, X., Lin, Y. and Rong, Z.
(2020). Cell-cell contact-induced gene editing/activation in mammalian cells using a
synNotch-CRISPR/Cas9 system. *Protein & Cell* **11**, 299–303.

Johnson, M. H. and Ziomek, C. A. (1983). Cell interactions influence the fate of mouse blastomeres
undergoing the transition from the 16- to the 32-cell stage. *Developmental Biology* **95**, 211–
218.

Kinoshita, M. and Smith, A. (2018). Pluripotency Deconstructed. *Development, Growth &*
Differentiation **60**, 44–52.

Kinoshita, N., Huang, A. J. Y., McHugh, T. J., Miyawaki, A. and Shimogori, T. (2020). Diffusible
GRAPHIC to visualize morphology of cells after specific cell–cell contact. *Scientific Reports* **10**,
14437.

Kondoh, G., Gao, X.-H., Nakano, Y., Koike, H., Yamada, S., Okabe, M. and Takeda, J. (1999). Tissue-
inherent fate of GPI revealed by GPI-anchored GFP transgenesis. *FEBS Letters* **458**, 299–303.

Luo, H., Wu, X., Sun, R., Su, J., Wang, Y., Dong, Y., Shi, B., Sun, Y., Jiang, H. and Li, Z. (2019). Target-
Dependent Expression of IL12 by synNotch Receptor-Engineered NK92 Cells Increases the
Antitumor Activities of CAR-T Cells. *Frontiers in Oncology* **9**,.

Ma, Q., Kintner, C. and Anderson, D. J. (1996). Identification of neurogenin, a Vertebrate Neuronal
Determination Gene. *Cell* **87**, 43–52.

Ma, Q., Chen, Z., Barrantes, I. del B., Pompa, J. L. de la and Anderson, D. J. (1998). neurogenin1 Is
Essential for the Determination of Neuronal Precursors for Proximal Cranial Sensory Ganglia.
Neuron **20**, 469–482.

Malaguti, M., Nistor, P. A., Blin, G., Pegg, A., Zhou, X. and Lowell, S. (2013). Bone morphogenic
protein signalling suppresses differentiation of pluripotent cells by maintaining expression of
E-Cadherin. *eLife* **2**, e01197.

Malaguti, M., Migueles, R. P., Blin, G., Lin, C.-Y. and Lowell, S. (2019). Id1 Stabilizes Epiblast Identity
by Sensing Delays in Nodal Activation and Adjusting the Timing of Differentiation. *Dev. Cell*
50, 462–477.e5.

Martin, G. R. (1981). Isolation of a pluripotent cell line from early mouse embryos cultured in
medium conditioned by teratocarcinoma stem cells. *Proceedings of the National Academy of*
Sciences **78**, 7634–7638.

Matsuda, M., Koga, M., Woltjen, K., Nishida, E. and Ebisuya, M. (2015). Synthetic lateral inhibition
governs cell-type bifurcation with robust ratios. *Nat Commun* **6**, 6195.

Matsunaga, S., Jeremiah, S. S., Miyakawa, K., Kurotaki, D., Shizukuishi, S., Watashi, K., Nishitsuji,
H., Kimura, H., Tamura, T., Yamamoto, N., et al. (2020). Engineering Cellular Biosensors
with Customizable Antiviral Responses Targeting Hepatitis B Virus. *iScience* **23**,.

McBurney, M. W., Sutherland, L. C., Adra, C. N., Leclair, B., Rudnicki, M. A. and Jardine, K. (1991).
The mouse Pgk-1 gene promoter contains an upstream activator sequence. *Nucleic Acids*
Research **19**, 5755–5761.

Morsut, L., Roybal, K. T., Xiong, X., Gordley, R. M., Coyle, S. M., Thomson, M. and Lim, W. A.
(2016). Engineering Customized Cell Sensing and Response Behaviors Using Synthetic Notch
Receptors. *Cell* **164**, 780–791.

Mort, R. L., Ford, M. J., Sakaue-Sawano, A., Lindstrom, N. O., Casadio, A., Douglas, A. T., Keighren,
M. A., Hohenstein, P., Miyawaki, A. and Jackson, I. J. (2014). Fucci2a: A bicistronic cell cycle
reporter that allows Cre mediated tissue specific expression in mice. *Cell Cycle* **13**, 2681–
2696.

Nishida-Aoki, N. and Gujral, T. S. (2019). Emerging approaches to study cell-cell interactions in
tumor microenvironment. *Oncotarget* **10**, 785–797.

Niwa, H., Yamamura, K. and Miyazaki, J. (1991). Efficient selection for high-expression transfectants
with a novel eukaryotic vector. *Gene* **108**, 193–199.

Nowotschin, S. and Hadjantonakis, A.-K. (2010). Cellular dynamics in the early mouse embryo: from
axis formation to gastrulation. *Current Opinion in Genetics & Development* **20**, 420–427.

Nowotschin, S., Eakin, G. S. and Hadjantonakis, A.-K. (2009). Dual transgene strategy for live
visualization of chromatin and plasma membrane dynamics in murine embryonic stem cells
and embryonic tissues. *genesis* **47**, 330–336.

Ombrato, L., Nolan, E., Kurelac, I., Mavousian, A., Bridgeman, V. L., Heinze, I., Chakravarty, P.,
Horswell, S., Gonzalez-Gualda, E., Matacchione, G., et al. (2019). Metastatic-niche labelling
reveals parenchymal cells with stem features. *Nature* **572**, 603–608.

Pera, M. F. and Tam, P. P. L. (2010). Extrinsic regulation of pluripotent stem cells. *Nature* **465**, 713–
720.

Perez, S. E., Rebelo, S. and Anderson, D. J. (1999). Early specification of sensory neuron fate
revealed by expression and function of neurogenins in the chick embryo. *Development* **126**,
1715–1728.

Pfaff, N., Lachmann, N., Ackermann, M., Kohlscheen, S., Brendel, C., Maetzig, T., Niemann, H.,
Antoniou, M. N., Grez, M., Schambach, A., et al. (2013). A ubiquitous chromatin opening
element prevents transgene silencing in pluripotent stem cells and their differentiated
progeny. *STEM CELLS* **31**, 488–499.

Pfeifer, A., Ikawa, M., Dayn, Y. and Verma, I. M. (2002). Transgenesis by lentiviral vectors: lack of
gene silencing in mammalian embryonic stem cells and preimplantation embryos.
Proceedings of the National Academy of Sciences of the United States of America **99**, 2140–
2145.

Posfai, E., Lanner, F., Mulas, C. and Leitch, H. G. (2021). All models are wrong, but some are useful:
Establishing standards for stem cell-based embryo models. *Stem Cell Reports* **16**, 1117–1141.

Rao, C., Malaguti, M., Mason, J. O. and Lowell, S. (2020). The transcription factor E2A drives neural
differentiation in pluripotent cells. *Development*.

Rhee, J. M., Pirity, M. K., Lackan, C. S., Long, J. Z., Kondoh, G., Takeda, J. and Hadjantonakis, A.-K.
(2006). In vivo imaging and differential localization of lipid-modified GFP-variant fusions in
embryonic stem cells and mice. *genesis* **44**, 202–218.

Rossant, J. and Tam, P. P. L. (2009). Blastocyst lineage formation, early embryonic asymmetries and
axis patterning in the mouse. *Development* **136**, 701–713.

Roybal, K. T., Williams, J. Z., Morsut, L., Rupp, L. J., Kolinko, I., Choe, J. H., Walker, W. J., McNally,
K. A. and Lim, W. A. (2016). Engineering T Cells with Customized Therapeutic Response
Programs Using Synthetic Notch Receptors. *Cell* **167**, 419–432.e16.

Sancho, M., Di-Gregorio, A., George, N., Pozzi, S., Sánchez, J. M., Pernaute, B. and Rodríguez, T. A.
(2013). Competitive Interactions Eliminate Unfit Embryonic Stem Cells at the Onset of
Differentiation. *Developmental Cell* **26**, 19–30.

Santorelli, M., Lam, C. and Morsut, L. (2019). Synthetic development: building mammalian
multicellular structures with artificial genetic programs. *Curr Opin Biotechnol* **59**, 130–140.

Schlissel, G. and Li, P. (2020). Synthetic Developmental Biology: Understanding Through
Reconstitution. *Annu. Rev. Cell Dev. Biol.* **36**, 339–357.

Schultz, R. M. (1985). Roles of Cell-to-Cell Communication in Development. *Biology of Reproduction*
32, 27–42.

Seibler, J., Küter-Luks, B., Kern, H., Streu, S., Plum, L., Mauer, J., Kühn, R., Brüning, J. C. and
Schwenk, F. (2005). Single copy shRNA configuration for ubiquitous gene knockdown in
mice. *Nucleic Acids Research* **33**, e67–e67.

Sekine, R., Shibata, T. and Ebisuya, M. (2018). Synthetic mammalian pattern formation driven by
differential diffusivity of Nodal and Lefty. *Nat Commun* **9**, 5456.

Sevcsik, E., Brameshuber, M., Fölser, M., Weghuber, J., Honigmann, A. and Schütz, G. J. (2015).
GPI-anchored proteins do not reside in ordered domains in the live cell plasma membrane.
Nature Communications **6**, 6969.

Sgodda, M., Alfken, S., Schambach, A., Eggenschwiler, R., Fidzinski, P., Hummel, M. and Cantz, T.
896   (2020). Synthetic Notch-Receptor-Mediated Transmission of a Transient Signal into
897   Permanent Information via CRISPR/Cas9-Based Genome Editing. *Cells* **9**, 1929.

898   **Shcherbo, D., Murphy, C. S., Ermakova, G. V., Solovieva, E. A., Chepurnykh, T. V., Shcheglov, A. S.,**
899   **Verkhusha, V. V., Pletnev, V. Z., Hazelwood, K. L., Roche, P. M., et al.** (2009). Far-red
900   fluorescent tags for protein imaging in living tissues. *Biochemical Journal* **418**, 567–574.

901   **Shioi, G., Kiyonari, H., Abe, T., Nakao, K., Fujimori, T., Jang, C.-W., Huang, C.-C., Akiyama, H.,**
902   **Behringer, R. R. and Aizawa, S.** (2011). A mouse reporter line to conditionally mark nuclei
903   and cell membranes for in vivo live-imaging. *Genesis* **49**, 570–578.

904   **Smithies, O., Gregg, R. G., Boggs, S. S., Koralewski, M. A. and Kucherlapati, R. S.** (1985). Insertion of
905   DNA sequences into the human chromosomal ß-globin locus by homologous recombination.
906   *Nature* **317**, 230–234.

907   **Srivastava, S., Salter, A. I., Liggitt, D., Yechan-Gunja, S., Sarvothama, M., Cooper, K., Smythe, K. S.,**
908   **Dudakov, J. A., Pierce, R. H., Rader, C., et al.** (2019). Logic-Gated ROR1 Chimeric Antigen
909   Receptor Expression Rescues T Cell-Mediated Toxicity to Normal Tissues and Enables
910   Selective Tumor Targeting. *Cancer Cell* **35**, 489–503.e8.

911   **Stern, C.** (1936). Somatic Crossing Over and Segregation in Drosophila Melanogaster. *Genetics* **21**,
912   625–730.

913   **Tam, P. P. L. and Loebel, D. A. F.** (2007). Gene function in mouse embryogenesis: get set for
914   gastrulation. *Nature Reviews Genetics* **8**, 368–381.

915   **Tang, R., Murray, C. W., Linde, I. L., Kramer, N. J., Lyu, Z., Tsai, M. K., Chen, L. C., Cai, H., Gitler, A.**
916   **D., Engleman, E., et al.** (2020). A versatile system to record cell-cell interactions. *eLife* **9**,
917   e61080.

918   **Tchorz, J. S., Suply, T., Ksiazek, I., Giachino, C., Cloëtta, D., Danzer, C.-P., Doll, T., Isken, A.,**
919   **Lemaistre, M., Taylor, V., et al.** (2012). A Modified RMCE-Compatible Rosa26 Locus for the
920   Expression of Transgenes from Exogenous Promoters. *PLOS ONE* **7**, e30011.

921   **Toda, S., Blauch, L. R., Tang, S. K. Y., Morsut, L. and Lim, W. A.** (2018). Programming self-organizing
922   multicellular structures with synthetic cell-cell signaling. *Science* **361**, 156–162.

923   **Toda, S., McKeithan, W. L., Hakkinen, T. J., Lopez, P., Klein, O. D. and Lim, W. A.** (2020). Engineering
924   synthetic morphogen systems that can program multicellular patterning. *Science* **370**, 327–
925   331.

926   **Tosti, L., Ashmore, J., Tan, B. S. N., Carbone, B., Mistri, T. K., Wilson, V., Tomlinson, S. R. and Kaji,**
927   **K.** (2018). Mapping transcription factor occupancy using minimal numbers of cells in vitro
928   and in vivo. *Genome Research* **28**, 592–605.

929   **Toth, T., Balassa, T., Bara, N., Kovacs, F., Kriston, A., Molnar, C., Haracska, L., Sukosd, F. and**
930   **Horvath, P.** (2018). Environmental properties of cells improve machine learning-based
931   phenotype recognition accuracy. *Scientific Reports* **8**, 10085.

932   **Velkey, J. M. and O'Shea, K. S.** (2013). Expression of Neurogenin 1 in Mouse Embryonic Stem Cells
933   Directs the Differentiation of Neuronal Precursors and Identifies Unique Patterns of Down-
934   stream Gene Expression. *Developmental Dynamics* **242**, 230–253.

Wang, L., Long, J., Chen, H., Sun, S., Lv, K., Li, Q. and Wang, X. (2021). Manipulation of focal Wnt
activity via synthetic cells in a double-humanized zebrafish model of tumorigenesis.
International Journal of Cancer **148**, 2815–2824.

Whittingham, D. G. (1974). Embryo banks in the future of developmental genetics. *Genetics* **78**,
395–402.

Wu, G., Liu, N., Rittelmeyer, I., Sharma, A. D., Sgodda, M., Zaehres, H., Bleidißel, M., Greber, B.,
Gentile, L., Han, D. W., et al. (2011). Generation of Healthy Mice from Gene-Corrected
Disease-Specific Induced Pluripotent Stem Cells. *PLOS Biology* **9**, e1001099.

Yang, Z., Yu, Z., Cai, Y., Du, R. and Cai, L. (2020). Engineering of an enhanced synthetic Notch
receptor by reducing ligand-independent activation. *Communications Biology* **3**, 1–7.

Yang, B. A., Westerhof, T. M., Sabin, K., Merajver, S. D. and Aguilar, C. A. (2021). Engineered Tools
to Study Intercellular Communication. *Advanced Science* **8**, 2002825.

Yuan, L. and Hassan, B. A. (2014). Neurogenins in brain development and disease: An overview.
Archives of Biochemistry and Biophysics **558**, 10–13.

Zhu, I., Liu, R., Hyrenius-Wittsten, A., Piraner, D. I., Alavi, J., Israni, D. V., Khalil, A. S. and Roybal, K.
T. (2021). Design and modular assembly of synthetic intramembrane proteolysis receptors
for custom gene regulation in therapeutic cells. *bioRxiv* 2021.05.21.445218.

FIGURE LEGENDS

Figure 1. Diagram of SynNotch cell-cell interaction reporter ES cells.
(A) Sender cells express membrane-tethered extracellular eGFP from a ubiquitous promoter. (B)
Receiver cells contain a *TRE-mCherry* transgene, express *tagBFP-3xNLS* from a ubiquitous promoter,
and express a SynNotch receptor from a ubiquitous promoter. The SynNotch receptor is comprised
of an extracellular LaG17 anti-GFP nanobody, the core transmembrane region of Notch1 (containing
proteolytic cleavage sites), and an intracellular tTA. (C) Upon interaction of the eGFP on a sender cell
with the SynNotch receptor, the Notch1 core domain is cleaved, releasing the tTA, which can
translocate into the nucleus, bind the TRE element, and drive *mCherry* transcription.

Figure 2. Screening of clonal sender ESC lines.
(A-D) Diagram of the constructs used to generate eGFP sender cell lines. (E) Median eGFP intensity
and (G) percentage eGFP-positive cells in untagged eGFP sender cells. Parental wild-type cells are
included as a negative control. 5000 cells were analysed for each clone. (F) Median eGFP intensity
and (H) percentage eGFP-positive cells in HA- and Myc-tagged eGFP sender cells. Parental wild-type
cells are included as a negative control. 15000 cells were analysed for each clone. Analyses were
performed separately from those in (E), median intensities are not directly comparable. (I)
Comparison of eGFP distributions in parental wild-type cells, CmGP1 and CHmGMP19 sender cells.
85000 cells were analysed for each sample. (J-K) Immunofluorescence of (J) CmGP1 and (K)
CHmGMP19 sender cells. Scale bar: 30μm. NPC: nuclear pore complex. A.F.U.: arbitrary fluorescence
units.

Figure 3. Generation of clonal cell-cell interaction reporter STC receiver cells.
(A) Summary of transgenes stably integrated into the genome of STC clonal ESC lines. (B-C) Flow
cytometry analysis of mCherry and tagBFP-3xNLS expression in (B) wild-type and (C) control PSNBB-E
cells containing all SyNPL receiver constructs except for the TRE-mCherry cassette. These control cell

lines were used to set gates for tagBFP- and mCherry-positivity. (D) Immunofluorescence of STC
clones A1, B1 and B2 co-cultured with CmGP1 sender cells for 24 hours (1:3 sender:receiver cell
ratio). Scale bar: 30μm. (E-H) Flow cytometry analysis of mCherry and tagBFP-3xNLS expression in
STC receiver cells cultured alone or in the presence of indicated eGFP-expressing cell lines (9:1
sender:receiver cell ratio). The mCherry-HI gate displayed in figure was set based on mCherry
distribution in STC receiver cells cultured alone.
For panels (B,C,E-H) percentages of cells in each gate are indicated. 11000 cells were analysed for
each sample. Diagrams depicting the transgenes present in each cell type and expected mCherry and
tagBFP-3xNLS expression patterns are displayed next to flow cytometry plots.

Figure 4. Kinetics of mCherry induction in STC receiver cells.
(A) Stills from Movie 1 displaying mCherry and eGFP expression in STC clone A1 receiver cells co-
cultured with CmGP1GH1 sender cells (1:1 sender:receiver cell ratio). Immuofluorescence of cells
after 24 hours of filming is displayed as a 24 hour timepoint, and includes tagBFP-3xNLS signal in
place of a brightfield image. Scale bar: 30μm. Yellow arrowheads: initial STC receiver cell contact
with sender cells, onset of mCherry expression, cell descendants at 24 hours. Magenta arrowheads:
STC receiver cell not making contact with sender cells, cell descendants at 24 hours. Cyan
arrowhead: initial STC receiver cell contact with sender cell, cell descendant at 24 hours.
(B) Experimental setup to analyse kinetics of mCherry upregulation in STC receiver cells by flow
cytometry.
(C) Percentage of mCherry-HI STC receiver cells following co-culture with CmGP1 sender cells for the
indicated amount of time (9:1 sender:receiver cell ratio). Data presented as mean ± standard
deviation of three independent experiments. A minimum of 8000 cells were analysed for each
sample. The mCherry-HI gate was set based on mCherry distribution in STC receiver cells cultured
alone.
(D) Distribution of mCherry fluorescence in STC clone A1 receiver cells following co-culture with
CmGP1 sender cells for the indicated amount of time (9:1 sender:receiver cell ratio). Data from a
single experiment, representative of three biological replicates. STC clone A1 cells cultured alone
("0h") are displayed as a shaded black histogram in all panels. 10000 cells were analysed for each
sample.
(E) Stills from Movie 2 displaying mCherry and eGFP expression in STC clone A1 receiver cells co-
cultured with CmGP1GH1 sender cells (1:1 sender:receiver cell ratio). Immuofluorescence of cells
after 24 hours of filming is displayed as a 24 hour timepoint, and includes tagBFP-3xNLS signal in
place of a brightfield image. Scale bar: 30μm. White arrowheads label an STC receiver cell in contact
with sender cells for 7 hours, which then loses contact with sender cells between the 8 and 20 hour
timepoints whilst its levels of mCherry keep increasing.(F) Experimental setup to analyse kinetics of
mCherry upregulation in STC receiver cells by flow cytometry, allowing time for protein maturation.
Following sender:receiver cell co-culture for 0-24 hours, 1μg/ml doxycycline (dox) was added to the
culture medium for a further 16 hours in order to inhibit tTA-mediated *mCherry* transcription, and
allow translation and folding of previously transcribed *mCherry*.
(G) Percentage of mCherry-HI STC receiver cells after co-culture with CmGP1 sender cells for the
indicated amount of time and following a further 16 hours of doxycycline treatment (9:1
sender:receiver cell ratio). Data presented as mean ± standard deviation of three independent
experiments. A minimum of 8000 cells were analysed for each sample. The mCherry-HI gate was set
based on mCherry distribution in STC receiver cells cultured alone in doxycycline for 16 hours.
(H) Distribution of mCherry fluorescence in STC clone A1 receiver cells following co-culture with
CmGP1 sender cells for the indicated amount of time and 16 hours doxycycline treatment (9:1
sender:receiver cell ratio). Data from a single experiment, representative of three biological
replicates. STC clone A1 cells plated with CmGP1 sender cells in doxycycline-containing medium for
16 hours ("0h") are displayed as a shaded black histogram in all panels. 10000 cells were analysed
for each sample.
1027

Figure 5. Contact-mediated induction of mCherry in chimaeric blastocysts.
(A) Chimaeric blastocysts containing STC clone B2 receiver cells and/or CmGP1GH1 sender cells. The images of the three embryos were taken from the same z-plane of a confocal stack and come from a single field of view. Nuclei were counterstained with DRAQ7. Scale bars: 30µm. (B) Quantification of embryos containing cells expressing readily detectable levels of mCherry ("mCherry-HI") across all experiments. Embryos containing both sender and receiver cells were used as a reference for scoring sender-only chimaeras, receiver-only chimaeras, and wild-type embryos.

Figure 6. Contact-mediated synthetic patterning of gene expression and fate programming.
(A) Diagram illustrating synthetic patterning strategy: sender and receiver cells are grown to confluence in separate chambers of a multi-chamber culture insert. The insert is removed and cells are allowed to proliferate until they come into contact, which induces transgene expression in receiver cells in a stripe pattern. (B) Synthetic striped pattern of mCherry induction in STC clone B1 receiver ESCs in contact with CmGP1 sender ESCs. eGFP, tagBFP, mCherry immunofluorescence. (C) Summary of transgenes stably integrated into the genome of *Neurog1*-inducible STN clonal ESC lines. (D) Synthetic striped pattern of neuronal differentiation of STN receiver ES cells in contact with CmGP1 sender ESCs. eGFP, tagBFP, Tubb3 immunofluorescence. Scale bars: 30µm.

FIGURE 1

A **Sender cells**

B **Receiver cells**

C **Sender cells in direct contact with receiver cells**

eGFP ligand

● Membrane-tethered
  Extracellular eGFP

SynNotch receptor

⊻ GFP Nanobody (LaG17)

Minimal Notch core (mNC)

tTA

FIGURE 2

A CAG | Signal | eGFP | TM | IRES | Pac | pA

B CAG | Signal | HA | eGFP | Myc | TM | IRES | Pac | pA

C PGK | Signal | eGFP | TM | IRES | Pac | pA

D PGK | Signal | HA | eGFP | Myc | TM | IRES | Pac | pA

E Median eGFP intensity

F Median eGFP intensity

G Percentage eGFP+

H Percentage eGFP+

I Flow analysis
— Parental
— CmGP1
— CHmGMP19

J CmGP1 (CAG-membrane-tethered-eGFP-IRES-Pac clone 1)
eGFP | NPC | eGFP+NPC

K CHmGMP19 (CAG-HA-membrane-tethered-eGFP-Myc-IRES-Pac clone 19)
eGFP | Myc | NPC | eGFP+Myc | eGFP+NPC

FIGURE 3

A

STC receiver cells (SynNotch-TRE-mCherry)

B Wild-type

C EV control "receivers"

Flow analysis of GFP- cells

D STC receivers + CmGP1 senders

eGFP | tagBFP 3XNLS | mCherry | Merge

STC cl. A1

STC cl. B1

STC cl. B2

E STC receivers

F STC receivers + CmGP1 senders

G STC receivers + E14GIP1 "senders"

H STC receivers + CmGP1GH1 senders

Flow analysis of tagBFP+ GFP- cells

STC cl. A1 | STC cl. B1 | STC cl. B2

FIGURE 4

FIGURE 5

A

Chimaeric blastocysts with STC receiver + CmGP1GH1 sender cells

| Senders only | Receivers only | Senders + receivers |

Nuclei eGFP tagBFP-3xNLS mCherry

B

	mCherry-HI embryos / total embryos			
	All clones	STC cl. A1	STC cl. B1	STC cl. B2
Wild-type	**0/6**	-	-	-
Senders only	**0/5**	-	-	-
Receivers only	**1/13**	1/7	0/3	0/3
Receivers+senders	**55/55**	21/21	23/23	11/11

FIGURE 6

A

Plate sender and receiver cells in separate wells of multiwell insert

Remove multiwell insert Give cells time to proliferate

Transgene expression induced at linear interface between sender and receiver cells

B mCherry induction in STC receivers at sender-receiver border

eGFP (sender cells) | tagBFP-3XNLS (receiver cells) | mCherry | Merge

CmGP1 senders

mCherry-inducible STC receivers

C

STN receiver cells (Neurog1-inducible)

D Forced neuronal differentiation of STN receivers at sender-receiver border

eGFP (sender cells) | tagBFP-3XNLS (receiver cells) | Tubb3 | Merge

CmGP1 senders

Neurog1-inducible STN receivers