LEAFY AND WEEDY SEADRAGON GENOMES CONNECT GENIC AND REPETITIVE DNA FEATURES TO THE EXTRAVAGANT BIOLOGY OF SYNGNATHID FISHES

Running Title: The genomes of two seadragon species

AUTHORS

Clayton M. Small[1,2=#], Hope M. Healey[1], Mark C. Currey[1], Emily A. Beck[1,2], Julian Catchen[3], Angela S. P. Lin[4], William A. Cresko[1,2#], and Susan Bassham[1=#]

[1] Institute of Ecology and Evolution, University of Oregon, Eugene, OR 97403
[2] Presidential Initiative in Data Science, University of Oregon, Eugene, OR 97403
[3] Department of Evolution, Ecology and Behavior, University of Illinois at Urbana-Champaign, Urbana, IL 61801
[4] Knight Campus for Accelerating Scientific Impact, University of Oregon, Eugene, OR 97403
[=] Contributed equally

ORCID IDs

C.M.S.	0000-0003-1615-7590
H.M.H.	0000-0001-9978-1553
M.C.C.	0000-0003-3688-7208
E.A.B.	0000-0001-9810-5094
J.C.	0000-0002-4798-660X
A.S.P.L.	0000-0001-5349-927X
W.A.C.	0000-0002-3496-8074
S.B.	0000-0002-7309-2095

[#]**Corresponding Authors**
Susie Bassham, William A. Cresko and Clayton M. Small

Author Contributions: S.B. and C.M.S. conceived of the project, and along with W.A.C. planned the work and provided general oversight of the project. S.B. and C.M.S. generated the seadragon sequencing data, and C.M.S. performed the genome assemblies. M.C.C. led the genome annotation and assisted C.M.S. with annotation and analysis of TEs. J.C. and S.B. performed the conserved synteny analysis. C.M.S. performed the global gene family evolution analyses. In-depth analyses of FGF/FGFR gene family evolution were performed by S.B., H.M.H. E.A.B. S.B. and A.S.P.L. prepared the weedy seadragon specimen for scanning, A.S.P.L performed Zeiss Xradia scans and initial image processing, and H.M.H. led image processing for figure composites. H.M.H. also performed the analysis of microRNAs, with initial data generation and processing performed by S.B and C.M.S. M.C.C. produced all fish illustrations for the figures. S.B. and C.M.S. took the lead in writing the manuscript, and all authors made writing contributions. The work was performed in the laboratory of W.A.C. at the University of Oregon with funding to W.A.C.

Keywords: genome sequencing, novel traits, evolution, miRNA, transposable elements, molecular evolution, fibroblast growth factors, syngnathids

Abstract

Seadragons are a remarkable lineage of teleost fishes, and they are members of the family Syngnathidae renowned for having evolved male pregnancy. Comprising three known species, seadragons are widely recognized and admired for their fantastical body forms and coloration, and their specific habitat requirements have made them flagship representatives for marine conservation and natural history interests. Until recently, a gap has been the lack of significant genomic resources for seadragons. We have produced gene-annotated, chromosome-scale genome models for the leafy and weedy seadragon to advance investigations into evolutionary innovation and elaboration of morphological traits in seadragons as well as their pipefish and seahorse relatives. We identified several interesting features specific to seadragon genomes, including divergent non-coding regions near a developmental gene important for integumentary outgrowth, a high genome-wide density of repetitive DNA, and recent expansions of transposable elements and a vesicular trafficking gene family. Surprisingly, comparative analyses leveraging the seadragon genomes and additional syngnathid and outgroup genomes revealed striking, syngnathid-specific losses in the family of fibroblast growth factors (FGFs), which likely involve re-organization of highly conserved gene regulatory networks in ways that have not previously been documented in natural populations. The resources presented here serve as important tools for future evolutionary studies of developmental processes in syngnathids and will be a key resource for conservation studies of the extravagant seadragons and their relatives.

INTRODUCTION

Seadragons are phenotypic outliers in an already exceptional clade of teleost

fishes (Family Syngnathidae) that also includes seahorses and pipefishes. For this

reason, seadragons are often a colorful, flagship group in discussions of adaptation and

evolutionary innovation. Seadragons demonstrate strikingly derived characters

compared to their pipefish and seahorse relatives, including "leafy appendages,"

extreme curvature of the spine (kyphosis and lordosis), especially elongated craniofacial

bones, and large body size (Figure 1)(Dawson, 1985; Stiller et al., 2015). Substantial

differences in these traits exist even among the three known extant species:

Phycodurus eques (leafy seadragon), *Phyllopteryx taeniolatus* (weedy, or common,

seadragon), and the recently described *Phyllopteryx dewysea* (ruby seadragon) (Stiller

et al., 2015).

In addition to being a focus for evolutionary studies, seadragons are of

significant cultural and conservation interest (Connolly et al., 2002; Martin-Smith &

Vincent, 2006; Stiller et al., 2017). Presumed adaptations for crypsis including the leafy

appendages, unique body plan, and elaborate skin coloration contribute to the status of

seadragons as distinguished and valued cultural symbols for the people of Australia,

where seadragon species are endemic. Because seadragon distributions are specific to

temperate Australian macroalgal reefs, and their population sizes are relatively small,

seadragons are likely susceptible to negative human impacts including global climate

change. Furthermore, recent population genomic studies documenting significant

population structure (Klanten et al., 2020; Stiller et al., 2021) in these species are

especially relevant to conservation decisions. The unique evolutionary innovations,

cultural importance, and conservation challenges all elevate the need to better

understand and conserve seadragon species.

To improve our understanding of highly derived phenotypic traits and genomic

features within seadragons, as well as those that are shared but derived among the

Syngnathidae, we sequenced and annotated chromosome-scale assemblies for a male

leafy seadragon and a female weedy seadragon. In addition to the production of these

genomic resources we carried out several comparative analyses among five syngnathid

and many other teleost genomes to determine changes in genome organization and

content, including a more detailed analysis of a few key gene families and regulatory

elements that may be involved in the development of syngnathid phenotypic

innovations. Lastly, we performed high-resolution 3D X-ray microscope scans of an

adult, male weedy seadragon to more precisely view seadragon morphological

innovations.

Our work reveals a number of seadragon-specific genomic features, including

divergent conserved non-coding elements (CNEs) near key developmental genes, a

unique microRNA gene repertoire, and expanded gene families related to immunity and

vesicular trafficking. We also found that the seadragon genomes are highly repetitive for

their sizes, with unique repeat abundance distributions. Because the seadragon lineage

occupies a region of the syngnathid phylogeny that is relatively basal to most of the

species diversity, we leveraged their phylogenetic position to identify several genomic

synampomorphies of the family. These genomic features include the striking loss of

several highly conserved fibroblast growth factor (FGF) genes, expansions and

contractions of gene families related to immunity and potentially male pregnancy, and

syngnathid-specific transposable element (TE) expansion.

With these new genome models and rich accompanying data, we grow the

existing collection of high-quality genomic tools and insights for several syngnathid

groups including genera *Syngnathus* (Roth et al., 2020; Small et al., 2016),

Hippocampus (Li et al., 2021; Lin et al., 2016), *Microphis* (Zhang et al., 2020), and most

recently (published as of the writing of this paper) *Phyllopteryx* and *Syngnathoides* (Qu

et al., 2021). We add to this quiver of syngnathid genomes useful in illuminating the

evolution and development of puzzling syngnathid novelties such as male pregnancy

and leaf-like appendages. These genomic resources will also support ongoing efforts to

understand and conserve sensitive syngnathid populations, including phylogenetic

umbrella species like the seadragons.

91   **MATERIALS AND METHODS**

Seadragon genome assemblies

We isolated high molecular weight genomic DNA from tissues of an adult male

leafy seadragon (*Phycodurus eques*) and from an adult female common ("weedy")

seadragon (*Phyllopteryx taeniolatus*). We then generated PacBio libraries for both

species and sequenced 49.12 and 80.80 Gb, respectively. We also generated

"shotgun" whole-genome sequencing (WGS) Illumina libraries for both species,

sequencing 57.48 and 105.79 Gb, to estimate genome size and polish the PacBio

assemblies (see Supplemental Methods for all software versions and non-default

parameters). We assembled both genomes with Flye (Kolmogorov et al., 2019), using all

PacBio data excluding "scraps" with default parameters and an estimated genome size

of 600 Mb. We performed two rounds of polishing on the primary Flye assemblies with

the tool arrow (Chin et al., 2013), using the PacBio reads and performed an additional

two rounds of polishing for each genome with WGS Illumina data, using pilon (Walker et

al., 2014). To organize Flye assemblies into putative chromosome-scale genome

models we generated Hi-C libraries using Phase Genomics Proximo Animal kits, then

scaffolded using the 3D-DNA pipeline (Dudchenko et al., 2017) with breaking of original

scaffolds disabled, followed by visualization and manual editing using Juicer and

Juicebox (Dudchenko et al., 2018; Durand et al., 2016). We evaluated assembly quality

and completeness using Quast (Gurevich et al., 2013) and BUSCO (Waterhouse et al.,

2018).

Draft short-read genome assemblies for *Doryrhamphus excisus* and *Synchiropus*

splendidus

To supplement our comparative analyses with additional syngnathid genomes

and a close outgroup we isolated high molecular weight DNA and generated linked-

reads assemblies for the bluestripe pipefish (*Doryrhamphus excisus*) and the Mandarin

dragonet (*Synchiropus splendidus*). These assemblies were performed using 10x

Genomics Chromium technology and the Supernova assembly software (Weisenfeld et

al., 2017). Details for these assembly methods are as described in Stervander and

Cresko (2021).

mRNA-seq

To generate mRNA-seq libraries, we extracted total RNA from tissues of the

same *P. eques* and *P. taeniolatus* individuals as were used for PacBio genome

sequencing. From the *P. eques* specimen, we dissected testis, leafy appendage, eye,

and gill tissues. From *P. taenoiolatus* we dissected ovary, leafy appendage, eye, and

liver tissue (see Supplemental Methods). We used the Roche KAPA HyperPrep Kit to

generate indexed, stranded mRNA-seq libraries for Illumina sequencing to obtain

301.52 million paired-end 100 bp reads. We trimmed Illumina adaptors and low-quality

regions from reads using process_shortreads from the Stacks software suite (Catchen

et al., 2013; Catchen et al., 2011) and aligned cleaned RNA-seq reads from both

seadragon species to both *P. taeniolatus* and *P. eques* genome assemblies using STAR

aligner (Dobin et al., 2013).

miRNA-seq

To generate small RNA-seq reads we purified a small fraction of RNA from each

tissue above using Zymo DirectZol columns and generated indexed sequencing

libraries using the NextFlex Small RNA-seq Kit v3. We ran BBMap (Bushnell, 2014) on

the weedy and leafy seadragon genomes to align sequencing reads. We annotated

miRNAs in the leafy seadragon based on sequence conservation with included species'

miRNAs (Antarctic blackfin icefish (*Chaenocephalus aceratus*), platyfish (*Xiphophorus*

maculatus), European perch (*Perca fluviatilis*), and threespine stickleback (*Gasterosteus*

aculeatus), zebrafish (*Danio rerio*)*, spotted gar (*Lepisosteus oculatus*), and medaka

(*Oryzias latipes*)) as described in the Prost! (Desvignes et al., 2019) manual (see

Supplemental Methods). Additionally, to run Prost!, we supplied threespine stickleback

noncoding sequences (Desvignes et al., 2019). We ran Prost! for the weedy seadragon

by adding the leafy seadragon's annotations to the included species pool. We

distinguished ohnologous miRNAs for the seadragon species using synolog (Catchen et

al., 2009) synteny software, which aligned the leafy seadragon genome with the

stickleback genome. Any miRNA not initially identified in the sequencing reads were

searched for in the leafy seadragon genome with BLASTN and vista plots.

Genome annotation

To facilitate the analysis of comparable annotations among genome assemblies,

we performed new annotations for leafy seadragon (*P. eques*), weedy seadragon (*P.*

taeniolatus), Greater pipefish (*Syngnathus acus*), bluestripe pipefish (*Doryrhamphus*

excisus), Mandarin dragonet (*Synchiropus splendidus*), and Pacific bluefin tuna (Smit et

al.) (*Thunnus orientalis*) (see Supplemental Methods for all public sequence data

accessions). Assemblies were soft-masked for repetitive elements and areas of low

complexity with RepeatMasker (Smit et al.) using custom repeat libraries made by

combining a teleost-specific library extracted from RepeatModeler2 (Flynn et al., 2020)

with species-specific repeat libraries produced by running RepeatModeler on each

species listed above. We aligned all RNA-seq data (including new and previously

published reads) as described above and supplied the .bam files to BRAKER2 (Bruna et

al., 2021) for genome annotation.

We evaluated the final BRAKER2 set of annotated genes for each genome using

InterProScan (Jones et al., 2014) and retained a final, "filtered" version of the annotation

if they showed InterProScan evidence other than disorder-based (i.e. MobiDB-lite). We

subjected remaining amino acid sequences to blastp searches against the NCBI

RefSeq protein database and retained those in the filtered annotation if they returned a

hit (e-value threshold of 0.001). For all downstream analyses of protein-coding

sequences based on these and other annotations, we selected the longest transcript

per locus.

Repeat annotation

We characterized the repetitive content of 16 teleost genome assemblies (Figure

S1; see Supp. Methods for accession numbers). With one exception (tiger tail seahorse

(*Hippocampus comes*)), we focused exclusively on assemblies produced by long-read

(i.e., PacBio or Oxford Nanopore) and/or linked-read (i.e., 10x Genomics) technologies.

All genomes were subject to a unified repeat library generation and annotation

workflow. Briefly, we identified repeats *de novo* for each assembly using 1.

RepeatModeler2 (Flynn et al., 2020) and 2. TransposonPSI (Haas). We combined those

repeat predictions with teleost repeats extracted from RepeatMasker (Smit et al.)

libraries and all sequences from the FishTEDB database (Shao et al., 2018)

http://www.fishtedb.org). These sequences (576,007 in total) were classified using the

RepeatClassifier module of RepeatModeler. We also clustered repeats at 80%

sequence identity using USEARCH (Edgar, 2010) as a way to group and ultimately

enumerate repeats based on sequence divergence (see Supp. Methods). We ran

RepeatMasker on all 16 genome assemblies using all 576,007 sequences and

integrated the aforementioned RepeatClassifier and USEARCH cluster IDs into the

RepeatMasker output. Finally, we used RepeatClassifier taxonomy and USEARCH

cluster membership as alternative grouping mechanisms to characterize the distribution

of repeats within and among genomes (e.g., using heatmaps and rank abundance

plots), and to ordinate genomes in repeat space (i.e., using PCA). Regional repeat

abundance distributions for target gene families were compared to those from random

samples of genes across the genome using resampling-style hypothesis tests. All

downstream repeat analysis and visualizations were carried out using the R statistical

language (R_Core_Team, 2019).

Gene family evolution

We used protein annotations from 21 teleost genomes (Figure 2; see Supp.

Methods for accession numbers) to better understand gene family size evolution in

seadragon and syngnathid lineages. We defined putative gene families via all-by-all

blastp (Altschul et al., 1990) and clustering with mcl (Enright et al., 2002), then we

conducted a series of gene family size evolution analyses using CAFE 5 (Mendes et al.,

2020). Briefly, we first fit an error model to account for artifactual (e.g., genome

assembly or annotation error) family size variation, which was applied to subsequent

CAFE runs. We fit a model assuming a single rate of gene family expansion/contraction

(λ) to the data and identified gene families evolving especially rapidly using CAFE's

internal likelihood ratio tests (LRTs).

Of those gene families showing evidence for rapid evolution, we identified

subsets for which branch-specific LRTs suggested extreme λs along branches of

interest: the terminal *P. eques* and *P. taeniolatus* branches, the internal branch leading

to the seadragon lineage, and the internal branch leading to the syngnathid lineage. We

also assigned each gene family to a KEGG orthology (KO) ID if possible, and we used

this information to perform KEGG pathway overrepresentation analysis using

ClusterProfiler (Yu et al., 2012), with respect to the branch-specific rapidly evolving

gene families. Lastly, we fit several multiple-λ models to the data in order to test

hypotheses of overall λ differences (from a global λ) for the internal seadragon branch

and the internal syngnathid branch. To test these hypotheses we used CAFE to perform

100 gene family simulations, fit the aforementioned models to the simulated data sets,

and compared observed likelihood ratios (LRs) from the data to the LR distributions

from the simulations. For details, see the Supplementary Methods.

***fgf* and *fgfr* gene family characterization**

We collected fibroblast growth factor (FGF) and receptor (FGFR) amino acid

sequences from Ensembl of several percomorph species (see Supplementary

Methods). We aligned orthologs and screened the genome assemblies of syngnathids

and outgroups based on Hidden Markov Model (HMM) profiles from the alignments,

according to methods described in Small et al. (Small et al., 2016). In some cases, we

supplemented these sensitive searches with regional RNA-seq read alignments to

correctly define exon boundaries. We conducted targeted, lineage-specific tests of

positive selection using branch site-models in PAML (see Supplemental Methods) and

tested for deleterious mutations using Provean (Choi & Chan, 2015).

X-ray tomography and 3D model reconstruction

We obtained a euthanized adult male weedy seadragon, fixed in neutral buffered

formalin from the Tennessee Aquarium (Chattanooga, TN), and scanned it using a

resolution of 54 µm voxels on a Zeiss XRadia 620 Versa X-ray microscope at the

University of Oregon Knight Campus for Accelerating Scientific Impact's X-ray Imaging

Core Facility. The anterior of the fish including the head and cleithrum were scanned

again at 17 µm voxel resolution. Composite virtual three-dimensional reconstructions of

the unstained specimen were generated using Dragonfly Pro and Dragonfly software

(Object Research Systems). After completing scans of the unstained specimen, which

allowed high contrast visualization of electron-dense bony structures, the fish was then

dehydrated through an ethanol series, stained with an iodine-based X-ray contrast

agent to enhance imaging of soft tissues, and a section of the rostral part of the stained

tail that includes a pair of leafy appendages was scanned at 27 µm voxel resolution

(Figure 1; Figure S2; Supplemental Methods).

RESULTS

PacBio assembly with Hi-C scaffolding yields chromosome-level genome models

for the leafy and weedy seadragon

We estimated the haploid genome sizes for leafy and weedy seadragons to be

644.0 Mb and 597.3 Mb, respectively, based on k-mer frequency analysis of Illumina

WGS data (Vurture et al., 2017). From this analysis we also estimated genome-wide

heterozygosity for each individual to be 0.27 and 0.33%. Polished PacBio assemblies

were 664.24 and 650.38 Mb, with scaffold N50s of 19.59 and 9.90 Mb. The longest Flye

scaffolds were 38.02 and 29.83 Mb, and BUSCO completeness frequencies were 95%

for both genomes. After scaffolding both polished PacBio assemblies using *P. eques*

Hi-C reads, we obtained 23 putative chromosome models for each genome (Figure S3),

which reflect 93.22% and 96.10% of the total length for final leafy and weedy

seadragon genome models. These assembly and completeness metrics, our ability to

annotate 22,256 and 22,043 protein-coding genes in the respective genomes, and

extensive evidence for conserved synteny between the two seadragon assemblies

(Figure S4), all support that these genome models are of high quality relative to

available resources for teleost fishes. These genomes will therefore serve as useful

tools for a variety of current comparative genomic analyses.

Seadragon karyotypes are conserved relative to other syngnathid genomes but

lack one of two chromosome fusions observed in *Syngnathus* and *Hippocampus*

273   Haploid chromosome number in syngnathid fishes, as assessed by karyotyping

and genetic mapping, is reported to be 22 or 24 in seahorse (*Hippocampus*) species

(Vitturi & Catalano, 1988; Vitturi et al., 1998) and 22 in Gulf pipefish (*Syngnathus*

scovelli) (Small et al., 2016). In Gulf pipefish, the reduction in chromosome number from

24, the putative ancestral number in ray-finned fishes (Mank & Avise, 2006) to 22 likely

resulted from fusion of two pairs of chromosomes orthologous to Chromosomes 1 and

24 and to Chromosomes 14 and 23 in platyfish (*X. maculatus*), a representative

outgroup percomorph with 24 chromosomes (Small et al., 2016). Though a haploid

number of 24 chromosomes in seahorse was reported (with some published confusion

about in which species this was observed (Vitturi & Catalano, 1988; Vitturi et al., 1998)

the reference genome for tiger tail seahorse (*H. comes*) (Lin et al., 2016), provides

conserved gene synteny evidence that both ancestral chromosome fusions were

already present in the common ancestor of seahorses and *Syngnathus* pipefish (Figure

S5). As stated, our inferred seadragon haploid chromosome number of 23 is based on

the size distribution of Hi-C scaffolds (Figure S3). The seadragon lineage apparently

shares only the ancestral Chromosome 1 to Chromosome 24 fusion with the

seahorse+*Syngnathus* pipefish ancestor, leaving one to one orthologs of platyfish

Chromosomes 14 and 23 (Figure S5).

Seadragon genomes are highly repetitive for their compact size, with large

contributions from relatively recent TE expansions

A correlation between eukaryotic genome size and the proportion of a genome

classifiable as TEs has led to an appreciation that TEs can be an important driver of

genome size evolution (Kidwell, 2002). The relationship is particularly apparent among

some of the well characterized genomes of teleost fish model species. Genome size

and TE proportion share a positive, tightly linear relationship in a comparison of green

spotted puffer (*Tetraodon nigroviridis*), threespine stickleback (*G. aculeatus*), medaka

(*Oryzias latipes*), and zebrafish (*Brachydanio rerio*), whose genomes span a range from

roughly 358 Mb to over 1.37 Gb (Gao et al., 2016). *Syngnathus* pipefish have genomes

that rival green spotted puffer in genomic compactness, with assembly lengths of 307.0

Mb for the Gulf pipefish (Small et al., 2016) (*S. scovelli*), and 324.33 Mb for the greater

pipefish (*S. acus*) (NCBI RefSeq Genome GCF_901709675.1). In the spectrum of

vertebrate genomes, known seahorse genomes are also diminutive, estimated at 421

Mb for the lined seahorse (*H. erectus*) (Li et al., 2021) and 494 Mb for the tiger tail

seahorse (*H. comes*) (Lin et al., 2016).

Our final genome assembly sizes for seadragons were 666.5 Mb for leafy

seadragon and 652.2 Mb for weedy seadragon, which are appreciably larger than those

of *Syngnathus* and *Hippocampus* but still much smaller than many other teleost

genomes. Surprisingly, as much as 58.7% of the leafy and 57.9% of the weedy

seadragon genome is composed of repetitive sequences, as classified by our workflow

(see Methods). To assess whether seadragons or syngnathids are exceptional in their

repetitive DNA characteristics, we measured the proportion of repetitive sequence in

sixteen teleost genomes (Figure S1) using an in-common repeat reference library to

create a standardized basis for comparison. Species were chosen for taxonomic

breadth and because their genomes were assembled from long- or linked-read

sequence data (except tiger tail seahorse). Genomes of five syngnathids were

represented, including a seahorse and tail-brooding *Syngnathus* pipefish, the

seadragons, and a basal lineage relative to these, the abdominal-brooding bluestripe

pipefish (*D. excisus*).

We confirmed a strong, positive relationship between repeat content and

genome size for the 16 genomes using phylogenetic generalized least squares (PGLS)

regression ($t_{16,14}$=4.88; *p*=0.0002; *b*=0.00048), but seadragons are notable outliers with

large, positive residuals (Figure 3). Both seadragon genomes are unusually bloated with

repetitive DNA among teleosts for their relatively compact size, a feature either derived

specifically in the seadragon lineage or an ancestral lineage. This latter hypothesis is

not yet testable given currently available genome assemblies but can be addressed

with the addition of syngnathid genomes at key phylogenetic positions.

Repeat density was high across seadragon chromosomes, in stark contrast with

the greater pipefish (Figure 4), which shows repeat "hotspots" almost exclusively near

chromosome ends, and a close outgroup to syngnathids, common dragonet

(*Callionymus lyra*), which has relatively uniform, low repeat density in general (Figure

S6).

Several TE classes contribute notably to the large repeatomes of seadragons.

The *Tc1* family of the *Tc1/Mariner* superfamily of transposases is a major contributor to

repeatome composition variation among teleosts as revealed by PCA based on within-

repeatome relative class proportions, with seadragons, platyfish (*X. maculatus*), and

Northern pike (*Esox lucius*) genomes influenced heavily by abundant *Tc1* repeats

(Figure 3). Phylogenetic patterns of repeat abundance among the fish lineages we

analyzed suggest *Tc1* expansion in the syngnathid lineage, given lower *Tc1*

abundances in the close outgroups of common dragonet, (*C. lyra*), and Pacific bluefin

tuna (*T. orientalis*). Among syngnathids, *Tc1* repeats compose a disproportionately

large fraction of seadragon repeatomes (Figure 3). This class of "cut-and-paste" DNA

transposons is widespread in animals and especially common in teleost fishes, with

high abundance and variability among species (Gao et al., 2017; Gao et al., 2016). In

fact, phylogenetic evidence suggests *Tc1* transposons are still active and recently

expanding in some neoteleosts, such as threespine stickleback (*G. aculeatus*) (Gao et

al., 2017).

The second most abundant, classifiable TE category in the seadragon genomes

was the *BovB* family of non-LTR LINE retrotransposons (Figure 3; Figure S7). This TE

family is restricted to animal taxa, where its members are patchily distributed among

lineages and have been inferred via phylogenetic analysis to populate animal genomes

through horizontal transfer, perhaps via metazoan parasites (Ivancevic & Chuong, 2020;

Ivancevic et al., 2018). *BovB* density variation along seadragon chromosomes (Figure

S7) was largely concordant with overall repeat density patterns (Figure S6), suggesting

common mechanisms or constraints for the expansion of at least some TE families.

We also discovered an apparent expansion of *Tigger* transposases in the

syngnathid clade, which are members of the *pogo* superfamily closely related to

Tc1/Mariner (Gao et al., 2020). *Tigger* repeats are overrepresented in syngnathids

compared to outgroups but proportionally are conserved among the five syngnathid

genomes we analyzed (Figure 3). Unlike *Tc1* and *BovB*, *Tigger* repeat density variation

along seadragon and *Syngnathus* pipefish chromosomes deviates from the collective

pattern of repeat density (Figure S8), with the highest-density *Tigger* regions more

centrally located, although the potential significance of this is unclear.

Because the grouping and enumeration of repeats according to classification

alone comes with a loss of evolutionary resolution and precision, we also analyzed

repeats as clusters of sequences with ≥ 80% sequence identity. PCA based on this

cluster-wise treatment of repeats revealed that leafy and weedy seadragon genomes

are quite divergent in repeat space from the other teleost species, including the three

other syngnathids compared (Figure 5; Figure S9). In particular four repeat clusters are

heavily influential in this regard, one of which was not classifiable but especially

abundant in scaffolds unassigned to chromosomes (Cluster 20902), two of which

belong to the *BovB* LINEs mentioned above (Clusters 10278 and 10626), and the last

(Cluster 14395) belonging to the *Tc1* group of transposases mentioned above (Figure 5;

Supplemental File 1).

We also generated rank abundance distributions (RADs) for repeat clusters in

the 16 fish genomes, an approach from ecology used to understand community

evenness and major and rare constituents (Whittaker, 1965). Interestingly, leafy and

weedy seadragon RADs show markedly high repeat cluster abundances relative to the

other genomes for repeat ranks 1-20 (Figure 5). Among the highest-ranking repeats for

both seadragon species are the four repeat clusters mentioned above as major

contributors to the distinctiveness of seadragon repeatomes. These findings, along with

the observation that members of the top-ranking seadragon repeat clusters are rare in

the other genomes (Supplemental File 1), suggest that the unique repetitive features of

seadragon genomes are driven largely by recent expansions of *BovB* and *Tc1*, and a

yet-to-be-classified cluster.

Gene family contractions in Syngnathidae involve innate immunity, and

seadragon-specific expansions are associated with vesicular trafficking

We identified 290 total gene families as having expanded or contracted at a rate

(λ) significantly higher than the background λ among 21 teleost species. Of these, 109

showed evidence for rapid size evolution along the leafy seadragon branch, 113 along

the weedy seadragon branch, 38 along the internal seadragon branch, and 31 along the

internal branch leading to syngnathids (Figure 2; Supplemental File 2). Based on 100

simulations of gene family evolution along the tree, we also inferred that λs for the

internal seadragon and syngnathid branches are respectively distinct from the global λ

for the tree, and likely distinct from one another (Figure S10).

Among the gene families most likely to have evolved rapidly in size in the

ancestral syngnathid lineage, at least seven families related to innate immunity

experienced contractions (Supplemental File 2), in contrast to multiple lines of evidence

for expansion of inflammation and innate immunity gene families in teleosts relative to

other vertebrates (Balla et al., 2020; Boudinot et al., 2011; Chang et al., 2021; Howe et

al., 2016; Mattingsdal et al., 2018). Specifically, we found evidence for syngnathid-

specific contractions in NACHT, TRIM, and GIMAP gene families, consistent with some

of the immunity and detoxification pathway gene families depleted in the genome of the

Manado pipefish *(Microphis manadensis)* (Zhang et al., 2020).

Several functional categories (KEGG pathways) were overrepresented among

gene families with large size changes in the seadragon lineage, including cancer,

cardiomyopathy, and immunity (Supplemental File 2), primarily due to contraction

events. In terms of gene family expansions along the branch leading to seadragons,

however, two families with roles in vesicular trafficking - *Vacuolar Protein Sorting-*

Associated Protein 13B (*vps13b*) and *Coatomer Protein Complex Subunit Beta 2*

(*copb2*) – are notable. Sequences with high similarity to *copb2*, which encodes one

subunit of a Golgi budding and vesicular trafficking protein complex (Waters et al.,

1991), are especially abundant in seadragon genomes relative to other syngnathids and

teleosts (Supplemental File 2).

Though tuna (*T. orientalis*) and platyfish (*X. latipes*), for example, have two

paralogs of *copb2*, we could find evidence for only a single gene copy in the genomes

of seahorse (*H. comes*) and pipefish (*S. acus*). By contrast, we detected the presence of

at least nine and six copies (on the 26 longest Hi-C scaffolds) in leafy and weedy

seadragon genomes (Supplemental File 3). We also found many sequences matching

copb2 on several short scaffolds, suggesting that the repetitive nature of this region

prevented these from being incorporated into chromosome-scale scaffolds, or, but less

likely (Figure S11), that they could be redundant artifacts. One of the *copb2* paralogs

(on Pequ Hi-C scaffold 13, Ptae Hi-C scaffold 14) is the likely ortholog of platyfish

copb2 on the orthologous chromosome (Xmac 6), with the remaining copies likely

expanding secondarily via an unknown mechanism.

Given the repetitive nature of these regions, we hypothesized that TE activity

may have played a role in seadragon *copb2* expansion. Specifically, we first tested

whether the seadragon-expanded TE classes of *BovB* and *Tc1* are overrepresented in

the immediate vicinity (1-kb flanking both sides) of *copb2* copies, relative to 499

randomly resampled gene groups. Both *BovB* (4.24-fold enriched; $p < 0.002$) and *Tc1*

(1.39-fold enriched; $p < 0.002$) repeat classes were significantly overrepresented in

copb2 regions (Figure S11; Supplemental File 4). We secondarily performed naïve

hypothesis tests with the false discovery rate (FDR) controlled at 0.05 to identify other

TE classes with potential enrichment in these regions, revealing *Rex-Babar* LINEs, and

Tigger and *Charlie* transposons as additional candidates (Supplemental File 4).

Syngnathid fishes have lost several FGF family genes, most notably *fgf3* and *fgf4*

The FGF and FGFR gene families include well-studied ligand and receptor

signaling molecules central to vertebrate craniofacial, limb, dermal appendage, sensory

placode, and hindbrain development, among many other functions (reviewed in Xie et

al., 2020). Because of the prominence of FGF signaling in the morphogenesis of traits

that are distinctively modified in syngnathids, we explored whether FGF ligands or their

receptors are exceptional in seadragons and in three other representative lineages of

syngnathids: flagtail pipefishes (the most basal lineage of the four), seahorses, and

Syngnathus pipefishes (Figure 2).

The FGF and FGFR families in seadragons and other syngnathid fishes conform,

with two notable exceptions mentioned below, to a broadly typical complement of gene

paralogs relative to other percomorphs. Within the Syngnathidae, there are several

lineage-specific losses of FGF and FGFR genes (Table S1). We did not detect *fgf9* in

any syngnathid. Seadragons and *Syngnathus* pipefish have apparently lost *fgf4-like*,

while tiger tail seahorse (*Hippocampus comes*) and bluestripe pipefish (*Doryrhamphus*

excisus) have retained it. *fgf17* is missing birds (Abramyan, 2015), in seadragons and

bluestripe pipefish but is present in tiger tail seahorse and greater pipefish. Because

flagtail pipefishes, the lineage to which bluestripe pipefish belongs, are basal to the

clade containing seadragons, seahorses, and *Syngnathus* pipefish (Longo et al., 2017),

fgf17 therefore must have been lost at least twice. Evolutionary loss of *fgf17*, though

uncommon*,* is not unique to syngnathid lineages; orthologs of this gene have been

independently erased from other distant taxa, such as the medaka genus *Oryzias*

(Canestro et al., 2007). In addition, at least two clades of *fgf5* and *fgfr1b* are present in

seadragons, tiger tail seahorse, and blue-stripe pipefish, but these genes were not

detected in the two *Syngnathus* pipefish genomes (*S. acus* and *S. scovelli*).

The most startling gene losses from the FGF family are shared across the

seadragon, tiger tail seahorse, greater pipefish, and bluestripe pipefish lineages: all are

missing *fgf3* and *fgf4* (Figure 6; Table S1). In percomorph outgroups to the syngnathids,

fgf3, *fgf4*, and *fgf19* are clustered, with *fgf3* flanked by *ccnd1* and *lto1,* genes that are

also missing from seadragons, seahorses, and greater pipefish (but bluestripe pipefish

has retained *lto1* (Figure 6). The expected neighbors normally on the *fgf19* side of the

cluster (*zgc:153993* and *ano1*) are present in seadragons and both lineages of

pipefishes, but bluestripe pipefish has lost *fgf19*. Close outgroups to the syngnathids,

the razorfish (*Aeoliscus strigatus*) and the mandarinfish (*Synchiropus splendidus*) (Longo

et al., 2017) retain intact *fgf3/4/19* clusters, making it likely that drastic alteration to the

cluster is a syngnathid synapomorphy (Figure S12). In the common ancestor of lobe-

finned and ray-finned fishes, *fgf3*, *fgf4*, and *fgf19* were likely already clustered, and

remain tandemly arrayed in representatives from jawed vertebrate lineages as

separated as mammals and teleosts (Oulion et al., 2012). The syngnathid lineage has

therefore experienced a degree of change in this cluster that is perhaps unprecedented

throughout gnathostome evolution.

Exceptional absences of FGF and FGFR members in extant syngnathid

genomes suggest that some property of their syntenic neighborhoods is inherently

volatile. We tested whether the local repeat landscape flanking significant regions of

FGF, FGFR, and other developmental gene loss in syngnathids – the *fgf3/4/19* cluster,

fgf4l, *fgf5*, *fgf17*, *fgfr1b, eve1* (Small et al., 2016), and *tbx4* (Lin et al., 2016; Small et al.,

2016) - differs from that of genes in general, by using the leafy seadragon genome and

the close syngnathid outgroup *C. lyra* (common dragonet) as a pre-loss comparator.

While no individual repeat class was significantly enriched in these regions (of either

genome) after FDR adjustment, several show nominal evidence for enrichment and are

at high density in the leafy seadragon regions, including *Tc1*, *BovB*, *hAT-Ac*, and

additional LINEs *Rex-Babar*, and *L2* (Figure S13; Supplemental File 4). Furthermore, we

found support for the alternative hypothesis that *overall* repeat density in general is

enriched in these regions (Figure S13; Supplemental File 4), but only for the leafy

seadragon (Pequ: 1.45-fold enrichment; p = 0.022), and not the dragonet genome (Clyr:

1.11-fold enrichment; p = 0.27).

Limited evidence for compensatory evolution in the FGF pathway

FGF paralogs sometimes overlap in their expression and ability to compensate,

partially or fully, for loss-of-function of one another in specific developmental contexts.

In zebrafish, for example, knock down of either *fgf3* or *fgf10* causes relatively mild

effects in lateral line migration and neuromast maturation, but there are severe lateral

line defects when expression of both paralogs is depleted (Nechiporuk & Raible, 2008).

We explored whether those paralogs most recently diverged from *fgf3* and *fgf4*, or with

overlapping developmental roles, showed evidence for compensatory evolution prior to

the syngnathid radiation.

Syngnathid and close outgroup sequences are largely conserved for *fgf4-like*,

fgf6a, *fgf7*, *fgf8a*, *fgf8b*, *fgf10a*, *fgf10b*, *fgf19*, *fgf20a, fgf20b*, *fgf22,* and *fgf24*, and

exhibit no evidence of positive selection along the syngnathid branch (Table S2). We did

find potential evidence for positive selection in *fgf6b*, although this test result was within

the range of false discovery (Figure S14; Table S2). Second, although the gene was lost

in the seadragons and blue striped pipefish lineages, we found the strongest evidence

for positive selection on *fgf17*, which is retained in seahorse and *Syngnathus* pipefish

(Table S2).

Like the ligands, the syngnathid FGF receptors are largely conserved, and we

found no evidence of syngnathid lineage-specific positive selection for non-canonical

fgfrl1a and *fgfrl1b*, nor for canonical *fgfr1a, fgfr2*, *fgfr3*, or *fgfr4*. In contrast, *fgfr1b*

presented some evidence for positive selection (although not robust to false discovery),

displaying surprising syngnathid-specific substitutions (Table S2). These include a likely

deleterious substitution within the activation loop of the kinase domain, a position

conserved not only across vertebrate Fgfr1 proteins, but across all canonical FGFR

paralogy groups (Figure S15) (provean score= -6.123).

Although we did not find evidence for lineage-specific positive selection using

codon-based models, leafy and weedy seadragons share a derived six amino acid

deletion in a conserved region of the Fgf16 protein (Figure S16), while seahorse has

amino acid substitutions in this same motif. Other representative percomorphs,

chicken, and human Fgf16 protein sequences are identical across this region indicating

a high level of conservation, though no function has yet been ascribed to this protein

domain. Divergence of seadragon *fgf16* is not limited to the coding sequence. A

putative regulatory change is hinted at by the absence in both dragons of an

approximately 240 bp conserved non-coding element (CNE) that is well preserved

across percomorph fishes 5' of *fgf16* (Figure S17).

miRNA-seq data from seadragons reveal loss of conserved microRNAs

Although microRNAs (miRNAs) are important developmental regulators (Bizuayehu

& Babiak, 2014), there are currently no annotations of these genes in syngnathids. In

the leafy and weedy seadragons, we identified 261 and 251 miRNA genes that produce

331 and 318 unique mature miRNAs (Supplemental File 5). We found absences of

numerous conserved miRNA genes including *mir10a* and *mir196b,* hox miRNAs genes

that were reported missing in the Gulf pipefish genome assembly (Small et al., 2016).

Six of the additional highly conserved missing miRNAs belong to two miRNA clusters

from the miR-130 family. These absences include *mir130a*, *mir301a*, *mir130b*, *mir301b*,

mir130c-2, and *mir454b*. While *mir130a, mir301a, and mir301b* are convergently

missing in platyfish (*X. maculatus*) and medaka (*O. latipes*), absences of *mir130b,*

mir130c-2, and mir454b have not yet been reported in other vertebrate species (Figure

S18) (Desvignes et al., 2019; Desvignes et al., 2021; Kelley et al., 2021). In some

tetrapods and teleosts, *mir301a* is located in the first intron of *ska2*. The seadragons,

Gulf and greater pipefishes, and tiger tail seahorse are missing *ska2* as well as *mir301a*

and *mir130a.* Throughout vertebrates, *mir130b* and *mir301b* (and, in teleosts and

Coelacanth, additionally *mir130c-2* and *mir454b*) are located in the intergenic region

between *sdf2l1* and *top3b*. Though these adjacent protein coding genes and the

immediate syntenic neighborhood are conserved in seadragons, Gulf and greater

pipefishes, and tiger tail seahorse, this cluster of microRNAs is not.

DISCUSSION

Exploring the seadragon genomes in a comparative phylogenetic context has lifted

a veil on the evolution of seadragon-specific traits and has also revealed intriguing

evolutionary facets of this unusual vertebrate family, the Syngnathidae, as a whole. We

found that both leafy and weedy seadragon genomes stand out among their relatives in

having a surprisingly large contingent of repetitive DNA. This pattern appears to have

been driven largely by recent expansions of *BovB* and *Tc1*, and a yet-to-be-classified

cluster of repetitive sequences specific to the seadragon lineage. One possible

explanation for the large difference we observed between seadragon and *Syngnathus*

pipefish repeatome size and genomic distribution could be a difference in historical

effective population size (N_e). In this case, negative selection to remove rapidly

expanding repeats from a population would be less effective in the face of strong

genetic drift (small N_e) (Lynch, 2007; Lynch & Conery, 2003). Differences in k-mer-based

heterozygosity estimates from individual WGS data are at least consistent with this

idea. A heterozygosity estimate based on the Gulf pipefish data (Small et al., 2016) is

roughly three times the seadragon estimates (1.01% versus 0.27% and 0.33% for leafy

and weedy seadragons, respectively).

The explanation presented above for TE expansion assumes that deleterious

effects are common, likely through interruption of coding regions or promoters, or by

regional silencing via chromatin changes (Liu et al., 2018). However, TEs can also

contribute new genes or gene regulatory sequences when a host genome co-opts

("domesticates") these exogenous genetic elements (Bejerano et al., 2006). Some

classes of TE provide, for example, binding sites for master transcriptional regulators

NANOG and OCT4 throughout the mouse and human genomes but in largely non-

overlapping sets of loci between the taxa, potentially restructuring, in lineage-specific

ways, the transcriptional networks for developmental pluripotency (Kunarso et al.,

2010). TEs and repetitive DNA in general can fuel gene family expansion or contraction

by precipitating unequal crossover events; Hahn et al. (2007) suggest an "explosion" of

TEs in the primate lineage could be linked to its accelerated gene content evolution in

primates. Repeatome expansion, such as what we here observe in seadragons, could

have had disruptive impacts on gene regulatory networks and gene content, subject to

subsequent evolution via negative *or* positive selection.

Using global and targeted approaches we explored expansion and contraction

of gene families in seadragons and their relatives, and possible connections of gene

content changes to observed TE distributions. Perhaps the most obvious trend we

observed for the syngnathid lineage in general was contraction of particular immunity

and detoxification pathway gene families, some of which have previously been

described (Lin et al., 2016; Zhang et al., 2020). GTPase of the immunity-associated

protein genes (GIMAP), for example, was among the contracted families in seadragons

that was also detected in the seahorse genome (Lin et al., 2016), and our more

comprehensive analysis supports that the GIMAP contraction occurred prior to the

radiation of syngnathids. This family exists as a single eight-member cluster in

mammals (Krucken et al., 2004). GIMAP genes have multiplied in other teleost lineages,

numbering up to nearly 190 genes. Balla et al. (2020) found zebrafish GIMAP genes

respond to pathogenic viral exposure and suggest the gene expansion could have been

evolutionarily favored by the relatively long period that hatchlings must rely on innate

immunity before the development of a functional adaptive immune system. Male-

brooded syngnathid embryos might enjoy a luxury not available to free-spawned

progeny like those of zebrafish, namely pathogen climate control afforded by the

paternal immune system before their own adaptive immunity develops. The male's

immune system strikes a balance of defending against foreign agents but without

rejecting his own brood. The syngnathid contraction of the GIMAP family is also

interesting, therefore, in light of the possibility that *gimap4*, which does persist in the

lineage, could promote immunologic tolerance to embryos in brood pouch tissues (Roth

et al., 2020). Roth et al. (2020) based this assertion on the observation that *gimap4* is

up-regulated in *Syngnathus* pregnancy tissues, where it could contribute to local

suppression of the lymphocyte population (Roth et al., 2020; Schnell et al., 2006).

We detected seadragon-specific copy number expansion of a coatamer

complex gene *copb2*. Variants of the *BovB* and *Tc1* transposable elements were

enriched surrounding the supernumerary gene copies, suggesting a TE-driven

mechanism for the expansion. Copb2 forms part of a protein complex involved in

retrograde vesicle budding from the Golgi apparatus and secretion of macromolecule

cargo, such as collagen, which is critical for bone and connective tissue development.

Mice and fish developing with deficits in *copb2* have delayed bone mineralization and

low bone density, as well as defects in type II collagen trafficking and secretion (Marom

et al., 2021). Zebrafish mutants of *copb2* develop mispatterned, kinked notochords with

poorly formed vacuoles and a disorganized perinotochordal basement membrane,

which is secreted by notochord sheath cells (Coutinho et al., 2004). In teleosts, the

notochord, particularly its sheath, plays an instructive role in patterning the vertebrae

(Gray et al., 2014; Peskin et al., 2020).

Given the elaborated bony exoskeleton in seadragons, their stiff bodies with

connective tissue-dense leafy ornaments, and their kinked axial skeletons with varied

and regionalized vertebral forms (Figure 1), the proliferation of *copb2* gene sequences

in their genomes ignites curiosity about the possible evolutionary developmental

consequences of this expansion for the seadragons' unique exo- and endoskeletons.

Kinked vertebral columns in the guppy (*Poecelia reticulata*) mutant *curveback* are

characterized by wedge-shaped vertebrae (Gorman et al., 2007). Similar to the guppy

mutant and to the developmental malformation of vertebrae in human Scheuermann's

kyphosis sufferers, vertebrae are keystone-shaped in the weedy seadragon at locations

of spinal curvature (Figure 1; Figure S2).

One of the most surprising gene family reductions we uncovered is shared by all

of the syngnathid lineages we explored; it is the loss of *fgf3* and *fgf4*. Loss of these two

fibroblast growth factor genes in syngnathids is striking because their orthologs in other

vertebrates have long been hypothesized to play nearly indispensable pleiotropic

developmental roles in the pharyngeal arches, teeth, brain, cranial placodes, epidermal

appendages, limbs, and the segmental axis (Anderson et al., 2020; Boulet et al., 2004;

Cooper et al., 2017; Cooper et al., 2018; Crump et al., 2004; David et al., 2002;

Jackman et al., 2013; Leger & Brand, 2002; Lu et al., 2006; Maves et al., 2002; Miyake

& Itoh, 2013; Nechiporuk & Raible, 2008; Prykhozhij & Neumann, 2008; Reuter et al.,

2019; Walshe et al., 2002; Wang et al., 2007). It is reasonable to weigh whether losing

these two multifunctional signaling ligands could have had significant and broad

consequences to both deeply conserved developmental pathways and their

morphological readouts. Another possibility is that these developmental pathways had

diverged neutrally, or through changes in other pathway members, from anciently

conserved functions along the syngnathid lineage, permitting genes that had once been

critical to become expendable. It is nevertheless valuable to note that syngnathid fishes

share peculiarities in many features from the constellation of vertebrate traits that *fgf3*

and *fgf4* are known to help pattern.

Dermal integuments of syngnathid fishes are bony plates, and in several

syngnathid lineages including the seadragons these have been elaborated to

magnificence, sometimes independently. A subset of these plates in seadragons bear

blunted struts of bone that end in fleshy paddle-shaped ornaments, the "leaves" and

"weeds" of the species featured in this report (Figure 1). Another percomorph clade, the

pufferfishes, are adorned with bony spines likely evolved from elasmoid scales. Shono

et al., (Shono et al., 2019), showed that *fgf3* (and *fgf1a*) are expressed in developing

pufferfish dermal spines. Given this and a trove of other evidence for an FGF signaling

role in the development and diversification of scales, spines, and denticles in ray-finned

and chondrichthyan fishes (Albertson et al., 2018; Aman et al., 2018; Cooper et al.,

2017; Cooper et al., 2018; Daane et al., 2016; Kim et al., 2019; Shono et al., 2019),

absence of *fgf3* and *fgf4* in the often heavily armored, elaborately spined syngnathids

clearly suggests that derived mechanisms for integumentary bone development are at

play in this lineage.

Syngnathids have evolved elongated faces with an unusual hyoid apparatus

integral to specialized suction feeding (Figure 1) (Leysen et al., 2010), and they are

toothless. These craniofacial features arise developmentally from complex interactions

between - at least - endoderm, mesoderm, and neural crest components. Both *fgf3* and

fgf4 are expressed in the pharyngeal arches, which form skeletal elements of the jaw,

hyoid, and, in fish, the gill supports (David et al., 2002; Herzog et al., 2004; Niswander &

Martin, 1992) *fgf3* plays an essential role in craniofacial development. When *fgf3* is

knocked out in zebrafish, mutants die within 7-9 days; craniofacial defects (particularly

the full loss of cartilage in gill arches) are predicted to be the cause of their early

mortality (Herzog et al., 2004). Additionally, when *fgf3* expression is disrupted in the

mesendoderm in developing zebrafish, an "inverted" backward directed ceratohyal

cartilage is formed (David et al., 2002). It is tempting to speculate that evolutionary loss

of these genes could have led to altered craniofacial architecture of the elongated

syngnathid face, either directly or through the effects of genetic compensation

percolating through this signaling pathway.

Syngnathid toothlessness is also particularly interesting given *fgf3* and *fgf4*

expression in zebrafish dental epithelium and their suspected roles in tooth

morphogenesis (Jackman et al., 2004). Gilbert et al. (2019) propose a model for

patterning of dentition in ray-finned fishes in which a tooth primordium acts as an

organizer that induces the development of subsequent teeth, likely via secretion of Fgf3

and Fgf4. Other tooth developmental genes are known to be lost or reduced in copy

number in syngnathids, including *eve1* (Small et al., 2016); see above) and P/Q-rich

SCPP enamel/enameloid matrix genes (Lin et al., 2016; Qu et al., 2021; Zhang et al.,

2020). These losses imply an erosion of tooth development pathways spanning

induction to mineralization, with our discovery of *fgf3/4* loss enlarging the pool of

candidate causative genes for edentulism in syngnathids.

The syngnathid central nervous system features its own peculiarities. Bennedetti

(1991) described greater pipefish (*S. acus*) and long-snouted seahorse (*Hippocampus*

guttulatus) to have highly modified or no discernable Mauthner neurons, the large,

rhombomere 4 (r4) reticulospinal neurons critical for the rapid "C-start" escape

response in many fishes and some amphibians (reviewed in Korn & Faber, 2005).

Syngnathids are reputed also to lack mechanosensory lateral line neuromasts, from

which the Mauthner cells receive synaptic inputs (Benedetti, 1991; Korn & Faber, 1975).

In zebrafish, joint impairment of *fgf3* and *fgf8* impacts segmental identity of

rhombomeres 3 and 5 and their reticulospinal neurons (Maves et al., 2002; Walshe et

al., 2002). Furthermore, depletion of *fgf3* and *fgf10* reduced the number of zebrafish

posterior lateral line neuromasts and inhibited migration of the placode down the length

of the body (Nechiporuk & Raible, 2008). These observations present a compelling case

for future interrogation of a link between *fgf3* loss and derived hindbrain and sensory

development in syngnathids.

Though it is known that paralogous FGF ligands can compensate for one another

in some experimental contexts (Maves et al., 2002; Walshe et al., 2002), in general, we

did not find sweeping evidence for adaptive evolution of paralogous FGF proteins or

their receptors in the wake of syngnathid *fgf3* and *fgf4* gene losses. Correlated changes

could instead have included evolution of non-coding, regulatory sequences of

FGF/FGFR genes or changes to other gene families that interact with FGF signaling.

Syngnathids, we found, have lost six deeply conserved miRNAs in the miR-130 family.

Biological implications for these missing miRNA genes are uncertain, though it is

possible that their losses could relate to derived syngnathid-specific traits and gene

pathway changes. For instance, angiogenesis and tissue remodeling are critical

components in syngnathid male pregnancy tissues (Stolting & Wilson, 2007), and

mir130a is connected to vascular repatterning in mammals and teleosts (Chen & Gorski,

2008; Singh et al., 2017) mir130 genes are also known to affect FGF signaling. In

chicken, knockdowns of *mir130* lead to increased expression of *fgf8* and FGF pathway

member, *Erk 1/2* (Bobbs et al., 2012; Lopez-Sanchez et al., 2015). There is also a

suspected regulatory relationship between FGF receptors and these miRNAs; in human

differentiating stem cells, nuclear located FGFR1, nFGFR1, binds to the promoters of

mir301b and mir130b (Stachowiak & Stachowiak, 2016; Terranova et al., 2015). Loss of

miRNAs that regulate and are potentially regulated by FGFs and FGFRs could indicate

a further extension of the derived restructuring of FGF signaling pathways in

syngnathids.

Evidence for positive selection as a compensatory consequence of having lost

fgf3 and *fgf4* in a syngnathid ancestor proved to be scarce. *fgf16*, however, provides a

possible example of a separate evolutionary scenario: derived functional change in the

seadragon lineage, at both the protein and cis-regulatory levels. Leafy and weedy

seadragon fgf16 proteins share an unusual deletion in a deeply conserved motif, and

the seadragon gene also appears to have lost a 5' non-coding sequence that is

otherwise conserved among syngnathids and distantly related percomorphs. In

zebrafish, this gene is necessary for outgrowth of the pectoral fin, upstream of *fgf4* and

fgf8 (Nomura et al., 2006), and we show it is expressed similarly in fin margins in a

representative percomorph, threespine stickleback (Figure S19) (*Gasterosteus*

aculeatus). The eponymous "leaves" of seadragons that are fleshy extensions borne on

bony supports extending from the dermal plates, are apparently stiffened by a core of

collagenous tissue rather than ossified structures such as fin rays (Figure 1). Homology

of the leafy ornaments with fins might pertain only at the level of shared genetic

pathways, such as in the case of a cooption or re-deployment of FGF-signaling to elicit

proliferation and outgrowth of these superficially fin-like structures. A role in scale or

dermal plate patterning for *fgf16* is not known from teleosts, though in birds the

orthologous gene, potentially acting with *fgfr1*, can suppress both *shh* expression and

downy feather elongation when locally overexpressed (Chen et al., 2016). The

seadragon-specific changes in a known AER and integument-patterning gene are

intriguing in the context of this morphological oddity, the leafy "appendages". The

seahorse fgf16 protein, we found, bears substitutions in the same amino acid motif

deleted in seadragons. The fact that two lineages that have evolved elaborate bony and

fleshy ornaments (seadragons and seahorses) both show divergence in a deeply

conserved motif of this gene is tantalizing and warrants further comparative work.

753   **SUMMARY**

The genome models of the leafy and weedy seadragon provide a first step

towards unlocking how the fantastical forms of these animals and their close relatives

have evolved. Even with our preliminary analyses we have identified likely recent

expansions of transposable elements, with hypothesized involvement in significant

changes to gene families of known consequence, the most surprising being the loss of

fgf3 and fgf4 in the sygnathid lineage. Repetitive DNA, including that generated by the

activity of TEs, can precipitate genomic rearrangements such as deletions, duplications,

and inversions (reviewed in Bourque et al., 2018; Schrader & Schmitz, 2019). It is

therefore likely that historical and possibly continuing TE activity in genomes of the

syngnathid lineage has resulted in alterations to genome structure and content, with

extreme consequences for developmental genetic programs that are otherwise deeply

conserved across vertebrates.

ACKNOWLEDGEMENTS

We are truly grateful for the dedicated efforts of L. Matsushige and the seadragon
husbandry staff of the Birch Aquarium at Scripps, who preserved *P. eques* samples
crucial for this work. We also thank the Tennessee Aquarium, and particularly Aquarist
K. Hurt for generous sharing of *P. taeniolatus* samples. We are indebted to M.
Weitzman and D. Turnbull from the UO GC3F for critical assistance with library
preparation and sequencing. We would like to acknowledge staff at Carl Zeiss
Microscopy, LLC, particularly J. Mancuso, A. Browning, K. Skinner, and R. White for
collaborating to scan the Gulf pipefish in Figure S2. We are grateful for T. Desvignes for
guidance on the miRNA annotations. We especially thank C. Kimmel for his helpful
comments on the manuscript. This work was funded by National Institutes of Health
grants RR032670 and P50GM098911 (to WAC), National Science Foundation grant
OPP-2015301 (to WAC, SB, and CMS), and Oregon Research Excellence Funds (to
WAC). HMH was supported by the Genetics Training Program (NIH T32GM007413) at
UO. Purchase of the Zeiss Xradia 620 Versa at the University of Oregon was made
possible with funding support from the M.J. Murdock Charitable Trust Grant SR-
201812008 (to WAC).

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

788
Data Accessibility: All raw sequencing data associated with this study will be available
upon peer-reviewed publication via the NCBI Sequencing Read Archive (SRA), under
BioProjects PRJNA765699 and PRJNA765702. Annotation and summary files not in the
Supplementary Information will be available upon peer-reviewed publication in the
Dryad repository.

FIGURES

Figure 1. The anatomy of the weedy seadragon includes remarkably elongated facial
features terminating in toothless, upturned jaws, an unusual hyoid apparatus
specialized for suction feeding, a bony exoskeleton with elaborate spines that support
fleshy "leaves", and a sinusoidal spine of rib-less vertebrae that vary in shape and size.
A) a lateral view of the skeleton of *Phyllopterix taeniolatus* reconstructed by X-ray
microscopy. B) detail of the head (the ceratohyal, c, and urohyal, u, of the hyoid
apparatus are noted). C) Detail of the pectoral region (lateral view) showing a dorsal,
unpaired "leafy" appendage support surrounded by other dermal plates with much
shorter spines. D) optical cross section of the tail through a pair of leafy appendage
spines. E) optical cross section through the same appendages as in D) but with a
contrast agent that reveals the fleshy leaves, seen edge on (denoted by l). F) lateral view
shows keystone-shaped vertebrae at curvatures - both kyphosis and lordosis - of the

820 **Figure 2.** Genomes of 21 teleost species provide phylogenetic context for gene family
821 evolution in seadragons and syngnathid fishes. Represented are evolutionary

relationships among a sample of morphologically diverse teleosts, according to a time-calibrated phylogenetic tree adapted from Rabosky et al. (2018). We used published and newly generated protein-coding gene annotations for the species pictured here to understand putative gene family expansions and contractions in lineages of interest, specifically the seadragons. The number of gene families with statistical evidence for expansion (top values) and contraction (bottom values) along all terminal branches and internal syngnathiform branches are shown. Note that the four-letter species symbols in the legend are used throughout this article.

Figure 3. Seadragon genomes demonstrate a large fraction of repetitive DNA relative to other teleost fishes, characterized by substantial contributions from TcMar-Tc1 transposons and unclassified constituents. A) Scatterplot showing the strong, positive relationship between genome assembly size (x-axis) and the proportion of the genome annotated as repetitive for 16 recent teleost genome assemblies. Note that seadragon genomes (pequ and ptae) are especially repetitive (~60%) given their relatively small size (~650 Mb). Dashed line shows a phylogenetic generalized least squares (PGLS) regression fit, and syngnathid genomes are in red. B) A Principal Components Analysis (PCA) biplot shows similarity of the 16 genomes based on relative frequencies of repeat classes. TcMar-Tc1 and "Unknown" repeat classes load especially heavily on PC1 and PC2, respectively, as indicated by vectors (purple arrows) drawn in the space, and these contribute to the distinctiveness of syngnathid genomes (in red). C) Barplots

855
856
Figure 4. Leafy and weedy seadragon chromosomes are densely and non-uniformly
populated by repetitive DNA. A) Orthologous pairs of the 23 seadragon chromosomes,

ordered ascending from shortest to longest leafy seadragon sequence. For each pair
the leafy seadragon chromosome is on top, and the weedy seadragon ortholog is
below. Cases in which the orientation of the weedy seadragon chromosome has been
reversed to align with the leafy ortholog are denoted by "(-)". B) Chromosome models of
the greater pipefish (*Syngnathus acus*), also ascending from shortest to longest. Lines
connect seadragon and pipefish chromosomes with large regions of orthology, as
inferred via conserved synteny analysis. Dashed lines reflect the fusion of two ancestral
syngnathid chromosomes that is derived in the lineage leading to *Syngnathus* and
Hippocampus. Overall repeat basepair occupation of 100 kb windows (expressed as a
proportion) is presented as a heatmap.
869
870
871

873
873
Figure 5. Relatively recent repeat expansions in the seadragon lineage drive the
uniqueness of seadragon repeatomes. Repeat clusters defined at the 80% sequence
identity level and quantified by the proportion of total genome length they occupy were
used to conduct principal components analysis (PCA). A) A bi-plot of the first two PCA
axes, showing extreme separation of seadragons from the other 14 species in
repeatome space, particularly along the leading axis of variation (x-axis). Purple arrows
represent individual repeat clusters, and how strongly (and in what direction) they
influence the position of genomes in this repeatome space. Four repeat clusters (shown
in a box) with especially large loadings on PC1 are strongly associated with seadragon
repeatome uniqueness. B) Rank abundance distributions for the top 100 repeat clusters
in each of the 16 species included in the repeat analysis. The top 20 clusters for each
seadragon species are consistently elevated in abundance, relative to corresponding
ranks in the other 14 fish genomes. Shown in a box are the top 15 leafy seadragon
clusters, their ranks in weedy seadragon, and the repeat class to which they likely
belong. Note that the top 3 (and 5th) clusters in the list correspond to the boxed clusters
in panel A.
890
891
892

Figure 6. The *fgf3/4/19* cluster locus has experienced surprising gene losses in the syngnathid lineage. While retaining the same immediate gene neighbors, the FGF cluster became inverted in the percomorph fishes relative to outgroups like zebrafish and tetrapods. Several genes (grey rectangles) appear to have been deleted from the locus in the syngnathids, though not all of the losses are shared. *spg21*, *shcbp1*, and *b3galt2*, which neighbor the locus in the syngnathid lineage, are separated from the FGF cluster by other genes in non-syngnathid percomorphs like platyfish. Genes that appear lost from the locus cannot be found anywhere else in the genome assemblies of pipefishes, seahorses or seadragons. Arrows show gene order and orientation; grey rectangles represent apparent gene losses.

REFERENCES

Abramyan, J. (2015). Lineage-specific loss of FGF17 within the avian orders Galliformes and Passeriformes. *Gene*, *563*(2), 180-189. https://doi.org/10.1016/j.gene.2015.03.027

Albertson, R. C., Kawasaki, K. C., Tetrault, E. R., & Powder, K. E. (2018). Genetic analyses in Lake Malawi cichlids identify new roles for Fgf signaling in scale shape variation. *Commun Biol*, *1*, 55. https://doi.org/10.1038/s42003-018-0060-4

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *J Mol Biol*, *215*(3), 403-410. https://doi.org/10.1016/S0022-2836(05)80360-2

Aman, A. J., Fulbright, A. N., & Parichy, D. M. (2018). Wnt/beta-catenin regulates an ancient signaling network during zebrafish scale development. *Elife*, *7*. https://doi.org/10.7554/eLife.37001

Anderson, M. J., Magidson, V., Kageyama, R., & Lewandoski, M. (2020). Fgf4 maintains Hes7 levels critical for normal somite segmentation clock function. *Elife*, *9*. https://doi.org/10.7554/eLife.55608

Balla, K. M., Rice, M. C., Gagnon, J. A., & Elde, N. C. (2020). Linking virus discovery to immune responses visualized during zebrafish infections. *Curr Biol*, *30*(11), 2092-2103 e2095. https://doi.org/10.1016/j.cub.2020.04.031

Bejerano, G., Lowe, C. B., Ahituv, N., King, B., Siepel, A., Salama, S. R., Rubin, E. M., Kent, W. J., & Haussler, D. (2006). A distal enhancer and an ultraconserved exon are derived from a novel retroposon. *Nature*, *441*(7089), 87-90. https://doi.org/10.1038/nature04696

Benedetti. (1991). Mauthner neurons in syngnathid bony fish.

Bizuayehu, T. T., & Babiak, I. (2014). MicroRNA in teleost fish. *Genome Biol Evol*, *6*(8), 1911-1937. https://doi.org/10.1093/gbe/evu151

Bobbs, A. S., Saarela, A. V., Yatskievych, T. A., & Antin, P. B. (2012). Fibroblast growth factor (FGF) signaling during gastrulation negatively modulates the abundance of microRNAs that regulate proteins required for cell migration and embryo patterning. *J Biol Chem*, *287*(46), 38505-38514. https://doi.org/10.1074/jbc.M112.400598

Boudinot, P., van der Aa, L. M., Jouneau, L., Du Pasquier, L., Pontarotti, P., Briolat, V., Benmansour, A., & Levraud, J. P. (2011). Origin and evolution of TRIM proteins: new insights from the complete TRIM repertoire of zebrafish and pufferfish. *PLoS ONE*, *6*(7), e22022. https://doi.org/10.1371/journal.pone.0022022

Boulet, A. M., Moon, A. M., Arenkiel, B. R., & Capecchi, M. R. (2004). The roles of Fgf4 and Fgf8 in limb bud initiation and outgrowth. *Dev Biol*, *273*(2), 361-372. https://doi.org/10.1016/j.ydbio.2004.06.012

Bourque, G., Burns, K. H., Gehring, M., Gorbunova, V., Seluanov, A., Hammell, M., Imbeault, M., Izsvak, Z., Levin, H. L., Macfarlan, T. S., Mager, D. L., & Feschotte,

C. (2018). Ten things you should know about transposable elements. *Genome Biol*, *19*(1), 199. https://doi.org/10.1186/s13059-018-1577-z

Bruna, T., Hoff, K. J., Lomsadze, A., Stanke, M., & Borodovsky, M. (2021). BRAKER2: automatic eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein database. *NAR Genom Bioinform*, *3*(1), lqaa108. https://doi.org/10.1093/nargab/lqaa108

Bushnell, B. (2014). BBMap: A fast, accurate, splice-aware aligner. *sourceforge.net/projects/bbmap/*.

Canestro, C., Yokoi, H., & Postlethwait, J. H. (2007). Evolutionary developmental biology and genomics. *Nat Rev Genet*, *8*(12), 932-942. https://doi.org/10.1038/nrg2226

Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: an analysis tool set for population genomics. *Mol Ecol*, *22*(11), 3124-3140. https://doi.org/10.1111/mec.12354

Catchen, J. M., Amores, A., Hohenlohe, P., Cresko, W., & Postlethwait, J. H. (2011). Stacks: building and genotyping Loci de novo from short-read sequences. *G3 (Bethesda)*, *1*(3), 171-182. https://doi.org/10.1534/g3.111.000240

Catchen, J. M., Conery, J. S., & Postlethwait, J. H. (2009). Automated identification of conserved synteny after whole-genome duplication. *Genome Res*, *19*(8), 1497-1505. https://doi.org/10.1101/gr.090480.108

Chang, M. X., Xiong, F., Wu, X. M., & Hu, Y. W. (2021). The expanding and function of NLRC3 or NLRC3-like in teleost fish: Recent advances and novel insights. *Dev Comp Immunol*, *114*, 103859. https://doi.org/10.1016/j.dci.2020.103859

Chen, C. K., Ng, C. S., Wu, S. M., Chen, J. J., Cheng, P. L., Wu, P., Lu, M. Y., Chen, D. R., Chuong, C. M., Cheng, H. C., Ting, C. T., & Li, W. H. (2016). Regulatory differences in natal down development between altricial zebra finch and precocial chicken. *Molecular Biology and Evolution*, *33*(8), 2030-2043. https://doi.org/10.1093/molbev/msw085

Chen, Y., & Gorski, D. H. (2008). Regulation of angiogenesis through a microRNA (miR-130a) that down-regulates antiangiogenic homeobox genes GAX and HOXA5. *Blood*, *111*(3), 1217-1226. https://doi.org/10.1182/blood-2007-07-104133

Chin, C. S., Alexander, D. H., Marks, P., Klammer, A. A., Drake, J., Heiner, C., Clum, A., Copeland, A., Huddleston, J., Eichler, E. E., Turner, S. W., & Korlach, J. (2013). Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods*, *10*(6), 563-569. https://doi.org/10.1038/nmeth.2474

Choi, Y., & Chan, A. P. (2015). PROVEAN web server: a tool to predict the functional effect of amino acid substitutions and indels. *Bioinformatics*, *31*(16), 2745-2747. https://doi.org/10.1093/bioinformatics/btv195

Connolly, R., Melville, A., & Keesing, J. (2002). Abundance, movement and individual identification of leafy seadragons, *Phycodurus eques* (Pisces: Syngnathidae). *Marine and Freshwater Research*, *53*, 777-780.

Cooper, R. L., Martin, K. J., Rasch, L. J., & Fraser, G. J. (2017). Developing an ancient epithelial appendage: FGF signalling regulates early tail denticle formation in sharks. *Evodevo*, *8*, 8. https://doi.org/10.1186/s13227-017-0071-0

Cooper, R. L., Thiery, A. P., Fletcher, A. G., Delbarre, D. J., Rasch, L. J., & Fraser, G. J. (2018). An ancient Turing-like patterning mechanism regulates skin denticle development in sharks. *Sci Adv*, *4*(11), eaau5484. https://doi.org/10.1126/sciadv.aau5484

Coutinho, P., Parsons, M. J., Thomas, K. A., Hirst, E. M., Saude, L., Campos, I., Williams, P. H., & Stemple, D. L. (2004). Differential requirements for COPI transport during vertebrate early development. *Dev Cell*, *7*(4), 547-558. https://doi.org/10.1016/j.devcel.2004.07.020

Crump, J. G., Maves, L., Lawson, N. D., Weinstein, B. M., & Kimmel, C. B. (2004). An essential role for Fgfs in endodermal pouch formation influences later craniofacial skeletal patterning. *Development*, *131*(22), 5703-5716. https://doi.org/10.1242/dev.01444

Daane, J. M., Rohner, N., Konstantinidis, P., Djuranovic, S., & Harris, M. P. (2016). Parallelism and epistasis in skeletal evolution identified through use of phylogenomic mapping strategies. *Molecular Biology and Evolution*, *33*(1), 162-173. https://doi.org/10.1093/molbev/msv208

David, N. B., Saint-Etienne, L., Tsang, M., Schilling, T. F., & Rosa, F. M. (2002). Requirement for endoderm and FGF3 in ventral head skeleton formation. *Development*, *129*(19), 4457-4468. https://doi.org/10.1242/dev.129.19.4457

Dawson, C. E. (1985). *Indo-Pacific pipefishes (Red Sea to the Americas)*. Gulf Coast Research Laboratory.

Desvignes, T., Batzel, P., Sydes, J., Eames, B. F., & Postlethwait, J. H. (2019). miRNA analysis with Prost! reveals evolutionary conservation of organ-enriched expression and post-transcriptional modifications in three-spined stickleback and zebrafish. *Sci Rep*, *9*(1), 3913. https://doi.org/10.1038/s41598-019-40361-8

Desvignes, T., Sydes, J., Montfort, J., Bobe, J., & Postlethwait, J. H. (2021). Evolution after whole-genome duplication: teleost microRNAs. *Molecular Biology and Evolution*, *38*(8), 3308-3331. https://doi.org/10.1093/molbev/msab105

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., & Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15-21. https://doi.org/10.1093/bioinformatics/bts635

Dudchenko, O., Batra, S. S., Omer, A. D., Nyquist, S. K., Hoeger, M., Durand, N. C., Shamim, M. S., Machol, I., Lander, E. S., Aiden, A. P., & Aiden, E. L. (2017). De novo assembly of the Aedes aegypti genome using Hi-C yields chromosome-length scaffolds. *Science*, *356*(6333), 92-95. https://doi.org/10.1126/science.aal3327

Dudchenko, O., Shamim, M. S., Batra, S. S., Durand, N. C., Musial, N. T., Mostofa, R., Pham, M., Glenn St Hilaire, B., Yao, W., Stamenova, E., Hoeger, M., Nyquist, S. K., Korchina, V., Pletch, K., Flanagan, J. P., Tomaszewicz, A., McAloose, D., Pérez Estrada, C., Novak, B. J., Omer, A. D., & Aiden, E. L. (2018). The

Juicebox Assembly Tools module facilitates *de novo* assembly of mammalian
genomes with chromosome-length scaffolds for under $1000. *bioRxiv*. <Go to
WoS>://WOS:000255412600014

Durand, N. C., Robinson, J. T., Shamim, M. S., Machol, I., Mesirov, J. P., Lander, E. S.,
& Aiden, E. L. (2016). Juicebox provides a visualization system for Hi-C contact
maps with unlimited zoom. *Cell Syst*, *3*(1), 99-101.
https://doi.org/10.1016/j.cels.2015.07.012

Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST.
Bioinformatics, *26*(19), 2460-2461. https://doi.org/10.1093/bioinformatics/btq461

Enright, A. J., Van Dongen, S., & Ouzounis, C. A. (2002). An efficient algorithm for large-
scale detection of protein families. *Nucleic Acids Res*, *30*(7), 1575-1584.
https://doi.org/10.1093/nar/30.7.1575

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A.
F. (2020). RepeatModeler2 for automated genomic discovery of transposable
element families. *Proc Natl Acad Sci U S A*, *117*(17), 9451-9457.
https://doi.org/10.1073/pnas.1921046117

Gao, B., Chen, W., Shen, D., Wang, S., Chen, C., Zhang, L., Wang, W., Wang, X., &
Song, C. (2017). Characterization of autonomous families of Tc1/mariner
transposons in neoteleost genomes. *Mar Genomics*, *34*, 67-77.
https://doi.org/10.1016/j.margen.2017.05.003

Gao, B., Shen, D., Xue, S., Chen, C., Cui, H., & Song, C. (2016). The contribution of
transposable elements to size variations between four teleost genomes. *Mob
DNA*, *7*, 4. https://doi.org/10.1186/s13100-016-0059-7

Gao, B., Wang, Y., Diaby, M., Zong, W., Shen, D., Wang, S., Chen, C., Wang, X., &
Song, C. (2020). Evolution of pogo, a separate superfamily of IS630-Tc1-mariner
transposons, revealing recurrent domestication events in vertebrates. *Mob DNA*,
11, 25. https://doi.org/10.1186/s13100-020-00220-0

Gibert, Y., Samarut, E., Ellis, M. K., Jackman, W. R., & Laudet, V. (2019). The first
formed tooth serves as a signalling centre to induce the formation of the dental
row in zebrafish. *Proc Biol Sci*, *286*(1904), 20190401.
https://doi.org/10.1098/rspb.2019.0401

Gorman, K. F., Tredwell, S. J., & Breden, F. (2007). The mutant guppy syndrome
curveback as a model for human heritable spinal curvature. *Spine (Phila Pa
1976)*, *32*(7), 735-741. https://doi.org/10.1097/01.brs.0000259081.40354.e2

Gray, R. S., Wilm, T. P., Smith, J., Bagnat, M., Dale, R. M., Topczewski, J., Johnson, S.
L., & Solnica-Krezel, L. (2014). Loss of col8a1a function during zebrafish
embryogenesis results in congenital vertebral malformations. *Dev Biol*, *386*(1),
72-85. https://doi.org/10.1016/j.ydbio.2013.11.028

Gurevich, A., Saveliev, V., Vyahhi, N., & Tesler, G. (2013). QUAST: quality assessment
tool for genome assemblies. *Bioinformatics*, *29*(8), 1072-1075.
https://doi.org/10.1093/bioinformatics/btt086

Haas, B. J. TransposonPSI. *http://transposonpsi.sourceforge.net*.

Hahn, M. W., Demuth, J. P., & Han, S. G. (2007). Accelerated rate of gene gain and loss
in primates. *Genetics*, *177*(3), 1941-1949.
https://doi.org/10.1534/genetics.107.080077

Herzog, W., Sonntag, C., von der Hardt, S., Roehl, H. H., Varga, Z. M., &
Hammerschmidt, M. (2004). Fgf3 signaling from the ventral diencephalon is
required for early specification and subsequent survival of the zebrafish
adenohypophysis. *Development*, *131*(15), 3681-3692.
https://doi.org/10.1242/dev.01235

Howe, K., Schiffer, P. H., Zielinski, J., Wiehe, T., Laird, G. K., Marioni, J. C., Soylemez,
O., Kondrashov, F., & Leptin, M. (2016). Structure and evolutionary history of a
large family of NLR proteins in the zebrafish. *Open Biol*, *6*(4), 160009.
https://doi.org/10.1098/rsob.160009

Ivancevic, A., & Chuong, E. B. (2020). Transposable elements teach T cells new tricks.
Proc Natl Acad Sci U S A, *117*(17), 9145-9147.
https://doi.org/10.1073/pnas.2004493117

Ivancevic, A. M., Kortschak, R. D., Bertozzi, T., & Adelson, D. L. (2018). Horizontal
transfer of BovB and L1 retrotransposons in eukaryotes. *Genome Biol*, *19*(1), 85.
https://doi.org/10.1186/s13059-018-1456-7

Jackman, W. R., Davies, S. H., Lyons, D. B., Stauder, C. K., Denton-Schneider, B. R.,
Jowdry, A., Aigler, S. R., Vogel, S. A., & Stock, D. W. (2013). Manipulation of Fgf
and Bmp signaling in teleost fishes suggests potential pathways for the
evolutionary origin of multicuspid teeth. *Evol Dev*, *15*(2), 107-118.
https://doi.org/10.1111/ede.12021

Jackman, W. R., Draper, B. W., & Stock, D. W. (2004). Fgf signaling is required for
zebrafish tooth development. *Dev Biol*, *274*(1), 139-157.
https://doi.org/10.1016/j.ydbio.2004.07.003

Jones, P., Binns, D., Chang, H. Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H.,
Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A. F., Sangrador-Vegas,
A., Scheremetjew, M., Yong, S. Y., Lopez, R., & Hunter, S. (2014). InterProScan
5: genome-scale protein function classification. *Bioinformatics*, *30*(9), 1236-1240.
https://doi.org/10.1093/bioinformatics/btu031

Kelley, J. L., Desvignes, T., McGowan, K. L., Perez, M., Rodriguez, L. A., Brown, A. P.,
Culumber, Z., & Tobler, M. (2021). microRNA expression variation as a potential
molecular mechanism contributing to adaptation to hydrogen sulphide. *J Evol
Biol*, *34*(6), 977-988. https://doi.org/10.1111/jeb.13727

Kidwell, M. G. (2002). Transposable elements and the evolution of genome size in
eukaryotes. *Genetica*, *115*(1), 49-63. https://doi.org/10.1023/a:1016072014259

Kim, D. I., Kai, W., Hosoya, S., Sato, M., Nozawa, A., Kuroyanagi, M., Jo, Y., Tasumi,
S., Suetake, H., Suzuki, Y., & Kikuchi, K. (2019). The genetic basis of scale-loss
phenotype in the rapid radiation of Takifugu fishes. *Genes (Basel)*, *10*(12).
https://doi.org/10.3390/genes10121027

Klanten, O. S., Gaither, M. R., Greaves, S., Mills, K., O'Keeffe, K., Turnbull, J.,
McKinnon, R., & Booth, D. J. (2020). Genomic and morphological evidence of
distinct populations in the endemic common (weedy) seadragon *Phyllopteryx*

taeniolatus (Syngnathidae) along the east coast of Australia. *PLoS ONE*, *15*(12), e0243446. https://doi.org/10.1371/journal.pone.0243446

Kolmogorov, M., Yuan, J., Lin, Y., & Pevzner, P. A. (2019). Assembly of long, error-prone reads using repeat graphs. *Nat Biotechnol*, *37*(5), 540-546. https://doi.org/10.1038/s41587-019-0072-8

Korn, H., & Faber, D. S. (1975). Inputs from the posterior lateral line nerves upon the goldfish Mauthner cell. I. Properties and synaptic localization of the excitatory component. *Brain Research*, *96*(2), 342-348. https://doi.org/10.1016/0006-8993(75)90745-3

Korn, H., & Faber, D. S. (2005). The Mauthner cell half a century later: a neurobiological model for decision-making? *Neuron*, *47*(1), 13-28. https://doi.org/10.1016/j.neuron.2005.05.019

Krucken, J., Schroetel, R. M., Muller, I. U., Saidani, N., Marinovski, P., Benten, W. P., Stamm, O., & Wunderlich, F. (2004). Comparative analysis of the human gimap gene cluster encoding a novel GTPase family. *Gene, 341*, 291-304. https://doi.org/10.1016/j.gene.2004.07.005

Kunarso, G., Chia, N. Y., Jeyakani, J., Hwang, C., Lu, X., Chan, Y. S., Ng, H. H., & Bourque, G. (2010). Transposable elements have rewired the core regulatory network of human embryonic stem cells. *Nat Genet*, *42*(7), 631-634. https://doi.org/10.1038/ng.600

Leger, S., & Brand, M. (2002). Fgf8 and Fgf3 are required for zebrafish ear placode induction, maintenance and inner ear patterning. *Mech Dev, 119*(1), 91-108. https://doi.org/10.1016/s0925-4773(02)00343-x

Leysen, H., Jouk, P., Brunain, M., Christiaens, J., & Adriaens, D. (2010). Cranial architecture of tube-snouted gasterosteiformes (*Syngnathus rostellatus* and *Hippocampus capensis*). *J Morphol, 271*(3), 255-270. https://doi.org/10.1002/jmor.10795

Li, C., Olave, M., Hou, Y., Qin, G., Schneider, R. F., Gao, Z., Tu, X., Wang, X., Qi, F., Nater, A., Kautt, A. F., Wan, S., Zhang, Y., Liu, Y., Zhang, H., Zhang, B., Zhang, H., Qu, M., Liu, S., Chen, Z., Zhong, J., Zhang, H., Meng, L., Wang, K., Yin, J., Huang, L., Venkatesh, B., Meyer, A., Lu, X., & Lin, Q. (2021). Genome sequences reveal global dispersal routes and suggest convergent genetic adaptations in seahorse evolution. *Nat Commun*, *12*(1), 1094. https://doi.org/10.1038/s41467-021-21379-x

Lin, Q., Fan, S., Zhang, Y., Xu, M., Zhang, H., Yang, Y., Lee, A. P., Woltering, J. M., Ravi, V., Gunter, H. M., Luo, W., Gao, Z., Lim, Z. W., Qin, G., Schneider, R. F., Wang, X., Xiong, P., Li, G., Wang, K., Min, J., Zhang, C., Qiu, Y., Bai, J., He, W., Bian, C., Zhang, X., Shan, D., Qu, H., Sun, Y., Gao, Q., Huang, L., Shi, Q., Meyer, A., & Venkatesh, B. (2016). The seahorse genome and the evolution of its specialized morphology. *Nature*, *540*(7633), 395-399. https://doi.org/10.1038/nature20595

Liu, N., Lee, C. H., Swigut, T., Grow, E., Gu, B., Bassik, M. C., & Wysocka, J. (2018). Selective silencing of euchromatic L1s revealed by genome-wide screens for L1 regulators. *Nature*, *553*(7687), 228-232. https://doi.org/10.1038/nature25179

Longo, S. J., Faircloth, B. C., Meyer, A., Westneat, M. W., Alfaro, M. E., & Wainwright,
P. C. (2017). Phylogenomic analysis of a rapid radiation of misfit fishes
(Syngnathiformes) using ultraconserved elements. *Mol Phylogenet Evol*, *113*, 33-
48. https://doi.org/10.1016/j.ympev.2017.05.002

Lopez-Sanchez, C., Franco, D., Bonet, F., Garcia-Lopez, V., Aranega, A., & Garcia-
Martinez, V. (2015). Negative Fgf8-Bmp2 feed-back is regulated by miR-130
during early cardiac specification. *Dev Biol*, *406*(1), 63-73.
https://doi.org/10.1016/j.ydbio.2015.07.007

Lu, P., Minowada, G., & Martin, G. R. (2006). Increasing Fgf4 expression in the mouse
limb bud causes polysyndactyly and rescues the skeletal defects that result from
loss of Fgf8 function. *Development*, *133*(1), 33-42.
https://doi.org/10.1242/dev.02172

Lynch, M. (2007). *The Origins of Genome Architecture*. Oxford University Press.

Lynch, M., & Conery, J. S. (2003). The origins of genome complexity. *Science*,
302(5649), 1401-1404. https://doi.org/10.1126/science.1089370

Mank, J. E., & Avise, J. C. (2006). Phylogenetic conservation of chromosome numbers
in Actinopterygiian fishes. *Genetica*, *127*(1-3), 321-327.
https://doi.org/10.1007/s10709-005-5248-0

Marom, R., Burrage, L. C., Venditti, R., Clement, A., Blanco-Sanchez, B., Jain, M., Scott,
D. A., Rosenfeld, J. A., Sutton, V. R., Shinawi, M., Mirzaa, G., DeVile, C.,
Roberts, R., Calder, A. D., Allgrove, J., Grafe, I., Lanza, D. G., Li, X., Joeng, K.
S., Lee, Y. C., Song, I. W., Sliepka, J. M., Batkovskyte, D., Washington, M.,
Dawson, B. C., Jin, Z., Jiang, M. M., Chen, S., Chen, Y., Tran, A. A., Emrick, L.
T., Murdock, D. R., Hanchard, N. A., Zapata, G. E., Mehta, N. R., Weis, M. A.,
Scott, A. A., Tremp, B. A., Phillips, J. B., Wegner, J., Taylor-Miller, T., Gibbs, R.
A., Muzny, D. M., Jhangiani, S. N., Hicks, J., Stottmann, R. W., Dickinson, M. E.,
Seavitt, J. R., Heaney, J. D., Eyre, D. R., Undiagnosed Diseases, N., Westerfield,
M., De Matteis, M. A., & Lee, B. (2021). COPB2 loss of function causes a
coatopathy with osteoporosis and developmental delay. *Am J Hum Genet*,
108(9), 1710-1724. https://doi.org/10.1016/j.ajhg.2021.08.002

Martin-Smith, K. M., & Vincent, A. C. J. (2006). Exploitation and trade of Australian
seahorses, pipehorses, sea dragons and pipefishes (Family Syngnathidae).
Oryx, *40*(2), 141-151. https://doi.org/10.1017/s003060530600010x

Mattingsdal, M., Jentoft, S., Torresen, O. K., Knutsen, H., Hansen, M. M., Robalo, J. I.,
Zagrodzka, Z., Andre, C., & Gonzalez, E. B. (2018). A continuous genome
assembly of the corkwing wrasse (*Symphodus melops*). *Genomics*, *110*(6), 399-
403. https://doi.org/10.1016/j.ygeno.2018.04.009

Maves, L., Jackman, W., & Kimmel, C. B. (2002). FGF3 and FGF8 mediate a
rhombomere 4 signaling activity in the zebrafish hindbrain. *Development*,
129(16), 3825-3837. https://www.ncbi.nlm.nih.gov/pubmed/12135921

Mendes, F. K., Vanderpool, D., Fulton, B., & Hahn, M. W. (2020). CAFE 5 models
variation in evolutionary rates among gene families. *Bioinformatics*.
https://doi.org/10.1093/bioinformatics/btaa1022

Miyake, A., & Itoh, N. (2013). Fgf22 regulated by Fgf3/Fgf8 signaling is required for
zebrafish midbrain development. *Biol Open*, *2*(5), 515-524.
https://doi.org/10.1242/bio.20134226

Nechiporuk, A., & Raible, D. W. (2008). FGF-dependent mechanosensory organ
patterning in zebrafish. *Science*, *320*(5884), 1774-1777.
https://doi.org/10.1126/science.1156547

Niswander, L., & Martin, G. R. (1992). Fgf-4 expression during gastrulation, myogenesis,
limb and tooth development in the mouse. *Development*, *114*(3), 755-768.
https://www.ncbi.nlm.nih.gov/pubmed/1618140

Nomura, R., Kamei, E., Hotta, Y., Konishi, M., Miyake, A., & Itoh, N. (2006). Fgf16 is
essential for pectoral fin bud formation in zebrafish. *Biochem Biophys Res*
Commun, *347*(1), 340-346. https://doi.org/10.1016/j.bbrc.2006.06.108

Oulion, S., Bertrand, S., & Escriva, H. (2012). Evolution of the FGF Gene Family. *Int J*
Evol Biol, *2012*, 298147. https://doi.org/10.1155/2012/298147

Peskin, B., Henke, K., Cumplido, N., Treaster, S., Harris, M. P., Bagnat, M., & Arratia, G.
(2020). Notochordal signals establish phylogenetic identity of the teleost spine.
Curr Biol, *30*(14), 2805-2814 e2803. https://doi.org/10.1016/j.cub.2020.05.037

Prykhozhij, S. V., & Neumann, C. J. (2008). Distinct roles of Shh and Fgf signaling in
regulating cell proliferation during zebrafish pectoral fin development. *BMC Dev*
Biol, *8*, 91. https://doi.org/10.1186/1471-213X-8-91

Qu, M., Liu, Y., Zhang, Y., Wan, S., Ravi, V., Qin, G., Jiang, H., Wang, X., Zhang, H.,
Zhang, B., Gao, Z., Huysseune, A., Zhang, Z., Zhang, H., Chen, Z., Yu, H., Wu,
Y., Tang, L., Li, C., Zhong, J., Ma, L., Wang, F., Zheng, H., Yin, J., Witten, P. E.,
Meyer, A., Venkatesh, B., & Lin, Q. (2021). Seadragon genome analysis provides
insights into its phenotype and sex determination locus. *Sci Adv*, *7*(34).
https://doi.org/10.1126/sciadv.abg5196

R_Core_Team. (2019). A language and environment for statistical computing. R
Foundation for Statistical Computing,

. *Vienna, Austria. URL https://www.R-project.org/*.

Rabosky, D. L., Chang, J., Title, P. O., Cowman, P. F., Sallan, L., Friedman, M.,
Kaschner, K., Garilao, C., Near, T. J., Coll, M., & Alfaro, M. E. (2018). An inverse
latitudinal gradient in speciation rate for marine fishes. *Nature*, *559*(7714), 392-
395. https://doi.org/10.1038/s41586-018-0273-1

Reuter, I., Jackels, J., Kneitz, S., Kuper, J., Lesch, K. P., & Lillesaar, C. (2019). Fgf3 is
crucial for the generation of monoaminergic cerebrospinal fluid contacting cells in
zebrafish. *Biol Open*, *8*(6). https://doi.org/10.1242/bio.040683

Roth, O., Solbakken, M. H., Torresen, O. K., Bayer, T., Matschiner, M., Baalsrud, H. T.,
Hoff, S. N. K., Brieuc, M. S. O., Haase, D., Hanel, R., Reusch, T. B. H., & Jentoft,
S. (2020). Evolution of male pregnancy associated with remodeling of canonical
vertebrate immunity in seahorses and pipefishes. *Proc Natl Acad Sci U S A*,
117(17), 9431-9439. https://doi.org/10.1073/pnas.1916251117

Schnell, S., Demolliere, C., van den Berk, P., & Jacobs, H. (2006). Gimap4 accelerates
T-cell death. *Blood*, *108*(2), 591-599. https://doi.org/10.1182/blood-2005-11-4616

Schrader, L., & Schmitz, J. (2019). The impact of transposable elements in adaptive
evolution. *Mol Ecol*, *28*(6), 1537-1549. https://doi.org/10.1111/mec.14794

Shao, F., Wang, J., Xu, H., & Peng, Z. (2018). FishTEDB: a collective database of
transposable elements identified in the complete genomes of fish. *Database*
(Oxford), *2018*. https://doi.org/10.1093/database/bax106

Shono, T., Thiery, A. P., Cooper, R. L., Kurokawa, D., Britz, R., Okabe, M., & Fraser, G.
J. (2019). Evolution and developmental diversity of skin spines in pufferfishes.
iScience, *19*, 1248-1259. https://doi.org/10.1016/j.isci.2019.06.003

Singh, B. N., Tahara, N., Kawakami, Y., Das, S., Koyano-Nakagawa, N., Gong, W.,
Garry, M. G., & Garry, D. J. (2017). Etv2-miR-130a-Jarid2 cascade regulates
vascular patterning during embryogenesis. *PLoS ONE*, *12*(12), e0189010.
https://doi.org/10.1371/journal.pone.0189010

Small, C. M., Bassham, S., Catchen, J., Amores, A., Fuiten, A. M., Brown, R. S., Jones,
A. G., & Cresko, W. A. (2016). The genome of the Gulf pipefish enables
understanding of evolutionary innovations. *Genome Biol*, *17*(1), 258.
https://doi.org/10.1186/s13059-016-1126-6

Smit, A. F. A., Hubley, R., & Green, P. RepeatMasker at http://repeatmasker.org.

Stachowiak, M. K., & Stachowiak, E. K. (2016). Evidence-based theory for integrated
genome regulation of ontogeny—an unprecedented role of nuclear FGFR1
signaling. *J Cell Physiol*, *231*(6), 1199-1218. https://doi.org/10.1002/jcp.25298

Stervander, M., & Cresko, W. A. (2021). A highly contiguous nuclear genome assembly
of the mandarinfish *Synchiropus splendidus* (Syngnathiformes:

Callionymidae). *G3 (Bethesda)*, *in press*. https://doi.org/10.1093/g3journal/jkab306

Stiller, J., da Fonseca, R. R., Alfaro, M. E., Faircloth, B. C., Wilson, N. G., & Rouse, G.
W. (2021). Using ultraconserved elements to track the influence of sea-level
change on leafy seadragon populations. *Mol Ecol*, *30*(6), 1364-1380.
https://doi.org/10.1111/mec.15744

Stiller, J., Wilson, N. G., Donnellan, S., & Rouse, G. W. (2017). The leafy seadragon,
Phycodurus eques, a flagship species with low but structured genetic variability.
J Hered, *108*(2), 152-162. https://doi.org/10.1093/jhered/esw075

Stiller, J., Wilson, N. G., & Rouse, G. W. (2015). A spectacular new species of
seadragon (Syngnathidae). *R Soc Open Sci*, *2*(2), 140458.
https://doi.org/10.1098/rsos.140458

Stolting, K. N., & Wilson, A. B. (2007). Male pregnancy in seahorses and pipefish:
beyond the mammalian model. *BioEssays*, *29*(9), 884-896.
https://doi.org/10.1002/bies.20626

Terranova, C., Narla, S. T., Lee, Y. W., Bard, J., Parikh, A., Stachowiak, E. K.,
Tzanakakis, E. S., Buck, M. J., Birkaya, B., & Stachowiak, M. K. (2015). Global
developmental gene programing involves a nuclear form of fibroblast growth
factor receptor-1 (FGFR1). *PLoS ONE*, *10*(4), e0123380.
https://doi.org/10.1371/journal.pone.0123380

Vitturi, R., & Catalano, E. (1988). Karyotypes in two species of the genus *Hippocampus*
(Pisces: Syngnatiformes). *Marine Biology*, *99*(1), 119-121.
https://doi.org/10.1007/bf00644985

Vitturi, R., Libertini, A., Campolmi, M., Calderazzo, F., & Mazzola, A. (1998).
Conventional karyotype, nucleolar organizer regions and genome size in five
Mediterranean species of Syngnathidae (Pisces, Syngnathiformes). *Journal of*
Fish Biology, *52*(4), 677-687. https://doi.org/10.1111/j.1095-8649.1998.tb00812.x

Vurture, G. W., Sedlazeck, F. J., Nattestad, M., Underwood, C. J., Fang, H., Gurtowski,
J., & Schatz, M. C. (2017). GenomeScope: fast reference-free genome profiling
from short reads. *Bioinformatics*, *33*(14), 2202-2204.
https://doi.org/10.1093/bioinformatics/btx153

Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo, C.
A., Zeng, Q., Wortman, J., Young, S. K., & Earl, A. M. (2014). Pilon: an
integrated tool for comprehensive microbial variant detection and genome
assembly improvement. *PLoS ONE*, *9*(11), e112963.
https://doi.org/10.1371/journal.pone.0112963

Walshe, J., Maroon, H., McGonnell, I. M., Dickson, C., & Mason, I. (2002).
Establishment of hindbrain segmental identity requires signaling by FGF3 and
FGF8. *Curr Biol*, *12*(13), 1117-1123. https://doi.org/10.1016/s0960-
9822(02)00899-0

Wang, X. P., Suomalainen, M., Felszeghy, S., Zelarayan, L. C., Alonso, M. T., Plikus, M.
V., Maas, R. L., Chuong, C. M., Schimmang, T., & Thesleff, I. (2007). An
integrated gene regulatory network controls stem cell proliferation in teeth. *PLoS*
Biol, *5*(6), e159. https://doi.org/10.1371/journal.pbio.0050159

Waterhouse, R. M., Seppey, M., Simao, F. A., Manni, M., Ioannidis, P., Klioutchnikov,
G., Kriventseva, E. V., & Zdobnov, E. M. (2018). BUSCO Applications from
quality assessments to gene prediction and phylogenomics. *Molecular Biology*
and Evolution, *35*(3), 543-548. https://doi.org/10.1093/molbev/msx319

Waters, M. G., Serafini, T., & Rothman, J. E. (1991). 'Coatomer': a cytosolic protein
complex containing subunits of non-clathrin-coated Golgi transport vesicles.
Nature, *349*(6306), 248-251. https://doi.org/10.1038/349248a0

Weisenfeld, N. I., Kumar, V., Shah, P., Church, D. M., & Jaffe, D. B. (2017). Direct
determination of diploid genome sequences. *Genome Res*, *27*(5), 757-767.
https://doi.org/10.1101/gr.214874.116

Whittaker, R. H. (1965). Dominance and diversity in land plant communities: numerical
relations of species express the importance of competition in community function
and evolution. *Science*, *147*(3655), 250-260.
https://doi.org/10.1126/science.147.3655.250

Xie, Y., Su, N., Yang, J., Tan, Q., Huang, S., Jin, M., Ni, Z., Zhang, B., Zhang, D., Luo,
F., Chen, H., Sun, X., Feng, J. Q., Qi, H., & Chen, L. (2020). FGF/FGFR
signaling in health and disease. *Signal Transduct Target Ther*, *5*(1), 181.
https://doi.org/10.1038/s41392-020-00222-7

Yu, G., Wang, L. G., Han, Y., & He, Q. Y. (2012). clusterProfiler: an R package for
comparing biological themes among gene clusters. *OMICS, 16*(5), 284-287.
https://doi.org/10.1089/omi.2011.0118

Zhang, Y.-H., Ravi, V., Qin, G., Dai, H., Zhang, H.-X., Han, F.-M., Wang, X., Liu, Y.-H.,
Yin, J.-P., Huang, L.-M., Venkatesh, B., & Lin, Q. (2020). Comparative genomics
reveal shared genomic changes in syngnathid fishes and signatures of genetic
convergence with placental mammals. *National Science Review, 7*(6), 964-977.
https://doi.org/10.1093/nsr/nwaa002

1342