The PhoPQ two-component system promotes carbapenem tolerance in Enterobacterales.

Andrew N. Murtha[*1,2], Misha I. Kazi[*3], Richard D. Schargel[3], Trevor Cross[1,2], Conrad Fihn[4], Erin E. Carlson[4,5,6,7], Joseph M. Boll[#3], Tobias Dörr[#1,2,8]

[1] Weill Institute for Cell and Molecular Biology, Cornell University, Ithaca, NY 14853, USA

[2] Department of Microbiology, Cornell University, Ithaca NY 14853, USA

[3] Department of Biology, University of Texas Arlington, Arlington, TX, USA

[4] Department of Medicinal Chemistry, University of Minnesota, Minneapolis, Minnesota 55454, USA

[5] Department of Chemistry, University of Minnesota, Minneapolis, Minnesota 55455, USA

[6] Department of Biochemistry, Molecular Biology, and Biophysics, University of Minnesota, Minneapolis, Minnesota 55454, USA

[7] Department of Molecular Pharmacology and Therapeutics, University of Minnesota, Minneapolis, Minnesota 55454, USA

[8] Cornell Institute of Host-Microbe Interactions and Disease, Cornell University, Ithaca NY 14853, USA

corresponding authors: tdoerr@cornell.edu, joseph.boll@uta.edu

*contributed equally

Abstract

Antibiotic tolerance is an understudied potential contributor to antibiotic treatment failure and the emergence of multidrug-resistant bacteria. The molecular mechanisms governing tolerance remain poorly understood. A prominent type of β-lactam tolerance relies on the formation of cell wall-deficient spheroplasts, which maintain structural integrity via their

outer membrane (OM), an asymmetric lipid bilayer consisting of phospholipids on the

inner leaflet and a lipid-linked polysaccharide (lipopolysaccharide, LPS) enriched in the

outer monolayer on the cell surface. How a membrane structure like LPS, with its reliance

on mere electrostatic interactions to maintain stability, is capable of countering internal

turgor pressure is unknown. Here, we have uncovered a novel role for the PhoPQ two-

component system in tolerance to the β-lactam antibiotic meropenem in

Enterobacterales. We found that PhoPQ is induced by meropenem treatment and

promotes an increase in 4-amino-4-deoxy-L-aminoarabinose [L-Ara4N] modification of

lipid A, the membrane anchor of LPS. L-Ara4N modifications enhance structural integrity,

and consequently tolerance to meropenem, in several Enterobacterales species.

Importantly, mutational inactivation of the negative PhoPQ regulatory element *mgrB*

(commonly selected for during clinical therapy with the last-resort antibiotic colistin, an

antimicrobial peptide [AMP]) results in dramatically enhanced tolerance, suggesting that

AMPs can collaterally select for meropenem tolerance via stable overactivation of PhoPQ.

Lastly, we identify histidine kinase inhibitors (including an FDA-approved drug) that inhibit

PhoPQ-dependent LPS modifications and consequently potentiate meropenem to

enhance killing of tolerant cells. In summary, our results suggest that PhoPQ-mediated

LPS modifications play a significant role in stabilizing the OM, promoting survival when

the primary integrity maintenance structure, the cell wall, is removed.

Significance

Treating an infection with an antibiotic often fails, resulting in a tremendous public health

burden. One understudied likely reason for treatment failure is the development of

"antibiotic tolerance", the ability of bacteria to survive normally lethal exposure to an

antibiotic. Here, we describe a molecular mechanism promoting tolerance. A bacterial

stress sensor (PhoPQ) is activated in response to antibiotic (meropenem) treatment and

consequently strengthens a bacterial protective "shell" to enhance survival. We also

identify inhibitors of this mechanism, opening the door to developing compounds that help

antibiotics work better against tolerant bacteria.

Introduction

The rapid rise of antibiotic treatment failure threatens our ability to prevent and

control bacterial infections. Antibiotic resistance, the continued proliferation of bacteria in

the presence of the antibiotic, can often explain failure of clinical therapy. However, the

response to an antibiotic is oftentimes more nuanced than a simple dichotomy of

resistance vs. susceptibility. Bacteria can survive treatment in a non- or slowly-

proliferating state, readily reverting to healthy growth after removal of the antibiotic (such

as the end of a treatment course), and this is typically referred to as "antibiotic tolerance"

(1-3). Importantly, tolerance to antibotics has been shown to enhance the evolution of

outright resistance mechanisms (4-6), and can thus serve as both a direct and indirect

contributor to treatment failure.

β-lactams are the most widely prescribed antibiotic class used to treat bacterial

infections. The β-lactam ring inhibits the activity of the penicillin-binding proteins (PBPs)

through covalent modification of a catalytic residue. PBPs are enzymes that synthesize

the cell wall, an essential structure composed mainly of the polysaccharide peptidoglycan

(PG). In many well-studied model organisms, PBP inhibition induces cell wall degradation

and often subsequent lysis through the action of cell-wall degrading enzymes (collectively

referred to as "autolysins") in a poorly-understood manner (2). While lysis is the canonical

response of model organisms like *Escherichia coli* K12, many susceptible clinical isolates

of Gram-negative pathogens (including prominent Enterobacterales clinical isolates like

Klebsiella spp. and *Enterobacter spp.*) exhibit a unique type of β-lactam tolerance. Like

E. coli, these cells digest their PG upon exposure to β-lactams. However, instead of

lysing, these pathogens survive antibiotic-induced cell wall degradation by forming viable,

non-dividing, cell wall-deficient spheroplasts, which presumably rely on the outer

membrane to counter their internal turgor (7-9). Interestingly, spheroplasts do not

absolutely require osmotic stabilization and form in diverse types of growth media,

including human serum (9). This cell wall-deficient phenotype is reminiscent of so-called

L-forms (10-12), with the notable distinction that spheroplasts do not divide in the

presence of the antibiotic.

Remarkably, spheroplasts formed in response to the carbapenem antibiotic

meropenem readily resume growth and revert to wild-type rod shape when the β-lactam

is removed from the growth medium (7, 9). Little is known about the molecular

mechanisms that facilitate spheroplast formation and survival. In *Vibrio cholerae*, the two-

component system (TCS) VxrAB is essential for spheroplast recovery by upregulating cell

wall synthesis and downregulating iron uptake into the cells, mitigating toxic free iron

levels induced by β-lactam treatment and allowing the cell to avoid damage by oxidative

stress (13, 14). Many questions remain, however, as to how the cell envelope maintains

its integrity without a cell wall, the essential structure canonically thought to protect the

cell against immense turgor pressure.

In this study, we investigated genetic factors that contribute to the ability of

Enterobacter cloacae to tolerate meropenem, which is used as a last-resort β-lactam to

treat multidrug resistant bacterial infections (15-17). We first show that tolerance is

dependent on outer membrane modifications (specifically 4-amino-4-deoxy-L-

aminoarabinose [L-Ara4N]) induced by the PhoPQ TCS, an important cell envelope stress

sensor that has previously been shown to respond to magnesium limitation, cationic

antimicrobial peptide exposure, osmotic challenge, and pH changes (18). Both PhoPQ

regulon transcription and resulting lipid A modifications are induced by meropenem

treatment, suggesting a specific response to perturbations of PG synthesis in *E. cloacae*.

These findings represent a novel mechanism of β-lactam tolerance in clinically relevant

Enterobacterales, as well as an expanded role for the PhoPQ TCS.

Results

The PhoPQ TCS system regulates carbapenem tolerance

We previously showed that many Gram-negative pathogens are highly tolerant to

meropenem. Upon treatment, tolerant cells do not appreciably lyse. Instead, they form

viable, enlarged, non-replicating spheroplasts that are devoid of detectable cell wall

material (9). Meropenem-induced spheroplast formation is quantifiable as an OD_{600}

increase (**Fig. 1A**) and concomitant with only a moderate decrease in survival, as

measured by colony-forming units (CFU) (**Fig. 1B**). In contrast, non-tolerant bacteria like

many *E. coli* isolates rapidly lyse in the presence of meropenem, indicated by a decrease

in both OD_{600} and survival (**Fig. 1AB**). Since spheroplast integrity is presumably

maintained by the outer membrane, rather than the cell wall, we hypothesized that the

strength of the outer membrane might correlate with tolerance. To test this, we repeated

the killing experiments in the presence of the known outer membrane fortifying agents

Mg^{2+} and Ca^{2+}, which link adjacent lipopolysaccharide molecules by forming ionic bridges

between phosphate groups on the lipid A domain (19, 20). Addition of either divalent

cation (Mg^{2+} and Ca^{2+}) supported spheroplast formation with a concomitant reduction in

lysis during meropenem treatment in a concentration-dependent manner (**Fig. S1AB**).

Furthermore, combinatorial addition of excess Ca^{2+} and Mg^{2+} completely prevented lysis

(**Fig. S1A**). Thus, divalent cations prevent lysis during meropenem treatment, potentially

through increasing the mechanical load-bearing capacity of the outer membrane through

LPS crosslinking to protect the spheroplast structure.

When divalent cations are limiting, many Gram-negative pathogens induce outer

membrane modifications (hyperacylation and/or increasing the positive charge of lipid A)

to functionally substitute for divalent ionic bridges between lipopolysaccharide molecules

(21). In *E. cloacae* (22) and other Enterobacterales (23), the PhoPQ TCS directly

regulates expression of the *arn* operon, which synthesizes and transfers positively

charged L-Ara4N moieties to lipid A, and *pagP*, which encodes an outer membrane

acyltransferase (24). To test whether the PhoPQ TCS contributes to meropenem

tolerance, we measured spheroplast formation in Δ*phoPQ.* Strikingly, OD_{600} declined

sharply in Δ*phoPQ* relative to wild type, which could be fully complemented by ectopic

expression of *phoPQ* (**Fig. 2A**). The decline in spheroplast formation strongly correlated

with a robust 10-fold decrease in Δ*phoPQ* viability (CFU/mL) relative to wild type after 24

hours of treatment (**Fig. 2B**). Step-wise titration of Ca^{2+} and/or Mg^{2+} markedly enhanced

Δ*phoPQ* tolerance (**Fig. S1CD**), suggesting decreased tolerance (spheroplast formation)

in $\Delta phoPQ$ is due to its inability to crosslink adjacent LPS molecules in the outer

membrane. To corroborate the involvement of PhoPQ, we sought to either enhance or

reduce its activity and measured the effect of such perturbations on tolerance. PhoQ is

antagonized by the small periplasmic MgrB protein; $mgrB$ overexpression is thus

expected to result in suppression of PhoPQ induction (25). Indeed, $mgrB$ overexpression

from a plasmid reduced spheroplast formation (proxied by OD_{600} measurements) (**Fig.**

2C) in wild type, closely resembling the $\Delta phoPQ$ phenotype (**Fig. 2A**). Conversely,

exposing a strain deleted in $mgrB$ to meropenem resulted in an increase in OD_{600} (**Fig.**

S2A) that coincided with an approximately 1000-fold increased survival to meropenem

relative to wild type (**Fig. 2B**). Modified lipid A structures in $\Delta mgrB$ were confirmed using

matrix-assisted laser desorption ionization-time of flight mass spectrometry (MALDI-TOF

MS) (**Fig. S3AB**). We also phenotypically validated our $\Delta mgrB$ mutant by assessing

colistin MIC, which, as expected, was higher (≥128 μg/mL) than wild type (16 μg/mL).

Collectively, our data suggest that the intensity of the PhoPQ response positively

correlates with meropenem tolerance.

To dissect individual contributions of PhoPQ-regulated genes to tolerance, we

created mutants in the $pagP$ and arn loci. While $\Delta pagP$ displayed spheroplast formation

levels similar to wild type (**Fig. 2D**), $\Delta arnT$ exhibited a drop in OD_{600} reminiscent of

$\Delta phoPQ$ (**Fig. 2D**) and a concomitant 100-fold decrease in CFU/mL relative to wild type

(**Fig. 2B)**. Notably, the survival defect was consistently more pronounced in $\Delta arnT$ vs.

$\Delta phoPQ$ (**Fig. 2B**), suggesting either that residual L-Ara4N modification is retained in the

absence of PhoPQ through basal expression of $arnT$, or that $phoPQ$ induction in the

absence of $arnT$ is detrimental for an unknown reason. However, we observed notable

variation in the extent of *arnT* tolerance measured by OD_{600}; in some experiments,

tolerance was reduced slightly compared to WT (**Fig. S2B**), while in others, the mutant

approached Δ*phoPQ* levels (**Fig. 2D)**. Thus, L-Ara4N modification appears to be the

principal, but not the only, PhoPQ-regulated tolerance factor. Variability might reflect

stochastic variation in another tolerance factor that at least partially mitigates the effects

of *arnT* deletion.

Meropenem induces *arn* transcription in a PhoPQ-dependent manner

Expression of the *arn* operon in *E. cloacae* is directly regulated by phosphorylated

PhoP (22). Because we observed that PhoPQ was necessary for *arn*-mediated tolerance,

we asked whether meropenem induced *arn* transcription in a PhoPQ-dependent manner.

To test this, *E. cloacae* cells were exposed to meropenem for 30 minutes, after which

arnB, *pagP*, and *phoP* transcript levels were quantified (**Fig. 3A**). Relative expression

was calculated using 16s rRNA as an internal control. *phoP*, which is autoregulated (26),

showed 2.5-fold higher expression in meropenem-treated cells relative to untreated.

Additionally, *pagP* expression was 5-fold higher, and *arnB* expression was 6-fold higher

after meropenem treatment. These results support a model where PhoPQ signaling, as

well as transcription of its regulon, is induced in response to meropenem treatment.

To corroborate the qRT PCR findings, we also constructed a fluorescent

transcriptional reporter, fusing the *arnB* promoter (the first gene in the PhoPQ-regulated

arn operon (22, 27)) with msfGFP, followed by fluorescence measurements upon

exposure to meropenem (**Fig. 3B**). As a control, we first exposed cells to EDTA, which

chelates divalent cations to destabilize the outer membrane and consequently activates

PhoPQ (28). As expected, P_{arnB}:msfGFP was induced by EDTA treatment in a *phoPQ*-

dependent way (**Fig. 3B**). Interestingly, meropenem treatment also robustly activated the

P_{arnB}:msfGFP reporter, where a significant 3-fold fluorescence increase was measured,

comparable to EDTA treatment. In contrast, meropenem only slightly (but reproducibly)

increased *arnB* expression in the Δ*phoPQ* mutant under the same conditions. Thus, *arn*

transcription is induced by meropenem in a primarily PhoPQ-dependent manner.

Meropenem treatment promotes formation of* L-Ara4N-*modified lipid A species in

a PhoPQ-dependent manner

The LPS lipid A domain is modified with L-Ara4N in a PhoPQ-dependent manner

when Mg^{2+} is limiting (22), presumably to fortify the outer membrane during magnesium

limitation stress. To determine if the *E. cloacae* lipid A structure is modified with L-Ara4N

in response to meropenem treatment, we isolated lipid A from treated and untreated

cultures. A distinct shift in lipid A structures was evident following 3 hours of meropenem

treatment (**Fig. 4AB**), where increased ratios of L-Ara4N modified vs. unmodified forms

were produced. Notably, the lipid A species that dominated before treatment (hexa-

acylated, bis-phosphorylated, m/z = 1825.25) decreased in abundance in favor of Arn-

and PagP-modified lipid A. Doubly Arn/PagP-modified lipid A (m/z = 2114.14) was also

produced following treatment. Notably, while L-Ara4N modification of lipid A was PhoPQ-

dependent, PagP-dependent lipid A acylation was not (see below for discussion).

Quantitative thin-layer chromatography supported the MS results and revealed a 12.21

(+/- 1.13)-fold increase in single-modified L-Ara4-N lipid A in a PhoPQ-dependent manner

(**Fig. 4C**).

E. cloacae encodes a principal putative PagP acyltransferase that we denoted as

PagP1 (Ecl_03072) and PagP2 (Ecl_04468), a second apparent paralog. Lipid A

extracted from Δ*pagP1* lacked the acyl chain induced upon meropenem treatment (**Fig.**

S4A), suggesting that PagP1 removes palmitate ($C_{16:0}$) from surface-exposed

glycerophospholipids and transfers it to lipid A, as previously shown (29, 30), and

demonstrating that PagP2 does not contribute to hyperacylation under these conditions.

We also confirmed that meropenem-induced L-Ara4N modification was dependent on the

arn operon (**Fig. S4B**). Furthermore, MS analysis of Δ*phoPQ* Δ*pagP1* (**Fig. S4C**) and

Δ*arn* Δ*pagP1* (**Fig. S4D**) lipid A revealed that the mutants produced lipid A structures

lacking all modifications following meropenem treatment, confirming that PagP1 and the

arn operon products coordinate *E. cloacae* lipid A modifications in response to

meropenem treatment. Interestingly, meropenem-induced hyperacylation was absent in

Δ*mgrB* cells (**Fig. S3A**). We propose that meropenem-induced hyperacylation might

actually be reflective of enhanced outer membrane glycerophospholipid bilayer

accumulation in spheroplasts, a condition known for its likely ability to post-translationally

activate PagP1 enzymatic activity (28, 31). This model is also consistent with our

observation that while PagP1 is under genetic control of PhoPQ (22), meropenem-

induced hyperacylation is independent of PhoPQ (**Fig. 4AC**). The absence of PagP1-

dependent modification in Δ*mgrB* might indicate increased outer membrane strength (and

concomitant reduction in glycerophospholipid in the outer leaflet of the outer membrane)

in this background.

Based on the structural studies and our genetic evidence, we suggest PhoPQ-

dependent tolerance is primarily mediated via L-Ara4N addition to lipid A. Presumably, L-

Ara4N lipid A modification increases the structural integrity of the outer membrane

through stabilization of lateral lipopolysaccharide interactions, which protects

spheroplasts from internal turgor.

Colistin exposure primes* E. cloacae *for meropenem tolerance

Cationic antimicrobial peptides (CAMPs) are known inducers of the PhoPQ TCS

(32). Since our data above suggest that PhoPQ induction promotes tolerance, we

hypothesized that pre-exposure to the CAMP colistin "primes" *E. cloacae* for tolerance to

meropenem, either by inducing PhoPQ or by selecting for cells that have a higher baseline

level of PhoPQ induction. To test this, we measured the extent to which *E. cloacae* was

killed by meropenem with and without prior growth in colistin-containing media.

Interestingly, after pre-exposure to colistin, the fraction of cells surviving meropenem

treatment was approximately 10-fold greater **(Fig. 5)**. This suggests that CAMPs have the

potential to induce tolerance to other antibiotics, but that the temporal conditions of

treatment may determine the extent to which this effect is significant.

Outer membrane modifications are conserved β-lactam tolerance determinants in

other Enterobacterales

We next sought to establish whether outer membrane modifications might promote

tolerance in other Enterobacterales. We first turned to *Klebsiella pneumoniae* and used

transposon insertion mutants in *phoP* and *phoQ* from the Manoil laboratory's arrayed

transposon mutant library (33) in killing experiments. After 6 hours of meropenem

exposure, both the *phoQ*::Tn and *phoP*::Tn mutants exhibited 20- to 50-fold lower survival

compared to the wild-type (**Fig. 6A**). We next used an *E. coli* K12 variant (WD101)

engineered to constitutively upregulate the PmrAB two-component system (34) (which

regulates L-Ara4N modification of lipid A in *E. coli*) to test the hypothesis that outer

membrane modifications increase tolerance in *E. coli* (**Fig. 6B**). WD101 exhibited a

dramatic, 10,000-fold increase in survival after 24 hours of meropenem exposure

compared to the wild-type parental strain, further supporting a role for outer membrane

modifications in meropenem tolerance.

A small molecule inhibitor of PhoQ synergizes with meropenem and colistin in vitro

Tolerance is likely an under-appreciated contributor to antibiotic treatment failure.

Antibiotic adjuvants that promote killing of tolerant cells thus have the potential to find a

prominent place in our antibiotic armamentarium. Since histidine kinases like PhoQ are

in principle targetable by small molecules, we tested whether his-kinase inhibitors

synergized with meropenem. To this end, we turned to a previously developed suite of

small molecules with potent histidine kinase inhibitory activity (35) and tested them in

combination with meropenem. The anti-Amyotrophic Lateral Sclerosis (ALS) drug

Riluzole, as well as its derivative Rilu-2, exhibited potent, concentration-dependent

synergy in lysing tolerant *E. cloacae* cells *in vitro* (**Fig. 7A, Fig. S5A**). We also verified

that Rilu-2 inhibited the formation of PhoPQ-dependent lipid A modifications using

MALDI-TOF MS (**Fig. 7B**). Since the PhoPQ system is primarily recognized for its

contribution to CAMP resistance in many Enterobacterales, we next tested the Rilu

compounds' ability to synergize with colistin. As expected, Rilu-2 and Riluzole indeed

potentiated colistin-mediated killing (**Fig. S5B**), lending additional support to a PhoQ-

inhibitory role of these compounds and also confirming previous results in *Salmonella*

(36). Importantly, Riluzole is an FDA-approved treatment for ALS and could thus readily

serve as an adjuvant against both meropenem-tolerant and colistin-resistant

Enterobacterales.

284   **Discussion**

While much work has been done in Enterobacterales to elucidate mechanisms of

antibiotic resistance and persistence, the genetic and molecular determinants of

tolerance, and especially spheroplast formation, have remained poorly understood. In

contrast to resistance (continued growth) and persistence (dormancy), carbapenem-

tolerant populations are initially susceptible to treatment (i.e., lose their cell wall). This

phenotype is reminiscent of so-called "L-forms" (10), with the notable difference that

spheroplasts do not replicate in this state, while L-forms do. This is likely a consequence

of L-forms being able to "escape" the outer membrane to then proliferate through

stochastic membrane blebs (37, 38).

The remarkable ability of spheroplasts to survive without their PG layer lends support to

the recent realization that the outer membrane has load-bearing capabilities (39) and

prompted us to interrogate the molecular mechanism of outer membrane stabilization

during antibiotic exposure. Our data suggest that L-Ara4N addition to lipid A is a key factor

in spheroplast integrity. We propose that modifying lipid A molecules with a positive

charge might increase outer membrane stability due to the enhancement of electrostatic

interactions between adjacent LPS molecules on the surface-exposed face of the outer

membrane. Of note, the L-Ara4N modification has been previously implicated in

resistance to CAMPs, such as colistin (22). In this context, it is worrisome that colistin

therapy can select for *mgrB* mutations (40-42), which we demonstrate here also confers

high meropenem tolerance in addition to colistin resistance. Thus, treatment with

antimicrobial peptide analogs (and, although we did not test this directly, potentially innate

AMPs) can select for bacteria that are stably tolerant to subsequent therapy with a β-

lactam. Our data also suggest that colistin (and, by extension, potentially AMPs of the

innate immune system) can induce transient β-lactam tolerance, likely through induction

of the PhoPQ system and perhaps other protective stress responses. However, the

relationship between PhoPQ induction and tolerance is not absolute; a significant

proportion of both colistin-pretreated and *mgrB*-deleted cells still lyse in the presence of

meropenem. We speculate that β-lactam tolerance and colistin resistance, while relying

on the same basic outer membrane modification, may each require a specific fraction of

LPS molecules to be modified. Thus, within a sample, there may be a limited subset of

cells exhibiting the correct amount of modification to enable both colistin resistance and

meropenem tolerance (or even just optimal meropenem tolerance). This would explain

why we observe only partial overlap between these two phenomena.

In summary, this work demonstrates a novel genetic determinant of carbapenem

tolerance in clinically relevant Enterobacterales. Despite being a well-known regulator of

polymyxin resistance, the PhoPQ two-component system was not previously known to

respond or mediate tolerance to carbapenem treatment. As tolerance (and spheroplast

formation in particular) is a possible culprit for antibiotic treatment failure (2, 3, 43), our

results suggest a potential for combination therapies with histidine kinase inhibitors to

increase the efficacy of carbapenems.

Acknowledgements

329 Research on tolerance in the Dörr lab is supported by National Institutes of Health

330 (NIH/NIAID) grant R01AI143704. Research in the Boll lab is supported by National

331 Institutes of Health (NIH/NIGMS) grant GM143053. Histidine kinase inhibitor research in

332 the Carlson lab supported by National Institutes of Health (NIH/NIGMS) grant GM134538-

333 01A1 and the UMN Office of Academic Clinical Affairs.

Materials and Methods

Bacterial strains and growth

337 All strains/plasmids and primers used in this study are listed in **Table S1 and S2**. All

338 strains were initially grown from freezer stocks on solid agar at 37°C. Isolated colonies

339 were used to inoculate Luria-Bertani (LB), Brain heart infusion (BHI) or N-minimal medium

340 at 37°C. Where required, kanamycin was used at 50 μg/mL, meropenem and colistin were

341 used at 10 μg/mL, unless noted otherwise.

Meropenem killing experiments

Unless noted otherwise, killing experiments were conducted in 100-well honeycomb

plates in a Bioscreen C growth curve analyzer (Growth Curves USA, Piscataway NJ).

Overnight cultures were diluted 10-fold into fresh LB medium containing meropenem (10

µg/mL, 300x MIC) and transferred to honeycomb plates (200 µL volume/culture). OD_{600}

was measured by plate reader; at indicated timepoints, the experiment was paused and

an aliquot was removed for CFU/mL determination or microscopy. Rilu compounds were

dissolved in DMSO as 50 mM (Rilu-2) or 500 mM (Riluzole) stocks and added directly to

the LB medium containing meropenem at the indicated concentrations.

Colistin MIC experiment

Cultures were grown overnight at 37°C shaking, then diluted 1000-fold into fresh LB.

Subcultures were grown for 1 hour at 37°C shaking before being diluted 1000-fold again

into fresh LB to create a "seed culture". 100 µL of seed culture was subsequently diluted

2-fold into a 96-well plate containing colistin concentrations ranging 0.25 – 128 µg/mL.

Reported values are medians of 4 technical replicates.

qRT-PCR

Relative-abundance quantitative PCR (qPCR) was performed as previously described

(44, 45). In brief, the Sybr Fast One-Step qRT-PCR kit (Kapa Biosystems) was used with

16S rDNA as the internal reference. The PCR reaction was performed with Applied

Systems RNA-Ct one-step system. Relative expression levels were calculated using the

ΔΔCt method (46), with normalization of gene targets to16S rDNA signals.

Flow cytometry GFP measurements

Cultures of strains harboring transcriptional P_{arnB}:msfGFP fusions were grown overnight

in LB supplemented with 10 mM $MgSO_4$. Overnight cultures were then washed 2x in fresh

LB before 10-fold dilution into fresh LB medium containing $MgSO_4$ (10 mM),

Ethylenediaminetetraacetic acid (1 mM), or meropenem (10 µg/mL). Cultures were

incubated statically for 3 hours at 37°C. Then, 500 µL of culture was harvested and run

through a C6 Accuri flow cytometer (BD Biosciences) until 100,000 events (cells) had

been analyzed. Mean green fluorescence as measured by the FL1-A channel was used

as a readout for GFP.

Mutant construction

E. cloacae subsp. *cloacae* 13047 mutant strains (*phoPQ*, *arn* and *arnT*) were constructed

as previously described using recombineering with the plasmid pKOBEG (22, 47). Briefly,

linear PCR products were amplified from pKD3 and transformed into *E. cloacae* ATCC

13047/pKOBEG strain by electroporation and plated on chloramphenicol selective media.

Selected clones were transformed with pCP20 to cure the antibiotic resistance cassette.

All mutants were verified by PCR.

pagP1 was deleted using the Wanner method as described previously (22). Briefly, the

chl resistance cassette was amplified from pKD3 using primers TDP1532/TDP1533,

which contain 75 bp flanking homology overhangs. The resulting PCR product was

electroporated into *E. cloacae* ATCC13047 expressing lambda red recombinase from

pACBSR-hyg (48) (a hygromycin-resistant derivative of pKD46 (49)). Mutants were

selected on chloramphenicol (100 µg/mL) and verified by PCR.

Other mutants were constructed using either lambda red recombinase (49) or the suicide

vector pTox (50). The *mgrB* gene was deleted using the suicide plasmid pTox5 as

described in (50). ~700 bp upstream and downstream flanking homology regions were

amplified from ATCC13047 using primers TDP1767/68 and TDP1769/70, and cloned into

pTox5 (digested with EcoRV) using isothermal assembly (51). Successful pTox5Δ*mgrB*

were conjugated into ATCC13047 using the *E. coli* donor strain MFD lambda *pir*;

successful recombinants were selected on plates containing 100 µg/mL chloramphenicol.

Upon single colony purification, colonies were directly streaked out on an M9 minimal

medium plate containing 0.2 % casamino acids and 1 % rhamnose, followed by

incubation at 30 ˚C for 24 – 36 hours. Mutants were tested using primers TDP1771/72.

Lipid A isolation and mass spectrometry

Isolation of lipid A for analysis was performed as previously described (52) with

slight modifications. To analyze lipid A after meropenem treatment, overnight cultures

grown in BHI broth were diluted 1:10 in pre-warmed media with or without meropenem

statically for 3 h. To assess Rilu-dependent modification of lipid A, 12.5 mL of *E. cloacae*

was grown to OD_{600} 1.0. Rilu-2 was used at a final concentration of 200 µM. Bacteria

were harvested and lipid A extraction was carried out by mild-acid hydrolysis as

previously described (53). For mass spectrometry (MS), data were collected on a MALDI-

TOF (Axima Confidence, Shimadzu) mass spectrometer in the negative mode, as

previously done (22).

For quantification of lipid A, cultures were grown with 2.5 µCi/mL of ^{32}P ortho-

phosphoric acid (^{32}P) (Perkin Elmer) and lipid A was extracted. Thin layer

chromotagraphy was done in a pyridine, chloroform, 88% formic acid, aqueous

(50:50:16:5 v/v) tank for 3 hours. Plates were exposed to a phosphor screen, imaged,

and densitometry was used to calculate the percentage of each lipid species. Reported

densitometry was calculated using 2 replicates +/- standard deviation.

Figure Legends

Figure 1: *Enterobacter cloacae* is highly meropenem tolerant. (A) Representative

experiment demonstrating changes in OD_{600} measurements following meropenem

treatment in *E. cloacae* relative to *E. coli.* Error bars represent the average of 3 technical

replicates +/- standard deviation. (B) Survival was calculated as CFU/mL from the experiment depicted in (A). The red line denotes the limit of detection. Phase images from the same experiment show cells before and after meropenem exposure to illustrate spheroplast formation in *E. cloacae* and lysis in *E. coli* (only cell debris is visible in phase image)*.* Scale bars, 2 μm.

Figure 2: The PhoPQ system promotes meropenem tolerance in *E. cloacae.*

(A) Spheroplast formation in response to meropenem treatment. Overnight cultures were diluted 10-fold into fresh LB medium containing 10 μg/mL meropenem and OD_{600} was

measured. (B) Fraction of population surviving after 24 hours of meropenem exposure

from experiments as described in (A). (C) MgrB overexpression reduces tolerance. Cells

were treated as described in (A), but with the addition of 0.2 % arabinose (inducer).

pHERD; empty vector. (D) The *arn* operon is required for meropenem tolerance.

Experiments were conducted as described in (A). Data in each graph represent the

average of 3 replicates +/- standard deviation; additional biological replicates are shown

in Fig. S2B. Statistical significance for survival fraction determined by one-way ANOVA

of log transformed data, followed by Tukey's correction for multiple comparisons (ns, not

significant; ****, $p \leq 0.0001$).

Figure 3: Expression of *arn*B in response to meropenem treatment is dependent on PhoPQ. (A) Relative expression reverse-transcription quantitative PCR (qRT-PCR) of PhoPQ regulon transcripts after meropenem exposure. Each experiment was independently replicated three times (individual data points of the three experiments are

reported here). (B) Strains carrying transcriptional P_{arnB}:msGFP fusions were exposed to

indicated conditions and analyzed on a C6 Accuri flow cytometer. Statistical difference

between populations was determined with a one-way ANOVA followed by Tukey's

correction for multiple comparisons (ns, not significant; ****, $p \leq 0.0001$).

Figure 4: Analysis of *E. cloacae* lipid A after meropenem treatment. (A) MALDI-TOF

MS analysis of lipid A extracted from wild-type or Δ*phoPQ E. cloacae* strains. L-Ara4N

modifications are illustrated in red, while $C_{16:0}$ additions are green. Numbered labels that

are both red and green contain both modifications. Each experiment was independently

replicated three times, and one representative data set was reported. (B) Predicted lipid

A chemical structures in wild type and Δ*phoPQ E. cloacae.* (C) [32]P-radiolabelled lipid A

was extracted from treated or untreated wild-type or Δ*phoPQ E. cloacae.* Lipids were

separated based on hydrophobicity using thin-layer chromatography. Red circle denotes

L-Ara4N modification, while the green line indicates $C_{16:0}$ addition.

Figure 5: Colistin primes *E. cloacae* for meropenem tolerance. Overnight cultures

were grown in LB +/- 20 µg/mL colistin, then diluted 10-fold into LB +/- 20 µg/mL colistin.

Colistin pretreatment value represents total time exposed to colistin before exposure to

10 µg/mL meropenem. Cultures were then treated like other meropenem killing

experiments (see methods). Survival fraction was calculated by dividing CFU/mL at 3h

meropenem exposure by CFU/mL prior to meropenem exposure. Each bar represents

the mean of 9 biological replicates, error bars represent standard deviation. Statistical

significance determined by one-way ANOVA of log transformed data, followed by Tukey's

correction for multiple comparisons (ns, not significant; *, $p \leq 0.05$; **, $p \leq 0.01$).

Figure 6: A conserved mechanism for meropenem tolerance in Enterobacterales.

(A) *Klebsiella pneumoniae* KPNIH1 and its *phoP*::Tn and *phoQ*::Tn derivatives were

diluted 10-fold into LB medium containing 10 μg/mL meropenem. Survival fraction is the

CFU/mL after 6 hours of meropenem treatment divided by CFU/mL before treatment.

Data represent averages of 6 biological replicates +/- standard error (B) *E. coli* strain

W3110 (WT) or strain WD101, which has a constitutively active chromosomal copy of

pmrA (*pmrA^C*), were cultured in N-minimal medium and treated with or without

meropenem. Cultures were incubated statically at 37°C. CFU were enumerated at 0, 3,

6, 12 and 24 hours. Error bars indicate standard deviation. Each experiment was

independently replicated three times in triplicate, and one representative data set was

reported.

Figure 7: Rilu compounds synergize with meropenem to expedite *E. cloacae* killing.

(A) Rilu compounds potentiate meropenem-induced lysis against *E. cloacae.* Overnight

cultures were diluted 10-fold into fresh growth medium containing meropenem (10 µg/mL)

and increasing concentrations of Riluzole or its derivative Rilu-2. Data represent the

average of 3 technical replicates +/- standard deviation. (B) MALDI-MS analysis of lipid A

isolated from untreated wild-type *E. cloacae*, cells treated with RILU-2 or Δ*phoPQ*. *m/z*

corresponding with L-Ara4N modifications are illustrated in red. Relevant lipid A chemical

structures are shown. Each experiment was independently replicated three times, and

one representative data set is reported.

Figure S1: Addition of divalent cations prevents spheroplast lysis. Wild type (WT)

(A-B) or its $\Delta phoPQ$ derivative (C-D) were treated as described in Fig. 1A with addition of

the indicated concentrations of (A,C) $MgSO_4$ (Mg^{2+}) or (B,D) $CaCl_2$ (Ca^{2+}). Data represent

the average of 3 replicates +/- standard deviation.

Figure S2: Independent biological replicates of experiments shown in Fig. 1. (A) An

mgrB mutation promotes a moderate increase in mass increase during meropenem

exposure. (B) Experiments were conducted as described in Fig. 1A legend; each graph

represents experiments conducted on a different day. In addition, data in each graph

represent the average of 3 biological replicates +/- standard deviation.

Figure S3: Analysis of lipid A from Δ*mgrB*. (A) MALDI-MS analysis of lipid A isolated

from *E. cloacae* Δ*mgrB*. *m/z* corresponding with L-Ara4N modifications are illustrated in

red. Each experiment was independently replicated three times, and one representative

data set was reported. (B) Relevant lipid A chemical structures are shown.

Figure S4: Analysis of lipid A from *E. cloacae* mutants. (A) MALDI-MS analysis of

lipid A isolated from Δ*pagP1,* (B) Δ*arn*, (C) Δ*phoPQ* Δ*pagP1 and* (D) Δ*arn* Δ*pagP1*. *m/z*

corresponding with L-Ara4N modifications are illustrated in red, while structures with

altered acyl chain patterns are illustrated in green. Each experiment was independently

replicated three times, and one representative data set was reported.

Figure S5: Rilu synergizes with colistin to enhance killing. (A) Rilu compounds do not cause lysis, but (B) potentiate colistin mediate killing. Experiments were conducted as described in Fig. 1A legend. Data represent the average of 3 replicates +/- standard deviation.

1. A. Brauner, O. Fridman, O. Gefen, N. Q. Balaban, Distinguishing between resistance, tolerance and persistence to antibiotic treatment. *Nat Rev Microbiol* **14**, 320-330 (2016).
2. T. Dorr, Understanding tolerance to cell wall-active antibiotics. *Ann N Y Acad Sci* **1496**, 35-58 (2021).
3. L. F. Westblade, J. Errington, T. Dorr, Antibiotic tolerance. *PLoS Pathog* **16**, e1008892 (2020).
4. I. Levin-Reisman, A. Brauner, I. Ronin, N. Q. Balaban, Epistasis between antibiotic tolerance, persistence, and resistance mutations. *Proc Natl Acad Sci U S A* **116**, 14734-14739 (2019).
5. I. Levin-Reisman *et al.*, Antibiotic tolerance facilitates the evolution of resistance. *Science* **355**, 826-830 (2017).
6. J. Liu, O. Gefen, I. Ronin, M. Bar-Meir, N. Q. Balaban, Effect of tolerance on the evolution of antibiotic resistance under drug combinations. *Science* **367**, 200-204 (2020).
7. T. Dorr, B. M. Davis, M. K. Waldor, Endopeptidase-mediated beta lactam tolerance. *PLoS Pathog* **11**, e1004850 (2015).
8. L. G. Monahan *et al.*, Rapid conversion of Pseudomonas aeruginosa to a spherical cell morphotype facilitates tolerance to carbapenems and penicillins but increases

susceptibility to antimicrobial peptides. *Antimicrob Agents Chemother* **58**, 1956-
1962 (2014).

9. T. Cross *et al.*, Spheroplast-Mediated Carbapenem Tolerance in Gram-Negative
Pathogens. *Antimicrob Agents Chemother* **63** (2019).

10. J. Errington, K. Mickiewicz, Y. Kawai, L. J. Wu, L-form bacteria, chronic diseases
and the origins of life. *Philos Trans R Soc Lond B Biol Sci* **371** (2016).

11. R. Mercier, Y. Kawai, J. Errington, General principles for the formation and
proliferation of a wall-free (L-form) state in bacteria. *Elife* **3** (2014).

12. J. Errington, Cell wall-deficient, L-form bacteria in the 21st century: a personal
perspective. *Biochem Soc Trans* **45**, 287-295 (2017).

13. J. H. Shin *et al.*, A multifaceted cellular damage repair and prevention pathway
promotes high-level tolerance to beta-lactam antibiotics. *EMBO Rep* **22**, e51790
(2021).

14. T. Dorr *et al.*, A cell wall damage response mediated by a sensor kinase/response
regulator pair enables beta-lactam tolerance. *Proc Natl Acad Sci U S A* **113**, 404-
409 (2016).

15. D. L. Paterson, Recommendation for treatment of severe infections caused by
Enterobacteriaceae producing extended-spectrum beta-lactamases (ESBLs). *Clin*
Microbiol Infect **6**, 460-463 (2000).

16. J. A. Torres, M. V. Villegas, J. P. Quinn, Current concepts in antibiotic-resistant
Gram-negative bacteria. *Expert Review of Anti-infective Therapy* **5**, 833-843
(2007).

17. D. L. Paterson, R. A. Bonomo, Extended-spectrum beta-lactamases: a clinical
update. *Clin Microbiol Rev* **18**, 657-686 (2005).

18. E. A. Groisman, A. Duprey, J. Choi, How the PhoP/PhoQ System Controls
Virulence and Mg(2+) Homeostasis: Lessons in Signal Transduction,
Pathogenesis, Physiology, and Evolution. *Microbiol Mol Biol Rev* **85**, e0017620
(2021).

19. S. Snyder, D. Kim, T. J. McIntosh, Lipopolysaccharide Bilayer Structure: Effect of
Chemotype, Core Mutations, Divalent Cations, and Temperature. *Biochemistry* **38**,
10758-10767 (1999).

20. P. Garidel *et al.*, Divalent cations affect chain mobility and aggregate structure of
lipopolysaccharide from Salmonella minnesota reflected in a decrease of its
biological activity. *Biochim Biophys Acta* **1715**, 122-131 (2005).

21. J. C. Henderson *et al.*, The Power of Asymmetry: Architecture and Assembly of
the Gram-Negative Outer Membrane Lipid Bilayer. *Annual Review of Microbiology*
70, 255-278 (2016).

22. K. N. Kang *et al.*, Colistin heteroresistance in Enterobacter cloacae is regulated by
PhoPQ-dependent 4-amino-4-deoxy-l-arabinose addition to lipid A. *Mol Microbiol*
111, 1604-1616 (2019).

23. A. Y. Mitrophanov, M. W. Jewett, T. J. Hadley, E. A. Groisman, Evolution and
Dynamics of Regulatory Architectures Controlling Polymyxin B Resistance in
Enteric Bacteria. *PLOS Genetics* **4**, e1000233 (2008).

24. R. E. Bishop, S. H. Kim, A. El Zoeiby, Role of lipid A palmitoylation in bacterial
pathogenesis. *J Endotoxin Res* **11**, 174-180 (2005).

25. A. M. Lippa, M. Goulian, Feedback Inhibition in the PhoQ/PhoP Signaling System
by a Membrane Peptide. *PLOS Genetics* **5**, e1000788 (2009).
26. F. C. Soncini, E. G. Véscovi, E. A. Groisman, Transcriptional autoregulation of the
Salmonella typhimurium phoPQ operon. *Journal of Bacteriology* **177**, 4364-4371
(1995).
27. J. S. Gunn *et al.*, PmrA–PmrB-regulated genes necessary for 4-aminoarabinose
lipid A modification and polymyxin resistance. *Molecular Microbiology* **27**, 1171-
1182 (1998).
28. W. Jia *et al.*, Lipid trafficking controls endotoxin acylation in outer membranes of
Escherichia coli. *J Biol Chem* **279**, 44966-44975 (2004).
29. P. M. Hwang *et al.*, Solution structure and dynamics of the outer membrane
enzyme PagP by NMR. *Proceedings of the National Academy of Sciences* **99**,
13560-13565 (2002).
30. R. E. Bishop *et al.*, Transfer of palmitate from phospholipids to lipid A in outer
membranes of Gram-negative bacteria. *The EMBO Journal* **19**, 5071-5080 (2000).
31. J. C. Malinverni, T. J. Silhavy, An ABC transport system that maintains lipid
asymmetry in the gram-negative outer membrane. *Proc Natl Acad Sci U S A* **106**,
8009-8014 (2009).
32. M. W. Bader *et al.*, Recognition of Antimicrobial Peptides by a Bacterial Sensor
Kinase. *Cell* **122**, 461-472 (2005).
33. B. Ramage *et al.*, Comprehensive Arrayed Transposon Mutant Library of Klebsiella
pneumoniae Outbreak Strain KPNIH1. *Journal of Bacteriology* **199**, e00352-00317
(2017).
34. M. S. Trent *et al.*, Accumulation of a polyisoprene-linked amino sugar in polymyxin-
resistant Salmonella typhimurium and Escherichia coli: structural characterization
and transfer to lipid A in the periplasm. *J Biol Chem* **276**, 43132-43144 (2001).
35. K. E. Wilke, S. Francis, E. E. Carlson, Inactivation of Multiple Bacterial Histidine
Kinases by Targeting the ATP-Binding Domain. *ACS Chemical Biology* **10**, 328-
335 (2015).
36. M. K. Thielen, C. K. Vaneerd, M. Goswami, E. E. Carlson, J. F. May, 2-
Aminobenzothiazoles Inhibit Virulence Gene Expression and Block Polymyxin
Resistance in Salmonella enterica. *Chembiochem* **21**, 3500-3503 (2020).
37. R. Mercier, Y. Kawai, J. Errington, Excess Membrane Synthesis Drives a Primitive
Mode of Cell Proliferation. *Cell* **152**, 997-1007 (2013).
38. G. Billings *et al.*, De novo morphogenesis in L-forms via geometric control of cell
growth. *Molecular Microbiology* **93**, 883-896 (2014).
39. E. R. Rojas *et al.*, The outer membrane is an essential load-bearing element in
Gram-negative bacteria. *Nature* **559**, 617-621 (2018).
40. L. Poirel *et al.*, The mgrB gene as a key target for acquired resistance to colistin in
Klebsiella pneumoniae. *Journal of Antimicrobial Chemotherapy* **70**, 75-80 (2014).
41. A. Cannatelli *et al.*, *In Vivo* Emergence of Colistin Resistance in Klebsiella
pneumoniae Producing KPC-Type Carbapenemases Mediated by Insertional
Inactivation of the PhoQ/PhoP *mgrB* Regulator. *Antimicrobial Agents and*
Chemotherapy **57**, 5521-5526 (2013).

42.   E. Lopez-Camacho *et al.*, Genomic analysis of the emergence and evolution of
642   multidrug resistance during a Klebsiella pneumoniae outbreak including
643   carbapenem and colistin resistance. *J Antimicrob Chemother* **69**, 632-636 (2014).
644   43.   D. Roberts, E. Higgs, A. Rutman, P. Cole, Isolation of spheroplastic forms of
645   Haemophilus influenzae from sputum in conventionally treated chronic bronchial
646   sepsis using selective medium supplemented with N-acetyl-D-glucosamine:
647   possible reservoir for re-emergence of infection. *Br Med J (Clin Res Ed)* **289**, 1409-
648   1412 (1984).
649   44.   B. W. Davies, R. W. Bogard, J. J. Mekalanos, Mapping the regulon of Vibrio
650   cholerae ferric uptake regulator expands its known network of gene regulation.
651   *Proc Natl Acad Sci U S A* **108**, 12467-12472 (2011).
652   45.   J. M. Boll *et al.*, A penicillin-binding protein inhibits selection of colistin-resistant,
653   lipooligosaccharide-deficient *Acinetobacter baumannii*. *Proceedings of the*
654   *National Academy of Sciences* **113**, E6228-E6237 (2016).
655   46.   K. J. Livak, T. D. Schmittgen, Analysis of Relative Gene Expression Data Using
656   Real-Time Quantitative PCR and the 2−ΔΔCT Method. *Methods* **25**, 402-408
657   (2001).
658   47.   F. Guérin, C. Isnard, V. Cattoir, J. C. Giard, Complex Regulation Pathways of
659   AmpC-Mediated β-Lactam Resistance in Enterobacter cloacae Complex.
660   *Antimicrobial Agents and Chemotherapy* **59**, 7753-7761 (2015).
661   48.   T. W. Huang *et al.*, Capsule deletion via a lambda-Red knockout system perturbs
662   biofilm formation and fimbriae expression in Klebsiella pneumoniae MGH 78578.
663   *BMC Res Notes* **7**, 13 (2014).
664   49.   K. A. Datsenko, B. L. Wanner, One-step inactivation of chromosomal genes in
665   Escherichia coli K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**, 6640-
666   6645 (2000).
667   50.   J. E. Lazarus *et al.*, A New Suite of Allelic-Exchange Vectors for the Scarless
668   Modification of Proteobacterial Genomes. *Appl Environ Microbiol* **85** (2019).
669   51.   D. G. Gibson *et al.*, Enzymatic assembly of DNA molecules up to several hundred
670   kilobases. *Nature Methods* **6**, 343-345 (2009).
671   52.   K. N. Kang *et al.*, Septal Class A Penicillin-Binding Protein Activity and ld-
672   Transpeptidases Mediate Selection of Colistin-Resistant Lipooligosaccharide-
673   Deficient Acinetobacter baumannii. *mBio* **12** (2021).
674   53.   Z. Zhou, S. Lin, R. J. Cotter, C. R. Raetz, Lipid A modifications characteristic of
675   Salmonella typhimurium are induced by NH4VO3 in Escherichia coli K12.
676   Detection of 4-amino-4-deoxy-L-arabinose, phosphoethanolamine and palmitate.
677   *J Biol Chem* **274**, 18503-18514 (1999).
678