Simultaneous acoustic monitoring uncovers evidence of biodiversity loss and overlooked temporal variation in a threatened Amazonian bird community

W. Alexander Hopping[1], Holger Klinck[1]

[1] K. Lisa Yang Center for Conservation Bioacoustics, Cornell Lab of Ornithology, Cornell University, Ithaca, NY, USA

Corresponding Author:
W. Alexander Hopping[1]
Email address: wah63@cornell.edu

Abstract

1. Birds are frequently used as indicator taxon in ecology, but traditional surveys are prone to error and largely inadequate in the dense, speciose tropical forests where they are most needed. Passive acoustic monitoring (PAM) has the potential to address many of these pitfalls, providing unique advantages for both long-term monitoring and rapid biodiversity assessment. Here, we evaluate the effectiveness of PAM for characterizing spatiotemporal variation in avian species richness, diversity, vocal activity, and community composition in the Amazon basin global biodiversity hotspot.

2. We employed an array of ten autonomous recording units (ARUs) spanning edge, degraded, and interior forest at a reserve on the deforestation frontier of Madre de Dios, Peru, a region featuring some of the richest, most complex avian assemblages on Earth. Recordings from 21 dawn-hours across three days at seven sites were manually annotated by a single observer.

3. ARUs and manual annotation performed well as an inventory method, especially for midstory and understory species. Sites in edge and degraded habitats featured significantly lower avian species richness, Shannon diversity, and vocal activity levels than in the forest interior, patterns that replicate the findings of prior high-effort avian censuses in the region. We observed significant temporal variation between days; at all featured sites, vocal activity was highest on January 20th and lowest on January 31st.

4. We demonstrated that novel annotation-generated metrics can work as effective proxies for abundance data and per-capita song rate, correlating with diversity indices and efficiently

characterizing habitats at a level of detail unobtainable with human observers. Generalist
species were significantly overrepresented in the soundscapes of interior forest, relative to
their contributed share of species richness, and interior specialists were underrepresented,
suggesting that the entire 191ha site is degraded. The between-days temporal variation that
we documented, almost certainly overlooked without simultaneous monitoring, may
obfuscate or distort the results of traditional surveys.
45
46 **5. Synthesis & Applications.** We propose that PAM should become a cornerstone of
47 biodiversity research. The standardization advantages demonstrated in this study,
48 permanent storage, multi-taxa applications, and potential of automated identification make
49 bioacoustics an ideal methodological avenue for guiding management and policy under
50 rapid global change.
51
52
53 **Keywords:** avian biodiversity, bioacoustics, deforestation and degradation, edge effects,
54 fragmentation, passive acoustic monitoring, Peruvian Amazon, spatiotemporal variation.
55
56

1 Introduction

58
The tropics account for an overwhelming share of Earth's biodiversity, featuring more
than 75% of all species and over 90% of terrestrial birds (Barlow *et al.* 2018). Of particular
importance is the Amazon basin, the largest intact tropical system on the planet (Allan *et al.*
2017) and the epicenter of global biodiversity (Jenkins *et al.* 2013). Like most of the tropics, the
Amazon is undergoing major ecological change (Edelman 2014), primarily from the region's
rapidly expanding anthropogenic land use footprint (Laurence *et al.* 2014), as well as the
increasing impacts of climate change (Lovejoy 2013). Despite their importance, however, the
tropics are largely neglected in biodiversity, and ecosystem function literature (Wilson *et al.*
2016), and assumptions derived from studies in temperate regions have been erroneously used to
characterize tropical systems where they do not apply (Clarke *et al.* 2017). If we are to meet the
international community's sustainable development goals, more biodiversity research focused
specifically on tropical systems is necessary (United Nations 2015).
71
Land use change in Amazonia, primarily the result of agriculture, mining, and expanding
road networks, is the most significant threat to biodiversity in the basin (Laurance *et al.* 2014)
and likely an underestimated driver of climate change (Mahowald *et al.* 2017), compounding the
importance of its study. In Madre de Dios, Peru, these pressures have driven accelerated rates of
deforestation, fragmentation, and degradation over recent decades, making the region a global
deforestation hotspot (Nicolau *et al.* 2019, Sánchez-Cuervo *et al.* 2020), and inducing ecosystem
decay and biodiversity loss in remaining habitat patches (Canaday & Rivadeneyra 2001,

Laurance *et al.* 2002, Haddad *et al.* 2015, Stouffer 2020), but the full impacts of these changes
remain unknown. *Várzea* floodplain forest, the focus of this study, is particularly vulnerable to
these dynamics due to its limited distribution, easy access from river-based transport, and high
concentration of endemic bird species, many of which are restricted to particular successional
stages in this forest type (Foster *et al.* 1994, Robinson & Terborgh 1997). Even in intact
Amazonian systems, long-term studies have recently uncovered baseline shifts and population
declines in avian communities, perhaps due to climate change (Blake & Loiselle 2015, Stouffer
et al. 2020). However, such assessments are complicated by the difficulty of tropical field
surveys, leading to flawed baseline species occurrence data (Boakes *et al.* 2010, Lees *et al.* 2014,
Robinson *et al.* 2018) and overlooked biodiversity losses in Amazonia (Socolar *et al.* 2019),
which interfere with conservation efforts in the region.
90
Birds, which are generally accessible, identifiable, and better understood than other
Neotropical groups, are presently the most cost-effective study taxa for monitoring Amazonian
biodiversity (Gardner *et al.* 2008, Lees *et al.* 2014). Birds are ubiquitous and can be highly
responsive to habitat conditions, making them excellent proxies for understanding ecosystem
health and global change (Remsen 1994, Blackburn & Gaston 1998, Bibby 1999). Despite these
advantages, the Amazon's avifauna remains poorly documented and subject to significant
knowledge gaps (Barlow *et al.* 2011, Lees *et al.* 2014). While high-quality, standardized,
scalable data on birds and other taxonomic groups is necessary for guiding conservation
priorities in the tropics, such data are difficult to generate with traditional survey methods.
Overwhelming avian species richness, including many rare and similar species, challenging
logistics, and poor viewing conditions typical of dark, densely vegetated rainforest interiors (Fig.
1) contribute to the unreliability of avian surveys in Amazonia, where upwards of 95% of birds
are heard but never seen by a field observer (Robinson *et al.* 2018). Amazonian soundscapes
include thousands of vocalization types, inducing high rates of false-negative and false-positive
identification errors (Remsen 1994), and it can take months for skilled observers to reach a
comfort level attainable in just one to two weeks of field experience in temperate systems
(Parker 1991). Additionally, neotropical soundscapes are subject to high levels of spatiotemporal
heterogeneity, with species singing in narrow time windows (Parker 1991, Hart *et al.* 2015), at
seasonally variable rates (Blake 1992), and as a function of fine-grain floristic, geographic, and
spatial variables, which are still poorly understood (Menger *et al.* 2017). As a result, studies
relying on point counts and other non-simultaneous survey methods in Amazonia may require
substantial sample sizes, which are often infeasible to obtain (Esquivel & Peris 2008, Rodriguez
et al. 2014).

114
115 **Figure 1: Site 07, typical *várzea* habitat in the reserve.**
116

Autonomous Recordings Units (ARUs) effectively address many of the shortfalls of
traditional point counts in tropical systems. Simultaneous and continuous monitoring of many
locations has the capacity to completely eliminate temporal bias in surveys (Venier *et al.* 2012,
Tegeler *et al.* 2012), data collection is independent of observer skill level, and permanent records
can be reviewed, distributed, and verified by experts at will, enabling independent verification,
reducing identification errors (Rempel *et al.* 2005, Rempel *et al.* 2019), and serving as ecological
time capsules (Sugai & Llusia 2019). ARUs can generate significantly greater data volume than
traditional methods without increasing field time or cost (Hobson *et al.* 2002, Acevedo &
Villanueva-Rivera 2006, Tegeler *et al.* 2012), and bioacoustic data contains vocalizations of non-
target taxa and other soundscape elements, opening the door for more holistic research (Newson
et al. 2017), while studies focused exclusively on birds can come at the expense of developing
knowledge of other groups (Gardner *et al.* 2008). Human presence in the field during survey
periods, which can bias results, is unnecessary (Jorge *et al.* 2018), and habitat structure
influences bioacoustic surveys less than point counts (Darras *et al.* 2018). Perhaps most
importantly, acoustic data introduces the possibility of automated detection and identification,
which would have a transformative impact on biodiversity monitoring (Kahl *et al.* 2020). ARUs
provide equivalent or superior data to point counts, especially in tropical forests like the
Amazon, where the number of experienced observers is limited, access is difficult, the vast
majority of species are best detected aurally, and sites are highly heterogenous over space and
time (Haselmayer & Quinn 2000, Acevedo & Villanueva-Rivera 2006, Cellis-Murillo *et al.*
2012, Shonfield & Bayne 2017, Darras *et al.* 2018, Darras *et al.* 2019, Pillay *et al.* 2019, Blake
2021).
139

140   Here, we used ARUs to explore spatiotemporal variation in avian species richness,
141 Shannon diversity, vocal activity, and community composition between sites at a threatened
142 Amazonian reserve. Situated 15 km east of the rapidly growing frontier town of Puerto
143 Maldonado and adjacent to an unmarked road, Inkaterra Reserva Amazónica (ITRA), our study
144 site, faces intense land use pressure (Fig. 2). Likely inducing compositional changes and
145 biodiversity loss across the reserve (Canaday and Rivadeneyra 2001, Laurance *et al.* 2002,
146 Stouffer 2020), these dynamics are expected to intensify in coming years (Bird *et al.* 2012),
147 making high-quality, baseline bioacoustic data especially valuable. Additionally, fully annotated
148 Amazonian dawn choruses are critical for training automated soundscape identification
149 algorithms, for which this dataset has already been used (Kahl *et al.* 2020).

151   We asked three overarching questions in our analyses. (1) Performance of acoustic
152 sampling: how well do ARUs document avian community structure within the reserve, and can
153 acoustic data serve as a proxy for abundance data, replicating the results of historic census efforts
154 in Amazonia? (2) Temporal analysis: was there significant temporal variation between days that
155 non-simultaneous survey methods might overlook? (3) Spatial analysis: do differences in habitat
156 type and quality, particularly between edge and interior habitats, significantly impact avian
157 species richness, diversity, community composition, and vocal activity? Can bioacoustic data be
158 used to efficiently characterize habitats, without relying on manual estimates of per-capita song
159 rate?

162 **2 Materials & Methods**

164 **2.1 Site Selection**
165   The data for this study was collected at Inkaterra Reserva Amazónica (ITRA) in the
166 Madre de Dios Department of southeastern Peru (12°32'07.8" S, 69°02'58.2" W). This 191 ha
167 private reserve is located at an elevation of ~200 m along the Madre de Dios River, directly
168 across from the Reserva Nacional de Tambopata, in one of the most biodiverse regions on the
169 planet (Foster *et al.* 1994) (Fig. 2). The habitat at ITRA primarily consists of *várzea* floodplain
170 forest, the most endangered forest type in the southwestern Amazon (Phillips *et al.* 1994),
171 interspersed with seasonally flooded *Mauritia* palm swamp forest.

Figure 2: Swift recorder layout and progression of deforestation at the study site. Recorder placements covered
a range of habitats including edge (Site 01), degraded forest (Site 10), and intact *várzea* in the reserve's interior

2.2 Data Collection

Data were collected January 14-February 2, 2019, using ten ARUs (Swift recorder, K. Lisa Yang Center for Conservation Bioacoustics, Cornell Lab of Ornithology, Ithaca, NY, USA). ARUs were deployed across the entire reserve, at a minimum distance of 350 m from each other and 450 m from the river, to limit overlapping detection (pseudo-replication) and background noise (Ralph *et al.* 1995). Deployment locations were chosen to represent a gradient of intactness and forest habitat type, including edge (Site 01), degraded (Site 10), mature *várzea* in the reserve's interior (Site 04 and Site 06), and mixed palm swamp-*várzea* habitat (Site 02 and Site 05) (FIG. 2). Placed at a height of 1.5 m above the ground, the 10 ARUs recorded continuously (mono, WAV format) throughout the deployment using a sampling rate of 48 kHz (16-bit resolution) and a gain setting of 35 dB. The microphone sensitivity of the Swift was -44 dBV/Pa (+/- 3 dB) and featured a flat frequency response (+/- 3 dB) in the frequency range of the vocalizations of interest. The clipping level of the analog-to-digital (ADC) converter was +/- 0.9 V.

2.3 Acoustic Analysis (Annotation Process)

Twenty-one total dawn-hours, from 05:00-06:00 PET (10:00-11:00 UTC), representing 7 of the 10 sites on three days, were manually annotated by the lead author using the Raven Pro Sound Analysis Software® (version 1.5; K. Lisa Yang Center for Conservation Bioacoustics, Cornell Lab of Ornithology, Ithaca, NY, USA). Six of these sites were used for analyses, two in each habitat type: fringe (edge and degraded), mixed, and *várzea*, with Site 08 excluded. For some analyses, mixed and *várzea* were grouped together as interior habitats. The three days, January 16 (Day A), January 20 (Day B), and January 31 (Day C), 2019, were randomly chosen from a subset of days that featured less than 5 minutes of heavy rain during the dawn-hour. The dawn-hour was selected because it contains the highest level of vocal activity and diversity among neotropical birds (Berg *et al.* 2006), including many species that only sing in narrow time windows around first light (Parker 1991, Blake 1992). Additionally, the dawn soundscape in Amazonia features far less anthropogenic activity and insect noise than other periods, reducing interference with avian species detection (Hart *et al.* 2015, Metcalf *et al.* 2020).

Spectrograms were viewed in 60-second increments split over two rows (Hann window, 3dB bandwidth = 100 Hz, 690 FFT window size, 1024-sample DFT, 95% temporal overlap). To reduce bias introduced by improved sound identification knowledge developed throughout the project, the order in which site days were annotated was determined using a random number generator. Vocalizations were labeled for both species and a handful of broader groups with confusing or indistinguishable calls, e.g., "TRSP" as "trogon species" (*Trogonidae sp.*). Background vocalizations, audible but not clearly visible on a spectrogram, were denoted with a '1' and excluded from most analysis. To account for inter-specific variation in call volume and

frequency, consistency for applying 'background' designation was approached on a species-
specific basis. In cases where vocalizations from the same species were separated by fewer than
5 seconds, they were included as part of the same annotation, while annotations were split if the
gap between vocalizations exceeded 5 seconds. An assortment of unlabeled sound clips,
featuring both identified and unidentified species, were sent to regional experts for secondary
verification. Annotations were reviewed repeatedly until unidentified vocalizations made up less
than 10% of the total avian vocal activity on a given recording. Considerable effort was made to
accurately label as many foreground and background vocalizations as possible, with the goal of
improving the value of this dataset for training soundscape-based automated identification
algorithms.

2.4 Statistical Analyses

For each site, day, and individual recording, four metrics were calculated: species
richness, Shannon diversity, vocal prevalence, and Bray-Curtis dissimilarity. Species richness
and Shannon diversity were included as standard measures of ecological diversity (Okansen *et*
al. 2019, Metcalf *et al.* 2020). Vocal prevalence (VP), the number of 15-second blocks in a
recording containing at least one foreground vocalization from a species or taxa, was used to
quantify vocal activity levels or soundscape abundance. Bray-Curtis dissimilarity was calculated
between days, sites, and individual site days, ranging from 0 (no difference) to 1 (completely
different).

Certain taxa were excluded from most analysis for data quality reasons: parrots
(*Psittacidae sp.*), because they occurred almost exclusively as calling flyovers, along with
Columbiformes (pigeons and doves) and three congeneric tinamou species, Cinereous Tinamou
(*Crypturellus cinereus*), Little Tinamou (*C. soui*), and Bartlett's Tinamou (*C. bartletti*), which
had overlapping vocalizations and were inconsistently identified with confidence. Non-species
taxa were excluded from calculations of species richness and Shannon diversity but included in
total vocal prevalence. Species were assigned to broad habitat-preference groups: interior
(primary forest) specialists, edge (secondary forest) specialists, and generalists, based on the
Birds of Peru field guide (Schulenberg *et al.* 2010), the Cornell Lab of Ornithology's *Birds of the*
World (Billerman *et al.* 2020), and personal field experience (Supporting Information).

The scientific questions posed in the introduction were addressed as follows:

1: Performance of acoustic sampling. The list of species identified during this study was
compared with data submitted from the reserve to the citizen science database eBird during the
Dec-Feb season, over 13 years through 2020 (Sullivan *et al.* 2009). Because the relative utility of
ARUs vs field observers varies across groups, species were placed in one of 10 guilds to
facilitate more detailed comparisons: soaring raptors, hummingbirds, parrots, aerial insectivores

(swallows and swifts), nocturnal species, open habitat specialists, aquatic habitat specialists (including flooded forest obligates), and canopy, midstory, and understory forest species. Correlation analysis (Pearson's *r*) was employed to assess the relationship between vocal prevalence, species richness, and Shannon diversity. Using vocal prevalence as a stand-in for abundance data, a rank-abundance curve was plotted and compared with the results of Terborgh *et al.* 1990, a historical high-effort census conducted in the region using mist-netting and point counts, to evaluate the performance of ARUs as an alternative to traditional approaches. The total number of species recorded was summarized in a table for each site and day, along with average species richness.

2: Temporal analysis. Paired T-tests were used to compare variation in species richness, Shannon diversity, and vocal prevalence across the three days, and to compare variation in Bray-Curtis dissimilarity between parings of days.

3: Spatial analysis. Two-way repeated-measures ANOVA followed by Tukey's HSD tests for multiple comparisons were conducted in R (R Core Team, 2019), to assess differences in Shannon diversity, species richness, and vocal prevalence between habitat types. To evaluate whether or not community composition differed at edge sites compared to the forest interior, the Bray-Curtis dissimilarity between the edge Site (01) and all other sites was compared to the Bray-Curtis dissimilarity between non-edge sites. Relative soundscape representation (RSR) was assessed for each site day and habitat type by comparing the share of total vocal prevalence to the share of species richness contributed by habitat-preference groups. RSR has the potential to serve as a proxy for per-capita song rate, which is positively correlated with density and territory quality in birds (Pillay *et al.* 2019), therefore aiding in the characterization of habitat type and condition. Paired T-tests were used to assess the significance of representation. Patterns in community composition across sites were assessed using non-metric multidimensional scaling ordination (NMDS). NMDS analyses were calculated using the "bray" method and metaMDS function from the "vegan" package in R (Okansen *et al.* 2019), using vocal prevalence in place of abundance, for all species, interior specialists, and edge specialists, respectively. The NMDS ordinations were performed on two axes and checked to ensure that the output met minimum stress requirements (Kruskal 1964).

3 Results

3.1 Summary Information

We identified 134 species over 21 dawn-hours; 18,963 individual bounding-box annotations were drawn in Raven Pro, of which 94.8% were identified to the species or taxon level. At individual sites, the percent of total Vocal Prevalence that was not identified to a species or taxa group ranged from 4.1% (Site 06) to 7.1% (Site 02) between sites, and from 5.1% (Day C) to 5.4% (Day A) between days. Most annotations occurred in the frequency range 0.5-5

kHz, which features the highest diversity of Amazonian bird songs at dawn, and the least interference from insects, which contribute to significant signal masking between 4-12 kHz (Hart *et al.* 2015, Alvarez-Berríos *et al.* 2016, Aide *et al.* 2017, Metcalf *et al.* 2020). Seven species were recorded only as background vocalizations, and seven more were restricted to Site 08, which was excluded from the majority of analysis because of sustained interference from a Mealy Parrot (*Amazona farinose*), which obscured other vocalizations on Day B and severely impacted overall detections at the site (Supporting Information).

Of the 120 foreground species detected outside of Site 08, two, Black-faced Antthrush (*Formicarius analis*) and Buff-throated Woodcreeper (*Xiphorhynchus guttatus*) were detected on all 18 site days, and six more were detected on at least two-thirds of site days. Most species were rare; more than half (66, 55%) were detected on 4 or fewer site days, and roughly a quarter (27, 23%) on only a single recording (Fig. 3). In terms of total vocal prevalence, the most abundant species were Hauxwell's Thrush (*Turdus hauxwelli*), Black-faced Antthrush, and Thrush-like Wren (*Camplyothynchs turdinus*), which were also the three most common species on eBird.

Figure 3: Number of site-days on which species were detected.

3.2 Performance of Acoustic Sampling

A total of 134 species were identified from recordings across all sites, including background vocalizations and species only detected at Site 08. Through 2020, field observers in the reserve reported 360 species to the citizen science database eBird, spanning 253 complete checklists from December-February. Four species were detected exclusively with ARUs:

Cinereous Mourner (*Laniocera hypopyrra*), Royal Flycatcher (*Onychorhynchus coronatus*), Striped Cuckoo (*Tapera naevia*), and Blue-crowned Manakin (*Lepidothrix 11oronate*), a first record for the reserve, bringing the total species list from both methods to 364. Species reported on >2% and >5% of eBird checklists were considered "regular" and "expected," respectively, and the majority of species reported to eBird but missed by ARUs were neither (Table 1).

	Guild	All Species			Regular Species (>2% of eBird Checklists)			Expected Species (>5% of eBird Checklists)		
		ARUs	eBird	% Coverage	ARUs	eBird	% Coverage	ARUs	eBird	% Coverage
Target Groups	Canopy	29	91	31.9	28	61	45.9	21	40	52.5
	Midstory	36	52	69.2	28	34	82.4	22	25	88.0
	Understory	55	90	61.1	36	48	75.0	29	31	93.5
Non-target Groups	Parrots	5	16	31.3	5	11	45.5	5	6	83.3
	Aquatic	5	35	14.3	3	8	37.5	2	5	40.0
	Open	1	11	9.1	0	4	0.0	0	1	0.0
	Nocturnal	2	10	20.0	1	2	50.0	1	1	100.0
	Aerial Insectivores	0	16	0.0	0	7	0.0	0	2	0.0
	Hummingbirds	0	16	0.0	0	6	0.0	0	2	0.0
	Soaring Raptors	1	23	4.3	1	12	8.3	0	4	0.0
	Total	**134**	**360**	**37.2**	**102**	**193**	**52.8**	**80**	**117**	**68.4**

Table 1: Performance of ARUs vs eBird data by guild and rarity level. eBird detections are based on 253 complete checklists submitted within the reserve from December-February.

ARUs performed best with midstory and understory species, detecting 63% of all species in these foraging strata, including 91% of expected species, but were considerably less effective for canopy foragers, and, as expected, missed the vast majority of species in non-target groups (Table 1). There was a significant positive relationship between vocal prevalence and both species richness ($r = 0.83$, $p = 1.9e\text{-}05$) and Shannon diversity ($r = 0.75$, $p = 3e\text{-}04$) (Fig. 4). Using vocal prevalence as a stand-in for abundance, a rank-order curve was plotted; in line with the abundance-based results of Terborgh *et al.* (1990), we found that vocal prevalence was distributed equitably, fitting a log distribution ($R^2 = 0.84$), with a high proportion of total vocal prevalence contributed by only a few species (Fig. 5).

Figure 4: Vocal prevalence vs Species Richness (A) and Shannon Diversity (B).

Figure 5: Rank-order curves for vocal prevalence (A), using our data, and no. individuals/100 ha (B), using the data from Terborgh *et al.* (1990), an avian census conducted with traditional methods over 12 person-months of effort at Cocha Cashu, another reserve in Madre de Dios.

3.3 Temporal Variation

We captured significant temporal variation in estimated avian species richness, Shannon diversity, vocal prevalence and inter-site community dissimilarity (Table 2). The three diversity and activity measures averaged highest on Day B, and temporal variation was especially consistent for vocal prevalence; at all six sites, vocal prevalence was highest on Day B and

lowest on Day C. Paired t-tests found that Day B had significantly higher average species
richness ($p = 0.024$), Shannon diversity ($p = 0.015$), and VP ($p = 0.0027$) than Day C, which also
had significantly lower average VP than Day A ($p = 0.0026$). Day B also featured consistently
higher values than Day A for species richness ($p = 0.094$), Shannon diversity ($p = 0.092$) and VP
($p = 0.059$), but these results were not statistically significant. The compositional similarity
between sites also differed across days; the average Bray-Curtis dissimilarity between Days A
and B, 0.750, was significantly greater than the Bray-Curtis dissimilarity between Days A and C,
0.715 ($p = 0.022$), and between Days B and C, 0.710, ($p = 0.044$). The highest species total
detected in an individual recording came from Site 02, Day B, which featured 41 species, 46%
more than the 28 species identified at the site on Day A and C; adjacent Site 01 also featured
significantly more species on Day B (32) than on Days A and C (18).

Metric	Day	Total	Mean/pt	SE/pt	Range	Highest Site	Lowest Site
Vocal Prevalence	A	4546	757.7	25	677-844	Site 06	Site 01
	B	5738	956.3	79	794-1335	Site 02	Site 10
	C	3225	537.5	56	315-671	Site 02	Site 01
Species Richness	A	57	27.5	2.1	18-32	Site 06	Site 01
	B	78	33	1.9	28-41	Site 02	Site 01
	C	56	25.7	2.5	18-35	Site 06	Site 01
Shannon Diversity	A	3.50	2.46	0.084	2.10-2.72	Site 06	Site 01
	B	3.57	2.69	0.068	2.53-2.98	Site 02	Site 10
	C	3.26	2.39	0.084	2.15-2.73	Site 06	Site 01

Table 2: Summary of Temporal Variation

3.4 Spatial Variation

Sites in fringe habitats averaged significantly lower species richness ($F_{1, 14} = 10.31$, $p =$
0.0063), Shannon diversity ($F_{1, 14} = 6.98$, $p = 0.019$), and vocal prevalence ($F_{1, 14} = 6.56$, $p =$
0.023) than interior sites (Table 3). Site 01, in edge habitat, was the only site to feature
significantly higher inter-site Bray-Curtis dissimilarity (0.83) than the average between other
sites (0.66, $p = 2.2\text{e-}16$) (Fig. 6), and NMDS ordination showed clustering of interior sites in
terms of species composition (Fig. 7).

Metric	Site	Habitat	Total	Mean/day	SE/day	Range	Highest Day	Lowest Day	UnID%
Vocal Prevalence	01	Fringe	1947	649	185.3	315-955	Day B	Day C	6.5 (206)
	10	Fringe	1939	646.3	108.8	434-794	Day B	Day C	4.2 (119)
	02	Mixed	2767	922.3	208.0	677-1335	Day B	Day C	7.1 (244)
	05	Mixed	2211	737	73.8	604-859	Day B	Day C	4.5 (138)
	04	Várzea	2261	753.7	100.3	560-896	Day B	Day C	5.2 (154)
	06	Várzea	2384	794.7	78.5	641-899	Day B	Day C	4.1 (141)
Species Richness	01	Fringe	43	22.7	4.7	18-32	Day B	Day A/C	
	10	Fringe	42	26.3	2.7	21-30	Day A	Day C	
	02	Mixed	50	32.3	4.3	28-41	Day B	Day A/C	
	05	Mixed	49	29	2.5	24-32	Day B	Day A/C	
	04	Várzea	46	28	1.2	26-30	Day B	Day C	
	06	Várzea	55	34	1.0	32-35	Day B	Day A	
Shannon Diversity	01	Fringe	2.52	2.26	0.112	2.10-2.53	Day B	Day A	
	10	Fringe	2.65	2.45	0.115	2.17-2.64	Day B	Day C	
	02	Mixed	2.89	2.59	0.163	2.32-2.98	Day B	Day A	
	05	Mixed	2.64	2.44	0.075	2.28-2.60	Day B	Day C	
	04	Várzea	2.74	2.56	0.029	2.53-2.63	Day A	Day C	
	06	Várzea	3.02	2.76	0.032	2.72-2.84	Day B	Day A	

Table 3: Summary of Spatial Variation

Figure 6: Bray-Curtis dissimilarity between the edge Site (01) and all other sites vs the Bray-Curtis dissimilarity between non-edge Sites. Same-site comparisons between days were excluded.

Figure 7: Nonmetric multi-dimensional scaling (NMDS) ordinations using Bray-Curtis dissimilarity, with vocal prevalence in place of abundance for interior specialists (A), edge specialists (B), and all species (C). Significantly representative species ($p < 0.02$) are shown with vectors on (C).

Relative soundscape representation (RSR) (Fig. 8) indicated that interior specialists were significantly underrepresented as a share of vocal prevalence (23.8%) relative to their share of overall species richness (41.2%) across the reserve ($p = 1.56e-06$). While consistent across all habitat types, this effect was strongest in fringe habitats, where interior specialists made up 25.8% of average species richness but only 4.7% of average vocal prevalence ($p = 5.28e-05$), followed by mixed habitats (S% = 43.0, VP% = 18.6, $p = 0.0011$) and *várzea* (S% = 54.9, VP% = 48.1, $p = 0.74$). Across all interior habitats, interior specialists made up 49% of species richness and 33.4% of VP ($p = 0.0009$). In contrast with interior specialists, generalists were relatively overrepresented in ITRA's soundscape, making up 38.7% of average species richness and 51.5% of vocal prevalence across all site days ($p = 0.0017$). In fringe habitats, generalists

were proportionally represented, making up 39.7% of species richness and 44.6% of VP, but they were significantly overrepresented in the soundscape of interior habitats, making up 38.2% of species richness and 55.0% of VP ($p = 0.00054$). Soundscape overrepresentation of generalists was strongest in mixed habitats, where they made up 64% of VP compared to 41.1% of species richness ($p = 0.008$). Generalists were also significantly overrepresented in the soundscape of *várzea* sites, accounting for 35.3% of species richness and 45.9% of VP ($p = 0.02$). Edge specialists made up 22.9% of VP and 17.3% of species richness across all sites, but this difference was not statistically significant ($p = 0.10$). In fringe habitats, however, edge specialists made up a significantly higher share of average vocal prevalence, 49.0%, than species richness, 29.4% ($p = 0.012$). Edge specialists contributed 11.2% of species richness and 9.8% of VP at interior sites.

Figure 8: Relative soundscape representation (RSR) of interior (primary forest) specialists (A), edge (secondary forest) specialists (B), and generalists (C) by habitat type. Shaded areas represent one standard deviation of intra-site variation, averaged for each habitat preference group.

4 Discussion

4.1 Performance of Acoustic Sampling

ARUs and manual annotation performed exceptionally well for understory and midstory species in the reserve, but was less effective for canopy foragers. With only 3 days of dawn recordings, we detected almost half (27/63) of the rare (<2% of checklists) midstory and understory species reported to eBird over 13 years, but only one of the 30 rare canopy species, Chestnut-winged Foliage-gleaner (*Philydor erythropterum*). The high-pitched calls of many Amazonian canopy-foraging passerines are extremely challenging to identify to species; even experienced observers are typically unable to use them as a primary identification method in the field. Such calls are poorly represented in acoustic databases like Xeno-canto and the Macaulay Library, making them even harder to learn. Additionally, detection of canopy species was almost certainly impacted by the 1.5 m average placement height of ARUs, especially in a region where canopy heights regularly exceed 30 m (Asner *et al.* 2014). Stevens *et al.* (2019) used ARUs at heights of 2-5m and documented decreased detection rates of canopy-dwelling mixed flock species relative to point counts, but found no such effects in habitats with shorter, smaller trees, supporting the idea that the distance between recorder and canopy height meaningfully influences detection rates of canopy species. Placing recorders in both the canopy and understory would likely improve the detection of species across multiple foraging strata (Bradfer-Lawrence *et al.* 2018, Darras *et al.* 2018, Castro *et al.* 2019). Including annotations from later in the morning is also likely to increase detection rates of canopy foragers, which are known to reach their vocal activity peak in the second or third hour after dawn, and time windows with lower activity would be easier to annotate (Blake 1992, Woltmann 2005, Esquivel & Peris 2008).

Passive acoustic monitoring and manual annotation were an effective avian census method for the forest species that were the focus of this study, detecting 75% of the canopy, midstory, and understory species present on >5% of the 253 December-February eBird checklists in only 21 hours of acoustic data (Table 1). The majority of missed species were rare at the study site or belonged to non-target groups, e.g., species of aquatic and open habitats. A larger sample of habitat types would capture more of these species, but we restricted our ARU placements for consistency. Detections of hummingbirds, aerial insectivores, and soaring raptors, which are primarily visual, are notoriously random in Amazonia and poorly suited for ARUs; only 8/55 (15%) of these species were featured on more than 5% of eBird checklists at ITRA, mostly coming from open areas and the reserve's canopy tower, compared to 96/236 (41%) of canopy, midstory and understory forest species. Additionally, soaring raptors and aerial insectivores have large foraging ranges and are observed primarily as flyovers, meaning that their detection rate can have more to do with visibility conditions than the characteristics of a site on the ground (Kułaga & Budka 2019). As a result, flyover species are of only limited value for assessing differences between sites, and regularly excluded from general point count studies, even though their detectability is one of the biggest advantages of visual survey methods over ARUs (Ralph

et al. 1995, Van Wilgenburg *et al.* 2017). Avian censuses in Amazonia typically find that a high
percentage of species occupy only a small share of sites, which our results reinforced (Terborgh
et al., 1990, Menger *et al.,* 2017, Blake 2021) (Fig. 3). That 23% (27/120) of the species in our
study were featured on just one recording site, and 32% and 27% at one site and day,
respectively, suggests that ARU performance should be expected to improve continuously over a
larger sample (Wood *et al.* 2021). Annotating many short-duration time blocks, representing a
broader range of temporal windows than contiguous dawn-hours, can reduce requisite analysis
time or improve results, including higher detection rates of uncommon and rare species, more
complete estimates of alpha and gamma diversity, and fewer false absences (Shonfield & Bayne
2017, Metcalf *et al.* 2021).
460
Vocal prevalence correlated strongly with estimates of both species richness ($r = 0.83$, p
$= 1.9e$-5) and Shannon diversity ($r = 0.75$, $p = 3e$-4). VP is a better indicator of overall activity
levels than raw call count, total number of annotations, or total annotation time, because it
smooths out natural differences in vocalization patterns across species. While metrics that
require species-level ID are highly contingent on observer skill level, driving inconsistencies in
neotropical bird surveys (Robinson *et al.* 2018), total VP supports the use of broader taxonomic
groups for challenging identifications, lowering time and experience barriers for researchers.
Skill-driven variation in results from bioacoustic surveys is further reduced by the ability to
replay, review, and distribute recordings to other experts, which is impossible to do with point
counts. Substituting 'number of individuals/100 ha' with VP, we found that a rank-order curve of
our data closely mirrored the results of Terborgh *et al.* (1990), a traditional, high-effort avian
census conducted in Madre de Dios by skilled Neotropical field ornithologists (Fig. 5). This
comparison validated the effectiveness of VP a proxy for abundance, enabling the calculation of
abundance-based indices like Bray-Curtis dissimilarity and Shannon diversity with acoustic data.
475
476 **4.2 Temporal Variation**
477
We documented significant temporal variation between days, which was controlled for
with simultaneous monitoring but could distort the results of traditional surveys. While temporal
variation over the course of a morning is well documented in the tropics (Blake 1992, Woltmann
2005, Esquivel & Peris 2008, Blake 2021), the variation that we observed between days was
much greater and more consistent than expected. Though evident in all featured metrics, these
effects were strongest and most uniform for VP, which was highest on Day B and lowest on Day
C at all six sites. Day B was major outlier at adjacent Sites 01 and 02, likely the result of a single
large mixed flock that moved past both sites during the same hour. Plots comparing VP to
Shannon diversity and species richness showed clear clustering by day (Fig. 4).
487
Stochastic vocal activity levels across days mean that temporal variation may obscure, or
even exceed, differences in richness and diversity between sites and habitat types, complicating

the interpretation of point count studies. That temporal variation was strongest for VP is relevant
for point counts because aural detection probability for a field observer is a direct function of
VP, which reflects the probability that a species vocalizes during a given 15-s window. Standard
duration point counts are much more likely to record false absences on days with lower total VP,
even with static species presence. Esquivel & Peris (2008) estimated that it would take four 10-
min visits per point, strategically spread out over different hours of the morning, to sufficiently
cover the hourly variability in avian activity levels, even without considering variation between
days, between different observers (Robinson *et al.* 2008), and travel time between sites. As a
result, Amazonian point count studies seeking to do more than compile species lists may require
sample sizes that are not practical, or risk drawing erroneous conclusions.
500
Many Amazonian bird species vocalize only in brief temporal windows, especially those
surrounding dawn and dusk (Scott *et al.* 1986, Parker 1991, Metcalf *et al.* 2020, pers. obs.).
These effects also apply to broader groups: for example, canopy species are known to reach their
activity peak later in the morning than understory species (Blake 1992, Esquivel & Peris 2008),
and species with high sensitivity to habitat fragmentation may have proportionately higher
detection rates in pre-dawn surveys than less sensitive species (Woltmann 2005). More detailed
data on the specific temporal activity patterns of species and groups will be informative for
selecting annotation windows, estimating avian presence/absence and abundance, and
interpreting the results of point count studies. ARUs are uniquely suited for answering these
questions, because their capacity for simultaneous monitoring can precisely quantify temporal
variation between days, meaning that bioacoustic surveys can draw comparable conclusions with
less field time and/or higher precision than point counts.
513
514   **4.3 Spatial Variation**
515
Intact forest outperformed edge and degraded sites ("fringe" habitats) in all featured
diversity and activity metrics. On all three days, the edge Site (01) featured the lowest species
richness, and the lowest values for vocal prevalence and Shannon diversity came from either Site
01 or Site 10, in degraded forest. These results are consistent with the findings of previously
conducted Amazonian avian inventories using traditional methods, such as point counts and
mist-netting (Terborgh *et al.* 1990, Laurence 2004), demonstrating the viability of ARUs as an
alternative method. This habitat usage pattern differs from observed dynamics in temperate
systems, where avian diversity tends to be higher in edge habitats than forest interiors (Baldi
1996, Lindell *et al.* 2007). Acoustic data is likely superior to point counts for comparing tropical
habitat types with variable visibility conditions; visual detection rate is a direct function of
vegetation density and other habitat characteristics, which can lead to relative overestimation of
avian diversity and abundance in more open habitats (Kułaga & Budka 2019), obscuring the
magnitude of degradation-induced biodiversity loss in the tropics (Celis-Murillo *et al.* 2012,
Darras *et al.* 2018, Darras *et al.* 2019). While sound transmission and auditory detection rate are

also influenced by habitat structure, this impacts both point counts and acoustic monitoring
(Darras *et al.* 2016, Castro *et al.* 2019). Mist-netting surveys are likewise impacted by habitat;
Karr (1981) found that capture rates of canopy-foraging bird species were higher in successional
habitats with low canopy heights than in tall primary forest (Remsen & Parker 1983, Remsen
1996).
535

We found that relative soundscape representation (RSR), comparing the share of VP to
the share of species richness contributed by habitat-preference groups, was a viable and
significantly more efficient alternative to per-capita song rate. Bird species are known to vocalize
more frequently in preferred habitats and where they are abundant (Pillay *et al.* 2019); song rate
is used by females as an indicator of territory quality (Møller 1991, Gil & Gahr 2002), is
positively associated with density-dependent competition between males (Penteriani 2003,
Goretskaia 2004), and may also increase conspecific immigration from surrounding areas
(Fletcher 2007, Laiolo & Tella 2008). While per-capita song rate can be calculated with ARUs
through careful, manual analysis of individual variation between vocalizations (Pillay *et al.*
2019), doing so is labor-intensive and difficult to scale. Mirroring expected patterns for per-
capita song rates, RSR was highest in preferred habitats of both interior and edge specialists.
Even in the reserve's most intact *várzea* forest, however, interior specialists made up a smaller
share of VP than their share of species richness, and they were significantly underrepresented
across all interior sites ($p = 0.0009$). Generalists, conversely, were overrepresented in the
soundscapes of interior sites, an indicator of fragmentation; Amazonian fragments tend to
disproportionately feature generalists, which possess strong dispersal abilities and high tolerance
for deforested habitats (Sekercioglu *et al.* 2002, Lees & Peres 2008). Site 01 was the only site to
significantly differ in community composition compared to all other sites (Fig. 6). The lack of
significant compositional differences between interior sites is another potential symptom of
degradation; Burivalova *et al.* (2019) found that the soundscapes of selectively logged forests in
Borneo were less heterogenous than the soundscapes of a never-logged forest, likely the result of
subtractive homogenization. Together, the soundscape overrepresentation of generalists,
soundscape underrepresentation of interior specialists, and relative compositional homogeneity
of interior sites suggest that the entire 191 ha reserve is degraded, and in the process of becoming
a fragment.
561

Edge effects are known to pervasively impact Amazonian bird communities and
ecosystem function, stretching deep into forest interiors (Terborgh *et al.* 1990, Gascon *et al.*
2000, Laurence 2004, Barlow *et al.* 2006, Broadbent *et al.* 2008, Laurence *et al.* 2011,
Whitworth *et al.* 2015, Wolfe *et al.* 2015, Haddad *et al.* 2015, Moura *et al.* 2016, Stouffer 2020),
but we did not expect RSR to illustrate these dynamics so clearly. Based on its performance in
this study, we suggest that RSR, which can only be calculated with ARUs, has the potential to be
a useful, efficient tool for remotely characterizing habitat features and quality. RSR has
applications for both long-term biodiversity monitoring and rapid assessment, without requiring

established baseline data. Though frequently misrepresented as homogenous, Amazonian
habitats are highly heterogeneous (Pires & Prance 1985, Tuomisto *et al.* 1995), and important
elements of their vegetational diversity remain poorly understood (Milliken *et al.* 2010, Macía
2011, Borges 2013). Interactions between habitat features and avian community structure in the
Amazon are extremely complex, influenced by variables like soil type (Borges 2013), riverine
sediment concentration (Laranjeiras *et al.* 2021), the condition of surrounding habitat (Stouffer
2006, Latta *et al.* 2011, Laurence *et al.* 2011, Woltmann 2012, Laurence *et al.* 2012, Wolfe *et al.*
2015, Barlow 2016, Menger *et al.* 2017, Hernández-Palma & Stouffer 2018), the presence of
large animals (Redford 1992, Estes *et al.* 2011), and climate change (Blake & Loiselle 2015,
Stouffer *et al.* 2020). As a result, Amazonian habitat assessments, which are generally reliant on
features discernable by field observers, can be overly simplistic. Under global change, RSR and
other approaches that assess nuanced habitats based on their use by species may be particularly
valuable.

5 Synthesis & Applications

Developing a better understanding of the effects that deforestation, fragmentation, and
degradation have on biodiversity in tropical biodiversity hotspots is a major international
conservation priority (Gardner *et al.* 2008, Barlow *et al.* 2016, Stouffer 2020), particularly
analyses that go beyond presence/absence and that can be applied at landscape scales
(Hernández-Palma & Stouffer 2018). The results of this study support the potential of
simultaneous passive acoustic monitoring as a cornerstone of biodiversity assessment, for both
birds and other taxa. We found that annotated acoustic data can replicate the results of historical
avian censuses, characterizing community structure, species richness, Shannon diversity, and
vocal activity levels across habitat types. We documented significant temporal variation between
days, which may be missed with non-simultaneous survey methods, distorting their results.

Acoustic data can generate informative, novel metrics that are impossible to produce with
human observers, including effective proxies for abundance data and per-capita song rate. Such
metrics, namely VP and RSR, can efficiently characterize complex habitats based on species
usage patterns, rather than relying on visible features. In the future, passive acoustic monitoring's
scalability, permanent storage, and ability to integrate with new and developing technologies
could have a transformative impact on global conservation efforts (Kahl *et al.* 2020). Rampant
habitat destruction and climate change threaten to destabilize the Amazon basin, and long-term,
perhaps cascading declines in biodiversity and ecosystem function have already been
documented, even in structurally intact habitats (Bird & Lees 2011, Laurence *et al.* 2012, Blake
& Loiselle 2015, Brienen *et al.* 2015, Lovejoy & Nobre 2019, Stouffer 2020). If we wish to save,
or at least document, these imperiled systems, furthering cutting-edge approaches like
bioacoustics should be a central research priority.

Authors' Contributions

Both authors designed the methodology. H.K. helped to secure the funding to conduct the study; W.A.H. collected and annotated the recordings, analyzed the data, and led the writing of the manuscript. Both authors contributed critically to the drafts and gave final approval for publication.

Acknowledgements

We would like to thank the Inkaterra Association (ITA) staff for providing logistical support and excellent field station facilities, particularly Noe Huaraca, Dennis Osorio, and Kevin Jiménez Gonzales, who helped set up recorders. Noe, John Fitzpatrick, Fernando Angulo, Will Sweet, Ken Rosenburg, and Alex Wiebe helped identify unknown vocalizations, and Ken Rosenburg, Andrew Farnsworth, and Connor Wood provided comments on the draft. Viviana Ruiz-Gutierrez and Amanda Rodewald gave general guidance, and Sidd Srinivasan assisted with Python. Funding for equipment was provided by the K. Lisa Yang Center for Conservation Bioacoustics at the Cornell Lab of Ornithology, with support from Innóvate Perú, CORBIDI, and the Inkaterra Association. Travel expenses were funded by the Cornell Lab of Ornithology.

Data Availability Statement

All data collected is available on request and will be made publicly available upon acceptance.

References

Acevedo, M. A., & Villanueva-Rivera, L. J. (2006). Using Automated Digital Recording Systems as Effective Tools for the Monitoring of Birds and Amphibians. *Wildlife Society Bulletin*, *34*(1), 211–214. https://doi.org/10.2193/0091-7648(2006)34[211:uadrsa]2.0.co;2

Aide, T. M., Hernández-Serna, A., Campos-Cerqueira, M., Acevedo-Charry, O., & Deichmann, J. L. (2017). Species richness (of insects) drives the use of acoustic space in the tropics. *Remote Sensing*, *9*(11), 1–12. https://doi.org/10.3390/rs9111096

Allan, J. R., Venter, O., & Watson, J. E. M. (2017). Temporally inter-comparable maps of terrestrial wilderness and the Last of the Wild. *Scientific Data*, *4*, 1–8. https://doi.org/10.1038/sdata.2017.187

Alvarez-Berríos, N., Campos-Cerqueira, M., Hernández-Serna, A., Delgado, C. J. A., Román-Dañobeytia, F., &
Aide, T. M. (2016). Impacts of small-scale gold mining on birds and anurans near the Tambopata Natural
Reserve, Peru, assessed using passive acoustic monitoring. *Tropical Conservation Science*, *9*(2), 832–851.
https://doi.org/10.1177/194008291600900216

Asner, G. P., Anderson, C. B., Martin, R. E., Knapp, D. E., Tupayachi, R., Sinca, F., & Malhi, Y. (2014).
Landscape-scale changes in forest structure and functional traits along an Andes-to-Amazon elevation
gradient. *Biogeosciences*, *11*(3), 843–856. https://doi.org/10.5194/bg-11-843-2014

Baldi, A. (1996). Edge effects in tropical versus temperate forest bird communities: Three alternative hypotheses for
the explanation of differences. *Acta Zoologica Academiae Scientiarum Hungaricae*, *42*(3), 163–172.

Barlow, J., Peres, C. A., Henriques, L. M. P., Stouffer, P. C., & Wunderle, J. M. (2006). The responses of
understorey birds to forest fragmentation, logging and wildfires: An Amazonian synthesis. *Biological
Conservation*, *128*(2), 182–192. https://doi.org/10.1016/j.biocon.2005.09.028

Barlow, J., Ewers, R. M., Anderson, L., Aragao, L. E. O. C., Baker, T. R., Boyd, E., … Gardner, T. A. (2011). Using
learning networks to understand complex systems: A case study of biological, geophysical and social research
in the Amazon. *Biological Reviews*, *86*(2), 457–474. https://doi.org/10.1111/j.1469-185X.2010.00155.x

Barlow, J., Lennox, G. D., Ferreira, J., Berenguer, E., Lees, A. C., Nally, R. Mac, … Gardner, T. A. (2016).
Anthropogenic disturbance in tropical forests can double biodiversity loss from deforestation. *Nature*,
535(7610), 144–147. https://doi.org/10.1038/nature18326

Barlow, J., França, F., Gardner, T. A., Hicks, C. C., Lennox, G. D., Berenguer, E., … Graham, N. A. J. (2018). The
future of hyperdiverse tropical ecosystems. *Nature*, *559*(7715), 517–526. https://doi.org/10.1038/s41586-018-
0301-1

Berg, K. S., Brumfield, R. T., & Apanius, V. (2006). Phylogenetic and ecological determinants of the neotropical
dawn chorus. *Proceedings of the Royal Society B: Biological Sciences*, *273*(1589), 999–1005.
https://doi.org/10.1098/rspb.2005.3410

Bibby, C. J. (1999). Making the most of birds as environmental indicators. *Ostrich*, *70*(1), 81–88.
https://doi.org/10.1080/00306525.1999.9639752

Billerman, S.M., Keeney, B. K., P. G. Rodewald, and T. S. Schulenberg (Editors) (2020). Birds of the World.
Cornell Laboratory of Ornithology, Ithaca, NY, USA. https://birdsoftheworld.org/bow/home

Bird, J. P., & Lees, A. C. (2011). Acknowledging an Amazon avifauna in crisis. *Neotropical Birding*, *11*(July 2012),
74–77.

Bird, J. P., Buchanan, G. M., Lees, A. C., Clay, R. P., Develey, P. F., Yépez, I., & Butchart, S. H. M. (2012).
Integrating spatially explicit habitat projections into extinction risk assessments: A reassessment of
Amazonian avifauna incorporating projected deforestation. *Diversity and Distributions*, *18*(3), 273–281.
https://doi.org/10.1111/j.1472-4642.2011.00843.x

Blackburn, T. M., & Gaston, K. J. (1998). Some methodological issues in macroecology. *American Naturalist*,
151(1), 68–83. https://doi.org/10.1086/286103

Blake, J. G. (1992). Temporal Variation in Point Counts of Birds in a Lowland Wet Forest in Costa Rica. *The*
Condor, *94*(1), 265–275. https://doi.org/10.2307/1368816

Blake, J. G., & Loiselle, B. A. (2015). Enigmatic declines in bird numbers in lowland forest of eastern Ecuador may
be a consequence of climate change. *PeerJ*, *2015*(8), 1–20. https://doi.org/10.7717/peerj.1177
700

701   Blake, J. G. (2021). Acoustic monitors and direct observations provide similar but distinct perspectives on bird
702   assemblages in a lowland forest of eastern Ecuador. *PeerJ*, (January). https://doi.org/10.7717/peerj.10565
703

Boakes, E. H., McGowan, P. J. K., Fuller, R. A., Chang-Qing, D., Clark, N. E., O'Connor, K., & Mace, G. M.
(2010). Distorted views of biodiversity: Spatial and temporal bias in species occurrence data. *PLoS Biology*,
8(6). https://doi.org/10.1371/journal.pbio.1000385

Borges, S. H. (2013). Bird species distribution in a complex Amazonian landscape: Species diversity, compositional
variability and biotic-environmental relationships. *Studies on Neotropical Fauna and Environment*, *48*(2),
106–118. https://doi.org/10.1080/01650521.2013.841627

Bradfer-Lawrence, T., Gardner, N., & Dent, D. H. (2018). Canopy bird assemblages are less influenced by habitat
age and isolation than understory bird assemblages in Neotropical secondary forest. *Ecology and Evolution*,
8(11), 5586–5597. https://doi.org/10.1002/ece3.4086
713

714   Brienen, R. J. W., Phillips, O. L., Feldpausch, T. R., Gloor, E., Baker, T. R., & Lloyd, J. (2015). This is a repository
715   copy of Long-term decline of the Amazon carbon sink . White Rose Research Online URL for this paper :
716   Version : Accepted Version Article : Brienen , RJW orcid . org / 0000-0002-5397-5755 , Phillips , OL Long-
717   term decline of the Amazon. *Nature*, *519*(7543), 344–348.
718

719   Broadbent, E. N., Asner, G. P., Keller, M., Knapp, D. E., Oliveira, P. J. C., & Silva, J. N. (2008). Forest
720   fragmentation and edge effects from deforestation and selective logging in the Brazilian Amazon. *Biological*
721   *Conservation*, *141*(7), 1745–1757. https://doi.org/10.1016/j.biocon.2008.04.024
722

723   Burivalova, Z., Purnomo, Wahyudi, B., Boucher, T. M., Ellis, P., Truskinger, A., … Game, E. T. (2019). Using
724   soundscapes to investigate homogenization of tropical forest diversity in selectively logged forests. *Journal of*
725   *Applied Ecology*, *56*(11), 2493–2504. https://doi.org/10.1111/1365-2664.13481
726

727   Canaday, C., & Rivadeneyra, J. (2001). Initial effects of a petroleum operation on Amazonian birds: Terrestrial
728   insectivores retreat. *Biodiversity and Conservation*, *10*(4), 567–595.
729   https://doi.org/10.1023/A:1016651827287
730

731   Castro, I., De Rosa, A., Priyadarshani, N., Bradbury, L., & Marsland, S. (2019). Experimental test of birdcall
732   detection by autonomous recorder units and by human observers using broadcast. *Ecology and Evolution*,
733   *9*(5), 2376–2397. https://doi.org/10.1002/ece3.4775
734

735   Cellis-Murillo, A., Deppe, J., & Ward, M. P. (2012). Evaluating the use of acoustic monitoring for surveying
736   tropical birds. *Journal of Field Ornithology*, *83(2)*(May 2012), 166–179. https://doi.org/10.2307/23256481
737

Clarke, D. A., York, P. H., Rasheed, M. A., & Northfield, T. D. (2017). Does Biodiversity–Ecosystem Function
Literature Neglect Tropical Ecosystems? *Trends in Ecology and Evolution*, *32*(5), 320–323.
https://doi.org/10.1016/j.tree.2017.02.012

Darras, K., Pütz, P., Fahrurrozi, Rembold, K., & Tscharntke, T. (2016). Measuring sound detection spaces for
acoustic animal sampling and monitoring. *Biological Conservation*, *201*, 29–37.
https://doi.org/10.1016/j.biocon.2016.06.021

Darras, K., Batáry, P., Furnas, B., Celis-Murillo, A., Van Wilgenburg, S. L., Mulyani, Y. A., & Tscharntke, T.
(2018). Comparing the sampling performance of sound recorders versus point counts in bird surveys: A meta-
analysis. *Journal of Applied Ecology*, *55*(6), 2575–2586. https://doi.org/10.1111/1365-2664.13229

Darras, K., Batáry, P., Furnas, B. J., Grass, I., Mulyani, Y. A., & Tscharntke, T. (2019). Autonomous sound
recording outperforms human observation for sampling birds: a systematic map and user guide. *Ecological
Applications*, *29*(6). https://doi.org/10.1002/eap.1954

Edelman, A. et al. State of the Tropics: 2014 Report (James Cook Univ., Cairns, 2014).Available at:
https://researchonline.jcu.edu.au/35471/1/SOTT_ExcecutiveSummary_Print1.pdf

Esquivel M., Alberto & Peris, S. (2008). Influence of time of day, duration and number of counts in point count
sampling of birds in an atlantic forest of Paraguay. *Ornitologia Neotropical*, *19*(2), 229–242.

Estes, J. A., Terborgh, J., Brashares, J. S., Power, M. E., Berger, J., Bond, W. J., … Wardle, D. A. (2011). Trophic
downgrading of planet earth. *Science*, *333*(6040), 301–306. https://doi.org/10.1126/science.1205106

Fletcher, R. J. (2007). Species interactions and population density mediate the use of social cues for habitat
selection. *Journal of Animal Ecology*, *76*(3), 598–606. https://doi.org/10.1111/j.1365-2656.2007.01230.x

Foster, R. B., Parker III, T. A., Gentry, A. H., Emmons, L. H., Chicchón, A., Schulenberg, T., … Wust, W. (1994).
The Tambopata-Candamo Reserved Zone of Southeastern Perú : A Biological Assessment Conservation
Priorities : The Role of RAP. In *Conservation Biology*. Retrieved from
https://www.conservation.org/docs/default-source/publication-pdfs/rap06_tambopata_candamo_peru_nov-
1994.pdf?Status=Master&sfvrsn=72f3b47a_3

Gardner, T. A., Barlow, J., Araujo, I. S., Ávila-Pires, T. C., Bonaldo, A. B., Costa, J. E., … Peres, C. A. (2008). The
cost-effectiveness of biodiversity surveys in tropical forests. *Ecology Letters*, *11*(2), 139–150.
https://doi.org/10.1111/j.1461-0248.2007.01133.x

Gascon, C., Williamson, G. B., & Da Fonseca, G. A. B. (2000). Receding forest edges and vanishing reserves.
Science, *288*(5470), 1356–1358. https://doi.org/10.1126/science.288.5470.1356

Gil, D., & Gahr, M. (2002). The honesty of bird song: Multiple constraints for multiple traits. Trends in Ecology and
Evolution, 17, 133–141. https://doi.org/10.1016/S0169-5347(02)02410-2

Goretskaia, Maria I. (2004). Song Structure and Singing Behaviour of Willow Warbler Phylloscopus in Populations
of Low and High Density. *Bioacoustics*, (November 2012), 183–195.
https://doi.org/10.1080/09524622.2004.9753524

Haddad, N. M., Brudvig, L. A., Clobert, J., Davies, K. F., Gonzalez, A., Holt, R. D., … Townshend, J. R. (2015).
Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances*, *1*(2), 1–10.
https://doi.org/10.1126/sciadv.1500052

Hart, P. J., Hall, R., Ray, W., Beck, A., & Zook, J. (2015). Cicadas impact bird communication in a noisy tropical
rainforest. *Behavioral Ecology*, *26*(3), 839–842. https://doi.org/10.1093/beheco/arv018

Haselmayer, J., & Quinn, J. S. (2000). A comparison of point counts and sound recording as bird survey methods in
amazonian southeast Peru. *Condor*, *102*(4), 887–893. https://doi.org/10.2307/1370317
790

791 Hernández-Palma, A., & Stouffer, P. C. (2018). Matrix and area effects on the nutritional condition of understory
792   birds in Amazonian rainforest fragments. *Perspectives in Ecology and Conservation*, *16*(3), 139–145.
793   https://doi.org/10.1016/j.pecon.2018.06.003
794

795 Hobson, K. A., Rempel, R. S., Greenwood, H., Turnbull, B., & Van Wilgenburg, S. L. (2002). Acoustic surveys of
796   birds using electronic recordings: New potential from an omnidirectional microphone system. *Wildlife Society*
797   *Bulletin*, *30*(3), 709–720.
798

799 Jenkins, C. N., Pimm, S. L., & Joppa, L. N. (2013). Global patterns of terrestrial vertebrate diversity and
800   conservation. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(28),
801   E2603–E2610. https://doi.org/10.1073/pnas.1302251110
802

803 Jorge, F. C., Machado, C. G., da Cunha Nogueira, S. S., & Nogueira-Filho, S. L. G. (2018). The effectiveness of
804   acoustic indices for forest monitoring in Atlantic rainforest fragments. *Ecological Indicators*, *91*(August
805   2017), 71–76. https://doi.org/10.1016/j.ecolind.2018.04.001
806

Kahl, S., Clapp, M., Hopping, W. A., Goëau, H., Glotin, H., Planqué, R., Vellinga,W. P., Joly, A. (2020). Overview
of BirdCLEF 2020: Bird sound recognition in complex acoustic environments. In: CLEF task overview 2020,
CLEF: Conference and Labs of the Evaluation Forum, Sep. 2020, Thessaloniki, Greece.

Kahl, S., Wood, C. M., Eibl, M., & Klinck, H. (2021). BirdNET: A deep learning solution for avian diversity
monitoring. *Ecological Informatics*, *61*(December 2020), 101236.
https://doi.org/10.1016/j.ecoinf.2021.101236

Kułaga, K., & Budka, M. (2019). Bird species detection by an observer and an autonomous sound recorder in two
different environments: Forest and farmland. *PLoS ONE*, *14*(2), 1–19.
https://doi.org/10.1371/journal.pone.0211970
816

817 Laiolo, P., & Tella, J. L. (2008). Social determinants of songbird vocal activity and implications for the persistence
818   of small populations. *Animal Conservation*, *11*(5), 433–441. https://doi.org/10.1111/j.1469-
819   1795.2008.00202.x
820

821 Laranjeiras, T. O., Naka, L. N., Leite, G. A., Cohn-Haft, M. (2021). Effects of a major Amazonian river confluence
822   on the distribution of floodplain forest avifauna, *Journal of Biogeography*, 48, 4, (847-860).
823   https://doi.org/10.1111/jbi.14042
824

825 Latta, S. C., Tinoco, B. A., Astudillo, P. X., & Graham, C. H. (2011). Patterns and magnitude of temporal change in
826   avian communities in the ecuadorian andes. *Condor*, *113*(1), 24–40. https://doi.org/10.1525/cond.2011.090252
827

828 Laurance, W. F., Lovejoy, T. E., Vasconcelos, H. L., Bruna, E. M., Didham, R. K., Stouffer, P. C., … Sampaio, E.
829   (2002). Ecosystem decay of Amazonian forest fragments: A 22-year investigation. *Conservation Biology*,
830   *16*(3), 605–618. https://doi.org/10.1046/j.1523-1739.2002.01025.x

831
832 Laurance, S. G. W. (2004). Responses of understory rain forest birds to road edges in central Amazonia. *Ecological*
833   *Applications*, *14*(5), 1344–1357. https://doi.org/10.1890/03-5194
834

835 Laurance, W. F., Camargo, J. L. C., Luizão, R. C. C., Laurance, S. G., Pimm, S. L., Bruna, E. M., … Lovejoy, T. E.
836   (2011). The fate of Amazonian forest fragments: A 32-year investigation. *Biological Conservation*, *144*(1),
837   56–67. https://doi.org/10.1016/j.biocon.2010.09.021
838

839 Laurance, W. F., Useche, D. C., Rendeiro, J., Kalka, M., Bradshaw, C. J. A., Sloan, S. P., … Campbell, M. (2012).
840   Averting biodiversity collapse in tropical forests. *Nature*, *489*, 290–294.
841

842 Laurance, W. F., Sayer, J., & Cassman, K. G. (2014). Agricultural expansion and its impacts on tropical nature.
843   *Trends in Ecology and Evolution*, *29*(2), 107–116. https://doi.org/10.1016/j.tree.2013.12.001
844

845 Lees, A. C., & Peres, C. A. (2006). Rapid avifaunal collapse along the Amazonian deforestation frontier. *Biological*
846   *Conservation*, *133*(2), 198–211. https://doi.org/10.1016/j.biocon.2006.06.005
847

848 Lees, A. C., Naka, L. N., Aleixo, A., Cohn-Haft, M., Piacentini, V. de Q., Santos, M. P. D., & Silveira, L. F. (2014).
849   Conducting rigorous avian inventories: Amazonian case studies and a roadmap for improvement. *Revista*
850   *Brasileira de Ornitologia*, *22*(2), 107–120.
851

852 Lindell, C. A., Riffell, S. K., Kaiser, S. A., Battin, A. L., Smith, M. L., & Sisk, T. D. (2007). Edge Responses of
853   Tropical and Temperate Birds. *The Wilson Journal of Ornithology*, *119*(2), 205–220.
854   https://doi.org/10.1676/05-133.1
855

856 Lovejoy, T. (2013). Tropical forests under a changing climate and innovations in tropical forest management.
857   *Tropical Conservation Science*, *6*(3), 311–314. https://doi.org/10.1177/194008291300600301
858

Lovejoy, T. E., & Nobre, C. (2019). Amazon tipping point: Last chance for action. *Science Advances*, *5*(12), 4–6.
https://doi.org/10.1126/sciadv.aba2949

Macía, M. J. (2011). Spatial distribution and floristic composition of trees and lianas in different forest types of an
Amazonian rainforest. *Plant Ecology*, *212*(No. 7 (July 2011)), 1159–1177. https://doi.org/10.1007/s11258-
011-9895-1

Mahowald, N. M., Ward, D. S., Doney, S. C., Hess, P. G., & Randerson, J. T. (2017). Are the impacts of land use on
warming underestimated in climate policy? *Environmental Research Letters*, *12*(9).
https://doi.org/10.1088/1748-9326/aa836d
867

868 Menger, J., Magnusson, W. E., Anderson, M. J., Schlegel, M., Pe'er, G., & Henle, K. (2017). Environmental
869   characteristics drive variation in Amazonian understorey bird assemblages. *PLoS ONE*, *12*(2), 1–20.
870   https://doi.org/10.1371/journal.pone.0171540
871

872 Metcalf, O. C., Barlow, J., Devenish, C., Marsden, S., Berenguer, E., & Lees, A. C. (2020). Acoustic indices
873   perform better when applied at ecologically meaningful time and frequency scales. *Methods in Ecology and*
874   *Evolution*, *2020*(October), 1–11. https://doi.org/10.1111/2041-210x.13521
875

876 Metcalf, O. C., Barlow, J., Marsden, S., Moura, N. G. De, Berenguer, E., Ferreira, J., & Lees, A. C. (2021).
877   Optimizing tropical forest bird surveys using passive acoustic monitoring and high temporal resolution
878   sampling. *Remote Sensing in Ecology and Conservation*, 1–12. https://doi.org/10.1002/rse2.227
879

880 Milliken, W., Zappi, D., Sasaki, D., Hopkins, M., & Pennington, R. T. (2010). Amazon vegetation: How much don't
881   we know and how much does it matter? *Kew Bulletin*, *65*(4), 691–709. https://doi.org/10.1007/s12225-010-
882   9236-x
883

884 Møller, A. P. (1991). Why mated songbirds sing so much: Mate guarding and male announcement of mate fertility
885   status. The American Naturalist, 138, 994–1014. https ://doi.org/10.1086/285264
886

887 Moura, N. G., Lees, A. C., Aleixo, A., Barlow, J., Berenguer, E., Ferreira, J., … Gardner, T. A. (2016). Idiosyncratic
888   responses of Amazonian birds to primary forest disturbance. *Oecologia*, *180*(3), 903–916.
889   https://doi.org/10.1007/s00442-015-3495-z
890

891 Newson, S. E., Bas, Y., Murray, A., & Gillings, S. (2017). Potential for coupling the monitoring of bush-crickets
892   with established large-scale acoustic monitoring of bats. *Methods in Ecology and Evolution*, *8*(9), 1051–1062.
893   https://doi.org/10.1111/2041-210X.12720
894

895 Nicolau, A. P., Herndon, K., Flores-Anderson, A., & Griffin, R. (2019). A spatial pattern analysis of forest loss in
896   the Madre de Dios region, Peru. *Environmental Research Letters*, *14*(12). https://doi.org/10.1088/1748-
897   9326/ab57c3
898

899 Oksanen, J., Guillaume Blanchet, F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'hara, R.
900   B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H. (2019). vegan: Community
901   ecology package. R package version 2.5-6. Retrieved from https:// CRAN.R-project.org/packa ge=vegan
902

903 Parker, T. (1991). On the Use of Tape Recorders in Avifaunal Surveys. *The Auk: Ornithological Advances*, *108*(02),
904   443–444. https://doi.org/10.1093/auk/108.2.443
905

906 Penteriani, V. (2003). Breeding density affects the honesty of bird vocal displays as possible indicators of
907   male/territory quality. *Ibis*, *145*(3), 513–513. https://doi.org/10.1046/j.1474-919x.2003.01734.x
908

909 Phillips, O., Gentry, A. H., Reynel, C., Wilkin, P., & Galvez-Durand B, C. (1994). Quantitative Ethnobotany and
910   Amazonian Conservation. *Conservation Biology*, *8*(1), 225–248. https://doi.org/10.1046/j.1523-
911   1739.1994.08010225.x
912

913 Pillay, R., Fletcher, R. J., Sieving, K. E., Udell, B. J., & Bernard, H. (2019). Bioacoustic monitoring reveals shifts in
914   breeding songbird populations and singing behaviour with selective logging in tropical forests. *Journal of
915   Applied Ecology*, *56*(11), 2482–2492. https://doi.org/10.1111/1365-2664.13492
916

917 Pires, J.M., Prance, G. 1985. The vegetation types of Brazilian Amazon. In: Prance GT, Lovejoy TE, editors. Key
918   Environment – Amazonia. New York (NY): Pergamon Press. p. 109–145.
919

920 R Core Team. (2019). R: A language and environment for statistical comput- ing. R Foundation for Statistical
921   Computing. Retrieved from https:// www.R-proje ct.org/
922

923 Ralph, C. J., Sauer, J. R., & Droege, S. (1995). Monitoring Bird Populations by Point Counts. *The Journal of*
924   *Wildlife Management*, *61*(4), 1453. https://doi.org/10.2307/3802161
925

926 Redford, K. H. (1992). The Empty Forest. *BioScience*, *42*(No. 6 (Jun., 1992)), 412–422.
927   https://doi.org/https://doi.org/10.2307/1311860
928

929 Rempel, R. S., Hobson, K. A., Holborn, G., Van Wilgenburg, S. L., & Elliott, J. (2005). Bioacoustic monitoring of
930   forest songbirds: interpreter variability and effects of configuration and digital processing methods in the
931   laboratory. *Journal of Field Ornithology*, *76*(1), 1–11. https://doi.org/10.1648/0273-8570-76.1.1
932

933 Rempel, R. S., Jackson, J. M., Van Wilgenburg, S. L., & Rodgers, J. A. (2019). A multiple detection state
934   occupancy model using autonomous recordings facilitates correction of false positive and false negative
935   observation errors. *Avian Conservation and Ecology*, *14*(2). https://doi.org/10.5751/ACE-01374-140201
936

937 Remsen, J. V, & Parker III, T. A. (1983). Contribution of River-Created Habitats to Bird Species Richness in
938   Amazonia. *Biotropica*, *15*(3), 223–231.
939

940 Remsen, J. V. (1994). Use and Misuse of Bird Lists in Community Ecology and Conservation. *The Auk*, *111*(1),
941   225–227. https://doi.org/10.2307/4088531
942

Remsen, J. V., & Good, D. A. (1996). Misuse of Data from Mist-Net Captures to Assess Relative Abundance in
Bird Populations. *The Auk*, *113*(2), 381–398.

Robinson, S. K., & Terborgh, J. (1997). Bird Community Dynamics along Primary Successional Gradients of an
Amazonian Whitewater River. *Ornithological Monographs*, (48), 641–672. Retrieved from
https://www.jstor.org/stable/40157559

Robinson, W. D., Lees, A. C., & Blake, J. G. (2018). Surveying tropical birds is much harder than you think: a
primer of best practices. *Biotropica*, *50*(6), 846–849. https://doi.org/10.1111/btp.12608
950

951 Rodriguez, A., Gasc, A., Pavoine, S., Grandcolas, P., Gaucher, P., & Sueur, J. (2014). Temporal and spatial
952   variability of animal sound within a neotropical forest. *Ecological Informatics*, *21*, 133–143.
953   https://doi.org/10.1016/j.ecoinf.2013.12.006
954

955 Sánchez-Cuervo, A. M., de Lima, L. S., Dallmeier, F., Garate, P., Bravo, A., & Vanthomme, H. (2020). Twenty
956   years of land cover change in the southeastern Peruvian Amazon: implications for biodiversity conservation.
957   *Regional Environmental Change*, *20*(1). https://doi.org/10.1007/s10113-020-01603-y
958

959 Schulenberg, Thomas S., et al. *Birds of Peru: Revised and Updated Edition.* Revised and Updated Edition ed.
960   Princeton University Press, 2010. *Project MUSE* muse.jhu.edu/book/30246.
961

962 Scott, J. M., S. Mountainspring, F. L. Ramsey, and C. B. Kepler. 1986. Forest bird communities of the Hawaiian
963   Islands: their dynamics, ecology, and conservation. Stud. Avian Biol. No. 9.
964

965 Sekercioglu, Ç. H., Ehrlich, P. R., Daily, G. C., Aygen, D., Goehring, D., & Sandí, R. F. (2002). Disappearance of
966   insectivorous birds from tropical forest fragments. *Proceedings of the National Academy of Sciences of the*
967   *United States of America*, *99*(1), 263–267. https://doi.org/10.1073/pnas.012616199
968

969   Shonfield, J., & Bayne, E. M. (2017). Autonomous recording units in avian ecological research: current use and
970   future applications. *Avian Conservation and Ecology*, *12*(1). https://doi.org/10.5751/ace-00974-120114
971

Socolar, J. B., Valderrama Sandoval, E. H., & Wilcove, D. S. (2019). Overlooked biodiversity loss in tropical
smallholder agriculture. *Conservation Biology*, *33*(6), 1338–1349. https://doi.org/10.1111/cobi.13344

Stevens, H. C., Metz, E. M., Saboya Del Castillo, P., Alván, J. D., & Bowler, M. T. (2019). Use of autonomous
audio recordings for the rapid inventory of birds in the white-sand forests of the Peruvian Amazon. *Journal of*
Field Ornithology, *90*(1), 70–79. https://doi.org/10.1111/jofo.12279

Stouffer, P. C. (2020). Birds in fragmented Amazonian rainforest: Lessons from 40 years at the Biological
Dynamics of Forest Fragments Project. *The Condor*, *122*, 1–15. https://doi.org/10.1093/condor/duaa005
979

980   Stouffer, P. C., Jirinec, V., Rutt, C. L., Bierregaard, R. O., Hernández-Palma, A., Johnson, E. I., … Lovejoy, T. E.
981   (2020). Long-term change in the avifauna of undisturbed Amazonian rainforest: ground-foraging birds
982   disappear and the baseline shifts. *Ecology Letters*. https://doi.org/10.1111/ele.13628
983

984   Sugai, L. S. M., & Llusia, D. (2019). Bioacoustic time capsules: Using acoustic monitoring to document
985   biodiversity. *Ecological Indicators*, *99*(December 2018), 149–152.
986   https://doi.org/10.1016/j.ecolind.2018.12.021
987

988   Sullivan, B. L., Wood, C. L., Iliff, M. J., Bonney, R. E., Fink, D., & Kelling, S. (2009). eBird: A citizen-based bird
989   observation network in the biological sciences. *Biological Conservation*, *142*(10), 2282–2292.
990   https://doi.org/10.1016/j.biocon.2009.05.006
991

992   Tegeler, A. K., Morrison, M. L., & Szewczak, J. M. (2012). Using extended-duration audio recordings to survey
993   avian species. *Wildlife Society Bulletin*, *36*(1), 21–29. https://doi.org/10.1002/wsb.112
994

995   Terborgh, J., Robinson, S. K., Parker III, T. A., & Munn, C. A. (1990). Structure and Organization of an Amazonian
996   Forest Bird Community. *Ecological Monographs*, *60*(2), 213–238.
997

998   Tuomisto, H., Ruokolainen, K., Kalliola, R., Linna, A., Danjoy, W., & Rodriguez, Z. (1995). Dissecting amazonian
999   biodiversity. *Science*, *269*(5220), 63–66. https://doi.org/10.1126/science.269.5220.63
1000

1001   United Nations General Assembly. Transforming our world: the 2030 Agenda for Sustainable Development (United
1002   Nations, New York, 2015). Available at
1003   https://sustainabledevelopment.un.org/content/documents/21252030%20Agenda%20for%20Sustainable%20D
1004   evelopment%20web.pdf
1005

1006   Van Wilgenburg, S. L., Sólymos, P., Kardynal, K. J., & Frey, M. D. (2017). Paired sampling standardizes point
1007   count data from humans and acoustic recorders. *Avian Conservation and Ecology*, *12*(1).
1008   https://doi.org/10.5751/ace-00975-120113
1009

1010   Venier, L. A., Holmes, S. B., Holborn, G. W., McIlwrick, K. A., & Brown, G. (2012). Evaluation of an automated
1011   recording device for monitoring forest birds. *Wildlife Society Bulletin*, *36*(1), 30–39.
1012   https://doi.org/10.1002/wsb.88
1013

1014 Whitworth, A., Beirne, C., Rowe, J., Ross, F., Acton, C., Burdekin, O., & Brown, P. (2015). The response of faunal
1015  biodiversity to an unmarked road in the Western Amazon. *Biodiversity and Conservation*, *24*(7), 1657–1670.
1016  https://doi.org/10.1007/s10531-015-0883-y

1017

1018 Wilson, K. A., *et al*. (2016). Conservation Research Is Not Happening Where It Is Most Needed. *PLoS Biology*,
1019  *14*(3), 1–5. https://doi.org/10.1371/journal.pbio.1002413

1020

1021 Wolfe, J. D., Stouffer, P. C., Mokross, K., Powell, L. L., & Anciaes, M. M. (2015). Island vs. countryside
1022  biogeography: An examination of how Amazonian birds respond to forest clearing and fragmentation.
1023  *Ecosphere*, *6*(12), 1–14. https://doi.org/10.1890/ES15-00322.1

1024

1025 Woltmann, S. (2005). Patterns of daily temporal variation in detectability of forest birds in Bolivia. *Ornitologia*
1026  *Neotropical*, *16*(3), 337–346.

1027

1028 Woltmann, S., Kreiser, B. R., & Sherry, T. W. (2012). Fine-scale genetic population structure of an understory
1029  rainforest bird in Costa Rica. *Conservation Genetics*, *13*(4), 925–935. https://doi.org/10.1007/s10592-012-
1030  0341-2

1031

1032 Wood, C.M., Kahl, S., Chaon, P., Peery, M.Z., Klinck, H. (2021). Survey coverage, record-
1033  ing duration and community composition affect observed species richness in passive
1034  acoustic surveys. *Methods in Ecology and Evolution* 12(5), 885–896. https://doi.org/10.1111/2041-
1035  210X.13571