1

KLC4 shapes axon arbors during development and mediates adult behavior

Elizabeth M. Haynes[1,2,3,4], Korri H. Burnett[1,3], Jiaye He[4,5], Marcel Jean-Pierre[1,3], Martin
Jarzyna[1,3], Kevin W. Eliceiri[2,4], Jan Huisken[1,4,6], and Mary C. Halloran[1,3]

1.Department of Integrative Biology, University of Wisconsin-Madison

2. Center for Quantitative Cell Imaging, University of Wisconsin-Madison

3. Department of Neuroscience, University of Wisconsin-Madison

4. Morgridge Institute for Research, Madison, WI

5. National Innovation Center for Advanced Medical Devices, Shenzen, China.

6. Dept. of Biology and Psychology, Georg-August-University Göttingen, Germany

*Corresponding author:
Mary C. Halloran
Department of Integrative Biology
1117 W. Johnson Street
Madison, WI 53706
mchalloran@wisc.edu
608-263-7875

23
24

Abstract

Development of elaborate and polarized neuronal morphology requires precisely regulated transport of cellular cargos by motor proteins such as kinesin-1. Kinesin-1 has numerous cellular cargos which must be delivered to unique neuronal compartments. The process by which this motor selectively transports and delivers cargo to regulate neuronal morphogenesis is poorly understood, although the cargo-binding kinesin light chain (KLC) subunits contribute to specificity. Our work implicates one such subunit, KLC4, as an essential regulator of axon branching and arborization pattern of sensory neurons during development. Using live imaging approaches in *klc4* mutant zebrafish, we show that KLC4 is required for stabilization of nascent axon branches, proper microtubule (MT) dynamics, and endosomal transport. Furthermore, KLC4 is required for proper tiling of peripheral axon arbors: in *klc4* mutants, peripheral axons showed abnormal fasciculation, a behavior characteristic of central axons. This result suggests that KLC4 patterns axonal compartments and helps establish molecular differences between central and peripheral axons. Finally, we find that *klc4* mutant larva are hypersensitive to touch and adults show anxiety-like behavior in a novel tank test, implicating *klc4* as a new gene involved in stress response circuits.

Introduction

Proper development of complex neuronal morphology and assembly of neural circuits requires tight regulation of axon growth and branching. Precise control over axon branch formation is essential not only for development but also for plasticity and regeneration after injury. Branching allows one neuron to innervate numerous targets, and in many cases, neurons extend multiple axons or branches that navigate to distinct targets. Many molecular signaling pathways can influence branching (Gallo 2011; Kalil and Dent 2014), although the mechanisms that control branch formation are still not fully understood. Highly arborized vertebrate sensory neurons, including Rohon-Beard (RB) neurons, serve as an ideal model to study mechanisms of branching and neuronal morphogenesis. Sensory neurons must extend separate axon branches to targets in the periphery and to the CNS to relay sensory information. Peripheral and central axons have different capacity for branching (E. Andersen, Asuri, and Halloran 2011), and behave differently when contacting other axons—central axons fasciculate with one another while peripheral axons show contact repulsion (Liu and Halloran 2005; Sagasti et al. 2005). Contact repulsion of peripheral axons is necessary for establishment of proper axon arbor patterns that "tile" with arbors from other neurons to effectively innervate the sensory field. Despite the importance of establishing axons with distinct character and branching patterns, the mechanisms regulating molecular compartmentalization of separate axons from one neuron are poorly understood.

Molecular compartmentalization of neurons depends on precisely controlled transport of vesicle and organelle cargos to specific cell locations. Cargos are transported by motor proteins that move in a directional manner along MT tracks. Kinesin-1, the founding member of the kinesin superfamily of motor proteins, is a heterotetramer composed of homodimers of kinesin heavy chains (KHCs), which contain the MT binding and ATPase motor domains, and of kinesin light chains (KLCs), which interact with cargos. Kinesin-1 can mediate long-distance, anterograde transport of multiple cargos including mitochondria, synaptic vesicle precursors, endosomes, lysosomes and RNA granules (Hirokawa, Niwa, and Tanaka 2010; Morfini et al. 2016). Kinesin-1 also has roles in organizing MT tracks. The forces it exerts can physically influence the MT lattice (Shima et al. 2018; Triclin et al. 2021). It can also cross-link MTs by combined interactions of the motor domain and a second MT binding site in the KHC tail, and slide MTs along one another, thereby influencing cell shape (Lu and Gelfand 2017). While it is clear that kinesin-1 is essential for proper development of the nervous system, the mechanisms by which

75 it achieves cell-context specificity of its multiple functions and how its cargo-binding and MT-
76 organizing functions are regulated have yet to be fully defined.
77
The specific activities of kinesin-1 in a given context are likely governed in part by KLCs. KLCs
are known to mediate binding between the kinesin motor and cellular cargos, and also to
activate the motor by releasing KHC autoinhibition (Morfini et al. 2016; K. Wong et al. 2010;
Wozniak and Allan 2006). Vertebrates have four *klc* genes (*klc1-4*), with *klc1, 2* and *4*
expressed in the nervous system, but our knowledge of the different roles played by individual
KLCs in neurons remains very limited. In *Drosophila*, mutants for the single *klc* gene are lethal
in larval stages and have defects in neural development (Gindhart et al. 1998; Mochizuki et al.
2011), indicating essential neural functions for KLC. Growing evidence supports the idea that
molecular functions are divided among vertebrate KLCs and that each has unique roles.
Structural and biochemical studies show that KLC1 and KLC2 have differing affinities for cargo
adaptor proteins such as JIP1 and JIP3 (Cockburn et al. 2018; Pernigo et al. 2018; Zhu et al.
2012). Research to date has begun to define distinct neurophysiological functions of KLC1 and
KLC2. Studies of mouse *klc1* mutants revealed functions for KLC1 in regulating motor behavior
(Rahman et al. 1999), transport of amyloid precursor protein into axons (Kamal et al. 2000),
phagosome transport in retinal pigment epithelium (Jiang et al. 2015), and axon guidance and
cannabinoid receptor transport during neural development (Saez et al. 2020). Mouse *klc2*
mutants have hearing loss, cochlear hair cell loss and mislocalized mitochondria in hair cells (Fu
et al. 2021). A study using a peptide inhibitor of KLC2 phosphorylation showed that KLC2
regulates synaptic plasticity, AMPAR trafficking and mood associated behavior (Du et al. 2010).
Mutations in human *klc2* can cause SPOAN syndrome, which is characterized by progressive
spastic paraplegia, optic atrophy and neuropathy with onset in infancy (Bazvand et al. 2021;
Melo et al. 2015). On the whole, previous work supports the idea that individual KLCs have
unique, indispensable functions, but has only begun to reveal the specific cellular functions for
individual KLCs.
102
In contrast to KLC1 and KLC2, KLC4 has been the subject of few studies. Mutation of *KLC4* in
humans causes a type of hereditary spastic paraplegia (HSP) that manifests in early childhood
(Bayrakli et al. 2015), indicating essential developmental functions. To date, the only cellular
function shown for KLC4 is that it supports mitochondrial function and confers resistance to
radiation in cancer cells (Baek et al. 2018). A clear cellular function for KLC4 in neurons and
therefore insight into the *KLC4* mutation-related pathology is lacking.
109
In this study, we reveal critical functions for KLC4 in sensory axon branching and axon arbor
patterning during development, and in regulation of animal behavior. Using live imaging in *klc4*
mutant zebrafish, we found that KLC4 is required for stabilization of nascent peripheral sensory
axon branches. The branching defects do not reflect a general growth defect, as axons grew
faster in *klc4* mutants and did not show signs of early degeneration. Live imaging also showed
that *klc4* mutant peripheral axons have defects in mutual repulsion and instead often fasciculate
with one another, a behavior characteristic of central axons, suggesting KLC4 may contribute to
the molecular compartmentalization that defines differential behavior of these axons. Using
high-speed, *in vivo* imaging of endosomal vesicle transport and MT dynamics, we showed that
klc4 mutant axons have altered vesicle transport and MT dynamics. Nascent branches in *klc4*
mutants have less acetylated tubulin, suggesting less stable MTs. Finally, we investigated the
ramifications of KLC4 loss on larval and adult behavior. Larval *klc4* mutant zebrafish show
hyperactivity in response to touch, implying heightened sensitivity. Adult *klc4* mutants exhibit
increased anxiety-like behavior, indicating essential roles for KLC4 in neural circuit formation
and/or function in adults.
125

Results
klc4 is expressed during early development
klc genes encode proteins with conserved functional domains including the heptad repeat region that mediates binding to KHC, six cargo-binding tetratricopeptide repeats (TPRs), a C-terminal lipid-binding amphipathic helix, and an LFP motif that mediates KLC auto-inhibition (Figure 1A) (Anton 2021; Cockburn et al. 2018; Morfini et al. 2016; Pernigo et al. 2018; Zhu et al. 2012). As a first step toward understanding developmental functions of KLC4, we investigated the expression of zebrafish *klc4* at different stages of development. Zebrafish *klc4* has two alternative splice isoforms, one long and one short, both of which are also present in other vertebrates including mice and humans. In zebrafish, the long *klc4* transcript encodes a full length protein of 620 amino acids while the short isoform is predicted to encode a 185 amino acid protein, ending just before the LFP motif (Figure 1 – figure supplement Figure1A). To determine whether both transcripts are expressed during development, we performed non-quantitative reverse transcriptase-PCR with transcript-specific primers on RNA extracted from embryos during early development (Figure 1B, Figure 1 - figure supplement 1A). We found that both isoforms are expressed throughout early development, with expression detectable at 2 hours post fertilization (hpf) and continuing through the initial period of neuronal axon outgrowth, pathfinding and branching, between 16 and 28 hpf (Figure 1B). We next used *in situ* hybridization to determine the spatial expression pattern of *klc4* during development, using a riboprobe that binds both transcripts. At 24 hpf, *klc4* is expressed in clusters of neurons in the telencephalon, epiphysis, diencephalon, ventral midbrain and hindbrain, and in the trigeminal and the posterior lateral line (pLL) ganglia (Figure1 C,D). In the spinal cord, *klc4* is strongly expressed in the Rohon-Beard sensory neurons, and at lower levels in other neuron populations (Figure 1E). At 48 hpf, *klc4* continues to be expressed broadly in the nervous system with high expression levels in the trigeminal ganglia, pLL ganglia, and RB neurons (Figure 1 F,G,H). A riboprobe that only binds the long isoform shows a similar expression pattern to the probe that recognizes both isoforms (Figure 1I,J,K), indicating the full-length *klc4* mRNA is present in developing neurons. The early expression of *klc4* in diverse neural populations in both the central and peripheral nervous system suggests it has functions in neural development.

Generation of klc4 mutants
To assess the role of KLC4 during neural development, we used CRISPR/Cas9 to generate *klc4* mutants. We used a guide RNA targeting exon 3 of *klc4*, which encodes a part of the N-terminal heptad repeat region. One resulting allele, *klc4^uw314^*, has a 25bp deletion in exon 3 from 404bp to 428bp in the heptad repeat region that leads to an early stop after 9 missense amino acids (Figure 1A, figure supplement 1B). Heterozygous *klc4^uw314^* mutants were outcrossed to wild type (AB strain) fish for three generations to remove any off-target mutations from the background. *In situ* hybridization analysis of homozygous *klc4^uw314^* mutant embryos showed virtually no *klc4* mRNA expression, suggesting that the mutant transcript undergoes nonsense-mediated decay (Figure 1 L,M,N).

klc4^uw314^ mutants show reduced branching of RB peripheral sensory axons
Because *klc4* is highly expressed in spinal RB sensory neurons, we first analyzed the effects of *klc4* mutation on their axon growth and morphology. We crossed the *klc4^uw314^* mutants with a transgenic line in which RB neurons are labeled with membrane-targeted GFP [*Tg(-3.1ngn1:gfp-caax)*] (E. Andersen, Asuri, and Halloran 2011) and used light sheet microscopy to image RB axon development across the whole embryo (Figure 2A,B). Inspection of mutant embryos revealed that RB peripheral axon branches appeared less dense and displayed less overall innervation of the sensory field compared to wild type (Figure 2A,B, Video 1, Video 2).

To quantify these differences we labeled sensory neurons with anti-HNK1 antibody at 24 hpf
(Figure 2C,D,E). We counted the number of axons that crossed 8 regions of interest drawn at
standardized intervals from the central axon fascicle, proceeding from dorsal to ventral (Figure 2
– figure supplement 2). Consistent with our visual impression of reduced branching, mutant
embryos had consistently fewer axons at each ROI (Figure 2F). The reduction in branches at
the most dorsal levels in the mutant embryos may indicate an effect on initiation of the
peripheral axon, which most often forms as a branch from the central axon near the cell body
(E. Andersen, Asuri, and Halloran 2011).
184
As an alternative approach to understand the role of KLCs in axon branching, and to develop a
more time-resolved means to disrupt KLC function, we took a pharmacological approach.
Specifically, embryos were treated with Kinesore, a small molecule that has been shown to
interfere with the interaction between the KLC TPR repeat domain and cargo adaptor proteins,
while also relieving KLC auto-inhibition and activating kinesin-1's ability to cross-link MTs and
regulate MT dynamics (Randall et al. 2017). We found that treatment of wild type embryos with
100 µM Kinesore during early stages of peripheral axon arborization, from 17 hpf to 24 hpf,
caused a reduction in axon branch number in ventral regions at 24 hpf of a similar magnitude to
that seen in $klc4^{uw314}$ mutants (Figure 2G,H,I), though more dorsal branching appears
unaffected. These results give further evidence that properly regulated KLC-cargo interactions
or kinesin-1-MT interactions during axon outgrowth are important for successful axon branching.
196
KLC4 is required for stabilization of axon branches
To investigate which specific dynamic processes during branching are affected by *klc4* loss of
function, we used live confocal imaging. Zebrafish RB axons branch by two mechanisms:
growth cone bifurcation (GCB) and interstitial back branching (IB) (Gibson and Ma 2011). To
determine if GCB, IB or both were defective in mutants, we used the *Tg(-3.1ngn1:gfp-caax)*
transgenic line (E. Andersen, Asuri, and Halloran 2011). Z-stacks of RB peripheral axon arbors
were collected every 1-1.5 minutes for 2-4 hours beginning from approximately 18 hpf. The
number of branch initiation and retraction events during a two-hour time window were counted
and categorized as either a GCB (Figure 3A, white arrows) or IB event (Figure 3A, magenta
arrows). Surprisingly, we found no difference in the rate of branch initiation between $klc4^{uw314}$
mutant and wild type embryos (Figure 3C). In both conditions, new branches were initiated via
GCB more frequently than IB, but there was no difference in the proportion of GCB versus IB
branching events between wild type and $klc4^{uw314}$ mutants.
210
Nevertheless, the live imaging immediately suggested an explanation for the branch number
deficit: failure of branches to persist. That is, following formation, while most of the branches in
wild type animals were relatively stable and continued growing (Video 3), those in the mutants
often retracted after formation (Figure 3B, Video 4). Quantification showed that in $klc4^{uw314}$
mutants, 57.7% of GCB branches and 61.9% of IB branches retracted after initiating, compared
to 14.2% of GCB and 11.8% of IB branches in wild type (Figure 3D). Mutants showed no
significant directional bias to the retracted branches (on average, 52.5% of branches lost per
neuron were anteriorly-directed, and 47.2% were posteriorly-directed. p=0.6588). This
phenotype cannot be explained as a general axon growth defect, as $klc4^{uw314}$ mutant axons
grew at a faster rate than wild type axons (Figure 3E). Thus, these data suggest that KLC4 is
required specifically for stabilization of axon branches.
222
Sensory axon arborization pattern is altered in $klc4^{uw314}$ mutants
Peripheral sensory axons develop a distinctive arborization pattern to successfully innervate the
sensory field. During this process, axons mutually repel each other upon contact (Sagasti et al.
2005; Liu and Halloran 2005). Precise regulation of contact repulsion as well as branching are

critical for effective tiling of the entire skin. To determine whether mutants have altered
arborization and to quantify branching pattern in an unbiased manner, we developed an image
analysis pipeline to measure axon directional orientation and overall axon abundance (Figure
4). We used OrientationJ (Rezakhaniha et al. 2011) to measure the fluorescence signal of
aligned pixels at each orientation (Figure 4A,B) and created an axon directionality "profile" that
represents the average signal at each 1.0 degree angle interval across multiple embryos (Figure
4C). In wild type embryos, most axons are oriented in ventral-posterior and ventral-anterior
directions, with a higher proportion of axon signal in posteriorly directed angles, peaking
approximately between -50° and -85° (Figure 4 A,C, figure supplement 4A). Little signal was
detected in the horizontal (0 degrees) orientation. In contrast, $klc4^{uw314}$ mutants had reduced
overall axon signal compared to their wild type cousins (Figure 4B, C, figure supplement 4B),
consistent with the reduced branch density. Mutants showed an overall directional profile similar
to wild type (Figure 4B,C). However, because differences in axon orientation between wild type
and mutants may be masked by the overall difference in signal, we normalized the data and
displayed it as a percentage of total signal at each angle. We found a small but significant
increase in axons oriented in the posterior direction in $klc4^{uw314}$ mutants, specifically those
oriented between -20.5° and -50.5° (Figure 4D). There was no difference in the proportion of
anterior-directed axons between wild type and $klc4^{uw314}$ mutants (Figure 4 – figure supplement
4C).
246
To investigate what changes in axon behavior could generate the posterior orientation bias in
mutants, we turned to live imaging. We analyzed the growth direction of actively extending
axons by measuring the orientation angle of the distal-most axon shaft at each time point over
2-4 hour movies (Figure 5A) and displaying the distribution of growth angles as rose plots. We
found $klc4^{uw314}$ mutant axons had a preference for posterior growth paths, while wild type axons
had no anterior-posterior bias (Figure 5B, C). We then measured the angles between newly
bifurcating sister branches. Prior to bifurcation, axons commonly grow in a posterior direction in
both wild type and mutant embryos (Figure 5D). As growth cones bifurcate in wild type embryos,
the sister branches grow away from each other (Figure 5D',E, Video 5), resulting in an average
branch angle of 96.5 degrees 15 minutes after branch formation, and 91.7 degrees 75 minutes
after branch formation (Figure 5E, G). This behavior results in axon arbors with an even
distribution of axon orientation angles in wild type embryos (Fig 4 C). In $klc4^{uw314}$ mutants, the
newly formed branches did not separate as much, resulting in a shallower average angle of 79.7
degrees between sister branches 15 minutes after branch formation, and narrowing to 60.2
degrees after 75 minutes (Figure 5 F,G, Video 6). In fact, branches in $klc4^{uw314}$ mutants at times
failed to repel at all, instead appearing to collapse onto one another (see Figure 6B). These
results suggest that KLC4 may be involved in mediating mutual repulsion between sister axon
branches. Moreover, the posterior growth bias detected by automated orientation analysis in
$klc4^{uw314}$ mutant embryos (Figure 4C) is likely the result of branch separation failure. Because
axons initially tend toward posterior growth, reduced branch separation or collapse of anteriorly
directed branches onto the more posterior axon shaft leads to axon arbors with an overall
increase in posterior orientation angles. Thus, our automated, population-scale method of axon
orientation profiling led to the discovery of a branch separation defect, demonstrating the utility
of the profiling method for revealing alterations in axon arbor pattern.
271
Peripheral axons in $klc4^{uw314}$ mutants exhibit abnormal fasciculation behavior
Live imaging also revealed a broader defect in axon repulsion: normally, in wild type embryos
when a peripheral axon growth cone contacts another axon (either a branch of the same neuron
or of another neuron) it retracts or changes direction to grow away from the other axon, thus
minimizing contact between axons (Sagasti et al. 2005; Liu and Halloran 2005) (Figure 6A).
However, in $klc4^{uw314}$ mutants, axons often fasciculate after such contacts and grow together for

substantial distances (Figure 6B,C, Video 7). In some cases, axons were fasciculated so tightly
that they only became distinguishable as two axons when pulled apart by the force of a
developing branch (Figure 6C, arrows, Video 7, minutes 41-47, Video 8, minutes 24-44). In wild
type embryos, all observed fasciculations (n=3) resolved by retraction or defasciculation, but
only 46.5% of fasciculations (n=19) in $klc4^{uw314}$ mutants were resolved in this manner.
Fasciculation behavior was quantified by measuring the frequency and length of fasciculation
events occurring during our live imaging period. We found that fasciculation events were rare in
wild type embryos but occurred relatively frequently in mutants (Figure 6D). Fasciculations
ranged from 7.5-60.5 μm in length in $klc4^{uw314}$ mutants (Figure 6E). Together, these results
show that KLC4 is important for the mutual repulsion needed to shape sensory arbors. The
mutual repulsion normally exhibited by peripheral axons is in contrast to the fasciculation
behavior of the central axons from the same neuron. The failure of $klc4^{uw314}$ mutant peripheral
axons to show repulsion suggests that KLC4 may be involved in the localization of factors
controlling central versus peripheral axon behavior.
292
RB neurons have aberrant apical axons in $klc4^{uw314}$ mutants
RB neurons in $klc4^{uw314}$ mutants also showed abnormal morphology suggestive of altered cell
polarity. We found that some RB neurons in $Tg(-3.1ngn1:gfp-caax);klc4^{uw314}$ mutant embryos
and Kinesore treated embryos projected a process from the apical cell body that crossed the
midline within the dorsal spinal cord (Figure 7A,B, Video 9, 10). Small apical protrusions
sometimes form on RB neurons, especially in the anterior portion of the spinal cord, though they
can be found nearly anywhere along the spinal cord. These processes are typically short,
dynamic, and are most common early in development (18-24 hpf) (Video 11). In wild type
embryos, they rarely cross the midline of the spinal cord. We counted the number of midline
crossing apical processes from images of anti-HNK-1-labeled embryos captured in the spinal
cord overlying the yolk extension and found that 4% of neurons (n=1509 neurons in 51
embryos) in $klc4^{uw314}$ mutants had midline crossing processes compared to 1% in wild type
embryos (n=608 neurons in 21 embryos) (Figure 7C). Similar results were obtained following
treatment with Kinesore from 18-24 hpf, with 7% of Kinesore-treated neurons (n=433 neurons in
17 embryos) extending an apical protrusion that crossed the midline, compared to 2% of
neurons in vehicle treated control (n=436 neurons in 15 embryos) (Figure 7C). To further
assess neuronal morphology and the pathways of the aberrant projections, we sparsely labeled
individual RB neurons by injecting DNA encoding TagRFP-caax driven by the -3.1ngn1
promoter at the 1-cell stage. This mosaic cell labeling showed that the apical processes grew to
and fasciculated with the contralateral central axon fascicle in both $klc4^{uw314}$ mutant and
kinesore treated embryos (Figure 7D,F), suggesting they may respond to cues directing central
axon behavior despite emerging from an ectopic apical location. These results suggest that
KLCs have a role in determining the location of axon emergence from the cell body.
316
Endosomal vesicle transport is altered in $klc4^{uw314}$ mutants
To begin exploring the mechanisms by which KLC4 affects axon arbor development, we imaged
endosomal vesicle transport. We previously found that a properly functioning endosomal
trafficking system is necessary for RB axon branching (Ponomareva et al. 2014; Ponomareva,
Eliceiri, and Halloran 2016). Moreover, we showed that the kinesin adaptor Clstn1 is required for
branching and that it functions in part by trafficking Rab5-labeled endosomal vesicles to branch
points (Ponomareva et al. 2014). We used a previously described zebrafish Rab5c construct
(Clark et al. 2011) to label Rab5-containing endosomes *in vivo*. We expressed GFP-Rab5c in
individual RB neurons by injecting *ngn1:GFP-Rab5c* DNA into embryos at the 1-cell stage
(Figure 8A). We imaged vesicle movement using a high-speed swept field confocal microscope
that allows imaging of rapid events in 3D. We captured z-stacks of RB axon arbors at 2-5
second intervals for 7-10 minute periods and used kymography to quantify endosome dynamics

(Figure 8A'). We found some alterations in transport. In *klc4*[uw314] mutants, a smaller proportion
of vesicles moved in the anterograde direction compared to wild type (Figure 8B), and the
maximum anterograde vesicle run length was reduced (Figure 8C), suggesting that some
aspects of Rab5 vesicle transport or a subset of Rab5 vesicles rely on KLC4 function. However,
there was no significant difference in velocity of the Rab5 vesicles that were moving, suggesting
these vesicles can attach to the motor without KLC4 (Figure 8D).
335
To determine whether KLC4 regulates Rab5 vesicular transport from the cell body specifically to
axon branch points, we labeled Rab5c with photoactivatable GFP (PA-GFP). We co-injected
ngn1:PA-GFP-Rab5c and *ngn1:TagRFP-caax* to label individual RB neurons. We
photoactivated the GFP in the cell body during stages when peripheral axons are actively
growing and branching, and captured images at ~1-2 minute intervals for 30-40 minutes after
photoactivation (Figure 8—figure supplement 8A). We previously reported that in wild type
embryos, PA-GFP-Rab5c vesicles originating in the cell body appeared to accumulate at the
peripheral axon initiation point and in secondary branch points within minutes of photoactivation
(Ponomareva et al. 2014). To quantify vesicle accumulations at branch points over time, we
divided the movies into 4 time periods (t1-t4), made maximum intensity projections of the z-
stacks for t1-t4, and measured the fluorescent signal at branch points as a percentage of the
signal at the first time period (t1). In both wild type and *klc4* mutant neurons, the intensity
increased similarly during this period, suggesting KLC4 does not play a significant role in vesicle
delivery from the cell body to branch points (Figure 8—figure supplement 8B).
350
Microtubule dynamics are altered in *klc4*[uw314] mutants
Because local changes in MT dynamics are associated with axon branch formation, and MT
invasion of nascent axon branches is required for branches to be stabilized and maintained (Yu,
Ahmad, and Baas 1994; Dent and Kalil 2001; Yu et al. 2008; Gallo 2011; Armijo-Weingart and
Gallo 2017), we assessed whether the MT cytoskeleton is affected in *klc4*[uw314] mutants. To
visualize MT dynamics *in vivo,* we used the MT plus-tip binding protein EB3 fused to GFP
(Stepanova et al. 2003). EB3-GFP binds to the plus ends of actively polymerizing MTs, which
appear as moving GFP puncta or "comets". We expressed EB3-GFP in RB neurons by injecting
ngn1:EB3-GFP DNA, and imaged with a swept field confocal microscope. We captured z-stacks
at 2-5 second intervals for periods ranging from 7-10 minutes and analyzed microtubule
dynamics using kymography (Figure 9A, Video 12). In axons, the majority of comets move in the
anterograde direction, reflecting the plus-end-distal organization of MTs. We found no change in
the percentage of retrograde comets between wild type and *klc4*[uw314] mutants, suggesting KLC4
is not necessary for organization of MT polarity (Figure 9 B,C). However, we found a significant
increase in the frequency of anterograde comets in peripheral axons, but not central axons, in
klc4[uw314] mutant RB neurons (Figure 9B,D, figure supplement 9A), suggesting increased MT
dynamics in peripheral axons. In addition, we found that the velocity of EB3-GFP comets, both
anterograde and retrograde, was increased in peripheral and central axons in *klc4*[uw314] mutants
(Figure 9E, figure supplement 9B). There was no difference in the distance traveled by EB3-
GFP comets in any condition (Figure 9—figure supplement 9C,D), though duration of
anterograde comet runs was reduced in peripheral axons, consistent with their increased
velocity (Figure 9—figure supplement 9E,F). Thus, the length of new MT polymer growth at plus
ends is not affected in mutants, but mutant MTs polymerize faster, suggesting a role for KLC4 in
regulating the growth rate of MT plus ends.
375
MTs in nascent branches are less stable in *klc4*[uw314] mutants
To ask whether MT dynamics are altered in newly forming branches, we co-injected *ngn:EB3-
GFP* DNA with *ngn:tagRFP-caax* DNA to label cell membranes, and imaged EB3-GFP comets
in nascent branches (Figure 10A). We found no difference in the frequency of comets in nascent

branches between wild type and mutants (Figure 10B). EB3 comet frequency also did not differ
between mutant and wild type when separated by branch type (IB vs GCB, data not shown,
p>0.7 by t-test). However, because EB3-GFP comet frequency in peripheral axons overall is
increased in *klc4*^{uw314} mutants, the equivalent comet frequencies between mutant and wild type
in nascent branches may represent a net reduction in MT polymerization into nascent branches
in mutants.
386
As another approach to assess MT stability in new branches, we fixed embryos at a stage of
active branching (24 hpf) and double-labeled neurons with anti-HNK1 and an antibody to
acetylated tubulin, a modification found on stable MTs (Webster and Borisy 1989)(Figure
10C,D). We analyzed terminal branches up to 10 μm in length formed by interstitial branching or
bifurcation, then scored those with acetylated tubulin signal. We found a significant reduction in
the percentage of nascent branches with acetylated tubulin in *klc4*^{uw314} mutants (Figure 10E).
This finding suggests that KLC4 is required for MTs to stabilize in newly formed branches.
394
Larval and adult *klc4*^{uw314} fish show behavioral defects
The formation of functional neural circuits ultimately depends on the precise orchestration of cell
biological processes during neural development. To ask whether KLC4 loss affects circuit
function, we analyzed larval escape behavior in response to touch, a behavior mediated by
sensory neurons. Larva at 3 days post-fertilization (dpf) were given a light tail touch, evoking a
swim response (Figure 11A). The time of the swim response and number of swim bouts
completed was measured for each larva. *klc4*^{uw314} mutant larva swam for a longer total time
(wild type median = 0.65 seconds, *klc4*^{uw314} 1.49 seconds) and performed more swim bouts
(wild type median = 1 bout, *klc4*^{uw314} mutants = 2 bouts) after touch stimulus than wild type larva
(Figure 11B,C), suggesting KLC4 loss leads to hypersensitivity to touch.
405
The cell biological defects found during development in *klc4*^{uw314} mutants seemed likely to lead
to deficits in adult fish as well. Indeed, we found that *klc4*^{uw314} mutants are adult viable and
fertile but are smaller in size than their wild type siblings (Figure 11—figure supplement 11A).
To measure body size in a controlled manner, we in-crossed F4 homozygous *klc4*^{uw314} mutants
from both AB and mixed AB/TL backgrounds and separately in-crossed their wild type (AB)
siblings. The F5 embryos were born on the same day, raised side-by-side in an identical
environment, and fed the same diet. We measured the body length of these fish at 7 weeks and
at 9 weeks (Figure 11—figure supplement 11A). *klc4*^{uw314} mutants were on average 83.3% the
size of their wild type cousins at 7 weeks. This difference remained at 9 weeks, despite both the
wild type and *klc4*^{uw314} mutant fish increasing in size, with *klc4*^{uw314} mutant fish measured at
84.6% the length of their wild type cousins. Smaller body size has also been reported in *klc1*
mutant mice (Rahman et al. 1999), supporting the idea that KLCs are important for post-
embryonic growth of the animal.
419
Multiple neural systems are affected in the human disease caused by *KLC4* mutation; patients
show defects in vision, hearing, movement and cognition (Bayrakli et al. 2015). Thus, we
analyzed adult behavior to more broadly investigate KLC4's roles in neural circuit function. To
ask if the mutant's small size reflected a defect in feeding behavior, we assayed feeding ability
(Howe, McIntyre, and Wolman 2018). Adult zebrafish were isolated in individual 1L tanks, then
50 larval zebrafish were added to each tank. The fish were allowed to feed on the larvae for one
hour, then removed from the tank to allow counting of the remaining larvae. Wild type fish were
effective feeders, eating on average 90% of available zebrafish larvae (Figure 11 – figure
supplement 11C). In comparison, *klc4*^{uw314} mutants ate on average only 60% of the larvae
available. We noted that the mutant fish that performed the worst in the feeding assay

underwent bouts of freezing behavior during the assay. Upon introduction of the larvae to the
tank, the fish ceased movement for extended periods.
432
Freezing behavior in zebrafish in response to a novel environment has been previously
described and interpreted as increased anxiety (K. Wong et al. 2010; Maximino et al. 2010; Ziv
et al. 2013; Khan et al. 2017). To test whether *klc4*^{uw314} mutant fish show anxiety-like behavior,
we used a novel tank diving test, which is similar in principle to the open field test used in
rodents (K. Wong et al. 2010; Maximino et al. 2010). In this test, fish are introduced to an
unfamiliar tank and allowed to explore. Zebrafish typically prefer to stay in the lower zone of the
tank initially, and once they become acclimated, they begin to explore the upper regions of the
tank (Figure 11D). We introduced adult animals (7 months old) to a novel tank and filmed their
behavior for 3 minutes. Analysis revealed that 90.9% of wild type fish explored the upper zone
of the tank during the 3 minute window, compared to only 18.8% of the *klc4*^{uw314} mutant fish
(Figure 11E). Of the animals that did explore the upper zone, both wild type and *klc4*^{uw314}
mutant animals took an average of 100 seconds before their first cross into the upper zone
(Figure 11F). Wild type animals spent an average of 14 seconds in the upper region of the tank,
compared to just 3.26 seconds for *klc4*^{uw314} mutants (Figure 11G). Notably, freezing behavior,
defined for this experiment as cessation of all movement except for gills for at least 3 seconds,
was observed in an average of 43% of mutant fish, and only 3% of wild type fish (Figure 11H).
15.4% of *klc4*^{uw314} mutant fish, but no wild type animals, froze for the full 3 minute duration of the
experiment (Figure 11I).
451
Studies in both zebrafish and rodents have shown that repeated exposure to the unfamiliar
environment leads to habituation of the anxiety behavior. In zebrafish, habituation manifests as
increased exploration of the upper zone and reduced freezing behavior. Fish that do not
habituate to the novel environment or that worsen over multiple exposures are thought to
demonstrate a depression-like phenotype (Ziv et al. 2013). Behaviors driven by increased
anxiety are expected to lessen with repeat exposures. We tested the ability of *klc4*^{uw314} mutants
to habituate to a novel environment after multiple exposures. In addition, because KLC4 and
other kinesin-1 subunits have been implicated in neurodegenerative diseases, we sought to
determine whether the anxiety-like behavior worsened with age. We challenged 1.25-1.5 year
old *klc4*^{uw314} mutants and their wild type cousins by repeated exposure to a novel tank for 10
minutes, twice a week for 3 weeks in succession. The first exposure allowed us to test the
effects of age. We found that increased age improved freezing response of *klc4*^{uw314} mutants,
with only 29.2% (8 out of a total of 23) of individuals tested freezing for more than 3 seconds
during first 3 minutes of exposure to the novel tank (Figure 12A). Aged wild-type animals
worsened in performance, with 13.3% (3 out of 23) undergoing a bout of freezing behavior
during first exposure. These numbers remained similar when the entire 10 minutes was
analyzed (wt = 16.7%, mut = 29.2%, Figure 12B, figure supplement 12A). Mutant fish spent a
longer period of time frozen than wild type fish and underwent more total bouts of freezing per
fish (Figure 12C, figure supplement 12B). Upon subsequent exposures to the same tank,
klc4^{uw314} mutant freezing behavior worsened (Figure 12B, figure supplement 12A,B), with 50%
of *klc4*^{uw314} mutants freezing by the sixth exposure, indicating that mutants fail to habituate
freezing behavior. Mutants consistently froze for longer periods of time than wild type animals
throughout all exposures (Figure 12B,C). Wild type animals performed approximately equally
through subsequent exposures, undergoing less frequent freezing bouts and spending less time
frozen than mutants (Figure 12B,C, figure supplement 12A,B). Wild type fish also swam faster
than *klc4*^{uw314} mutants and showed similar swim velocities when their total swim velocity
(including pauses) was compared to their mobile swim velocity (with pauses in swimming
subtracted) (Figure 12D). *klc4*^{uw314} swim mobile swim velocity was faster than their total swim
velocity, reflecting that they spent a greater amount of time immobile (Figure 12D).

481
Interestingly, both wild type and *klc4*[uw314] mutants did show habituation as measured by the
percentage of time spent in the upper zone of the tank (Figure 12E). During the first exposure,
mutant animals only spent 8.7% of the ten minute experiment window in the upper zone and
took on average 170.2 seconds before exploring the upper zone. Wild type animals explored the
upper zone after an average of 80.0 seconds and spent 17.6% of the first exposure period in the
upper zone (Figure 12E,F). At the time of the sixth exposure, however, both wild type and
klc4[uw314] mutants increased their time spent in the upper zone, and mutants entered the upper
zone faster than in their first exposure (Figure 12E,F). Wild type animals took on average 79.7
seconds to explore the upper zone and spent an average of 25.4% of the experiment period in
the upper zone. *klc4*[uw314] mutants performed similarly, spending 23.1% of the experiment period
in the upper zone, but taking on average 129.0 seconds to explore the upper zone (Figure 12E,
F). Thus, the failure of *klc4*[uw314] mutants to habituate freezing behavior is specific rather than
reflecting a general deficit in habituation.
495
496 **Discussion**
497
The association of KLC4 with human disease demonstrates its critical importance, yet its
cellular and developmental functions are poorly understood. In this study, we used automated
analysis of sensory axon arbors and live, *in vivo* imaging of axon behavior and intracellular MT
dynamics to define new roles for KLC4 in shaping neuronal morphology. We found that KLC4 is
required for sensory axon branching and for repulsion between peripheral axons, behaviors
which are important for proper innervation and tiling of the sensory field. *klc4* mutant neurons do
not have a general growth defect, in that their axons grew faster than wild type axons. In
addition, *klc4* mutants are adult viable and showed no signs of early neurodegeneration. These
results suggest that KLC4 has a specific function in patterning developing neurons rather than
simply being required for general neuronal health. We found that mutant larvae were
hypersensitive to touch stimuli, suggesting the altered sensory arborization pattern affects
behaviors mediated by sensory neurons. Behavioral analysis also showed that *klc4* mutant
adults exhibit increased anxiety-like behavior, indicating roles for KLC4 in adult circuit function
and identifying KLC4 as a novel gene associated with anxiety behavior. This work sheds light on
potential mechanisms of human HSPs and provides a new animal model uniquely suited to
further investigation of the cell biological and circuit mechanisms underlying disease.
514
Precise regulation of axon branching is critical for neurons to develop diverse and complex
architecture. We identified KLC4 as a novel regulator of axon branching and show that it is
required for stabilization of nascent branches. Axon branching requires extensive cytoskeletal
rearrangements (Kalil and Dent 2014; Gallo 2011; Winkle et al. 2016). Branches begin as actin-
based filopodial protrusions, which must be invaded by MTs to mature into branches. MT
severing by proteins such as katanin and spastin is required to facilitate invasion (Qiang et al.
2010; Yu, Ahmad, and Baas 1994; Yu et al. 2008) and branch points are hot spots of increased
MT dynamics (Dent and Kalil 2001). Local unbundling of MTs in the axon shaft occurs at sites of
interstitial branch formation (Kalil and Dent 2014; Ketschek et al. 2015). Kinesin-1 can bind MTs
via the KHC C-terminal tail and crosslink or bundle MTs (Seeger and Rice 2010; Navone et al.
1992). Binding of KLC to KHC releases the autoinhibition of KHC and can tip the balance
between kinesin-1's cargo transport activities and the ability of the KHC tail region to bind MTs
(Y.L. Wong and Rice 2010; Cai et al. 2007; Chiba et al. 2021). Our finding that MT dynamics are
affected in *klc4* mutants is consistent with a model in which KLC4 promotes the localized MT
dynamics or severing that allows MTs to invade new branches, either by transporting/localizing
MT regulators or by facilitating MT unbundling. The increase in MT velocity could suggest an
increase in MT sliding or altered regulation or localization of polymerization-promoting MAPs

such as CRMP2 (Nakamura, Ohshima, and Goshima 2020). CRMP2 has been reported to
influence MT polymerization and stabilization and has been found in complex with KLC4,
supporting its potential role in KLC4-mediated effects on MTs (Liz et al. 2014; Koh et al. 2021).
The model of KLC4 as a mediator of microtubule dynamics is also supported by the similarities
in phenotype between *klc4* mutants and kinesore treatment. Kinesore blocks binding between
KLCs and cargo adaptors, and also releases KLC autoinhibition and drives KHC toward
increased MT bundling function (Randall et al. 2017). MTs that invade nascent branches in *klc4*
mutants have less acetylation, suggesting KLC4 may influence post-translational modifications
of MTs and thereby affect stabilization of new branches.
541
In addition to cytoskeletal remodeling, regulated organelle/cargo transport and membrane
remodeling are important for axon branching (Winkle et al. 2016), although little is known about
how individual regulators of trafficking contribute to branching patterns. We previously found
that a properly functioning endosomal trafficking system is necessary for RB axon branching
(Ponomareva et al. 2014; Ponomareva, Eliceiri, and Halloran 2016). Moreover, we showed that
the kinesin adaptor Clstn1 is required for branching of peripheral RB axons and that it functions
in part by trafficking endosomal vesicles to branch points (Ponomareva et al. 2014). KLC4 loss
has a relatively minor effect on Rab5 endosome transport, which is not likely to fully explain the
branch stabilization phenotype. KLC4 could potentially influence transport or localization of
other necessary organelle cargos, such as mitochondria, which must be localized to branch
points to support branch growth (Courchet et al. 2013; Spillane et al. 2013; Tao, Matsuki, and
Koyama 2014; H.H. Wong et al. 2017), or organelles that support mitochondria, such as ER,
which is a critical regulator of mitochondrial function (Csordas, Weaver, and Hajnoczky 2018;
Rowland and Voeltz 2012; Krols, Bultynck, and Janssens 2016). An important future research
goal is to determine whether KLC4 localizes specific organelles to branch points.
557
To maximize coverage of an innervation field, the axonal or dendritic arbors of neuronal
populations that share a similar function must repel one another on contact to "tile" the field
(Grueber and Sagasti 2010; Spead and Poulain 2020). Our results show that KLC4 is an
important player in repulsion between sensory axons, and to our knowledge KLC4 is the first
motor protein shown to be involved in tiling. Several cell surface or secreted proteins that
meditate dendritic tiling have been identified, including protocadherins (Kostadinov and Sanes
2015; Lefebvre et al. 2012), DSCAM (Matthews et al. 2007; Soba et al. 2007), netrin (Smith et
al. 2012), and slit/robo signaling (Gibson et al. 2014). Less is known about the mechanisms of
axon tiling, although similar molecules appear to be involved. For example, in *Drosophila*,
DSCAM mediates sister branch separation in mushroom body axons (Wang et al. 2002), and
Dscam2 mediates repulsion between axons in the visual system (Millard et al. 2007). An
atypical cadherin, Flamingo, is required for spacing of photoreceptor axon arbors (Senti et al.
2003), and functions together with the transmembrane protein Golden goal (Tomasi et al. 2008;
Hakeda-Suzuki et al. 2011). More recent studies have shown that the protocadherin PcdhaC2
mediates repulsion between axons of serotonergic neurons in the mammalian brain (Katori et al.
2017; Chen et al. 2017). Thus, the field has seen multiple advances in knowledge of the
membrane proteins that mediate repulsion. However, the mechanisms that regulate their
localization to the correct cellular compartments are unknown. Our work can drive new
exploration into such mechanisms. For example, it will be interesting to determine whether
KLC4 mediates transport of contact repulsion molecules and their specific targeting to
peripheral versus central sensory axons.
579
The formation of polarized neuronal structure is a critical component of proper neural circuit
development. An important early step in neuronal polarity development is determination of the
location of axon emergence from the cell body. The ectopic apical axons we see in *klc4* mutants

suggest that KLC4 also has roles in the polarity of axon initiation location. RB neuron axons
normally emerge from basal edge of cell body (E. Andersen, Asuri, and Halloran 2011), and
their emergence correlates with centrosome movement to the basal region of the cell (E.F.
Andersen and Halloran 2012). We previously found that *laminin-1* mutants also have ectopic
apical axons in RB neurons and these ectopic axons correlate with apically displaced
centrosomes (E.F. Andersen and Halloran 2012). Laminin has been shown to influence the
location of axon emergence from the cell body in zebrafish RGCs (Randlett et al. 2011), and
can direct the positioning of the centrosome in neurons *in vitro* and *in vivo* (Randlett et al. 2011;
Gupta et al. 2010). KLC4 may play a role in coordinating the response to extracellular signals
such as laminin that help define site of axon formation.
593
Overall, the developmental phenotypes seen in *klc4^{uw314}* mutants are striking in that they appear
to reflect problems with specific patterning of neuronal structure rather than problems in bulk
cargo transport. Neurons appear healthy in the mutants, show no signs of axon degeneration
during development, and in fact axons grow faster in mutants than in wild type embryos.
Moreover, the *klc4^{uw314}* mutants are adult viable. Interestingly, *klc1* knockout mice are also adult
viable and show specific defects in transport of cannabinoid receptors in developing axons
(Saez et al. 2020). Thus our findings may reflect a general theme, that the function of the KLC
subunits of the kinesin-1 motor are to provide specificity to the process of localizing cargos to
particular cell regions. Our results also give insight into mechanisms of the human disease
caused by *KLC4* mutation, which may be caused by defects in developmental patterning. It will
be important to fully define the unique cellular functions of individual KLCs in order to better
understand mechanisms of disease.
606
Adult circuit function can be affected by both defects in circuit development and/or later defects
in the function of neurons within circuits. Our finding that *klc4^{uw314}* mutant adults show freezing
and reduced exploration in the novel tank assay implicates KLC4 in the circuits that regulate
stress response and that are affected in anxiety and depression disorders. The hypothalamic-
pituitary-adrenal (HPA) axis is an important system that mediates stress responses, and
zebrafish mutants with disruptions to the HPA also exhibit anxiety behavior in the novel tank
test. Mutation of zebrafish glucocorticoid receptor, a target of the stress hormone cortisol,
results in increased freezing and decreased exploratory behavior. Like *klc4^{uw314}* mutants,
glucocorticoid receptor mutants fail to habituate freezing behavior after repeated exposures to
the novel tank (Ziv et al. 2013). However, unlike *klc4^{uw314}* mutants, glucocorticoid receptor
mutants do not habituate exploratory behavior over multiple exposures. This suggests that
freezing behavior and exploratory behavior are controlled by different mechanisms.
Interestingly, zebrafish with mutation in a chemokine gene expressed in multiple brain regions,
sam2, freeze during the novel tank test, but show normal tank exploration (Choi et al. 2018),
providing additional evidence that these two behaviors can be uncoupled. The *sam2* mutants
have disruptions in corticotropin releasing hormone neurotransmission, linking the mutation to
HPA disruption, but not excluding contribution from other circuits. Our *klc4^{uw314}* mutants provide
another tool to further investigate the circuit changes underlying anxiety disorders.
625
Whether developmental defects in RB neurons contribute to adult circuit function is unknown.
RB neurons are traditionally thought to be a transient cell population that undergoes
programmed cell death during larval stages. However recent evidence has shown that 40% of
RBs are maintained for at least the first 2 weeks of development, suggesting they may be
retained (Williams and Ribera 2020). Moreover, RB neurons have been shown to play
significant roles in regulating larval behaviors (Knafo et al. 2017). Hypersensitivity to touch is a
common trait of autism spectrum disorder (ASD), although the relationship of somatosensory
neurons to states like anxiety, depression, and hyperstimulation is not well defined.

Interestingly, deletion of the ASD risk genes Mecp2 and Gabrb3 in mouse primary
somatosensory neurons during development led to a reduction in exploratory behavior in the
open field test, while deletion of these genes only during adulthood caused no such impairment
or anxiety behavior (Orefice et al. 2016). In addition to sensory neurons, *klc4* is expressed in
many cell types in the developing nervous system and thus likely has important roles in
development and function of other neurons, which may include neurons that are directly
involved in circuits mediating adult behavior. A future research goal is to test the functions of
KLC4 in neurons of stress response circuits, such as the HPA axis.
642
643   **Methods**
644

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
gene (*Danio rerio*)	klc4	Ensembl	ENSDARG 00000008 69 85	
strain, strain background (*Danio rerio*)	AB	ZIRC	ZL1	Wildtype Strain of Zebrafish
strain, strain background (*Danio rerio*)	TL	ZIRC	ZL86	Wildtype Strain of Zebrafish
genetic reagent (*Danio rerio*)	Tg(-3.1ngn1:gfp-caax)	Andersen et al. 2011	Tg(ngn:gfp-caax)	Transgenic line labelling sensory axons, AB/TL background
Recombinant DNA reagent	3.1ngn1:trfp-caax	Andersen et al. 2011	Tg(ngn:trfp-caax)	Plasmid for labelling sensory axons
Recombinant DNA reagent	3.1ngn1-mcherry-caax	Andersen et al. 2011	Tg(ngn:mcherry-caax)	Plasmid for labelling sensory axons

Recombinant DNA reagent	3.1ngn1:eb3-caax	Stepanova et al. 2003, Ponomareva et al. 2014	Tg(ngn:eb3-gfp)	Plasmid for labelling microtubule plus ends
Recombinant DNA reagent	3.1ngn1:GFP-rab5c	Clark et al. 2011, Ponomareva et al 2014	Tg(ngn:GFP-rab5c)	Plasmid for labelling Rab5c vesicles
Recombinant DNA reagent	3.1ngn1:PA-gfp-Rab5c	Ponomareva et al. 2014		Plasmid for labelling Rab5c vesicles
Sequence-based reagent	klc4 *in situ* probe	This paper		Sequences listed in methods
Sequence-based reagent	klc4 long isoform specific *in situ* probe	This paper		Sequences listed in methods
Sequence-based reagent	short isoform RT primers	This paper		Sequences listed in methods
Sequence-based reagent	long isoform RT primers	This paper		Sequences listed in methods
Sequence-based reagent	klc4 gRNA	This paper		Sequence listed in methods
Commercial reagent	Cas9 protein with NLS	PNA Bio	NC0789474	

Commercial Reagent	Trizol	Thermo-Fisher	15596026	For RNA extraction
Chemical compound, drug	Kinesore	Chembridge Hit2Lead library	#6233307	Kinesin1 targeting small molecule
Commercial assay or kit	Superscript III	Invitrogen	18080093	
Commercial assay or kit	DIG RNA labelling kit	Roche	11175025910	
Commercial Reagent	Oligo dT	Invitrogen	18418020	
Commercial Reagent	Random Hexamers	Invitrogen	N8080127	
antibody	anti-HNK1 (mouse monoclonal)	ZIRC	ZN-12	IF(1:250)
antibody	anti-acetylated tubulin (Mouse monoclonal)	Sigma	T6793	IF(1:1000)
antibody	anti-mouse IgG AlexaFluor 488	Invitrogen	A-21121	IF(1:1000)

antibody	anti-mouse IgG2b AlexaFluor 647	Invitrogen	A-21141	IF(1:1000)
Software, algorithm	CRISPRscan	Moreno-Mateos et al. 2015		https://www.crisprscan.org/
Software, algorithm	CRISP-ID	Dehairs et al. 2016		http://crispid.gbiomed.kuleuven.be/
Software, algorithm	ToxTrac	Rodriguez et al. 2018		Animal Tracking Freeware
Software, algorithm	OrientationJ	Rezakhaniha et al. 2011		Used to quantify orientation of axons
Software, algorithm	Graphpad Prism			Graph making software
Software, algorithm	Chemotaxis Tool	Ibidi		https://ibidi.com/chemotaxis-analysis/171-chemotaxis-and-migration-tool.html

Animals

Adult zebrafish (Danio rerio) were kept in a 14/10 h light/dark cycle. Embryos were maintained at 28.5°C and staged as hours postfertilization (hpf) as described previously (Kimmel et al. 1995). Wild type AB strain or transgenic Tg(−3.1ngn1:GFP-CAAX) (E. Andersen, Asuri, and Halloran 2011) (AB/TL background) embryos were used for all experiments. In experiments using the *Tg(−3.1ngn1:GFP-CAAX)* line as a background, wild type is used to refer to *Tg(−3.1ngn1:GFP-CAAX), klc4$^{+/+}$* animals. All animals in these studies were handled in accordance with the National Institutes of Health Guide for the care and use of laboratory animals, and the University of Wisconsin Institutional Animal Care and Use Committee (IACUC). These studies were approved by the University of Wisconsin IACUC.

CRISPR/Cas9 mutagenesis

A guide RNA targeting zebrafish *klc4* (5'-GAGCGTGGCACAGCTGGAGGAGG-3') was designed by using CRISPRScan (crisprscan.org) and synthesized as previously described (Shah et al. 2015; Moreno-Mateos et al. 2015). To generate *klc4^{uw314}* mutants, embryos at the

very early 1 cell stage were injected with 374.5 ng/µl of the guide RNA targeted to zebrafish *klc4*
complexed with 800 ng/µl of Cas9 protein with NLS (PNA Bio). Injection volume was
approximately 1nL. F0 adult fish were pair-mated to determine individuals carrying germ line
mutations. A subset of individuals with germ line mutations were outcrossed to wild type AB fish.
F1 fish were raised and genotyped by PCR and sequencing, using CRISP-ID
(http://crispid.gbiomed.kuleuven.be/) to assist in sequence unmixing (Dehairs et al. 2016). The
klc4^{uw314} allele was found to have a 25 bp deletion. Fish heterozygous for *klc4^{uw314}* were
outcrossed to wild type AB fish until the F3 generation. F3 *klc4^{uw314}* heterozygotes were in-
crossed to generate F4 *klc4^{uw314}* homozygotes. All homozygous mutants in this study were F5-
F6 generation mutants from homozygous parents, unless otherwise specified. Adults and
embryos described as "wild type cousins" of *klc4^{uw314}* mutants were generated by crossing F4
klc4$^{+/+}$ siblings from the F3 *klc4$^{uw314/+}$* in-cross.
673
RT-PCR
RNA was extracted from zebrafish embryos at specified ages with Trizol (Thermo-Fisher),
followed by chloroform extraction and isopropanol precipitation. cDNA was generated from
embryo-derived RNA using the Superscript III reverse transcriptase kit and a 50:50 mixture of
oligo dT and random hexamer primers (Invitrogen). We designed PCR primers to amplify
specific *klc4* isoforms. A reverse primer unique to the short isoform's 3' UTR was designed for
the *klc4* short isoform, with the forward primer in exon 2 (Fwd: 5'-
GCGCAAATCAGTGGAGATGATAGAGC-3', Rev: 5'-GCATAGTGGCGCGGTAGGTA-3'). A set
of primers for exons exclusive to the long isoform of *klc4* (spanning exons 7 through 12) was
designed for the long isoform (Fwd: 5'- CAGAGAGAAGGTGCTGGG -3', Rev: 5'-
GGTGGTGTTCACTGTTGG-3'). PCR was then run to test for presence of the isoforms in cDNA
from each embryonic age.
686
in situ **hybridization**
A digoxigenin-labeled riboprobe complementary to a region spanning exons 2-3 of the *klc4*
mRNA, which are contained in both long and short isoform mRNAs, was synthesized by in vitro
transcription using T7 RNA polymerase with a DIG RNA labeling kit (Roche). The template for
transcription was made using PCR on zebrafish cDNA with the following primers: Fwd: 5'-
GCGCAAATCAGTGGAGATGATAGAGC-3', Rev: 5'-
TAATACGACTCACTATAGCTCCAGATGTTTCTTCTCCTCC-3'. A second probe recognizing a
region in exons 8-11, found only in the long isoform, was synthesized using PCR on zebrafish
cDNA with the following primers: Fwd: 5'-CAG AGA GAA GGT GCT GGG-3', Rev: 5'-
TAATACGACTCACTATAG GGTGGTGTTCACTGTTGG-3'. Whole-mount in situ hybridization
was performed as previously described (Halloran et al. 1999).
698
Immunohistochemistry
Embryos were fixed overnight at room temperature in 4% paraformaldehyde in 0.1M PO$_4$ buffer,
then rinsed with PBS/0.1% Tween-20 (PBST) and blocked with incubation buffer (IB: 2mg/mL
BSA, 1% Sheep Serum, 1% DMSO, 0.5% Triton X-100 in PBS) for at least 1 hour. Embryos
were incubated overnight at 4°C with monoclonal anti-HNK1 antibody in IB (ZN-12, 1:250;
Zebrafish International Resource Center), rinsed in PBST, then incubated with anti-mouse
AlexaFluor 488 secondary antibody (1:1000; Invitrogen) for at least 2 hours at room temperature
or overnight at 4°C. For double labeling with anti-acetylated tubulin (T6793, 1:1000, Sigma), the
IgG subtype-specific secondary antibodies anti-mouse IgG1 AlexaFluor 488 and anti-mouse
IgG2b AlexaFluor 647 (1:1000, Invitrogen) were used.
709
Microinjection and Mosaic Expression

DNA expression constructs were made using the Multisite Gateway Cloning System (Invitrogen)
into Tol2 vectors (Kwan et al. 2007). Zebrafish embryos at the 1-cell stage were injected with
~10 pg -*3.1ngn1:EB3-GFP* DNA (Lee et al. 2017) to visualize microtubule dynamics. For
experiments to image individual neuron morphology, embryos were injected with ~12pg
3.1ngn1:mCherry-CAAX DNA (E. Andersen, Asuri, and Halloran 2011).
716
717 **Kinesore treatment**
718 Kinesore was acquired from the Hit-2-Lead library (Chembridge, compound #6233307) and
719 suspended in DMSO for a stock concentration of 100 mM. Embryos treated with Kinesore were
720 dechorionated at 17 hpf and placed in E3 containing 150 µM Kinesore and 2% DMSO. Controls
721 were treated with 2% DMSO in E3 for the same time period.
722
Live Imaging
Light sheet imaging was performed on a custom multi-view light sheet microscope as described
previously (Daetwyler et al. 2019). The detection objective was either an Olympus UMPLFLN
10x W (NA 0.3) or a Nikon CFI75 Apochromat 25xC W (NA 1.1) with a 400mm tube lens
(TL400-A) for 50x total magnification. Briefly, the Tg(*3.1ngn1:GFP-CAAX*) wild type or mutant
embryos were mounted in an FEP tube (0.8□mm inner diameter, 1.2 outer diameter, Bola)
suspended in 0.02% tricaine in E3 with an agarose plug to prevent leakage.
730
For live confocal imaging, embryos were anesthetized in 0.02% tricaine and mounted in 1% low
melting agarose in 10 mM HEPES E3 medium as described previously (Andersen et al., 2010).
Axon outgrowth was imaged in Tg(*3.1ngn1:GFP-CAAX*) embryos with either an Olympus
FV1000 laser-scanning confocal microscope using a UPlanSApo 20x Oil objective (NA 0.85) or
an Opterra Swept-Field confocal (Bruker Fluorescence Microscopy) with a Nikon S Fluor 40x Oil
immersion objective (NA 1.30). Imaging was started between 18 and 19 hpf, when peripheral
RB axons are extending out of the spinal cord. Optical sections (1-1.25 µm) covering a 30–36
µm Z depth were captured every 60-90 s for 2-4 hours.
739
High speed imaging of EB3-GFP comets was performed with the Opterra Swept-Field confocal
microscope equipped with a Nikon Plan Apo VC 100x oil immersion objective (NA 1.40).
Embryos were imaged at stages between 18 and 26 hpf, during peripheral axon outgrowth.
EB3-GFP comets in nascent branches were imaged at stages between 22 and 25 hpf with a
Nikon Plan Apo 60x oil immersion objective (NA 1.40). Z-stacks of 5–30 1-µm optical sections
were captured at 2–5 s time intervals, for total durations between 7 and 10 min. High speed
imaging of Rab5 vesicle dynamics was done with the Opterra Swept-Field confocal and the
Nikon Plan Apo 100x oil objective (NA 1.40). Embryos were imaged between 22 and 26 hpf. Z-
stacks of 5-30 µm optical sections were captured at 2-5 s time intervals for a total duration of 7-
10 min. Photoactivation and imaging of PA-GFP-Rab5 were performed on the Olympus FV1000
confocal using a 60X UPlanSApo oil immersion objective (NA 1.35). A region of interest
encompassing the cell body was activated with the 405 nm laser during the scan of one Z-plane
in the cell body. Z-stacks of 1 µm optical sections were captured every 30-170 s for durations
ranging from 26-34 minutes (average 29.22 minutes, median 28.84 minutes).
754
Axon Outgrowth Analysis
To measure the speed of axon outgrowth, the area immediately behind the growth cone was
measured using the Manual Tracking plugin in FIJI (Fabrice Cordelières). The generated data
was loaded into the Chemotaxis and Migration Tool (Ibidi, https://ibidi.com/chemotaxis-
analysis/171-chemotaxis-and-migration-tool.html) to generate velocity measurements and rose
plot histograms. Rose plots were re-plotted in MATLAB (MathWorks) using secplot (Matthew
Foster, https://www.mathworks.com/matlabcentral/fileexchange/14174-secplot).

Branch Analysis

Custom macros for FIJI (Schindelin et al. 2012) were written to automate branch analysis. Briefly, stacks were processed to be maximum intensity projections and oriented with anterior to the left. A rectangular ROI was selected over the posterior region of the yolk tube extension such that the top of the selection was aligned with the bottom of the central RB axons. A tubeness filter was applied to reduce non-axonal background. A histogram of axon directionality was generated for the ROI using OrientationJ (Rezakhaniha et al. 2011). To measure differences in axon directional distribution, we computed the area under the curve (AUC) for the regions to be compared. We then performed a t-test using the resulting mean and standard error to determine significance. To measure axon branch density, eight ROIs measuring 3-μm in height were generated at regularly spaced intervals from dorsal to ventral (Figure 2 – figure supplement 2). These selections were individually thresholded, binarized, and the number of objects contained in each ROI was determined using the Analyze Particles function. To analyze acetylated tubulin signal in nascent branches, terminal branches up to 10 μm in length were scored based on the presence or absence of visible acetylated tubulin signal. Total proportion of nascent branches with signal was compared between wild type and mutant embryos.

Kymography

Hyperstacks of EB3-GFP or GFP-Rab5c images were assembled using FIJI (Schindelin et al. 2012) and segments of axons were traced for kymography using the segmented line tool. For peripheral axons, a segment was defined as an area of axon between two branch nodes, or between the first visible portion of axon and the first node of branching. Growth cones and filopodia were excluded from analysis. For analysis of EB3-GFP in nascent branches, only branches <10 μm long were analyzed. For unbranched central axons, a segment was defined as the entire visible length of either the ascending or descending central axon within each field of view. Kymographs were made using the Multi Kymograph plugin (developed by J. Rietdorf and A. Seitz, European Molecular Biology Laboratory, 2004). Comet or vesicle parameters were measured from the resulting kymographs. To measure Rab5 accumulation at branch points, movies were divided into 4 time periods (t1-t4), each containing 7-8 minutes. Maximum intensity projections were made for t1-t4 and the fluorescent signal at branch points was measured for each maximum intensity projection. Background signal from the branch point prior to activation was subtracted from measured t1-t4 values, and then all values were normalized to the signal at the t1 period. Comparisons between experimental conditions were made using the Mann-Whitney test for non-parametric data.

Adult Body Size Measurements

Zebrafish were raised for this experiment by in-crossing wildtype or $klc4^{uw314}$ homozygous siblings. Larva were placed in 3L tanks side-by-side and subjected to the same light cycle and feeding schedule during rearing. At 7 weeks and 9 weeks, fish were briefly anesthetized with 0.02% tricaine, placed on white waterproof surface, measured with a ruler, and photographed.

Larval Touch Assay

To assess responsiveness to touch, 3 dpf zebrafish larva were individually placed in a 6 cm dish and allowed to acclimate for 2 minutes. A tail touch was administered to the end of the tail using a gel loading tip attached to a long applicator stick. The tail touch was given from a side angle to avoid looming stimulus. A high-speed camera was used to acquire 50 frames per second for a total of 1050 frames (21 sec). The response time of each larva was measured from the first movement after a touch to the cessation of movement for more than 15 frames (inter-bout pauses only lasted ~7-8 frames). Some wild type and *klc4* mutant larva did not respond to an initial touch (10 wild type, 14 mutant), and these samples were excluded from analysis.

813
Feeding Experiment
Feeding experiments were conducted as in Howe et al., 2018. Briefly, adult zebrafish were isolated in individual 1L tanks without food for 48 hours, then 50 larval zebrafish were added to each tank. The fish were allowed to feed on the larvae for one hour, then removed from the tank to allow counting of the remaining larvae. All tested animals were of AB background.

Novel Tank Diving Test
The evening before an experiment, fish were moved to the behavior experiment room and allowed to acclimate overnight. Exposure to novel tank was carried out in the morning, and the order of which genotype received exposure first was randomized. The novel tank had 3 opaque sides to block potential views of other fish and one clear side for imaging. The water in the novel tank was changed between groups. Fish were netted from their 3L home tank, released into the novel tank, and allowed to explore for 3 minutes before being removed and placed into a recovery tank. All tested animals were of AB background.

Habituation Experiment
Two weeks prior to an experiment, adults were sorted into separate 3L tanks with 5-6 individuals of the same genotype in each. Experiment was conducted in a similar manner as described above, but fish were allowed to explore for 10 minutes before removal. Novel tank exposure was repeated on the same groups twice a week for 3 weeks. Exposures were conducted a minimum of two days apart. The experiment was repeated for another 3 weeks with an additional two groups of 6 fish per genotype. All tested animals were of AB background. When possible, groups were divided into similar ratios of male and female fish for a total of 11 females and 12 males within each genotype. The group compositions were as follows: **wild type**: group 1 (3 females [F], 2 males [M]) group 2, (3 F, 3 M), group 3 (3 F, 3 M), group 4 (2 F, 4 M), *klc4^{uw314}*: group 1 (2 F, 3 M), group 2, (3 F, 3 M), group 3 (3 F, 3 M), group 4 (3 F, 3 M).

Automated Tracking of Adults
For the habituation experiment, automated tracking of zebrafish through ToxTrac (Rodriguez et al. 2018) was used to quantify swim velocity and percent of time spent in the upper zone. When necessary for accurate tracking, background subtraction was performed on movies prior to using ToxTrac. This was accomplished by subtracting the first frame of each movie (showing only the empty tank) from subsequent frames in the movie via FIJI (Schindelin et al. 2012). Operations were performed using Clij2 where possible to speed up processing time (Haase et al. 2019).

Statistics
Statistical analyses and graphs were generated with Graphpad Prism. See figure legends for details on sample sizes, specific statistical tests used, error bars and p values. Data were checked for normality using Anderson-Darling test, D'Agostino & Pearson test, and Shapiro-Wilk test before deciding on a statistical test. Non-normal data were tested for significance using a Mann-Whitney test. In some cases, the N values were too low to perform a statistical test (Fig. 6E, Fig. 11E,H, Fig. 12A). In Fig. 6E, the N for wild type is low because fasciculation events are very rare in wild type and obtaining a value for fasciculation length is contingent on the event occurring. For the behavioral data shown in Fig. 11E,H and Fig. 12A, multiple animals were tested in each trial, but N is the number of individual behavioral trial days (N=3, which is too low to test for normality). In all these cases, individual data points are shown for comparison. Data tested using the Mann-Whitney test were independent of each other and comprised ordinal or continuous numbers.

Acknowledgements

The authors would like to thank the members of the Huisken lab for technical expertise and intellectual contributions, particularly Kurt Weiss, Joe Li, and Alyssa Graves. We are grateful to Katie Drerup and Bill Bement for critical reading and feedback on this manuscript, which greatly improved the final work. We also give our heartfelt thanks to the current and former members of the Halloran lab for their support of and contributions to this work, especially Conlin Bass for valuable work during the initial characterization of *klc4* expression pattern, and Daniel North for fish care and maintenance.

Funding

This work was supported by funds to Mary Halloran from the National Institutes of Health (R01 NS086934, R21 NS116326) and funds to Elizabeth Haynes from the National Institutes of Health (F32 NS098689) and the Morgridge Institute for Research. The Morgridge Institute for Research provided support for the Huisken lab and the light sheet microscope. The Center for Quantitative Cell Imaging provided support for the Opterra Confocal.

Ethics Statement

This study was performed in accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. Animals were handled according to approved institutional animal care and use committee protocols of the University of Wisconsin.

Competing interests

K. Eliceiri is a consultant for Bruker, the manufacturer of the Opterra Swept Field Confocal microscope used in this work.

Figure 1. *klc4* mRNA is expressed in developing zebrafish embryos. **A)** Diagram of the highly conserved domains of KLC4. Arrows indicate where mutation occurs in humans and in our zebrafish mutant. **B)** Qualitative RT-PCR showing both long and short *klc4* isoforms are expressed throughout development. **C-D)** *In situ* hybridization for *klc4* mRNA shown in lateral view (**C**) or dorsal view (**D**) of the head in wild type 24 hpf embryos. **E)** Lateral view of trunk showing *klc4* expression in Rohon-Beard sensory neurons in a 24 hpf wild type embryo. **F-G)** *klc4* expression in the lateral view (**F**) or dorsal view (**G**) of the head of a 48 hpf wild type embryo. **H)** Lateral view of trunk showing *klc4* expression in the spinal cord of a 48 hpf wild type embryo. **I-K)** Lateral (**I,K**) and dorsal (**J**) views of *klc4* long isoform specific *in situ* hybridization in wild type 24 hpf embryos. Lateral (**L,N**) and dorsal (**M**) views of *klc4 in situ* hybridization in 24 hpf *klc4*^uw314^ mutant embryo head (**L,M**) and trunk (**N**) showing little expression of *klc4*. te = telencephalon, dien = diencephalon, mb = midbrain, hb = hindbrain, tg = trigeminal ganglion, pllg = posterior lateral line ganglion, rb = rohon beard neuron.

Figure 2. *klc4*^uw314^ mutants have reduced Rohon-Beard peripheral axon branching. **A-B)** Stills taken at approximately 24 hpf from long term light sheet movies of wild type (**A**) or *klc4*^uw314^ mutant (**B**) embryo development. **C)** Diagram of a 24 hpf embryo indicating the area over the yolk tube extension imaged for axon analysis, with the standard region

used for analysis outlined by the white dashed rectangle. **D-E)** Representative
examples of HNK-1 stained embryos showing peripheral axon branching in wild type (**D**)
and *klc4^{uw314}* (**E**) embryos. **F-G)** Quantification of axon branching across three technical
replicates in wild type (n = 32) and *klc4^{uw314}* mutant (n=32) embryos (**F**) and DMSO (n =
12) and kinesore (n = 17) treated embryos (**G**). Error bars = SEM. Statistical
significance was measured by comparing the area under the curve for **F-G** using t-test.
For **F**, *p=0.0378 for entire graph. For **G**, *p=0.0484 for shaded region. **H-I)**
Representative examples of HNK-1 stained embryos showing peripheral axon
branching in DMSO (**H**) and kinesore treated (**I**) embryos. All scale bars = 50 µm.

Figure 3. Branches initiate but do not stabilize in *klc4^{uw314}* mutants. **A)** A wild type axon
forms stable branches by growth cone bifurcation (GCB, white arrows) and interstitial
branching (IB, magenta arrows) over the course of 38' minutes. **B)** A *klc4^{uw314}* mutant
axon initiates a branch via GCB (0 minutes, white arrow), but one branch begins to
retract and is nearly fully gone after 66' minutes (panel 3, white arrow). 20 µm scale bar.
C) Quantification of axon branch initiations per neuron per hour in wild type and mutant
embryos (wild type GCB mean = 2.22 branches/neuron/hr, N = 11 neurons from 4
embryos. *klc4^{uw314}* GCB = 2.13 branches/neuron/hr, N = 14 neurons from 4 embryos.
wild type IB = 0.99 branches/neuron/hr, N = 10 neurons from 4 embryos. *klc4^{uw314}* IB =
0.65 branches/neuron/hr, N = 8 neurons from 4 embryos). Error bars = SEM. There was
no significant difference in branch initiation between wild type and mutant axons. GCB p
= 0.73, IB p = 0.15, Mann–Whitney test. **D)** Quantification of branches that retracted
after initiation in wild type and *klc4^{uw314}* embryos. Data are displayed as a branch loss
ratio (branches retracted/total branching events). Branches created by either GCB or IB
were both more likely to retract in *klc4^{uw314}* embryos (wild type GCB mean = 0.13, N =
11 neurons from 4 embryos; *klc4^{uw314}* GCB = 0.58, N = 14 neurons from 4 embryos; wild
type IB = 0.10, N = 10 neurons from 4 embryos; *klc4^{uw314}* IB = 0.62, N = 8 neurons from
4 embryos). **p=0.0055, ****p<0.0001, Mann–Whitney test. Error bars = SEM. **E)**
Quantification of axon growth velocity. Axons in *klc4^{uw314}* mutants grow faster than in
wild type. Wild type N = 19 axons from 3 embryos. *klc4^{uw314}* N = 22 axons from 4
embryos. ***p=0.0001, Mann-Whitney test. Error bars = SEM.

Figure 4. Axon orientation profiling reveals increased posterior directionality of
peripheral axons in *klc4^{uw314}* mutants. **A-B)** A representative region of axon analysis for
a 24 hpf wild type (**A**) and *klc4^{uw314}* (**B**) embryo. The axons are color coded for
directionality according to the legend in the upper left corner of (**A**). **C)** Plot of axon
directionality in 24 hpf wild type (green) and *klc4^{uw314}* mutants (magenta). **D)** Axon
directionality data was normalized to total signal and posterior angles are displayed to
compare directional bias. Area under the curve was measured for shaded area and
tested for statistical significance using a t-test. **p = 0.0047. For **C-D**, error bars = 95%
CI. Wild type N = 40 embryos, *klc4^{uw314}* N = 23 embryos.

Figure 5. Live imaging shows that *klc4^{uw314}* mutants have a bias toward posterior
growth direction and defective sister branch separation. **A)** Illustration of how axon
growth direction was measured. The angle between an axon's position in each frame
and the frame preceding it was measured. **B-C)** These angles were plotted as rose

plots to show growth direction distributions. Wild type axons grow evenly between
anterior and posterior directions, while *klc4uw314* mutant axons show a preference for
posterior directed growth. Wild type N = 19 axons from 3 embryos. *klc4uw314* N = 22
axons from 4 embryos. **D)** Stills from a live capture of wild type axon outgrowth,
showing that axons often grow posteriorly, but anterior directed branches form over time
(**D',** white arrows). **E)** As a branch forms in a wild type peripheral axon, the angle
between sister branches remains consistent as axons grow apart. **F)** In *klc4uw314*
mutants, sister branches do not grow as far apart and the sister branch angle narrows
over time compared to wild type. All scale bars = 20 μm. **G)** Quantification of the angle
between newly formed sister branches at 15 and 75 minutes after branch creation. Wild
type 15' N = 37 branches, *klc4uw314* 15' N = 36 branches, Wild type 75' N = 25 branches,
klc4uw314 75' N = 18 branches. 4 embryos per genotype. *p=0.021, ***p=0.0003, One-
way ANOVA. Error bars = 5-95th percentile.
970
Figure 6. Loss of KLC4 results in aberrant axon fasciculation. **A)** In wild type embryos,
the most common response to self or non-self contact between axons is retraction. Two
neighboring axons in a wild type embryo make contact (time 0') and within 30 minutes,
one of the axons has retracted. **B-C)** When axons make contact in *klc4uw314* embryos,
they often fasciculate instead of retracting. This happens with both self-contacts (**B-C,**
sf) and non-self-contacts (**C**, nsf). All scale bars = 20 μm. **D)** Fasciculations per neuron,
separated by self or non-self, as well as all fasciculations combined. Wild type N = 35
neurons in 4 embryos. *klc4uw314* N = 37 neurons in 5 embryos. Error bars = SEM. **E)**
Fasciculation length was measured from the last frame of each 2 hour imaging period.
Wild type N = 3 fasciculations, *klc4uw314* N = 19 fasciculations. Mean with SEM displayed
over individual data points. Because fasciculation events are rare in wild type embryos
and a length measurement is conditional on the event occurring, we could not apply a
statistical test to assess the significance of the difference in fasciculation length.
984
Figure 7. Midline crossing of apical protrusions occurs more frequently when KLC
function is disrupted. **A-B)** Confocal projections (dorsal views) of partial z-stacks
containing only the optical sections in the plane of the RB cell bodies within the spinal
cord in a wild type Tg*(-3.1ngn1:gfp-caax)* embryo **(A)**, or a *klc4uw314* Tg*(-3.1ngn1:gfp-*
caax) embryo **(B),** with an apical protrusion from a neuron crossing the midline and
reaching the contralateral side (white arrowhead). Scale bars = 20 μm. **C)** The
percentage of neurons per embryo that had an apical protrusion crossing the midline of
the spinal cord was quantified. **p<0.0052, Mann–Whitney test. Mean with SEM
displayed over individual data points. Wild type N = 608 neurons from 21 embryos,
klc4uw314 N = 1496 neurons from 47 embryos, 2% DMSO N = 429 neurons from 15
embryos, Kinesore N = 595 neurons from 21 embryos. **D-F)** Mosaic labeling of
individual neurons by injection with -3.1ngn1:gfp-caax DNA into *klc4uw314* mutants (**D**),
DMSO-treated embryos (**E**) or kinesore-treated embryos (**F**) showing that apical
protrusions (magenta arrowheads) can fasciculate onto the contralateral central axon.
999
1000 **Figure 8.** Endosomal trafficking dynamics are disrupted in *klc4314* mutants. **A)** Live RB
1001 neuron expressing eGFP-Rab5c. Scale bar = 10μm. **A')** Representative kymograph of
1002 axon segment boxed in magenta in (**A**). **B)** Percentage of anterogradely moving

vesicles out of all motile Rab5 vesicles in central and peripheral axons. The proportion of anterogradely moving vesicles is decreased in $klc4^{314}$ mutants, **p=0.0087 by Mann-Whitney test; error bars = mean with SD. Wild type N = 18 embryos, $klc4^{314}$ N = 20 embryos. **C)** Average Rab5 vesicle run length and maximum anterograde vesicle run length in peripheral axons. *p=0.0358, ns p=0.5359, Mann-Whitney test. **D)** Rab5 vesicle velocity in peripheral axons. NS by Mann-Whitney test (anterograde p = 0.6505, retrograde p = 0.0792), wild type N=12 embryos, $klc4^{314}$ N=11 embryos. For all data, one neuron per embryo was analyzed. Error bars in box and whisker plots represent 5-95th percentile.

Figure 9. Microtubule dynamics are increased in $klc4^{uw314}$ mutants. **A-A")** Live RB neuron expressing EB3-GFP. Magenta box in (**A**) indicates axon segment selected for kymograph in (**A'**). **A")** EB3 comets identified as anterogradely (green) or retrogradely (magenta) directed based on their angle. **B)** Representative kymographs from peripheral axon segments of wildtype (top) or $klc4^{uw314}$ (bottom). **C)** Quantification of the percentage of retrograde comets out of all comets imaged for both peripheral and central axons. NS by Mann-Whitney test. **D)** Peripheral axon comet frequency. ***p=0.0003, Mann-Whitney test. **E)** Peripheral axon comet velocity. ***p=0.0004, *p=0.042, Mann-Whitney test. For all data, error bars = SEM. Wild type central N = 8 segments, $klc4^{uw314}$ central N = 15 segments, Wild type peripheral N = 26 segments, $klc4^{uw314}$ peripheral N = 48 segments. Wild type data from 10 neurons in 7 embryos. $Klc4^{uw314}$ data from 13 neurons in 8 embryos. All comets in a kymograph were measured.

Figure 10. Nascent RB axon branches show reduced microtubule stabilization in $klc4uw^{314}$ mutants. **A)** Still image of nascent RB branches expressing EB3-GFP in a live wild type embryo at 23hpf. **A')** Zoom in of the branches outlined in white in (**A**). **A")** EB3-GFP signal in nascent branches in (**A'**). **A''')** Representative kymograph generated from nascent branch outlined in magenta in (**A'**). **B)** Quantification of peripheral axon comet frequency. NS by Mann-Whitney test, p=0.8410. Wild type N = 22 nascent branches from 9 embryos, $klc4^{314}$ N = 25 nascent branches from 8 embryos. **C-D")** Representative examples of HNK-1 and acetylated tubulin stained embryos at 24hpf. White arrows in (**C'-C''**) indicate nascent branches with tubulin acetylation. Magenta arrows in (**D'-D''**) indicate nascent branches lacking tubulin acetylation. **E)** Percentage of nascent branches containing acetylated tubulin out of all nascent branches. Fewer nascent branches contained acetylated tubulin in $klc4^{314}$ mutants. ****p<0.0001 by Mann-Whitney test. Wild type N=132 nascent branches from 15 embryos, and $klc4^{314}$ N=143 nascent branches from 16 embryos. Scale bars = 20μm Error bars in box and whisker plots represent 5-95th percentile.

Figure 11. Larval and adult $klc4^{uw314}$ mutants have behavioral defects **A)** Example of touch assay administration**.** A 3 dpf larva is touched on the tail using a gel loading tip, provoking a swim response. **B)** Larval response time was measured starting from beginning of swim to cessation of motion. **C)** The number of swim bouts performed by each larva during total post-touch swim time was counted. For B-C, each point represents an individual larva. ***p=0.0005, Mann-Whitney test. Error bars = SEM. Wild type N = 45, $klc4^{uw314}$ N = 41. 55 total larva were tested for each genotype, but larva that did not respond were not included in these data. **D)** Schematic of the novel tank

test design, with upper and lower zones demarcated by the center dotted line. **E)**
Percentage of fish per experiment that entered the upper zone of the tank during a 3-
minute exposure to a novel tank. Error bars = SEM. Each point represents one
experiment day: wild type N = 3 experiments with 30 total fish, *klc4*^{uw314} N = 3
experiments with 39 total fish. **F)** The time it took for individual fish to first cross into the
upper zone during a 3-minute exposure to a novel tank. Points represent individual fish,
error bars = SEM. Wild type N = 24, *klc4*^{uw314} N = 8. Fish that did not cross into the
upper zone are not represented in this data. NS by t-test, p=0.7766. **G)** The total time
fish spent in the upper zone of the novel tank during a 3-minute exposure. Points
represent individual fish, ***p=0.0002 by t-test, error bars = SEM. Wild type N = 30,
klc4^{uw314} N = 39. **H)** Percentage of fish to freeze per experiment during a 3-minute
exposure to the novel tank. Error bars = SEM. Each point represents one experiment
day: wild type N = 3 experiments with 30 total fish, *klc4*^{uw314} N = 3 experiments with 39
total fish. **I)** The total time fish spent frozen during a 3-minute exposure to a novel tank.
Points represent individual fish, ****p <0.0001 by Mann-Whitney test, error bars = SEM.
Wild type N = 30, *klc4*^{uw314} N = 39.

1066

Figure 12. Adult *klc4*^{uw314} mutant anxiety-like behavior does not fully habituate after
repeat exposure to a novel environment. **A)** Percentage of fish per experiment that froze
during the first 3 minutes of a 10-minute initial exposure to a novel tank. **B)** Plot of the
percentage of fish that froze during the course of six 10-minute exposures to the same
novel tank. **C)** Plot of the average time spent frozen during the course of six 10-minute
exposures to the same novel tank. **D)** Plot of the average total and mobile-only swim
velocities of fish during each of six 10-minute exposures to the same novel tank. Mobile-
only velocity excludes periods of time the fish spent at a velocity of 0 cm/sec (frozen). **E)**
Plot of the percentage of time that fish spent in the upper zone of the tank during the
course of six 10-minute exposures to the same novel tank. **E)** Plot of the average time it
took for a fish to first enter the upper zone of the tank during the course of six 10-minute
exposures to the same novel tank. For all data, N = 4 groups per genotype, 23 total fish
per genotype. Error bars = SEM. For figures D-F, videos that could not be automatically
tracked were omitted (for WT, 6/138, for *klc4*^{uw314}, 4/138)

1081

Figure 1 -- Supplemental Figure 1. A) Diagram of the primer design for the RT-PCR of
the long and short isoforms of *klc4*. For the long isoform, the primers were designed for
regions of exon 8 (fwd) and exon 12 (rev), which are not contained in the short isoform.
The amplified product is 421 bp. For the short isoform, the primers were designed
against exon 2 (contained in long isoform) and the short isoform 3'UTR (not contained
in long isoform). The amplified product is 523 bp. **B)** The sequence alignment around
the 25 bp deletion within exon 3 contained in *klc4*^{uw314} mutants.

1089

Figure 2 -- Supplemental Figure 2. Schematic describing our branch analysis method.
A standard sized ROI (ROI$_{trunk}$) from the trunk over the yolk tube extension is selected
with the top of the ROI$_{trunk}$ aligned to the bottom of the central axon fascicle. Eight 3 μm
ROIs (white and green, ROI$_n$, n=1-8) are selected at regularly spaced intervals along
ROI$_{trunk}$. Each ROI$_n$ is duplicated and then thresholded into a binary image. This binary
image is run through Particle Analysis in ImageJ, which counts the number of particles

present with gating for size and shape characteristics. We use the number of particles
counted to represent the number of branches passing through ROI_n.
1098
Figure 4 -- Supplemental Figure 4. Additional data for branch directionality profiling.
A) All individual data points for the 40 wild type embryos analyzed (dots), with the mean
overlayed (dark green line). **B)** All individual data points for the 23 $klc4^{uw314}$ mutant
embryos analyzed (dots), with the mean overlayed (magenta line). **C)** Axon
directionality data normalized to total signal with posterior and anterior data displayed.
Error bars = 95% CI.
1105
Figure 8 -- Supplemental Figure 8.
A) Representative image series of PA-GFP-Rab5 photoactivation. A region outlining the
cell body is selected and photoactivated using a 405nm laser (middle panel). Rab5
vesicles can then be visualized moving into the axon and into branch points (lower
panel). **B)** We quantified the accumulation of Rab5 at branch points by dividing the
acquired images into 4 time periods (t1-t4) comprised of 7-8 minute intervals and
measuring the signal at branch points from maximum intensity projections of each time
period (t1-t4). Signal was normalized to the signal at t1. There was no significant
difference in the area under the curve between $klc4+/+$ and $klc4^{uw314}$ mutants
(p=0.1319, t-test, $klc4+/+$ N= 42 branches from 20 embryos, $klc4^{uw314}$ N = 18 branches
from 8 embryos). Error bars = SEM.
1117
Figure 9 -- Supplemental Figure 9. A) Central axon comet frequency. NS by Mann-
Whitney test. (anterograde p=0.4664, retrograde p=0.7268) **B)** Central axon comet
velocity. *p<0.05, Mann-Whitney test. **C)** Average distance traveled by peripheral axon
comets. NS by Mann-Whitney test (anterograde p=0.1476, retrograde p=0.9288) **D)**
Average distance traveled by central axon comets. NS by Mann-Whitney test
(anterograde p=0.3738, retrograde p=0.8836) **E)** Average comet run time for peripheral
axon comets. NS by Mann-Whitney test (***p= 0.0001, NS p= 0.1389) **F)** Average
comet run time for central axon comets. NS by Mann-Whitney test (anterograde
p=0.3363, retrograde p= 0.3196) For all data, error bars = SEM. Wild type central N = 8
segments, $klc4^{uw314}$ central N = 15 segments, Wild type peripheral N = 26 segments,
$klc4^{uw314}$ peripheral N = 48 segments. Wild type data from 10 neurons in 7 embryos.
$Klc4^{uw314}$ data from 13 neurons in 8 embryos.
1130
1131 **Figure 11 -- Supplemental Figure 11. A)** Age-matched adult wild type and $klc4^{uw314}$
1132 mutant fish. Ruler tick marks represent 1mm. **B)** Body length measurements for age
1133 matched adult fish at 7 weeks and 9 weeks old. ****p<0.0001, **p=0.0024, *p=0.013, t-
1134 test. N is given in figure as individual fish. **C)** Percentage of larvae consumed over an
1135 hour. **p=0.0090, t-test. Wild type N = 31, $klc4^{uw314}$ N = 30.
1136
Figure 12 -- Supplemental Figure 12. Additional data for novel tank habituation
experiments. **A)** The average number of freezing bouts per fish over the course of 6
exposures. **B)** The number of fish that froze out of the total number of fish in the
experiment. * indicates data for 3 fish lost from one exposure due to video equipment

malfunction. These data could not be generated again since fish had already been
exposed to the tank, so the group N for this exposure is 20 instead of 23.
1143
1144 **Video 1**
1145 A wild type Tg*(-3.1ngn1:gfp-caax)* embryo imaged by light sheet microscopy over 11.6
1146 hours to show sensory axon development, from approximately 18-29.6 hpf. Anterior is
1147 to the left.
1148
1149 **Video 2**
1150 A *klc4^{uw314}* mutant Tg*(-3.1ngn1:gfp-caax)* embryo imaged by light sheet microscopy
1151 over 8.6 hours to show sensory axon development, from approximately 19-27.6 hpf.
1152 Anterior is to the left.
1153
1154 **Video 3**
1155 Branches in a wild type Tg*(-3.1ngn1:gfp-caax)* embryo growing and spreading over 15
1156 minutes.
1157
1158 **Video 4**
1159 An example of a growth cone bifurcation in a *klc4^{uw314}* mutant Tg*(-3.1ngn1:gfp-caax)*
1160 embryo where the left branch begins to retract at approximately 23 minutes.
1161
1162 **Video 5**
1163 Example of a wild type peripheral axon growing out while maintaining a wide branch
1164 angle between sister branches.
1165
1166 **Video 6**
1167 Peripheral axon growth in a *klc4^{uw314}* mutant embryo demonstrating both fasciculation
1168 between axons from the same neuron (self-fasciculation) and the narrowing of the angle
1169 between sister axons over time.
1170
1171 **Video 7**
1172 A peripheral axon in a *klc4^{uw314}* mutant embryo contacts a neighboring peripheral axon
1173 and fasciculates with it.
1174
1175 **Video 8**
1176 Tight fasciculation is revealed in a *klc4^{uw314}* mutant peripheral axon as a gap opens
1177 between the two axons as an interstitial branch forms.
1178
1179 **Video 9**
1180 Light sheet microscopy of an embryo treated with 100µM Kinesore, showing what
1181 appear to be apical protrusions crossing the midline of the spinal cord. This behavior
1182 was seen in both Kinesore treated and *klc4^{uw314}* mutant embryos. Anterior is to the right.
1183
1184 **Video 10**

1185   High-speed, high-magnification light sheet microscopy of a *klc4^uw314* mutant embryo
1186   showing an apical protrusion growing across the midline with a dynamic tip that shows
1187   growth-cone like behavior. Anterior is to the top right corner.
1188
1189   **Video 11**
1190   A cropped portion of Video 1 showing normal behavior of apical protrusions in the spinal
1191   cord of a wild type embryo. Video has been rotated to show the spinal cord in a more
1192   horizontal position. Anterior to the left.
1193
1194

References
Andersen, E, N. Asuri, and M Halloran. 2011. "In vivo imaging of cell behaviors and F-actin
reveals LIM-HD transcription factor regulation of peripheral versus central sensory axon
development." *Neural Dev* 6: 27.
Andersen, E. F., and M. C. Halloran. 2012. "Centrosome movements in vivo correlate with
specific neurite formation downstream of LIM homeodomain transcription factor
activity." *Development* 139 (19): 3590-9.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita
tion&list_uids=22899847
Anton, Z., et al. . 2021. "Molecular mechanisms for kinesin-1 direct membrane recognition."
bioRxiv 2021.01.20.427326.
https://doi.org/https://doi.org/10.1101/2021.01.20.427326.
Armijo-Weingart, Lorena, and Gianluca Gallo. 2017. "It takes a village to raise a branch: Cellular
mechanisms of the initiation of axon collateral branches." *Molecular and Cellular*
Neuroscience 84: 36-47. https://doi.org/10.1016/j.mcn.2017.03.007.
https://reader.elsevier.com/reader/sd/pii/S1044743116302093?token=4A9CDFAB7F5B
3C900E6293DE9EA59222CD47B95B5A9A8044341CB6EE6A873FABE4C793C2969CD7BF7
FA48876918DF603&originRegion=us-east-1&originCreation=20210722045428.
Baek, J. H., J. Lee, H. S. Yun, C. W. Lee, J. Y. Song, H. D. Um, J. K. Park, I. C. Park, J. S. Kim, E. H.
Kim, and S. G. Hwang. 2018. "Kinesin light chain-4 depletion induces apoptosis of
radioresistant cancer cells by mitochondrial dysfunction via calcium ion influx." *Cell*
Death Dis 9 (5): 496. https://doi.org/10.1038/s41419-018-0549-2.
https://www.ncbi.nlm.nih.gov/pubmed/29717133.
Bayrakli, F., H. G. Poyrazoglu, S. Yuksel, C. Yakicier, B. Erguner, M. S. Sagiroglu, B. Yuceturk, B.
Ozer, S. Doganay, B. Tanrikulu, A. Seker, F. Akbulut, A. Ozen, H. Per, S. Kumandas, Y.
Altuner Torun, Y. Bayri, M. Sakar, A. Dagcinar, and I. Ziyal. 2015. "Hereditary spastic
paraplegia with recessive trait caused by mutation in KLC4 gene." *J Hum Genet* 60 (12):
763-8. https://doi.org/10.1038/jhg.2015.109.
https://www.ncbi.nlm.nih.gov/pubmed/26423925.
Bazvand, Fatemeh, Mohammad Keramatipour, Hamid Riazi-Esfahani, and Alireza Mahmoudi.
2021. "SPOAN syndrome: a novel mutation and new ocular findings; a case report." *BMC*
Neurology 21 (1): 24. https://doi.org/10.1186/s12883-021-02051-9.
https://doi.org/10.1186/s12883-021-02051-9.

Cai, Dawen, Adam D. Hoppe, Joel A. Swanson, and Kristen J. Verhey. 2007. "Kinesin-1 structural
organization and conformational changes revealed by FRET stoichiometry in live cells."
Journal of Cell Biology 176 (1): 51-63. https://doi.org/10.1083/jcb.200605097.
https://doi.org/10.1083/jcb.200605097.
Chen, Weisheng V., Chiamaka L. Nwakeze, Christine A. Denny, Sean O'Keeffe, Michael A. Rieger,
George Mountoufaris, Amy Kirner, Joseph D. Dougherty, René Hen, Qiang Wu, and Tom
Maniatis. 2017. "Pcdhαc2 is required for axonal tiling and assembly of serotonergic
circuitries in mice." *Science* 356 (6336): 406-411.
https://doi.org/10.1126/science.aal3231.
https://science.sciencemag.org/content/sci/356/6336/406.full.pdf.
Chiba, Kyoko, Kassandra M. Ori-McKenney, Shinsuke Niwa, and Richard J. McKenney. 2021.
"Reconstitution of Kinesin-1 Activation." *bioRxiv*: 2021.03.12.434960.
https://doi.org/10.1101/2021.03.12.434960.
http://biorxiv.org/content/early/2021/03/12/2021.03.12.434960.abstract.
Choi, Jung-Hwa, Yun-Mi Jeong, Sujin Kim, Boyoung Lee, Krishan Ariyasiri, Hyun-Taek Kim,
Seung-Hyun Jung, Kyu-Seok Hwang, Tae-Ik Choi, Chul O. Park, Won-Ki Huh, Matthias
Carl, Jill A. Rosenfeld, Salmo Raskin, Alan Ma, Jozef Gecz, Hyung-Goo Kim, Jin-Soo Kim,
Ho-Chul Shin, Doo-Sang Park, Robert Gerlai, Bradley B. Jamieson, Joon S. Kim, Karl J.
Iremonger, Sang H. Lee, Hee-Sup Shin, and Cheol-Hee Kim. 2018. "Targeted knockout of
a chemokine-like gene increases anxiety and fear responses." *Proceedings of the
National Academy of Sciences* 115 (5): E1041.
https://doi.org/10.1073/pnas.1707663115.
http://www.pnas.org/content/115/5/E1041.abstract.
Clark, B. S., M. Winter, A. R. Cohen, and B. A. Link. 2011. "Generation of Rab-based transgenic
lines for in vivo studies of endosome biology in zebrafish." *Dev Dyn* 240 (11): 2452-65.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita
tion&list_uids=21976318
Cockburn, J. J. B., S. J. Hesketh, P. Mulhair, M. Thomsen, M. J. O'Connell, and M. Way. 2018.
"Insights into Kinesin-1 Activation from the Crystal Structure of KLC2 Bound to JIP3."
Structure 26 (11): 1486-1498 e6. https://doi.org/10.1016/j.str.2018.07.011.
https://www.ncbi.nlm.nih.gov/pubmed/30197037.
Courchet, J., T. L. Lewis, Jr., S. Lee, V. Courchet, D. Y. Liou, S. Aizawa, and F. Polleux. 2013.
"Terminal axon branching is regulated by the LKB1-NUAK1 kinase pathway via
presynaptic mitochondrial capture." *Cell* 153 (7): 1510-25.
https://doi.org/10.1016/j.cell.2013.05.021.
http://www.ncbi.nlm.nih.gov/pubmed/23791179.
Csordas, G., D. Weaver, and G. Hajnoczky. 2018. "Endoplasmic Reticulum-Mitochondrial
Contactology: Structure and Signaling Functions." *Trends Cell Biol* 28 (7): 523-540.
https://doi.org/10.1016/j.tcb.2018.02.009.
https://www.ncbi.nlm.nih.gov/pubmed/29588129.
Daetwyler, Stephan, Ulrik Günther, Carl D. Modes, Kyle Harrington, and Jan Huisken. 2019.
"Multi-sample SPIM image acquisition, processing and analysis of vascular growth in
zebrafish." *Development* 146 (6). https://doi.org/10.1242/dev.173757.
https://doi.org/10.1242/dev.173757.

Dehairs, Jonas, Ali Talebi, Yacine Cherifi, and Johannes V. Swinnen. 2016. "CRISP-ID: decoding
CRISPR mediated indels by Sanger sequencing." *Scientific Reports* 6 (1): 1-5.
https://doi.org/doi:10.1038/srep28973. https://www.nature.com/articles/srep28973.
Dent, E. W., and K. Kalil. 2001. "Axon branching requires interactions between dynamic
microtubules and actin filaments." *J Neurosci* 21 (24): 9757-69.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita
tion&list_uids=11739584
Du, J., Y. Wei, L. Liu, Y. Wang, R. Khairova, R. Blumenthal, T. Tragon, J. G. Hunsberger, R.
Machado-Vieira, W. Drevets, Y. T. Wang, and H. K. Manji. 2010. "A kinesin signaling
complex mediates the ability of GSK-3beta to affect mood-associated behaviors." *Proc
Natl Acad Sci U S A* 107 (25): 11573-8. https://doi.org/10.1073/pnas.0913138107.
https://www.ncbi.nlm.nih.gov/pubmed/20534517.
Fu, Xiaolong, Yachun An, Hongyang Wang, Peipei Li, Jing Lin, Jia Yuan, Rongyu Yue, Yecheng Jin,
Jiangang Gao, and Renjie Chai. 2021. *Molecular Neurobiology* 58 (9): 4376-4391.
https://doi.org/doi:10.1007/s12035-021-02422-w.
https://link.springer.com/article/10.1007/s12035-021-02422-w.
Gallo, G. 2011. "The cytoskeletal and signaling mechanisms of axon collateral branching." *Dev
Neurobiol* 71 (3): 201-20.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita
tion&list_uids=21308993
Gibson, D. A., and L. Ma. 2011. "Developmental regulation of axon branching in the vertebrate
nervous system." *Development* 138 (2): 183-95.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita
tion&list_uids=21177340
Gibson, D. A., S. Tymanskyj, R. C. Yuan, H. C. Leung, J. L. Lefebvre, J. R. Sanes, A. Chédotal, and L.
Ma. 2014. "Dendrite self-avoidance requires cell-autonomous slit/robo signaling in
cerebellar purkinje cells." *Neuron* 81 (5): 1040-1056.
https://doi.org/10.1016/j.neuron.2014.01.009.
Gindhart, J. G., Jr., C. J. Desai, S. Beushausen, K. Zinn, and L. S. Goldstein. 1998. "Kinesin light
chains are essential for axonal transport in Drosophila." *J Cell Biol* 141 (2): 443-54.
https://doi.org/10.1083/jcb.141.2.443.
https://www.ncbi.nlm.nih.gov/pubmed/9548722.
Grueber, Wesley B., and Alvaro Sagasti. 2010. "Self-avoidance and Tiling: Mechanisms of
Dendrite and Axon Spacing." *Cold Spring Harbor Perspectives in Biology* 2 (9).
https://doi.org/10.1101/cshperspect.a001750.
http://cshperspectives.cshlp.org/content/2/9/a001750.abstract.
Gupta, Shailesh Kumar, Karina F. Meiri, Kashif Mahfooz, Upasna Bharti, and Shyamala Mani.
2010. "Coordination between Extrinsic Extracellular Matrix Cues and Intrinsic Responses
to Orient the Centrosome in Polarizing Cerebellar Granule Neurons." *The Journal of
Neuroscience* 30 (7): 2755. https://doi.org/10.1523/JNEUROSCI.4218-09.2010.
http://www.jneurosci.org/content/30/7/2755.abstract.
Haase, Robert, Loic A. Royer, Peter Steinbach, Deborah Schmidt, Alexandr Dibrov, Uwe Schmidt,
Martin Weigert, Nicola Maghelli, Pavel Tomancak, Florian Jug, and Eugene W. Myers.
2019. "CLIJ: GPU-accelerated image processing for everyone." *Nature Methods* 17 (1): 5-

6. https://doi.org/doi:10.1038/s41592-019-0650-1.
https://www.nature.com/articles/s41592-019-0650-1.

Hakeda-Suzuki, Satoko, Sandra Berger-Müller, Tatiana Tomasi, Tadao Usui, Shin-ya Horiuchi,
Tadashi Uemura, and Takashi Suzuki. 2011. "Golden Goal collaborates with Flamingo in
conferring synaptic-layer specificity in the visual system." *Nature Neuroscience* 14 (3):
314-323. https://doi.org/10.1038/nn.2756. https://doi.org/10.1038/nn.2756.

Halloran, M. C., S. M. Severance, C. S. Yee, D. L. Gemza, J. A. Raper, and J. Y. Kuwada. 1999.
"Analysis of a zebrafish semaphorin reveals potential functions in vivo." *Developmental*
Dynamics 214: 13-25.

Hirokawa, N., S. Niwa, and Y. Tanaka. 2010. "Molecular motors in neurons: transport
mechanisms and roles in brain function, development, and disease." *Neuron* 68 (4): 610-
38.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita
tion&list_uids=21092854

Howe, Hollis B., Peter B. McIntyre, and Marc A. Wolman. 2018. "Adult zebrafish primarily use
vision to guide piscivorous foraging behavior | Elsevier Enhanced Reader." *Behavioural*
Processes 157: 230-237. https://doi.org/10.1016/j.beproc.2018.10.005.
https://reader.elsevier.com/reader/sd/pii/S0376635718302584?token=BD74AF6F73B3
B85BEBC8FD1C88FA245B60B1CB9801D386E4D1CBDD9784118B6DE94FC10C34E4DFB7
D4FBE44EFC68CA09&originRegion=us-east-1&originCreation=20210722050736.

Jiang, M., J. Esteve-Rudd, V. S. Lopes, T. Diemer, C. Lillo, A. Rump, and D. S. Williams. 2015.
"Microtubule motors transport phagosomes in the RPE, and lack of KLC1 leads to AMD-
like pathogenesis." *J Cell Biol* 210 (4): 595-611. https://doi.org/10.1083/jcb.201410112.
https://www.ncbi.nlm.nih.gov/pubmed/26261180.

Kalil, K., and E. W. Dent. 2014. "Branch management: mechanisms of axon branching in the
developing vertebrate CNS." *Nat Rev Neurosci* 15 (1): 7-18.
https://doi.org/10.1038/nrn3650. https://www.ncbi.nlm.nih.gov/pubmed/24356070.

Kamal, A., G. B. Stokin, Z. Yang, C. H. Xia, and L. S. Goldstein. 2000. "Axonal transport of amyloid
precursor protein is mediated by direct binding to the kinesin light chain subunit of
kinesin-I." *Neuron* 28 (2): 449-59. https://doi.org/10.1016/s0896-6273(00)00124-0.
https://www.ncbi.nlm.nih.gov/pubmed/11144355.

Katori, Shota, Yukiko Noguchi-Katori, Atsushi Okayama, Yoshimi Kawamura, Wenshu Luo, Kenji
Sakimura, Takahiro Hirabayashi, Takuji Iwasato, and Takeshi Yagi. 2017. "Protocadherin-
αC2 is required for diffuse projections of serotonergic axons." *Scientific Reports* 7 (1):
15908. https://doi.org/10.1038/s41598-017-16120-y. https://doi.org/10.1038/s41598-
017-16120-y.

Ketschek, A., S. Jones, M. Spillane, F. Korobova, T. Svitkina, and G. Gallo. 2015. "Nerve growth
factor promotes reorganization of the axonal microtubule array at sites of axon
collateral branching." *Dev Neurobiol* 75 (12): 1441-61.
https://doi.org/10.1002/dneu.22294.
https://www.ncbi.nlm.nih.gov/pubmed/25846486.

Khan, K. M., A. D. Collier, D. A. Meshalkina, E. V. Kysil, S. L. Khatsko, T. Kolesnikova, Y. Y.
Morzherin, J. E. Warnick, A. V. Kalueff, and D. J. Echevarria. 2017. "Zebrafish models in
neuropsychopharmacology and CNS drug discovery." *Br J Pharmacol* 174 (13): 1925-

1944. https://doi.org/10.1111/bph.13754.
https://www.ncbi.nlm.nih.gov/pubmed/28217866.
Kimmel, C. B., W. W. Ballard, S. R. Kimmel, B. Ullmann, and T. F. Schilling. 1995. "Stages of
embryonic development of the zebrafish." *Dev Dyn* 203 (3): 253-310.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita
tion&list_uids=8589427
Knafo, Steven, Kevin Fidelin, Andrew Prendergast, Po-En Brian Tseng, Alexandre Parrin, Charles
Dickey, Urs Lucas Böhm, Sophie Nunes Figueiredo, Olivier Thouvenin, Hugues Pascal-
Moussellard, and Claire Wyart. 2017. "Mechanosensory neurons control the timing of
spinal microcircuit selection during locomotion."
https://doi.org/doi:10.7554/eLife.25260. https://elifesciences.org/articles/25260.
Koh, S. Y., J. T. Cain, H. Magee, K. A. White, M. Rechtzigel, B. Meyerink, H. Leppert, D. J. Timm, J.
P. Morgan, T. B. Johnson, B. Grove, R. Khanna, K. Hensley, J. Brudvig, and J. M. Weimer.
2021. "A CLN6-CRMP2-KLC4 complex regulates anterograde ER-derived vesicle
trafficking in cortical neurites." *bioRxiv*: 2021.09.16.460653.
https://doi.org/10.1101/2021.09.16.460653.
http://biorxiv.org/content/early/2021/09/18/2021.09.16.460653.abstract.
Kostadinov, Dimitar, and Joshua R Sanes. 2015. "Protocadherin-dependent dendritic self-
avoidance regulates neural connectivity and circuit function."
https://doi.org/doi:10.7554/eLife.08964. https://elifesciences.org/articles/8964.
Krols, M., G. Bultynck, and S. Janssens. 2016. "ER-Mitochondria contact sites: A new regulator
of cellular calcium flux comes into play." *J Cell Biol* 214 (4): 367-70.
https://doi.org/10.1083/jcb.201607124.
https://www.ncbi.nlm.nih.gov/pubmed/27528654.
Kwan, K. M., E. Fujimoto, C. Grabher, B. D. Mangum, M. E. Hardy, D. S. Campbell, J. M. Parant,
H. J. Yost, J. P. Kanki, and C. B. Chien. 2007. "The Tol2kit: a multisite gateway-based
construction kit for Tol2 transposon transgenesis constructs." *Dev Dyn* 236 (11): 3088-
99.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita
tion&list_uids=17937395
Lee, T. J., J. W. Lee, E. M. Haynes, K. W. Eliceiri, and M. C. Halloran. 2017. "The Kinesin Adaptor
Calsyntenin-1 Organizes Microtubule Polarity and Regulates Dynamics during Sensory
Axon Arbor Development." *Front Cell Neurosci* 11: 107.
https://doi.org/10.3389/fncel.2017.00107.
https://www.ncbi.nlm.nih.gov/pubmed/28473757.
Lefebvre, Julie L., Dimitar Kostadinov, Weisheng V. Chen, Tom Maniatis, and Joshua R. Sanes.
2012. "Protocadherins mediate dendritic self-avoidance in the mammalian nervous
system." *Nature* 488 (7412): 517-521. https://doi.org/10.1038/nature11305.
https://doi.org/10.1038/nature11305.
Liu, Y., and M. C. Halloran. 2005. "Central and peripheral axon branches from one neuron are
guided differentially by Semaphorin3D and transient axonal glycoprotein-1." *J Neurosci*
25 (45): 10556-63.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita
tion&list_uids=16280593.

Liz, Márcia A, Fernando M Mar, Telma E Santos, Helena I Pimentel, Ana M Marques, Marlene M
Morgado, Sílvia Vieira, Vera F Sousa, Hayley Pemble, Torsten Wittmann, Calum
Sutherland, James R Woodgett, and Mónica M Sousa. 2014. "Neuronal deletion of
GSK3β increases microtubule speed in the growth cone and enhances axon regeneration
via CRMP-2 and independently of MAP1B and CLASP2." *BMC Biology* 12 (1): 1-19.
https://doi.org/doi:10.1186/1741-7007-12-47.
https://bmcbiol.biomedcentral.com/articles/10.1186/1741-7007-12-47.
Lu, W., and V. I. Gelfand. 2017. "Moonlighting Motors: Kinesin, Dynein, and Cell Polarity."
Trends Cell Biol 27 (7): 505-514. https://doi.org/10.1016/j.tcb.2017.02.005.
https://www.ncbi.nlm.nih.gov/pubmed/28284467.
Matthews, Benjamin J., Michelle E. Kim, John J. Flanagan, Daisuke Hattori, James C. Clemens, S.
Lawrence Zipursky, and Wesley B. Grueber. 2007. "Dendrite Self-Avoidance Is
Controlled by Dscam." *Cell* 129 (3): 593-604.
https://doi.org/https://doi.org/10.1016/j.cell.2007.04.013.
https://www.sciencedirect.com/science/article/pii/S0092867407004709.
Maximino, Caio, Thiago Marques de Brito, Annanda Waneza da Silva Batista, Anderson Manoel
Herculano, Silvio Morato, and Amauri Gouveia. 2010. "Measuring anxiety in zebrafish: A
critical review." *Behavioural Brain Research* 214 (2): 157-171.
https://doi.org/https://doi.org/10.1016/j.bbr.2010.05.031.
https://www.sciencedirect.com/science/article/pii/S0166432810003931.
Melo, U. S., L. I. Macedo-Souza, T. Figueiredo, A. R. Muotri, J. G. Gleeson, G. Coux, P. Armas, N.
B. Calcaterra, J. P. Kitajima, S. Amorim, T. R. Olavio, K. Griesi-Oliveira, G. C. Coatti, C. R.
Rocha, M. Martins-Pinheiro, C. F. Menck, M. S. Zaki, F. Kok, M. Zatz, and S. Santos. 2015.
"Overexpression of KLC2 due to a homozygous deletion in the non-coding region causes
SPOAN syndrome." *Hum Mol Genet* 24 (24): 6877-85.
https://doi.org/10.1093/hmg/ddv388.
https://www.ncbi.nlm.nih.gov/pubmed/26385635.
Millard, S. Sean, John J. Flanagan, Kartik S. Pappu, Wei Wu, and S. Lawrence Zipursky. 2007.
"Dscam2 mediates axonal tiling in the Drosophila visual system." *Nature* 447 (7145):
720-724. https://doi.org/10.1038/nature05855. https://doi.org/10.1038/nature05855.
Mochizuki, H., H. Toda, M. Ando, M. Kurusu, T. Tomoda, and K. Furukubo-Tokunaga. 2011.
"Unc-51/ATG1 controls axonal and dendritic development via kinesin-mediated vesicle
transport in the Drosophila brain." *PLoS One* 6 (5): e19632.
https://doi.org/10.1371/journal.pone.0019632.
https://www.ncbi.nlm.nih.gov/pubmed/21589871.
Moreno-Mateos, Miguel A, Charles E Vejnar, Jean-Denis Beaudoin, Juan P Fernandez, Emily K
Mis, Mustafa K Khokha, and Antonio J Giraldez. 2015. "CRISPRscan: designing highly
efficient sgRNAs for CRISPR-Cas9 targeting in vivo." *Nature Methods* 12 (10): 982-988.
https://doi.org/doi:10.1038/nmeth.3543.
https://www.nature.com/articles/nmeth.3543.
Morfini, G., N. Schmidt, C. Weissmann, G. Pigino, and S. Kins. 2016. "Conventional kinesin:
Biochemical heterogeneity and functional implications in health and disease." *Brain Res*
Bull 126 (Pt 3): 347-353. https://doi.org/10.1016/j.brainresbull.2016.06.009.
https://www.ncbi.nlm.nih.gov/pubmed/27339812.

Nakamura, Fumio, Toshio Ohshima, and Yoshio Goshima. 2020. "Collapsin Response Mediator
Proteins: Their Biological Functions and Pathophysiology in Neuronal Development and
Regeneration." *Frontiers in Cellular Neuroscience* 14.
https://doi.org/10.3389/fncel.2020.00188.
https://www.frontiersin.org/articles/10.3389/fncel.2020.00188.
Navone, F., J. Niclas, N. Hom-Booher, L. Sparks, H. D. Bernstein, G. McCaffrey, and R. D. Vale.
1992. "Cloning and expression of a human kinesin heavy chain gene: interaction of the
COOH-terminal domain with cytoplasmic microtubules in transfected CV-1 cells."
Journal of Cell Biology 117 (6): 1263-1275. https://doi.org/10.1083/jcb.117.6.1263.
https://doi.org/10.1083/jcb.117.6.1263.
Orefice, Lauren L., Amanda L. Zimmerman, Anda M. Chirila, Steven J. Sleboda, Joshua P. Head,
and David D. Ginty. 2016. "Peripheral Mechanosensory Neuron Dysfunction Underlies
Tactile and Behavioral Deficits in Mouse Models of ASDs." *Cell* 166 (2): 299-313.
https://doi.org/10.1016/j.cell.2016.05.033. https://doi.org/10.1016/j.cell.2016.05.033.
Pernigo, S., M. S. Chegkazi, Y. Y. Yip, C. Treacy, G. Glorani, K. Hansen, A. Politis, S. Bui, M. P.
Dodding, and R. A. Steiner. 2018. "Structural basis for isoform-specific kinesin-1
recognition of Y-acidic cargo adaptors." *Elife* 7. https://doi.org/10.7554/eLife.38362.
https://www.ncbi.nlm.nih.gov/pubmed/30320553.
Ponomareva, O. Y., K. W. Eliceiri, and M. C. Halloran. 2016. "Charcot-Marie-Tooth 2b associated
Rab7 mutations cause axon growth and guidance defects during vertebrate sensory
neuron development." *Neural Dev* 11: 2. https://doi.org/10.1186/s13064-016-0058-x.
https://www.ncbi.nlm.nih.gov/pubmed/26791407.
Ponomareva, O. Y., I. C. Holmen, A. J. Sperry, K. W. Eliceiri, and M. C. Halloran. 2014.
"Calsyntenin-1 regulates axon branching and endosomal trafficking during sensory
neuron development in vivo." *The Journal of neuroscience : the official journal of the
Society for Neuroscience* 34 (28): 9235-48. https://doi.org/10.1523/JNEUROSCI.0561-
14.2014. http://www.ncbi.nlm.nih.gov/pubmed/25009257.
Qiang, L., W. Yu, M. Liu, J. M. Solowska, and P. W. Baas. 2010. "Basic fibroblast growth factor
elicits formation of interstitial axonal branches via enhanced severing of microtubules."
Mol Biol Cell 21 (2): 334-44.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita
tion&list_uids=19940015
Rahman, A., A. Kamal, E. A. Roberts, and L. S. Goldstein. 1999. "Defective kinesin heavy chain
behavior in mouse kinesin light chain mutants." *J Cell Biol* 146 (6): 1277-88.
https://doi.org/10.1083/jcb.146.6.1277.
https://www.ncbi.nlm.nih.gov/pubmed/10491391.
Randall, Thomas S., Yan Y. Yip, Daynea J. Wallock-Richards, Karin Pfisterer, Anneri Sanger,
Weronika Ficek, Roberto A. Steiner, Andrew J. Beavil, Maddy Parsons, and Mark P.
Dodding. 2017. "A small-molecule activator of kinesin-1 drives remodeling of the
microtubule network." https://doi.org/10.1073/pnas.1715115115.
https://www.pnas.org/content/114/52/13738.
Randlett, Owen, Lucia Poggi, Flavio R Zolessi, and William A Harris. 2011. *Neuron* 70 (2): 266-
280. https://doi.org/10.1016/j.neuron.2011.03.013.
https://doi.org/10.1016/j.neuron.2011.03.013.

Rezakhaniha, R., A. Agianniotis, J. T. C. Schrauwen, A. Griffa, D. Sage, C. V. C. Bouten, F. N. van
de Vosse, M. Unser, and N. Stergiopulos. 2011. "Experimental investigation of collagen
waviness and orientation in the arterial adventitia using confocal laser scanning
microscopy." *Biomechanics and Modeling in Mechanobiology* 11 (3): 461-473.
https://doi.org/doi:10.1007/s10237-011-0325-z.
https://link.springer.com/article/10.1007/s10237-011-0325-z.
Rodriguez, Alvaro, Hanqing Zhang, Jonatan Klaminder, Tomas Brodin, Patrik L. Andersson, and
Magnus Andersson. 2018. "ToxTrac: A fast and robust software for tracking organisms."
Methods in Ecology and Evolution 9 (3): 460-464. https://doi.org/10.1111/2041-
210X.12874. https://besjournals.onlinelibrary.wiley.com/doi/10.1111/2041-
210X.12874.
Rowland, A. A., and G. K. Voeltz. 2012. "Endoplasmic reticulum-mitochondria contacts: function
of the junction." *Nat Rev Mol Cell Biol* 13 (10): 607-25.
https://doi.org/10.1038/nrm3440. https://www.ncbi.nlm.nih.gov/pubmed/22992592.
Saez, T. M. M., I. Fernandez Bessone, M. S. Rodriguez, M. Alloatti, M. G. Otero, L. E. Cromberg,
V. M. Pozo Devoto, G. Oubina, L. Sosa, M. G. Buffone, D. M. Gelman, and T. L. Falzone.
2020. "Kinesin-1-mediated axonal transport of CB1 receptors is required for
cannabinoid-dependent axonal growth and guidance." *Development* 147 (8).
https://doi.org/10.1242/dev.184069.
https://www.ncbi.nlm.nih.gov/pubmed/32265198.
Sagasti, A., M. R. Guido, D. W. Raible, and A. F. Schier. 2005. "Repulsive interactions shape the
morphologies and functional arrangement of zebrafish peripheral sensory arbors." *Curr
Biol* 15 (9): 804-14.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita
tion&list_uids=15886097.
Schindelin, Johannes, Ignacio Arganda-Carreras, Erwin Frise, Verena Kaynig, Mark Longair,
Tobias Pietzsch, Stephan Preibisch, Curtis Rueden, Stephan Saalfeld, Benjamin Schmid,
Jean-Yves Tinevez, Daniel James White, Volker Hartenstein, Kevin Eliceiri, Pavel
Tomancak, and Albert Cardona. 2012. "Fiji: an open-source platform for biological-image
analysis." *Nature Methods* 9 (7): 676-682. https://doi.org/doi:10.1038/nmeth.2019.
https://www.nature.com/articles/nmeth.2019.
Seeger, Mark A., and Sarah E. Rice. 2010. "Microtubule-associated Protein-like Binding of the
Kinesin-1 Tail to Microtubules." *Journal of Biological Chemistry* 285 (11): 8155-8162.
https://doi.org/10.1074/jbc.M109.068247. https://doi.org/10.1074/jbc.M109.068247.
Senti, Kirsten-André, Tadao Usui, Karin Boucke, Urs Greber, Tadashi Uemura, and Barry J.
Dickson. 2003. "Flamingo Regulates R8 Axon-Axon and Axon-Target Interactions in the
Drosophila Visual System." *Current Biology* 13 (10): 828-832.
https://doi.org/10.1016/S0960-9822(03)00291-4. https://doi.org/10.1016/S0960-
9822(03)00291-4.
Shah, Arish N, Crystal F Davey, Alex C Whitebirch, Adam C Miller, and Cecilia B Moens. 2015.
"Rapid reverse genetic screening using CRISPR in zebrafish." *Nature Methods* 12 (6):
535-540. https://doi.org/doi:10.1038/nmeth.3360.
https://www.nature.com/articles/nmeth.3360.

Shima, Tomohiro, Manatsu Morikawa, Junichi Kaneshiro, Taketoshi Kambara, Shinji Kamimura, Toshiki Yagi, Hiroyuki Iwamoto, Sotaro Uemura, Hideki Shigematsu, Mikako Shirouzu, Taro Ichimura, Tomonobu M. Watanabe, Ryo Nitta, Yasushi Okada, and Nobutaka Hirokawa. 2018. "Kinesin-binding–triggered conformation switching of microtubules contributes to polarized transport." *Journal of Cell Biology* 217 (12): 4164-4183. https://doi.org/10.1083/jcb.201711178. https://doi.org/10.1083/jcb.201711178.

Smith, Cody J., Joseph D. Watson, Miri K. VanHoven, Daniel A. Colón-Ramos, and David M. Miller. 2012. "Netrin (UNC-6) mediates dendritic self-avoidance." *Nature Neuroscience* 15 (5): 731-737. https://doi.org/10.1038/nn.3065. https://doi.org/10.1038/nn.3065.

Soba, Peter, Sijun Zhu, Kazuo Emoto, Susan Younger, Shun-Jen Yang, Hung-Hsiang Yu, Tzumin Lee, Lily Yeh Jan, and Yuh-Nung Jan. 2007. "Drosophila Sensory Neurons Require Dscam for Dendritic Self-Avoidance and Proper Dendritic Field Organization." *Neuron* 54 (3): 403-416. https://doi.org/https://doi.org/10.1016/j.neuron.2007.03.029. https://www.sciencedirect.com/science/article/pii/S0896627307002887.

Spead, Olivia, and Fabienne E. Poulain. 2020. "Trans-Axonal Signaling in Neural Circuit Wiring." *International Journal of Molecular Sciences* 21 (14). https://doi.org/10.3390/ijms21145170.

Spillane, M., A. Ketschek, T. T. Merianda, J. L. Twiss, and G. Gallo. 2013. "Mitochondria coordinate sites of axon branching through localized intra-axonal protein synthesis." *Cell reports* 5 (6): 1564-75. https://doi.org/10.1016/j.celrep.2013.11.022. http://www.ncbi.nlm.nih.gov/pubmed/24332852.

Stepanova, T., J. Slemmer, C. C. Hoogenraad, G. Lansbergen, B. Dortland, C. I. De Zeeuw, F. Grosveld, G. van Cappellen, A. Akhmanova, and N. Galjart. 2003. "Visualization of microtubule growth in cultured neurons via the use of EB3-GFP (end-binding protein 3-green fluorescent protein)." *J Neurosci* 23 (7): 2655-64. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12684451

Tao, K., N. Matsuki, and R. Koyama. 2014. "AMP-activated protein kinase mediates activity-dependent axon branching by recruiting mitochondria to axon." *Dev Neurobiol* 74 (6): 557-73. https://doi.org/10.1002/dneu.22149. http://www.ncbi.nlm.nih.gov/pubmed/24218086.

Tomasi, Tatiana, Satoko Hakeda-Suzuki, Stephan Ohler, Alexander Schleiffer, and Takashi Suzuki. 2008. "The Transmembrane Protein Golden Goal Regulates R8 Photoreceptor Axon-Axon and Axon-Target Interactions." *Neuron* 57 (5): 691-704. https://doi.org/10.1016/j.neuron.2008.01.012. https://doi.org/10.1016/j.neuron.2008.01.012.

Triclin, Sarah, Daisuke Inoue, Jérémie Gaillard, Zaw Min Htet, Morgan E. DeSantis, Didier Portran, Emmanuel Derivery, Charlotte Aumeier, Laura Schaedel, Karin John, Christophe Leterrier, Samara L. Reck-Peterson, Laurent Blanchoin, and Manuel Théry. 2021. "Self-repair protects microtubules from destruction by molecular motors." *Nature Materials* 20 (6): 883-891. https://doi.org/10.1038/s41563-020-00905-0. https://doi.org/10.1038/s41563-020-00905-0.

Wang, J., C. T. Zugates, I. H. Liang, C. H. Lee, and T. Lee. 2002. "Drosophila Dscam is required for divergent segregation of sister branches and suppresses ectopic bifurcation of axons."

Neuron 33 (4): 559-71.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita
tion&list_uids=11856530.
Webster, D. R., and G. G. Borisy. 1989. "Microtubules are acetylated in domains that turn over
slowly." *Journal of Cell Science* 92 (1): 57-65. https://doi.org/10.1242/jcs.92.1.57.
https://doi.org/10.1242/jcs.92.1.57.
Williams, Kristina, and Angeles B. Ribera. 2020. "Long-lived zebrafish Rohon-Beard cells."
Developmental Biology 464 (1): 45-52.
https://doi.org/https://doi.org/10.1016/j.ydbio.2020.05.003.
https://www.sciencedirect.com/science/article/pii/S0012160620301470.
Winkle, C. C., K. L. Taylor, E. W. Dent, G. Gallo, K. F. Greif, and S. L. Gupton. 2016. "Beyond the
cytoskeleton: The emerging role of organelles and membrane remodeling in the
regulation of axon collateral branches." *Dev Neurobiol* 76 (12): 1293-1307.
https://doi.org/10.1002/dneu.22398.
https://www.ncbi.nlm.nih.gov/pubmed/27112549.
Wong, H. H., J. Q. Lin, F. Strohl, C. G. Roque, J. M. Cioni, R. Cagnetta, B. Turner-Bridger, R. F.
Laine, W. A. Harris, C. F. Kaminski, and C. E. Holt. 2017. "RNA Docking and Local
Translation Regulate Site-Specific Axon Remodeling In Vivo." *Neuron* 95 (4): 852-868 e8.
https://doi.org/10.1016/j.neuron.2017.07.016.
https://www.ncbi.nlm.nih.gov/pubmed/28781168.
Wong, Keith, Marco Elegante, Brett Bartels, Salem Elkhayat, David Tien, Sudipta Roy, Jason
Goodspeed, Chris Suciu, Julia Tan, Chelsea Grimes, Amanda Chung, Michael Rosenberg,
Siddharth Gaikwad, Ashley Denmark, Andrew Jackson, Ferdous Kadri, Kyung Min Chung,
Adam Stewart, Tom Gilder, Esther Beeson, Ivan Zapolsky, Nadine Wu, Jonathan Cachat,
and Allan V. Kalueff. 2010. "Analyzing habituation responses to novelty in zebrafish
(Danio rerio)." *Behavioural Brain Research* 208 (2): 450-457.
https://doi.org/https://doi.org/10.1016/j.bbr.2009.12.023.
https://www.sciencedirect.com/science/article/pii/S0166432809007499.
Wong, Y. L., and S. E. Rice. 2010. "Kinesin's light chains inhibit the head- and microtubule-
binding activity of its tail." *Proc Natl Acad Sci U S A* 107 (26): 11781-6.
https://doi.org/10.1073/pnas.1005854107.
https://www.ncbi.nlm.nih.gov/pubmed/20547877.
Wozniak, M. J., and V. J. Allan. 2006. "Cargo selection by specific kinesin light chain 1 isoforms."
EMBO J 25 (23): 5457-68. https://doi.org/10.1038/sj.emboj.7601427.
https://www.ncbi.nlm.nih.gov/pubmed/17093494.
Yu, W., F. J. Ahmad, and P. W. Baas. 1994. "Microtubule fragmentation and partitioning in the
axon during collateral branch formation." *J Neurosci* 14 (10): 5872-84.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita
tion&list_uids=7931550
Yu, W., L. Qiang, J. M. Solowska, A. Karabay, S. Korulu, and P. W. Baas. 2008. "The microtubule-
severing proteins spastin and katanin participate differently in the formation of axonal
branches." *Mol Biol Cell* 19 (4): 1485-98.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Cita
tion&list_uids=18234839

Zhu, H., H. Y. Lee, Y. Tong, B. S. Hong, K. P. Kim, Y. Shen, K. J. Lim, F. Mackenzie, W. Tempel, and
H. W. Park. 2012. "Crystal structures of the tetratricopeptide repeat domains of kinesin
light chains: insight into cargo recognition mechanisms." *PLoS One* 7 (3): e33943.
https://doi.org/10.1371/journal.pone.0033943.
https://www.ncbi.nlm.nih.gov/pubmed/22470497.
Ziv, L., A. Muto, P. J. Schoonheim, S. H. Meijsing, D. Strasser, H. A. Ingraham, M. J. Schaaf, K. R.
Yamamoto, and H. Baier. 2013. "An affective disorder in zebrafish with mutation of the
glucocorticoid receptor." *Mol Psychiatry* 18 (6): 681-91.
https://doi.org/10.1038/mp.2012.64.
https://www.ncbi.nlm.nih.gov/pubmed/22641177.

1633

Fig. 1

A heptad repeats — KHC binding | LFP | TPR domains (1 2 3 4 5 6) | CTD — Lipid Binding

Autoinhibition — Cargo-Binding

klc4^uw314 25 bp mutation

H. sapiens 19 bp deletion

WT NQWLRDELANTQHKVQKSEQSVAQLEEEKKHL
uw314 NQWLRDELANTQHKVQKRRRRNIWSS*

B klc4

2 6 12 18 22 24 28 hpf

short

long

Fig. 2

A klc4+/+

B klc4uw314

C

D klc4+/+

E klc4uw314

F

# of Axons Crossing Region

klc4+/+
klc4uw314
*

Dorsal 1 2 3 4 5 6 7 8 Ventral

G

# of Axons Crossing Region

2% DMSO
Kinesore
*

Dorsal 1 2 3 4 5 6 7 8 Ventral

H 2% DMSO

I Kinesore

Fig. 3

Fig. 4

A klc4$^{+/+}$ 0° 0° 45° -45° 90° A P

B klc4^{uw314} A P

C klc4$^{+/+}$ klc4^{uw314} 600 A.U. 400 200 ventral ventral 89.5 50 0 -50 -89.5 anterior posterior **Angle**

D klc4$^{+/+}$ klc4^{uw314} ** 1.5 1.0 0.5 0.0 ventral 0 -20 -40 -60 -80 **posterior angles**

Fig. 5

Fig. 7

A klc4+/+
tg(ngn:gfp-caax)

B klc4uw314
tg(ngn:gfp-caax)

C

% of neurons/embryo with midline crossing

klc4+/+   klc4uw314   DMSO   Kinesore

D klc4uw314

E DMSO

F Kinesore

Fig 8.

A

A'

Distance →

Time ↓

Prox   Dist

B % of Motile Vesicles Traveling Anterogradely

**

% of Motile Vesicles

$klc4^{+/+}$ $klc4^{uw314}$

+/+ Central
+/+ Peripheral
uw314 Central
uw314 Peripheral

C Anterograde Run Length (Peripheral)

Distance (μm)

ns   *

+/+ uw314   +/+ uw314
Average   Maximum

D Peripheral

Velocity (μm/sec)

ns   ns

+/+ uw314   +/+ uw314
Anterograde   Retrograde

Fig. 9

Fig 10.

A, A', A'' tRFP-caax, EB3-GFP

A''' Distance / Prox Dist / Time / EB3-GFP

B — Comet frequency (comets/um/min) — ns — $klc4^{+/+}$, $klc4^{uw314}$

C, C', C'' AcTub, HNK-1, +/+, AcTub

D, D', D'' uw314

E — % Nascent Branches with AcTub Signal — **** — $klc4^{+/+}$, $klc4^{uw314}$

Fig. 11

Fig. 12