The longevity response to warm temperature is neurally controlled via regulation of

collagen genes

Sankara Naynar Palani[1,#], Durai Sellegounder[1,#], and Yiyong Liu[1,2*]

[1]Department of Translational Medicine and Physiology, Elson S. Floyd College of Medicine,

Washington State University, Spokane, Washington, USA

[2]Genomics Core, Washington State University, Spokane, Washington, USA

[#]These authors contributed equally

*Correspondence: yiyong.liu@wsu.edu

ABSTRACT

Lower temperatures have been associated with longer lifespan, while higher temperatures with

shorter lifespan. These inverse effects of temperature on longevity were traditionally explained

by the "rate of living theory", which posits that higher temperatures increase chemical reaction

rates, thus speeding up the aging process. Recent studies have identified specific molecules and

cells that mediate the longevity response to temperature, suggesting that the temperature effects

on aging are not simply thermodynamic but regulated processes. However, the mechanisms

underlying such regulation are not well understood. In the current study, we found that

Caenorhabditis elegans lacking NPR-8, a neuronal GPCR related to the mammalian

neuropeptide Y receptors, exhibited extended lifespan at warm temperature. Such lifespan

extension can be suppressed by re-expression of NPR-8 in the amphid sensory neurons AWB

and AWC or by inactivation of NPR-8-regulated collagen genes. These results suggest that the

warm temperature effect on longevity is regulated by the nervous system through controlling

collagen gene expression. Our study potentially provides mechanistic insights into understanding

the relationship between temperature and longevity, which could prove useful in mitigating

negative impacts of increasing temperature due to global warming.

INTRODUCTION

Various studies in many species have associated lower temperatures with longer lifespan and

higher temperatures with shorter lifespan, aside from extreme temperatures that would severely

harm organisms' survival (1). The inverse effects of temperature on longevity were first recorded

a century ago in poikilotherms such as the fruit fly *Drosophila melanogaster* (2) and the

cladoceran crustacean *Daphnia magna* (3). This phenomenon was later also found in the

nematode *Caenorhabditis elegans* (4), the South American fish *Cynolebias* (5), as well as in

mice (6) and humans (7), although with some obscurity in the mammals (1). Nonetheless, there

is a clear trend for increases in lifespan from decreases in temperature across species. These

inverse effects of temperature on longevity were traditionally explained by the "rate of living

theory", which posits that lower temperatures reduce chemical reaction rates, thus slowing down

the aging process, whereas higher temperatures do the opposite (8). Recent studies have

identified specific molecules and cells that mediate the longevity responses to temperature,

suggesting that the temperature effects on aging are not simply thermodynamic but regulated

processes (9). For example, in *C. elegans*, the cold-sensitive TRP channel TRPA-1 detects

temperature drop in the environment to extend lifespan (10), but it also mediates lifespan

shortening when the worm was subjected to low-temperature treatment at the larval stage (11).

The thermosensory AFD neurons and the AIY interneurons are required for maintaining *C.*

elegans lifespan at warm temperature (12), which was later found to be controlled by the worm

cyclic AMP-responsive element binding protein (CREB) (13). Co-chaperone p23 regulates the

C. elegans longevity responses to both cold temperatures and warm temperatures (14). A

transcriptional coregulator, Mediator 15, controls lipid homeostasis to regulate low-temperature-

induced longevity and proteostasis (15). These studies suggest that the longevity response to

temperature is a complex and highly regulated process and warrants major research efforts to

elucidate the underlying mechanisms.

In the current study, we found that *C. elegans* lacking NPR-8, a G protein-coupled receptor

(GPCR) related to the mammalian neuropeptide Y receptors, exhibited extended lifespan at

warm temperature. This extended lifespan phenotype was rescued by re-expressing NPR-8 in the

amphid sensory neurons AWB and AWC or by inactivating NPR-8-regulated collagen genes.

Our study suggests that the warm temperature effect on longevity is regulated by the nervous

system through controlling the expression of collagen genes.

RESULTS

Functional loss of NPR-8 extends *C. elegans* lifespan at warm temperature

Three temperatures (15°C, 20°C, and 25°C) are routinely used for *C. elegans* propagation in the

laboratory, and their inverse effects on lifespan have been well documented, i.e., a 75% increase

in lifespan from a 5°C drop in temperature (4). During our work on neural regulation of *C.*

elegans defense against pathogen infection (16), we unexpectedly observed that under

noninfectious conditions, *C. elegans* lacking NPR-8 ((*npr-8(ok1439)* animals) lived longer than

wild-type *N2* animals at 25°C, whereas the mutant and wild-type animals had the same lifespan

at 20°C or 15°C (Fig. 1). To determine whether the observed extended lifespan phenotype at

warm temperature was due to the deletion mutation in *npr-8*, we performed lifespan assays at

25°C using the JRS17 rescue strain in which *npr-8* is re-expressed under its native promoter in

npr-8(ok1439) animals. Our results showed that re-expression of *npr-8* in JRS17 rescued the

extended lifespan phenotype of the mutants (Fig. S1), indicating that the deletion mutation in

npr-8 is indeed responsible for the extended longevity at warm temperature. These results

suggest that NPR-8 suppresses lifespan at warm temperature, contributing to the inhibitory effect

of warm temperature on longevity.

NPR-8-dependent longevity response to warm temperature is neurally regulated

NPR-8 is a neuronal GPCR that is expressed in the amphid sensory neurons AWB, ASJ and

AWC (16). Our previous study showed that NPR-8 functions in these neurons to regulate *C.*

elegans defense response to pathogen infection (16). Is NPR-8-dependent longevity response to

warm temperature also neurally controlled? To address this question, we took advantage of our

established rescue strains that specifically express NPR-8 in the individual neurons under cell-

specific promoters in *npr-8(ok1439)* animals. Lifespan assays showed that NPR-8 rescue in

AWB (strain JRS18) and AWC neurons (strain JRS19) shortened the lifespan of *npr-8(ok1439)*

animals at 25°C, whereas the rescue in ASJ neurons (strain JRS20) had no effects on the

mutants' lifespan (Fig. 2). These results indicate that the lack of NPR-8 in AWB and AWC

neurons is responsible for the mutants' extended lifespan at 25°C. In other words, NPR-8

functions in AWB and AWC neurons to regulate the longevity response to warm temperature.

NPR-8-regulated collagen genes contribute to the warm temperature-induced longevity

response

Previously, we demonstrated that NPR-8 suppresses the expression of collagen genes in wild-

type animals under both normal (i.e., fed worm food *Escherichia coli* strain OP50) and infectious

(i.e. exposed to *P. aeruginosa*) conditions, and that NPR-8-regulated collagens positively

influence *C. elegans* defense against infection (16). In the current study, we wanted to determine

whether NPR-8-regulated collagen genes also play a role in warm temperature-induced longevity

response. To this end, we inactivated NPR-8-regulated collagen genes either by RNA

interference or by loss-of-function mutants and then examined the effects of such inactivation on

worm lifespan at 25°C. Our results showed that RNAi of *col-49* or *col-77* fully rescued the

extended lifespan of *npr-8(ok1439)* animals to the wild-type level, while a null mutation in *col-*

179 shortened the lifespan of *npr-8* mutant animals to a level lower than the wild-type level (Fig.

3). These results indicate that NPR-8-regulated collagen genes contribute to the extended

lifespan of *npr-8(ok1439)* animals at 25°C, and that NPR-8 controls the longevity response to

warm temperature by regulating collagen genes.

DISCUSSION

In the current study, we showed that *C. elegans* lacking NPR-8 had longer lifespan than wild-

type animals at 25°C but had the same lifespan as wild type at 20°C or 15°C, and that NPR-8

expression in amphid sensory neurons AWB and AWC as well as NPR-8-regulated collagen

genes suppress the lifespan extension of *npr-8* mutant animals at 25°C. An NPR-8-dependent

neural-longevity regulatory circuit has emerged whereby NPR-8 functions in AWB and AWC

neurons to control the longevity response to warm temperature by regulating the expression of

collagen genes. Many questions remain unanswered in this regulatory circuit. For example, how

do NPR-8-regulated collagen genes influence longevity at warm temperature? What other

neurons, in addition to AWB and AWC, are involved in the regulation? These are the subjects of

our future investigation.

Previously, we discovered an NPR-8-dependent neural-immune regulatory circuit that controls

C. elegans defense against pathogen infection (16). This circuit shares many components with

the above-described neural-longevity regulatory circuit, including NPR-8, AWB and AWC

neurons, and NPR-8-regulated collagen genes. Because both warm temperature and pathogen

infection are stressful for *C. elegans*, the two regulatory circuits might function in the same

pathway to maintain worm lifespan in response to stressful conditions, and they both achieve

extended lifespan by regulating the expression of collagen genes. Indeed, recent research has

shown that both collagen production and function decline with age, and that elevated collagen

expression is a shared feature of multiple conserved longevity pathways as well as essentially

every longevity intervention (17-20). These studies indicate that collagen expression might play

a causal role in slowing down the aging process under both normal and stressful conditions. This

notion is in direct contrast to the general belief that collagen decline is consequential to aging.

Such concept switch has profound implications for how we study aging and develop lfespan-

extending treatments. For example, as wrinkled skin is an indicator of collagen decline, healthy

skin can be used as an indicator of collagen homeostasis. Because collagen expression is causal

to the slowdown of aging, healthy skin could be used as a readout of an individual's overall good

health. This opens avenues for adapting a diverse array of therapies, ingredients, and

technologies that have been established in the skin care industry to restore skin to its optimal

condition into anti-aging strategies.

MATERIALS AND METHODS

Worm strains

Worm strains used in the study were Wild-type (N2 Bristol), *npr-8 (ok1439), npr-8 (ok1449),*

ttx-1 (p767), ttx-3 (ks5), npr-8;ttx-1(JRS--). NPR-8 translational rescue strains were JRS17 [*npr-*

8p(2kb)::npr-8 gDNA::gfp] (native NPR-8 rescue strain), JRS18 [*str-1p(1.3kb)::npr-8*

gDNA::gfp] (AWB NPR-8 rescue strain), JRS19 [*str-2p(2kb)::npr-8 gDNA::gfp*] (AWC NPR-8

rescue strain), and JRS20 [*trx-1p(1.1kb)::npr-8 gDNA::gfp*] (ASJ NPR-8 rescue strain) also used

in this study. All strains were maintained at 20°C and fed with *E.coli* OP50 on SB medium.

Lifespan assay

Temperature-associated lifespan assays were carried out at 15°C, 20°C, and 25°C. Normally

worms were maintained at 20°C. Worms used for lifespan assays at 15°C and 25°C were

maintained two generations at respective temperature before they are used for egg

synchronization. For worm synchronization, gravid adult worms were allowed to lay eggs for 1

hr at their assay temperature. 3x20 gravid adult worms were used for egg synchronization per

worm strain. Egg to adult developmental time window was followed as 90hr at 15°C, 65hr at

20°c, for all strains. 48hr at 25°c for all strains except *npr-8 (ok1439)*. At 25°C egg to adult

developmental time window was set as 56hr for *npr-8 (ok1439). E. coli OP50* lawn was prepared

by spreading a 50µl drop of an overnight culture of bacteria on SB medium (3.5-cm-diameter

petri plates used). Plates were incubated at 37°C for 16 hours and then cooled down at room

temperature for at least 1 hour before seeding with synchronized worms. A lifespan assay starts

with synchronized population of 20 adult worms per strain on live *E.coli* OP50. All assays were

performed in triplicates for each worm strain. Worms counted and transferred to fresh live *E. coli*

OP50 culture every day till they die.

RNA interference

The impact of NPR-8-regulated collagen genes expression on *C. elegans* lifespan was examined

by RNA interference by feeding. Collagen genes *col-49, col-77, col-139, col-179,* and *rol-1* were

silenced in wild-type and *npr-8 (ok1439)* animals at 20°C and 25°C. The RNAi protocol was

performed as previously described (16). Temperature-specific worm cultivation, egg-

synchronization time window, and temperature-specific egg to adult development time window

were followed as mentioned in lifespan assay.

Statistical analysis

Lifespan assay curves were plotted using GraphPad PRISM (version 8) computer software.

Lifespan was considered different from the appropriate control indicated in the main text when P

<0.05. PRISM uses the product limit or Kaplan-Meier method to calculate survival fractions and

the log-rank test, which is equivalent to the Mantel-Haenszel test, to compare survival curves.

All experiments were repeated at least three times, unless otherwise indicated.

ACKNOWLEDGEMENTS

We thank Dr. Jingru Sun at Washington State University for providing us with experimental

reagents, worm strains, and technical support. Some worm strains were provided by the

Caenorhabditis Genetics Center, which is funded by the NIH Office of Research Infrastructure

Programs (P40 OD010440).

FUNDING

This work was supported by the Department of Translational Medicine and Physiology, Elson S.

Floyd College of Medicine, WSU-Spokane (Startup to Y.L.). The funder had no role in study

design, data collection and interpretation, or the decision to submit the work for publication.

AUTHOR CONTRIBUTIONS

S.N.P., D.S., and Y.L. designed and performed experiments and analyzed data. S.N.P. and Y.L.

wrote the paper.

COMPETING INTERESTS

The authors declare that they have no competing interests.

DATA AND MATERIALS AVAILABILITY

All data needed to evaluate the conclusions in the paper are present in the paper and/or the

Supplementary Materials. Additional data related to this paper may be requested from the

authors. The *C. elegans* strains and plasmids constructed by us and primer sequences used for

their construction are available upon request.

REFERENCES

1.   Keil, G., Cummings, E., and de Magalhaes, J. P. (2015) Being cool: how body

temperature influences ageing and longevity. *Biogerontology* **16**, 383-397

2.   Loeb, J., and Northrop, J. H. (1916) Is There a Temperature Coefficient for the Duration

of Life? *Proceedings of the National Academy of Sciences of the United States of*

America **2**, 456-457

3.   Macarthur, J. W., and Baillie, W. H. (1929) Metabolic activity and duration of life. I.

209   Influence of temperature on longevity in Daphnia magna. *Journal of Experimental*

210   *Zoology* **53**, 221-242

211   4.   Van Voorhies, W. A., and Ward, S. (1999) Genetic and environmental conditions that

212   increase longevity in Caenorhabditis elegans decrease metabolic rate. *Proceedings of the*

213   *National Academy of Sciences of the United States of America* **96**, 11399-11403

214   5.   Walford, R. L., and Liu, R. K. (1965) Husbandry, life span, and growth rate of the annual

215   fish, Cynolebias adloffi E. Ahl. *Experimental Gerontology* **1**, 161-168

216   6.   Conti, B., Sanchez-Alavez, M., Winsky-Sommerer, R., Morale, M. C., Lucero, J.,

217   Brownell, S., Fabre, V., Huitron-Resendiz, S., Henriksen, S., Zorrilla, E. P., de Lecea, L.,

218   and Bartfai, T. (2006) Transgenic mice with a reduced core body temperature have an

219   increased life span. *Science* **314**, 825-828

220   7.   Rikke, B. A., and Johnson, T. E. (2004) Lower body temperature as a potential

221   mechanism of life extension in homeotherms. *Exp Gerontol* **39**, 927-930

222   8.   Demetrius, L. A. (2013) Boltzmann, Darwin and Directionality theory. *Physics Reports*

223   **530**, 1-85

224   9.   Kim, B., Lee, J., Kim, Y., and Lee, S. V. (2020) Regulatory systems that mediate the

225   effects of temperature on the lifespan of Caenorhabditis elegans. *J Neurogenet* **34**, 518-

226   526

227 10.   Xiao, R., Zhang, B., Dong, Y., Gong, J., Xu, T., Liu, J., and Xu, X. Z. (2013) A genetic

228   program promotes C. elegans longevity at cold temperatures via a thermosensitive TRP

229   channel. *Cell* **152**, 806-817

230   11.   Zhang, B., Xiao, R., Ronan, E. A., He, Y., Hsu, A. L., Liu, J., and Xu, X. Z. (2015)

231   Environmental Temperature Differentially Modulates C. elegans Longevity through a

232   Thermosensitive TRP Channel. *Cell reports* **11**, 1414-1424

233   12.   Lee, S. J., and Kenyon, C. (2009) Regulation of the longevity response to temperature by

234   thermosensory neurons in Caenorhabditis elegans. *Current biology : CB* **19**, 715-722

235   13.   Chen, Y. C., Chen, H. J., Tseng, W. C., Hsu, J. M., Huang, T. T., Chen, C. H., and Pan,

236   C. L. (2016) A C. elegans Thermosensory Circuit Regulates Longevity through crh-

237   1/CREB-Dependent flp-6 Neuropeptide Signaling. *Developmental cell* **39**, 209-223

238   14.   Horikawa, M., Sural, S., Hsu, A. L., and Antebi, A. (2015) Co-chaperone p23 regulates

239   C. elegans Lifespan in Response to Temperature. *PLoS genetics* **11**, e1005023

240   15.   Lee, D., An, S. W. A., Jung, Y., Yamaoka, Y., Ryu, Y., Goh, G. Y. S., Beigi, A., Yang, J.

241   S., Jung, G. Y., Ma, D. K., Ha, C. M., Taubert, S., Lee, Y., and Lee, S. V. (2019) MDT-

242   15/MED15 permits longevity at low temperature via enhancing lipidostasis and

243   proteostasis. *PLoS biology* **17**, e3000415

244   16.   Sellegounder, D., Liu, Y., Wibisono, P., Chen, C. H., Leap, D., and Sun, J. (2019)

245   Neuronal GPCR NPR-8 regulates *C. elegans* defense against pathogen infection. *Sci Adv*

246   **5**, eaaw4717

247   17.   Myllyharju, J., and Kivirikko, K. I. (2004) Collagens, modifying enzymes and their

248   mutations in humans, flies and worms. *Trends in genetics : TIG* **20**, 33-43

249   18.   Toyama, B. H., and Hetzer, M. W. (2013) Protein homeostasis: live long, won't prosper.

250   *Nature reviews. Molecular cell biology* **14**, 55-61

251   19.   Varani, J., Dame, M. K., Rittie, L., Fligiel, S. E., Kang, S., Fisher, G. J., and Voorhees, J.

252   J. (2006) Decreased collagen production in chronologically aged skin: roles of age-

dependent alteration in fibroblast function and defective mechanical stimulation. *Am J*

Pathol **168**, 1861-1868

20.   Ewald, C. Y., Landis, J. N., Porter Abate, J., Murphy, C. T., and Blackwell, T. K. (2015)

Dauer-independent insulin/IGF-1-signalling implicates collagen remodelling in

longevity. *Nature* **519**, 97-101

FIGURE LEGENDS

Fig. 1. Functional loss of NPR-8 extends lifespan at 25°C but has no effect on lifespan at

20°C or 15°C. WT and *npr-8(ok1439)* animals were grown on *E. coli* strain OP50 at 15°C **(A)**,

20°C **(B)**, or 25°C **(C),** and scored for survival over time. The graphs are the combined results of

three independent experiments. Each experiment included $n = 60$ adult animals per strain. *p*-

values represent the significance level of the mutants relative to the WT, $p = 0.7416$ in (A), $p =$

0.1460 in (B), and $p < 0.0001$ in (C).

Fig. 2. NPR-8 functions in neurons to regulate the longevity response to warm temperature.

WT, *npr-8(ok1439)*, and rescue animals were grown at 25°C and scored for survival over time.

The graphs are the combined results of three independent experiments. Each experiment

included $n = 60$ adult animals per strain. JRS18, *npr-8* rescue in AWB neurons; JRS19, *npr-8*

rescue in AWC neurons; JRS20, *npr-8* rescue in ASJ neurons; JRS21, *npr-8* pan-neuronal

rescue. *p*-values are relative to *npr-8(ok1439):* WT, $p < 0.0001$; JRS18, $p < 0.0001$; JRS19, $p <$

0.0001; JRS20, $p = 0.3476$.

Fig. 3. NPR-8-regulated collagen genes are required for warm temperature-induced

lifespan extension in *npr-8* mutant animals. (A) WT and *npr-8(ok1439)* animals were grown

on dsRNA for empty vector (control), *col-49*, *col-77*, or *col-139* at 25°C and scored for survival

over time. The graphs are the combined results of three independent experiments. Each

experiment included $n = 60$ adult animals per strain. *p*-values are relative to *npr-8(ok1439)*+EV:

WT+EV, $p < 0.0001$; *npr-8 (ok1439)+col-49*, $p < 0.0001$; *npr-8 (ok1439)+col-77*, $p < 0.0001$;

npr-8 (ok1439)+col-139, $p = 0.0527$. **(B)** WT, *npr-8 (ok1439)*, *col-179 (ok3010)*, and *col-*

179;npr-8(JRS30) animals were grown on at 25°C and scored for survival over time. The graphs

are the combined results of three independent experiments. Each experiment included $n = 60$

adult animals per strain. *p*-values are relative to *npr-8 (ok1439)*: WT, $p < 0.0001$; *col-179*

(ok3010), $p < 0.0001$; *col-179;npr-8 (JRS30)*, $p < 0.0001$.

SUPPLEMENTARY DATA

Fig. S1. Functional loss of NPR-8 extends lifespan at 25°C. WT, *npr-8(ok1439)*, JRS17 (*npr-8*

expression restored in *npr-8(ok1439)* under its native promoter) animals were grown on *E. coli*

strain OP50 at 25°C and scored for survival over time. The graphs are the combined results of

three independent experiments. Each experiment included $n = 60$ adult animals per strain. *p*-

values are relative to *npr-8(ok1439)*: WT, $p < 0.0001$; JRS17, $p < 0.0001$.

FIGURES

Fig. 1

Fig. 1. Functional loss of NPR-8 extends lifespan at 25°C but has no effect on lifespan at

20°C or 15°C. WT and *npr-8(ok1439)* animals were grown on *E. coli* strain OP50 at 15°C **(A)**,

20°C **(B)**, or 25°C **(C)**, and scored for survival over time. The graphs are the combined results of

three independent experiments. Each experiment included *n* = 60 adult animals per strain. *p*-

values represent the significance level of the mutants relative to the WT, *p* = 0.7416 in (A), *p* =

0.1460 in (B), and *p* < 0.0001 in (C).

Fig. 2
315

316
Fig. 2. NPR-8 functions in neurons to regulate the longevity response to warm temperature.
WT, *npr-8(ok1439)*, and rescue animals were grown at 25°C and scored for survival over time.
The graphs are the combined results of three independent experiments. Each experiment
included *n* = 60 adult animals per strain. JRS18, *npr-8* rescue in AWB neurons; JRS19, *npr-8*
rescue in AWC neurons; JRS20, *npr-8* rescue in ASJ neurons; JRS21, *npr-8* pan-neuronal
rescue. *p*-values are relative to *npr-8(ok1439):* WT, $p < 0.0001$; JRS18, $p < 0.0001$; JRS19, $p <$
0.0001; JRS20, $p = 0.3476$.
324
325

Fig. 3

Fig. 3. NPR-8-regulated collagen genes are required for warm temperature-induced lifespan extension in *npr-8* mutant animals. (A) WT and *npr-8(ok1439)* animals were grown on dsRNA for empty vector (control), *col-49*, *col-77*, or *col-139* at 25°C and scored for survival over time. The graphs are the combined results of three independent experiments. Each experiment included $n = 60$ adult animals per strain. *p*-values are relative to *npr-8(ok1439)*+EV: WT+EV, $p < 0.0001$; *npr-8 (ok1439)+col-49*, $p < 0.0001$; *npr-8 (ok1439)+col-77*, $p < 0.0001$; *npr-8 (ok1439)+col-139*, $p = 0.0527$. **(B)** WT, *npr-8 (ok1439)*, *col-179 (ok3010)*, and *col-179;npr-8(JRS30)* animals were grown on at 25°C and scored for survival over time. The graphs are the combined results of three independent experiments. Each experiment included $n = 60$ adult animals per strain. *p*-values are relative to *npr-8 (ok1439)*: WT, $p < 0.0001$; *col-179 (ok3010)*, $p < 0.0001$; *col-179;npr-8 (JRS30)*, $p < 0.0001$.

344 **SUPPLEMENTARY DATA**
345
346 **Fig. S1**
347

348
349 **Fig. S1. Functional loss of NPR-8 extends lifespan at 25°C.** WT, *npr-8(ok1439)*, JRS17 (*npr-8*
350 expression restored in *npr-8(ok1439)* under its native promoter) animals were grown on *E. coli*
351 strain OP50 at 25°C and scored for survival over time. The graphs are the combined results of
352 three independent experiments. Each experiment included *n* = 60 adult animals per strain. *p*-
353 values are relative to *npr-8(ok1439)*: WT, p < 0.0001; JRS17, p < 0.0001.
354
355
356