Pyramidal cell types drive functionally distinct cortical activity patterns during decision-making

Simon Musall[1,2,8], Xiaonan R. Sun[3,4,8], Hemanth Mohan[3,5], Xu An[3,5], Steven Gluf[3], Shujing Li[3], Rhonda Drewes[3],

Emma Cravo[2], Irene Lenzi[1,2], Chaoqun Yin[6], Björn M. Kampa[2,7], Anne K. Churchland[3,6]

[1]Institute of Biological Information Processing (IBI-3), Forschungszentrum Jülich, Jülich, Germany

[2]Department of Neurophysiology, Institute for Zoology, RWTH Aachen University, Aachen, Germany

[3]Cold Spring Harbor Laboratory, Neuroscience, Cold Spring Harbor, NY, USA

[4]Department of Neurosurgery, Zucker School of Medicine, Hofstra University, Hempstead, NY, USA

[5]Department of Neurobiology, Duke University Medical Center, Durham, NC

[6]Department of Neurobiology, David Geffen School of Medicine, University of California, Los Angeles, CA, USA.

[7]JARA Brain, Institute for Neuroscience and Medicine, Forschungszentrum Jülich, Germany

[8]These authors contributed equally

Abstract

Understanding how cortical circuits generate complex behavior requires investigating the cell types that comprise them. Functional differences across pyramidal neuron (PyN) types have been observed within cortical areas, but it is not known whether these local differences extend throughout the cortex, nor whether additional differences emerge when larger-scale dynamics are considered. We used genetic and retrograde labeling to target pyramidal tract (PT), intratelencephalic (IT) and corticostriatal projection neurons and measured their cortex-wide activity. Each PyN type drove unique neural dynamics, both at the local and cortex-wide scale. Cortical activity and optogenetic inactivation during an auditory decision task also revealed distinct functional roles: all PyNs in parietal cortex were recruited during perception of the auditory stimulus, but, surprisingly, PT neurons had the largest causal role. In frontal cortex, all PyNs were required for accurate choices but showed distinct choice-tuning. Our results reveal that rich, cell-type-specific cortical dynamics shape perceptual decisions.

Introduction

The neocortex is organized into discrete layers that form a vertically-arranged microcircuit motif. This core circuit is largely conserved across cortical areas with each layer consisting of distinct excitatory and inhibitory cell types that can be categorized based on genetic markers, cell morphology, anatomical projections or developmental lineage[1]. The precise interplay between these cell types is crucial for accurate cortical circuit function and their respective functional roles are the subject of intense study. Tremendous progress has been made particularly for cortical interneurons, where the availability of specific mouse driver lines has revealed the functional arrangement of inhibitory circuit motifs[2–4], e.g. for network synchronization[5–7] and state-dependent sensory processing[8–11]. However, the roles of glutamatergic pyramidal neuron (PyN) types are less well established, although PyNs comprise ~80% of all cortical neurons and form almost all long-range projections that enable the communication between local cortical circuits and other brain areas.

While often treated as a monolithic group, PyNs appear to be far more diverse than interneurons with at least one hundred putative subtypes indicated by RNA sequencing[12]. These molecular signatures are critical for categorizing PyNs and go far beyond layer identity as different

subtypes are often intermingled within the same layers[13–17]. PyNs can also be classified based on
their projection target. Long-range projection neurons are broadly categorized into two major
types: intratelencephalic (IT) neurons, projecting to other cortical structures and the striatum, and
pyramidal tract (PT) neurons, projecting to subcortical structures, such as the superior colliculus
(SC), thalamus, the pons and the striatum. Furthermore, PT and IT neurons differ in their
electrophysiological properties, dendritic morphology, local connectivity, and responsiveness to
sensory stimulation[15–17]. Recent studies in sensory cortex also showed that only PT but not IT
neurons are important for active perception of tactile or visual stimuli, suggesting that PT and IT
neurons encode separate streams of information[18,19]. Similar results have been found in
secondary motor cortex (M2), where specific PT neurons are involved in motor generation[13,20].
This suggests that the functional divergence of PyN types could be key for understanding cortical
microcircuits, with PT and IT neurons forming functionally-distinct, parallel subnetworks that
independently process different information. However, the functional tuning of individual PyNs
in frontal cortex is still best predicted by cortical area location instead of laminar location or
projection type[21]. Since PyN subtype activity has only been studied in single areas, it is therefore
not known whether PyN-specific subcircuits exist throughout the cortex or only in a subset of
cortical regions.

An ideal method to address this question is widefield calcium imaging, which allows measuring
neural activity across the dorsal cortex with cell-type specificity[22–24]. Interneuron-specific
widefield imaging revealed clear differences in the spatiotemporal dynamics of different
inhibitory cell types during an odor detection task[25]. However, cortex-wide studies of different
PyN types are lacking, in part due to the limited availability of PyN-specific driver lines[26–28].
Here, we used two novel knock-in mouse driver lines targeting PT or IT neurons[26] and
performed widefield Ca^{2+} imaging to measure PyN subtype-specific activity while animals
performed a perceptual decision-making task. Moreover, we developed a retrograde labeling
approach to selectively measure the activity of corticostriatal projection (CStr) neurons
throughout the dorsal cortex. Dimensionality-reduction and clustering analyses revealed unique
cortex-wide dynamics for each PyN subtype, suggesting the existence of specialized subcircuits.
Cortical dynamics of different PyNs were further segregated based on their role in decision-
making, with encoder and decoder approaches revealing the strongest stimulus- and choice-
related modulation in sensory, parietal and frontal cortices. This was confirmed by PyN-type-
specific inactivation experiments. In parietal cortex, PT neurons were most important for sensory
processing while all PyN types in frontal cortex were needed for choice formation and retention.
Taken together, our results demonstrate that different PyN-types exhibit functionally distinct,
cortex-wide neural dynamics with separate roles during perceptual decision-making.

Results

To monitor PyN-type specific neural activity throughout the dorsal cortex, we used two knock-in
inducible CreER lines that target developmentally-distinct classes of excitatory cortical neurons:
Fezf2-CreER targeting PT neurons and PlexinD1-CreER targeting IT neurons[26]. For PyN-type
specific GCaMP6s expression, both lines were crossed with the Ai162 reporter line[29] and CreER
activity was induced at four weeks of age. In each mouse line, GCaMP expression was restricted
to specific cortical layers and PyN types (Fig 1a,b). In Fezf2 mice, expression was concentrated
in layer 5b (Fig. 1a). We observed axonal projections to multiple subcortical regions, such as
striatum and the corticospinal tract, confirming that Fezf2 is a reliable marker for corticofugal PT
neurons. In PlexinD1 mice, GCaMP expression was restricted to layer 2/3 and layer 5a and
axonal projections were visible in the corpus callosum and the striatum (Fig. 1b). These results

are in agreement with earlier reports[26], confirming that PlexinD1 is a reliable marker for
intracortical and corticostriatal IT neurons.

After confirming non-overlapping expression patterns of PT and IT neurons, we measured cell-
type-specific cortical activity with widefield calcium imaging. All imaging data was aligned to
the Allen Common Coordinate Framework v3 (CCF)[25,30] to compare activity across individuals
and PyN types. Both lines yielded robust GCaMP-dependent fluorescence and we observed rich
neural dynamics throughout the dorsal cortex (Supp. Movies 1-3). We first used retinotopic
mapping to assess sensory responses of different PyN types and the resulting spatial arrangement
of visual areas. Both PT and IT neurons robustly responded to visual stimulation and we could
construct retinotopic maps that reliably indicated the location of known visual areas (Fig. 1c)[31].
Retinotopic maps were similar to those observed in an Ai93D;Emx-Cre;LSL-tTA mouse line
(EMX) that expressed GCaMP6f in all excitatory cortical neurons[32], suggesting that the
functional architecture of visual areas is comparable across PyN types. However, clear
differences were apparent in the modulation of cortical regions in the absence of visual
stimulation (Fig. 1d, same individual mice as in 1c). For example, the total variance of cortex-
wide activity was largest in parietal and frontal regions in PT neurons (Fig. 1d, left) while
variance was highest in visual and somatosensory regions of IT neurons (Fig. 1d, center). This
was comparable to variance in EMX mice, which showed additional modulation in retrosplenial
(RS) cortex (Fig. 1d, right). These different patterns were also highly consistent across individual
mice of the same PyN type (Supp. Fig. 1).

Figure 1. Knock-in mouse lines enable PyN subtype-specific recordings of cortex-wide neural activity.
a) GCaMP6s expression in Fezf2-CreER;Ai162 mice. Cortical labeling is largely confined to layer 5b. Axonal
projections were found in multiple subcortical regions such as the striatum and the corticospinal tract. (Scale bars:
100 μm). **b)** GCaMP6s expression in PlexinD1-CreER; Ai162 mice is widespread throughout the cortex and
restricted to superficial layers 2/3 and layer 5a. Axonal projections were found in striatum and corpus callosum but
absent in the corticospinal tract. **c)** Visual sign maps from retinotopic mapping experiments. IT and PT populations
showed clear retinotopic responses in primary and secondary visual areas where boundaries largely resembled
known areas that were also observed in nonspecific PyNs (EMX mice). **d)** Total variance maps from same mice as
in c), showing most modulated cortical regions in each PyN type.

The observed differences in variance could indicate general differences in the spatiotemporal
dynamics of cortex-wide activity across PyN types. To describe such differences, we performed
semi-nonnegative matrix factorization (sNMF), reducing the imaging data to a small number of
spatial- and temporal components that capture nearly all variance (>99%)[23,33]. Here,
dimensionality (the number of required components) is a measure of the overall complexity of
cortical dynamics. Surprisingly, PT neurons had much lower dimensionality compared to IT
neurons (Fig. 2a), suggesting that PT neurons are more functionally-homogeneous. This was also

seen in the correlations between cortical regions, which were consistently highest for PT neurons
(Supp. Fig. 2). A potential reason for this difference could be that IT neurons encompass a larger
number of specialized subtypes than PT neurons and may thus support a wider range of
functions[34].
130

131
132
Figure 2. PyN subtypes exhibit unique cortical activity patterns.
a) Number of sNMF components accounting for 99% of cortical variance in each PyN type. Violin plots show
individual mice, dots represent individual sessions. **b)** Examples of spatial sNMF components. Components from
different mice from the same PyN subtype (colored rectangles) strongly resembled each other. **c)** UMAP embedding
of spatial sNMF components for EMX (red), IT (green) and PT (blue) mice. Maps show example spatial
components for each type and their respective UMAP locations. Different markers denote individual mice. Blow-
ups show examples of PT-specific regions. Additional clustering for individual mice is sometimes evident (top left
region), but was generally weaker than PyN-specific clustering. **d)** Accuracy of a PyN type classifier based on
neighbor identity of individual components in UMAP space. Each data point represents the mean classification
accuracy over all components in one session. **e)** Map of seed regions used for LocaNMF analysis. Example spatial
components are compact and mostly confined to the appropriate seed region. **f)** Number of LocaNMF components,
accounting for 99% of cortical variance in each PyN type. Conventions as in (a). **g)** UMAP projection embedding of
spatial LocaNMF components. Conventions as in (c). Left: UMAP shows clustering of LocaNMF components from
similar regions (same 24 regions as shown in e). Right: Components within individual regions are further divided for

different PyN types. Maps show example LocaNMF components (I-III). **h)** Accuracy of a PyN type classifier, based
on individual LocaNMF components. Conventions as in (d). **i)** Top: Peak normalized distributions of area size for
PyN-type-predictive (blue) versus unspecific (red) LocaNMF components. PyN-predictive components are smaller
than unspecific components (PyN-predictive: median = 0.59 mm^2, n = 6317 components, nonspecific: median =
0.68mm^2, n = 18938 components; ranksum test: $p < 10^{-10}$). Bottom: Examples of PyN-predictive (I, II) and
unspecific (III) components in right parietal cortex.

Since PT neurons are a subset of EMX neurons, their lower dimensionality could indicate that
PT components are also a subset of EMX components. We therefore quantified to which extent
components from different PyN types are linearly overlapping or independent from each other.
For each PyN type, we computed the amount of variance that could be explained with
components from other mice of either the same or different PyN type (Supp. Fig. 3a). For all
groups, components from a similar PyN type explained significantly more variance (ΔR^2) than
the maximal amount that could be explained by other types (EMX, mean $\Delta R^2 = 3.3\%$, *t*-test: $p <$
10^{-10}; IT, mean $\Delta R^2 = 1.5\%$, $p < 10^{-10}$; PT, mean $\Delta R^2 = 1.2\%$, $p < 10^{-10}$). Each group therefore
contains additional PyN-type-specific variance that is independent from other types. This was
also visible in reconstructed data: PT components captured larger fluctuations in IT activity
(Supp. Movie 4) but failed to represent spatial details in the cortical activity patterns (Supp. Fig.
3b). Despite the similar dimensionality of IT and EMX activity (Fig. 2a), EMX ΔR^2 was also
similar for IT and PT components ($\Delta R^2_{EMX-IT} = 3.4\%$, $\Delta R^2_{EMX-PT} = 3.3\%$, $p = 0.1$), demonstrating
that EMX data was not largely dominated by either IT or PT neurons.

To directly compare spatial activity patterns, we then focused on spatial sNMF components.
Spatial components represent cortex-wide maps of positively correlated areas and we wondered
whether these correlation patterns would differ between PyN subtypes. Indeed, individual spatial
components from different mice of the same PyN subtype often strongly resembled each other
but differed from other PyN subtypes (Fig. 2b). To assess if most spatial components (or only a
small subset) was PyN-type-specific, we performed a UMAP projection of the first 20
components from all recordings and PyN types, non-linearly embedding the pixels of each
component in a 2-dimensional space (Fig. 2c)[35]. If neural activity in PT, IT and EMX mice
tended to exhibit similar correlation patterns, these spatial components would be mixed together.
Instead, in agreement with the notion that components of the same PyN type resembled each
other, we found that components formed clusters that were largely dominated by either PT or IT
neurons (green/blue markers). EMX neurons formed a third set of non-overlapping clusters,
likely reflecting the combined cortical dynamics from diverse PyN types beyond PT and IT
neurons that were contained in this larger group (red markers). A simple classifier reliably
identified each group, based on the nearest neighbors in a UMAP projection that was exclusively
based on data from other animals. Remarkably, prediction accuracy was high even when the
classification was based on just a single spatial component (Fig. 2d). The presence of such clear
clusters is striking given that the spatial components used in the analysis were pooled over many
sessions and mice. This confirms that UMAP clusters reflect consistent PyN-type-specific
activity patterns, rather than idiosyncratic differences within individual sessions or mice. Taken
together, these results clearly demonstrate that PyN types differ in the complexity of cortical
dynamics, contain independent variance, and exhibit unique cortex-wide correlation patterns.

An important concern is that variations in the density of Cre expression in each mouse line could
result in non-uniform GCaMP expression patterns which might contribute to PyN-specific spatial
components. However, GCaMP-related fluorescence was largely uniform across cortex in all
mice and we observed no clear relationship between raw widefield fluorescence patterns and
PyN-specific spatial components (Supp. Fig. 4a, b). Moreover, despite some fluctuations in the
density of Cre-expressing neurons across areas in both mouse lines, Cre-expression patterns did
not account for spatial components or specific differences between PyN types (Supp. Fig. 4c, d).

Nevertheless, particularly distinct activity of each PyN type in specific cortical areas could lead
to cortex-wide correlation patterns that are either dominated by highly active areas or where
inactive areas are 'missing'. In this case, the relatively low dimensionality of PT neurons might
be due to a lower number of active cortical areas. We therefore used a localized form of sNMF
(LocaNMF)[33] which extracts components that are dense and spatially restricted to a specific
cortical "seed" region (Fig. 2e). Analyzing LocaNMF components therefore allowed us to reveal
if PyN-specific differences mostly occur on a cortex-wide level (reflecting the interactions
between cortical areas) or further extend to the local level of individual cortical areas (e.g. due to
PyN-specific differences in the shape and localization of cortical areas).
205
First, we again compared the number of components required to explain at least 99% of the
variance in each PyN type. As expected, the number of components was greater than sNMF and,
interestingly, was also more similar across PyN subtypes (Fig. 2f). This shows that PyN-type-
specific differences in cortex-wide correlation patterns are not due to a lack of activity in some
cortical areas (which would have resulted in a lower number of required components, e.g. in PT
mice) and instead result from PyN-type specific differences in the coordinated activation of
multi-area cortical networks. Moreover, as with sNMF components, LocaNMF components from
mice of the same PyN type explained more variance than components from other types (Supp.
Fig. 3c).
215
Repeating the UMAP embedding for these local components also uncovered PyN-type-specific
clustering (Fig. 2g), which could be used to accurately identify each PyN type (Fig. 2h).
Classification accuracy was high across most cortical regions (Supp. Fig. 4d). Specificity of
LocaNMF components could either be due to the presence of specific 'subregions', where PyN
types are most active in smaller parts of a given cortical area, or larger 'superregions', where the
activity of a specific PyN type extends across known area borders. We therefore compared the
size of LocaNMF components that accurately predicted their respective PyN type (classifier
accuracy > 99%) versus nonspecific components. Interestingly, PyN-predictive clusters were
significantly smaller than nonspecific clusters (Fig. 2i), suggesting that different PyNs might be
most active in distinct subregions instead of larger multi-area components. This indicates that
smaller, PyN-specific subregions may reside within the coarser, traditionally-defined cortical
areas.
We next assessed how cortical dynamics of PyN subpopulations are related to decision-making
by imaging PyN subtype activity in animals trained on an auditory decision-making task (Fig.
3a)[36]. Mice initiated trials by touching small handles, which triggered the simultaneous
presentation of sequences of clicking sounds to their left and right side. After a short delay
period, mice reported decisions by licking one of two water spouts and were rewarded for
choosing the side where more clicks were presented. To reduce temporal correlations between
task events, such as trial initiation and stimulus onset, the durations of the initiation, stimulus and
delay periods were randomly varied across trials. In all mice, decisions varied systematically
with stimulus strength (Fig. 3b) and were equally affected by click sounds throughout the
stimulus period (Supp. Fig. 5).
As with spatial components, trial-averaged temporal sNMF and LocaNMF components showed
pronounced clustering, suggesting that different PyN-types also exhibit distinct task-related
temporal dynamics (Supp. Fig. 6). Correspondingly, we found clear differences in trial-averaged
neural activity between PyN subtypes, especially during stimulus presentation (Fig. 3c). Here,
cortical activity was uniformly suppressed in EMX mice, partially suppressed in somatosensory
and visual areas in IT mice, and uniformly increased in PT mice (Fig. 3d). In all PyN types,

activity was largely symmetric across the left and right hemispheres, even when only analyzing
trials where stimuli were presented on the left side and the animal made a corresponding leftward
choice (Fig. 3c; Supp. Fig. 7). Moreover, responses to sensory stimuli were much weaker than
movement-related responses, such as trial initiation or licking (Fig. 3d, gray bar versus dashed
lines). A potential explanation is that lateralized, task-related activity is obscured by cortical
activity due to animal movements[32,37–39].

Figure 3. An auditory decision-making task reveals distinct functional activity patterns in each PyN type.
a) Auditory discrimination task structure of an example trial. Mice touched paw handles to initiate randomized click
sequences on the left and/or right side. After a delay period, a lick response on the correct side was rewarded with
water. The episode duration was randomly varied in individual trials. **b)** Psychometric functions fit to behavioral
data from the discrimination task in (a) of individual EMX (red), IT (green) and PT (blue) mice. **c)** Trial-averaged
response maps for all correct, leftward trials in different PyN types. Shown are averages for the 'Initiation',
'Stimulus', 'Delay' and 'Response' periods shown in (a). **d)** Averaged activity in auditory (left) and frontal cortex
(right) for each PyN type. Averages were separately aligned to each of the four trial periods, indicated by short gaps.
Left dashed line: time of initiation, gray box: stimulus presentation, right dashed line: animal's response. Traces
show standard deviation units (SDU). White dashed circles in (c) show respective area locations. Colors as in (b).
Shading shows s.e.m.; n = 4 EMX/IT mice and 5 PT mice.

To isolate task-related activity from movements we used a linear encoding model[32]. The model
combines many task- and movement-variables to predict single-trial fluctuations in cortical
activity (Fig. 4a, top). Task variables were related to the learned behavior, and included past and

current choices, and the presentation of a sensory stimulus. Movement variables included licking
and touching the handles, alongside facial movements (whisking or nose movements) obtained
from two video cameras of the animal's face and body (a complete list of all task- and movement
variables is shown in Supp. Table 1). All variables were combined into a single design matrix
and we used ridge regression to fit the model to the imaging data. This produced a time-varying
event kernel for each variable, allowing us to isolate how specific task variables (e.g. the sensory
stimulus, Fig. 4a, bottom) are related to neural activity while separating them from the impact of
other factors, such as licking movements, which are captured by other model variables.

To assess how well the model predicts cortical activity, we first computed the tenfold cross-
validated R^2 (cvR^2). For all PyN types, the model successfully captured a large fraction of single-
trial variance throughout the cortex (Fig. 4b), confirming the validity of our approach. Predicted
variance was higher in IT and PT mice compared to EMX mice, with over 90% explained
variance in frontal cortex of PT mice. Consistent with earlier results[32,37], movements consistently
captured more variance than task variables across all PyNs (Fig. 4c). The impact of movements
was also apparent when computing the unique contributions from movement or task variables by
removing either set of variables from the full model, demonstrating that movements are an
important driver of cortex-wide activity in all PyN types (Supp. Fig. 8a). Interestingly, unique
contributions of many task variables were higher for IT, further indicating that they might be
more functionally diverse than PT neurons (Supp. Fig. 8b).

Figure 4. An encoding model uncovers task-specific differences across PyN types.
a) Schematic of the encoding model. Top: Task- and movement-variables account for fluctuations in cortical
activity. Bottom: Weights for each variable define a spatiotemporal event kernel, revealing cortical activity in
response to a specific event (example shows right stimulus kernel in EMX mice). **b)** Average maps of cvR^2. The
model accurately predicted cortical variance for all PyNs. **c)** cvR^2 from two models, using only movement
('Movement') or task variables ('Task'). In all groups, movements were more predictive than task variables and
accounted for the majority of the full models explained variance (dashed lines). Circles denote sessions. **d)**
Contributions of movements, stimulus and choice to the model's total explained variance. Although movements
contributed the most, stimulus and choice also made sizable contributions.

To gain insight into the cortical dynamics related to stimulus and choice, we focused on their
respective event kernels learned by the model for each PyN type. To ensure that both task
variables still accounted for a sizable amount of the observed neural activity (instead of all
variance being explained by movements), we first computed how much variance the stimulus
and choice kernels contribute to the full model compared to the sum of all movement variables

(Fig. 4d). Here, although the combined movement variables made the largest contributions
(~60% of the model's total explained variance), both stimulus and choice also made sizable
contributions to the model (10-20% explained variance in all PyN types). This demonstrates that
stimulus and choice remain important for understanding cortical activity patterns and can be
leveraged to selectively isolate task-related activity.

We then focused on cortical responses to the auditory stimulus. In contrast to trial averages of
ΔF/F (Fig. 3c), the stimulus kernels in EMX mice uncovered clear sensory-locked and lateralized
responses in auditory, parietal and frontal cortex (Fig. 4a, bottom; Fig. 5a, top). Other areas, such
as somatosensory and visual cortex, were inhibited. Sensory-locked responses were also present
in auditory, parietal and frontal cortices of PT and IT mice (Fig. 5a, center and bottom).
However, the cortex-wide response patterns were not identical: for instance, no inhibition was
apparent in PT mice. Sensory responses were particularly distinct in the parietal cortex (Fig. 5b).
Responses in EMX and IT mice were more anterolateral in parietal area A while PT responses
were surprisingly prominent and more posteromedial at the border between area AM and RS.

Figure 5. PyN-specific differences are evident in the location and specificity of cortical stimulus responses.
a) Left: Response kernels for contralateral stimuli over all EMX (red), IT (green), and PT mice (blue), averaged
between 0 and 200 ms. Right: Stimulus-evoked activity in auditory (blue), parietal (red), and frontal cortex (yellow).
Dashed circles in the left stimulus maps show locations of respective cortical areas. **b)** Magnified view on parietal
regions of the stimulus maps in (a). PyNs differed in the location of sensory responses. Arrows show location of
parietal areas A, AM and the RS. **c)** Side-specific stimulus responses, computed as the difference between contra-
and ipsilateral stimulus kernels. Conventions as in (a). **d)** Magnified view on parietal regions of side-specific maps
in (c). IT neurons show clear, side-specific responses that were weaker in EMX and absent for PT neurons.

PyN types also differed in their response dynamics following the sensory stimulus. While some
cortical areas, such as auditory cortex, preferentially responded to contralateral stimuli, PT

neurons in parietal cortex were activated bilaterally in response to ipsi- or contralateral stimuli.
To assess such side-specificity, we computed the difference between the contra- and ipsilateral
stimulus kernels (Fig. 5c, colors indicate regions for which responses differed for contra- versus
ipsilateral stimulus kernels). In EMX mice, we found lateralized responses in auditory, frontal,
and to a lesser extent, parietal cortex (Fig. 5c,d). Lateralized IT activity was found in auditory
and parietal but not frontal cortex. In contrast, PT mice showed clear side-specific responses in
auditory and frontal but not in parietal cortex. Such differences in unilateral versus bilateral
responses in PT and IT neurons may also reflect different functional roles, with unilateral
responses encoding the spatial location of sensory information and bilateral responses signaling
its importance for guiding subsequent decisions.

Having identified PyN-type dependent activity for sensory stimuli, we then examined choice-
dependent activity and again observed clear differences across PyN types. In EMX mice, a
number of regions showed choice-related activity, particularly in the frontal cortex, while
sensory and parietal regions were only weakly modulated (Fig. 6a). We also found choice signals
in somatosensory areas of the whiskers and nose that slowly increased over the course of the
trial, even before the stimulus onset (Supp. Fig. 9a). These signals could indicate that mice used
their whiskers to probe the location of the spouts or performed subtle, choice-predictive facial
movements throughout the trial[40]. In contrast, choice-specific activity in frontal cortex strongly
increased after stimulus onset and remained elevated as the decision progressed from sampling
the stimulus to the subsequent delay period (Fig. 6a, yellow trace). We found equally prominent
choice signals in frontal cortex of PT mice while little choice-predictive activity was seen in IT
mice (Fig. 6b, Supp. Fig. 9b,c). In EMX and PT mice, positive signals for contralateral choices
were concentrated in the medial part of secondary motor cortex (M2) while parts of the primary
motor cortex (M1) were inhibited. This could indicate accumulation of sensory evidence and
motor preparation in M2, and inhibition in parts of M1 when early lick responses must be
witheld[41].

Although the choice kernels revealed differences between PyN types, choice-related activity only
accounted for a small amount of the total neural variance (Fig. 4d, Supp. Fig. 8). Since the
encoding model is designed to capture as much variance as possible, we hypothesized that this
approach might miss subtle choice signals that are specific but low in magnitude. The linear
model's ridge penalty also enforces choice-related variance to be distributed over all correlated
model variables, which might 'diffuse' weaker choice-related activity from the choice to other
model kernels.

To selectively isolate all choice-related activity, we therefore built a decoding model, using a
logistic regression choice classifier with L1 penalty (see Methods). In contrast to the encoding
model, this decoder approach isolates the cortical signals that are best suited to predict the
animal's choices, regardless of their magnitude. For all PyN types, the decoder predicted the
animal's choices with high accuracy, confirming that cortical activity reliably reflects trial-by-
trial choices (Supp. Fig. 10a). When analyzing the decoder weights, we found comparable
patterns to the encoding model's choice kernels but with much clearer separation of cortical
areas (compare top row 'Delay' in Fig. 6c to Fig. 6a, left). Here, positive decoder weights denote
areas that are most predictive for contralateral choices but, importantly, this does not suggest that
these areas are necessarily the most active. For all PyN types, we found significant choice signals
in multiple areas of the anterior cortex that evolved during decisions (Fig. 6c, Supp. Figs. 10,
11). In EMX and PT mice (top and bottom rows), large parts of M2 were again highly choice
predictive, including the anterior lateral motor cortex (ALM) and the medial motor cortex
(MM)[21]. M2 choice weights strongly increased immediately after stimulus onset and remained
elevated during the subsequent delay period (Fig. 6c,d). Cortical choice signals also persisted

after removing movement-related activity from the data, suggesting that they are not explained by choice-predictive animal movements but instead reflect the formation of sensory-driven decisions in frontal cortex (Supp. Fig. 12).

Figure 6. The temporal dynamics of choice-related activity differ across PyN types.
a) Left: Averaged choice kernels for EMX mice during the delay period. Positive weights indicate increased activity for contralateral choices, negative weights indicate choice-related reduction in activity. Right: Choice-related activity in auditory (blue), parietal (red), and frontal cortex (yellow). Traces are re-aligned to the initiation, stimulus, delay and response periods, indicated by gaps in weight traces. **b)** Zoomed-in map for frontal choice kernels of EMX, IT and PT mice during the delay period. **c)** Cortical maps of contralateral choice weights for different trial episodes. Several areas in anterior cortex showed clear choice signals. **d)** Baseline-corrected decoder weights in parietal (left) and frontal cortex (right) throughout the trial. Conventions as in (a). Dashed circles in the delay maps of (c) show the parietal and frontal locations that were used to compute the traces. **e)** Zoomed-in map for frontal decoder weights of EMX, IT and PT mice during the delay period. Dashed circle shows location of ALM.

Surprisingly, we also found a mild ipsilateral choice-preference for M2 in IT mice, despite strong
bilateral activation of frontal cortex during the delay period (Fig. 3c; Supp. Fig. 7). This
ipsilateral choice signal evolved more slowly during the stimulus and delay period (Fig. 6c,d)
and was more spatially restricted to ALM (Fig. 6e). In line with our previous results, no choice
signals were seen in parietal cortex of any PyN type (Fig. 6d, left), suggesting that parietal cortex
is mostly involved in sensory processing instead of choice formation or motor execution[42,43].

Compared to the encoding model, the decoding model recovered more finely structured choice
maps, especially in the frontal regions (compare Fig. 6e to 6b), revealing strong contralateral
choice signals in PT and ipsilateral choice signals in IT mice. One possible cause of this
unexpected inversion may be different choice-selectivity of specific IT-subtypes: intracortical
versus corticostriatal (CStr) projection neurons. Earlier work suggested an even distribution of
ipsi- and contralateral choice selectivity in frontal intracortical projection neurons[20,21], and we
thus hypothesized that our observed IT choice selectivity is due to the activity of CStr neurons.
To address this directly, we used an intersectional approach that exclusively labels large
populations of CStr neurons by performing multiple bilateral striatal injections of a retrograde
virus expressing Cre (CAV2-Cre) in GCaMP6s reporter mice (Fig. 7a), inducing widespread
expression in CStr neurons (Fig. 7b). As expected, GCaMP6 expression in CStr neurons was
largely confined to layer 5 with sparse expression in deeper layers.

We then used widefield calcium imaging to selectively measure activity from CStr neurons. As
with PT and IT mice, we observed robust fluorescence signals throughout the dorsal cortex
(Supp. Movie 5) and identified visual areas using retinotopic mapping (Supp. Fig. 13a). sNMF
showed that the dimensionality of CStr mice was intermediate between PT and IT activity, and
that the spatial components did not strongly overlap with other PyN types (Supp. Fig. 13b,c).
This clear difference between IT and CStr mice strongly suggests that imaging signals from IT
mice are not solely dominated by IT-positive CStr neurons but represent a mixture of
intracortical- and corticostriatal-projecting IT neurons with distinct activity patterns.

We then trained CStr mice in the auditory discrimination task. Trial-averaged dynamics partially
resembled IT mice, for example in frontal cortex, but also showed clear differences, such as a
lack of pre-stimulus suppression in sensory cortex (Fig. 7c,d; orange versus gray traces).
Differences between CStr and IT mice were also visible in the stimulus kernels from the
encoding model. In CStr mice, stimulus-related activity in parietal cortex was stronger than in
sensory and frontal cortex but the activated parietal regions were more medial than in IT mice
(Fig. 7e). Interestingly, these stimulus-driven parietal regions (Fig. 7f, top) closely resembled
cortical areas that form anatomical and functional connections to the dorsomedial striatum[44,45].
As with PT neurons (Fig. 5b,d), parietal CStr responses equally responded to contra- and
ipsilateral stimulation (Fig. 7f, bottom).

To determine whether CStr activity contributed to ipsilateral-preferring IT choice signals, we
repeated the choice decoder analysis. The decoder predicted animals' choices with equally high
accuracy as for PT and IT mice (Supp. Fig. 13d). We then extracted choice weights for each task
episode. Here, CStr activity was overall similar to that of IT mice, with an even stronger
ipsilateral-choice preference in frontal cortex that started after stimulus onset and lasted
throughout the delay and response periods (Fig. 7g, right). As in IT mice, this inversion from
contra- to ipsilateral choice preference was prominent in area ALM but did not extend to MM,
strongly suggesting that ipsilateral choice preference is driven by IT-CStr neurons.

Figure 7. An intersectional approach to measure cortex-wide activity of CStr neurons.
a) Schematic of the retrograde labeling approach. Multiple bilateral injections of retrograde CAV2-Cre virus in
reporter mice robustly induced expression of GCaMP6s in CStr neurons. **b)** Left: GCaMP6s-expression was robust
throughout the dorsal cortex. Scale bars are 100 μm. Right: Brain slices show robust cortex-wide expression. **c)**
Cortical dynamics of CStr neurons in the auditory discrimination task. Shown are trial-averages over all correct,
leftward trials in different trial episodes. No clear lateralization was observed in trial averages. **d)** Averaged activity
in auditory (left) and frontal cortex (right) for CStr (orange) and IT (gray) mice. Dashed lines show times of
initiation and response periods, gray areas the stimulus period. Traces show standard deviation units (SDU). **e)** Left:
Contralateral stimulus kernel, averaged over all CStr mice, between 0 and 200 ms after stimulus onset. Right:
Colored traces show changes in sensory (blue), parietal (red) and frontal cortex (yellow). Locations for each area are
indicated by dashed circles in the weight map. **f)** Top: Weight map from (e), zoomed in on parietal cortex. Bottom:
Difference of contra- versus ipsilateral stimulus kernels. **g)** Left: Cortical weight map from choice decoder during
the delay period, averaged over all CStr mice. Right: Baseline-corrected decoder weights in parietal (left) and
frontal cortex (right) for CStr (orange) and IT mice (gray). Traces are re-aligned to the initiation, stimulus, delay and
response periods, indicated by gaps in weight traces. **h)** Weight map from (g), zoomed in on frontal cortex. **i)**
Example field-of-view from two-photon imaging. Left panel: GCaMP6s-expression in all PyNs (green). Center
panel: retrogradely-labeled CStr neurons, expressing tdTomato (red). Right panels: zoomed-in maps for both
channels and a merged image. **j)** Left: Overview of significantly choice-tuned neurons in ALM (top) and MM
(bottom), imaged between 200-400 μm. Orange line shows CStr neurons, gray lines are unlabeled PyNs. AUC
values below 0.5 indicate stronger responses for ipsilateral choices. Right: Trial-averaged activity for all choice-
selective neurons, separated for ipsi- (red) versus contralateral choices (blue). CStr neurons in ALM (top right) show
higher activity for ipsilateral choices. **k)** Fraction of cells responding selectively for ipsi- versus contralateral
choices. Top: A higher fraction of CStr neurons in ALM are ipsi-selective (CStr$_{Ipsi}$: 20.4%, CStr$_{Contra}$: 15.5%,
binomial test: $p = 0.0018$) while unlabeled neurons are more contra-selective (Unlabeled$_{Ipsi}$: 14.3%, Unlabeled$_{Contra}$:
17.2%, binomial test: $p = 3.5 \times 10^{-10}$). Bottom: The majority of CStr as well as unlabeled neurons in MM are selective
for contralateral choices (CStr$_{Ipsi}$: 10.2%, CStr$_{Contra}$: 19.1%, binomial test: $p = 2.7 \times 10^{-8}$; Unlabeled$_{Ipsi}$: 9.3%,
Unlabeled$_{Contra}$: 19.6%, binomial test: $p < 1 \times 10^{-10}$).

To confirm these results on a single-cell level, we then used two-photon calcium imaging in
Camk2α-tTA;G6s2 mice and labeled CStr neurons through striatal injections of a retrograde
adeno-associated virus (AAVrg-tdTomato) (Fig. 7i). This allowed us to compare the activity of
all PyNs in frontal cortex with tdTomato-labeled CStr neurons in the same animal. Comparing
the choice-tuning of CStr and unlabeled PyNs revealed a specific difference in ipsi- versus
contralateral choice preference in ALM (Fig. 7j). Here, significantly more choice-selective CStr
neurons preferred ipsilateral choices, whereas unlabeled PyNs were mildly contra-selective (Fig.
7k). In agreement with our widefield results, these differences were only seen in ALM but not
MM. Interestingly, this ipsilateral choice preference was also restricted to superficial IT-CStr
neurons at a cortical depth between 200-400 µm. Infragranular CStr neurons (400-600 µm),
which are also often PT cells[17], showed strong contralateral choice tuning that was similar to
unlabeled cells (Supp. Fig. 14a). Lastly, we also tested if the neuropil exhibits distinct choice
signals that may have masked somatic activity in our PyN-specific widefield measures (Supp.
Fig. 14b). In all regions and depths, neuropil choice tuning was largely similar to somatic activity
of unlabeled neurons, confirming that widefield measures indeed represented local somatic
activity. Here, IT-specific widefield signals matched the mixed choice-selectivity of superficial
layers, while PT-specific imaging was well-aligned with the clear contralateral choice tuning in
deeper cortical layers.

The observed differences between PyN types suggest that each type may drive distinct aspects of
decision-making. To causally test the functional role of different PyN types, we performed PyN-
specific optogenetic inactivation in different cortical areas. We induced Cre-dependent
expression of the inhibitory opsin stGtACR2[46] by injecting an AAV in the parietal and frontal
cortex of Fezf2-, PlexinD1-, or EMX-Cre mice (Fig. 8a, b, Supp. Table 2). To express stGtACR2
in CStr neurons, we used a combined viral receptor complementation and intersectional approach
to maximize the efficiency of retrograde expression[47]. First, we simultaneously expressed the
coxsackievirus and adenovirus receptor (CAR) and transduced Cre-dependent stGtACR2 in
cortical neurons. Two weeks later, we performed retrograde-mediated stGtACR2 expression
through striatal CAV2-Cre injections. CAR mediates efficient axonal uptake of CAV2 and
prevents potential viral tropism in subsets of CStr neurons. The coordinates of auditory, parietal,
and frontal injections and subsequent inactivation were determined from our stimulus and choice
analyses (Fig. 5a, 6a; dashed circles). To test whether optogenetic effects are area-specific, we
also targeted the primary visual cortex (V1) in a subset of EMX mice as controls. To simplify the
behavioral task during the optogenetic inactivation experiments, we only presented unilateral
stimuli in each trial and kept the stimulus duration and delay periods constant (1 and 0.5 seconds,
respectively).

In agreement with our imaging results, bilateral optogenetic inactivation of auditory, parietal or
frontal cortex of EMX mice strongly impaired auditory decision accuracy (Fig. 8c). Performance
was unaffected by V1 inactivation, confirming that these areas are selectively required for
auditory decisions. We then illuminated each area for 0.5 seconds during four different task
episodes: Early and late stimulus (the first and last 0.5 seconds of the stimulus period), delay and
response (Fig. 8d). In agreement with our observation that auditory and parietal cortex reflects
stimulus driven activity (Fig. 5a, red and blue traces), inactivation of either area strongly reduced
animal's performance, particularly during the stimulus period (Fig. 8e, f). Behavioral
impairments (computed as the normalized difference between performance in non-optogenetic
trials and chance) were weaker during the subsequent trial periods, indicating that these areas are
most important for early processing of auditory stimuli. As expected, impairments were strongest
in EMX mice where all PyNs were affected (gray trace).

In agreement with earlier work[48], inhibiting CStr neurons in A1 also affected auditory decisions
(Fig. 8e, orange trace). However, the effects were more transient and weaker compared to
inactivation of EMX neurons, suggesting that CStr neurons were not exclusively required for
accurate task performance. This was even more evident in parietal cortex where impairments
were surprisingly weak when inhibiting either IT or CStr neurons (Fig. 8f, green/orange traces)
while inactivating PT neurons resulted in a more robust impairment (blue trace). This suggests
that the subcortical PT projection pathways in the parietal cortex have a larger causal impact on
sensory processing than intracortical IT or CStr projections, pointing to a role for PT neurons
that extends beyond preparation and execution of movements.

Figure 8. Temporally restricted, PyN-specific inactivation of parietal and frontal cortex disrupts decisions.
516
a) Left: Schematic of injection scheme to induce stGtACR2 expression in EMX, IT, or PT neurons. Injections were
performed in auditory, parietal and frontal cortex of transgenic mice to target different PyN types. V1 injections
were performed in a subset of control EMX mice. Right: Intersectional viral approach for targeting CStr neurons.
AAV-DJ-hSYN-DIO-hCAR{off} induced expression of CAR in cortical neurons to promote uptake of CAV2-Cre
virus from the striatum. A second cortical injection induced Cre-dependent stGtACR2 expression in all CStr
neurons. **b)** Layer-specific expression of stGtACR2-FusionRed in EMX, PT, IT and CStr neurons. **c)** Behavioral
performance (% correct) of EMX mice during inactivation of V1, auditory, parietal or frontal cortex. Inhibition of
auditory, parietal and frontal cortex, but not V1, significantly reduced task performance. Circles indicate individual
mice, error bars show the s.e.m. **d)** Schematic of optogenetic inactivation paradigm. 0.5-s long optogenetic
inhibition was performed during the first or last part of the stimulus period, the subsequent delay or the response
period. Light power ramped down after 0.3 seconds. **e)** Behavioral impairment (% change from control
performance) with inhibition of EMX or CStr neurons in auditory cortex. Circles show mean impairments, error bars
95% confidence intervals. **f)** Behavioral impairment with inhibition of different PyN types in parietal cortex.
Conventions as in e). **g)** Behavioral impairment with inhibition of different PyN types in frontal cortex. Conventions
as in f). Stars in all panels indicate significance (bonferroni-corrected p < 0.01, binomial test).

Frontal inactivation resulted in a particularly strong impairment, especially for nonspecific PyN
inactivation (EMX mice, Fig. 8g, gray trace), consistent with a role for frontal cortex in sensory
integration and working memory[20,43,49]. IT and CStr inactivation resulted in comparable
behavioral impairments during the stimulus and delay periods (Fig. 8g, green and orange traces).
Choice impairments in IT mice are therefore not solely due to the disruption of intracortical
processing[20] but also involve alterations of CStr neurons. Inactivating PT neurons equally
impaired animal performance during the stimulus and delay period but showed stronger effects
during the final response period. Impairments in the response period were similar for EMX and

PT mice, suggesting that PT neurons are particularly involved in licking responses. These results
show that multiple PyN types in frontal cortex are involved in the formation and maintenance of
choices, despite clear differences in their respective choice tuning. Lastly, we also analyzed
licking patterns to test if optogenetic inhibition broadly disrupted animal movements. Frontal
inactivation in the delay period had a mild effect on response latency but did not affect response
probability or licking patterns, arguing against a strong motor impairment (Supp. Fig. 15). This
shows that PyN-specific inhibition selectively reduced the animals' response accuracy rather
than broadly disrupting their ability for movement initiation and execution.

Discussion

We measured and manipulated PyN subtypes to determine whether they play distinct roles in
decision-making. Cortex-wide activity patterns were PyN subtype-specific, each reflecting
distinct neural dynamics at multiple spatial scales. Functional specificity across PyN subtypes
was also evident during auditory decision-making: activity of each PyN subtype exhibited unique
cortical localization and spatial specificity associated with stimulus and choice. These response
patterns were not seen when imaging from PyNs nonspecifically. Optogenetic inactivation
confirmed that PyN types in parietal and frontal cortex have distinct functional roles,
highlighting the importance of subcortical projection pathways for sensory processing and choice
formation. Taken together, our results suggest that different PyN types form parallel subnetworks
throughout the cortex, are functionally-distinct, and perform separate roles during auditory
decision-making.

Dimensionality reduction of cortical dynamics[33,50,51] revealed that nearly all spatial components
were also PyN-subtype-specific. Differences between PyNs are therefore not only present within
parcellated cortical regions[18–20] but also evident in large-scale activity patterns. This has
important implications for studies of cortex-wide neural dynamics, which are often based on
indirect measures of neural activity[52,53] or pooled activity from all PyNs[22,25,54,55]. Earlier work
revealed intricate circuit motifs and functional modules that span the entire cortex[50,54–57] and
largely follow intracortical connectivity patterns[58,59]. Our results point to the existence of
additional circuit motifs when specific PyN population are isolated, especially subcortical
projection types, such as PT or CStr neurons. Furthermore, most PyN-specific LocaNMF
components consisted of spatially precise subregions that were smaller than Allen CCF areas.
Detailed analysis of PyN-specific activity might therefore reveal more detailed cortical structures
than could be observed with nonspecific measures. For future studies, a particularly promising
approach to achieve this goal would be to combine large-scale measures of multiple PyN types
with multi-color widefield imaging[24,60] and directly observe interactions between PyN-specific
cortical dynamics within the same animal.

Sensory responses were pronounced in auditory, parietal and frontal cortices[42,43,61,62] and
revealed unique response patterns for each PyN type. This functional segregation is in line with
recent results from primary sensory areas, such as somatosensory[18,63] and visual cortex[19], and
strongly suggests that different PyN types play separate roles during sensory processing.
Moreover, the clear differences in magnitude, localization, and lateralization of sensory
responses in parietal and frontal cortex demonstrate that the functional specialization of different
PyN subtypes is a general feature of cortical circuit function.

We selectively inactivated different PyN types across cortical areas to directly test their
involvement in the behavioral task. In agreement with earlier work, inhibiting CStr neurons in
auditory cortex impaired task performance[48], suggesting that projections from cortex to striatum
are important for accurate auditory decisions. Moreover, inactivation of parietal cortex also

caused strong behavioral impairments, especially during sensory stimulation. Surprisingly,
inactivation of CStr or IT neurons caused only weak behavioral effects. This shows that the
importance of CStr projections does not generalize from auditory to parietal cortex and also
argues against models in which task-relevant sensory information is directly transmitted from
parietal to frontal cortex to promote decision formation[42,62,64]. Instead, inactivation of parietal PT
neurons during the stimulus presentation strongly disrupted eventual decisions, highlighting the
importance of subcortical projection pathways for auditory performance. A likely subcortical
target is the SC, which receives inputs from PT neurons[18,26] and has been implicated with
somatosensory[18] and visual[65,66] perception, and decision-making[39,67,68].

The strong behavioral effects with inactivation of parietal cortex are at odds with earlier studies
in rats that showed very little or no impact of parietal inactivation in auditory task
performance[69,70]. Conversely, other work in head-fixed mice had found robust behavioral effects
in visual[43,71–73] and auditory[62] task performance. While a potential reason for these contradictory
results could be species differences between rats and mice, the precise location of parietal
inactivation most likely plays an important role. Different sensory modalities are processed along
a mediolateral gradient in parietal cortex, emphasizing the need to precisely target specific
parietal areas to obtain a modality-specific behavioral effect[62]. We also found that parietal
responses of PT and CStr neurons were located more medially than IT neurons, further
emphasizing the need for precise localization. Another important factor is the specific behavioral
task. Our task design required accumulation of sensory evidence and working memory which
engage a wider range of cortical regions[72] and might therefore explain the involvement of
parietal cortex for accurate auditory decisions.

The accumulation and memory requirements of the decisions we studied might also explain why
we found clear cortical choice signals, especially in frontal cortex, whereas recent cortex-wide
studies reported little choice-selectivity[39,74]. The lack of side-specific choice tuning in IT
population data is consistent with earlier work, showing that intracortical projection neurons in
frontal cortex include an equal amount of contra- and ipsilateral choice-preferring cells[20,21]. In
contrast to other PyNs, CStr neurons were more selective for ipsilateral choices. Using two-
photon imaging, we confirmed that this ipsilateral choice preference was also present in
individual CStr neurons but not in unlabeled IT neurons. This confirms that our PyN-specific
widefield results indeed reflect selective somatic activity and not just superficial neuropil signals,
in line with earlier findings that widefield signals are strongly correlated to spiking of cortical
neurons[44]. Ipsilateral choice signals in CStr neurons were restricted to superficial layers of area
ALM, which is mostly implicated in movement generation[13,49]. A recent study showed that CStr
projections from the anterior cingulate cortex can inhibit striatal activity and motor behavior[75]. A
potential reason for the reduced activity of CStr neurons during the response period could
therefore be to disinhibit striatal circuits and release a targeted licking response.

In agreement with our imaging results, frontal inactivation strongly impaired animal behavior
during the stimulus and delay period, suggesting an important role for the translation of sensory
inputs into behavior[21,74,76–78]. Impairments were largely similar for all frontal PyN types, which
appear to be equally required for choice formation and retention. Frontal PyNs may thus be more
reliant on each other to maintain accurate function than in sensory areas[18,19,48]. As the only
exception, PT neurons were more important during the response period, consistent with a
specific role of brainstem-projecting PT neurons for motor execution[13].

Our work offers a new perspective on cortex-wide dynamics by viewing them through the lens of
different PyN types and strongly supports the view that cortical circuits perform parallel
computations, even within the same cortical layer[13,18,19,79]. Future work to reveal the circuit logic

of cortex-wide activity patterns in different behavioral contexts should therefore include PyN
type information to resolve the functional heterogeneity that is often encountered when studying
cortical decision circuits. A powerful tool to achieve this goal are novel mouse lines, such as our
temporally inducible knock-in lines that allowed us to reliably target PT and IT neurons with
high specificity. These new mouse lines also overcome several known problems of earlier
transgenic approaches, unstable expression patterns across cortical areas or mixtures of cell types
due to interactions with surrounding genetic elements[26,80]. Future studies could explore PyN type
function at even higher granularity by using more specific mouse lines, such as the Cux1-line to
selectively isolates intracortical projection neurons[26]. Moreover, combining genetic mouse lines
with retrograde labeling will enable the targeting of specific PyN subtypes, such as projection-
specific PT neurons[12,13], that might serve a large array of functions from sensory processing, to
working memory and motor function[13,18].

Acknowledgements

We thank Z. Josh Huang for providing the Fezf2-2A-CreER and PlexinD1-2A-CreER mouse
line and for countless insightful conversations; Shreya Saxena and Joao Couto for technical
advice and scientific discussions; Anup Khanal for assistance with animal perfusions; Sandra
Brill, Maria Perez and Alice Despatin for animal training; Adam Kepecs for providing the viral
vector AAV-DJ-hSYN-DIO-hCAR{off}; Katherine S. Matho for invaluable help with Fezf2 and
Plexin cell density quantification. This work was supported in part by NIH grants R01EY022979
and W911NF-16-1-0368 as part of the collaboration between the US DOD, the UK MOD and
the UK Engineering and Physical Research Council under the Multidisciplinary University
Research Initiative to A.K.C. S.M. was funded by the Deutsche Forschungsgemeinschaft (DFG,
German Research Foundation) - 368482240/GRK2416 and the Helmholtz association (VH-NG-
1611). X.R.S. was supported by the NRSA F32 MH120888.

Methods

Mouse lines

All surgical and behavioral procedures conformed to the guidelines established by the National
Institutes of Health and were approved by the Institutional Animal Care and Use Committee of
Cold Spring Harbor Laboratory. Experiments were conducted with male mice between the ages of
8 to 25 weeks. No statistical methods were used to pre-determine sample sizes but our sample sizes
are similar to those reported in previous publications[22,25]. All mouse strains were acquired from
the Jackson Laboratory, the Allen Brain Institute, or generated at Cold Spring Harbor Laboratory.
Transgenic strains crossed to generate double- and triple-transgenic mice used for imaging: Emx-
Cre (JAX 005628), LSL-tTA (JAX 008600), Ai93D (JAX 024103), Ai162 (JAX 031562), TRE-
GCaMP6s (G6s2, JAX 024742) and H2B-eGFP (JAX 006069). EMX mice, used for calcium
imaging, were bred as Ai93D;Emx-Cre;LSL-tTA. To avoid potential aberrant cortical activity
patterns, EMX mice were on a doxycycline-containing diet (DOX), preventing GCaMP6s
expression until they were 6 weeks or older[22,25].

For widefield imaging of PT and IT neurons, inducible knock-in drivers Fezf2-2A-CreER and
PlexinD1-2A-CreER (Supp. Table 2), respectively, were crossed with Ai162 reporter mice to drive
cortex-wide GCaMP6s expression. Cre activity was induced through two doses of intra-peritoneal
injections of tamoxifen (200 mg/kg; 20 mg/ml solution dissolved in corn oil) at P28 and P32.
Histologic characterization revealed pyramidal neuron expression patterns consistent with prior
reports[26]. For widefield imaging of corticostriatal neurons, we first crossed Ai162 with G6s2 to

create a new double-transgenic reporter strain Ai162;G6s2 with two hemizygous copies of
GCaMP6s under tetO control. Since LSL-tTA is incorporated in-tandem to the reporter gene in
the Ai162 strain[29], this hybrid reporter line permits Cre-dependent expression of GCaMP6s at
higher levels than Ai162 hemizygotes while avoiding potential leaky reporter gene expression that
is sometimes found in Ai162 homozygotes. Next, to achieve widespread GCaMP6s expression in
corticostriatal neurons, we performed striatal injections of retrograde virus (CAV2-Cre) in the
hybrid Ai162;G6s2 reporter line (see section: viral injections).

For two-photon imaging experiments, GCaMP6s expression in pyramidal neurons were generated
using the hybrid strain Camk2α-tTA;G6s2.

General surgical procedures

All surgeries were performed under 1-2% isoflurane in oxygen anesthesia. After induction of
anesthesia, 1.2 mg/kg of meloxicam was injected subcutaneously and sterile lidocaine ointment
was applied topically to the skin incision site. After making a midline cranial incision, the skin
was retracted laterally and fixed in position with tissue adhesive (Vetbond, 3M). We then built an
outer wall using dental cement (C&B Metabond, Parkell; Ortho-Jet, Lang Dental) along the lateral
edge of the dorsal cranium (frontal and parietal bones) to maximize the area of exposed skull. A
custom titanium skull post was then attached to the dental cement. For skull clearing, debris and
periosteum were thoroughly cleaned from the skull followed by the application of a thin layer of
cyanoacrylate (Zap-A-Gap CA+, Pacer technology).

To prepare mice for two-photon imaging, a circular craniotomy (ø = 3 mm) centered over the right
frontal cortex (1.75 mm lateral to midline and 1.75 mm rostral to bregma), was made using a
biopsy punch. A circular coverslip (ø = 3 mm) was then lowered to the surface of the brain and
Vetbond and Metabond were used to seal the window to the skull. Lastly, a titanium skull post
was implanted as described above.

Viral injections

After induction with isoflurane anesthesia, animals were placed in a stereotaxic frame (David Kopf
Instruments). The skull was leveled along the rostral-caudal and medio-lateral axis, allowing for
accurate and reproducible targeting. All injections were made using a programmable nanoliter
injector (Nanoject III, Drummond Scientific, PA). For *widefield imaging* of CStr mice, widespread
corticostriatal GCaMP6s expression was generated in Ai162;G6s2 reporter mice by performing
bilateral stereotaxic injections of CAV-2-Cre (at 3-4 weeks of age) into the dorsal striatum at three
targets per hemisphere spanning the rostro-caudal axis. The target coordinates (relative to bregma
and dura, in mm) are (1) RC +0.75, ML ±1.8, DV 3.0; (2) RC 0, ML ±2.2, DV 3.1; (3) RC -0.75,
ML ±2.9, DV 3.1. For each striatal target, a burr hole was created using a small dental burr
followed by injection of 1.8×109 purified particles (pp) of CAV-2-Cre using pipettes with long
taper tips pulled from borosilicate capillaries (3.5" Drummond # 3-000-203-G/X, Drummond
Scientific, PA). For *two-photon imaging* experiments, CStr neurons were labeled through striatal
injections of AAV-2-retro-CAG-tdTomato (using the same approach and coordinates as described
above) in Camk2α-tTA;G6s2 mice.

For cell type-specific *optogenetic silencing* experiments, we performed bilateral injections in
frontal, parietal, and auditory cortex (coordinates relative to bregma, in mm: frontal cortex: RC
+2.5, ML ±1.5; parietal cortices: RC -1.7, ML ±2.5; auditory cortex: RC -2.5, ML ±4.6) to
induce expression of Cre-dependent stGtACR2 (AAV1-hSyn-SIO-stGtACR2-FusionRed, Upenn
Vector Core, Supp. Table 2). Cortical injections were performed in P42 to P56 Fezf2-2A-CreER

(PT), PlexinD1-2A-CreER (IT), and EMX-Cre (nonspecific PyNs) reporter mice. In mice with ligand-dependent Cre recombinase activity, intraperitoneal tamoxifen was administered one week after viral injections. Injections were made at two depths (300 and 600 μm) per cortical target. In two EMX-Cre mice, bilateral injections were performed in the frontal and visual cortex (RC -4, ML ±2.5). To express stGtACR2 in CStr neurons, viral injections were performed in C57BL/6J mice in two stages. In stage one, we utilized a viral receptor complementation strategy[47] by injecting both AAV-DJ-hSYN-DIO-hCAR{off} and AAV1-SIO-hSyn1-stGtACR2-FusionRed in different cortical locations (targets as described above) in P21-P28 mice. In stage two, we performed bilateral striatal CAV-2-Cre injections 6 weeks after the initial cortical injections (see above). hCAR is expressed in all transfected neurons in a Cre-OFF manner, meaning that Cre expression stops expression of hCAR while inducing expression of stGtACR2.

Optical fiber implantation

For optogenetic silencing, we expressed the soma-targeted anion-conducting channelrhodopsin stGtACR2 in target neuronal populations[46]. Optical fibers (NA = 0.36, ø = 0.4mm, FT400UMT, Thorlabs) were glued into metal or ceramic ferrules (ø = 1.25 mm, Thorlabs) and secured above the cortex following viral injections at each target. Ferrule-enclosed optical fiber implantations immediately followed cortical AAV injections in Fezf2, PlexinD1, and Emx mice and striatal CAV-2-Cre injections in CStr mice. One polished end of the optical fiber was positioned extradural to the site of cortical injections and interfaced with thinned skull using cyanoacrylate. Next, the fiber was fixed to the skull using light-cured glass ionomer (Vitrebond, 3M). Additional layers of dental cement and dental acrylic (Lang Dental Jet Repair Acrylic; Part#1223MEH) were applied around the fiber implant and the skull to reinforce for durability and long-term stability. After all layers were cured, a final outer coating of cyanoacrylate and nail polish were applied.

Behavioral training

The behavioral setup was controlled with a microcontroller-based (Arduino Due) finite state machine (Bpod r0.5, Sanworks) using custom Matlab code (2015b, Mathworks) running on a Linux PC. Servo motors (Turnigy TGY-306G-HV) and touch sensors were controlled by microcontrollers (Teensy 3.2, PJRC) running custom code. Fifty-four mice were trained on a delayed, spatial discrimination task. Mice initiated trials by placing their forepaws on at least one of the two handles, which were mounted on servo motors that rotated out of reach during the inter-trial period. Upon trial initiation, animals placed their forepaws on the handles and, after a variable duration of 0.25-0.75 seconds of continuous contact, the auditory stimulus was presented. Auditory stimuli consisted of a sequence of Poisson-distributed, 3-ms long auditory click sounds[36], presented from either a left and/or right speaker for a variable duration between 1 and 1.5 seconds. The stimulus period was followed by a variable delay of up to 1 second, then the servo motors moved two lick spouts into close proximity of the animal's mouth. If the animal licked twice on the side where more clicks were presented, a drop of water reward was dispensed. The amount of water rewarded per trial (typically 1.5 to 3 μl) was constant within a single session but was sometimes adjusted daily based on the animal's body weight. After one spout has been licked twice, the contralateral spout moved out of reach to force the animal to commit to its decision.

All trained mice were housed in groups of two or more under reverse light cycle (12- hour dark and 12-hour light) and trained during their active dark cycle. Animals were trained over the course of approximately 30-60 days. After 2-3 days of restricted water access, animals began habituation to head fixation and received water from spouts in the behavior chamber. During these sessions, unilateral auditory stimuli were presented followed by a droplet of water dispensed freely from the

ipsilateral water spout. After several habituation sessions, animals were then required to touch the handles to trigger stimulus presentation. Once mice could reliably reach for the handles, the required touch duration was progressively increased up to 0.75 seconds. During the next stage of training, self-performed trials, where both spouts moved within reach of the animal following stimulus presentation, were progressively introduced. An animal was considered trained when its detection performance across two or more sessions was above 80%.

Behavioral monitoring

Data was collected from multiple sensors in the behavioral setup to measure different aspects of animal movement. Touch sensors using a grounding circuit on handles and lick spouts detected contact with the animal's forepaws and tongue, respectively. A piezo sensor (1740, Adafruit LLC) below the animal's trunk was used for monitoring body and hindlimb movements. Two webcams (C920 and B920, Logitech) were used for video recording of animal movements. Cameras were positioned to capture the animal's face (side view) and the ventral surface of the body (ventral view).

Widefield imaging

Widefield imaging was done as reported previously[23,32,81] using an inverted tandem-lens macroscope in combination with an sCMOS camera (Edge 5.5, PCO) running at 30 fps. The top lens had a focal length of 105 mm (DC-Nikkor, Nikon) and the bottom lens 85 mm (85M-S, Rokinon). The field of view was 12.5 x 10.5 mm^2 and the imaging resolution was 640 x 540 pixels after 4x spatial binning, resulting in a spatial resolution of ~20 μm per pixel. To capture GCaMP fluorescence, a 525 nm band-pass filter (#86-963, Edmund optics) was placed in front of the camera. Using excitation light at two different wavelengths, we isolated Ca^{2+}-dependent fluorescence and corrected for intrinsic signals (e.g., hemodynamic responses)[22,25]. Excitation light was projected on the cortical surface using a 495 nm long-pass dichroic mirror (T495lpxr, Chroma) placed between the two macro lenses. The excitation light was generated by a collimated blue LED (470 nm, M470L3, Thorlabs) and a collimated violet LED (405 nm, M405L3, Thorlabs) that were coupled into the same excitation path using a dichroic mirror (#87-063, Edmund optics). We alternated illumination between the two LEDs from frame to frame, resulting in one set of frames with blue and the other with violet excitation at 15 fps each. Excitation of GCaMP at 405 nm results in non-calcium dependent fluorescence[82], allowing us to isolate the true calcium-dependent signal by rescaling and subtracting frames with violet illumination from the preceding frames with blue illumination. All subsequent analysis was based on this differential signal. The imaging data was then rigidly aligned to the Allen common coordinate framework (CCF), using four anatomical landmarks: the left, center and right points where anterior cortex meets the olfactory bulbs, and the medial point at the base of retrosplenial cortex. Retinotopic visual mapping experiments[31,83] confirmed accurate CCF alignment and showed high correspondence between functionally identified visual areas and the CCF across PyN types (Fig. 1c).

Two-photon imaging

We used a two-photon resonant scanning microscope (Moveable Objective Microscope, Sutter Instruments) for session-long, continuous image acquisition at 30.9 Hz. A 16X, 0.8 NA Nikon objective lens was used for single-plane imaging with a 512x512 pixels (575 μm x 575 μm) fields of view. Mode-locked illumination at 930 nm was delivered using a Ti:Sapphire laser (Ultra II, Coherent). Imaging frames were aligned with behavior control events synchronized acquisition of analog galvanometric and Bpod output signals. Depth of focal planes was 200-400 μm below the dura. Emission was collected using band-pass red (670/50 nm) and green (525/50 nm) filters

(Chroma Technologies). MScan software (Sutter Instruments) was used for image acquisition.
Recordings were performed in ALM (2.5 mm rostra and 1.5 mm lateral to bregma) or MM (1.5
mm anterior and 1 mm lateral to bregma) in randomized order across mice. Across imaging
session, we selected planes that differed from those of prior sessions in order to maximize the
number of unique neurons by.

Raw images were processed using the Suite2P package[84] which performed motion correction,
model-based ROI detection, correction for neuropil contamination, and spike deconvolution.
Somatic and non-somatic (neuropil) ROI identification was performed through a combination of
pre-trained classifier and manual curation. Somata with tdTomato expression were identified in a
two-step process. First, potential green channel bleed-through was subtracted from the red channel
using non-rigid regression with individual channels being divided into smaller blocks. Next, all
sessions were then manually inspected to identify a conservative red fluorescence threshold, which
was subsequently applied to all sessions. Analyses of neural activity were based on deconvolved
values ("inferred spiking activity"). Since the deconvolved values do not represent absolute firing
rates, we performed z-score normalization for each neuron before computing trial-averages across
cells. The total number of recorded neurons per session was 396 ± 105 (mean $\pm$ standard deviation).

Optogenetic inactivation

Photostimulation was performed using a 470 nm high-power LED (M470F3, Thorlabs) with a
power density of 25 mW/mm^2. Stimuli consisted of a square wave stimulus that ramped down in
power for 200 milliseconds, to avoid an excitatory post-illumination rebound due to sudden release
of inhibition[85]. To prevent animals' visual detection of photostimulation, either through external
leakage from light-insulated mating sleeves or transmission to the retina across the brain, an
external LED with matching wavelength placed at the center of the animal's visual field was
flashed throughout the duration of every trial. Photoinhibition was performed in 20% of total trials
and randomly interleaved between light-off trials. During each session, only bilateral frontal,
parietal or visual cortex inhibition was performed. Once an animal was habituated and able to
complete detection behavior trials with > 90% accuracy, optogenetic inactivation trials were
introduced. During these initial sessions, optogenetic inhibition was performed from the beginning
of the stimulus epoch until the end of the delay epoch. Additionally, we performed 0.5-second
inhibition during four pre-defined epochs of the detection behavior trials: (1) first half of the
stimulus, (2) second half of the stimulus, (3) delay, (4) response.

Immunohistology, microscopy and image analysis

For a given animal, after all experiments were concluded, we performed transcardial perfusion
with PBS followed by fixation with 4% PFA in 0.1 M PB. Brains were post-fixed in 4% PFA for
an additional 12-18 hours at 4°. Prior to sectioning, brains were rinsed three times in PBS and
embedded in 4% agarose-PBS. Slices 50 μm in thickness were made using a vibrating microtome
(Leica, VT100S). Sections were then suspended in blocking solution (10% Normal Goat Serum
and 0.1% Triton-X100 in 1X PBS) for 1 hour followed by overnight incubation at 4°C with the
primary antibody. Next, sections were washed with PBS, incubated for 1 h at room temperature
with the secondary antibody at 1:500 dilution. For visualization of GCaMP6s, we used primary
goat polyclonal anti-GFP antibody (Abcam) and secondary donkey anti-goat Alexa Fluor 488
(Abcam). Sections were then dry-mounted on slides using Vectashield (Vector Labs, H1000) prior
to imaging. No immunostaining was performed for the visualization of FusionRed or tdTomato.
Imaging was performed using upright fluorescence macroscope and microscope (Olympus BX61).
Images were acquired using Ocular Scientific Image Acquisition Software (Teledyne Imaging)
and visualization and analysis were performed using ImageJ/FIJI software packages.

Quantification cortex-wide gene expression

Quantification of cell counts across the dorsal cortex was performed using publicly available serial two-photon tomography (STPT) datasets. (http://www.brainimagelibrary.org/)[26]. Cre expression patterns for IT and PT neurons were characterized with data from eight mice, expressing either Cre-dependent GFP (PlexinD1-2A-CreER;Snap25-LSL-2A-EGFP) or tdTomato (Fezf2-2A-CreER;Ai14), respectively. Cell counting was then performed via automated soma detection, using a trained convolutional neural network[86]. STPT datasets were then registered to the Allen CCF v3 using the elastix toolbox[87]. To obtain the density of Cre-expressing neurons for individual cortical areas, we used the area outlines from the Allen CCF and computed the average sum of detected IT or PT neuron in each area, normalized by its surface area.

Preprocessing of neural data

We first performed motion correction on each imaging frame, using a rigid-body image registration method implemented in the frequency domain[88] that aligned each frame to the median over all frames in the first trial. To reduce the computational cost of subsequent analyses, we then computed the 200 highest-variance components using singular value decomposition (SVD). These components accounted for at least 95% of the total variance in each recording, whereas computing 500 components accounted for little additional variance (~0.15%). SVD reduces the raw imaging data Y to a matrix of 'spatial components' U (of size pixels by components), 'temporal components' V^T (of size components by frames) and singular values S (of size components by components) to scale temporal components to the original data. The resulting decomposition has the form $Y = USV^T$. All subsequent analysis in the time domain (such as the encoder and decoder models described below) were then performed on the product SV^T and the respective results were later multiplied with U, to recover results for the original pixel space. To avoid slow drift in the imaging data, SVT was high-pass filtered above 0.1 Hz using a zero-phase, second-order Butterworth filter.

To compute trial averages and perform choice decoder analysis (see below), imaging data in individual trials were aligned to the four trial periods, each marked by a specific event. This was required because the duration of different trial events was randomized to reduce temporal correlations, e.g. between trial initiation, the stimulus presentation and subsequent lick responses. The first period (Initiate) was aligned to the time when animal initiated a trial by touching the handles, the second (Stimulus) was aligned to the stimulus onset, the third (Delay) to the end of the stimulus sequence, and the fourth (Response) to the time when spouts were moved in to allow a lick response. After alignment, the total trial duration was 2 seconds and durations of respective trial episodes were 0.5 (Initiate), 1 (Stimulus), 0.2 (Delay), and 0.3 seconds (Response).

Spatial clustering and classification

To obtain more interpretable spatial components and assess the dimensionality of cortical activity in different PyN types, we used semi-nonnegative matrix factorization (sNMF). As with SVD, sNMF also creates spatial and temporal components for each session and for each mouse but enforces spatial components to be strictly positive. The reason why temporal components were not also enforced to be non-negative is that our hemodynamic correction resulted in temporal dynamics that can be either positive or negative, relative to baseline. We used the LocaNMF toolbox by Saxena et al[33] (https://github.com/ikinsella/locaNMF) to transform the spatial and temporal components U and SV^T into two corresponding matrices A and C, where A is a matrix of non-negative spatial components (of size pixels by components) and C the corresponding temporal components (of size components by frames). In addition to regular sNMF, the LocaNMF

toolbox can be initialized with spatial constraints that are based on the Allen CCF. To obtain
spatially restricted localized LocaNMF components, we constructed a map of larger seed regions
by merging several smaller areas in the Allen CCF together (Fig. 2e). This region map is then used
to enforce that each component in A is sparse outside the boundary of a given region. The amount
of possible overlap between regions is specified by a localization threshold which specifies the
percentage of a given component that is constrained to be inside a single region's boundary. To
obtain dense spatial components that were mostly driven by the local correlations between pixels
and could also lie at the border between seed regions, we used a localization threshold of 50%. To
obtain unconstrained sNMF spatial components, we also used the LocaNMF toolbox but only
provided a single region that spanned the entire cortex. This resulted in cortex-wide components,
similar to vanilla sNMF, while ensuring that all other analysis steps were done identically for
sNMF and LocaNMF components. In both cases, we determined how many components in A and
C were needed to explain 99% of the variance of Y (with Y=AC) after the initial SVD.

To compare spatial sNMF and LocaNMF components from different PyN types, we embedded
them in a 2-dimensional space, using Uniform Manifold Approximation and Projection (UMAP)
analysis (Fig. 2c,g). UMAP analysis was performed with the UMAP toolbox by McInnes et al.[35]
(https://github.com/lmcinnes/umap). For each recording, the first 20 spatial components in A
(either from sNMF or region-constrained LocaNMF) were downsampled by a factor of 2,
smoothed with a 2-D gaussian filter (5 x 5 pixels, 2 pixel standard deviation) and peak-normalized.
Components from all recordings and animals were then combined into a larger matrix (of size
pixels by components) and we used UMAP to project the first (pixel) dimension into two,
maximally separating non-linear dimensions. Each point in the two dimensional space (Fig. 2c,g)
therefore reflects a single component from one animal in a given imaging session.

The same UMAP approach was used for temporal sNMF and LocaNMF components. Before the
UMAP projection, we first computed the trial-averaged and z-scored activity of each component
to achieve temporal dynamics that are comparable across sessions and individual mice.

To identify PyN types based on individual spatial components (Fig. 2d, h), we performed a separate
UMAP projection for each mouse. Each of these projections excluded all components from the
test animal, ensuring that the UMAP projection was not shaped by potential noise patterns or other
unknown features of the test components that could affected the classifier result. We then tested
the first 20 components of each session of the test animal with 100 repetitions per component. In
every repetition, 1000 components from each PyN type were randomly selected from the pre-
computed UMAP space and we then assigned the PyN type of the test component based on the
identity of its 10 nearest neighbors in UMAP space. For LocaNMF components, we performed the
same procedure but additionally ensured to use an equal number of components from each seed
region and PyN type to prevent PyN types with a larger number of components in a given region
from biasing the classifier result. Classifier accuracy for each session (Fig. 2d,h) was then
computed as the mean probability over all repetitions to accurately identify the PyN type.

To determine the size of PyN-predictive LocaNMF components, we selected all spatial
components that achieved a classification accuracy of 99% percent or higher (all other components
were assigned as non-specific) and thresholded each component above 0.2 to obtain a binary
image. The size of each component was then computed as the square root of the sum of all pixels
and converted to square millimeters.

Linear encoding model

The linear encoding model was based on a combination of task- and movement-related variables,
as described previously[32]. Each variable consisted of multiple regressors that were combined into
a larger design matrix. Binary regressors contained a single pulse that signaled the occurrence of
specific events, such as the stimulus onset, and additional regression copies that were shifted
forward or backward in time to account for changes in cortical activity before or after the respective
event. For auditory stimuli, the time-shifted copies spanned all frames from the onset of the
auditory sequence until the end of the trial. Individual click sounds were also captured by an
additional regressor set that spanned the 2 seconds after click onset. For motor events, like licking
or whisking, the time-shifted copies spanned the frames from 1 s before until 2 s after each event.
Lastly, for some variables, such as the previous choice, the time-shifted copies spanned the whole
trial. Other variables were analog, such as measures from the piezo sensor or the pupil diameter,
and also contained the 200 highest temporal components of video information from both cameras
(using SVD as described above). This ensured that the model could account for animal movements
and accurately isolate task-related activity. Movement and task variables were additionally
decorrelated due to the variable durations of the initiation, stimulus and delay period. The model
was fit using ridge regression to allow for similar contributions from different correlated variables.
To determine the regularization penalty λ for each column of the widefield data, we used marginal
maximum likelihood estimation (MLE)[89]. MLE expresses the encoding model as a Bayesian linear
model and determines the ridge penalty λ by maximizing the marginal likelihood $\pi(D \mid \lambda)$ of the
model, given data D. This was done iteratively by testing different λ values to determine a global
minimum for the negative log-likelihood $-\log \pi(D \mid \lambda)$. The main advantage of this approach is
that λ can be determined without computationally expensive cross-validation procedures, resulting
in a ~50-fold decrease in required compute time on a regular work station. Moreover, the faster
MLE approach allows adjusting λ values for individual widefield data components which results
in higher cross-validated explained variance of the encoding model, compared to a regular cross-
validation approach (Supp. Fig. 16).

Variance analysis

Explained variance (cvR^2) was obtained using 10-fold cross-validation. This was done by fitting
the model weights to a continuous 90%-large section of the imaging data and then computing the
explained variance in the remaining 10% of the data. The procedure was repeated for 10 times,
while shifting the training and test data to ensure that each part of the recording was used in the
test data in one of the folds. To assess unique explained variance by individual variables (ΔR^2),
we created reduced models in which all regressors of a specific variable were shuffled in time.
Shuffling of each regressor was done within individual trials to account for a potential impact of
very slow temporal correlations due the kinetics of the calcium indicator. The difference in
explained variance between the full and the reduced model yielded the unique contribution ΔR^2 of
that model variable that could not be explained by other variables in the model. The same approach
was used to compute unique contributions for groups of variables, i.e., 'Movements' and 'Task'.
Here, all variables that corresponded to a given group were shuffled at once.

Decoding model

To predict animal's left/right choices from widefield data, we trained logistic regression decoders
with an L1 penalty on the temporal component matrix SV^T in each session. The L1 penalty was
defined as the inverse of the number of observations in the test dataset during cross-validation,
which yielded a good balance between the cross-validated prediction accuracy of the decoder and
the number of non-zero model regressors. When decoding choice, we randomly removed trials
until there was an equal amount of correct and incorrect trials where mice chose the left and the
right side. By balancing left/right choice sides and correct/incorrect trials, we ensured that the

decoder would not predict choices due to corresponding sensory information or would be influenced by potential side biases. The logistic regression model was implemented in Matlab using the 'fitclinear' function and run repeatedly for each time point in individual trials after re-aligning them to trial periods as described above. In each session, all decoder runs were performed with the same amount of trials (at least 250 trials) and we used 10-fold cross-validation to compute decoder accuracy at each time point in the trial. Beta weights were averaged from all models created during cross-validation and convolved with the spatial component matrix U to create cortical maps of the choice decoder weights.

Receiver operating characteristic (ROC) analysis

We computed the area under the ROC curve (AUC) to quantify choice preference of single neurons obtained from two-photon imaging. AUCs were computed by comparing the mean neural activity during the stimulus and delay period in all trials with ipsilateral versus contralateral choices. AUC values denote the specificity of the neural activity to ipsi- or contralateral choices, with values below 0.5 signifying ipsilateral choice-selectivity and AUC values above 0.5 contralateral choice-selectivity. To identify statistically significant choice-selective neurons, AUC values were also computed for shuffled trial labels (randomly assigning ipsi- and contralateral choices across trials) for each neuron. This procedure was repeated 100 times to create a distribution of shuffled AUC values for each neuron. A neuron's choice selectivity was then deemed significant if the probability of obtaining the actual AUC from the shuffled AUC distribution was less than 0.05.

Code availability statement: Code for preprocessing (e.g., hemodynamic correction and dimensionality reduction) is available here: https://github.com/churchlandlab/WidefieldImager/tree/master/Analysis. A cloud-based option for many of these steps is available at Neurocaas: http://neurocaas.org. To simplify the interaction with the online platform, a graphical user interface, dedicated to launching analysis and retrieving results from the NeuroCAAS platform is available here: (https://github.com/churchlandlab/wfield). Code for the ridge regression used in the encoding model is available on the lab's GitHub page: https://github.com/churchlandlab/ridgeModel.

Data availability statement: We will make preprocessed datasets from all animals available at time of publication. As in previous publications, we will also include a "readme" statement that explains how the data are stored. An example dataset has already been shared (DOI: 10.5281/zenodo.5834513) on Zenodo to demonstrate the feasibility of this pipeline: https://zenodo.org/record/5834513#.Yd3jHViZPX8.

1028 **Supplementary Table 1: Overview of different model variables**

1029

Variable name	Description	Variable type	Category
Hindlimb	Piezo sensor below the animal	Analog + Event kernel	Movement
Handles (Left / Right)	Touch events from handle sensors	Event kernel	Movement
Licks (Left / Right)	Lick events from spout sensors	Event kernel	Movement
Pupil	Pupil diameter, extracted from face camera	Analog + Event kernel	Movement
Nose	Nose movements, extracted from face camera	Analog + Event kernel	Movement
Whisking	Whisker movements, extracted from face camera	Analog + Event kernel	Movement
Body	Average motion energy across all body camera pixels	Analog + Event kernel	Movement
Video	Video dimensions from both cameras (SVD)	Analog	Movement
Video ME	Video dimensions from motion energy in both cameras	Analog	Movement
Choice (Left / Right)	All frames in either a left- or a rightward choice trial	Event kernel	Task
Previous choice	Every trial after a leftward choice trial	Event kernel	Task
Previous modality	Every trial after a visual trial	Event kernel	Task
Previous success	Every trial after a successful trial	Event kernel	Task
Success	All successful trials	Event kernel	Task
Water given	All frames after a water reward was given	Event kernel	Task
Auditory stimulus (Left / Right)	All frames after a left- or rightward auditory stimulus	Event kernel	Task
Visual stimulus (Left / Right)	All frames after a left- or rightward visual stimulus	Event kernel	Task

1030

1031 **Supplementary Table 2: Number of mice included for each experiment**

	Optogenetics	Widefield	Two-photon
Emx	9	4	-
CStr	9	4	-
Fezf2	8	5	-
PlexinD1	8	4	-
Camk2α-tTA;G6s2	-	-	3

1032

1033 **Supplementary Table 3: Resources**

Reagent/resource	Source	Identifier
Antibodies		
Goat polyclonal anti-GFP	Abcam	ab6673
Donkey Anti-Goat Alexa Fluor 488	Abcam	ab150129
Viral strains		
CAV-2-Cre	Plateforme de Vectorologie de Montpellier	N/A
AAV1-hSyn1-SIO-stGtACR2-FusionRed	Penn Vector Core	105677-AAV1
AAVrg-CAG-tdTomato	Penn Vector Core	59462-AAVrg
AAV-DJ-hSYN-DIO-hCAR{off} (Titer: 5.7e12 vg/ml)	Laboratory of Adam Kepecs	
Experimental Models		
Mouse: Emx1-IRES-Cre: Emx1$^{tm1(cre)Krj}$	The Jackson Laboratory	JAX#005628
Mouse: ROSA:LNL:tTA: Gt(ROSA)26Sor$^{tm1(tTA)Roos}$	The Jackson Laboratory	JAX#008600
Mouse: Camk2α-tTA: Tg(Camk2a-tTA)1Mmay	The Jackson Laboratory	JAX#003010
Mouse: Ai93(TITL-GCaMP6f)-D (Ai93D): Igs7$^{tm93.1(tetO-GCaMP6f)Hze}$	The Jackson Laboratory	JAX#024103
Mouse: Ai162(TIT2L-GC6s-ICL-tTA2)-D (Ai162D): Igs7$^{tm162.1(tetO-GCaMP6s,CAG-tTA2)Hze}$	H. Zeng, Allen Institute for Brain Science	JAX#031562
Mouse: TRE-GCaMP6s (G6s2): Tg(tetO-GCaMP6s)2Niell	The Jackson Laboratory	JAX#024742
Mouse: H2B-eGFP: Tg(HIST1H2BB/EGFP)1Pa	The Jackson Laboratory	JAX#006069
Mouse: Fezf2-2A-CreER: Fezf2$^{tm1.1(cre/ERT2)Zjh}$	The Jackson Laboratory	JAX#036296
Mouse: PlexinD1-2A-CreER: Plxnd1$^{tm2.1(flpo)Zjh}$	The Jackson Laboratory	JAX#036295
Software		
MATLAB 2018B	Mathworks	
Python 3.6.10	Python Software Foundation	
Other		
Bpod State Machine r0.5	Sanworks	N/A

1034

1035

References

1. Harris, K. D. & Mrsic-Flogel, T. D. Cortical connectivity and sensory coding. *Nature* **503**, 51–58 (2013).
2. Pfeffer, C. K., Xue, M., He, M., Huang, Z. J. & Scanziani, M. Inhibition of Inhibition in Visual Cortex: The Logic of Connections Between Molecularly Distinct Interneurons. *Nat Neurosci* **16**, 1068–1076 (2013).
3. Keller, A. J. *et al.* A Disinhibitory Circuit for Contextual Modulation in Primary Visual Cortex. *Neuron* **108**, 1181-1193.e8 (2020).
4. Tremblay, R., Lee, S. & Rudy, B. GABAergic Interneurons in the Neocortex: From Cellular Properties to Circuits. *Neuron* **91**, 260–292 (2016).
5. Cardin, J. A. *et al.* Driving fast-spiking cells induces gamma rhythm and controls sensory responses. *Nature* **459**, 663–667 (2009).
6. Veit, J., Hakim, R., Jadi, M. P., Sejnowski, T. J. & Adesnik, H. Cortical gamma band synchronization through somatostatin interneurons. *Nature Neuroscience* **20**, 951–959 (2017).
7. Sohal, V. S., Zhang, F., Yizhar, O. & Deisseroth, K. Parvalbumin neurons and gamma rhythms enhance cortical circuit performance. *Nature* **459**, 698–702 (2009).
8. Reimer, J. *et al.* Pupil fluctuations track fast switching of cortical states during quiet wakefulness. *Neuron* **84**, 355–362 (2014).
9. Fu, Y. *et al.* A Cortical Circuit for Gain Control by Behavioral State. *Cell* **156**, 1139–1152 (2014).
10. Zhou, M. *et al.* Scaling down of balanced excitation and inhibition by active behavioral states in auditory cortex. *Nature Neuroscience* **17**, 841–850 (2014).
11. Polack, P.-O., Friedman, J. & Golshani, P. Cellular mechanisms of brain state–dependent gain modulation in visual cortex. *Nature Neuroscience* **16**, 1331–1339 (2013).
12. Tasic, B. *et al.* Shared and distinct transcriptomic cell types across neocortical areas. *Nature* **563**, 72–78 (2018).
13. Economo, M. N. *et al.* Distinct descending motor cortex pathways and their roles in movement. *Nature* **563**, 79–84 (2018).
14. Chen, J. L., Carta, S., Soldado-Magraner, J., Schneider, B. L. & Helmchen, F. Behaviour-dependent recruitment of long-range projection neurons in somatosensory cortex. *Nature* **499**, 336–340 (2013).
15. Glickfeld, L. L., Andermann, M. L., Bonin, V. & Reid, R. C. Cortico-cortical projections in mouse visual cortex are functionally target specific. *Nat Neurosci* **16**, 219–226 (2013).
16. Kim, T. *et al.* Cortically projecting basal forebrain parvalbumin neurons regulate cortical gamma band oscillations. *PNAS* **112**, 3535–3540 (2015).
17. Harris, K. D. & Shepherd, G. M. G. The neocortical circuit: themes and variations. *Nature Neuroscience* **18**, 170–181 (2015).
18. Takahashi, N. *et al.* Active dendritic currents gate descending cortical outputs in perception. *Nature Neuroscience* 1–9 (2020) doi:10.1038/s41593-020-0677-8.
19. Tang, L. & Higley, M. J. Layer 5 Circuits in V1 Differentially Control Visuomotor Behavior. *Neuron* **105**, 346-354.e5 (2020).
20. Li, N., Chen, T.-W., Guo, Z. V., Gerfen, C. R. & Svoboda, K. A motor cortex circuit for motor planning and movement. *Nature* **519**, 51–56 (2015).
21. Chen, T.-W., Li, N., Daie, K. & Svoboda, K. A Map of Anticipatory Activity in Mouse Motor Cortex. *Neuron* **94**, 866-879.e4 (2017).
22. Wekselblatt, J. B., Flister, E. D., Piscopo, D. M. & Niell, C. M. Large-scale imaging of cortical dynamics during sensory perception and behavior. *Journal of Neurophysiology* **115**, 2852–2866 (2016).

23. Couto, J. *et al.* Chronic, cortex-wide imaging of specific cell populations during behavior.
Nat Protoc 1–25 (2021) doi:10.1038/s41596-021-00527-z.
24. Cardin, J. A., Crair, M. C. & Higley, M. J. Mesoscopic Imaging: Shining a Wide Light on
Large-Scale Neural Dynamics. *Neuron* **108**, 33–43 (2020).
25. Allen, W. E. *et al.* Global Representations of Goal-Directed Behavior in Distinct Cell Types
of Mouse Neocortex. *Neuron* **94**, 891-907.e6 (2017).
26. Matho, K. S. *et al.* Genetic dissection of the glutamatergic neuron system in cerebral cortex.
Nature **598**, 182–187 (2021).
27. Gerfen, C. R., Paletzki, R. & Heintz, N. GENSAT BAC cre-recombinase driver lines to
study the functional organization of cerebral cortical and basal ganglia circuits. *Neuron* **80**,
1368–1383 (2013).
28. Harris, J. A. *et al.* Anatomical characterization of Cre driver mice for neural circuit mapping
and manipulation. *Front Neural Circuits* **8**, 76 (2014).
29. Daigle, T. L. *et al.* A Suite of Transgenic Driver and Reporter Mouse Lines with Enhanced
Brain-Cell-Type Targeting and Functionality. *Cell* **174**, 465-480.e22 (2018).
30. Wang, Q. *et al.* The Allen Mouse Brain Common Coordinate Framework: A 3D Reference
Atlas. *Cell* **181**, 936-953.e20 (2020).
31. Garrett, M. E., Nauhaus, I., Marshel, J. H. & Callaway, E. M. Topography and Areal
Organization of Mouse Visual Cortex. *J. Neurosci.* **34**, 12587–12600 (2014).
32. Musall, S., Kaufman, M. T., Juavinett, A. L., Gluf, S. & Churchland, A. K. Single-trial
neural dynamics are dominated by richly varied movements. *Nat Neurosci* **22**, 1677–1686
(2019).
33. Saxena, S. *et al.* Localized semi-nonnegative matrix factorization (LocaNMF) of widefield
calcium imaging data. *PLOS Computational Biology* **16**, e1007791 (2020).
34. Condylis, C. *et al.* Dense functional and molecular readout of a circuit hub in sensory cortex.
Science **375**, eabl5981 (2022).
35. McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and
Projection for Dimension Reduction. *arXiv:1802.03426 [cs, stat]* (2018).
36. Brunton, B. W., Botvinick, M. M. & Brody, C. D. Rats and Humans Can Optimally
Accumulate Evidence for Decision-Making. *Science* **340**, 95–98 (2013).
37. Salkoff, D. B., Zagha, E., McCarthy, E. & McCormick, D. A. Movement and Performance
Explain Widespread Cortical Activity in a Visual Detection Task. *Cereb Cortex* **30**, 421–437
(2020).
38. Orsolic, I., Rio, M., Mrsic-Flogel, T. D. & Znamenskiy, P. Mesoscale cortical dynamics
reflect the interaction of sensory evidence and temporal expectation during perceptual
decision-making. *Neuron* **109**, 1861-1875.e10 (2021).
39. Steinmetz, N. A., Zatka-Haas, P., Carandini, M. & Harris, K. D. Distributed coding of
choice, action and engagement across the mouse brain. *Nature* **576**, 266–273 (2019).
40. Bergmann, R. *et al.* Predicting behavior from eye movement and whisking asymmetry.
2021.02.11.430785 Preprint at https://doi.org/10.1101/2021.02.11.430785 (2021).
41. Esmaeili, V. *et al.* Rapid suppression and sustained activation of distinct cortical regions for
a delayed sensory-triggered motor response. *Neuron* **109**, 2183-2201.e9 (2021).
42. Hanks, T. D. *et al.* Distinct relationships of parietal and prefrontal cortices to evidence
accumulation. *Nature* **520**, 220–223 (2015).
43. Goard, M. J., Pho, G. N., Woodson, J. & Sur, M. Distinct roles of visual, parietal, and frontal
motor cortices in memory-guided sensorimotor decisions. *eLife* **5**, e13764 (2016).
44. Peters, A. J., Fabre, J. M. J., Steinmetz, N. A., Harris, K. D. & Carandini, M. Striatal activity
topographically reflects cortical activity. *Nature* 1–6 (2021) doi:10.1038/s41586-020-03166-
8.
45. Oh, S. W. *et al.* A mesoscale connectome of the mouse brain. *Nature* **508**, 207–214 (2014).

46. Mahn, M. *et al.* High-efficiency optogenetic silencing with soma-targeted anion-conducting
channelrhodopsins. *Nature Communications* **9**, 4125 (2018).
47. Li, S.-J., Vaughan, A., Sturgill, J. F. & Kepecs, A. A Viral Receptor Complementation
Strategy to Overcome CAV-2 Tropism for Efficient Retrograde Targeting of Neurons.
Neuron **98**, 905-917.e5 (2018).
48. Znamenskiy, P. & Zador, A. M. Corticostriatal neurones in auditory cortex drive decisions
during auditory discrimination. *Nature* **497**, 482–485 (2013).
49. Guo, Z. V. *et al.* Flow of Cortical Activity Underlying a Tactile Decision in Mice. *Neuron*
81, 179–194 (2014).
50. MacDowell, C. J. & Buschman, T. J. Low-Dimensional Spatiotemporal Dynamics Underlie
Cortex-wide Neural Activity. *Current Biology* **30**, 2665-2680.e8 (2020).
51. Stringer, C. *et al.* Spontaneous behaviors drive multidimensional, brainwide activity. *Science*
364, eaav7893 (2019).
52. Logothetis, N. K. *et al.* Hippocampal-cortical interaction during periods of subcortical
silence. *Nature* **491**, 547–553 (2012).
53. Rogers, B. P., Morgan, V. L., Newton, A. T. & Gore, J. C. Assessing Functional
Connectivity in the Human Brain by FMRI. *Magn Reson Imaging* **25**, 1347–1357 (2007).
54. Vanni, M. P. & Murphy, T. H. Mesoscale Transcranial Spontaneous Activity Mapping in
GCaMP3 Transgenic Mice Reveals Extensive Reciprocal Connections between Areas of
Somatomotor Cortex. *J. Neurosci.* **34**, 15931–15946 (2014).
55. Vanni, M. P., Chan, A. W., Balbi, M., Silasi, G. & Murphy, T. H. Mesoscale Mapping of
Mouse Cortex Reveals Frequency-Dependent Cycling between Distinct Macroscale
Functional Modules. *J. Neurosci.* **37**, 7513–7533 (2017).
56. Macé, É. *et al.* Whole-Brain Functional Ultrasound Imaging Reveals Brain Modules for
Visuomotor Integration. *Neuron* **100**, 1241-1251.e7 (2018).
57. Matsui, T., Murakami, T. & Ohki, K. Transient neuronal coactivations embedded in globally
propagating waves underlie resting-state functional connectivity. *PNAS* **113**, 6556–6561
(2016).
58. Huang, L. *et al.* BRICseq Bridges Brain-wide Interregional Connectivity to Neural Activity
and Gene Expression in Single Animals. *Cell* **182**, 177-188.e27 (2020).
59. Mohajerani, M. H. *et al.* Spontaneous cortical activity alternates between motifs defined by
regional axonal projections. *Nat Neurosci* **16**, 1426–1435 (2013).
60. Lohani, S. *et al. Dual color mesoscopic imaging reveals spatiotemporally heterogeneous*
coordination of cholinergic and neocortical activity. 2020.12.09.418632
https://www.biorxiv.org/content/10.1101/2020.12.09.418632v1 (2020)
doi:10.1101/2020.12.09.418632.
61. Gilad, A. & Helmchen, F. Spatiotemporal refinement of signal flow through association
cortex during learning. *Nature Communications* **11**, 1–14 (2020).
62. Gallero-Salas, Y. *et al.* Sensory and Behavioral Components of Neocortical Signal Flow in
Discrimination Tasks with Short-Term Memory. *Neuron* **109**, 135-148.e6 (2021).
63. Park, J. M. *et al.* Deep and superficial layers of the primary somatosensory cortex are critical
for whisker-based texture discrimination in mice. *bioRxiv* 2020.08.12.245381 (2020)
doi:10.1101/2020.08.12.245381.
64. Gilad, A., Gallero-Salas, Y., Groos, D. & Helmchen, F. Behavioral Strategy Determines
Frontal or Posterior Location of Short-Term Memory in Neocortex. *Neuron* **99**, 814-828.e7
(2018).
65. Jun, E. J. *et al.* Causal role for the primate superior colliculus in the computation of evidence
for perceptual decisions. *Nat Neurosci* (2021) doi:10.1038/s41593-021-00878-6.

66. Wang, L., McAlonan, K., Goldstein, S., Gerfen, C. R. & Krauzlis, R. J. A Causal Role for Mouse Superior Colliculus in Visual Perceptual Decision-Making. *J. Neurosci.* **40**, 3768–3782 (2020).

67. Felsen, G. & Mainen, Z. F. Neural substrates of sensory-guided locomotor decisions in the rat superior colliculus. *Neuron* **60**, 137–148 (2008).

68. Duan, C. A. *et al.* A cortico-collicular pathway for motor planning in a memory-dependent perceptual decision task. *Nature Communications* **12**, 2727 (2021).

69. Erlich, J. C., Brunton, B. W., Duan, C. A., Hanks, T. D. & Brody, C. D. Distinct effects of prefrontal and parietal cortex inactivations on an accumulation of evidence task in the rat. *eLife Sciences* **4**, e05457 (2015).

70. Raposo, D., Kaufman, M. T. & Churchland, A. K. A category-free neural population supports evolving demands during decision-making. *Nat Neurosci* **17**, 1784–1792 (2014).

71. Harvey, C. D., Coen, P. & Tank, D. W. Choice-specific sequences in parietal cortex during a virtual-navigation decision task. *Nature* **484**, 62–68 (2012).

72. Pinto, L. *et al.* Task-Dependent Changes in the Large-Scale Dynamics and Necessity of Cortical Regions. *Neuron* **104**, 810-824.e9 (2019).

73. Licata, A. M. *et al.* Posterior Parietal Cortex Guides Visual Decisions in Rats. *J. Neurosci.* **37**, 4954–4966 (2017).

74. Zatka-Haas, P., Steinmetz, N. A., Carandini, M. & Harris, K. D. Sensory coding and the causal impact of mouse cortex in a visual decision. *eLife* **10**, e63163 (2021).

75. Kim, J.-H., Ma, D.-H., Jung, E., Choi, I. & Lee, S.-H. Gated feedforward inhibition in the frontal cortex releases goal-directed action. *Nat Neurosci* **24**, 1452–1464 (2021).

76. Wu, Z. *et al.* Context-Dependent Decision Making in a Premotor Circuit. *Neuron* **106**, 316-328.e6 (2020).

77. Coen, P., Sit, T. P. H., Wells, M. J., Carandini, M. & Harris, K. D. *Mouse frontal cortex mediates additive multisensory decisions*. 2021.04.26.441250 https://www.biorxiv.org/content/10.1101/2021.04.26.441250v2 (2021) doi:10.1101/2021.04.26.441250.

78. Park, J., Phillips, J. W., Martin, K. A., Hantman, A. W. & Dudman, J. T. *Descending neocortical output critical for skilled forelimb movements is distributed across projection cell classes*. 772517 https://www.biorxiv.org/content/10.1101/772517v3 (2021) doi:10.1101/772517.

79. Callaway, E. M. *et al.* A multimodal cell census and atlas of the mammalian primary motor cortex. *Nature* **598**, 86–102 (2021).

80. Laboulaye, M. A., Duan, X., Qiao, M., Whitney, I. E. & Sanes, J. R. Mapping Transgene Insertion Sites Reveals Complex Interactions Between Mouse Transgenes and Neighboring Endogenous Genes. *Frontiers in Molecular Neuroscience* **11**, 385 (2018).

81. Ratzlaff, E. H. & Grinvald, A. A tandem-lens epifluorescence macroscope: hundred-fold brightness advantage for wide-field imaging. *J. Neurosci. Methods* **36**, 127–137 (1991).

82. Lerner, T. N. *et al.* Intact-Brain Analyses Reveal Distinct Information Carried by SNc Dopamine Subcircuits. *Cell* **162**, 635–647 (2015).

83. Marshel, J. H., Garrett, M. E., Nauhaus, I. & Callaway, E. M. Functional Specialization of Seven Mouse Visual Cortical Areas. *Neuron* **72**, 1040–1054 (2011).

84. Pachitariu, M. *et al.* Suite2p: beyond 10,000 neurons with standard two-photon microscopy. *bioRxiv* 061507 (2016) doi:10.1101/061507.

85. Chuong, A. S. *et al.* Noninvasive optical inhibition with a red-shifted microbial rhodopsin. *Nat Neurosci* **17**, 1123–1129 (2014).

86. Kim, Y. *et al.* Brain-wide Maps Reveal Stereotyped Cell-Type-Based Cortical Architecture and Subcortical Sexual Dimorphism. *Cell* **171**, 456-469.e22 (2017).

87. Klein, S., Staring, M., Murphy, K., Viergever, M. A. & Pluim, J. P. W. elastix: A Toolbox
for Intensity-Based Medical Image Registration. *IEEE Transactions on Medical Imaging* **29**,
196–205 (2010).
88. Reddy, B. S. & Chatterji, B. N. An FFT-based technique for translation, rotation, and scale-
invariant image registration. *IEEE Trans Image Process* **5**, 1266–1271 (1996).
89. Karabatsos, G. Marginal maximum likelihood estimation methods for the tuning parameters
of ridge, power ridge, and generalized ridge regression. *Communications in Statistics -*
Simulation and Computation (2017).
1240