Comammox *Nitrospira* bacteria outnumber canonical nitrifiers irrespective of electron donor mode and availability.

Katherine J. Vilardi[1], Irmarie Cotto[1], Maria Sevillano Rivera[1], Zihan Dai[2,3], Christopher L. Anderson[1], Ameet Pinto[4]

[1] Department of Civil and Environmental Engineering, Northeastern University, Massachusetts, MA, USA
[2] Key Laboratory of Drinking Water Science and Technology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing, China
[3] University of Chinese Academy of Sciences, Beijing, China
[4] School of Civil and Environmental Engineering, Georgia Institute of Technology, Atlanta, GA, USA.

Corresponding author email: apinto36@gatech.edu

Keywords: comammox bacteria, nitrification, drinking water biofiltration, electron donor

47
<u>Abstract</u>

Complete ammonia oxidizing bacteria coexist with canonical ammonia and nitrite oxidizing

bacteria in a wide range of environments. Whether this coexistence is due to competitive or

cooperative interactions between the three guilds, or a result of niche separation is not yet clear.

Understanding the factors driving coexistence of nitrifying guilds is critical to effectively manage

nitrification processes occurring in engineered and natural ecosystems. In this study, microcosms-

based experiments were used to investigate the impact of electron donor mode (i.e., ammonia and

urea) and loading on the population dynamics of nitrifying guilds in drinking water biofilter media.

Shotgun sequencing of DNA from select time points followed by co-assembly and re-construction

of metagenome assembled genomes (MAGs) revealed multiple clade A2 and one clade A1

comammox bacterial populations coexisted in the microcosms alongside *Nitrosomonas-like*

ammonia oxidizers and *Nitrospira*-like nitrite oxidizer populations. Clade A2 comammox bacteria

were likely the primary nitrifiers within the microcosms and increased in abundance over canonical

ammonia and nitrite oxidizing bacteria irrespective of electron donor mode or nitrogen loading

rates. This suggests that comammox bacteria will outnumber nitrifying communities sourced from

oligotrophic environments irrespective of variable nitrogen regimes. Changes in comammox

bacterial abundance were not correlated with either ammonia or nitrite oxidizing bacterial

abundance in urea amended systems where metabolic reconstruction indicated potential cross

feeding between ammonia and nitrite oxidizing bacteria. In contrast, comammox bacterial

abundance demonstrated a negative correlation with that of nitrite oxidizers in ammonia amended

systems. This suggests that potentially weaker synergistic relationships between ammonia and

nitrite oxidizers might enable comammox bacteria to displace nitrite oxidizers from complex

nitrifying communities.

Importance

A deeper understanding of the interactions between nitrifying microorganisms, including comammox bacteria, is crucial to effectively managing nitrogen in engineered and natural systems. Despite their ubiquitous detection, our understanding of interactions between these coexisting nitrifying groups is limited. Here, we investigate the influence of differing sources of electron donor and loadings on a mixed nitrifier community which includes canonical ammonia and nitrite oxidizers, as well as comammox bacteria. Our results indicate that comammox bacteria will dominate nitrifier communities sourced from low nitrogen environments (e.g., drinking water) irrespective of fluctuations in nitrogen availability or source without directly competing with canonical ammonia oxidizers. However, canonical nitrite oxidizers are likely to get outcompeted by comammox bacteria in situations where metabolic interactions between ammonia and nitrite do not include cross feeding.

Introduction

Nitrification, the biological transformation of ammonia to nitrate via nitrite, is an important process in engineered and natural ecosystems. While nitrification mediated by ammonia oxidizing microorganisms (AOM) (1, 2), including ammonia oxidizing bacteria (AOB) and archaea (AOA), and nitrite oxidizing bacteria (NOB) (3) has been extensively investigated, complete ammonia oxidation (comammox) performed by comammox bacteria is understudied in large part due to its recent discovery. All known comammox bacteria belong to *Nitrospira* sub-lineage II (4-6) and are currently divided into two clades, A and B, with clade A further separated into sub-clades A1 and A2 (7). Due to close phylogenetic relatedness, comammox-*Nitrospira* cannot be distinguished from *Nitrospira*-NOB based on the 16S rRNA gene sequence or the marker genes for nitrite oxidation (*nxrAB*) (4). Thus, characterization of comammox bacteria has been largely enabled by shotgun DNA sequencing followed by reconstruction of assembled genomes (6, 8-12) and the development of primers targeting subuntis of comammox bacteria ammonia monooxygenase (*amo*) gene (13-18).

Within the engineered water cycle, clade A1 comammox bacteria have been primarily detected in wastewater treatment plants while clade A2 and B are been associated with drinking water treatment and distribution systems (7). It is unclear if this translates into physiological differences between the clades/sub-clades since there is only one comammox isolate and an enrichment enrichments whose kinetics have been characterized. To date, kinetic parameters of comammox bacteria are confined to two clade A representatives, cultured *Candidatus* N. inopinata and an enrichment of *Candidatus* N. kreftii (19, 20). Both demonstrate a high affinity for ammonia, with half-saturation constants orders of magnitude lower than strict AOB. Comparatively, the

Candidatus N. kreffti enrichment exhibited a higher affinity for nitrite compared to *Candidatus* N.

inopinata and partial inhibition of ammonia oxidation even at low ammonia concentrations (20).

This suggests that clade-specific comammox bacterial niche, if applicable, may be arise from a

combination of factors ranging from affinity to inhibition. Beyond clade specific traits, identifying

the potential environmental and physiological factors driving the coexistence of comammox

bacteria with canonical nitrifiers is also important to better understand comammox bacteria role in

complex nitrifying communities (21-29). Comammox bacteria have been detected along with their

canonical nitrifying counterparts in wastewater treatment plants (18, 25, 30-32), drinking water

systems (6, 9, 15, 33, 34) and soils (22, 28, 35, 36) at varying abundances over a wide range of

ammonium concentrations. While there is currently no quantitative estimate of the contribution of

comammox bacteria to nitrification compared to AOB and NOB, several studies have investigated

comammox bacterial dynamics in the context of mixed nitrifying communities. For instance,

DNA/RNA stable isotope probing combined with transcriptional analysis provided support for

comammox *Nitrospira* contributing to ammonia oxidation in lab-scale biofilters exposed to very

low ammonium concentrations (21). Soil microcosm amended with high ammonia concentrations

were enriched in AOB compared to those with lower ammonia concentrations where clade B

comammox bacteria proliferated (24, 28). Interestingly, in a lab-scale partial nitrification-

anammox reactor operating with incrementally increased ammonia loadings, comammox bacteria

initially dominated over strict AOB but its abundance significantly declined as loadings were

further increased (29).

Comammox bacteria may also acquire ammonia via urea degradation. Specifically, genes

encoding for urea transport and the urease enzyme are distributed among many *Nitrospira*

populations (37), including most comammox populations (38). While this may diversify potential

nitrogen sources for comammox bacteria (3), this could be a potential advantage for canonical

nitrifiers involved in a reciprocal feeding strategy as observed with co-cultured *Nitrospira*

moscoviensis converting urea to ammonia for *Nitrosomonas europaea* (37). The tight interplay

between canonical nitrifiers is well established; however, our understanding of comammox

competition (or lack thereof) with AOM and its impact on strict NOB in mixed communities is

limited. To better understand the comammox bacterial role within these complex nitrifying

communities, we investigated their population dynamics across two electron donor modes

(ammonia or urea) at three total nitrogen dosing strategies. The objectives of this study were (1)

to determine if comammox bacteria and canonical nitrifiers exhibit concentration and nitrogen

source dependent dynamics when subject to repeat nitrogen amendments and (2) to determine if

these dynamics are consistent or variable at the clade or population within each functional guild.

Results

Microbial community composition in microcosms and nitrogen biotransformation potential.

Microcosms consisting of granular activated carbon (GAC) from drinking water biofilters were

subject to intermittent amendments of nitrogen using two sources of electron donor (ammonia or

Urea) across three nitrogen concentrations (14, 3.5, and 1.5 mg-N/L). The conditions used in these

experiments are denoted as 14A, 3.5A, 1.5A, 14U, 3.5U and 1.5U where A or U represents

ammonia or urea amendments, respectively, and the number represents the concentration of

electron donor spike in mg/L as nitrogen. Two microcosms were sacrificed on a weekly basis over

the dotation of a eight week experiment (n=96 total microcosms). Extracted DNA from the inocula

and weeks four and eight were subject to shotgun DNA sequencing (n=13).

Initial assessment of taxonomic diversity in the samples based on analyses of metagenomic reads

mapping to the small subunit rRNA database (SILVA SSU NR99 version 138.1) indicated that the

GAC inocula largely consisted of bacteria with archaea and eukaryota constituting a small

proportion of the overall metagenomic reads (~0.002%). The bacterial community was primarily

composed of Gammaproteobacteria (30-20%), Alphaproteobacteria (25-31%) and Nitrospirota (8-

15%) (Figure 1A). *Nitrospira* and *Nitrosomonadaceae* were the only nitrifiers identified and

constituted 9-15% of the overall microbial community in samples. Full length 16S rRNA gene

sequences were assembled from each sample (n=13) resulting in a total of eight sequences with

closest matching SILVA database hits to uncultured *Nitrospira* bacteria (Accession numbers:

MF040566, AY328760, JN868922). Clustering of all eight *Nitrospira* 16S rRNA gene sequences

at 99% identity resulted in two *Nitrospira* operational taxonomic units (OTUs) with one cluster

composed of six sequences (Nitrospira OTU 1) and the other cluster with two sequences

(Nitrospira OTU 2). Phylogenetic placements of these OTUs revealed both clustered within

Nitrospira sublineage II (supplementary figure S1A). Diversity of *Nitrospira* was likely

underrepresented as full length *Nitrospira* 16S rRNA gene sequences could not be assembled from

some samples despite a large portion of extracted 16S rRNA gene reads mapping to *Nitrospira*

references in the SILVA database. Limited assembly of these reads could be due to several closely

related *Nitrospira* species/strains coexisting in the samples making re-construction of full length

sequences difficult. For canonical AOB, Nitrosomonas sp. AL212 (CP002552) was the closest

matching database hit to one assembled sequence while another six had hits closet to

Nitrosomonadaceae (Accession numbers: FPLP01009519, KJ807851, FPLK01002446) but could

not be further classified at the genus or species level. Phylogenetic placement of the single

Nitrosomonas OTU affiliated it with *Nitrosomonas* sp. AL212 and *Nitrosomonas ureae* (Figure

S1B).

Following co-assembly of metagenomic reads, predicted protein coding genes from scaffolds

associated with the nitrogen metabolism were taxonomically classified (Figure 1B). The majority

of methane/ammonia monooxygenase (*pmo-amo*) like genes (KEGG orthology: K10944, K10945,

K10946) were associated with either nitrifiers (i.e., *Nitrospira* or *Nitrosomonas*) or methanotrophs

(i.e., *Methylocystis*) (Figure 1C). While some *amoCAB* genes could not be classified to the genus

level using kaiju software, blastp searches against the NCBI non-redundant protein database

indicated these were closely related to *Nitrosomonas*. All retrieved *hao* sequences (KEGG

orthology: K10535) were associated with Nitrospira which is likely due to the low relative

abundance of *Nitrosomonas*-like populations and the resulting inability to assemble their *hao*

genes. Potential for ureolytic activity was detected across four phyla based on the urease alpha

subunit (*ureC*). *ureC* sequences associated with Nitrospirota and Gammaproteobacteria could be

classified at the genus level as *Nitrospira* and *Nitrosomonas*. Sequences identified as nitrate

reductase/nitrite oxidoreductase alpha and beta subunits (K00370, K00371) were subject to further

classification to differentiate between nitrite oxidoreductase genes belonging to NOB from nitrate

reductases belonging to other community members. Phylogenetic placement of most *Nitrospira*

nxrA sequences found in this study cluster within a branch containing both comammox and

Nitrospira-NOB species (Candidatus *N. inopinata*, Candidatus *N. nitrosa* and *N. defluvii*) (Figure

1D). While other sequences clustered on a separate branch with Candidatus *N. nitrificans*, a single

Nitrospira nxrA sequence clustered closely within a branch containing only *Nitrospira*-NOB

belonging to sublineage II.

<u>Phylogenomic placement of nitrifying populations and their metabolism.</u>

To further define the composition of nitrifiers, metagenome assembled genomes (MAGs) were

obtained from GAC microcosms after dereplication of MAGs from three binning approaches. All

204 MAGs were classified as bacteria, with 145 MAGs exhibiting completeness greater than 70%

and contamination less than 10% (Table S1). Approximately 62% of the metagenomic reads

mapped to these MAGs. Nine MAGs in total were classified as nitrifying bacteria belonging to

Nitrosomonas and *Nitrospira* (Table S2). Genome annotation confirmed that four *Nitrospira*

MAGs had key ammonia (ammonia monooxygenase and hydroxylamine oxidoreductase) and

nitrite (nitrite oxidoreductase) oxidation genes (Figure S2). Quality assessment for these

comammox MAGs indicated two high (Bin_49_2_2 and Bin_49_4) and one medium quality

(Bin_260) (Table S1) according to (39). A fourth comammox MAG (Bin_13) was assembled with

high completeness (89%) but also possessed high redundancy (18%) that could not be improved

with further manual refinement. The remaining two *Nitrospira* MAGs (Bin_7_1 and Bin_188),

which were likely strict NOB due to lack of ammonia oxidation genes, were less complete (38.04%

and 48.25%) with low redundancy (8.76% and 8.46%). The low completeness was likely not due

to their lower abundance, but potentially high level of strain heterogeneity which may have

affected the assembly of reads associated with *Nitrospira*-NOB. For example, RPKM-based

relative abundance estimated using all reads (total RPKM) showed the two *Nitrospira*-NOB

MAGs exhibited similar relative abundance to comammox bacteria MAGs Bin_49_2_2 and

Bin_49_4 (~7-10 total RPKM), but the CheckM estimated strain heterogeneities for Bin_7_1 and

Bin_188 were 40 and 75, respectively, compared to 0 for both Bin_49_2_2 and Bin_49_4. Two

MAGs classifying as *Nitrosomonas* were deemed high (Bin_83) and medium quality (Bin_168);

however, a third *Nitrosomonas* MAG was considered low quality.

A maximum likelihood tree based on 91 single copy core genes confirmed all *Nitrospira* MAGs

affiliated with sublineage II (Figure 2A). Four of the *Nitrospira* MAGs from this study clustered

within clade A comammox *Nitrospira* (Bin_49_2_2, Bin_49_4, Bin_260 and Bin_13) but were

separated into distinct groups on the phylogenomic tree; namely, forming three clusters with

MAGs obtained from tap water, drinking water filters, and freshwater. *amoA*-based phylogenetic

analysis corroborated their placement into clade A (Figure 2B); however, *hao*-based phylogeny

distinguished three of comammox MAGs (Bin_49_2_2, Bin_49_4, Bin_260) as clade A2 (Palomo

et al. 2019) while one clustered within clade A1 (Bin_13) (Figure 2C). Consistent across all trees,

Bin_49_2_2 and Bin_260 cluster closely with comammox MAGs *Nitrospira* sp. SG-bin2 and ST-

bin4 (ANI ~ 92%) derived from tap water metagenomes (9). Bin_49_4 clustered closely with

Nitrospirae bacterium Ga0074138 (ANI ~ 99%), which was previously detected in GAC from the

same drinking water treatment plant (6), along with other tap water and groundwater-fed rapid

sand filter MAGs (8, 9). Bin_13 associated with comammox MAGs obtained from freshwater,

UBA5698 and UBA5702 (40) (ANI ~ 90%); however, its high contamination (18%) likely renders

ANI comparison less accurate. Overall, the MAGs demonstrated less then 95% ANI to other

reference comammox bacterial MAGs (Figure S3) suggesting comammox bacteria detected in

GAC microcosms are distinct from one another and previously published comammox MAGs; as

a result, they are likely novel *Nitrospira* species. The two remaining *Nitrospira* MAGs, Bin_7_1

and Bin_188, clustered with strict *Nitrospira*-NOB MAGs recovered from tap water,

Nitrospira_sp_ST-bin5 (9) (ANI ~ 94%), and a rapid sand filter, Nitrospira CG24D (ANI ~ 87%)

(8) (Figure 2A and S3). Only two strict AOB MAGs (Bin_83 and Bin_168) from this study were

used for phylogenomic analysis due high redundancy and low completeness of the third (Bin_195).

Both Bin_83 and Bin_168 originate from *Nitrosomonas* cluster 6a and clustered closely with

Nitrosomonas ureae and *Nitrosomonas* sp. AL212 (Figure 2D). Bin_168 shares a high sequence

similarity to *N. ureae* (ANI ~ 98%) while Bin_83 shares less than 83% ANI to any of the references

on the tree including Bin_168.

All comammox MAGs demonstrated the potential for ureolytic activity with the presence of the

ureABC operon in addition to most genes for urease accessory proteins (Figure S2). *Nitrospira*-

NOB MAGs did not contain genes encoding for urease; however, two *ureC* sequences found on

assembled scaffolds that were classified as *Nitrospira* but were not binned into any of the

Nitrospira MAGs. Queries of these *ureC* genes against the NCBI non-redundant database revealed

one sequence shared the highest percent identity to *Nitrospira lenta* and *Nitrospira moscoviensis*

while top hits for the second sequence belonged to an unclassified *Nitrospira*. One *Nitrospira*-

NOB MAG (Bin_7_1) did harbor genes for the urea transport system permease proteins (*urtBC*),

urea transport system substrate-binding proteins (*urtA*) and urea transport system ATP-binding

proteins (*urtDE*). This suggests that the two unbinned *ureC* genes likely belonged to *Nitrospira*-

like NOB bacteria. *Nitrosomonas* MAGs Bin_168 and Bin_83 each contained the *ureCAB* operon

and some genes for urease accessory proteins and urea transport. A third *ureC* sequence found in

the metagenome classified as *Nitrosomonas* but was not binned into any *Nitrosomonas* MAGs.

The impact of nitrogen amendments on nitrifying populations.

To address concentration and nitrogen source-dependent dynamics of the three nitrifier

populations detected in our metagenomic analysis, qPCR-assays were used to estimate their

abundances over time in the nitrogen amended microcosms. In the high ammonia amendment

(14A), strict AOB relative abundance increased 2.4-fold from weeks 1-3 but remained below 2%

of total bacteria for the duration of the experiment whereas comammox relative abundance

increased markedly over time reaching 2.8% of total bacteria by end of the experiment (Figure

3B). Similar to strict AOB, *Nitrospira*-NOB relative abundance increased early on but thereafter

reduced from 4% at its peak in week two to 1.8% by week eight. Weekly measurements for

nitrogen concentrations taken alongside biomass samples indicated the presence of residual

ammonia and accumulated nitrite concentrations were highest during the first three weeks of the

experiment but gradually reduced over time with most inorganic nitrogen present as nitrate (Figure

S4). While comammox bacteria were always dominant, the abundance of strict AOB as a portion

of AOM was significantly higher when ammonia and nitrite accumulated in weeks 1-3 as

compared to weeks 5-8 (Welsh's t-test, p-value < 0.05) (Figure 3A).

The qPCR data was in concordance with metagenomic estimation of MAG abundance with clade

A2 comammox (Bin_49_2_2, Bin_49_2 and Bin_260) highly abundant compared to strict AOB

(Bin_83, Bin_168 and Bin_195) and clade A1 comammox (Bin_13) in the inocula and at weeks

four and eight in the high ammonia amendment (Figure 3C). In particular, clade A2 MAGs

Bin_49_2_2 and Bin_49_4 were the most dominant comammox populations while strict AOB was

dominated by Bin_83 at each time point. *Nitrospira*-NOB MAGs had comparable abundance to

clade A2 comammox MAGs but displayed limited variation in abundance in the high ammonia

amendments. This contrasts with the qPCR data, where *Nitrospira*-NOB were significantly more

abundant than comammox bacteria at earlier timepoints and then demonstrated a significant

decrease in abundance over time. This is likely due to the fact that the two assembled Nitrospira-

NOB MAG's do not represent the entirety of NOB diversity in the microcosms and the fact that

metagenomic data is only available for select timepoints as compared to qPCR data.

Nitrifier populations in mid and low ammonia amendments displayed similar dynamics to those

observed in high ammonia with comammox relative abundance increasing to 3% and 2.2% of total

bacteria by week eight, respectively. Interestingly, Bin_260, the least abundant clade A2

comammox MAG in the inocula, demonstrated significant increase in abundance in the low

ammonia amendment over the course of the experiment compared to its abundance in the other

ammonia amendments. Consistent with the ammonia amended microcosms, strict AOB in urea

amended microcosms increased in relative abundance only at earlier time points followed by low

but stable relative abundance (~2% of total bacteria). In the high urea amendment, relative

abundance of comammox bacteria remained largely unchanged at earlier time points followed by

an increase in abundance. Despite this, mean relative abundance of comammox bacteria compared

to strict AOB was still approximately 2-fold greater in all urea amendments. Similar to the

ammonia amendments, *Nitrospira*-NOB relative abundance did increase initially followed by a

decline in all urea amendments. Interestingly though, the relative abundance of comammox

bacteria and *Nitrospira*-NOB were similar in the later weeks of the experiment after *Nitrospira*-

NOB's initial rise in urea amendments. Clade A2 comammox MAG Bin_260 was consistently

lower in abundance than Bin_49_2_2 and Bin_49_2 in the urea amendments except for mid urea.

Abundance of the clade A1 comammox MAG remained lower than all clade A2 MAGs and

displayed minimal enrichment in all the urea amendments which was consistent with ammonia

amended microcosms. Bin_168, which showed high sequence similarity to *Nitrosomonas ureae*,

did not exhibit enrichment in any of the urea amendments and remained low in abundance with all

other strict AOB MAGs.

There was no significant difference in mean relative abundance of strict AOB or *Nitrospira*-NOB

between the high ammonia (14A) and urea amendments (14U) (Welch t-test, $p > 0.05$) but the

mean relative abundance of comammox bacteria was significantly greater in high ammonia than

in the high urea amendment (Welch t-test, $p < 0.05$). Comparatively, out of all nitrogen

amendments, mean relative abundance of comammox bacteria was the lowest in high urea (1.8%

of total bacteria). Comparisons between the mid ammonia (3.5A) and urea amendments (3.5U) as

well as the low ammonia (1.5A) and urea (1.5U) amendments revealed no significant difference

in mean relative abundance for any of the nitrifier populations (Welch t-test, $p > 0.05$).

Additionally, no significant differences were detected when testing the mean relative abundance

of the three nitrifier populations between high, mid, and low concentrations within each

amendment type (ANOVA, $p > 0.05$).

327
328
The relative abundance of the nitrifying groups were used to examine potential correlations

between the different populations in each of the nitrogen amendments. The ratio of comammox

bacteria as portion of AOM to comammox bacteria as a portion of total *Nitrospira* revealed a

strong positive relationship in all amendments (Pearson R = 0.75-0.87, $p < 0.001$) (Figure S5A),

however, the change in relative abundance of comammox bacteria was not directly correlated with

that of strict AOB in any of the nitrogen amendments (Figure S5B). Strict AOB and *Nitrospira*-

NOB abundances were strongly correlated for all urea amendments and high (14A) and mid

ammonia (3.5A) (Pearson r = 0.58-0.82, $p < 0.05$, Figure 4A) but exhibited a weaker relationship

in low ammonia (Pearson r = 0.42, $p > 0.05$). Interestingly, while comammox bacteria abundance

was significantly and negatively correlated with that of *Nitrospira*-NOB in ammonia amendments

(Pearson r = -0.37 to -0.61) (p < 0.05), there was no significant association between them in the

urea amendments (p > 0.05) (Figure 4B).

341
342
343   **Discussion**

344   **Key nitrifiers encompassing Nitrospira and Nitrosomonas-like bacteria share ureolytic**
345   **potential**
346
16S rRNA gene sequences assembled from short reads indicated *Nitrospira*- and *Nitrosomonas*-

like populations were the only nitrifiers present in the microcosms. The proportion of 16S rRNA

gene reads mapping to *Nitrospira*-like populations in this study suggested that they were highly

abundant in the inocula and nitrogen amendments. Surveys of other DWTP biofilters using 16S

rRNA gene amplicons have indicated that sublineage II *Nitrospira* account for a dominant portion

of the bacterial community (41) with further investigation confirming high contributions to its

abundance were from comammox-*Nitrospira* (8, 33). The strict AOB OTU found in this study was

affiliated with oligotrophic *Nitrosomonas* cluster 6a which exhibit maximum growth rates at

ammonia concentrations similar to the ones used for high and mid nitrogen amendments (42, 43).

Despite this, the proportions of SSU reads mapping to *Nitrosomonas*-like populations in all

nitrogen amendments were consistently low. Taxonomic classification of nitrogen cycling genes

revealed metabolic potential for nitrification processes were confined to *Nitrospi*ra- and

Nitrosomonas-like populations corroborating with assembled 16S rRNA gene sequences.

Additionally, phylogeny of *amoA* sequences found in the metagenome indicated ammonia

oxidation could be mediated by both *Nitrospira* and *Nitrosomonas*. Consequently, nitrate made

available from nitrification processes would be available for nitrate reduction by other community

members or as a source of nitrogen for biomass synthesis in oligotrophic environments.

We assembled a total of nine nitrifier MAGs which included comammox-*Nitrospira* (n=4),

Nitrospira-NOB-like (n=2), and *Nitrosomonas*-like (n=3) populations. Three of the four

comammox MAGs assembled were identified as clade A2 based on phylogenetic analyses of

hydroxylamine dehydrogenase (*hao*) which has previously been shown to dominate drinking water

biofilters along with comammox clade B (7). The remaining comammox MAG assembled from

biofilter media in this study was affiliated with clade A1 based on *hao* gene phylogeny, which

while atypical for drinking water biofilters is consistent with previously published metagenome

from the Ann Arbor drinking water filters (6). Similar coexistence of clade A1 and A2 comammox

bacteria with canonical nitrifiers has been observed in tertiary rotating biological contactors

treating municipal wastewater with low ammonium concentrations (44). However, phylogenomic

placement of clade A sub-groups in this study separated the comammox MAGs into distinct

clusters associated with freshwater (Bin_13, clade A1), groundwater biofilters (Bin_49_4, clade

A2) and tap water (Bin_260 and Bin_49_2_2, clade A2). Maintenance of high functional

redundancy for the complete ammonia oxidation pathway may rely on coexisting comammox

populations avoiding direct competition through distinct physiological niches. Additionally, the

innocula were sourced from low substrate conditions which may also allow for the coexistence of

multiple comammox populations. Strict AOB MAGs obtained in this study associated with low

ammonia adapted *Nitrosomonas* cluster 6a (45) which is consistent with the inocula source being

an oligotrophic environment (i.e., DWTPs). Furthermore, close relatives of *Nitrospira*-NOB

MAGs obtained in this study originated from a tap water source where *Nitrospira*-NOB also

coexisted with strict AOB and comammox bacteria under oligotrophic conditions (9). Our

findings, consistent with previous studies, confirm the nitrifier community encompassed multiple

populations capable of single and two-step nitrification within a single system. Further, assessment

of metabolic versatility revealed initiation of nitrification through urea degradation was possible by

all three nitrifying guilds. Though ureolytic activity is a widespread trait among cultured

comammox-*Nitrospira* representatives and curated MAGs, the capability is confined to only some

Nitrospira-NOB and *Nitrosomonas* species (37, 43). Here in particular, this a would allow

Nitrospira-NOB to play a role in nitrite production in urea microcosms by crossing feeding

ammonia from urea degradation to strict AOB, a mutualistic strategy which may not be active in

ammonia amended microcosms.

395
396   **Comammox bacterial abundance increased irrespective of nitrogen source or loading but**
397   **may compete with NOB depending on nitrogen source type.**
398
We tested the impact of nitrogen source and loading rates on temporal dynamics of a mixed

nitrifying community to determine whether comammox bacteria are outcompeted at higher

concentrations and/or favored in urea amendments due to their ureolytic activity. qPCR-based

abundance tracking revealed comammox bacteria demonstrated a preferential enrichment over

strict AOB in the nitrogen amendments irrespective of electron donor source or availability.

Additionally, strict AOB abundance did not exhibit any significant difference across the nitrogen

amendment types. This is in contrast to previous work in soil microcosms where AOB abundance

increased in response to high ammonia amendments (28). However, strict AOB populations in

these soil microcosms were primarily *Nitrosospira* compared to oligotrophic *Nitrosomonas* cluster

6a which were the primary AOB in this study. Here, both comammox bacteria and strict AOB

demonstrated increased abundance in all amendments during the earlier weeks of the experiment.

Ultimately, while comammox bacteria were enriched over time our findings demonstrated this

increased abundance was not associated with a decrease in the abundance of strict AOB in any of

the nitrogen amendments. This suggests a lack of direct competition between the two comammox

and strict AOB which could be attributable to the two ammonia oxidizers occupying separate

nitrogen availability niches (19, 46). Stable abundances of strict AOB compared to enrichment of

comammox could be due to a combination of factors ranging from (1) higher abundances of

comammox bacteria in the inocula and (2) significantly higher biomass yields per mole of

ammonia oxidizers for comammox bacteria compared to AOB (19).

Clade A2 associated comammox bacterial MAGs were dominant in the inocula and over the course

of the experiment showed increased abundance in all amendments. In contrast, comammox

bacteria belonging to clade A1 were lower in abundance and did not demonstrate significant

change over time in any amended microcosm. Though physiological differences between

comammox bacteria clades/sub-clades have yet to be established, earlier studies of DWTP

biofilters have observed higher abundances of clade B (15) or alternatively both clades found at

the same DWTP but within separate rapid sand filters, where clade B was more abundant in the

secondary filters receiving lower ammonia concentrations (34). In this study, the lack of clade A1

enrichment may also indicate distinct physiological niches within clades (i.e., subclade-level niche

differentiation). Future research is necessary to develop a clearer understanding of physiological

differences between comammox bacteria at the clade/sub-clade level. Since cultivability of

comammox bacteria remains an ongoing challenge, integrating multiple 'omics techniques (i.e.,

metatranscriptomics and metaproteomics) may be an appropriate strategy for examining ammonia

utilization and the expressed metabolisms of multiple coexisting comammox bacteria populations

alongside canonical nitrifiers.

The negative association between comammox bacteria and canonical NOB observed in ammonia

amendments could be a result of nitrite limitation resulting from complete nitrification driven by

comammox bacteria. Nitrite limitation driven competition between comammox bacteria and NOB

is supported by the fact the negative associations between the groups were stronger at medium (3.5

mg-N/l) and low (1.5 mg-N/l) nitrogen availability as compared to the high ammonia amendments

(i.e., 14 mg-N/l). In contrast, there was no significant association between the abundance of

comammox bacteria and Nitrospira-NOB in the urea amended systems irrespective of nitrogen

loading. We hypothesize that this variable observations between ammonia and urea amended

systems likely emerge from the extent of metabolic coupling between AOB and NOB and the

resultant ability of comammox to outcompete NOB. Specifically, while the rate of nitrite

availability for NOB in ammonia amended systems is largely dictated by ammonia oxidation

activity of AOB it is likely that nitrite availability in urea amended systems would be dictated by

a combination of both AOB activity and indirectly by NOB. In this case, the production of nitrite

could be mediated by *Nitrospira*-NOB capable of ureolytic activity by crossing feeding ammonia

to strict AOB who in turn provide nitrite at a rate at which *Nitrospira*-NOB. This tight coupling

between AOB and NOB is supported by stronger and more significant correlation between AOB

and NOB abundance in urea amended systems as compared to ammonia amended systems. Thus,

it appears that while comammox bacteria may outcompete *Nitrospira*-NOB in systems where AOB

abundances are low and nitrite availability is largely dictated by AOB activity, this competitive

exclusion may be limited in scenarios with established AOB-NOB cross feeding via urea where

nitrite availability is governed not only by AOB's ammonia oxidation rate but also by NOB's

ureolytic activity.

Altogether, our study demonstrates that comammox bacteria will dominate over canonical

nitrifiers in communities sourced from nitrogen limited environments irrespective of electron

donor type or loading rate without directly competing with canonical AOB. Further, our study also

indicates comammox bacteria and AOB may occupy independent niches in communities sources

from low nitrogen environments. Interestingly, we see evidence of potential competitive exclusion

of NOB by comammox bacteria governed by nitrogen source dependent metabolic coupling

between AOB and NOB.

Materials and Methods

Experimental design and execution: Granular activated carbon (GAC) with coexisting AOB,

NOB, and comammox bacterial populations from biofilters at the drinking water treatment plant

(DWTP) in Ann Arbor, (AA) Michigan was used as the inoculum for this experimental work (6).

Microcosms consisted of 3 grams of GAC supplemented with 10 mL of filter influent from AA

DWTP in 40 mL pre-sterilized glass vials (DWK Life Sciences – Fisher 033395C). A total of 96

glass microcosms were prepared such that two biological replicates for each of the three nitrogen

concentrations (1.5, 3.5 and 14 mg-N/L) for the two nitrogen sources (i.e., ammonium and urea)

were harvested weekly for analyses over the period of the 8-week experiment. Ammonium was

spiked in at 0.1, 0.25 and 1 mM (in the form of ammonium chloride solution), corresponding to

1.5, 3.5 and 14 mg-N/L. For urea, 0.05, 0.125 and 0.5 mM (in the form of urea solution) were used

to ensure similar concentrations of total nitrogen as the ammonium microcosms. Microcosms were

maintained by removing approximately 10 mL of spent filter influent and subsequently

replenishing them with 10 mL of fresh influent and the respective electron donor spike every two

days. Once a week, two microcosms per condition (i.e., nitrogen concentration and nitrogen

source) were sacrificed and two 0.5 g GAC samples from each microcosm were transferred to

Lysing Matrix E tubes (MP Biomedical Lysing Matrix E – Fisher MP116914100) and stored at -

80°C until further processing. Spent filter influent was filtered using 0.2 μM filtration (Sartorius

Minisart NML Syringe Filter – Fisher Scientific 14555269) for subsequent chemical analyses.

Hach Company Test n' Tube Vials were used to determine concentrations of ammonia-N (Hach,

Cat No. 2606945), nitrite-N (Hach, Cat No. 2608345) and nitrate-N (Hach, Cat No. 2605345) in

microcosms. All samples were analyzed on a Hach DR1900 photospectrometer (Hach – DR1900-

01H). Alkalinity of filtered liquid samples were measured using Hach Alkalinity Total TNTplus

Vials (Hach – TNT870).

DNA extraction and qPCR: GAC samples were subjected to DNA extraction using the DNAeasy

PowerSoil kit (Qiagen, Inc – Cat No.12888) on the QIAcube (Qiagen, Inc – Cat No. 9002160)

following manufacturer's instructions with a few modifications. Specifically, the lysing buffer

from the PowerBead tubes were transferred to the Lysing Matrix E tubes and C1 buffer was added.

Prior to bead beating, an equal volume of chloroform was added (610 μL). Bead beating consisted

of four rounds of 40 seconds on a FastPrep-24 instrument (MP Bio – 116005500) with bead

beading tubes placed on ice for two minutes between each bead beating. Samples were then

centrifuged at 10,000 g for 1 minute and 750 μL of aqueous phase used to purify DNA using the

QIAcube Protocol for the DNeasy PowerSoil Kit. Each round of extractions included a reagent

blank as a negative control. After extraction, DNA concentration was determined using a Qubit

instrument with the dsDNA Broad Range Assay (ThermoFisher Scientific – Cat No. Q32850)

(Table S3). DNA was stored in a -80°C freezer until future use.

503
504   qPCR assays were conducted using QuantStudio 3 Real-Time PCR System (ThermoFisher

505   Scientific – Cat. No. A28567). Primer sets targeting the 16S rRNA gene of AOB (47), 16S rRNA

506   gene of *Nitrospira* (48), *amoB* gene of clade A comammox bacteria (18) and 16S rRNA gene for

507 total bacteria (49) were used (Table S4). Previously published primer set for the *amoB* gene of

508 clade A comammox bacteria was updated based on metagenomic data generated as part of this

509 study (18). Based on alignments of *amoB* gene sequences from the comammox MAGs assembled

510 in this study, the previously published forward primer for comammox clade A *amoB* from Cotto

511 et. al 2020 had one mismatch with one of our bins. Thus, this forward primer was further modified

512 by changing the 13[th] position from G to a degenerate base S (seven base pairs from 3′-end). The

513 use of the modified primers resulted in increased abundance of comammox bacteria in this study

514 as shown in supplementary Figure S5, indicating the ability to capture comammox *amoB* gene

515 sequences not amplified by previous primer set.

517 The qPCR reactions were carried out in 20 μL volumes, which included 10 μL Luna Universal

518 qPCR Master Mix (New England Biolabs, Inc., Cat. No. NC1276266), 5 μL of 10-fold diluted

519 template DNA, primer concentrations are outlined in Table S4 and DNAse/RNAse free water

520 (Fisher Scientific, Cat. No. 10977015) to make up the remaining volume to 20 μL. Each sample

521 per assay was subject to qPCR in triplicate and qPCR plates were prepared using the epMotion

522 M5073 liquid handling system (Eppendorf, Cat. No. 5073000205D). The cycling conditions used

523 in this study were as follows: initial denaturing at 95°C for 1 minute, 40 cycles of denaturing at

524 95°C for 15 seconds, annealing temperatures and time used are listed in Table S4 and extension at

525 72°C for 1 minute. qPCR analysis proceeded with a negative control and 7-point standard curve

526 ranging from 10^3-10^9 copies of 16S rRNA gene of *Nitrosomonas europaea* for total bacteria

527 quantification, 10^2-10^8 copies of 16S rRNA genes of *Nitrosomonas europaea* and *Ca* Nitrospira

528 inopinata for AOB and *Nitrospira* quantification, respectively, and $10^2 – 10^8$ copies of *amoB* gene

529 of *Ca* Nitrospira inopinata for the quantification of comammox bacteria. The primer used to detect

the 16S rRNA gene of *Nitrospira* would inclusively track both comammox-*Nitrospira* and

Nitrospira-NOB. Thus, *Nitrospira*-NOB abundance was estimated by subtracting the copy number

of comammox bacteria *amoB* from the copy number of 16S rRNA gene of *Nitrospira*.

<u>Metagenomic analyses</u>:

A subset of samples were selected for metagenomic analysis including DNA extracted from the

initial GAC inoculum and samples from weeks four and eight (n=13) for all nitrogen dosing and

sources. DNA extracts from duplicated microcosms for each time point were pooled by in equal

mass proportion before sending DNA templates for sequencing at the Roy J. Carver Biotechnology

Center at University of Illinois Urbana-Champaign Sequencing Core. Two lanes of Illumina

NovaSeq were used to generate paired-end reads ranging from 29 to 68 million per sample (2x150-

bp read length) (Table S5). Raw paired-end reads were trimmed and quality filtered with fastp (50)

(Table S5). Filtered reads were mapped to the UniVec Database (National Center for

Biotechnology Information) using BWA (51) to remove potential vector contamination.

Subsequent unmapped reads were extracted, sorted and indexed using SAMtools v1.3.1 (52) then

converted back to FASTQ using bedtools v2.19.1 (53).

Small subunit rRNA sequence reconstruction from quality filtered short reads was carried out

using the Phyloflash pipeline v3.4 (54). Briefly, bbmap was used to map short reads against the

SILVA 138.1 NR99 database with the default minimum identity of 70% followed by assembly of

full-length sequences with Spades (kmers = 99,111,127) and detection of closest-matching

database sequences using usearch global within VSEARCH at a minimum identity of 70%. For

read pairs, taxonomic classification was performed by taking the lowest common ancestor of the

taxonomic strings of database hits using SILVA taxonomy (55). Assembled sequences from all

samples belonging to nitrifying bacteria were clustered at 99% identity using vsearch v2.15.2 (56).

Reference *Nitrospira* and *Nitrosomonadaceae* 16S rRNA reference sequences were obtained from

ARB-SILVA and aligned with assembled sequences using muscle v3.8.1551 (57). Construction of

16S rRNA phylogenetic trees for *Nitrospira* and *Nitrosomonadaceae* was performed using IQ-

TREE v1.6.12 (58) with model finder option (59) selecting TIM3+F+I+G4 and TPM2u+F+I+G4

as models for respective trees.

Quality filtered paired-end reads from all samples were co-assembled with metaSPAdes v3.11.1

(60) with k-mers lengths 21, 33, 55, 77, 99, and 119, and phred off-set of 33. Quality evaluation

of the assembled scaffolds was performed using Quast v5.0.2 (61) (Table S6). Open reading frames

(ORF) on scaffolds were predicted using Prodigal v2.6.2 (62) with the "meta" flag and functional

prediction of resulting protein sequences were determined by similarity searches of the KEGG

database (63) using kofamscan (64). Taxonomic classification of scaffolds harboring nitrogen

cycling genes was performed using kaiju v1.7.4 (65) against the NCBI nr database with default

parameters. CoverM v0.5.0 (www.github.com/wwood/CoverM) was used to calculate reads per

kilobase million (RPKM) of these scaffolds as a metric for estimating relative abundance in each

sample.

Scaffolds were binned into clusters and manually refined using Anvi'o (v5.1 and 5.5) (66) with

three binning algorithms including CONCOCT (67), Metabat2 v2.5 (68) and Maxbin2 v2.2.7 (69).

DAS_tool v1.1.2 (70) was used to merge bins from the three approaches to generate final

metagenome assembled genomes (MAGs). Completeness and contamination of the final set was

determined using CheckM v1.0.7 (71) followed by taxonomic classification using the Genome

Taxonomy Database Toolkit v1.2.0 with release 89 v04-RS89 (72). CoverM was used to calculate

RPKM for each bin. Similar to the annotation of the metagenome, functional prediction of bin

ORFs were determined by similarity searches against the KEGG database using kofamscan. The

annotation of genes of interest were further confirmed by querying protein sequences against the

NCBI-nr database using BLASTP. MAGs were also annotated using Prokka as a secondary

annotation method (73). The Up-to-date Bacterial Core Gene pipeline (UBCG) (74) with default

parameters was used to extract and align a set of 92 singe copy core genes from *Nitrospira* and

Nitrosomonas references genomes (Table S7) and nitrifier MAGs for phylogenomic tree

reconstruction. Maximum likelihood trees were generated based on the nucleotide alignment using

IQ-TREE with model finder selecting the GTR+F+R10 and GTR+F+R4 models for *Nitrospira*

and *Nitrosomonas* trees, respectively, with 1000 bootstrap iterations. For outgroups, two

Leptospirillum and three *Nitrosospira* genomes were used for *Nitrospira* and *Nitrosomonas* trees,

respectively. Pairwise alignments of comammox *amoA* and *hao* and Nitrospira *nxrA* protein

sequences were created using muscle. Maximum likelihood trees were inferred by IQ-TREE with

model finder selecting LG+G4 for the *amoA* tree and LG+I+G4 for *hao* and *nxrA* trees with 1000

bootstrap iterations for each tree. The *amoA* and *hao* protein sequences from *Nitrosomonas*

europaea and *Nitrosomonas oligotropha* were used as the outgroup for comammox trees. All trees

were visualized using the Interactive Tree of Life (itol) (75). Pairwise comparisons of average

nucleotide identity of 38 *Nitrospira* and 15 *Nitrosomonadaceae* genomes (Table S4) with nitrifier

MAGs obtained in this study was determined using FastANI v1.31 (76).

Statistical analysis

The relative abundance of each nitrifier population was tested to determine if significant

differences existed between concentration or source of electron donor types using ANOVA and

Welch t-tests, respectively, with R version 4.0.4. Shapiro Wilks tests were used to confirm

normality prior to these statistical tests. Linear regression and correlation analysis were used to

examine the relationship between the abundance of nitrifying guilds in each of the nitrogen

amendments over time.

Data availability

Funding sources

This research was supported by NSF Graduate Research Fellowship and Cochrane Fellowship to

KV and by NSF Award number: 1703089.

633
634 References
635
1. Kowalchuk GA, Stephen JR. 2001. Ammonia-Oxidizing Bacteria: A Model for
Molecular Microbial Ecology. Annu Rev Microbiol 55:485-529.
2. Stahl DA, de la Torre JR. 2012. Physiology and diversity of ammonia-oxidizing archaea.
Annu Rev Microbiol 66:83-101.
3. Daims H, Lucker S, Wagner M. 2016. A New Perspective on Microbes Formerly Known
as Nitrite-Oxidizing Bacteria. Trends Microbiol 24:699-712.
4. Daims H, Lebedeva EV, Pjevac P, Han P, Herbold C, Albertsen M, Jehmlich N,
Palatinszky M, Vierheilig J, Bulaev A, Kirkegaard RH, von Bergen M, Rattei T,
Bendinger B, Nielsen PH, Wagner M. 2015. Complete nitrification by Nitrospira
bacteria. Nature 528:504-9.
5. van Kessel MA, Speth DR, Albertsen M, Nielsen PH, Op den Camp HJ, Kartal B, Jetten
MS, Lucker S. 2015. Complete nitrification by a single microorganism. Nature 528:555-
9.
6. Pinto AJ, Marcus DN, Ijaz UZ, Bautista-de Lose Santos QM, Dick GJ, Raskin L. 2016.
Metagenomic Evidence for the Presence of Comammox Nitrospira-Like Bacteria in a
Drinking Water System. mSphere 1.
7. Palomo A, Dechesne A, Smets BF. 2019. Genomic profiling of Nitrospira species reveals
ecological success of comammox Nitrospira. bioRxiv doi:10.1101/612226.
8. Palomo A, Jane Fowler S, Gulay A, Rasmussen S, Sicheritz-Ponten T, Smets BF. 2016.
Metagenomic analysis of rapid gravity sand filter microbial communities suggests novel
physiology of Nitrospira spp. ISME J 10:2569-2581.
9. Wang Y, Ma L, Mao Y, Jiang X, Xia Y, Yu K, Li B, Zhang T. 2017. Comammox in
drinking water systems. Water Res 116:332-341.
10. Annavajhala MK, Kapoor V, Santo-Domingo J, Chandran K. 2018. Comammox
Functionality Identified in Diverse Engineered Biological Wastewater Treatment
Systems. Environ Sci Technol Lett 5:110-116.
11. Poghosyan L, Koch H, Lavy A, Frank J, van Kessel M, Jetten MSM, Banfield JF, Lucker
S. 2019. Metagenomic recovery of two distinct comammox Nitrospira from the terrestrial
subsurface. Environ Microbiol 21:3627-3637.
12. Camejo PY, Santo Domingo J, McMahon KD, Noguera DR. 2017. Genome-Enabled
Insights into the Ecophysiology of the Comammox Bacterium "Candidatus Nitrospira
nitrosa". mSystems 2.
13. Bartelme RP, McLellan SL, Newton RJ. 2017. Freshwater Recirculating Aquaculture
System Operations Drive Biofilter Bacterial Community Shifts around a Stable Nitrifying
Consortium of Ammonia-Oxidizing Archaea and Comammox Nitrospira. Front
Microbiol 8:101.
14. Pjevac P, Schauberger C, Poghosyan L, Herbold CW, van Kessel M, Daebeler A,
Steinberger M, Jetten MSM, Lucker S, Wagner M, Daims H. 2017. AmoA-Targeted
Polymerase Chain Reaction Primers for the Specific Detection and Quantification of
Comammox Nitrospira in the Environment. Front Microbiol 8:1508.
15. Fowler SJ, Palomo A, Dechesne A, Mines PD, Smets BF. 2018. Comammox Nitrospira
are abundant ammonia oxidizers in diverse groundwater-fed rapid sand filter
communities. Environ Microbiol 20:1002-1015.

16. Wang M, Huang G, Zhao Z, Dang C, Liu W, Zheng M. 2018. Newly designed primer
pair revealed dominant and diverse comammox amoA gene in full-scale wastewater
treatment plants. Bioresour Technol 270:580-587.
17. Beach NK, Noguera DR. 2019. Design and Assessment of Species-Level qPCR Primers
Targeting Comammox. Front Microbiol 10:36.
18. Cotto I, Dai Z, Huo L, Anderson CL, Vilardi KJ, Ijaz U, Khunjar W, Wilson C, De
Clippeleir H, Gilmore K, Bailey E, Pinto AJ. 2020. Long solids retention times and
attached growth phase favor prevalence of comammox bacteria in nitrogen removal
systems. Water Res 169:115268.
19. Kits KD, Sedlacek CJ, Lebedeva EV, Han P, Bulaev A, Pjevac P, Daebeler A, Romano
S, Albertsen M, Stein LY, Daims H, Wagner M. 2017. Kinetic analysis of a complete
nitrifier reveals an oligotrophic lifestyle. Nature 549:269-272.
20. Sakoula D, Koch H, Frank J, Jetten MSM, van Kessel M, Lucker S. 2020. Enrichment
and physiological characterization of a novel comammox Nitrospira indicates ammonium
inhibition of complete nitrification. ISME J doi:10.1038/s41396-020-00827-4.
21. Gulay A, Fowler SJ, Tatari K, Thamdrup B, Albrechtsen HJ, Al-Soud WA, Sorensen SJ,
Smets BF. 2019. DNA- and RNA-SIP Reveal Nitrospira spp. as Key Drivers of
Nitrification in Groundwater-Fed Biofilters. mBio 10.
22. Liu T, Wang Z, Wang S, Zhao Y, Wright AL, Jiang X. 2019. Responses of ammonia-
oxidizers and comammox to different long-term fertilization regimes in a subtropical
paddy soil. European Journal of Soil Biology 93.
23. Wang J, Wang J, Rhodes G, He JZ, Ge Y. 2019. Adaptive responses of comammox
Nitrospira and canonical ammonia oxidizers to long-term fertilizations: Implications for
the relative contributions of different ammonia oxidizers to soil nitrogen cycling. Sci
Total Environ 668:224-233.
24. Wang Z, Cao Y, Zhu-Barker X, Nicol GW, Wright AL, Jia Z, Jiang X. 2019.
Comammox Nitrospira clade B contributes to nitrification in soil. Soil Biology and
Biochemistry 135:392-395.
25. Zheng M, Wang M, Zhao Z, Zhou N, He S, Liu S, Wang J, Wang X. 2019.
Transcriptional activity and diversity of comammox bacteria as a previously overlooked
ammonia oxidizing prokaryote in full-scale wastewater treatment plants. Sci Total
Environ 656:717-722.
26. Gottshall EY, Bryson SJ, Cogert KI, Landreau M, Sedlacek CJ, Stahl DA, Daims H,
Winkler M. 2020. Sustained nitrogen loss in a symbiotic association of Comammox
Nitrospira and Anammox bacteria. bioRxiv doi:10.1101/2020.10.12.336248.
27. Wang X, Wang S, Jiang Y, Zhou J, Han C, Zhu G. 2020. Comammox bacterial
abundance, activity, and contribution in agricultural rhizosphere soils. Sci Total Environ
727:138563.
28. He S, Li Y, Mu H, Zhao Z, Wang J, Liu S, Sun Z, Zheng M. 2021. Ammonium
concentration determines differential growth of comammox and canonical ammonia-
oxidizing prokaryotes in soil microcosms. Applied Soil Ecology 157.
29. Shao YH, Wu JH. 2021. Comammox Nitrospira Species Dominate in an Efficient Partial
Nitrification-Anammox Bioreactor for Treating Ammonium at Low Loadings. Environ
Sci Technol 55:2087-2098.

30.   Yang Y, Daims H, Liu Y, Herbold CW, Pjevac P, Lin J-G, Li M, Gu J-D. 2020. Activity
724   and Metabolic Versatility of Complete Ammonia Oxidizers in Full-Scale Wastewater
725   Treatment Systems. mBio 11:e03175-19.
726   31.   Gonzalez-Martinez A, Rodriguez-Sanchez A, van Loosdrecht MCM, Gonzalez-Lopez J,
727   Vahala R. 2016. Detection of comammox bacteria in full-scale wastewater treatment
728   bioreactors using tag-454-pyrosequencing. Environ Sci Pollut Res Int 23:25501-25511.
729   32.   Roots P, Wang Y, Rosenthal AF, Griffin JS, Sabba F, Petrovich M, Yang F, Kozak JA,
730   Zhang H, Wells GF. 2019. Comammox Nitrospira are the dominant ammonia oxidizers
731   in a mainstream low dissolved oxygen nitrification reactor. Water Res 157:396-405.
732   33.   Tatari K, Musovic S, Gulay A, Dechesne A, Albrechtsen HJ, Smets BF. 2017. Density
733   and distribution of nitrifying guilds in rapid sand filters for drinking water production:
734   Dominance of Nitrospira spp. Water Res 127:239-248.
735   34.   Poghosyan L, Koch H, Frank J, van Kessel M, Cremers G, van Alen T, Jetten MSM, Op
736   den Camp HJM, Lucker S. 2020. Metagenomic profiling of ammonia- and methane-
737   oxidizing microorganisms in two sequential rapid sand filters. Water Res 185:116288.
738   35.   Prosser JI, Nicol GW. 2012. Archaeal and bacterial ammonia-oxidisers in soil: the quest
739   for niche specialisation and differentiation. Trends Microbiol 20:523-31.
740   36.   Shi X, Hu H-W, Wang J, He J-Z, Zheng C, Wan X, Huang Z. 2018. Niche separation of
741   comammox Nitrospira and canonical ammonia oxidizers in an acidic subtropical forest
742   soil under long-term nitrogen deposition. Soil Biology and Biochemistry 126:114-122.
743   37.   Koch H, Lucker S, Albertsen M, Kitzinger K, Herbold C, Spieck E, Nielsen PH, Wagner
744   M, Daims H. 2015. Expanded metabolic versatility of ubiquitous nitrite-oxidizing
745   bacteria from the genus Nitrospira. Proc Natl Acad Sci U S A 112:11371-6.
746   38.   Palomo A, Pedersen AG, Fowler SJ, Dechesne A, Sicheritz-Ponten T, Smets BF. 2018.
747   Comparative genomics sheds light on niche differentiation and the evolutionary history
748   of comammox Nitrospira. ISME J 12:1779-1793.
749   39.   Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-Smith M, Doud D, Reddy TBK,
750   Schulz F, Jarett J, Rivers AR, Eloe-Fadrosh EA, Tringe SG, Ivanova NN, Copeland A,
751   Clum A, Becraft ED, Malmstrom RR, Birren B, Podar M, Bork P, Weinstock GM,
752   Garrity GM, Dodsworth JA, Yooseph S, Sutton G, Glockner FO, Gilbert JA, Nelson WC,
753   Hallam SJ, Jungbluth SP, Ettema TJG, Tighe S, Konstantinidis KT, Liu WT, Baker BJ,
754   Rattei T, Eisen JA, Hedlund B, McMahon KD, Fierer N, Knight R, Finn R, Cochrane G,
755   Karsch-Mizrachi I, Tyson GW, Rinke C, Genome Standards C, Lapidus A, Meyer F,
756   Yilmaz P, Parks DH, et al. 2017. Minimum information about a single amplified genome
757   (MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and archaea. Nat
758   Biotechnol 35:725-731.
759   40.   Parks DH, Rinke C, Chuvochina M, Chaumeil PA, Woodcroft BJ, Evans PN, Hugenholtz
760   P, Tyson GW. 2017. Recovery of nearly 8,000 metagenome-assembled genomes
761   substantially expands the tree of life. Nat Microbiol 2:1533-1542.
762   41.   Gulay A, Musovic S, Albrechtsen HJ, Al-Soud WA, Sorensen SJ, Smets BF. 2016.
763   Ecological patterns, diversity and core taxa of microbial communities in groundwater-fed
764   rapid gravity filters. ISME J 10:2209-22.
765   42.   Bollmann A, French E, Laanbroek HJ. 2011. Isolation, cultivation, and characterization
766   of ammonia-oxidizing bacteria and archaea adapted to low ammonium concentrations.
767   Methods Enzymol 486:55-88.

43. Sedlacek CJ, McGowan B, Suwa Y, Sayavedra-Soto L, Laanbroek HJ, Stein LY, Norton
JM, Klotz MG, Bollmann A. 2019. A Physiological and Genomic Comparison of
Nitrosomonas Cluster 6a and 7 Ammonia-Oxidizing Bacteria. Microb Ecol 78:985-994.
44. Spasov E, Tsuji JM, Hug LA, Doxey AC, Sauder LA, Parker WJ, Neufeld JD. 2020.
High functional diversity among Nitrospira populations that dominate rotating biological
contactor microbial communities in a municipal wastewater treatment plant. ISME J
14:1857-1872.
45. Koops H-P, Purkhold U, Pommerening-Röser A, Timmermann G, Wagner M. 2006. The
Lithoautotrophic Ammonia-Oxidizing Bacteria, p 778-811, The Prokaryotes
doi:10.1007/0-387-30745-1_36.
46. Martens-Habbena W, Berube PM, Urakawa H, de la Torre JR, Stahl DA. 2009. Ammonia
oxidation kinetics determine niche separation of nitrifying Archaea and Bacteria. Nature
461:976-9.
47. Hermansson A, Lindgren PE. 2001. Quantification of ammonia-oxidizing bacteria in
arable soil by real-time PCR. Appl Environ Microbiol 67:972-6.
48. Graham DW, Knapp CW, Van Vleck ES, Bloor K, Lane TB, Graham CE. 2007.
Experimental demonstration of chaotic instability in biological nitrification. ISME J
1:385-93.
49. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ,
Fierer N, Knight R. 2011. Global patterns of 16S rRNA diversity at a depth of millions of
sequences per sample. Proc Natl Acad Sci U S A 108 Suppl 1:4516-22.
50. Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics 34:i884-i890.
51. Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics 25:1754-60.
52. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
Durbin R, Genome Project Data Processing S. 2009. The Sequence Alignment/Map
format and SAMtools. Bioinformatics 25:2078-9.
53. Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics 26:841-2.
54. Gruber-Vodicka HR, Seah BKB, Pruesse E. 2020. phyloFlash: Rapid Small-Subunit
rRNA Profiling and Targeted Assembly from Metagenomes. mSystems 5.
55. Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J, Glockner FO. 2007.
SILVA: a comprehensive online resource for quality checked and aligned ribosomal
RNA sequence data compatible with ARB. Nucleic Acids Res 35:7188-96.
56. Rognes T, Flouri T, Nichols B, Quince C, Mahe F. 2016. VSEARCH: a versatile open
source tool for metagenomics. PeerJ 4:e2584.
57. Edgar RC. 2004. MUSCLE: a multiple sequence alignment method with reduced time
and space complexity. BMC Bioinformatics 5:113.
58. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol
Evol 32:268-74.
59. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017.
ModelFinder: fast model selection for accurate phylogenetic estimates. Nat Methods
14:587-589.

60.   Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. metaSPAdes: a new versatile
814   metagenomic assembler. Genome Res 27:824-834.

815   61.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for
816   genome assemblies. Bioinformatics 29:1072-5.

817   62.   Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal:
818   prokaryotic gene recognition and translation initiation site identification. BMC
819   Bioinformatics 11:119.

820   63.   Hiroyuki Ogata SG, Kazushige Sato, Wataru Fujibuchi, Hidemasa Bono and Minoru
821   Kanehisa. 1999. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic Acids
822   Research 27.

823   64.   Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, Ogata H. 2020.
824   KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score
825   threshold. Bioinformatics 36:2251-2252.

826   65.   Menzel P, Ng KL, Krogh A. 2016. Fast and sensitive taxonomic classification for
827   metagenomics with Kaiju. Nat Commun 7:11257.

828   66.   Eren AM, Esen OC, Quince C, Vineis JH, Morrison HG, Sogin ML, Delmont TO. 2015.
829   Anvi'o: an advanced analysis and visualization platform for 'omics data. PeerJ 3:e1319.

830   67.   Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, Ijaz UZ, Lahti L, Loman
831   NJ, Andersson AF, Quince C. 2014. Binning metagenomic contigs by coverage and
832   composition. Nat Methods 11:1144-6.

833   68.   Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, Wang Z. 2019. MetaBAT 2: an
834   adaptive binning algorithm for robust and efficient genome reconstruction from
835   metagenome assemblies. PeerJ 7:e7359.

836   69.   Wu YW, Simmons BA, Singer SW. 2016. MaxBin 2.0: an automated binning algorithm
837   to recover genomes from multiple metagenomic datasets. Bioinformatics 32:605-7.

838   70.   Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, Banfield JF. 2018.
839   Recovery of genomes from metagenomes via a dereplication, aggregation and scoring
840   strategy. Nat Microbiol 3:836-843.

841   71.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM:
842   assessing the quality of microbial genomes recovered from isolates, single cells, and
843   metagenomes. Genome Res 25:1043-55.

844   72.   Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. 2019. GTDB-Tk: a toolkit to classify
845   genomes with the Genome Taxonomy Database. Bioinformatics
846   doi:10.1093/bioinformatics/btz848.

847   73.   Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics
848   30:2068-9.

849   74.   Na SI, Kim YO, Yoon SH, Ha SM, Baek I, Chun J. 2018. UBCG: Up-to-date bacterial
850   core gene set and pipeline for phylogenomic tree reconstruction. J Microbiol 56:280-285.

851   75.   Letunic I, Bork P. 2019. Interactive Tree Of Life (iTOL) v4: recent updates and new
852   developments. Nucleic Acids Res 47:W256-W259.

853   76.   Jain C, Rodriguez RL, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High throughput
854   ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat Commun
855   9:5114.

859
List of Figures.

Figure 1: A) Proportion of SSU reads mapping to the SILVA SSU NR99 database (version 138.1)

from the inocula and treated samples at weeks four and eight. The community primarily consisted

of Gammaproteobacteria, Alphaproteobacteria and Nitrospirota. Proteobacteria is broken down

by classes, Alphaproteobacteria and Gammaproteobacteria, though a small portion of

Proteobacteria reads could not be classified further. B) Taxonomic classification of genes for

nitrogen biotransformation present in the metagenome at the phyla level with Proteobacteria

presented by classes, Alphaproteobacteria and Gammaproteobacteria. C and D) Phylogenetic

placement of amoA-pmoA like sequences (C), and nxrA sequences (D) detected in the

metagenomes. Both maximum likelihood trees were constructed based alignments of protein

sequences of the respective genes. Sequences identified in this study are colored according to their

phylogenetic placement (red, green, blue and orange) while references are black. AOB = ammonia

oxidizing bacteria, MOB = methane oxidizing bacteria.

Figure 2: Phylogenomic tree for Nitrospira MAGs (blue and green) obtained in this study and 32

reference genomes (black). Blue label = comammox, green label = NOB. Branch colors represent

different Nitrospira sublineages. Two Leptospirillum reference genomes were used as the outgroup

for maximum likelihood tree construction. B) Maximum likelihood tree based on ammonia

monooxygenase subunit A (amoA) sequences from comammox-Nitrospira. C) Maximum

likelihood tree based on hydroxylamine oxidoreductase (hao) sequences of comammox-Nitrospira.

For B and C blue labels represent amoA/hao gene sequences found in comammox MAGs from

this study, while black labels are reference sequences. Environment of origin is denoted with

colored squares to the left of each tree. amoA and hao protein sequences from Nitrosomonas

europaea and Nitrosomonas oligotropha were used as the outgroup for comammox trees in B and

C, respectively. D) Phylogenomic tree for strict AOB MAGs (yellow) obtained in this study and

10 Nitrosomonas reference genomes (black). Three Nitrosospira genomes were used as the

outgroup.

Figure 3: A) The relative abundance of comammox bacteria (blue) and strict AOB (red) as a

proportion of all ammonia oxidizing microorganisms calculated using copy number of 16S rRNA

and amoB genes of strict AOB and comammox bacteria, respectively and dividing by the

combined copy number to represent total AOM for each time point. Data points are the average

ratio between duplicate samples. B) Relative abundance of comammox bacteria (blue), Nitrospira-

NOB (orange) and strict AOB (red) as a proportion of total bacteria using the ratio of copy number

of the respective nitrifier genes to copy number of total bacteria averaged between duplicate

samples. Initial abundances of comammox bacteria, Nitrospira-NOB, and strict AOB were 1.2, 1.7

and 0.9% of total bacteria, respectively. Panels display relative abundances of the nitrifiers subject

to varying nitrogen amendments. C) Reads per kilobase million (RPKM) calculated for all MAGs

identifying as comammox bacteria (blue), Nitrospira-NOB (orange) and strict AOB (red) at

selected time points.

Figure 4: A) Significant positive correlation between changes in AOB concentration and that of

Nitrospira-NOB as a proportion of total bacteria were found in most treatments expect low

ammonia (1.5A). B) Negative associations between changes in comammox bacteria concentration

and that of Nitrospira-NOB as a proportion of total bacteria existed in ammonia amendments with

statistically significance detected in 3.5A and 1.5A while no association existed in urea

amendments.