1 **Comammox *Nitrospira* bacteria outnumber canonical nitrifiers irrespective of nitrogen**
2 **source and availability.**
3

4 Katherine J. Vilardi[1], Irmarie Cotto[1], Maria Sevillano Rivera[1], Zihan Dai[2,3], Christopher L.
5 Anderson[1], Ameet Pinto[4]
6
[1] Department of Civil and Environmental Engineering, Northeastern University, Massachusetts,
MA, USA
[2] Key Laboratory of Drinking Water Science and Technology, Research Center for Eco-
Environmental Sciences, Chinese Academy of Sciences, Beijing, China
[3] University of Chinese Academy of Sciences, Beijing, China
[4] School of Civil and Environmental Engineering, Georgia Institute of Technology, Atlanta, GA,
USA.
14
15 Corresponding author email: apinto36@gatech.edu

17 Keywords: comammox bacteria, nitrification, drinking water biofiltration, electron donor

<u>Abstract</u>

Complete ammonia oxidizing bacteria coexist with canonical ammonia and nitrite oxidizing bacteria in a wide range of environments. Whether this is due to competitive or cooperative interactions, or a result of niche separation is not yet clear. Understanding the factors driving coexistence of nitrifiers is critical to manage nitrification processes occurring in engineered and natural ecosystems. In this study, microcosm-based experiments were used to investigate the impact of nitrogen source and loading on the population dynamics of nitrifiers in drinking water biofilter media. Shotgun sequencing of DNA followed by co-assembly and reconstruction of metagenome assembled genomes revealed clade A2 comammox bacteria were likely the primary nitrifiers within microcosms and increased in abundance over *Nitrsomonas-like* ammonia and *Nitrospira-like* nitrite oxidizing bacteria irrespective of nitrogen source type or loading. Changes in comammox bacterial abundance did not correlate with either ammonia or nitrite oxidizing bacterial abundance in urea amended systems where metabolic reconstruction indicated potential for cross feeding between ammonia and nitrite oxidizing bacteria. In contrast, comammox bacterial abundance demonstrated a negative correlation with nitrite oxidizers in ammonia amended systems. This suggests potentially weaker synergistic relationships between ammonia and nitrite oxidizers might enable comammox bacteria to displace nitrite oxidizers from complex nitrifying communities.

Introduction

Nitrification, the biological transformation of ammonia to nitrate via nitrite, is an important process in engineered and natural ecosystems. While nitrification mediated by ammonia oxidizing microorganisms (AOM) (Kowalchuk & Stephen, 2001, Stahl & de la Torre, 2012), including ammonia oxidizing bacteria (AOB) and archaea (AOA), and nitrite oxidizing bacteria (NOB) (Daims *et al.*, 2016) has been extensively investigated, complete ammonia oxidation (comammox) performed by comammox bacteria is understudied in large part due to its recent discovery. All known comammox bacteria belong to *Nitrospira* sub-lineage II (Daims *et al.*, 2015, van Kessel *et al.*, 2015, Pinto *et al.*, 2016) and are currently divided into two clades, A and B, with clade A further separated into sub-clades A1 and A2 (Palomo *et al.*, 2019). Due to close phylogenetic relatedness, comammox-*Nitrospira* cannot be distinguished from *Nitrospira*-NOB based on the 16S rRNA gene sequence or the marker genes for nitrite oxidation (*nxrAB*) (Daims *et al.*, 2015). Thus, characterization of comammox bacteria has been largely enabled by shotgun DNA sequencing followed by reconstruction of assembled genomes (Palomo *et al.*, 2016, Pinto *et al.*, 2016, Camejo *et al.*, 2017, Wang *et al.*, 2017, Annavajhala *et al.*, 2018, Poghosyan *et al.*, 2019) and the development of primers targeting subunits of comammox bacteria ammonia monooxygenase (*amo*) gene (Bartelme *et al.*, 2017, Pjevac *et al.*, 2017, Fowler *et al.*, 2018, Wang *et al.*, 2018, Beach & Noguera, 2019, Cotto *et al.*, 2020).

Within the engineered water cycle, clade A1 comammox bacteria have been primarily detected in wastewater treatment plants while clade A2 and B have been associated with drinking water treatment and distribution systems (Palomo *et al.*, 2019). It is unclear if this translates into physiological differences between the clades/sub-clades since there is only one comammox

isolate and an enrichment whose kinetic parametes have been reported. To date, kinetic

parameters of comammox bacteria are confined to two clade A representatives, cultured

Candidatus N. inopinata and an enrichment of *Candidatus* N. kreftii (Kits *et al.*, 2017, Sakoula *et*

al., 2020). Both demonstrate a high affinity for ammonia, with half-saturation constants orders of

magnitude lower than strict AOB. Comparatively, the *Candidatus* N. kreffti enrichment

exhibited a higher affinity for nitrite compared to *Candidatus* N. inopinata and demonstrated

partial inhibition of ammonia oxidation even at low ammonia concentrations (Sakoula *et al.*,

2020). This suggests that clade-specific comammox bacterial niche, if applicable, may be arise

from a combination of factors ranging from affinity to inhibition. Beyond clade specific traits,

identifying the potential environmental and physiological factors driving the coexistence of

comammox bacteria with canonical nitrifiers is also important to better understand comammox

bacteria role in complex nitrifying communities (Gulay *et al.*, 2019, Liu *et al.*, 2019, Wang *et al.*,

2019, Wang *et al.*, 2019, Zheng *et al.*, 2019, Gottshall *et al.*, 2020, Wang *et al.*, 2020, He *et al.*,

2021, Shao & Wu, 2021). Comammox bacteria have been detected along with their canonical

nitrifying counterparts in wastewater treatment plants (Gonzalez-Martinez *et al.*, 2016, Roots *et*

al., 2019, Zheng *et al.*, 2019, Cotto *et al.*, 2020, Yang *et al.*, 2020), drinking water systems

(Pinto *et al.*, 2016, Tatari *et al.*, 2017, Wang *et al.*, 2017, Fowler *et al.*, 2018, Poghosyan *et al.*,

2020) and soils (Prosser & Nicol, 2012, Shi *et al.*, 2018, Liu *et al.*, 2019, He *et al.*, 2021) at

varying abundances over a wide range of ammonium concentrations. While there is currently no

quantitative estimate of the contribution of comammox bacteria to nitrification compared to

AOB and NOB, several studies have investigated comammox bacterial dynamics in the context

of mixed nitrifying communities. For instance, DNA/RNA stable isotope probing provided

support for comammox *Nitrospira* contributing to ammonia oxidation in lab-scale biofilters

exposed to very low ammonium concentrations (Gulay *et al.*, 2019). Soil microcosms amended

with high ammonia concentrations were enriched in AOB compared to those with lower

ammonia concentrations where clade B comammox bacteria proliferated (Wang *et al.*, 2019, He

et al., 2021). Interestingly, in a lab-scale partial nitrification-anammox reactor operating with

incrementally increased ammonia loadings, comammox bacteria initially dominated over strict

AOB but its abundance significantly declined as loadings were further increased (Shao & Wu,

2021).

Comammox bacteria may also acquire ammonia via urea degradation. Specifically, genes

encoding for urea transport and the urease enzyme are distributed among many *Nitrospira*

populations (Koch *et al.*, 2015), including most comammox populations (Palomo *et al.*, 2018).

While this may diversify potential nitrogen sources for comammox bacteria (Daims *et al.*, 2016),

this could be a potential advantage for canonical nitrifiers involved in a reciprocal feeding

strategy as observed with co-cultured *Nitrospira moscoviensis* converting urea to ammonia for

Nitrosomonas europaea (Koch *et al.*, 2015). The tight interplay between canonical nitrifiers is

well established; however, our understanding of comammox competition (or lack thereof) with

AOM and its impact on strict NOB in mixed communities is limited.

To better understand the comammox bacterial role within these complex nitrifying communities,

we investigated their population dynamics across two nitrogen sources (ammonia or urea) at

three total nitrogen dosing strategies. Thus, the objectives of this study were (1) to determine if

comammox bacteria and canonical nitrifiers exhibit concentration and nitrogen source dependent

dynamics when subject to repeat nitrogen amendments and (2) to determine if these dynamics

are consistent or variable at the clade or population within each functional guild.

Characterization of microbial communities in biofilters at drinking water treatment plants has

revealed rich nitrifier diversity (Fowler *et al.*, 2018, Gulay *et al.*, 2019), making it an ideal

sample source for this study. Collectively, our microcosm-based study offers novel insights

regarding the ecophysiology of clade 2 associated comammox bacteria; information on this clade

are very limited. Further, while other microcosm studies have focused on competitive

interactions between comammox bacteria and strict AOB under controlled conditions (Wang *et*

al., 2019, He *et al.*, 2021), there is only limited assessment of NOB response to experimental

treatment. This study explicitly assesses the NOB dynamics in response to nitrogen source and

loading rates in the context of the broader nitrifying community.

Materials and Methods

<u>**Experimental design and execution**</u>: Granular activated carbon (GAC) with coexisting AOB,

NOB, and comammox bacterial populations from biofilters at the drinking water treatment plant

(DWTP) in Ann Arbor, (AA) Michigan was used as the inoculum for this experimental work

(Pinto *et al.*, 2016). Microcosms consisted of 3 grams of GAC supplemented with 10 mL of filter

influent from AA DWTP in 40 mL pre-sterilized glass vials (DWK Life Sciences – Fisher

033395C). A total of 96 glass microcosms were prepared such that two biological replicates for

each of the three nitrogen concentrations (1.5, 3.5 and 14 mg-N/L) for the two nitrogen sources

(i.e., ammonium (direct) and urea (indirect)) were harvested weekly for analyses over the period

of the 8-week experiment. Ammonium was spiked in at 0.1, 0.25 and 1 mM (in the form of

ammonium chloride solution), corresponding to final concentrations of 1.5, 3.5 and 14 mg-N/L.

For urea, 0.05, 0.125 and 0.5 mM (in the form of urea solution) were used to ensure similar

concentrations of total nitrogen as the ammonium microcosms. Microcosms were maintained by

removing approximately 10 mL of spent filter influent and subsequently replenishing them with

10 mL of fresh influent and the respective nitrogen source spike every two days. Once a week,

two microcosms per condition (i.e., nitrogen concentration and nitrogen source) were sacrificed

and two 0.5 g GAC samples from each microcosm were transferred to Lysing Matrix E tubes

(MP Biomedical Lysing Matrix E – Fisher MP116914100) and stored at -80°C until further

processing. Additionally, aqueous samples were collected and filtered through 0.2 μM filters

(Sartorius Minisart NML Syringe Filter – Fisher Scientific 14555269) for chemical analyses. The

sampled aqueous volume was replaced with fresh substrate to increase the ammonia and urea

concentrations to microcosm specific concentrations. Hach Company Test n' Tube Vials were

used to determine concentrations of ammonia-N (Hach, Cat No. 2606945), nitrite-N (Hach, Cat

No. 2608345) and nitrate-N (Hach, Cat No. 2605345) in microcosms. All samples were analyzed

on a Hach DR1900 photospectrometer (Hach – DR1900-01H). Alkalinity of filtered liquid

samples were measured using Hach Alkalinity Total TNTplus Vials (Hach – TNT870).

DNA extraction and qPCR: GAC samples were subjected to DNA extraction using the

DNAeasy PowerSoil kit (Qiagen, Inc – Cat No.12888) on the QIAcube (Qiagen, Inc – Cat No.

9002160) following manufacturer's instructions with a few modifications. Specifically, the

lysing buffer from the PowerBead tubes were transferred to the Lysing Matrix E tubes and C1

buffer was added. Prior to bead beating, an equal volume of chloroform was added (610 μL).

Bead beating consisted of four rounds of 40 seconds on a FastPrep-24 instrument (MP Bio –

116005500) with bead beading tubes placed on ice for two minutes between each bead beating.

Samples were then centrifuged at 10,000 g for 1 minute and 750 μL of aqueous phase used to

purify DNA using the QIAcube Protocol for the DNeasy PowerSoil Kit. Each round of

extractions included a reagent blank as a negative control. After extraction, DNA concentration

was determined using a Qubit instrument with the dsDNA Broad Range Assay (ThermoFisher

Scientific – Cat No. Q32850) (Table S1). DNA was stored in a -80$^{°}$C freezer until future use.

191
qPCR assays were conducted using QuantStudio 3 Real-Time PCR System (ThermoFisher

Scientific – Cat. No. A28567). Primer sets targeting the 16S rRNA gene of AOB (Hermansson

& Lindgren, 2001), 16S rRNA gene of *Nitrospira* (Graham *et al.*, 2007), *amoB* gene of clade A

comammox bacteria (Cotto *et al.*, 2020) and 16S rRNA gene for total bacteria (Caporaso *et al.*,

2011) were used (Table S2). Previously published primer set for the *amoB* gene of clade A

comammox bacteria was updated based on metagenomic data generated as part of this study

(Cotto *et al.*, 2020). Based on alignments of *amoB* gene sequences from the comammox MAGs

assembled in this study, the previously published forward primer for comammox clade A *amoB*

from Cotto et. al 2020 had one mismatch with one of our bins. Thus, this forward primer was

further modified by changing the 13th position from G to a degenerate base S (seven base pairs

from 3´-end). The use of the modified primers resulted in increased abundance of comammox

bacteria in this study as shown in supplementary Figure S1, indicating the ability to capture

comammox *amoB* gene sequences not amplified by previous primer set.

205
The qPCR reactions were carried out in 20 μL volumes, which included 10 μL Luna Universal

qPCR Master Mix (New England Biolabs, Inc., Cat. No. NC1276266), 5 μL of 10-fold diluted

template DNA, primer concentrations are outlined in Table S4 and DNAse/RNAse free water

(Fisher Scientific, Cat. No. 10977015) to make up the remaining volume to 20 μL. Each sample

per assay was subject to qPCR in triplicate and qPCR plates were prepared using the epMotion

M5073 liquid handling system (Eppendorf, Cat. No. 5073000205D). The cycling conditions used

in this study were as follows: initial denaturing at 95°C for 1 minute, 40 cycles of denaturing at

95°C for 15 seconds, annealing temperatures and time used are listed in Table S2 and extension

at 72°C for 1 minute. qPCR analysis proceeded with a negative control and 7-point standard

curve ranging from 10^3-10^9 copies of 16S rRNA gene of *Nitrosomonas europaea* for total

bacteria quantification, 10^2-10^8 copies of 16S rRNA genes of *Nitrosomonas europaea* and *Ca*

Nitrospira inopinata for AOB and *Nitrospira* quantification, respectively, and $10^2 – 10^8$ copies of

amoB gene of *Ca* Nitrospira inopinata for the quantification of comammox bacteria. The primer

used to detect the 16S rRNA gene of *Nitrospira* would inclusively track both comammox-

Nitrospira and *Nitrospira*-NOB. Thus, *Nitrospira*-NOB abundance was estimated by subtracting

the copy number of comammox bacteria *amoB* from the copy number of 16S rRNA gene of

Nitrospira.

Metagenomic analyses: A subset of samples were selected for metagenomic analysis including

DNA extracted from the initial GAC inoculum and samples from weeks four and eight (n=13)

for all nitrogen sources and dosing strategies. DNA extracts from duplicate microcosms for each

time point were pooled in equal mass proportion before sending DNA templates for sequencing

at the Roy J. Carver Biotechnology Center at University of Illinois Urbana-Champaign

Sequencing Core. Two lanes of Illumina NovaSeq were used to generate paired-end reads

ranging from 29 to 68 million per sample (2x150-bp read length) (Table S3). Raw paired-end

reads were trimmed and quality filtered with fastp (Chen *et al.*, 2018) (Table S5). Filtered reads

were mapped to the UniVec Database (National Center for Biotechnology Information) using

BWA (Li & Durbin, 2009) to remove potential vector contamination. Subsequent unmapped

reads were extracted, sorted and indexed using SAMtools v1.3.1 (Li *et al.*, 2009) then converted

back to FASTQ using bedtools v2.19.1 (Quinlan & Hall, 2010).

Small subunit rRNA sequence reconstruction from quality filtered short reads was carried out

using the Phyloflash v3.4 (Gruber-Vodicka *et al.*, 2020). Briefly, bbmap was used to map short

reads against the SILVA 138.1 NR99 database with the default minimum identity of 70%

followed by assembly of full-length sequences with Spades (kmers = 99,111,127) and detection

of closest-matching database sequences using usearch global within VSEARCH at a minimum

identity of 70%. For read pairs, taxonomic classification was performed by taking the lowest

common ancestor using SILVA taxonomy (Pruesse *et al.*, 2007). Assembled sequences from all

samples belonging to nitrifying bacteria were clustered at 99% identity using vsearch v2.15.2

(Rognes *et al.*, 2016). Reference *Nitrospira* and *Nitrosomonadaceae* 16S rRNA reference

sequences were obtained from ARB-SILVA and aligned with assembled sequences using muscle

v3.8.1551 (Edgar, 2004). Construction of 16S rRNA phylogenetic trees for *Nitrospira* and

Nitrosomonadaceae was performed using IQ-TREE v1.6.12 (Nguyen *et al.*, 2015) with model

finder option (Kalyaanamoorthy *et al.*, 2017) selecting TIM3+F+I+G4 and TPM2u+F+I+G4 as

models for respective trees.

Quality filtered paired-end reads from all samples were co-assembled with metaSPAdes v3.11.1

(Nurk *et al.*, 2017) with k-mers lengths 21, 33, 55, 77, 99, and 119, and phred off-set of 33.

Quality evaluation of the assembled scaffolds was performed using Quast v5.0.2 (Gurevich *et*

al., 2013) (Table S4). Open reading frames (ORF) on scaffolds were predicted using Prodigal

v2.6.2 (Hyatt *et al.*, 2010) with the "meta" flag and functional prediction of resulting protein

sequences were determined by similarity searches of the KEGG database (Hiroyuki Ogata, 1999)

using kofamscan (Aramaki *et al.*, 2020). Taxonomic classification of scaffolds harboring

nitrogen cycling genes was performed using kaiju v1.7.4 (Menzel *et al.*, 2016) against the NCBI

nr database with default parameters. CoverM v0.5.0 (www.github.com/wwood/CoverM) was

used to calculate reads per kilobase million (RPKM) of these scaffolds as a metric for estimating

relative abundance in each sample.

Scaffolds were binned into clusters and manually refined using Anvi'o (v5.1 and 5.5) (Eren *et*

al., 2015) with three binning algorithms including CONCOCT (Alneberg *et al.*, 2014), Metabat2

v2.5 (Kang *et al.*, 2019) and Maxbin2 v2.2.7 (Wu *et al.*, 2016). DAS_tool v1.1.2 (Sieber *et al.*,

2018) was used to merge bins from the three approaches to generate final metagenome

assembled genomes (MAGs). Completeness and contamination of the final set was determined

using CheckM v1.0.7 (Parks *et al.*, 2015) followed by taxonomic classification using the

Genome Taxonomy Database Toolkit v1.2.0 with release 89 v04-RS89 (Chaumeil *et al.*, 2019).

CoverM was used to calculate RPKM for each bin. Similar to the annotation of the metagenome,

functional prediction of bin ORFs were determined by similarity searches against the KEGG

database using kofamscan. The annotation of genes of interest were further confirmed by

querying protein sequences against the NCBI-nr database using BLASTP. MAGs were also

annotated using Prokka as a secondary annotation method (Seemann, 2014). The Up-to-date

Bacterial Core Gene pipeline (UBCG) (Na *et al.*, 2018) with default parameters was used to

extract and align a set of 92 singe copy core genes from *Nitrospira* and *Nitrosomonas* references

genomes (Table S5) and nitrifier MAGs for phylogenomic tree reconstruction. Maximum

likelihood trees were generated based on the nucleotide alignment using IQ-TREE with model

finder selecting the GTR+F+R10 and GTR+F+R4 models for *Nitrospira* and *Nitrosomonas* trees,

respectively, with 1000 bootstrap iterations. For outgroups, two *Leptospirillum* and three

Nitrosospira genomes were used for *Nitrospira* and *Nitrosomonas* trees, respectively. Pairwise

alignments of comammox *amoA* and *hao* and Nitrospira *nxrA* protein sequences were created

using muscle. Maximum likelihood trees were inferred by IQ-TREE with model finder selecting

LG+G4 for the *amoA* tree and LG+I+G4 for *hao* and *nxrA* trees with 1000 bootstrap iterations

for each tree. The *amoA* and *hao* protein sequences from *Nitrosomonas europaea* and

Nitrosomonas oligotropha were used as the outgroup for comammox trees. All trees were

visualized using the Interactive Tree of Life (itol) (Letunic & Bork, 2019). Pairwise

comparisons of average nucleotide identity of 38 *Nitrospira* and 15 *Nitrosomonadaceae*

genomes (Table S5) with nitrifier MAGs obtained in this study was determined using FastANI

v1.31 (Jain *et al.*, 2018).

Statistical analysis

The relative abundance of each nitrifier population was tested to determine if significant

differences existed between concentration or source of electron donor types using ANOVA and

Welch t-tests, respectively, with R version 4.0.4. Shapiro Wilks tests were used to confirm

normality prior to these statistical tests. Linear regression and correlation analysis were used to

examine the relationship between the abundance of nitrifying guilds in each of the nitrogen

amendments over time.

Results

Microbial community composition in microcosms and nitrogen biotransformation potential.

Microcosms consisting of granular activated carbon (GAC) from drinking water biofilters were subject to intermittent amendments of nitrogen using two nitrogen sources (ammonia or Urea) across three nitrogen concentrations (14, 3.5, and 1.5 mg-N/L). The conditions used in these experiments are denoted as 14A, 3.5A, 1.5A, 14U, 3.5U and 1.5U where A or U represents ammonia or urea amendments, respectively, and the number represents the concentration of nitrogen source spike in mg/L as nitrogen. Two microcosms were sacrificed on a weekly basis over the duration of a eight week experiment (n=96 total microcosms). Extracted DNA from the inocula and weeks four and eight were subject to shotgun DNA sequencing (n=13).

Initial assessment of taxonomic diversity in the samples based on analyses of metagenomic reads mapping to the small subunit rRNA database (SILVA SSU NR99 version 138.1) indicated that the GAC inocula largely consisted of bacteria with archaea and eukaryota constituting a small proportion of the overall metagenomic reads (~0.002%). The bacterial community was primarily composed of Gammaproteobacteria (20-30%), Alphaproteobacteria (25-31%) and Nitrospirota (8-15%) (Figure 1A). *Nitrospira* and *Nitrosomonadaceae* were the only nitrifiers identified and constituted 9-15% of the overall microbial community in samples. Full length 16S rRNA gene sequences were assembled from each sample (n=13) resulting in a total of eight sequences with closest matching SILVA database hits to uncultured *Nitrospira* bacteria (Accession numbers: MF040566, AY328760, JN868922). Clustering of all eight *Nitrospira* 16S rRNA gene sequences at 99% identity resulted in two *Nitrospira* operational taxonomic units (OTUs) with one cluster composed of six sequences (Nitrospira OTU 1) and the other cluster with two sequences

(Nitrospira OTU 2). Phylogenetic placements of these OTUs revealed both clustered within

Nitrospira sublineage II (supplementary figure S2A). Diversity of *Nitrospira* was likely

underrepresented as full length *Nitrospira* 16S rRNA gene sequences could not be assembled

from some samples despite a large portion of extracted 16S rRNA gene reads mapping to

Nitrospira references in the SILVA database. Limited assembly of these reads could be due to

several closely related *Nitrospira* species/strains coexisting in the samples making re-

construction of full length sequences difficult. For canonical AOB, Nitrosomonas sp. AL212

(CP002552) was the closest matching database hit to one assembled sequence while another six

had hits closet to *Nitrosomonadaceae* (Accession numbers: FPLP01009519, KJ807851,

FPLK01002446) but could not be further classified at the genus or species level. Phylogenetic

placement of the single Nitrosomonas OTU affiliated it with *Nitrosomonas* sp. AL212 and

Nitrosomonas ureae (Figure S2B).

Following co-assembly of metagenomic reads, predicted protein coding genes from scaffolds

associated with the nitrogen metabolism were taxonomically classified (Figure 1B). The majority

of methane/ammonia monooxygenase (*pmo-amo*) like genes (KEGG orthology: K10944,

K10945, K10946) were associated with either nitrifiers (i.e., *Nitrospira* or *Nitrosomonas*) or

methanotrophs (i.e., *Methylocystis*) (Figure 1C). While some *amoCAB* genes could not be

classified to the genus level using kaiju software, blastp searches against the NCBI non-

redundant protein database indicated these were closely related to *Nitrosomonas*. All retrieved

hao sequences (KEGG orthology: K10535) were associated with Nitrospira which is likely due

to the low relative abundance of *Nitrosomonas*-like populations and the resulting inability to

assemble their *hao* genes. Potential for ureolytic activity was detected across four phyla based on

the urease alpha subunit (*ureC*). *ureC* sequences associated with Nitrospirota and

Gammaproteobacteria could be classified at the genus level as *Nitrospira* and *Nitrosomonas*.

Sequences identified as nitrate reductase/nitrite oxidoreductase alpha and beta subunits (K00370,

K00371) were subject to further classification to differentiate between nitrite oxidoreductase

genes belonging to NOB from nitrate reductases belonging to other community members.

Phylogenetic placement of most *Nitrospira nxrA* sequences found in this study cluster within a

branch containing both comammox and *Nitrospira*-NOB species (Candidatus *N. inopinata*,

Candidatus *N. nitrosa* and *N. defluvii*) (Figure 1D). While other sequences clustered on a

separate branch with Candidatus *N. nitrificans*, a single *Nitrospira nxrA* sequence clustered

closely within a branch containing only *Nitrospira*-NOB belonging to sublineage II.

Phylogenomic placement of nitrifying populations and their metabolism.

Metagenome assembled genomes (MAGs) were obtained from GAC microcosms after

dereplication of MAGs from three binning approaches. All 204 MAGs were classified as

bacteria, with 145 MAGs exhibiting completeness greater than 70% and contamination less than

10% (Table S6). Approximately 62% of the metagenomic reads mapped to these MAGs. Nine

MAGs in total were classified as nitrifying bacteria belonging to *Nitrosomonas* and *Nitrospira*

(Table S7). Genome annotation confirmed that four *Nitrospira* MAGs had key ammonia

(ammonia monooxygenase and hydroxylamine oxidoreductase) and nitrite (nitrite

oxidoreductase) oxidation genes (Figure S3). Quality assessment for these comammox MAGs

indicated two high (Bin_49_2_2 and Bin_49_4) and one medium quality (Bin_260) (Table S1)

according to (Bowers *et al.*, 2017). A fourth comammox MAG (Bin_13) was assembled with

high completeness (89%) but also possessed high redundancy (18%) that could not be improved

with further manual refinement. The remaining two *Nitrospira* MAGs (Bin_7_1 and Bin_188),

which were likely strict NOB due to lack of ammonia oxidation genes, were less complete

(38.04% and 48.25%) with low redundancy (8.76% and 8.46%). The low completeness was

likely not due to their lower abundance, but potentially high level of strain heterogeneity which

may have affected the assembly of reads associated with *Nitrospira*-NOB. For example, RPKM-

based relative abundance estimated using all reads (total RPKM) showed the two *Nitrospira*-

NOB MAGs exhibited similar relative abundance to comammox bacteria MAGs Bin_49_2_2

and Bin_49_4 (~7-10 total RPKM), but the CheckM estimated strain heterogeneities for Bin_7_1

and Bin_188 were 40 and 75, respectively, compared to 0 for both Bin_49_2_2 and Bin_49_4.

Two MAGs classifying as *Nitrosomonas* were deemed high (Bin_83) and medium quality

(Bin_168); however, a third *Nitrosomonas* MAG was considered low quality.

A maximum likelihood tree based on 91 single copy core genes confirmed all *Nitrospira* MAGs

affiliated with sublineage II (Figure 2A). Four of the *Nitrospira* MAGs from this study clustered

within clade A comammox *Nitrospira* (Bin_49_2_2, Bin_49_4, Bin_260 and Bin_13) but were

separated into distinct groups on the phylogenomic tree; namely, forming three clusters with

MAGs obtained from tap water, drinking water filters, and freshwater. *amoA*-based phylogenetic

analysis corroborated their placement into clade A (Figure 2B); however, *hao*-based phylogeny

distinguished three of comammox MAGs (Bin_49_2_2, Bin_49_4, Bin_260) as clade A2

(Palomo et al. 2019) while one clustered within clade A1 (Bin_13) (Figure 2C). Consistent

across all trees, Bin_49_2_2 and Bin_260 cluster closely with comammox MAGs *Nitrospira* sp.

SG-bin2 and ST-bin4 (ANI ~ 92%) derived from tap water metagenomes (Wang *et al.*, 2017).

Bin_49_4 clustered closely with Nitrospirae bacterium Ga0074138 (ANI ~ 99%), which was

previously detected in GAC from the same drinking water treatment plant (Pinto *et al.*, 2016),

along with other tap water and groundwater-fed rapid sand filter MAGs (Palomo *et al.*, 2016,

Wang *et al.*, 2017). Bin_13 associated with comammox MAGs obtained from freshwater,

UBA5698 and UBA5702 (Parks *et al.*, 2017) (ANI ~ 90%); however, its high contamination

(18%) likely renders ANI comparison less accurate. Overall, the MAGs demonstrated less then

95% ANI to other reference comammox bacterial MAGs (Figure S4) suggesting comammox

bacteria detected in GAC microcosms are distinct from one another and previously published

comammox MAGs; as a result, they are likely novel *Nitrospira* species. The two remaining

Nitrospira MAGs, Bin_7_1 and Bin_188, clustered with strict *Nitrospira*-NOB MAGs recovered

from tap water, Nitrospira_sp_ST-bin5 (Wang *et al.*, 2017) (ANI ~ 94%), and a rapid sand filter,

Nitrospira CG24D (ANI ~ 87%) (Palomo *et al.*, 2016) (Figure 2A and S3). However, since

Bin_7_1 and Bin_188 were highly incomplete, the possibly they are novel comammox bacteria

cannot be excluded. Only two strict AOB MAGs (Bin_83 and Bin_168) from this study were

used for phylogenomic analysis due high redundancy and low completeness of the third

(Bin_195). Both Bin_83 and Bin_168 originate from *Nitrosomonas* cluster 6a and clustered

closely with *Nitrosomonas ureae* and *Nitrosomonas* sp. AL212 (Figure 2D). Bin_168 shares a

high sequence similarity to *N. ureae* (ANI ~ 98%) while Bin_83 shares less than 83% ANI to

any of the references on the tree including Bin_168.

All comammox MAGs demonstrated the potential for ureolytic activity with the presence of the

ureABC operon in addition to most genes for urease accessory proteins (Figure S2). *Nitrospira*-

NOB MAGs did not contain genes encoding for urease; however, two *ureC* sequences found on

assembled scaffolds that were classified as *Nitrospira* but were not binned into any of the

Nitrospira MAGs. Queries of these *ureC* genes against the NCBI non-redundant database

revealed one sequence shared the highest percent identity to *Nitrospira lenta* and *Nitrospira*

moscoviensis while top hits for the second sequence belonged to an unclassified *Nitrospira*. One

Nitrospira-NOB MAG (Bin_7_1) did harbor genes for the urea transport system permease

proteins (*urtBC*), urea transport system substrate-binding proteins (*urtA*) and urea transport

system ATP-binding proteins (*urtDE*). This suggests that the two unbinned *ureC* genes likely

belonged to *Nitrospira*-like NOB bacteria. *Nitrosomonas* MAGs Bin_168 and Bin_83 each

contained the *ureCAB* operon and some genes for urease accessory proteins and urea transport. A

third *ureC* sequence found in the metagenome classified as *Nitrosomonas* but was not binned

into any *Nitrosomonas* MAGs.

The impact of nitrogen amendments on nitrifying populations.

To address concentration and nitrogen source-dependent dynamics of the three nitrifier

populations detected in our metagenomic analysis, qPCR-assays were used to estimate their

abundances over time in the nitrogen amended microcosms. In the high ammonia amendment

(14A), strict AOB relative abundance increased 2.4-fold from weeks 1-3 but remained below 2%

of total bacteria for the duration of the experiment whereas comammox relative abundance

increased markedly over time reaching 2.8% of total bacteria by end of the experiment (Figure

3B). Similar to strict AOB, *Nitrospira*-NOB relative abundance increased early on but thereafter

reduced from 4% at its peak in week two to 1.8% by week eight. Weekly measurements for

nitrogen concentrations taken alongside biomass samples indicated the presence of residual

ammonia and accumulated nitrite concentrations were highest during the first three weeks of the

experiment but gradually reduced over time with most inorganic nitrogen present as nitrate

(Figure S5). While comammox bacteria were always dominant, qPCR-based abundance of strict

AOB as a portion of AOM was significantly higher when ammonia and nitrite accumulated in

weeks 1-3 as compared to weeks 5-8 (Welch's t-test, p-value < 0.05) (Figure 3A).

The qPCR data was in concordance with metagenomic estimation of MAG abundance with clade

A2 comammox (Bin_49_2_2, Bin_49_2 and Bin_260) highly abundant compared to strict AOB

(Bin_83, Bin_168 and Bin_195) and clade A1 comammox (Bin_13) in the inocula and at weeks

four and eight in the high ammonia amendment (Figure 4). In particular, clade A2 MAGs

Bin_49_2_2 and Bin_49_4 were the most dominant comammox populations while strict AOB

was dominated by Bin_83 at each time point. *Nitrospira*-NOB MAGs had comparable

abundance to clade A2 comammox MAGs but displayed limited variation in abundance in the

high ammonia amendments. This contrasts with the qPCR data, where *Nitrospira*-NOB were

significantly more abundant than comammox bacteria at earlier timepoints and then

demonstrated a significant decrease in abundance over time. This is likely due to the fact that the

two assembled Nitrospira-NOB MAG's do not represent the entirety of NOB diversity in the

microcosms as several *nxr* genes were not binned into MAGs and that metagenomic data is only

available for select timepoints as compared to qPCR data.

Nitrifier populations in mid and low ammonia amendments displayed similar dynamics to those

observed in high ammonia with comammox relative abundance increasing to 3% and 2.2% of

total bacteria by week eight, respectively. Interestingly, Bin_260, the least abundant clade A2

comammox MAG in the inocula, demonstrated significant increase in abundance in the low

ammonia amendment over the course of the experiment compared to its abundance in the other

ammonia amendments. Consistent with the ammonia amended microcosms, strict AOB in urea

amended microcosms increased in relative abundance only at earlier time points followed by low

but stable relative abundance (~2% of total bacteria). In the high urea amendment, relative

abundance of comammox bacteria remained largely unchanged at earlier time points followed by

an increase in abundance. Despite this, mean relative abundance of comammox bacteria

compared to strict AOB was still approximately 2-fold greater in all urea amendments. Similar to

the ammonia amendments, *Nitrospira*-NOB relative abundance did increase initially followed by

a decline in all urea amendments. Interestingly though, the relative abundance of comammox

bacteria and *Nitrospira*-NOB were similar in the later weeks of the experiment after *Nitrospira*-

NOB's initial rise in urea amendments. Clade A2 comammox MAG Bin_260 was consistently

lower in abundance than Bin_49_2_2 and Bin_49_2 in the urea amendments except for mid urea.

Abundance of the clade A1 comammox MAG remained lower than all clade A2 MAGs and

displayed minimal enrichment in all the urea amendments which was consistent with ammonia

amended microcosms. Bin_168, which showed high sequence similarity to *Nitrosomonas ureae*,

did not exhibit enrichment in any of the urea amendments and remained low in abundance with

all other strict AOB MAGs.

There was no significant difference in the mean qPCR-based relative abundance of strict AOB or

Nitrospira-NOB between the high ammonia (14A) and urea amendments (14U) (Welch t-test, p

> 0.05) but the mean relative abundance of comammox bacteria was significantly greater in high

ammonia than in the high urea amendment (Welch t-test, $p < 0.05$). Comparatively, out of all

nitrogen amendments, mean relative abundance of comammox bacteria was the lowest in high

urea (1.8% of total bacteria). Comparisons between the mid ammonia (3.5A) and urea

amendments (3.5U) as well as the low ammonia (1.5A) and urea (1.5U) amendments revealed no

significant difference in mean relative abundance for any of the nitrifier populations (Welch t-

test, $p > 0.05$). Additionally, no significant differences were detected when testing the mean

relative abundance of the three nitrifier populations between high, mid, and low concentrations

within each amendment type (ANOVA, $p > 0.05$).

The relative abundance of the nitrifying groups were used to examine potential correlations

between the different populations in each of the nitrogen amendments. The ratio of comammox

bacteria as portion of AOM to comammox bacteria as a portion of total *Nitrospira* revealed a

strong positive relationship in all amendments (Pearson R = 0.75-0.87, $p < 0.001$) (Figure S6A),

however, the change in relative abundance of comammox bacteria was not directly correlated

with that of strict AOB in any of the nitrogen amendments (Figure S6B). Strict AOB and

Nitrospira-NOB abundances were strongly correlated for all urea amendments and high (14A)

and mid ammonia (3.5A) (Pearson r = 0.58-0.82, $p < 0.05$, Figure 5A) but exhibited a weaker

relationship in low ammonia (Pearson r = 0.42, $p > 0.05$). Interestingly, while comammox

bacteria abundance was significantly and negatively correlated with that of *Nitrospira*-NOB in

ammonia amendments (Pearson r = -0.37 to -0.61) ($p < 0.05$), there was no significant

association between them in the urea amendments ($p > 0.05$) (Figure 5B).

Discussion

Key nitrifiers encompassing Nitrospira and Nitrosomonas-like bacteria share ureolytic

potential.

16S rRNA gene sequences assembled from short reads indicated *Nitrospira-* and *Nitrosomonas-*

like populations were the only nitrifiers present in the microcosms. The proportion of 16S rRNA

gene reads mapping to *Nitrospira*-like populations in this study suggested that they were highly

abundant in the inocula and nitrogen amendments. Surveys of other DWTP biofilters using 16S

rRNA gene amplicon sequencing have indicated that sublineage II *Nitrospira* account for a

dominant portion of the bacterial community (Gulay *et al.*, 2016) with further investigation

confirming high contributions to its abundance were from comammox-*Nitrospira* (Palomo *et al.*,

2016, Tatari *et al.*, 2017). The strict AOB OTU found in this study was affiliated with

oligotrophic *Nitrosomonas* cluster 6a which exhibit maximum growth rates at ammonia

concentrations similar to the ones used for high and mid nitrogen amendments (Bollmann *et al.*,

2011, Sedlacek *et al.*, 2019). Despite this, the proportions of SSU reads mapping to

Nitrosomonas-like populations in all nitrogen amendments were consistently low. Taxonomic

classification of nitrogen cycling genes revealed metabolic potential for nitrification processes

were confined to *Nitrospi*ra- and *Nitrosomonas*-like populations corroborating with assembled

16S rRNA gene sequences. Additionally, phylogeny of *amoA* sequences found in the

metagenome indicated ammonia oxidation could be mediated by both *Nitrospira* and

Nitrosomonas.

We assembled a total of nine nitrifier MAGs which included comammox-*Nitrospira* (n=4),

Nitrospira-NOB-like (n=2), and *Nitrosomonas*-like (n=3) populations. Three of the four

comammox MAGs assembled were identified as clade A2 based on phylogenetic analyses of

hydroxylamine dehydrogenase (*hao*) which has previously been shown to dominate drinking

water biofilters along with comammox clade B (Palomo *et al.*, 2019). The remaining comammox

MAG assembled from biofilter media in this study was affiliated with clade A1 based on *hao*

gene phylogeny, which while atypical for drinking water biofilters is consistent with previously

published metagenome from the Ann Arbor drinking water filters (Pinto *et al.*, 2016). Similar

coexistence of clade A1 and A2 comammox bacteria with canonical nitrifiers has been observed

in tertiary rotating biological contactors treating municipal wastewater with low ammonium

concentrations (Spasov *et al.*, 2020). However, phylogenomic placement of clade A sub-groups

in this study separated the comammox MAGs into distinct clusters associated with freshwater

(Bin_13, clade A1), groundwater biofilters (Bin_49_4, clade A2) and tap water (Bin_260 and

Bin_49_2_2, clade A2). Maintenance of high functional redundancy for the complete ammonia

oxidation pathway may rely on coexisting comammox populations avoiding direct competition

through distinct physiological niches. Additionally, the innocula were sourced from low

substrate conditions which may also allow for the coexistence of multiple comammox

populations. Strict AOB MAGs obtained in this study associated with low ammonia adapted

Nitrosomonas cluster 6a (Koops *et al.*, 2006) which is consistent with the inocula source being

an oligotrophic environment (i.e., DWTPs). Furthermore, close relatives of *Nitrospira*-NOB

MAGs obtained in this study originated from a tap water source where *Nitrospira*-NOB also

coexisted with strict AOB and comammox bacteria under oligotrophic conditions (Wang *et al.*,

2017). Our findings, consistent with previous studies, confirm the nitrifier community

encompassed multiple populations capable of single and two-step nitrification within a single

system with Nitrospira as the dominant nitrifier. However, the mechanism behind high abundances

of Nitrospira-NOB in biofilters is not yet completely understood. Further, assessment of metabolic

versality revealed initiation of nitrification through urea degradation was possible by all three

nitrifying guilds. Though ureolytic activity is a widespread trait among cultured comammox-

Nitrospira representatives and curated MAGs, the capability is confined to only some

Nitrospira-NOB and *Nitrosomonas* species (Koch *et al.*, 2015, Sedlacek *et al.*, 2019). Here in

particular, this a would allow *Nitrospira*-NOB to play a role in nitrite production in urea

microcosms by crossing feeding ammonia from urea degradation to strict AOB, a mutualistic

strategy which may not be active in ammonia amended microcosms.

Comammox bacterial abundance increased irrespective of nitrogen source or loading but

may compete with NOB depending on nitrogen source type.

We tested the impact of nitrogen source and loading rates on temporal dynamics of a mixed

nitrifying community to determine whether comammox bacteria are outcompeted at higher

concentrations and/or favored in urea amendments due to their ureolytic activity. qPCR-based

abundance tracking revealed comammox bacteria demonstrated a preferential enrichment over

strict AOB in the nitrogen amendments irrespective of nitrogen source or availability.

Additionally, strict AOB abundance did not exhibit any significant difference across the nitrogen

amendment types. This is in contrast to previous work in soil microcosms where AOB

abundance increased in response to high ammonia amendments (He *et al.*, 2021). However, strict

AOB populations in these soil microcosms were primarily *Nitrosospira* compared to oligotrophic

Nitrosomonas cluster 6a which were the primary AOB in this study. Here, both comammox

bacteria and strict AOB demonstrated increased abundance in all amendments during the earlier

weeks of the experiment. Ultimately, while comammox bacteria were enriched over time our

findings demonstrated this increased abundance was not associated with a decrease in the

abundance of strict AOB in any of the nitrogen amendments. This suggests a lack of direct

competition between the two comammox and strict AOB which could be attributable to the two

ammonia oxidizers occupying separate nitrogen availability niches (Martens-Habbena *et al.*,

2009, Kits *et al.*, 2017). Stable abundances of strict AOB compared to enrichment of comammox

could be due to a combination of factors ranging from (1) higher abundances of comammox

bacteria in the inocula and (2) significantly higher biomass yields per mole of ammonia oxidizers

for comammox bacteria compared to AOB (Kits *et al.*, 2017).

Clade A2 associated comammox bacterial MAGs were dominant in the inocula and over the

course of the experiment showed increased abundance in all amendments. In contrast,

comammox bacteria belonging to clade A1 were lower in abundance and did not demonstrate

significant change over time in any amended microcosm. Though physiological differences

between comammox bacteria clades/sub-clades have yet to be established, earlier studies of

DWTP biofilters have observed higher abundances of clade B (Fowler *et al.*, 2018) or

alternatively both clades found at the same DWTP but within separate rapid sand filters, where

clade B was more abundant in the secondary filters receiving lower ammonia concentrations

(Poghosyan *et al.*, 2020). In this study, the lack of clade A1 enrichment over the course of the

experiment may also indicate distinct physiological niches within clades (i.e., subclade-level

niche differentiation). Future research is necessary to develop a clearer understanding of

physiological differences between comammox bacteria at the clade/sub-clade level. Since

cultivability of comammox bacteria remains an ongoing challenge, integrating multiple 'omics

techniques (i.e., metatranscriptomics and metaproteomics) may be an appropriate strategy for

examining ammonia utilization and the expressed metabolisms of multiple coexisting

comammox bacteria populations alongside canonical nitrifiers.

The negative association between comammox bacteria and canonical NOB observed in ammonia

amendments could be a result of nitrite limitation resulting from complete nitrification driven by

comammox bacteria. The possibility of comammox bacteria being a source of leaked nitrite to

Nitrospira-NOB seems unlikely in this case as this would likely form a positive association

between the two. Nitrite limitation driven competition between comammox bacteria and NOB is

supported by the fact the negative associations between the groups were stronger at medium (3.5

mg-N/l) and low (1.5 mg-N/l) nitrogen availability as compared to the high ammonia

amendments (i.e., 14 mg-N/l). In contrast, there was no significant association between the

abundance of comammox bacteria and Nitrospira-NOB in the urea amended systems irrespective

of nitrogen loading. We hypothesize that this variable observations between ammonia and urea

amended systems likely emerge from the extent of metabolic coupling between AOB and NOB

and the resultant ability of comammox to outcompete NOB. Specifically, while the rate of nitrite

availability for NOB in ammonia amended systems is largely dictated by ammonia oxidation

activity of AOB it is likely that nitrite availability in urea amended systems would be dictated by

a combination of both AOB activity and indirectly by NOB. In this case, the production of nitrite

could be mediated by *Nitrospira*-NOB capable of ureolytic activity by crossing feeding ammonia

to strict AOB who in turn provide nitrite at a rate at which *Nitrospira*-NOB. This tight coupling

between AOB and NOB is supported by stronger and more significant correlation between AOB

and NOB abundance in urea amended systems as compared to ammonia amended systems. Thus,

it appears that while comammox bacteria may outcompete *Nitrospira*-NOB in systems where

AOB abundances are low and nitrite availability is largely dictated by AOB activity, this

competitive exclusion may be limited in scenarios with established AOB-NOB cross feeding via

urea where nitrite availability is governed not only by AOB's ammonia oxidation rate but also by

NOB's ureolytic activity. Since urea is used directly by urease-positive nitrifiers, variabilities in

their affinities for the substrate would play a role in the outcome of competition in urea

microcosms but was not assessed in this study.

Altogether, our study demonstrates that comammox bacteria will dominate over canonical

nitrifiers in communities sourced from nitrogen limited environments irrespective of nitrogen

source type or loading rate without directly competing with canonical AOB. Further, our study

also indicates comammox bacteria and AOB may occupy independent niches in communities

sources from low nitrogen environments. Interestingly, we see evidence of potential competitive

exclusion of NOB by comammox bacteria governed by nitrogen source dependent metabolic

coupling between AOB and NOB.

Data availability

Funding sources

This work was supported by NSF Graduate Research Fellowship and Cochrane Fellowship to

KV and by NSF Award number: 1703089.

669 References
670
Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, Ijaz UZ, Lahti L, Loman NJ,
Andersson AF & Quince C (2014) Binning metagenomic contigs by coverage and composition.
Nat Methods **11**: 1144-1146.
Annavajhala MK, Kapoor V, Santo-Domingo J & Chandran K (2018) Comammox Functionality
Identified in Diverse Engineered Biological Wastewater Treatment Systems. *Environ Sci*
Technol Lett **5**: 110-116.
Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S & Ogata H (2020)
KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score
threshold. *Bioinformatics* **36**: 2251-2252.
Bartelme RP, McLellan SL & Newton RJ (2017) Freshwater Recirculating Aquaculture System
Operations Drive Biofilter Bacterial Community Shifts around a Stable Nitrifying Consortium of
Ammonia-Oxidizing Archaea and Comammox Nitrospira. *Front Microbiol* **8**: 101.
Beach NK & Noguera DR (2019) Design and Assessment of Species-Level qPCR Primers
Targeting Comammox. *Front Microbiol* **10**: 36.
Bollmann A, French E & Laanbroek HJ (2011) Isolation, cultivation, and characterization of
ammonia-oxidizing bacteria and archaea adapted to low ammonium concentrations. *Methods*
Enzymol **486**: 55-88.
Bowers RM, Kyrpides NC, Stepanauskas R, *et al.* (2017) Minimum information about a single
amplified genome (MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and
archaea. *Nat Biotechnol* **35**: 725-731.
Camejo PY, Santo Domingo J, McMahon KD & Noguera DR (2017) Genome-Enabled Insights
into the Ecophysiology of the Comammox Bacterium "Candidatus Nitrospira nitrosa". *mSystems*
2.
Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, Fierer N &
Knight R (2011) Global patterns of 16S rRNA diversity at a depth of millions of sequences per
sample. *Proc Natl Acad Sci U S A* **108 Suppl 1**: 4516-4522.
Chaumeil PA, Mussig AJ, Hugenholtz P & Parks DH (2019) GTDB-Tk: a toolkit to classify
genomes with the Genome Taxonomy Database. *Bioinformatics*.
Chen S, Zhou Y, Chen Y & Gu J (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**: i884-i890.

Cotto I, Dai Z, Huo L*, et al.* (2020) Long solids retention times and attached growth phase favor prevalence of comammox bacteria in nitrogen removal systems. *Water Res* **169**: 115268.

Daims H, Lucker S & Wagner M (2016) A New Perspective on Microbes Formerly Known as Nitrite-Oxidizing Bacteria. *Trends Microbiol* **24**: 699-712.

Daims H, Lebedeva EV, Pjevac P*, et al.* (2015) Complete nitrification by Nitrospira bacteria. *Nature* **528**: 504-509.

Edgar RC (2004) MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics* **5**: 113.

Eren AM, Esen OC, Quince C, Vineis JH, Morrison HG, Sogin ML & Delmont TO (2015) Anvi'o: an advanced analysis and visualization platform for 'omics data. *PeerJ* **3**: e1319.

Fowler SJ, Palomo A, Dechesne A, Mines PD & Smets BF (2018) Comammox Nitrospira are abundant ammonia oxidizers in diverse groundwater-fed rapid sand filter communities. *Environ Microbiol* **20**: 1002-1015.

Gonzalez-Martinez A, Rodriguez-Sanchez A, van Loosdrecht MCM, Gonzalez-Lopez J & Vahala R (2016) Detection of comammox bacteria in full-scale wastewater treatment bioreactors using tag-454-pyrosequencing. *Environ Sci Pollut Res Int* **23**: 25501-25511.

Gottshall EY, Bryson SJ, Cogert KI, Landreau M, Sedlacek CJ, Stahl DA, Daims H & Winkler M (2020) Sustained nitrogen loss in a symbiotic association of Comammox Nitrospira and Anammox bacteria. *bioRxiv*.

Graham DW, Knapp CW, Van Vleck ES, Bloor K, Lane TB & Graham CE (2007) Experimental demonstration of chaotic instability in biological nitrification. *ISME J* **1**: 385-393.

Gruber-Vodicka HR, Seah BKB & Pruesse E (2020) phyloFlash: Rapid Small-Subunit rRNA Profiling and Targeted Assembly from Metagenomes. *mSystems* **5**.

Gulay A, Musovic S, Albrechtsen HJ, Al-Soud WA, Sorensen SJ & Smets BF (2016) Ecological patterns, diversity and core taxa of microbial communities in groundwater-fed rapid gravity filters. *ISME J* **10**: 2209-2222.

Gulay A, Fowler SJ, Tatari K, Thamdrup B, Albrechtsen HJ, Al-Soud WA, Sorensen SJ & Smets BF (2019) DNA- and RNA-SIP Reveal Nitrospira spp. as Key Drivers of Nitrification in Groundwater-Fed Biofilters. *mBio* **10**.

Gurevich A, Saveliev V, Vyahhi N & Tesler G (2013) QUAST: quality assessment tool for genome assemblies. *Bioinformatics* **29**: 1072-1075.

He S, Li Y, Mu H, Zhao Z, Wang J, Liu S, Sun Z & Zheng M (2021) Ammonium concentration determines differential growth of comammox and canonical ammonia-oxidizing prokaryotes in soil microcosms. *Applied Soil Ecology* **157**.

Hermansson A & Lindgren PE (2001) Quantification of ammonia-oxidizing bacteria in arable soil by real-time PCR. *Appl Environ Microbiol* **67**: 972-976.

Hiroyuki Ogata SG, Kazushige Sato, Wataru Fujibuchi, Hidemasa Bono and Minoru Kanehisa (1999) KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Research* **27**.

Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW & Hauser LJ (2010) Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* **11**: 119.

Jain C, Rodriguez RL, Phillippy AM, Konstantinidis KT & Aluru S (2018) High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* **9**: 5114.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A & Jermiin LS (2017) ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods* **14**: 587-589.

Kang DD, Li F, Kirton E, Thomas A, Egan R, An H & Wang Z (2019) MetaBAT 2: an adaptive
binning algorithm for robust and efficient genome reconstruction from metagenome assemblies.
PeerJ **7**: e7359.
Kits KD, Sedlacek CJ, Lebedeva EV*, et al.* (2017) Kinetic analysis of a complete nitrifier reveals
an oligotrophic lifestyle. *Nature* **549**: 269-272.
Koch H, Lucker S, Albertsen M, Kitzinger K, Herbold C, Spieck E, Nielsen PH, Wagner M &
Daims H (2015) Expanded metabolic versatility of ubiquitous nitrite-oxidizing bacteria from the
genus Nitrospira. *Proc Natl Acad Sci U S A* **112**: 11371-11376.
Koops H-P, Purkhold U, Pommerening-Röser A, Timmermann G & Wagner M (2006) The
Lithoautotrophic Ammonia-Oxidizing Bacteria. *The Prokaryotes,*p.^pp. 778-811.
Kowalchuk GA & Stephen JR (2001) Ammonia-Oxidizing Bacteria: A Model for Molecular
Microbial Ecology. *Annu Rev Microbiol* **55**: 485-529.
Letunic I & Bork P (2019) Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. *Nucleic Acids Res* **47**: W256-W259.
Li H & Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**: 1754-1760.
Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R &
Genome Project Data Processing S (2009) The Sequence Alignment/Map format and SAMtools.
Bioinformatics **25**: 2078-2079.
Liu T, Wang Z, Wang S, Zhao Y, Wright AL & Jiang X (2019) Responses of ammonia-oxidizers
and comammox to different long-term fertilization regimes in a subtropical paddy soil. *European*
Journal of Soil Biology **93**.
Martens-Habbena W, Berube PM, Urakawa H, de la Torre JR & Stahl DA (2009) Ammonia
oxidation kinetics determine niche separation of nitrifying Archaea and Bacteria. *Nature* **461**:
976-979.
Menzel P, Ng KL & Krogh A (2016) Fast and sensitive taxonomic classification for
metagenomics with Kaiju. *Nat Commun* **7**: 11257.
Na SI, Kim YO, Yoon SH, Ha SM, Baek I & Chun J (2018) UBCG: Up-to-date bacterial core
gene set and pipeline for phylogenomic tree reconstruction. *J Microbiol* **56**: 280-285.
Nguyen LT, Schmidt HA, von Haeseler A & Minh BQ (2015) IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* **32**: 268-
274.
Nurk S, Meleshko D, Korobeynikov A & Pevzner PA (2017) metaSPAdes: a new versatile
metagenomic assembler. *Genome Res* **27**: 824-834.
Palomo A, Dechesne A & Smets BF (2019) Genomic profiling of Nitrospira species reveals
ecological success of comammox Nitrospira. *bioRxiv.*
Palomo A, Jane Fowler S, Gulay A, Rasmussen S, Sicheritz-Ponten T & Smets BF (2016)
Metagenomic analysis of rapid gravity sand filter microbial communities suggests novel
physiology of Nitrospira spp. *ISME J* **10**: 2569-2581.
Palomo A, Pedersen AG, Fowler SJ, Dechesne A, Sicheritz-Ponten T & Smets BF (2018)
Comparative genomics sheds light on niche differentiation and the evolutionary history of
comammox Nitrospira. *ISME J* **12**: 1779-1793.
Parks DH, Imelfort M, Skennerton CT, Hugenholtz P & Tyson GW (2015) CheckM: assessing
the quality of microbial genomes recovered from isolates, single cells, and metagenomes.
Genome Res **25**: 1043-1055.

Parks DH, Rinke C, Chuvochina M, Chaumeil PA, Woodcroft BJ, Evans PN, Hugenholtz P &
Tyson GW (2017) Recovery of nearly 8,000 metagenome-assembled genomes substantially
expands the tree of life. *Nat Microbiol* **2**: 1533-1542.
Pinto AJ, Marcus DN, Ijaz UZ, Bautista-de Lose Santos QM, Dick GJ & Raskin L (2016)
Metagenomic Evidence for the Presence of Comammox Nitrospira-Like Bacteria in a Drinking
Water System. *mSphere* **1**.
Pjevac P, Schauberger C, Poghosyan L*, et al.* (2017) AmoA-Targeted Polymerase Chain
Reaction Primers for the Specific Detection and Quantification of Comammox Nitrospira in the
Environment. *Front Microbiol* **8**: 1508.
Poghosyan L, Koch H, Lavy A, Frank J, van Kessel M, Jetten MSM, Banfield JF & Lucker S
(2019) Metagenomic recovery of two distinct comammox Nitrospira from the terrestrial
subsurface. *Environ Microbiol* **21**: 3627-3637.
Poghosyan L, Koch H, Frank J, van Kessel M, Cremers G, van Alen T, Jetten MSM, Op den
Camp HJM & Lucker S (2020) Metagenomic profiling of ammonia- and methane-oxidizing
microorganisms in two sequential rapid sand filters. *Water Res* **185**: 116288.
Prosser JI & Nicol GW (2012) Archaeal and bacterial ammonia-oxidisers in soil: the quest for
niche specialisation and differentiation. *Trends Microbiol* **20**: 523-531.
Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J & Glockner FO (2007) SILVA:
a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data
compatible with ARB. *Nucleic Acids Res* **35**: 7188-7196.
Quinlan AR & Hall IM (2010) BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**: 841-842.
Rognes T, Flouri T, Nichols B, Quince C & Mahe F (2016) VSEARCH: a versatile open source
tool for metagenomics. *PeerJ* **4**: e2584.
Roots P, Wang Y, Rosenthal AF, Griffin JS, Sabba F, Petrovich M, Yang F, Kozak JA, Zhang H
& Wells GF (2019) Comammox Nitrospira are the dominant ammonia oxidizers in a mainstream
low dissolved oxygen nitrification reactor. *Water Res* **157**: 396-405.
Sakoula D, Koch H, Frank J, Jetten MSM, van Kessel M & Lucker S (2020) Enrichment and
physiological characterization of a novel comammox Nitrospira indicates ammonium inhibition
of complete nitrification. *ISME J*.
Sedlacek CJ, McGowan B, Suwa Y, Sayavedra-Soto L, Laanbroek HJ, Stein LY, Norton JM,
Klotz MG & Bollmann A (2019) A Physiological and Genomic Comparison of Nitrosomonas
Cluster 6a and 7 Ammonia-Oxidizing Bacteria. *Microb Ecol* **78**: 985-994.
Seemann T (2014) Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**: 2068-2069.
Shao YH & Wu JH (2021) Comammox Nitrospira Species Dominate in an Efficient Partial
Nitrification-Anammox Bioreactor for Treating Ammonium at Low Loadings. *Environ Sci*
Technol **55**: 2087-2098.
Shi X, Hu H-W, Wang J, He J-Z, Zheng C, Wan X & Huang Z (2018) Niche separation of
comammox Nitrospira and canonical ammonia oxidizers in an acidic subtropical forest soil under
long-term nitrogen deposition. *Soil Biology and Biochemistry* **126**: 114-122.
Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG & Banfield JF (2018)
Recovery of genomes from metagenomes via a dereplication, aggregation and scoring strategy.
Nat Microbiol **3**: 836-843.
Spasov E, Tsuji JM, Hug LA, Doxey AC, Sauder LA, Parker WJ & Neufeld JD (2020) High
functional diversity among Nitrospira populations that dominate rotating biological contactor
microbial communities in a municipal wastewater treatment plant. *ISME J* **14**: 1857-1872.

Stahl DA & de la Torre JR (2012) Physiology and diversity of ammonia-oxidizing archaea. *Annu*
Rev Microbiol **66**: 83-101.
Tatari K, Musovic S, Gulay A, Dechesne A, Albrechtsen HJ & Smets BF (2017) Density and
distribution of nitrifying guilds in rapid sand filters for drinking water production: Dominance of
Nitrospira spp. *Water Res* **127**: 239-248.
van Kessel MA, Speth DR, Albertsen M, Nielsen PH, Op den Camp HJ, Kartal B, Jetten MS &
Lucker S (2015) Complete nitrification by a single microorganism. *Nature* **528**: 555-559.
Wang J, Wang J, Rhodes G, He JZ & Ge Y (2019) Adaptive responses of comammox Nitrospira
and canonical ammonia oxidizers to long-term fertilizations: Implications for the relative
contributions of different ammonia oxidizers to soil nitrogen cycling. *Sci Total Environ* **668**:
224-233.
Wang M, Huang G, Zhao Z, Dang C, Liu W & Zheng M (2018) Newly designed primer pair
revealed dominant and diverse comammox amoA gene in full-scale wastewater treatment plants.
Bioresour Technol **270**: 580-587.
Wang X, Wang S, Jiang Y, Zhou J, Han C & Zhu G (2020) Comammox bacterial abundance,
activity, and contribution in agricultural rhizosphere soils. *Sci Total Environ* **727**: 138563.
Wang Y, Ma L, Mao Y, Jiang X, Xia Y, Yu K, Li B & Zhang T (2017) Comammox in drinking
water systems. *Water Res* **116**: 332-341.
Wang Z, Cao Y, Zhu-Barker X, Nicol GW, Wright AL, Jia Z & Jiang X (2019) Comammox
Nitrospira clade B contributes to nitrification in soil. *Soil Biology and Biochemistry* **135**: 392-
395.
Wu YW, Simmons BA & Singer SW (2016) MaxBin 2.0: an automated binning algorithm to
recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**: 605-607.
Yang Y, Daims H, Liu Y, Herbold CW, Pjevac P, Lin J-G, Li M & Gu J-D (2020) Activity and
Metabolic Versatility of Complete Ammonia Oxidizers in Full-Scale Wastewater Treatment
Systems. *mBio* **11:e03175-19**.
Zheng M, Wang M, Zhao Z, Zhou N, He S, Liu S, Wang J & Wang X (2019) Transcriptional
activity and diversity of comammox bacteria as a previously overlooked ammonia oxidizing
prokaryote in full-scale wastewater treatment plants. *Sci Total Environ* **656**: 717-722.
866
867
868
869