LORE homomerization is required for 3-OH-C10:0 induced immune signaling

Sabine Eschrig[1], Milena Schäffer[1], Tina Illig[1], Sonja Eibel[1,3], Lin-Jie Shu[1], Atiara Fernandez[1,2], Stefanie Ranf[1*]

[1]Phytopathology, TUM School of Life Sciences, Technical University of Munich, Freising-Weihenstephan, Germany
[2]Current address: Center for Plant Molecular Biology (ZMBP), University of Tübingen, Tübingen, Germany
[3]Current address: Bavarian State Research Center for Agriculture, Institute for Crop Science and Plant Breeding, Freising, Germany

*Correspondence: ranf@wzw.tum.de

ABSTRACT

Perception and processing of various internal and external signals is essential for all living organisms. Plants have an expanded and diversified repertoire of cell surface-localized receptor-like kinases (RLKs) that transduce signals across the plasma membrane. RLKs often assemble into higher-order receptor complexes with co-receptors, regulators and scaffolds to convert extracellular stimuli into cellular responses. To date, the only S-domain-RLK from *Arabidopsis thaliana* with a known ligand and function is *At*LORE, a pattern recognition receptor that senses bacterial 3-hydroxy fatty acids of medium chain length, such as 3-hydroxy decanoic acid (3-OH-C10:0), to activate pattern-triggered immunity. Here we show that *At*LORE forms receptor homomers, which is essential for 3-OH-C10:0-induced immune signaling. *At*LORE homomerization is mediated by the transmembrane and extracellular domain. We show natural variation in the perception of 3-OH-C10:0 within the Brassicaceae family. *Arabidopsis lyrata* and *Arabidopsis halleri* do not respond to 3-OH-C10:0, although they possess a putative LORE orthologue. We found that LORE orthologues of these 3-OH-C10:0 nonresponsive species have defective extracellular domains that can bind the 3-OH-C10:0 ligand but lack the ability to homomerize. Our findings shed light on the activation mechanisms of *At*LORE and explain natural variation of 3-OH-C10:0 perception within the Brassicaceae family.

KEYWORDS

Receptor-like kinase, S-domain, LORE, homomerization, pattern recognition receptor, pattern-triggered immunity, microbe-associated molecular pattern

INTRODUCTION

Perception, processing and integration of a wide range of environmental and cellular stimuli is fundamental to all living organisms. In plants, this is implemented, amongst others, by members of the super-families of receptor-like kinases (RLKs) and receptor-like proteins (RLPs) (Shiu and Bleecker, 2001, 2003; Hohmann et al., 2017; Jamieson et al., 2018; Dievart et al., 2020). They regulate various cellular processes such as growth, development, reproduction or immunity (Shiu and Bleecker, 2001; De Smet et al., 2009; Li and Yang, 2016; Boutrot and Zipfel, 2017; Gou and Li, 2020). Most RLKs comprise an extracellular, presumably ligand binding domain (ECD) with a variety of sequence motifs, a single-span transmembrane domain (TMD) and an intracellular domain (ICD) comprising a serine/threonine protein kinase domain for signal transduction (Shiu and Bleecker, 2001; Dievart et al., 2020). In many cases, additional tyrosine phosphorylation activity was observed. Presumably, RLKs are generally dual-specificity kinases, (Bojar et al., 2014; Macho et al., 2015), but pseudo kinases also exist. RLPs, lacking an intracellular kinase domain, are incapable of signal transduction on their own and require signaling competent partners (Jamieson et al., 2018). RLKs and RLPs are classified according to their ECD motifs, such as leucine-rich-repeat (LRR), L-type lectin (Lec), S-domain (SD) or lysin-motif (LysM) domains (Shiu and Bleecker, 2001, 2003; Dievart et al., 2020).

A common theme of RLK and RLP signaling is their association into higher order receptor complexes via homo- or hetero-oligomerization. They can act as ligand-binding receptors, co-receptors, scaffolds or positive/negative regulators to orchestrate and fine tune signaling (Ma et al., 2016b; Burkart and Stahl, 2017; Wan et al., 2019; Gou and Li, 2020). To date, we have gained a detailed molecular and structural understanding of ligand binding, activation mechanisms and the role of receptor complex formation for a few RLKs, which have become important receptor models. One prototypical heteromeric signaling complex is the heterodimer between LRR-RLK FLAGELLIN SENSING 2 (*At*FLS2) and its LRR-co-receptors of the SOMATIC EMBRYOGENESIS RECEPTOR-LIKE KINASE (*At*SERK) family, particularly BRI1-ASSOCIATED RECEPTOR KINASE 1/SERK3 (*At*BAK1) from *A. thaliana* (Gomez-Gomez and Boller, 2000; Chinchilla et al., 2006; Chinchilla et al., 2007; Roux et al., 2011). FLS2 functions as pattern recognition receptor (PRR) in plant immunity. PRRs perceive microbe-associated molecular patterns (MAMPs) to induce pattern-triggered immunity (PTI) (Boutrot and Zipfel, 2017). Perception of the ligand, the peptide epitope flg22 of bacterial flagellin, triggers association of the *At*FLS2 receptor with its co-receptors (Boller and Felix, 2009; Robatzek and Wirthmueller, 2013; Couto and Zipfel, 2016). Co-crystallization of the ECDs of both *At*FLS2 and *At*BAK1 with flg22 shows a series of intermolecular bonds that stabilize the heterodimer in both ligand- and receptor mediated manner (Sun et al., 2013b). State of the art imaging techniques enable real-time analysis of receptor dynamics in *vivo* and suggested that *At*FLS2/*At*BAK1 heterodimers can further associate into tetrameric complexes (Somssich et al., 2015). The multicomponent PRR complex for perception of the fungal MAMP chitin is characterized in detail as well. In *Arabidopsis*, three LysM-RLKs (LYK), *At*LYK5, *At*LYK4 and the CHITIN ELICITOR RECEPTOR KINASE 1 (*At*CERK1/ *At*LYK1) cooperate in chitin perception and signaling (Miya et al., 2007; Cao et al., 2014; Xue et al., 2019; Gong et al., 2020). Upon chitin binding, the pseudo-kinase *At*LYK5 hetero-dimerizes with *At*CERK1, triggering further homomerization of *At*CERK1. This leads to the formation of a sandwich-like receptor complex to enable downstream signaling (Couto and Zipfel, 2016; Gong et al., 2020). *At*LYK4 interacts with both *At*LYK5 and *At*CERK1 and was reported to have a role in scaffolding or ligand binding (Gong et al., 2020).

While ligand induced hetero-dimerization with co-receptors is essential for signaling in the
previous examples, the mode of action is different for S-LOCUS RECEPTOR KINASEs (SRKs)
from *Brassica rapa*. *Bra*SRKs belong to the family of S-domain-RLKs (SD-RLKs, also referred to
as G- or B-type lectin RLKs) which comprises about 40 members in Arabidopsis and more than
100 in rice (Shiu and Bleecker, 2001; Shiu et al., 2004; Vaid et al., 2012; Xing et al., 2013; Teixeira
et al., 2018). SRKs and their ligands, namely S-LOCUS CYSTEIN-RICH PEPTIDE (SCRs),
mediate self-incompatibility (SI) to maintain genetic variability by avoiding inbreeding (Ivanov et
al., 2010; Nasrallah and Nasrallah, 2014; Jany et al., 2019). Recognition of pollen-secreted SCRs
by SRKs expressed in the stigma enables the plant to detect and subsequently reject self-pollen
(Nasrallah and Nasrallah, 2014; Jany et al., 2019). The ECD of SRKs comprises two G-type lectin-
like, an epidermal-growth factor (EGF)-like and a plasminogen-apple-nematode (PAN) domain,
representing one typical domain architecture of SD-RLKs (Shiu and Bleecker, 2001; Xing et al.,
2013; Dievart et al., 2020). Spontaneous and ligand-independent homomerization of SRKs via the
ECD has been reported (Giranton et al., 2000; Naithani et al., 2007). However, the crystal
structures obtained from *Bra*SRK9 and engineered *Bra*SRK8 elucidated, that two SKRs and two
of the respective SCR peptides form a 2:2 hetero-tetrameric complex in a ligand- and receptor-
mediated manner (Ma et al., 2016a; Murase et al., 2020). Receptor homo-dimerization of SRKs
appears to be essential for SRK signaling. To date, no co-receptors are found to be involved in
SRK receptor complexes.

Apart from SRKs, only a few SD-RLKs from different plant species have been described in more
detail. They are involved in plant-pathogen interactions, symbiosis or abiotic stress responses
(Navarro-Gochicoa et al., 2003; Chen et al., 2006; Kanzaki et al., 2008; Kim et al., 2009b; Kim et
al., 2009a; Gilardoni et al., 2011; Chen et al., 2013; Cheng et al., 2013; Sun et al., 2013a; Zou et
al., 2015; Fan et al., 2018; Schnepf et al., 2018; Labbé et al., 2019; Jinjun et al., 2020; Pan et al.,
2020; Sun et al., 2020; Liu et al., 2021; Zhou et al., 2021). In *Glycine soja*, S-LOCUS LecRK
(*Gs*SRK) regulates osmotic homeostasis and plant architecture under salt stress (Sun et al.,
2013a; Sun et al., 2018). *Oryza sativa Os*Pi-d2 confers resistance to rice blast caused by the
fungus *Magnaporthe grisea* (Chen et al., 2006). Interestingly, natural variation in a single amino
acid position in the TMD appears to define resistant or susceptible alleles (Chen et al., 2006; Li et
al., 2015). Expression of the LARGE SPIKE S-DOMAIN RECEPTOR LIKE KINASE 1 (*Os*LSK1)
in rice is regulated by growth hormones and is associated with abiotic stress sensitivity and yield.
The ECD of *Os*LSK1 interacts with itself and heteromerizes with the ECDs of five homologous
SD-RLKs. Overexpression of truncated *Os*LSK1 lacking the ICD increases plant height and grain
yield, supposedly by exerting a dominant negative effect on endogenous *Os*LSK1 signaling (Zou
et al., 2015). Although a growing number of members of the large SD-RLK sub-family are being
characterized, their ligands, receptor complex formation and downstream signaling mechanisms
remain largely unknown. Biochemical characterization of SD-RLKs and identification of ligands is
proving to be particularly challenging. Large-scale expression and purification of SD-RLKs
appears to be especially difficult, likely because of their complex folding, high degree of
glycosylation, and tendency of protein aggregation (Murase et al., 2020; Sun et al., 2020).

In *A. thaliana*, we previously discovered the SD-RLK *At*LORE (LIPOOLIGOSACCHARIDE
SPECIFIC REDUCED ELICITATION or SD1-29) as PRR for bacterial medium-chain 3-
hydroxylated fatty acid (3-OH-FA) metabolites (Ranf et al., 2015; Kutschera et al., 2019). 3-OH-
FAs are sensed in a chain-length and hydroxylation specific manner (Kutschera et al., 2019). Only
3-OH-FAs with acyl chains comprising 8 – 12 carbon atoms activate *At*LORE signaling, with 3-
hydroxydecanoic acid (3-OH-C10:0) being the strongest elicitor in *A. thaliana*. 3-OH-C10:0 directly
binds to the *At*LORE ECD, as evidenced by microscale thermophoresis (MST) and ligand

depletion-binding assays (Kutschera et al., 2019; Shu et al., 2021). *At*LORE is the first SD-RLK in
A. thaliana with a known ligand and thus a controllable means of activation, making it an important
model for mechanistic studies of the SD-RLK family. Recently, downstream signaling components
of *At*LORE were elucidated. Sensing of 3-OH-C10:0 leads to phosphorylation of *At*LORE at Y600,
which is required for phosphorylation and activation of the receptor-like cytoplasmic kinase PBS1-
LIKE 34 (*At*PBL34). *At*LORE directly interacts with *At*PBL34 and its homologs *At*PBL35 and
*At*PBL36, which seem to redundantly facilitate downstream signaling in *At*LORE-mediated
immune responses (Luo et al., 2020). To date, no co-receptor has been found to be involved in
*At*LORE signaling and the steps of receptor complex formation are unknown.

Here, we show that *At*LORE forms receptor homomers via TMD and ECD, which is required for
3-OH-C10:0-induced immune signaling. Interestingly, we found natural variations in
responsiveness to 3-OH-C10:0 in some closely related Brassicaceae, which have LORE
orthologues with high protein sequence identity to *At*LORE. Functional analysis of these
orthologues and chimera with *At*LORE shows that orthologues from nonresponsive species have
defects in their ECDs. Interestingly, the defective LORE orthologues still bind the 3-OH-C10:0
ligand via their ECDs but are compromised in homomerization. This highlights that receptor
homomerization is essential for LORE-dependent immune signaling. Moreover, our results
indicate that in the case of LORE, ligand binding is independent of homomerization, in contrast to
ligand- and receptor-mediated homodimerization of *Bra*SRKs. Our results show that two
Brassicaceae species closely related to *A. thaliana* lack the ability to respond to 3-OH-C10:0,
which is explained by altered homomerization ability. This sheds light on the mechanistic aspects
of 3-OH-C10:0 sensing in Brassicaceae.

RESULTS

_At_LORE forms receptor homomers _in planta_

We performed multiple sequence alignments (MAFFT algorithm, Jalview) (**Fig. S1**) of the amino
acid sequence of the ECDs of *At*LORE (AT1G61380) and well-studied SRKs from *Brassica rapa*,
*Bra*SRK9 (BAA21132.1) and *Bra*SRK8 (BAF91375). Labeling of percentage identity shows a high
sequence similarity and conservation of disulfide bridge forming cysteine residues in the EGF and
PAN domain, indicating an overall similar domain architecture of *At*LORE and *Bra*SRKs. Transient
agrobacterium-mediated overexpression of kinase-active *At*LORE in *Nicotiana benthamiana*,
which lacks a putative LORE orthologue (Ranf et al., 2015), results in a cell death-like necrotic
phenotype. These cell death symptoms can be visualized by trypan blue staining (**Fig. S2E**) and
an increased chlorophyll fluorescence (**Fig. 1A**), which is associated with cell death (Landeo
Villanueva et al., 2021). This phenomenon is not observed upon expression of AtLORE-Km, which
is mutated at the conserved ATP-binding site (K516A) in the kinase domain and has been shown
to be signaling incompetent (Ranf et al., 2015). Therefore, we assume that cell death is caused
by spontaneous receptor activation. Based on the high protein sequence similarity of *At*LORE to
*Bra*SRKs and the receptor auto-activation of *At*LORE upon transient overexpression in *N.
benthamiana*, we hypothesized that *At*LORE may form homomers like *Bra*SRKs. Indeed, *At*LORE
homomerizes *in vivo* upon transient expression in *N. benthamiana*. GFP and mCherry fusion
proteins of kinase-active and kinase-mutated (Km, K516A) *At*LORE co-immunoprecipitate (Co-IP)
with each other (**Fig. 1B**). *At*LORE homomerization was also observed in bimolecular
fluorescence complementation (BiFC) assays using *At*LORE-split-YFP N-/C-terminal fragment
expression (SPYCE/SPYNE) fusion proteins. To avoid background auto-fluorescence caused by

the cell-death symptoms, *At*LORE-Km was used for fluorescence-based assays. Transient co-
expression of *At*LORE-Km fused to either SPYCE or SPYNE restores YFP fluorescence,
detectable via laser scanning microscopy and fluorescence quantification (**Fig. S2A-C**). We
performed Förster resonance energy transfer (FRET) fluorescence lifetime imaging (FLIM) to
proof direct physical LORE-LORE interaction (**Fig. 1C and Fig. S2D**). Co-expression of *At*LORE-
Km-GFP and *At*LORE-Km-mCherry results in a significant reduction in GFP fluorescence lifetime
τ compared to the FRET-donor only control *At*LORE-Km-GFP. mCherry-tagged Glutathione-S
transferase of *A. thaliana* (AT1G17170, *At*GST-mCherry), was used as co-expression control and
does not reduce GFP fluorescence lifetimes. An *At*LORE-Km-GFP-mCherry fusion protein serves
as FRET positive control. *At*LORE homomerization is not influenced by the presence of its ligand
3-OH-C10:0 in FRET-FLIM experiments (**Fig. 1D**). Thus, under the experimental conditions used,
*At*LORE forms homomers in the plasma membrane in a ligand-independent manner.

AtLORE homomerization is mediated by the extracellular- and transmembrane domain

To identify the region of *At*LORE that mediates its homomerization, we generated truncated
variants of *At*LORE containing different combinations of ICD, ECD or TMD (**Fig. 2A**). Full length
*At*LORE-Km-HA co-immunoprecipitates with full length *At*LORE-Km-GFP, *At*LORE-ECD-TMD-
GFP and weakly with *At*LORE-TMD-ICD-Km-GFP upon transient expression in *N. benthamiana*
leaves (**Fig. 2B**). No co-immunoprecipitation is detected with apoplastic *At*LORE-ECD-GFP,
cytosolic *At*LORE-ICD-Km-GFP or cytosolic GFP. This indicates that ECD and TMD contribute to
LORE homomerization. In FRET-FLIM experiments, both *At*LORE-ECD-TMD-mCherry and
*At*LORE-TMD-ICD-Km-mCherry significantly reduce the GFP fluorescence lifetime of the FRET
donor *At*LORE-Km-GFP (**Fig. 2C and S3**). Furthermore, *At*LORE-ECD-TMD lacking the ICD can
still homomerize and interact with *At*LORE-TMD-ICD (**Fig. 2D**). In the latter case, only the TMDs
can serve as an interaction interface, yet GFP fluorescence lifetime is reduced, highlighting the
contribution of both ECD and TMD to *At*LORE homomerization.

A. lyrata and A. halleri have LORE orthologues, but do not respond to 3-OH-C10:0

Phylogenetic analysis shows that *LORE* is restricted to the plant family of Brassicaceae (Ranf et
al., 2015). Putative LORE orthologues are encoded in the genomes of *Arabidopsis halleri*
(*Aha*LORE, Araha.6790s0007.1), *Arabidopsis lyrata* (*Alyr*LORE, AL2G04470) and *Capsella*
rubella (*Crub*LORE, CARUB_v10021901mg) (**Fig. S1**). Therefore, we analyzed these species for
the production of reactive oxygen species (ROS) (**Fig. 3A**), which is a typical PTI response to
application of 3-OH-C10:0 in *A.thaliana* (Kutschera et al., 2019). Interestingly, we found that, in
contrast to *C. rubella* and *A. thaliana*, *A. lyrata* and *A. halleri* do not respond to 3-OH-C10:0
elicitation. All four species produced ROS upon flg22 treatment (**Fig. S4B**). This suggests that
functional PTI signaling pathways (Yu et al., 2017) and components for ROS production are in
principle functional. That *A. lyrata* and *A. halleri* do not respond to 3-OH-C10:0 can originate from
diverse failures, such as lack of expression, morphological differences that hinder the access of
3-OH-C10:0 to the PRR or a dysfunctional LORE protein. For further analysis, we extracted RNA
from untreated leaf material of *C. rubella*, *A. lyrata* and *A. halleri* and transcribed it into cDNA to
clone the coding sequence (CDS) of *LORE*. We obtained full length CDS clones of all orthologues,
confirming that *LORE* is expressed in *A. lyrata* and *A. halleri* leaf tissue used for the ROS assay
(**Fig. 3A**). Amino acid sequences of *Alyr*LORE and *Aha*LORE slightly varied from publicly
available database sequences (**Fig. S8**). Furthermore, transient expression of GFP-fusions of

LORE orthologues in *N benthamiana* shows that they localize to the plasma membrane,
*Ahal*LORE-GFP was partially mislocalized (**Fig. S4A**). Solanaceous *N. benthamiana* has no
putative LORE orthologue and is therefore insensitive to 3-OH-C10:0 but can gain the function of
3-OH-C10:0 sensing by transient expression of *At*LORE (Ranf et al., 2015; Kutschera et al., 2019).
Therefore, we tested the cloned LORE orthologues for functionality in gain-of-function (GOF) ROS
assays in *N. benthamiana* (**Fig. 3B**). Expression of *At*LORE or *Crub*LORE, but not *Ahal*LORE or
*Alyr*LORE resulted in a ROS response to treatment with 3-OH-C10:0. Taken together, these data
support that the LORE orthologues of *A. lyrata* and *A. halleri* are the underlying factor in the
inability of these species to respond to 3-OH-C10:0.

LORE orthologues have elicitor binding capacity

To identify the region of *Alyr*LORE that causes its dysfunction, we created reciprocal domain
swaps (DS) between ICD and ECD of *At*LORE and *Alyr*LORE and tested those domain swaps
(DS) for their GOF-ROS responses in *N. benthamiana* (**Fig. 4A**). Interestingly, we could render
*At*LORE inactive by replacing the ECD and TMD with *Alyr*LORE-ECD-TMD. In contrast, *At*LORE
remains functional, when the *At*LORE-ICD is substituted by *Alyr*LORE-ICD. From this we can
conclude that *Alyr*LORE has a signaling-competent kinase domain and a dysfunctional ECD. Next,
we tested whether the *Alyr*LORE-ECD is compromised in ligand binding. We have previously
shown with ligand-depletion assays that *At*LORE and *Crub*LORE, but not closely related *At*SD1-
23 (AT1G61390) can bind 3-OH-C10:0 (Shu et al., 2021). Here, we additionally tested the 3-OH-
C10:0 binding capacities of the ECDs of *Alyr*LORE and *Ahal*LORE expressed and harvested from
N. benthamiana apoplasts in ligand-depletion assays. Equal amounts of total protein (**Fig. S5**)
were incubated with 3-OH-C10:0 and mixtures were filtered through membranes with a 30 kDa
molecular weight cut-off. The presence of unbound 3-OH-C10:0 in the filtrate was tested in a
bioassay by measuring the cytosolic calcium response of *At*LORE-overexpressing (OE) *A.*
thaliana seedlings. When the ECDs bind the elicitor, the filtrates do not contain 3-OH-C10:0 and
do not elicit a calcium response. Interestingly, the ECDs of all tested LORE orthologues bind 3-
OH-C10:0 and completely deplete 3-OH-C10:0 from the filtrates, as no calcium responses is
detectable after filtrate application (**Fig. 4B**). Since neither the kinase activity nor the ligand binding
of *Alyr*LORE appears to be affected, the impaired 3-OH-C10:0 perception of *Alyr*LORE must have
other causes.

LORE orthologues from 3-OH-C10:0 nonresponsive species cannot homomerize

Interestingly, unlike overexpression of *At*LORE or *Crub*LORE, overexpression of *Alyr*LORE or
*Ahal*LORE in *N. benthamiana* does not cause cell death, as determined by chlorophyll
fluorescence measurements (**Fig. 5A**). This indicates that receptor activation, relevant for the
autoimmunity phenotype, is affected in these species. Therefore, we hypothesized that *Alyr*LORE
and *Ahal*LORE may fail to homomerize and analyzed their ability to homomerize using FRET-
FLIM experiments in *N. benthamiana* (**Fig. 5B-D**). Indeed, *Alyr*LORE and *Ahal*LORE are unable
to form homomers in *vivo*, whereas signaling-competent *Crub*LORE homomerizes similarly to
*At*LORE. Thus, we conclude that homomerization of LORE is crucial for the activation of signal
transduction. *Alyr*LORE and *Ahal*LORE can bind 3-OH-C10:0 but do not homomerize. Thus,
ligand binding is independent of receptor homomerization and not sufficient to activate LORE-
dependent signaling. Furthermore, our data indicate that the cell death phenotype in *N.*

benthamiana can be used as indicator of LORE receptor functionality and requires both, an active
kinase domain and receptor homomerization.

Mapping of homomerization region using *At*LORE and *Alyr*LORE chimera

Our data suggest that loss of the homomerization ability of LORE renders two *Arabidopsis*
species, *A. lyrata* and *A. halleri,* insensitive to 3-OH-C10:0. While *Alyr*LORE and *Ahal*LORE share
a very high overall amino acid identity with *At*LORE, they have several single amino acid
polymorphisms (SAPs) in their ECDs and TMD compared to the *At*LORE-ECD, distributed across
all domains (**Fig. S1**). To narrow down the region of the ECD that causes the loss of *Alyr*LORE
homomerization, we generated chimera between *At*LORE and *Alyr*LORE ECDs and TMDs. We
either exchanged individual domains, such as lectin 1, lectin 2, EGF or PAN domain (L1, L2, E, P,
Fig. S7A) or combinations of domains (LL, EP, **Fig. 6A**). After verifying the expression and correct
localization in *N. benthamiana* epidermal cells (**Fig. S6**), we analyzed homomerization of these
chimera in FRET-FLIM experiments (**Fig. 6 and S7B**). Chimera with individual domain swaps all
retained their ability to homomerize (**Fig. S7B**). This indicates that loss of homomerization in
*Alyr*LORE is not caused by a single SAP or diversification of a single domain. Therefore, we tested
domain combinations, namely LL, EP and full ECD-TMD swaps. Interestingly, substitution of the
whole ECD of *At*LORE by *Alyr*LORE, but not partial ECD swaps (LL, EP) impair homomerization
in FRET-FLIM experiments (**Fig. 6B**). A similar outcome was observed, when we tested these
chimeras in 3-OH-C10:0-triggered ROS or chlorophyll fluorescence assays upon overexpression
in *N. benthamiana* (**Fig. 4A, 6C and D**). Chimera with partial ECD swaps remain signaling
competent, while those with ECD-TMD exchange do not. Hence, we could not pin down a specific
region of the *Alyr*LORE-ECD or TMD that causes the loss of homomerization. This might indicate
a large, ectodomain-spanning interaction interface of *At*LORE.

Homomerization is essential for *At*LORE downstream signaling

Collectively, our data suggests that LORE homomerization is essential for its activation and
downstream signaling. To confirm this, we performed competition experiments with the truncated
*At*LORE-ECD-TMD variant. We have shown that *At*LORE-ECD-TMD lacking the ICD can form
hetero-complexes with full length *At*LORE (**Fig. 2B and C**). The hetero-complexes are presumably
signaling incompetent as they contain only one kinase domain. To outcompete *At*LORE homo-
complexes, *At*LORE-ECD-TMD must be present in excess relative to *At*LORE. We exploited the
fact, that upon transient expression in *N. benthamiana At*LORE-ECD-TMD accumulates to higher
protein levels than full length *At*LORE (**Fig. 1B**). To enhance the effect, we increased the ratio of
Agrobacteria carrying the competitor expression plasmid (*At*LORE-ECD-TMD) to those with full
length *At*LORE to 5:1. Indeed, *At*LORE-ECD-TMD outcompetes signaling-competent full length
*At*LORE homomers in competition assays, as shown by impaired ROS production in *N.*
benthamiana (**Fig. 7**). This is not observed when cytosolic mCherry is co-expressed in excess.
*At*LORE-ECD-TMD thus exerts a dominant-negative effect on the ROS response upon elicitation
with 3-OH-C10:0. Hence, homomerization of kinase-active LORE is required to activate signaling
upon 3-OH-C10:0 elicitation.

DISCUSSION

*At*LORE is the first SD-RLK from *A. thaliana* with a known ligand and characterized function (Ranf et al., 2015; Kutschera et al., 2019), which makes it an important model for studying signaling mechanisms of the SD-RLK family. However, the mechanism of *At*LORE receptor activation remains largely unknown. Here we show that *At*LORE forms receptor homomers *in vivo* and that homomerization is required to activate immune signaling. In plants, numerous examples of receptor-coreceptor hetero-dimerization have been found, making it a predominant concept of receptor activation mechanisms (Burkart and Stahl, 2017; Wan et al., 2019; Gou and Li, 2020). In contrast, the receptor activation by homo-dimerization, as shown here for *At*LORE, seems to be common among representatives of the SD-RLK family. The SD-RLKs *Os*LSK1 homomerizes and heteromerizes with five close homologs *in yeast* and *in planta* (Zou et al., 2015). Since its ligand and physiological function are unknown, the mechanistic relevance of the observed interactions also remains unclear. Homo-dimerization of *Bra*SRKs has been studied in detail and was shown to be required for ligand binding and signal transduction (Giranton et al., 2000; Naithani et al., 2007; Shimosato et al., 2007; Ma et al., 2016a; Murase et al., 2020). To date, no co-receptors of any SD-RLK has been identified, which underlines the relevance of homo-dimerization for receptor activation. Protein homomerization analysis is particularly challenging due to uncontrollable homodimer stoichiometry. Most protein-protein interaction methods require two different protein epitope tags, whether for FRET, Co-IP, or BiFC (Xing et al., 2016). However, the combinations of protein-tags in a homomer/dimer is subject to random distribution. Statistically, only one-third of dimers have a combination of two different tags when expressed equimolar. Therefore, large proportions of homomers remain unconsidered or even influence the readout of the experiment. For example, in Co-IPs homodimers with identical tags compete with homodimers containing different tags for binding to the antibody trap, but only the latter are detectable on the immunoblot. In case of FLIM the stoichiometry of homo-dimerization results in formation of FRET-incapable (GFP-GFP/mCherry-mCherry) or FRET-capable (GFP-mCherry) pairs. The method does not provide the resolution to distinguish individual molecules. Rather, the measured average GFP lifetime of each pixel is the result of a random mixture of all possible dimers. This might explain, why the GFP lifetime is only moderately reduced for *At*LORE homomers compared to the FRET-positive control, as FRET-incapable *At*LORE-GFP homodimers diminish the effect of the FRET-capable *At*LORE-GFP–*At*LORE-mCherry dimers.

*Bra*SRKs are thought to pre-assemble into homo-dimers in a ligand-independent manner and these preformed complexes allow ligand binding and a rapid activation of the receptor (Giranton et al., 2000; Naithani et al., 2007; Shimosato et al., 2007). Ligand binding might enhance *Bra*SRK dimerization by rearrangement of these pre-assembled complexes (Shimosato et al., 2007). However, whether preformed receptor complexes exist prior to ligand binding under physiological conditions is controversially discussed. *At*CERK1 forms homo-dimers independent of its ligand chitin upon overexpression in *A. thaliana* (Liu et al., 2012), whereas under physiological conditions *At*CERK1 homo-dimerization was fully dependent on *At*LYK5 and chitin binding (Cao et al., 2014; Gong et al., 2020). Ligand-independent self-association of *At*FLS2 was also reported (Sun et al., 2012). FLIM analysis of *At*FLS2 at the membrane did not show homomerization, leading to the speculation that *At*FLS2 might only self-associate upon internalization in a ligand-independent manner (Somssich et al., 2015). Our FRET-FLIM data show that *At*LORE homomerization is ligand-independent, at least under the experimental conditions used. We could not detect any significant differences in the homomerization state of *At*LORE between 10-20 minutes after application of 3-OH-C10:0. Although receptor complex formation should be a rapid process, this timeframe was suggested for comparable FLIM studies (Somssich et al., 2015). The authors

discussed that a relatively large proportion of complexes need to accumulate to change the
average lifetime of one pixel in an FLIM image. To avoid strong overexpression which might lead
to artifacts as shown for *At*CERK1(Liu et al., 2012), we used an inducible promotor system for an
adjustable and more moderate protein expression. To date we can only speculate how *At*LORE
homomerization acts under physiological conditions as protein expression driven by the
endogenous LORE-promotor is below the detection limit for fluorescence-based assays. At
present our data do not allow us to draw conclusions about the oligomerization state of *At*LORE.
However, for *Bra*SRKs oligomerization into higher order complexes was suggested by cross-
linking experiments and velocity sedimentation on sucrose gradients (Giranton et al., 2000), but
its physiological relevance remains unclear. Such oligomerization dynamics of receptor complex
formation was also reported for other receptor complexes, such as AtFLS2-*At*BAK1 (Somssich et
al., 2015). The plant may maintain receptor complexes in a stable preformed steady-state to
ensure rapid signal transduction that is fine-tuned by multiple regulators in higher-order
complexes.

Our data suggest that homomerization of *At*LORE is mediated by the ECD and TMD. In Co-IPs,
neither the apoplastic ECD nor the cytosolic ICD could interact with full-length *At*LORE, suggesting
that membrane anchoring is required for dimerization. However, it is also possible that the
interaction surface of the membrane-bound full-length *At*LORE is not accessible for the soluble
truncations. Interestingly, *At*LORE-ECD-TMD and *At*LORE-TMD-ICD can interact in FRET-FLIM
experiments. In this combination, only the TMDs provide a potential interaction surface suggesting
that the TMD substantially contributes to *At*LORE homomerization. TMD helixes represent a
typical interaction interface in the hydrophobic environment of the membrane, which often contain
conserved TMD interaction motifs (Herrmann et al., 2009; Langosch and Arkin, 2009; Fink et al.,
2012). The relevance of TMDs for receptor complex formation is highly discussed in the animal
field (Westerfield and Barrera, 2020). Involvement of TMDs in dimerization has also been shown
for several plant RLKs. The TMD of the RLK SUPPRESSOR OF BRI1-1 (SOBIR1) possesses a
GxxxG-motif, which mediates high affinity TMD-TMD association and is crucial for its interaction
with the RLK Cf-4 (Bi et al., 2015). The TMD of the RLK Arabidopsis CRINKLY 4 (ACR4) homo-
dimerizes (Stokes and Gururaj Rao, 2008) and is specifically required for the interaction with the
RLK CLAVATA1 (CLV1) (Stahl et al., 2013). The relevance of TMDs in SD-RLK signaling is
underlined by *Os*Pi-d2, in which a single amino acid polymorphism in the TMD (I441M) determines
susceptible or resistant alleles towards rice blast. For *Bra*SRKs it was hypothesized that the TMDs
may contribute to the preformed SRK dimer, which is then strengthened by ligand binding
(Shimosato et al., 2007; Ma et al., 2016a). Interestingly, high-affinity binding of SCR8 peptides
was only observable for membrane-bound SRK8 variants containing the ECD, TMD and parts of
the juxta-membrane domain, but not for soluble ECDs (Takayama et al., 2001; Shimosato et al.,
2007). An artificially dimerizing form of the SRK8-ECD, achieved by integration of a helix-loop-
helix zipper domain, exhibited a high SCR binding affinity, which highlights the importance of
membrane anchorage for ligand binding and dimerization (Shimosato et al., 2007). *Bra*SRK homo-
dimerization was furthermore shown to be mediated by the ECD in yeast-two-hybrid studies, while
kinase domains do not self-associate (Naithani et al., 2007). Especially the PAN and partially the
EGF domains, but not the lectin domains exhibited homo-dimerization capacities in yeast (Naithani
et al., 2007). For *Bra*SRK9 and engineered *Bra*SRK8, crystal structures of their ECDs show both
ligand- and receptor mediated interaction (Ma et al., 2016a; Murase et al., 2020). In both cases,
an *Bra*SRK-ECD dimer co-crystallized with two *Bra*SCR peptide ligands, forming a tetrameric 2:2
receptor-ligand complex. Is it suggested that *Bra*SCRs majorly contribute to complex formation by
cross-linkage of the *Bra*SRK monomers. Although studies about *Bra*SRKs provide a detailed
insight into homo-dimerization, the comparability between *At*LORE and *Bra*SRKs is limited as

*At*LORE binds a very small chemical compound instead of a peptide ligand. So far, the binding
site of 3-OH-C10:0 in the *At*LORE-ECD is unknown. Due to the small size of the ligand, one must
assume a relatively small interface between the receptor and the ligand, which may not be
sufficient to crosslink two *At*LORE monomers or to enhance homo-dimerization in a manner
comparable to SRK-SCR complexes. Interestingly, we found that *Alyr*LORE and *Ahal*LORE
cannot homo-dimerize but bind the 3-OH-C10:0 ligand. Thus, 3-OH-C10:0 can bind to LORE
monomers. One may speculate that dimerization is stabilized by conformational changes rather
than ligand-mediated cross-linking. Taken together, it seems reasonable that *At*LORE may form
pre-assembled complexes in a ligand-independent manner. However, the underlying molecular
mechanism of receptor activation remains unclear.

*At*LORE homomerization is essential for receptor activation and downstream signaling. We
demonstrate that LORE orthologues of 3-OH-C10:0 insensitive species are impaired in their
homomerization capacity, and therefore cannot mediate downstream signaling. Truncated
*At*LORE containing only ECD and TMD exerts a dominant negative effect on 3-OH-C10:0 induced
signaling. We assume that this effect is caused by formation of non-functional heterodimers with
full-length *At*LORE, which lack downstream signaling capacities. Competition for ligand binding
seems unlikely given the high ligand concentrations used in our experiments. Similar competition-
based phenomena have been repeatedly described for other RLKs. Overexpression of truncated
*Os*LSK1 lacking the kinase domain resulted in increased plant height and yield in rice (Zou et al.,
2015). *Os*LSK1 was shown to interact with itself and five homologous proteins. Truncated *Os*LSK1
may interfere with signaling of *Os*LSK1 and the other interacting SD-RLKs or may compete for
ligand binding. Upon overexpression of *At*FLS2 truncations lacking the LRR domain a similar
effect was observable. The truncations associate with full-length *At*FLS2 and exert a dominant
negative effect on flg22-induced signaling (Sun et al., 2012). Taken together, these studies show
that analysis of a dominant negative effect, which is especially applicable for homomerizing RLKs,
provides valuable insight into receptor complexes and their signaling mechanisms.

In vitro phosphorylation assay of microsomal membranes shows that oligomerization of
membrane-bound recombinant *Bra*SRKs is essential for auto-phosphorylation (Giranton et al.,
2000) and that preformed *Bra*SRK oligomers exhibit a basal constitutive auto-phosphorylation.
We assume a similar mode of action for *At*LORE. Strong overexpression of *At*LORE in *N.*
benthamiana results in cell death. We can show that this phenotype depends on both, an active
kinase domain and dimerization ability, as it is completely absent in non-homomerizing LORE-
orthologues or kinase mutated *At*LORE. Induced cell death is usually known as a highly effective
mechanism of hypersensitive defense responses after pathogen recognition (Balint-Kurti, 2019).
However, the phenomenon of cell death-like symptoms upon RLKs overexpression has been
described repeatedly. Overexpression of *At*CERK1 in *N. benthamiana* resulted in cell death in a
kinase-dependent manner (Pietraszewska-Bogiel et al., 2013). It was hypothesized that
overproduction of *At*CERK1 induces ligand-independent homomerization, leading to auto-
activation and dysregulated signal transduction. However, this *At*CERK1 cell-death phenotype
was not apparent upon stable expression in *A. thaliana*. Two L-type lectin RLKs (LecRKs),
*At*LecRK-IX.1 and *At*LecRK-IX.2, exhibit cell death upon overexpression in both *A. thaliana* and
N. benthamiana (Wang et al., 2015). The cell death correlated with transgene expression levels
and was dependent on kinase activity, but not on extracellular lectin domains. Homomerization
was not shown for *At*LecRK-IX.1 and *At*LecRK-IX.2, yet is also shown for other *At*LecRKs (Guo
et al., 2018). Taken together, this suggests that homomerizing RLKs are in particular prone to
spontaneous auto-activation upon strong overexpression, resulting in cell death. In the case of

LORE, we found that this phenotype can serve as a proxy and simple read-out for signaling active
receptor complexes.

Interestingly, we found natural variation in 3-OH-C10:0 responsiveness within a subset of
Brassicaceae species. *A. lyrata* and *A. halleri* are closely related to *A. thaliana* and have LORE
orthologues with a high degree of sequence identity to *At*LORE. While trying to decipher the
underlying defects in these two 3-OH-C10:0-insensitive species, we found that their orthologues
can bind the 3-OH-C10:0 ligand but exhibit defective dimerization capacities, which render them
signaling incompetent. Since we found a number of SAPs across the *Alyr*LORE-ECD, we
investigated chimera of *At*LORE and *Alyr*LORE to identify the causal region or SAP. However,
only substitution by the entire *Alyr*LORE-ECD abolishes homomerization. We hypothesize that
several SAPs collectively cause the loss of homomerization. From an evolutionary perspective,
random single polymorphisms e.g. in the ligand binding site, seem more likely to lead to a loss-of-
function mutation than a series of SAPs accumulating in the ectodomain leading to a loss of
homomerization. This has been shown for example for FLS2. Flg22 binding capacities to putative
FLS2 orthologues were analyzed in several *A. thaliana* accessions. The data suggested a direct
correlation between the FLS2 binding capacities and the responsiveness to flg22 (Bauer et al.,
2001). While our findings were rather unexpected from an evolutionary perspective, they may
shed light on the evolutionary processes of diversification and neo-functionalization in the family
of SD-RLKs which is so far rather poorly understood (Xing et al., 2013). This supports that natural
variation provides a powerful molecular toolbox to improve our mechanistic understanding of RLK
signaling.

MATERIALS AND METHODS

Sequence alignments

Multiple amino acid sequence alignments (MSA) (Fig. S1) were performed with Jalview (Version
2.11.1.4 (Waterhouse et al., 2009)). Amino acid sequences of respective proteins were retrieved
from the Uniprot database (Consortium, 2020). For LORE, ECD regions excluding signal peptides
(SP) were selected according to domain annotations given on Uniprot and in (Naithani et al., 2007)
for LORE (see supplementary table 2). SP predictions of *Bra*SRKs were done by SignalP-5.0
server (Almagro Armenteros et al., 2019). MSA of ECDs without SPs was calculated by the
MAFFT (multiple alignment using fast Fourier transformation) algorithm (Katoh et al., 2019) with
default settings with the in-built alignment tool of Jalview. Amino acid sequence alignment of
*Alyr*LORE and *Ahal*LORE cloned from cDNA and the respective sequences from genome
database entries (Fig. S8) was performed using the in-built alignment tool of the SnapGene
software (Insightful Science; available at snapgene.com).

Plant material and cultivation

N. benthamiana, *A. thaliana* Col-0, *A. halleri* (N9852, from P. Falter-Braun, Helmholtz Center
Munich), *A. lyrata* (MN47, from M. Quint, MLU Halle) and *C. rubella* (N22697, from M. Quint, MLU
Halle) were sown on standard potting soil mixed with vermiculite (9:1). All Brassicaceae seeds
were stratified in the dark for at least 48 h at 4°C and then grown under short day conditions (8 h
light, 16 h darkness, 21°C, and 60% relative humidity). *N. benthamiana* was grown under long day
conditions (16 h light, 8 h darkness, 24°C, and 60% humidity). Generation of *A. thaliana* Col-0^{AEQ}
overexpressing *At*LORE (CaMV35S:*LORE*, LORE-OE) was previously described (Shu et al.,

2021). For cytosolic Ca^{2+}-measurements, seeds were surface-sterilized with chlorine gas (4 h)
and seedlings grown in liquid medium (0.5× Murashige & Skoog medium, including vitamins
(Duchefa), 0.25% sucrose, 1 mM MES, pH 5.7) under long day conditions as described above.

Molecular cloning

All plasmids used in this study were cloned using Golden-Gate (GG) (Engler et al., 2009; Weber
et al., 2011) or Gateway™ cloning techniques (Katzen, 2007). Cloning of *At*LORE (AT1G61380),
*At*LORE-Km, *Crub*LORE (CARUB_v10021901mg), *Crub*LORE-ECD, *At*SD1-23i-ECD
(AT1G61390) and apoplastic mCherry was described previously (Ranf et al., 2015; Shu et al.,
2021). *At*GST (AT1G17170) was cloned from pGEX-6P-1. For cloning of *Ahal*LORE
(Araha.6790s0007.1) and *Alyr*LORE (AL2G04470), total RNA was extracted from leaf material of
the respective species by the TRIzol RNA extraction method. RNA samples were treated with
DNAse I (Thermo Fisher Scientific) and reverse-transcribed into cDNA using RevertAid reverse
transcriptase, Ribolock RNase inhibitor and Oligo(dT)$_{18}$ primer, according to manufacturer's
instructions (Thermo Fisher Scientific). Full length coding sequences (CDS) of respective genes
were amplified with specific primers (Supplementary table 1) and ligated into a suitable GG-vector.
Sequences cloned from cDNA of *Ahal*LORE and *Alyr*LORE slightly varied from database entries
(see full sequence alignments figure S8). Conserved ATP binding sites of LORE kinase domains
(K516A, Km, primers see supplementary table 1) were modified by site-directed mutagenesis as
previously described (Ranf et al., 2015) (primers supplementary table 1). For *Ahal*LORE the first
intron of *At*LORE (109 bp) was amplified from genomic DNA and inserted into *Ahal*LORE CDSs
(at the 45th codon) to circumvent problematic read through in *E. coli* or *Agrobacterium tumefaciens*
(primers supplementary table 1). *At*LORE truncations and receptor chimera with *Alyr*LORE were
designed according to domain annotations from Uniprot and Naithani et al. (2007) (see
supplementary table 2). For cloning of truncations and chimera respective parts of the CDS were
amplified by PCR with BpiI linkers from vectors with full length CDS (backbone PCR) and re-
circularized by GG techniques (see Supplementary table 1 and 3). For all membrane bound
truncations, the *At*LORE signal peptide was integrated. CDSs excluding STOP codons were
combined with CaMV35S promotor and terminator, a C-terminal ten-glycine linker and respective
C-terminal epitope tags (GFP, mCherry, SPYCE, SPYNE, hemagglutinin (HA)) using GG
techniques. For FRET-FLIM experiments, CDSs were transferred via Gateway™ cloning (LR-
clonase™, Thermo Fisher Scientific, according to manufacturer's manual) into a final Gateway™
vector containing an estradiol inducible trans-activator cassette (XVE) and C-terminal GFP,
mCherry or GFP-mCherry fusion epitope tags (FLIM-vectors, previously described in (Bleckmann
et al., 2009)). Final expression vectors were transformed into *Agrobacteria tumefaciens* for
transient expression in *planta*.

Transient protein expression in N. benthamiana

N. benthamiana was transiently transformed by *Agrobacterium*-mediated transformation as
described previously (Shu et al., 2021). *Agrobacterium tumefaciens* (strain GV3101 pMP90)
carrying expression constructs were cultivated on LB agar with 30 µg/mL gentamycin, 10 µg/mL
rifampicin, and 50 µg/mL kanamycin, or 100 µg/mL spectinomycin for *Agrobacteria* carrying FLIM
expression constructs. OD$_{600}$ of *Agrobacteria* was adjusted to 0.5 for general purposes or 0.025
for gain-of-function ROS measurement. Prior to infiltration, *Agrobacteria* carrying desired
expression vectors or the silencing suppressor p19 were mixed in a 1:1 ratio. In case of co-

expression, the ratio was adjusted to 1:1:1. For ROS competition assays, the OD_{600} of
Agrobacteria carrying p19 was 0.15, for *At*LORE 0.025 and for *At*LORE-ECD-TMD or cytosolic
mCherry 0.125 (ratio p19:*At*LORE:competing component 6:1:5). Expression of FLIM constructs
driven by the inducible XVE trans-activator was induced by infiltration of 20 µM β-estradiol and
0.1% Tween20 into transformed leaf areas 24 h after infiltration of agrobacteria. Protein
expression was observable 16-24 h after induction.

Protein extraction

Plant material was harvested (60-70 leaf discs of Ø 4 mm), frozen in liquid nitrogen and ground to
fine powder (1 min at 30 Hz in a TissueLyzerII, Qiagen). Total protein was extracted by incubation
with extraction buffer (6 µL/leaf disc, 150 mM Tris-HCl pH 7.5, 150 mM NaCl, 10% glycerol,
1% Nonidet-P40, 10 mM EDTA, 1 mM Na_2MoO_4, 1 mM NaF, 1 mM DTT, 1% (w/v)
polyvinylpyrrolidon, 1% (v/v) protease inhibitor cocktail P9599 (Sigma-Aldrich)) for 1 h at 4°C. After
centrifugation (18000 g, 30 min, 4°C), the supernatant was used for verification of protein
expression via immunoblot or Co-IP experiments.

Co-immunoprecipitation

550 For Co-IPs, protein extracts were incubated for 1-2 h at 4°C with GFP-Trap_MA (magnetic beads,
551 ChromoTek, handling according to manufacturer's manual). Magnetic beads were washed three
552 times with buffer (150 mM Tris-HCL pH 7.5, 150 mM NaCl, 0.5% Nonidet-P40) and re-suspended
553 in 20 µL 1xSDS-sample buffer (5x SDS sample buffer: 60 mM Tris-HCL pH 6.8, 2% SDS,
554 10% glycerol, 5% β-mercaptoethanol, 0.01% bromophenolblue).

Immunoblot

Protein samples in 1x SDS sample buffer were denatured for 10 min at 95 °C and separated by
SDS-polyacrylamide gel electrophoresis (5% stacking gel, 10% resolving gel, 60-100 V,
1x Laemmli running buffer). Proteins were blotted onto a 0.2 µm Protran™ nitrocellulose
membrane (GE healthcare) using 1x transfer buffer (3.03 g/L Tris base, 14.4 g/L glycine, 20%
methanol, 0.05% SDS) and a semi dry transfer cell (Bio-Rad, 1 mA/cm², 1 h). Membrane was
blocked with protein free blocking solution T20 (Pierce) and proteins were detected with respective
antibodies diluted in blocking solution for primary antibodies or 1x TBS-T (6.06 g/L Tris base,
8.76 g/L NaCl, pH 6.7, 0.05% Tween20) for secondary antibodies (1:1000 anti-GFP 3H9
(ChromoTek) with 1:20000 anti-rat-HRP (A9542, Sigma-Aldrich); 1:500 c-Myc 9E10 (Santa Cruz)
with 1:5000 anti-mouse sc-2031 (Santa Cruz); 1:2000 anti-HA-HRP 3F10 (Sigma-Aldrich)). After
washing of membranes (3x 10 min in 1x TBS-T), chemiluminescence was detected with a CCD
camera system (Fusion SL System, Vilber Lourmat GmbH) upon incubation with a peroxidase
substrate (SuperSignal®West Femto Maximum Sensitivity Substrate (Fig. 1A, S2D) or
SuperSignal®West Dura Extended Duration Substrate (Pierce) (Fig. 2B)). When immunoblots
were analyzed with a second, different antibody, membranes were washed (1x TBS-T) after
detection and incubated in stripping buffer (2% SDS, 62.5 mM Tris/HCl pH 6.7, 100mM
β-mercaptoethanol) at 50°C for 30 minutes. Stripped membrane was blocked and immuno-
detected again. Membranes were stained for total protein with amido black (1 g/L amido black,
250 mL/L isopropyl, 100 mL/L acetic acid).

Bimolecular fluorescence complementation

SPYCE/SPYNE epitope tags for BiFCs were previously described (Walter et al., 2004) and adapted for GG cloning. Respective interaction candidates were transiently co-expressed in *N. benthamiana*. YFP fluorescence complementation was visually assessed two days post infiltration via confocal laser scanning microscopy (Leica TCS SP5, ex 514nm, em 525 – 550 nm, Z-stack of 22 planes). For YFP fluorescence quantification, leaf discs (Ø 4 mm) were floated on water in a black 96-well plate and fluorescence was measured with a plate reader (Tecan Infinite F200 PRO, excitation 485 nm, emission 535 nm, 25 flashes, integration time 20 µs, gain set to 70, 2x2 reads per well). Protein expression was validated by immunoblots.

Trypan blue staining

Leaf discs (Ø 2 cm) were harvested five days post transformation and incubated (95°C, 5 min) with 3 mL trypan blue staining solution (25% (v/v) lactic acid, 25% (v/v) phenol, 25% (v/v) glycerol, 25% (v/v) H_2O, 25% (w/v) trypan blue). Leafs were washed several times with destaining solution (250 g chloral hydrate in 100 mL H_2O) and rinsed with water before photo-documentation.

ROS measurement

Accumulation of ROS was measured with a luminol based reporter system and a microplate reader (Tecan Infinite F200 PRO or Luminoscan Ascent 2.1, Thermo Fisher Scientific) as describe previously (Ranf et al., 2015). Leaf discs (Ø 4 mm) of Brassicaceae (6-8 weeks old) or transiently transformed *N. benthamiana* (36 h post agro-infiltration) were floated on water in white 96-well plates for at least 6h (for *N. benthamiana*) or overnight (for Brassicaceae). Prior to measurements, water was replaced with 100 µL horseradish peroxidase-luminol solution (2 µg/mL horseradish peroxidase, 10 µM L-012 (WAKO Chemicals GmbH)). After addition of elicitors or respective controls (3-OH-C10:0 (Matreya LLC) in MeOH, flg22 (QRLSTGSRINSAKDDAAGLQIA in ddH₂O), luminescence was recorded in relative light units (RLU) in 1 minute intervals for up to 60 minutes. Total ROS accumulation was summed up for elicitor and control treatments.

Ligand depletion assay

Ligand depletion assay was performed as previously described (Shu et al., 2021). ECDs of respective proteins were transiently expressed in *N. benthamiana* apoplasts and harvested via apoplastic washing fluids (AWF). AWFs were concentrated and incubated with 3-OH-C10:0 (9:1, v/v). Unbound elicitor was separated by filtration (30 kDa molecular weight cut-off). Filtrates were used for elicitation of cytosolic calcium signaling in LORE-OE reporter lines as described before (Ranf et al., 2015; Shu et al., 2021). For each time point, luminescence was normalized to total luminescence counts remaining (L/L_{max}) and total accumulation of calcium signals was summed up from 3 to 30 minutes. 5 µL of concentrated AWF with a total protein concentration of 1.5 mg/mL was analyzed for protein expression by immunoblot.

Red light chlorophyll fluorescence measurement for cell death quantification

Red light emission of chlorophyll florescence can be measured for quantification of cell death (Landeo Villanueva et al., 2021). Leaf discs (Ø 4 mm) of transiently transformed *N. benthamiana* were floated on water in black 96-well plates and chlorophyll fluorescence was measured with a plate reader (Tecan Infinite F200 PRO, excitation 535 nm, emission 590 nm, 25 flashes, integration time 20 µs, 4x4 reads per well, gain set to 80) as relative fluorescence units (RFU). Values of all reads per well were summed up.

Microscopy

Confocal laser scanning microscopy was performed using either a Leica TCS SP5 (Argon laser, HyD2 detectors) or an Olympus FV3000 system (diode lasers, PMT detectors). Images were acquired with an image size of 512x512 pixels. In case of co-expression analysis, images were taken with a sequential scan to avoid 'bleed through'. GFP was excited at 488 nm and emission detected at 500-540 nm. mCherry was excited at 561 nm and emission detected at 570-620 nm. YFP was excited at 514 nm and emission detected at 525-550 nm. Images were imported and processed with OMERO (Allan et al., 2012).

Fluorescence lifetime imaging (FLIM)

GFP fluorescence lifetimes were measured via time-correlated single-photon counting (TCSPC) by an Olympus FV3000 system linked to a PicoQuant FCS/FLIM-FRET/rapidFLIM upgrade kit, based on previously described protocols (Weidtkamp-Peters and Stahl, 2017). TCSPC was performed with a 485 nm (LDH-D-C-485) pulsed laser, two TCSPC modules (TimeHarp 260 PICO Dual, TimeHarp 260 NANO Dual) and two photon counting PMA hybrid 40 detectors. Co-expressing cells were detected by confocal imaging via an Olympus FV3000 60x water immersion objective (UPLSAPO60XW 60x/NA 1.2/WD 0.28) and a selected measuring area was magnified (4x zoom). TCSPC was performed with a laser pulse rate of 40.00 MHz, a TCSPC resolution of 25.0 ps and an image size of 512x512 pixels. 500-1000 photon counts per pixel were acquired for each image. FLIM was analyzed by the Symphotime64 software of PicoQuant via n-exponential reconvolution and an internally calculated instrument response function (IRF). A two-parameter fitting (n=2) was performed in most cases. Only FLIM analysis with fitting coefficients (χ^2) between 1.0 and 2.0 were accepted. Intensity weighted average lifetimes т of membrane ROIs were determined for each FLIM image.

 FIGURES

A

B

C

D

Figure 1 *At*LORE forms homomers and has an auto-immunity phenotype in *N. benthamiana.* A Chlorophyll fluorescence measurements (relative fluorescence units, RFU) in *N. benthamiana* upon transient expression of candidates four days post transformation. Pooled data of two technical replicates from one biological replicate are shown. Data show mean with SD. n=32 leaf discs. Statistics analyzed by one-way ANOVA with Tukey's multiple comparisons test, α=0.01. Data not sharing the same letter are significantly different. **B** Anti-GFP and anti-mCherry immunoblot of co-immunoprecipitation (GFP-trap) after transient co-expression of interaction candidates in *N. benthamiana.* Total protein was stained with amido black. Km, kinase mutated (K516A). **C** FRET-FLIM of *At*LORE-Km-GFP and indicated candidates fused to an mCherry epitope tag, transiently co-expressed in *N. benthamiana.* Pooled data of five independent biological replicates. Data show mean with SD. n, number of analyzed cells. Statistics analyzed by one-way ANOVA with Tukey's multiple comparisons test, α=0.01. Data not sharing the same letter are significantly different. **D** FRET-FLIM of *At*LORE-Km-GFP with *At*LORE-mCherry, transiently co-expressed in *N. benthamiana* and treated with 5 µM 3-OH-C10:0 or H_2O as control. FLIM images were acquired 10-20 minutes after treatment. Pooled data from two independent biological replicates. n, number of analyzed cells. Data show mean with SD. Differences between treatments were analyzed by two-way ANOVA with Sidak's multiple comparisons test, α=0.01.

666
Figure 2 Homomerization of *At*LORE is mediated by the extracellular and trans-membrane
domain. A Scheme of *At*LORE truncation variants; ECD, extracellular domain; TMD,
transmembrane domain; ICD, intracellular domain. Created with BioRender.com. **B** Anti-GFP and
anti-mCherry immunoblot of co-immunoprecipitation (GFP-trap) after transient co-expression of
full length and truncated *At*LORE-Km interaction candidates in *N. benthamiana*. Total protein is
stained with amido black. **C, D** FRET-FLIM of *At*LORE-Km-GFP full length or *At*LORE-ECD-TMD-
GFP versus truncated *At*LORE-mCherry variants transiently expressed in *N. benthamiana*. Pooled
data from two independent biological replicates each. Data show mean with SD. n, number of
analyzed cells. Statistics analyzed by one-way ANOVA with Tukey's multiple comparisons test,
α=0.01, data not sharing the same letter are significantly different.

Figure 3 3-OH-C10:0 responsiveness of *A. lyrata*, *A. halleri* and *C. rubella* or *N. benthamiana*
expressing putative LORE orthologues. A Total ROS accumulation of *A. halleri*, *C. rubella*,
A. lyrata and *A. thaliana* leaf discs treated with 5 µM 3-OH-C10:0 or MeOH as control. Median
with minimum to maximum of total ROS between 3-60 minutes after elicitation is shown; n=8 leaf
discs. Significance between MeOH and 3-OH-C10:0 treatment tested with two-way ANOVA with
Sidak's multiple comparisons test, α=0.01. **B** Total ROS accumulation of *N. benthamiana* leaf
discs transiently overexpressing *At*LORE or putative LORE orthologues upon application of 1 µM
3-OHC10:0 or MeOH as control. Boxplot displays minimum to maximum with median of total ROS
between 3-45 minutes after treatment. n, number of leaf discs. Combined data of two independent
datasets from one biological replicate are shown. For *At*LORE and *At*LORE-Km data of both
datasets were pooled. Statistics was analyzed by two-way ANOVA with Sidak's multiple
comparisons test, α= 0.01.

Figure 4 The ECD of *Alyr*LORE is causal for defective 3-OH-C10:0 perception, but ligand
binding of all orthologues is functional. A Total ROS accumulation of *At*LORE, *Alyr*LORE and
chimera with either ECD-TMD or ICD domain swap (DS) elicited with 10 µM 3-OH-C10:0 or MeOH.
Median with minimum to maximum is shown; n=12 leaf discs. Statistics was analyzed by two-way
ANOVA with Sidak's multiple comparisons test, α= 0.01. **B** 3-OH-C10:0 binding to LORE-ECDs
was tested in a ligand depletion assay. Unbound 3-OH-C10:0 in filtrates was detected by cytosolic
Ca^{2+} measurements of LORE-OE lines. Ligand binding is indicated by full depletion of a Ca^{2+}
response. Data show mean and SD of total $[Ca^{2+}]_{cyt}$ from 3 to 30 minutes; n=3 seedlings.

Figure 5 Putative LORE orthologues from 3-OH-C10:0 unresponsive species are impaired in homomerization. A Chlorophyll fluorescence measurement of *N. benthamiana* leaves transiently overexpressing *At*LORE, *Alyr*LORE, *Ahal*LORE and *Crub*LORE. Mean and SD of pooled data from two technical replicates are shown. n=32 leave discs. Statistics analyzed by one-way ANOVA with Tukey's multiple comparisons test, α=0.01. Data not sharing data are significantly different. Experiment was repeated two times with similar outcome. **B-D** FRET-FLIM of putative LORE orthologous from *A. lyrata* (**B**) *A. halleri* (**C**) and *C. rubella* (**D**) transiently expressed in *N. benthamiana*. Km, kinase mutated; i, integrated LORE intron; n, number of analyzed cells, r, number of biological replicates. Data show mean with SD. Statistics analyzed by one-way ANOVA with Tukey's multiple comparisons test, α=0.01. Data not sharing the same letters are significantly different.

Figure 6 Mapping of dimerization region using receptor chimera of *At*LORE and *Alyr*LORE.
A Schematic overview of chimera between *Alyr*LORE and *At*LORE. DS, domain swap; LL, lectin domain 1 and lectin domain 2; EP, EGF and PAN domain. Created with BioRender.com. **B** FRET-FLIM of *At*LORE-*Alyr*LORE domain swaps (DSAlyr) transiently expressed in *N. benthamiana*. Mean and SD of pooled data are shown; r, number of biological replicates; n, number of analyzed cells. Statistics analyzed by one-Way ANOVA with Tukey's multiple comparisons test, α=0.01. Data not sharing the same letter are significantly different. **C** Gain of function ROS measurements of *At*LORE-*Alyr*LORE chimera transiently expressed in *N. benthamiana* upon elicitation with 10 µM 3-OH-C10:0 or MeOH as control. Median with minimum to maximum is shown; n=12 leaf discs. Statistics was analyzed by two-way ANOVA with Sidak's multiple comparisons test, α= 0.01. **D** Chlorophyll fluorescence measurements (in relative fluorescence units, RFU) in *N. benthamiana* upon transient expression of *At*LORE-*Alyr*LORE chimera four days post transformation. Mean and SD of pooled data from two technical replicates of one biological replicate are shown. n, number of leaf discs. Statistics analyzed by one-way ANOVA with Tukey's multiple comparisons test, α=0.01. Data not sharing the same letter are significantly different.

Figure 7 *At*LORE homomerization is essential for downstream signaling. Dominant negative effects of co-expressed *At*LORE-ECD-TMD-HA or cytosolic mCherry was analyzed by competitive ROS accumulation measurement in *N. benthamiana* transiently co-expressing indicated candidates upon elicitation with 5 µM 3-OH-C10:0 or MeOH as a control. Median with minimum to maximum of total ROS between 3 and 45 min from three technical replicates is shown; n=12 leaf discs. Differences between treatments was analyzed by two-way ANOVA and Sidak's multiple comparisons test (α=0.01).

ACKNOWLEGEMENTS

We acknowledge the Center for Advanced Light Microscopy (CALM) of imaging@TUM at the TUM School of Life Sciences for providing access to FLIM microscopes used in this study, Pascal Falter-Braun (Helmholtz center, Munich) and Marcel Quint (MLU Halle-Wittenberg) for providing plant material and Yvonne Stahl (HHU Düsseldorf) for providing FLIM vectors. Graphs and statistics were created by GraphPad Prism, microscopy images by OMERO and schemes by BioRender.com. We thank Ralph Hückelhoven and Martin Stegmann for critical discussion of data. Research in the Ranf lab is supported by the German Research Foundation (SFB924/TP-B10) and Emmy Noether program (RA2541/1).

REFERENCES

Allan, C., Burel, J.-M., Moore, J., Blackburn, C., Linkert, M., Loynton, S., MacDonald, D., Moore, W.J., Neves, C., Patterson, A., Porter, M., Tarkowska, A., Loranger, B., Avondo, J., Lagerstedt, I., Lianas, L., Leo, S., Hands, K., Hay, R.T., Patwardhan, A., Best, C., Kleywegt, G.J., Zanetti, G., and Swedlow, J.R. (2012). OMERO: flexible, model-driven data management for experimental biology. Nature Methods **9,** 245-253.

Almagro Armenteros, J.J., Tsirigos, K.D., Sønderby, C.K., Petersen, T.N., Winther, O., Brunak, S., von Heijne, G., and Nielsen, H. (2019). SignalP 5.0 improves signal peptide predictions using deep neural networks. Nature biotechnology **37,** 420-423.

Balint-Kurti, P. (2019). The plant hypersensitive response: concepts, control and consequences. Molecular Plant Pathology **20,** 1163-1178.

Bauer, Z., Gómez-Gómez, L., Boller, T., and Felix, G. (2001). Sensitivity of Different Ecotypes and Mutants ofArabidopsis thaliana toward the Bacterial Elicitor Flagellin Correlates with the Presence of Receptor-binding Sites *. Journal of Biological Chemistry **276,** 45669-45676.

Bi, G., Liebrand, T.W.H., Bye, R.R., Postma, J., Burgh, A.M., Robatzek, S., Xu, X., and Joosten, M.H.A.J. (2015). SOBIR1 requires the GxxxG dimerization motif in its transmembrane domain to form constitutive complexes with receptor-like proteins. Molecular Plant Pathology **17,** 96-107.

Bleckmann, A., Weidtkamp-Peters, S., Seidel, C., and Simon, R. (2009). Stem Cell Signaling in Arabidopsis Requires CRN to Localize CLV2 to the Plasma Membrane. Plant physiology **152**.

Bojar, D., Martinez, J., Santiago, J., Rybin, V., Bayliss, R., and Hothorn, M. (2014). Crystal structures of the phosphorylated BRI1 kinase domain and implications for brassinosteroid signal initiation. The Plant journal : for cell and molecular biology **78,** 31-43.

Boller, T., and Felix, G. (2009). A Renaissance of Elicitors: Perception of Microbe-Associated Molecular Patterns and Danger Signals by Pattern-Recognition Receptors. Annual Review of Plant Biology **60,** 379-406.

Boutrot, F., and Zipfel, C. (2017). Function, Discovery, and Exploitation of Plant Pattern Recognition Receptors for Broad-Spectrum Disease Resistance. Annual review of phytopathology **55,** 257-286.

Burkart, R.C., and Stahl, Y. (2017). Dynamic complexity: plant receptor complexes at the plasma membrane. Curr Opin Plant Biol **40,** 15-21.

Cao, Y., Liang, Y., Tanaka, K., Nguyen, C.T., Jedrzejczak, R.P., Joachimiak, A., and Stacey, G. (2014). The kinase LYK5 is a major chitin receptor in Arabidopsis and forms a chitin-induced complex with related kinase CERK1. eLife **3,** e03766.

Chen, L.J., Wuriyanghan, H., Zhang, Y.Q., Duan, K.X., Chen, H.W., Li, Q.T., Lu, X., He, S.J., Ma, B., Zhang, W.K., Lin, Q., Chen, S.Y., and Zhang, J.S. (2013). An S-domain

receptor-like kinase, OsSIK2, confers abiotic stress tolerance and delays dark-induced
leaf senescence in rice. Plant Physiol **163,** 1752-1765.
Chen, X., Shang, J., Chen, D., Lei, C., Zou, Y., Zhai, W., Liu, G., Xu, J., Ling, Z., Cao, G., Ma,
B., Wang, Y., Zhao, X., Li, S., and Zhu, L. (2006). A B-lectin receptor kinase gene
conferring rice blast resistance. The Plant Journal **46,** 794-804.
Cheng, X., Wu, Y., Guo, J., Du, B., Chen, R., Zhu, L., and He, G. (2013). A rice lectin receptor-
like kinase that is involved in innate immune responses also contributes to seed
germination. The Plant Journal **76,** 687-698.
Chinchilla, D., Bauer, Z., Regenass, M., Boller, T., and Felix, G. (2006). The Arabidopsis
receptor kinase FLS2 binds flg22 and determines the specificity of flagellin perception.
The Plant cell **18,** 465-476.
Chinchilla, D., Zipfel, C., Robatzek, S., Kemmerling, B., Nürnberger, T., Jones, J.D., Felix,
G., and Boller, T. (2007). A flagellin-induced complex of the receptor FLS2 and BAK1
initiates plant defence. Nature **448,** 497-500.
Consortium, T.U. (2020). UniProt: the universal protein knowledgebase in 2021. Nucleic acids
research **49,** D480-D489.
Couto, D., and Zipfel, C. (2016). Regulation of pattern recognition receptor signalling in plants.
Nature reviews. Immunology **16,** 537-552.
De Smet, I., Voß, U., Jürgens, G., and Beeckman, T. (2009). Receptor-like kinases shape the
plant. Nature Cell Biology **11,** 1166-1173.
Dievart, A., Gottin, C., Périn, C., Ranwez, V., and Chantret, N. (2020). Origin and Diversity of
Plant Receptor-Like Kinases. Annual Review of Plant Biology **71,** 131-156.
Engler, C., Gruetzner, R., Kandzia, R., and Marillonnet, S. (2009). Golden Gate Shuffling: A
One-Pot DNA Shuffling Method Based on Type IIs Restriction Enzymes. PLOS ONE **4,**
e5553.
Fan, J., Bai, P., Ning, Y., Wang, J., Shi, X., Xiong, Y., Zhang, K., He, F., Zhang, C., Wang,
R., Meng, X., Zhou, J., Wang, M., Shirsekar, G., Park, C.H., Bellizzi, M., Liu, W.,
Jeon, J.S., Xia, Y., Shan, L., and Wang, G.L. (2018). The Monocot-Specific Receptor-
like Kinase SDS2 Controls Cell Death and Immunity in Rice. Cell host & microbe **23,** 498-
510.e495.
Fink, A., Sal-Man, N., Gerber, D., and Shai, Y. (2012). Transmembrane domains interactions
within the membrane milieu: Principles, advances and challenges. Biochimica et
Biophysica Acta (BBA) - Biomembranes **1818,** 974-983.
Gilardoni, P.A., Hettenhausen, C., Baldwin, I.T., and Bonaventure, G. (2011). Nicotiana
attenuata LECTIN RECEPTOR KINASE1 suppresses the insect-mediated inhibition of
induced defense responses during Manduca sexta herbivory. The Plant cell **23,** 3512-
3532.
Giranton, J.L., Dumas, C., Cock, J.M., and Gaude, T. (2000). The integral membrane S-locus
receptor kinase of Brassica has serine/threonine kinase activity in a membranous
environment and spontaneously forms oligomers in planta. Proceedings of the National
Academy of Sciences of the United States of America **97,** 3759-3764.
Gomez-Gomez, L., and Boller, T. (2000). FLS2: an LRR receptor-like kinase involved in the
perception of the bacterial elicitor flagellin in Arabidopsis. Molecular cell **5,** 1003-1011.
Gong, B.-Q., Wang, F.-Z., and Li, J.-F. (2020). Hide-and-Seek: Chitin-Triggered Plant Immunity
and Fungal Counterstrategies. Trends in plant science **25,** 805-816.
Gou, X., and Li, J. (2020). Paired Receptor and Coreceptor Kinases Perceive Extracellular
Signals to Control Plant Development. Plant physiology **182,** 1667-1681.
Guo, X., Ye, J., Li, M., Lin, X., Lu, X., Liu, S., Li, H., and Zhang, C. (2018). LecRKIII.1 and
LecRKIII.2 formed homodimers to play physiological functions in Arabidopsis thaliana.
SDRP Journal of Plant Science **2,** 1-8.
Herrmann, J.R., Panitz, J.C., Unterreitmeier, S., Fuchs, A., Frishman, D., and Langosch, D.
(2009). Complex Patterns of Histidine, Hydroxylated Amino Acids and the GxxxG Motif

Mediate High-affinity Transmembrane Domain Interactions. Journal of Molecular Biology
385, 912-923.

Hohmann, U., Lau, K., and Hothorn, M. (2017). The Structural Basis of Ligand Perception and
Signal Activation by Receptor Kinases. Annu Rev Plant Biol **68,** 109-137.

Ivanov, R., Fobis-Loisy, I., and Gaude, T. (2010). When no means no: guide to Brassicaceae
self-incompatibility. Trends in plant science **15,** 387-394.

Jamieson, P.A., Shan, L., and He, P. (2018). Plant cell surface molecular cypher: Receptor-like
proteins and their roles in immunity and development. Plant Sci **274,** 242-251.

Jany, E., Nelles, H., and Goring, D.R. (2019). Chapter One - The Molecular and Cellular
Regulation of Brassicaceae Self-Incompatibility and Self-Pollen Rejection. In
International Review of Cell and Molecular Biology, L. Galluzzi, ed (Academic Press), pp.
1-35.

Jinjun, Z., Peina, J., Fang, Z., Chongke, Z., Bo, B., Yaping, L., Haifeng, W., Fan, C., and
Xianzhi, X. (2020). OsSRK1, an Atypical S-Receptor-Like Kinase Positively Regulates
Leaf Width and Salt Tolerance in Rice. Rice Science **27,** 133-142.

Kanzaki, H., Saitoh, H., Takahashi, Y., Berberich, T., Ito, A., Kamoun, S., and Terauchi, R.
(2008). NbLRK1, a lectin-like receptor kinase protein of Nicotiana benthamiana, interacts
with Phytophthora infestans INF1 elicitin and mediates INF1-induced cell death. Planta
228, 977-987.

Katoh, K., Rozewicki, J., and Yamada, K.D. (2019). MAFFT online service: multiple sequence
alignment, interactive sequence choice and visualization. Briefings in Bioinformatics **20,**
1160-1166.

Katzen, F. (2007). Gateway((R)) recombinational cloning: a biological operating system. Expert
opinion on drug discovery **2,** 571-589.

Kim, H.S., Jung, M.S., Lee, S.M., Kim, K.E., Byun, H., Choi, M.S., Park, H.C., Cho, M.J., and
Chung, W.S. (2009a). An S-locus receptor-like kinase plays a role as a negative
regulator in plant defense responses. Biochem Biophys Res Commun **381,** 424-428.

Kim, H.S., Jung, M.S., Lee, K., Kim, K.E., Yoo, J.H., Kim, M.C., Kim, D.H., Cho, M.J., and
Chung, W.S. (2009b). An S-locus receptor-like kinase in plasma membrane interacts
with calmodulin in Arabidopsis. FEBS Lett **583,** 36-42.

Kutschera, A., Dawid, C., Gisch, N., Schmid, C., Raasch, L., Gerster, T., Schaffer, M.,
Smakowska-Luzan, E., Belkhadir, Y., Vlot, A.C., Chandler, C.E., Schellenberger, R.,
Schwudke, D., Ernst, R.K., Dorey, S., Huckelhoven, R., Hofmann, T., and Ranf, S.
(2019). Bacterial medium-chain 3-hydroxy fatty acid metabolites trigger immunity in
Arabidopsis plants. Science **364,** 178-181.

Labbé, J., Muchero, W., Czarnecki, O., Wang, J., Wang, X., Bryan, A.C., Zheng, K., Yang,
Y., Xie, M., Zhang, J., Wang, D., Meidl, P., Wang, H., Morrell-Falvey, J.L., Cope, K.R.,
Maia, L.G.S., Ané, J.-M., Mewalal, R., Jawdy, S.S., Gunter, L.E., Schackwitz, W.,
Martin, J., Le Tacon, F., Li, T., Zhang, Z., Ranjan, P., Lindquist, E., Yang, X.,
Jacobson, D.A., Tschaplinski, T.J., Barry, K., Schmutz, J., Chen, J.-G., and Tuskan,
G.A. (2019). Mediation of plant–mycorrhizal interaction by a lectin receptor-like kinase.
Nature Plants **5,** 676-680.

Landeo Villanueva, S., Malvestiti, M.C., van Ieperen, W., Joosten, M.H.A.J., and van Kan,
J.A.L. (2021). Red light imaging for programmed cell death visualization and
quantification in plant–pathogen interactions. Molecular Plant Pathology **22,** 361-372.

Langosch, D., and Arkin, I.T. (2009). Interaction and conformational dynamics of membrane-
spanning protein helices. Protein Sci **18,** 1343-1358.

Li, H., and Yang, W.-C. (2016). RLKs orchestrate the signaling in plant male-female interaction.
Science China Life Sciences **59,** 867-877.

Li, J.B., Sun, Y.D., Liu, H., Wang, Y.Y., Jia, Y.L., and Xu, M.H. (2015). Natural variation of rice
blast resistance gene Pi-d2. Genetics and molecular research : GMR **14,** 1235-1249.

Liu, C., Dong, X., Xu, Y., Dong, Q., Wang, Y., Gai, Y., and Ji, X. (2021). Transcriptome and
DNA Methylome Reveal Insights Into Phytoplasma Infection Responses in Mulberry
(Morus multicaulis Perr.). Frontiers in Plant Science **12**.
Liu, T., Liu, Z., Song, C., Hu, Y., Han, Z., She, J., Fan, F., Wang, J., Jin, C., Chang, J., Zhou,
J.-M., and Chai, J. (2012). Chitin-Induced Dimerization Activates a Plant Immune
Receptor. Science **336,** 1160.
Luo, X., Wu, W., Liang, Y., Xu, N., Wang, Z., Zou, H., and Liu, J. (2020). Tyrosine
phosphorylation of the lectin receptor-like kinase LORE regulates plant immunity. The
EMBO journal **n/a,** e102856.
Ma, R., Han, Z., Hu, Z., Lin, G., Gong, X., Zhang, H., Nasrallah, J.B., and Chai, J. (2016a).
Structural basis for specific self-incompatibility response in Brassica. Cell Res **26,** 1320-
1329.
Ma, X., Xu, G., He, P., and Shan, L. (2016b). SERKing Coreceptors for Receptors. Trends in
plant science **21,** 1017-1033.
Macho, A.P., Lozano-Durán, R., and Zipfel, C. (2015). Importance of tyrosine phosphorylation
in receptor kinase complexes. Trends in plant science **20,** 269-272.
Miya, A., Albert, P., Shinya, T., Desaki, Y., Ichimura, K., Shirasu, K., Narusaka, Y.,
Kawakami, N., Kaku, H., and Shibuya, N. (2007). CERK1, a LysM receptor kinase, is
essential for chitin elicitor signaling in *Arabidopsis*. Proceedings of the National Academy
of Sciences **104,** 19613-19618.
Murase, K., Moriwaki, Y., Mori, T., Liu, X., Masaka, C., Takada, Y., Maesaki, R., Mishima,
M., Fujii, S., Hirano, Y., Kawabe, Z., Nagata, K., Terada, T., Suzuki, G., Watanabe,
M., Shimizu, K., Hakoshima, T., and Takayama, S. (2020). Mechanism of self/nonself-
discrimination in Brassica self-incompatibility. Nature Communications **11,** 4916.
Naithani, S., Chookajorn, T., Ripoll, D.R., and Nasrallah, J.B. (2007). Structural modules for
receptor dimerization in the *S*-locus receptor kinase extracellular domain. Proceedings of
the National Academy of Sciences **104,** 12211-12216.
Nasrallah, J.B., and Nasrallah, M.E. (2014). S-locus receptor kinase signalling. Biochemical
Society transactions **42,** 313-319.
Navarro-Gochicoa, M.-T., Camut, S., Timmers, A.C.J., Niebel, A., Herve, C., Boutet, E.,
Bono, J.-J., Imberty, A., and Cullimore, J.V. (2003). Characterization of four lectin-like
receptor kinases expressed in roots of Medicago truncatula. Structure, location,
regulation of expression, and potential role in the symbiosis with Sinorhizobium meliloti.
Plant physiology **133,** 1893-1910.
Pan, J., Li, Z., Wang, Q., Yang, L., Yao, F., and Liu, W. (2020). An S-domain receptor-like
kinase, OsESG1, regulates early crown root development and drought resistance in rice.
Plant Sci **290,** 110318.
Pietraszewska-Bogiel, A., Lefebvre, B., Koini, M.A., Klaus-Heisen, D., Takken, F.L.W.,
Geurts, R., Cullimore, J.V., and Gadella, T.W.J. (2013). Interaction of Medicago
truncatula Lysin Motif Receptor-Like Kinases, NFP and LYK3, Produced in Nicotiana
benthamiana Induces Defence-Like Responses. PLOS ONE **8,** e65055.
Ranf, S., Gisch, N., Schaffer, M., Illig, T., Westphal, L., Knirel, Y.A., Sanchez-Carballo, P.M.,
Zahringer, U., Huckelhoven, R., Lee, J., and Scheel, D. (2015). A lectin S-domain
receptor kinase mediates lipopolysaccharide sensing in Arabidopsis thaliana. Nat
Immunol **16,** 426-433.
Robatzek, S., and Wirthmueller, L. (2013). Mapping FLS2 function to structure: LRRs, kinase
and its working bits. Protoplasma **250,** 671-681.
Roux, M., Schwessinger, B., Albrecht, C., Chinchilla, D., Jones, A., Holton, N., Malinovsky,
F.G., Tör, M., de Vries, S., and Zipfel, C. (2011). The Arabidopsis leucine-rich repeat
receptor-like kinases BAK1/SERK3 and BKK1/SERK4 are required for innate immunity to
hemibiotrophic and biotrophic pathogens. The Plant cell **23,** 2440-2455.

Schnepf, V., Vlot, A.C., Kugler, K., and Hückelhoven, R. (2018). Barley susceptibility factor
RACB modulates transcript levels of signalling protein genes in compatible interaction
with Blumeria graminis f.sp. hordei. Molecular plant pathology **19,** 393-404.
Shimosato, H., Yokota, N., Shiba, H., Iwano, M., Entani, T., Che, F.-S., Watanabe, M.,
Isogai, A., and Takayama, S. (2007). Characterization of the SP11/SCR high-affinity
binding site involved in self/nonself recognition in brassica self-incompatibility. The Plant
cell **19,** 107-117.
Shiu, S.H., and Bleecker, A.B. (2001). Plant receptor-like kinase gene family: diversity,
function, and signaling. Science's STKE : signal transduction knowledge environment
2001, re22.
Shiu, S.H., and Bleecker, A.B. (2003). Expansion of the receptor-like kinase/Pelle gene family
and receptor-like proteins in Arabidopsis. Plant Physiol **132,** 530-543.
Shiu, S.H., Karlowski, W.M., Pan, R., Tzeng, Y.H., Mayer, K.F., and Li, W.H. (2004).
Comparative analysis of the receptor-like kinase family in Arabidopsis and rice. The Plant
cell **16,** 1220-1234.
Shu, L.-J., Schäffer, M., Eschrig, S., and Ranf, S. (2021). Low cost, medium throughput
depletion-binding assay for screening S-domain-receptor ligand interactions using *in*
planta protein expression. bioRxiv, 2021.2006.2016.448648.
Somssich, M., Ma, Q., Weidtkamp-Peters, S., Stahl, Y., Felekyan, S., Bleckmann, A.,
Seidel, C.A.M., and Simon, R. (2015). Real-time dynamics of peptide ligand–dependent
receptor complex formation in planta. Science Signaling **8,** ra76-ra76.
Stahl, Y., Grabowski, S., Bleckmann, A., Kühnemuth, R., Weidtkamp-Peters, S., Pinto,
Karine G., Kirschner, Gwendolyn K., Schmid, Julia B., Wink, René H., Hülsewede,
A., Felekyan, S., Seidel, Claus A.M., and Simon, R. (2013). Moderation of Arabidopsis
Root Stemness by CLAVATA1 and ARABIDOPSIS CRINKLY4 Receptor Kinase
Complexes. Current Biology **23,** 362-371.
Stokes, K.D., and Gururaj Rao, A. (2008). Dimerization properties of the transmembrane
domains of Arabidopsis CRINKLY4 receptor-like kinase and homologs. Archives of
biochemistry and biophysics **477,** 219-226.
Sun, M., Qian, X., Chen, C., Cheng, S., Jia, B., Zhu, Y., and Sun, X. (2018). Ectopic
Expression of GsSRK in Medicago sativa Reveals Its Involvement in Plant Architecture
and Salt Stress Responses. Frontiers in Plant Science **9.**
Sun, W., Cao, Y., Jansen Labby, K., Bittel, P., Boller, T., and Bent, A.F. (2012). Probing the
Arabidopsis flagellin receptor: FLS2-FLS2 association and the contributions of specific
domains to signaling function. The Plant cell **24,** 1096-1113.
Sun, X.L., Yu, Q.Y., Tang, L.L., Ji, W., Bai, X., Cai, H., Liu, X.F., Ding, X.D., and Zhu, Y.M.
(2013a). GsSRK, a G-type lectin S-receptor-like serine/threonine protein kinase, is a
positive regulator of plant tolerance to salt stress. Journal of plant physiology **170,** 505-
515.
Sun, Y., Qiao, Z., Muchero, W., and Chen, J.-G. (2020). Lectin Receptor-Like Kinases: The
Sensor and Mediator at the Plant Cell Surface. Frontiers in plant science **11,** 596301-
596301.
Sun, Y., Li, L., Macho, A.P., Han, Z., Hu, Z., Zipfel, C., Zhou, J.-M., and Chai, J. (2013b).
Structural Basis for flg22-Induced Activation of the *Arabidopsis* FLS2-BAK1 Immune
Complex. Science **342,** 624-628.
Takayama, S., Shimosato, H., Shiba, H., Funato, M., Che, F.-S., Watanabe, M., Iwano, M.,
and Isogai, A. (2001). Direct ligand–receptor complex interaction controls Brassica self-
incompatibility. Nature **413,** 534-538.
Teixeira, M.A., Rajewski, A., He, J., Castaneda, O.G., Litt, A., and Kaloshian, I. (2018).
Classification and phylogenetic analyses of the Arabidopsis and tomato G-type lectin
receptor kinases. BMC Genomics **19,** 239.

Vaid, N., Pandey, P.K., and Tuteja, N. (2012). Genome-wide analysis of lectin receptor-like
kinase family from Arabidopsis and rice. Plant molecular biology **80,** 365-388.
Walter, M., Chaban, C., Schütze, K., Batistic, O., Weckermann, K., Näke, C., Blazevic, D.,
Grefen, C., Schumacher, K., Oecking, C., Harter, K., and Kudla, J. (2004).
Visualization of protein interactions in living plant cells using bimolecular fluorescence
complementation. The Plant Journal **40,** 428-438.
Wan, W.-L., Fröhlich, K., Pruitt, R.N., Nürnberger, T., and Zhang, L. (2019). Plant cell surface
immune receptor complex signaling. Current Opinion in Plant Biology **50,** 18-28.
Wang, Y., Cordewener, J.H.G., America, A.H.P., Shan, W., Bouwmeester, K., and Govers,
F. (2015). Arabidopsis Lectin Receptor Kinases LecRK-IX.1 and LecRK-IX.2 Are
Functional Analogs in Regulating Phytophthora Resistance and Plant Cell Death.
Molecular Plant-Microbe Interactions® **28,** 1032-1048.
Waterhouse, A.M., Procter, J.B., Martin, D.M., Clamp, M., and Barton, G.J. (2009). Jalview
Version 2--a multiple sequence alignment editor and analysis workbench. Bioinformatics
(Oxford, England) **25,** 1189-1191.
Weber, E., Engler, C., Gruetzner, R., Werner, S., and Marillonnet, S. (2011). A Modular
Cloning System for Standardized Assembly of Multigene Constructs. PLOS ONE **6,**
e16765.
Weidtkamp-Peters, S., and Stahl, Y. (2017). The Use of FRET/FLIM to Study Proteins
Interacting with Plant Receptor Kinases. Methods in molecular biology (Clifton, N.J.)
1621, 163-175.
Westerfield, J.M., and Barrera, F.N. (2020). Membrane receptor activation mechanisms and
transmembrane peptide tools to elucidate them. The Journal of biological chemistry **295,**
1792-1814.
Xing, S., Li, M., and Liu, P. (2013). Evolution of S-domain receptor-like kinases in land plants
and origination of S-locus receptor kinases in Brassicaceae. BMC Evolutionary Biology
13, 69.
Xing, S., Wallmeroth, N., Berendzen, K.W., and Grefen, C. (2016). Techniques for the
Analysis of Protein-Protein Interactions in Vivo. Plant Physiology **171,** 727-758.
Xue, D.-X., Li, C.-L., Xie, Z.-P., and Staehelin, C. (2019). LYK4 is a component of a tripartite
chitin receptor complex in Arabidopsis thaliana. Journal of experimental botany **70,** 5507-
5516.
Yu, X., Feng, B., He, P., and Shan, L. (2017). From Chaos to Harmony: Responses and
Signaling upon Microbial Pattern Recognition. Annual review of phytopathology **55,** 109-
137.
Zhou, D., Godinez-Vidal, D., He, J., Teixeira, M., Guo, J., Wei, L., Van Norman, J.M., and
Kaloshian, I. (2021). A G-lectin Receptor Kinase is a Negative Regulator of Arabidopsis
Immunity Against Root-Knot Nematode Meloidogyne incognita. bioRxiv,
2021.2009.2007.459316.
Zou, X., Qin, Z., Zhang, C., Liu, B., Liu, J., Zhang, C., Lin, C., Li, H., and Zhao, T. (2015).
Over-expression of an S-domain receptor-like kinase extracellular domain improves
panicle architecture and grain yield in rice. Journal of experimental botany **66,** 7197-
7209.

1032