Key features of the genetic architecture and evolution of host-microbe interactions revealed by high-resolution genetic mapping of the mucosa-associated gut microbiome in hybrid mice

Shauni Doms[1,2], **Hanna Fokt**[1,2], **Malte Christoph Rühlemann**[3,4], **Cecilia J. Chung**[1,2], **Axel Künstner**[5], **Saleh Ibrahim**[5], **Andre Franke**[3], **Leslie M. Turner**[7]*† **and John F. Baines**[1,2]*†

[1] Max Planck Institute for Evolutionary Biology, Plön, Germany
[2] Section of Evolutionary Medicine, Institute for Experimental Medicine, Kiel University, Kiel, Germany
[3] Institute for Clinical Molecular Biology (IKMB), Kiel University, Kiel Germany
[4] Institute for Medical Microbiology and Hospital Epidemiology, Hannover Medical School, Hannover, Germany
[5] Institute of Experimental Dermatology, University of Lübeck, Lübeck, Germany
[6] Sharjah Institue of Medical Research, Sharjah, UAE
[7] Milner Centre for Evolution, Department of Biology & Biochemistry, University of Bath, United Kingdom

* Correspondence: l.m.turner@bath.ac.uk; baines@evolbio.mpg.de
† These authors contributed equally to this work.

Abstract: Determining the forces that shape diversity in host-associated bacterial communities is critical to understanding the evolution and maintenance of metaorganisms. To gain deeper understanding of the role of host genetics in shaping gut microbial traits, we employed a powerful genetic mapping approach using inbred lines derived from the hybrid zone of two incipient house mouse species. Further, we uniquely performed our analysis on microbial traits measured at the gut mucosal interface, which is in more direct contact with host cells and the immune system. A high number of mucosa-associated bacterial taxa have significant heritability estimates; heritabilities are greater for 16S rRNA transcript- compared to gene copy-based traits, and interestingly, are positively correlated with cospeciation rate estimates. Genome-wide association mapping identifies 443 loci influencing 123 taxa, with narrow genomic intervals pinpointing promising candidate genes and pathways. Importantly, we identified an enrichment of candidate genes associated with several human diseases, including inflammatory bowel disease, and functional categories including innate immunity and G-protein-coupled receptors. These results highlight key features of the genetic architecture of mammalian host-microbe interactions and how they diverge as new species form.

Keywords: microbiome; GWAS; cospeciation; codiversification; hybridization; phylosymbiosis

41 **Introduction**

42   The recent widespread recognition of the gut microbiome's importance to host
43 health and fitness represents a critical advancement of biomedicine. Host phenotypes
44 affected by the gut microbiome are documented in humans (Ley et al., 2006; Turnbaugh
45 et al., 2009; Lynch and Pedersen, 2016), laboratory animals (Backhed et al., 2004;
46 Turnbaugh et al., 2008; Rolig et al., 2015; Rosshart et al., 2017; Gould et al., 2018), and
47 wild populations (Suzuki, 2017; Roth et al., 2019; Suzuki et al., 2020; Hua et al., 2020),
48 and include critical traits such as aiding digestion and energy uptake (Rowland et al.,
49 2018), and the development and regulation of the immune system (Davenport, 2020).

50   Despite the importance of gut microbiome, community composition varies
51 significantly among host species, populations, and individuals (Benson et al., 2010;
52 Yatsunenko et al., 2012; Brooks et al., 2016; Rehman et al., 2016; Amato et al., 2019).
53 While a portion of this variation is expected to be selectively neutral, alterations of the
54 gut microbiome are on the one hand linked to numerous human diseases (Carding et al.,
55 2015; Lynch and Pedersen, 2016) such as diabetes (Qin et al., 2012), inflammatory bowel
56 disease (IBD) (Ott et al., 2004; Gevers et al., 2014) and mental disorders (Clapp et al.,
57 2017). On the other hand, there is evidence that the gut microbiome can play an
58 important role in adaptation on both recent- (Hehemann et al., 2010; Suzuki and Ley,
59 2020) and ancient evolutionary timescales (). Collectively, these phenomena suggest that
60 it would be evolutionarily advantageous for hosts to influence their microbiome.

61   An intriguing observation made in comparative microbiome research in the last
62 decade is that the pattern of diversification among gut microbiomes appears to mirror
63 host phylogeny (Ochman et al., 2010). This phenomenon, coined "phylosymbiosis"
64 (Brucker and Bordenstein, 2012a; Brucker and Bordenstein, 2012b; Lim and Bordenstein,
65 2020), is documented in a number of diverse host taxa (Brooks et al., 2016) and also
66 extends to the level of the phageome (Gogarten et al., 2021). Several non-mutually
67 exclusive hypotheses are proposed to explain phylosymbiosis (Moran and Sloan, 2015).
68 However, it is likely that vertical inheritance is important for at least a subset of taxa, as
69 signatures of co-speciation/-diversification are present among numerous mammalian
70 associated gut microbes (Moeller et al., 2016; Groussin et al., 2017; Moeller et al., 2019),
71 which could also set the stage for potential coevolutionary processes. Importantly,
72 experiments involving interspecific fecal microbiota transplants indeed provide
73 evidence of host adaptation to their conspecific microbial communities (Brooks et al.,
74 2016; Moeller et al., 2019). Further, cospeciating taxa were observed to be significantly
75 enriched among the bacterial species depleted in early onset IBD, an immune-related
76 disorder, suggesting a greater evolved dependency on such taxa (Papa et al., 2012;
77 Groussin et al., 2017). However, the nature of genetic changes involving host-microbe
78 interactions that take place as new host species diverge remains under-explored.

79   House mice are an excellent model system for evolutionary microbiome research, as
80 studies of both natural populations and laboratory experiments are possible (Suzuki,

2017; Suzuki et al., 2019). In particular, the house mouse species complex is comprised of
subspecies that hybridize in nature, enabling the potential early stages of
codiversification to be studied. We previously analyzed the gut microbiome across the
central European hybrid zone of *Mus musculus musculus* and *M. m. domesticus* (Wang et
al., 2015), which share a common ancestor ~ 0.5 million years ago (Geraldes et al., 2008).
Importantly, transgressive phenotypes (i.e. exceeding or falling short of parental values)
among gut microbial traits as well as increased intestinal histopathology scores were
common in hybrids, suggesting that the genetic basis of host control over microbes has
diverged (Wang et al., 2015). The same study performed an F_2 cross between wild-
derived inbred strains of *M. m. domesticus* and *M. m. musculus* and identified 14
quantitative trait loci (QTL) influencing 29 microbial traits. However, like classical
laboratory mice, these strains had a history of rederivation and reconstitution of their
gut microbiome, thus leading to deviations from the native microbial populations found
in nature (Rosshart et al., 2017; Org and Lusis, 2018), and the genomic intervals were too
large to identify individual genes.

In this study, we employed a powerful genetic mapping approach using inbred
lines directly derived from the *M. m. musculus* - *M. m. domesticus* hybrid zone, and
further focus on the mucosa-associated microbiota due to its more direct interaction
with host cells (Fukata and Arditi, 2013; Chu and Mazmanian, 2013), distinct functions
compared to the luminal microbiota (Wang et al., 2010; Vaga et al., 2020), and greater
dependence on host genetics (Spor et al., 2011; Linnenbrink et al., 2013). Previous
mapping studies using hybrids raised in a laboratory environment showed that high
mapping resolution is possible due to the hundreds of generations of natural admixture
between parental genomes in the hybrid zone (Turner and Harr, 2014; Pallares et al.,
2014; Škrabar et al., 2018). Accordingly, we here identify 443 loci contributing to
variation in 123 taxa, whose narrow genomic intervals (median <2Mb) enable many
individual candidate genes and pathways to be pinpointed. We identify a high
proportion of bacterial taxa with significant heritability estimates, and find that bacterial
phenotyping based on 16S rRNA transcript compared to gene copy-based profiling
yields an even higher proportion. Further, these heritability estimates also significantly
positively correlate with cospeciation rate estimates, suggesting a more extensive host
genetic architecture for cospeciating taxa. Finally, we identify numerous enriched
functional pathways, whose role in host-microbe interactions may be particularly
important as new species form.

115 **Results**

116 *Microbial community composition*

117   To obtain microbial traits for genetic mapping in the G2 mapping population, we

118 sequenced the 16S rRNA gene from caecal mucosa samples of 320 hybrid male mice

119 based on DNA and RNA (cDNA), which reflect bacterial cell number and activity,

120 respectively. After applying quality filtering and subsampling 10,000 reads per sample,

121 we identified a total of 4684 amplicon sequence variants (ASVs). For further analyses,

122 we established a "core microbiome" (defined in Methods), such that analyses were

123 limited to those taxa common and abundant enough to reveal potential genetic signal.

124 The core microbiome is composed of four phyla, five classes, five orders, eleven families,

125 27 genera, and 90 ASVs for RNA, and four phyla, five classes, six orders, twelve families,

126 28 genera and 46 ASVs for DNA. A combined total of 98 unique ASVs belong to the core,

127 of which 38 were shared between DNA and RNA (Suppl. Fig. 1). The most abundant

128 genus in our core microbiome is *Helicobacter* (Suppl. Fig. 2), consistent with a previous

129 study of the wild hybrid *M. m. musculus*/*M. m. domesticus* mucosa-associated

130 microbiome (Wang et al., 2015).

131 *Correlation between host genetic relatedness and microbiome structure*

132   To gain a broad sense of the contribution of genetic factors to the variability of

133 microbial phenotypes in our mapping population, we compared the kinship matrix

134 based on genotypes to an equivalent based on gut microbial composition, whereby ASV

135 abundances were used as equivalents of gene dosage. We found a significant correlation

136 between these matrices (P = .001; Suppl. Fig. 3), indicating a host genetic effect on the

137 diversity of the gut microbiota.

138 *SNP-based heritability*

139   Next, we used a SNP-based approach to estimate the proportion of variance

140 explained (PVE) of the relative abundance of taxa, also called the narrow-sense

141 heritability (h^2) or SNP-based heritability. Out of the 153 total core taxa, we identified 46

142 taxa for DNA and 69 taxa for RNA with significant heritability estimates (P_{RLRT} < .05),

143 with estimates ranging between 29 and 91% (see Fig. 1A-B and Suppl. Table 1). An

144 unclassified genus belonging to the phylum Bacteroidetes followed by ASV7 (genus

145 *Paraprevotella*), *Paraprevotella* and Paraprevotellaceae showed the highest heritability

146 among DNA-based traits (91.8%, 88.8%, 88.8%, and 87.1%, respectively; Fig. 1A), while

147 ASV97 (genus *Oscillibacter*), followed by Prevotellaceae, *Paraprevotella* and ASV7

148 (*Paraprevotella*) had the highest heritability among RNA-based traits (86.6%, 85.7%,

149 85.7%, and 85.6%, resp.; Fig. 1B). The heritability estimates for DNA- and RNA-based

150 measurements of the same taxa are significantly correlated (P = 5.013 x 10^{-8}, R^2=0.58,

151 Suppl. Fig. 4), and neither measure appears to be systematically more heritable than

152   another, i.e. some taxa display higher RNA-based heritability estimates and others
153   higher DNA-based estimates.

154   *Heritability estimates are correlated with predicted co-speciation rates*

155   In an important meta-analysis of the gut microbiome across diverse mammalian
156   taxa, Groussin et al. (2017) estimated co-speciation rates of individual bacterial taxa by
157   measuring the congruence of host and bacteria phylogenetic trees relative to the number
158   of host-swap events. We reasoned that taxa with higher co-speciation rates might also
159   demonstrate higher heritability, as these more intimate evolutionary relationships would
160   provide a greater opportunity for genetic aspects to evolve. Intriguingly, we observe a
161   significant positive correlation for RNA-based traits (P= .008, R^2=.46, Fig. 1D) and a
162   similar trend for DNA (P= 0.1; Fig. 1C). These results support the notion that
163   cospeciating taxa evolved a greater dependency on host genes, and further suggest that
164   bacterial activity may better reflect the underlying biological interactions.

165   **Figure 1**: (A-B) Heritability estimates for the relative abundance of bacterial taxa. Proportion
166   of variance explained for each taxon on DNA level (A), and RNA level (B) for all SNPs (GRM) in
167   green, mating pair identifier in blue and residual variance in grey. Only significant heritability
168   estimates are shown ($P < .05$). The text labels on the y-axis are colored according to taxonomic
169   level: ASV in black, genus in purple, family in light blue, order in red, class in green, and phylum
170   in yellow. (C-D) Relationship between the heritability estimates for the relative abundance of
171   bacterial taxa and co-speciation rate for the same genus calculated by Groussin et al. (2017). DNA
172   level (C), and RNA level (D). The blue line represents a linear regression fit to the data and the
173   grey area the corresponding confidence interval.

174   *Genetic mapping of host loci determining microbiome composition*

175   Next, we performed genome-wide association mapping of the relative abundances
176   of core taxa, in addition to two alpha-diversity measures (Shannon and Chao1 indices),
177   based on 32,625 SNPs. We included both additive and dominance terms in the model to
178   enable the identification of under- and over-dominance (see Methods). While we found
179   no significant associations for alpha diversity at either the DNA or RNA level ($P > 1.53 \times$
180 10^{-6}), a total of 1099 genome-wide significant associations were identified for individual
181   taxa ($P < 1.53 \times 10^{-6}$, Suppl. Table 2), of which 443 achieved study-wide significance (P
182 $< 1.29 \times 10^{-8}$). Apart from the X chromosome, all autosomal chromosomes contained
183   study-wide significant associations (Fig. 2). Out of the 153 mapped taxa, 123 had at least
184   one significant association (Table 1). For the remainder of our analyses, we focus on the
185   results using the more stringent study-wide threshold, and combined significant SNPs
186   within 10 Mb into significant regions (Suppl. Table 3). The median size of significant
187   regions is 1.91 Mb, which harbor a median of 14 protein-coding genes. On average, we
188   observe 10 significant mouse genomic regions per bacterial taxon.

189   Of the significant loci with estimated interval sizes, we find 73 intervals (16.5%) that
190   are smaller than one Mb (Suppl. Table 4). The smallest interval is only 231 bases and
191   associated with the RNA-based abundance of an unclassified genus belonging to
192   Deltaproteobacteria. It is situated in an intron of the C3 gene, a complement component
193   playing a central role in the activation of the complement system, which modulates
194   inflammation and contributes to antimicrobial activity (Ricklin et al., 2016).

Figure 2: Heatmap of significant host loci from association mapping of bacterial abundances. Karotype plot showing the number of significant loci found using a study-wide threshold, where (A) plots the significance intervals, and (B) the significant SNP markers on the chromosomes.

199 **Table. 1**. Overview of mapping statistics.

	DNA	RNA	Total
Mapped taxa	101	142	153
Taxa with significant loci	67	96	123
Median interval size (Mb)	1.52	2.29	1.91
Total significant loci	478	791	1269
Unique significant loci	179	313	443
Significant loci total P	91	167	233
Significant loci additive P	155	260	377
Significant loci dominance P	95	166	231
Median significant loci per trait	5	6	8
Median unique significant loci per trait	3	3	4
Median unique significant SNPs per locus	2	2.5	2
Median number of genes per locus	31	52	43
Median protein coding genes per locus	11	15	14

The significant genomic regions and SNPs are displayed in Figure 2A and 2B,
respectively. Individual SNPs were associated with up to 12 taxa, and significant
intervals with up to 30 taxa. The SNPs with the lowest P values were associated with the
genus *Dorea* and two ASVs belonging to *Dorea* (ASV184 and ASV293; Suppl. Fig. 5). At
the RNA level this involves two loci: mm10-chr4: 67.07 Mb, where the peak SNP is 13 kb
downstream of the closest gene *Tlr4* (UNC7414459, $P=2.31 \times 10^{-69}$, additive $P= 4.48 \times$
10^{-118}, dominance $P= 1.37 \times 10^{-111}$), and mm10-chr15: 94.4 Mb, where the peak SNP is
found within the *Adamts20* gene (UNC26145702, $P=4.51 \times 10^{-65}$, additive $P= 1.87 \times 10^{-113}$,
dominance $P= 1.56 \times 10^{-105}$; Fig. 2; Suppl. Fig. 5). Interestingly, the *Irak4* gene, whose
protein product is rapidly recruited after TLR4 activation, is also located 181 kb
upstream of *Adamts20*. The five taxa displaying the most associations were ASV19
(*Bacteroides*), *Dorea*, ASV36 (*Oscillibacter*), ASV35 (*Bacteroides*), and ASV98 (unclassified
Lachnospiraceae) (Suppl. Fig. 6).

Ancestry, dominance, and effect sizes

A total of 435 significant SNPs were ancestry informative between *M. m. musculus*
and *M. m. domesticus* (*i.e.* represent fixed differences between subspecies). To gain further
insight on the genetic architecture of microbial trait abundances, we estimated the
degree of dominance at each significant locus using the
d/a ratio (Falconer, 1996), where alleles with strictly recessive, additive, and dominant
effects have d/a values of -1, 0, and 1, respectively. As half of the SNPs were not ancestry
informative (Fig. 3A), it was not possible to consistently have a associated with one
parent/subspecies, hence we report $d/|a|$ such that it can be interpreted with respect to
bacterial abundance. For the vast majority of loci (83.53%), the allele associated with
lower abundance is dominant or partially dominant ($-1.25 < d/|a| < -0.75$; Fig. 3B). On
the basis of the arbitrary cutoffs we used to classify dominance, only a small proportion

239 of alleles are underdominant (0.22%; $d/|a| < -1.25$) or overdominant (0.15%; $d/|a| >$
240 1.25). However for one-third of the significant SNPs, the heterozygotes display
241 transgressive phenotypes, i.e. mean abundances that are either significantly lower (31%
242 of SNPs)- or higher (2% of SNPs) than those of both homozygous genotypes.
243 Interestingly, the *domesticus* allele was associated with higher bacterial abundance in
244 two-thirds of this subset (33.2% vs 16.3% *musculus* allele; Fig. 3A).

245 **Figure 3**: Genetic architecture of significant loci. A) Source of the allele with the highest phe-
246 notypic value. B) Histogram of dominance values d/a of significant loci reveals a majority of
247 loci acting recessive or partially recessive. C) Histogram showing the percentage of variance
248 explained (PVE) by the peak SNP for DNA (blue, left) and RNA (orange, right). D) Collective
249 PVE by lead SNPs of significant loci within a taxon. Values are calculated separately for each
250 *P* value type (total, additive, and dominance).

251 Next, we estimated phenotypic effect sizes by calculating the percentage variance
252 explained (PVE) by the peak SNP of each significant region. Peak SNPs explain between
253 3% and 64% of the variance in bacterial abundance, with a median effect size of 9.3%
254 (Fig. 3C). The combined effects of all significant loci for each taxon ranged from 4.9% to
255 259%, with a median of 41.8% (Fig. 3D). Note, combined effects for many taxa (33 out of
256 59) exceed SNP-heritability estimates (Fig. 1). While exceeding 100% explained variance
257 is biologically possible, as loci can have opposite phenotypic effects, many of these are
258 likely inflated due to the Beavis effect (Beavis, 1994).

Functional annotation of candidate genes

In order to reveal potential higher level biological phenomena among the identified
loci, we performed pathway analysis to identify interactions and functional categories
enriched among the genes in significant intervals. We used STRING (Szklarczyk et al.,
2019) to calculate a protein-protein interaction (PPI) network of 925 protein-coding
genes nearest to significant SNPs (upstream and/or downstream). A total of 768 genes
were represented in the STRING database, and the maximal network is highly
significant (PPI enrichment *P* value: 2.15×10^{-14}) displaying 668 nodes connected by 1797
edges and an average node degree of 4.68. After retaining only the edges with the
highest confidence (interaction score > 0.9), this results in one large network with 233
nodes, 692 edges and ten smaller networks (Fig. 4).

Next, we functionally annotated clusters using STRING's functional enrichment
plugin. The genes of the largest cluster are part of the G protein-coupled receptor
(GPCR) ligand binding pathway. GPCRs are the largest receptor superfamily and also
the largest class of drug targets (Sriram and Insel, 2018). We then calculated the top ten
hub proteins from the network based on Maximal Clique Centrality (MCC) algorithm
with CytoHubba to predict important nodes that can function as 'master switches'
(Suppl. Fig. 7). The top ten proteins contributing to the PPI network were GNG12,
MCHR1, NMUR2, PROK2, OXTR, XCR1, TACR3, CHRM3, PTGFR, and C3, which are
all involved in the GPCR signaling pathway.

Further, we performed enrichment analysis on the 925 genes nearest to significant
SNPs using the *clusterprofiler* R package. We found 14 KEGG pathways to be over-
represented: circadian entrainment, oxytocin signaling pathway, axon guidance, calcium
signaling, cAMP signaling, cortisol synthesis and secretion, cushing syndrome, gastric
acid secretion, glutamatergic synapse, mucin type O-glycan biosynthesis, inflammatory
mediator regulation of TRP channels, PD-L1 expression and the PD-1 checkpoint
pathway in cancer, tight junction, and the *Wnt* signaling pathway (Suppl. Table 5, Suppl.
Fig. 8-9). Finally, genes involved in five human diseases are enriched, among them
mental disorders (Suppl. Fig. 10).

Finally, due to the observation of a significant enrichment of cospeciating taxa
among the bacterial species depleted in early onset IBD (Groussin et al., 2017) and the
evidence that IBD is especially associated with a dysbiosis in mucosa-associated
communities (Yang et al., 2020a; Daniel et al., 2021), we specifically examined possible
over-representation of genes involved in IBD (Khan et al., 2021) among the 925 genes
neighboring significant SNPs. We found 14 out of the 289 IBD genes, which was
significantly more than expected by chance (10 000 times permuted mean: 2.7, simulated
P = .0001; Suppl. Table 6). Interestingly, SNPs in five out of the 14 genes are associated
with ASVs belonging to the genus *Oscillibacter*, a cospeciating taxon known to decrease
during the active state of IBD (Metwaly et al., 2020).

Figure 4: High confidence protein-protein interaction network of genes closest to SNPs significantly associated with bacterial abundances. Network clusters are annotated using STRING's functional enrichment (Doncheva et al., 2019). Nodes represent proteins and edges their respective interactions. Only edges with an interaction score higher than 0.9 are retained. The width of the edge line expresses the interaction score calculated by STRING. The color of the nodes describe the expression of the protein in the intestine where yellow is not expressed and purple is highly expressed.

Comparison of significant loci to published gut microbiome mapping studies

Next, we compiled a list of 648 unique confidence intervals of significant associations with gut bacterial taxa from seven previous mouse QTL studies (Benson et al., 2010; McKnite et al., 2012; Leamy et al., 2014; Wang et al., 2015; Org et al., 2015; Snijders et al., 2016; Kemis et al., 2019) and compared this list to our significance intervals for bacterial taxa at both the DNA and RNA level (346 unique intervals). Regions larger than 10Mb were removed from all studies. We found 434 overlapping intervals, which is significantly more than expected by chance (10 000 times permuted mean: 368, simulated *P*=.0073, see Methods). Several of our smaller significant loci overlapped with larger loci from previous studies and removing this redundancy left 186 significant loci with a median interval size of 0.78 Mb (Fig. 5). The most frequently identified locus is located on chromosome 2 169-171 Mb where protein coding genes *Gm11011, Znf217, Tshz2, Bcas1, Cyp24a1, Pfdn4, 4930470P17Rik,* and *Dok5* are situated.

Additionally, we collected genes within genome-wide significant regions reported in seven human microbiome GWAS (mGWAS) (Bonder et al., 2016; Turpin et al., 2016;

320 Goodrich et al., 2016; Wang et al., 2016; Hughes et al., 2020; Rühlemann et al., 2021;

321 Kurilshikov et al., 2021). However, no significant over-representation of genes was

322 found within our significance intervals ($P = .156$), nor within our list of genes closest to a

323 significant SNP ($P = .62$).

324 **Figure 5**: Heatmap showing the significant loci in this study that were previously

325 found in other QTL studies of the mouse gut microbiome. The genes present in

326 two repeatedly identified regions are depicted in boxes.

327 *Proteins differentially expressed in germ-free vs conventional mice*

328 To further validate our results, we compared the list of genes contained within

329 intervals of our study to a list of differentially expressed protein between germ-free and

330 conventionally raised mice (Mills et al., 2020). This comparison was made based on the

331 general expectation that genes associated with variation in microbial abundances would

332 be more likely to differ according to the colonization status of the host. Thus, we

333 examined the intersection between genes identified in our study and the proteins

334 identified as highly associated ($|\pi| > 1$) with the colonization state of the colon and the

335 small intestine (Mills et al., 2020). Out of the 373 over- or under-expressed proteins

336 according to colonization status, we find 198 of their coding genes to be among our

337 significant loci, of which 17 are the closest genes to a significant marker (*Iyd, Nln,*

338 *Slc26a3, Slc3a1, Myom2, Nebl, Tent5a, Fxr1, Cbr3, Chrodc1, Nucb2, Arhgef10l, Sucla2, Enpep,*

339 *Prkcq, Aacs,* and *Cox7c*). This is significantly more than expected by chance (simulated

340 $P=.0156$, 10 000 permutations). Further, analyzing the protein-protein interactions with

341 STRING results in a significant network ($P=1.73 \times 10^{-14}$, and average node degree 2.4,

342 Suppl. Fig. 11), with *Cyp2c65, Cyp2c55, Cyp2b10, Gpx2, Cth, Eif3k, Eif1, Sucla2,* and *Rpl17*
343 identified as hub genes (Suppl. Fig. 12).

344 Subsequently, we merged the information from Mills et al. (2020) and the seven
345 previous QTL mapping studies discussed above to further narrow down the most
346 promising candidate genes, and found 30 genes overlapping with our study. Of these 30
347 genes, six are the closest gene to a significant SNP. These genes are myomesine 2
348 (*Myom2*), solute carrier family 3 member 1 (*Slc3a1*), solute carrier family 26 member 3
349 (*Slc26a3*), nebulette (*Nebl*), carbonyl reductase 3 (*Cbr3*), and acetoacetyl-coA synthetase
350 (*Aacs*).

351 *Candidate genes influencing bacterial abundance*

352 Finally, all previously mentioned candidate genes were combined in one gene set of
353 304 genes and compiled in a highly significant PPI network ($P < 1.0 \times 10^{-16}$, average node
354 degree=4.85, see Methods 4.13). Guided by this network, we filtered out genes situated
355 in the same genomic region and kept the gene with the highest connectivity and
356 supporting information (original network see Suppl. Fig. 13). This gave a resulting gene
357 set of 80 candidate genes (Fig. 6 and Suppl. Table 7). The G protein, GNG12 and the
358 complement component 3 C3, are the proteins with the most edges in the network (30
359 and 25, respectively), followed by MCHR1, CXCL12, and NMUR2 with each 18 edges.
360 Of these 80 highly connected genes, 66 are associated with bacteria that are either co-
361 speciating (co-speciation rate > 0.5; Groussin et al., 2017) and/or have high heritability
362 (> 0.5) suggesting a functionally important role for these bacterial taxa (Suppl. Table 7).

Figure 6: Network of host candidate genes influencing bacterial traits using STRING (https://string-db.org). The nodes represent proteins and are colored according to a selection of enriched GO terms and pathways: G protein coupled receptor (GPCR) signaling (red), regulation of the immune system process (blue), response to nutrient levels (light green), fatty acid metabolic process (pink), glucose homeostasis (purple), response to antibiotic (orange), regulation of feeding behavior (yellow), positive regulation of insulin secretion (dark green), circadian entrainment (brown), and response to vitamin D (turquoise). The color of the edges represents the interaction type: known interactions from curated databases (turquoise) or experimentally determined (pink); predicted interactions from gene neighborhood (green), gene fusions (red), gene co-occurrence (blue); other interactions from text-mining (light green), co-expression (black), and protein homology (purple).

375   **Discussion**

376   Understanding the forces that shape variation in host-associated bacterial
377   communities within host species is key to understanding the evolution and maintenance
378   of meta-organisms. Although numerous studies in mice and humans demonstrate that
379   host genetics influences gut microbiota composition (McKnite et al., 2012; Leamy et al.,
380   2014; Goodrich et al., 2014; Org et al., 2015; Davenport et al., 2015; Wang et al., 2016;
381   Bonder et al., 2016; Goodrich et al., 2016; Kemis et al., 2019; Suzuki et al., 2019; Ishida et
382   al., 2020; Hughes et al., 2020; Rühlemann et al., 2021), our study is unique in a number of
383   important ways. First, the unique genetic resource of mice collected from a naturally
384   occurring hybrid zone together with their native microbes yielded extremely high
385   mapping resolution and the possibility to uncover ongoing evolutionary processes in
386   nature. Second, our study is the first to perform genetic mapping of 16S rRNA
387   transcripts in the gut environment, which was previously shown to be superior to DNA-
388   based profiling in a genetic mapping study of the skin microbiota (Belheouane et al.,
389   2017). Third, our study is one of the only to specifically examine the mucosa-associated
390   community. It was previously reasoned that the mucosal environment may better reflect
391   host genetic variation (Spor et al., 2011), and evidence for this hypothesis exists in nature
392   (Linnenbrink et al., 2013). Finally, by cross-referencing our results with previous
393   mapping studies and recently available proteomic data from germ-free versus
394   conventional mice, we curated a more reliable list of candidate genes and pathways.
395   Taken together, these results provide unique and unprecedented insight into the genetic
396   basis for host-microbe interactions.

397   Importantly, by using wild-derived hybrid inbred strains to generate our mapping
398   population, we gained insight into the evolutionary association between hosts and their
399   microbiota at the transition from within species variation to between species divergence.
400   Genetic relatedness in our mapping population significantly correlates with microbiome
401   similarity, supporting a basis for codiversification at the early stages of speciation. A
402   substantial proportion of microbial taxa are heritable, and heritability is correlated with
403   cospeciation rates. This suggests that (i) vertical transmission could enable greater host
404   adaptation to bacteria and/or (ii) the greater number of host genes associated with
405   cospeciating taxa could indicate a greater dependency on the host, such that survival
406   outside a specific host is reduced, making horizontal transmission less likely.

407   By performing 16S rRNA gene profiling at both the DNA and RNA level, we found
408   that 30% (DNA-based) to 45% (RNA-based) of bacterial taxa are heritable, which is
409   consistent with or higher than estimates reported in humans (~10%, Goodrich et al.,
410   2016; ~21%, Turpin et al., 2016) and previous mouse studies (Kovacs et al., 2011; McKnite
411   et al., 2012; Campbell et al., 2012; O'Connor et al., 2014; Carmody et al., 2015; Korach-
412   Rechtman et al., 2019;). The high proportion of heritable taxa, with estimates of up to
413   91%, is likely explained in part by several factors of our study design. First, mice were

raised in a controlled common environment, and heritability estimates in other
mammals were shown to be contingent on the environment (Grieneisen et al., 2021).
Further, bacterial communities were sampled from cecal tissue instead of fecal content
(Linnenbrink et al., 2013), and genetic variation was higher than in a typical mapping
study due to subspecies differences. For the RNA-based traits, heritability estimates
were significantly correlated with previously reported cospeciation rates in mammals
(Groussin et al., 2017). This pattern, as well as the higher proportion of heritable taxa in
RNA-based traits, suggest that host genetic effects are more strongly reflected by
bacterial activity than cell number.

Accordingly, we found a total of 179 and 313 unique significant loci for DNA-based
and RNA-based bacterial abundance, respectively, passing the conservative study-wide
significance threshold. Taxa had a median of five significant loci, suggesting a complex
and polygenic genetic architecture affecting bacterial abundances. We identify a higher
number of loci in comparison to previous QTL and GWAS studies in mice (Benson et al.,
2010; McKnite et al., 2012; Leamy et al., 2014; Wang et al., 2015; Org et al., 2015; Snijders
et al., 2016; Kemis et al., 2019), which may be due to a number of factors. The parental
strains of our study were never subjected to rederivation and subsequent reconstitution
of their microbiota, and natural mouse gut microbiota are more variable than the
artificial microbiota of laboratory strains (Kohl and Dearing, 2014; Weldon et al., 2015;
Suzuki, 2017; Rosshart et al., 2017;). Furthermore, as noted above, our mapping
population harbors both within- and between-subspecies genetic variation. We crossed
incipient species sharing a common ancestor ~ 0.5 million years ago, hence we may also
capture the effects of mutations that fixed rapidly between subspecies due to strong
selection, which are typically not variable within species (Walsh, 1998; Barton and
Keightley, 2002).

Importantly, our results also help to describe general features of the genetic
architecture of bacterial taxon activity. For the majority of loci, the allele associated with
lower relative abundance of the bacterial taxon was (partially) dominant. This suggests
there is strong purifying selection against a high abundance of any particular taxon,
which may help ensure high alpha diversity. The heterozygotes of one-third of
significant SNPs displayed transgressive phenotypes. This is consistent with previous
studies of hybrids (Turner et al., 2012; Turner and Harr, 2014; Wang et al., 2015;), for
example, wild-caught hybrids showed broadly transgressive gut microbiome
phenotypes. This pattern can be explained by over- or underdominance, or by epistasis
(Rieseberg et al., 1999).

Notably, many loci significantly associated with bacterial abundance in this study
were implicated in previous studies (Fig. 5). For example, chromosome 2 169-171 Mb is
associated with ASV23 (*Eisenbergiella), Eisenbergiella* and ASV32 (unclassified
Lachnospiraceae) in this study, and overlaps with significant loci from three previous
studies (Leamy et al., 2014; Snijders et al., 2016; Kemis et al., 2019). This region contains

eight protein-coding genes: *Gm11011, Znf217, Tshz2, Bcas1, Cyp24a1, Pfdn4,*
4930470P17Rik, and *Dok5.* Another hotspot is on chromosome 5 101-103 Mb. This locus is
significantly associated with four taxa in this study (Prevotellaceae, *Paraprevotella*, ASV7
genus *Paraprevotella* and *Acetatifactor*) and overlaps with associations for Clostridiales,
Clostridiaceae, Lachnospiraceae, and Deferribacteriaceae (Snijders et al., 2016). Protein-
coding genes in this region are: *Nkx6-1*, *Cds1*, *Wdfy3*, *Arhgap24*, and *Mapk10*. As previous
studies were based on rederived mouse strains, identifying significant overlap in the
identification of host loci suggests that some of the same genes and/or mechanisms
influencing major members of gut microbial communities are conserved even in the face
of community 'reset' in the context of re-derivation. The identity of the taxa is however
not always the same, which suggests that functional redundancy may contribute to
these observations, if *e.g.* several bacterial taxa fulfill the same function within the gut
microbiome (Moya and Ferrer, 2016; Tian et al., 2020). Additionally, there is significant
overlap of genes within loci identified in the current study and proteins differentially
expressed in the intestine of germ-free mice compared to conventionally raised mice
(Mills et al., 2020). Finally, by analyzing the functions of the genes closest to significant
SNPs, we found that 12 of the 14 significantly enriched KEGG pathways were shown to
be related to interactions with bacteria (Fonken et al., 2010; Thaiss et al., 2014; Neumann
et al., 2014; Thaiss et al., 2015a; Thaiss et al., 2015b; Castoldi et al., 2015; Erdman and
Poutahidis, 2016; Thaiss et al., 2016; Deaver et al., 2018; Wu et al., 2018; Peng et al., 2020;
Nagpal et al., 2020; Hollander and Kaunitz, 2020; Suppl. Table 5).

To improve the robustness of our results, we combined multiple lines of evidence to
prioritize candidates, resulting in a network of 80 genes (Suppl. Table 7). At the center of
this network is a set of 22 proteins involved in G-protein coupled receptor signaling (Fig.
6, red nodes). MCHR1, NMUR2, and TACR3 (Fig. 6, yellow) are known to regulate
feeding behavior (Saito et al., 1999; Cardoso et al., 2012; Smith et al., 2019), and CHRM3
to control digestion (Gautam et al., 2006; Tanahashi et al., 2009). Gut microbes can
produce GPCR agonists to elicit host cellular responses (Cohen et al., 2017; Colosimo et
al., 2019; Chen et al., 2019; Pandey et al., 2019). Thus, GPCRs may be key modulators of
communication between the gut microbiota and host. Another interesting group of
genes are those responding to nutrient levels (*Bmp7, Cd40, Aacs, Gclc, Nmur2, Cyp24a1,*
Adcyap1, Serpinc1, and *Wnt11*) (Sethi and Vidal-Puig, 2008; Peier et al., 2009; Townsend et
al., 2012; Yi and Bishop, 2015; Shi and Tu, 2015; Toderici et al., 2016; Yasuda et al., 2021;
Gastelum et al., 2021;), as gut microbiota affect host nutrient uptake (Chung et al., 2018).
In addition, CYP24A1, BMP7 and CD40 respond to vitamin D. Previous studies
identified vitamin D/the vitamin D receptor to play a role in modulating the gut
microbiota (Wang et al., 2016; Malaguarnera, 2020; Yang et al., 2020b; Singh et al., 2020),
and CD40 is known to induce a vitamin D dependent antimicrobial response through
IFN-γ activation (Klug-Micu et al., 2013).

Another important category of candidate genes are those involved in immunity.
Our most significant SNP was situated downstream of the *Tlr4* gene and was associated
with the genus *Dorea* and several *Dorea* species. *Dorea* is a known short chain fatty acid
producer (Taras et al., 2002; Reichardt et al., 2018) and interacts with tight junction
proteins *Claudin-2* and *Occludin* (Alhasson et al., 2017). *Tlr4* is a member of the Toll-like
receptor family, and has been linked with obesity, inflammation, and changes in the gut
microbiota (Velloso et al., 2015). These combined results reflect an important role for
Dorea in fatty acid harvesting and intestinal barrier integrity, both of which could act
systemically to activate TLR4 and to promote metabolic inflammation (Cani et al., 2008;
Delzenne et al., 2011; Nicholson et al., 2012). Moreover, the SNP with the second lowest
P value was associated with the same taxa and situated 181 kb upstream of *Irak4*. IRAK4
is rapidly recruited after TLR4 activation to enable downstream activation of the NFκB
immune pathway. *Irak4* has previously been associated with a change in bacterial
abundance using inbred mice (McKnite et al., 2012; Org et al., 2015).

Finally, we identified notable links between candidate genes and five human
diseases (mental disorders, blood pressure finding, systemic arterial pressure, substance-
related disorders, and atrial septal deficits; Suppl. Fig. 10). The connection to mental
disorders is intriguing as involvement of the gut microbiota is suspected (Kelly et al.,
2015; Foster et al., 2017; Cox and Weiner, 2018; Chen et al., 2019; Sarkar et al., 2020;
Parker et al., 2020; Flux and Lowry, 2020). Taken together with our finding of an
enriched set of GPCRs, this highlights the importance of host-microbial interplay along
the gut-brain axis. Moreover, we also identify a significant over-representation of IBD
genes (Khan et al., 2021) among the 925 genes nearest to significant SNPs (Suppl. Table
6). Interestingly, SNPs in five out of 14 genes are associated with ASVs belonging to the
genus *Oscillibacter,* a highly cospeciating taxon known to decrease during the active state
of IBD (Metwaly et al., 2020).

In summary, our study provides a number of novel insights into the importance of
host genetic variation in shaping the gut microbiome, in particular for cospeciating
bacterial taxa. These findings provide an exciting foundation for future studies of the
precise mechanisms underlying host-gut microbiota interactions in the mammalian gut
and should encourage future genetic mapping studies that extend analyses to the
functional metagenomic sequence level.

525 **Materials and Methods**

526 *Intercross design*

527 We generated a mapping population using partially inbred strains derived from
528 mice captured in the *M. m. musculus* - *M. m. domesticus* hybrid zone around Freising,
529 Germany in 2008 (Turner et al., 2012). Originally, four breeding stocks were derived
530 from 8-9 ancestors captured from one (FS, HA, TU) or two sampling sites (HO), and
531 maintained with four breeding pairs per generation using the HAN-rotation out-
532 breeding scheme (Rapp, 1972). Eight inbred lines (two per breeding stock) were
533 generated by brother/sister mating of the 8th generation lab-bred mice. We set up the
534 cross when lines were at the 5th-9th generation of brother-sister meeting, with
535 inbreeding coefficients of > 82%.

536 We first set up eight G1 crosses, each with one predominantly *domesticus* line (FS,
537 HO - hybrid index <50%; see below) and one predominantly *musculus* line (HA, TU -
538 hybrid index >50%); each line was represented as a dam in one cross and sire in another
539 (Suppl. Fig. 14). One line, FS5, had a higher hybrid index than expected, suggesting
540 there was a misidentification during breeding (see genotyping below). Next, we set up
541 G2 crosses in eight combinations (subcrosses), such that each G2 individual has one
542 grandparent from each of the initial four breeding stocks. We included 40 males from
543 each subcross in the mapping population.

544 This study was performed according to approved animal protocols and in-
545 stitutional guidelines of the Max Planck Institute. Mice were maintained and handled in
546 accordance with FELASA guidelines and German animal welfare law (Tierschutzgesetz
547 § 11, permit from Veterinäramt Kreis Plön: 1401-144/PLÖ-004697).

548 *Sample collection*

549 Mice were sacrificed at 91 ± 5 days by CO^2 asphyxiation. We recorded body weight,
550 body length and tail length, and collected ear tissue for genotyping. The caecum was
551 removed and gently separated from its contents through bisection and immersion in
552 RNAlater (Thermo Fisher Scientific, Schwerte, Germany). After overnight storage in
553 RNAlater at 4° C, the RNAlater was removed and tissue stored at -20° C.

554 *DNA extraction and sequencing*

555 We simultaneously extracted DNA and RNA from caecum tissue samples using
556 Qiagen (Hilden, Germany) Allprep DNA/RNA 96-well kits. We followed the
557 manufacturer's protocol, with the addition of an initial bead beating step using Lysing
558 matrix E tubes (MP Biomedical, Eschwege) to increase cell lysis. We used caecum tissue
559 because host genetics has a greater influence on the microbiota at this mucosal site than
560 on the lumen contents (Linnenbrink et al., 2013). We performed reverse transcription of
561 RNA with High-Capacity cDNA Transcription Kits from Applied Biosystems
562 (Darmstadt, Germany). We amplified the V1-V2 hypervariable region of the 16S rRNA

gene using barcoded primers (27F-338R) with fused MiSeq adapters and heterogeneity

spacers following (Rausch et al., 2016) and sequenced amplicons with 250 bp paired-

reads on the Illumina MiSeq platform.

16S rRNA gene analysis

We assigned sequences to samples by exact matches of MID (multiplex identifier, 10

nt) sequences processed 16S rRNA sequences using the DADA2 pipeline, implemented

in the DADA2 R package, version 1.16.0 (Callahan et al., 2016; Callahan, 2016). Briefly,

raw sequences were trimmed and quality filtered with the maximum two 'expected

errors' allowed in a read, paired sequences were merged and chimeras removed. For all

downstream analyses, we rarefied samples to 10,000 reads each. Due to the quality

filtering, we have phenotyping data for 286 individuals on DNA level, and 320 G2

individuals on RNA level. We classified taxonomy using the Ribosomal Database Project

(RDP) training set 16 (Cole et al., 2014). Classifications with low confidence at the genus

level (<0.8) were grouped in the arbitrary taxon 'unclassified_group'.

We used the phyloseq R package (version 1.32.0) to estimate alpha diversity using

the Shannon index and Chao1 index, and beta diversity using Bray-Curtis distance

(McMurdie and Holmes, 2013). We defined core microbiomes at the DNA- and RNA-

level, including taxa present in > 25% of the samples and with median abundance of

non-zero values > 0.2% for amplicon sequence variant (ASV) and genus; and >0.5% for

family, order, class and phylum.

Genotyping

We extracted genomic DNA from ear samples using Qiagen Blood and Tissue 96

well kits (Hilden, Germany), according to the manufacturer's protocol. We sent DNA

samples from 26 G0 mice and 320 G2 mice to GeneSeek (Neogen, Lincoln, NE) for

genotyping using the Giga Mouse Universal Genotyping Array (GigaMUGA; Morgan et

al., 2015), an Illumina Infinium II array containing 141,090 single nucleotide

polymorphism (SNP) probes. We quality-filtered genotype data using plink 1.9 (Chang

et al., 2015); we removed individuals with call rates <90% and SNPs that were: not bi-

allelic, missing in >10% individuals, with minor allele frequency <5%, or Hardy-

Weinberg equilibrium exact test P values <1e-10. A total of 64,103 SNPs and all but one

G2 individual were retained. Prior to mapping, we LD-filtered SNPs with r^2 >0.9 using a

window of 5 SNPs and a step size of 1 SNP. We retain 32,625 SNPs for mapping.

Hybrid index calculation

For each G0 and G2 mouse, we estimated a hybrid index – defined as the

percentage of *M. m. musculus* ancestry. We identified ancestry-informative SNP markers

by comparing GigaMUGA data from ten individuals each from two wild-derived

outbred stocks of *M. m. musculus* (Kazakhstan and Czech Republic) and two of *M. m.*

domesticus (Germany and France) maintained at the Max Planck Institute for

601 Evolutionary Biology (L.M. Turner and B. Payseur, unpublished data). We classified
602 SNPs as ancestry informative if they had a minimum of 10 calls per subspecies, the
603 major allele differed between *musculus* and *domesticus*, and the allele frequency
604 difference between subspecies was > 0.3. A total of 48,361 quality-filtered SNPs from the
605 G0/G2 genotype data were informative, including 8,775 SNPs with fixed differences
606 between subspecies samples.

607 *Correlation between host relatedness and microbiome structure*

608 To investigate if host relatedness is correlated with individual variation in
609 microbiome composition, we computed a centered relatedness matrix using the 32,625
610 filtered SNPs with GEMMA (v 0.98.1; Zhou and Stephens, 2012) and microbial
611 composition-based kinship matrix among individuals based on relative bacterial
612 abundances (Chen et al., 2018). The kinship matrix was calculated with the formula:

$$Kinship = 1/p \sum_{i=1}^{p} (x_i - 1_n \bar{x}_i)(x_i - 1_n \bar{x}_i)^T$$

613 where **X** denotes the $n \times p$ matrix of genotypes or relative abundances, x_i as its ith
614 column representing the genotypes of ith SNP or the relative abundance of the ith ASV,
615 $\bar{x}_i$ as the sample mean and 1_n as a $n \times 1$ vector of 1's. We used a Mantel test with the
616 Spearman's correlation to test for correlation between the host SNP-based kinship and
617 microbial composition-based kinship using 10,000 permutations.

618 *SNP-based heritability of microbial abundances*

619 We calculated SNP-based heritabilities for bacterial abundances using a linear
620 mixed model implemented in the lme4qtl R package (version 0.2.2; Ziyatdinov et al.,
621 2018). The SNP-based heritability is expressed as:

$$h^2 = \frac{\sigma_g^2}{\sigma_g^2 + \sigma_m^2 + \sigma_e^2}$$

622 where σ_g^2 is the genetic variance estimated by K_{SNP}, σ_m^2 variance of the mating pair
623 component, and σ_e^2 the variance due to residual environmental factors. We determined
624 significance of the heritability estimates using exact likelihood ratio tests, following
625 Supplementary Note 3 in Ziyatdinov et al., 2018, using the exactLRT() function of the R
626 package RLRsim (version 3.1-6; Fabian et al., 2008).

627 *Genome-wide association mapping*

628 Prior to mapping, we inverse logistic transformed bacterial abundances using the
629 inv.logit function from the R package gtools (version 3.9.2; Gregory R. Warnes, 2020).

630 We performed association mapping in the R package lme4qtl (version 0.2.2;
631 Ziyatdinov et al., 2018) with the following linear mixed model:

$$y_i = \mu + a_i X_{ij}^a + d_i X_{ij}^d + Wu + e$$

632

633   where y_j is the phenotypic value of the jth individual; μ is the mean, X_{aij} the additive
634 and X_{dij} the dominance genotypic index values coded as for individual j at locus i. a and
635 d indicate fixed additive and dominance effects, W indicates random effects mating pair
636 and kinship matrix, plus residual error e.

637   We estimated additive and dominance effects separately because we expected to
638 observe underdominance and overdominance in our hybrid mapping population, as
639 well as additive effects, and aimed to estimate their relative importance. To model the
640 additive effect (i.e. 1/2 distance between homozygous means), genotypes at each locus,
641 i, were assigned additive index values ($X^a \in 1, 0, -1$) for AA, AB, BB, respectively, with A
642 indicating the major allele and B the minor allele. To model dominance effects (i.e.
643 heterozygote mean - midpoint of homozygote means), genotypes were assigned
644 dominance index values ($X^d \in 0, 1$) for homozygotes and heterozygotes, respectively.

645   We included mating pair as a random effect to account for maternal effects and cage
646 effects, as male litter mates are kept together in a cage after weaning. We included
647 kinship coefficient as a random effect in the model to account for population and family
648 structure. To avoid proximal contamination, we used a leave-one-chromosome-out
649 approach, that is, when testing each single-SNP association we used a relatedness matrix
650 omitting markers from the same chromosome (Parker et al., 2014). Hence, for testing
651 SNPs on each chromosome, we calculated a centered relatedness matrix using SNPs
652 from all other chromosomes with GEMMA (v0.97; Zhou and Stephens, 2012). We
653 calculated P values for single-SNP associations by comparing the full model to a null
654 model excluding fixed effects. Code for performing the mapping is available at https:/
655 /github.com/sdoms/mapping_scripts.

656   We evaluated significance of SNP-trait associations using two thresholds; first, we
657 used a genome-wide threshold for each trait, where we corrected for multiple testing
658 across markers using the Bonferroni method (Abdi, 2007). Second, as bacteria interact
659 with each other within the gut as members of a community, bacterial abundances are
660 non-independent, so we calculated a study-wide threshold dividing the genome-wide
661 threshold by the number of effective taxa included. We used matSpDlite (Nyholt, 2019;
662 Li and Ji, 2005; Qin et al., 2020) to estimate the number of effective bacterial taxa based
663 on eigenvalue variance.

664   To estimate the genomic interval represented by each significant LD-filtered SNP,
665 we report significant regions defined by the most distant flanking SNPs in the full pre-
666 LD-filtered genotype dataset showing $r^2 > 0.9$ with each significant SNP. We combined
667 significant regions less than 10 Mb apart into a single region. Genes situated in

668 significant regions were retrieved using biomaRt (Steffen Durinck, 2009), and the mm10

669 mouse genome.

670 *Dominance analyses*

671 We classified dominance for SNPs with significant associations on the basis of the

672 d/a ratio (Falconer, 1996) where d is the dominance effect, a the additive effect. As the

673 expected value under purely additive effects is 0. As our mapping population is a multi-

674 parental-line cross, and not all SNPs were ancestry-informative with respect to *musculus/*

675 *domesticus*, the sign of *a* effects is defined by the major allele within our mapping

676 population, which lacks clear biological interpretation. To provide more meaningful

677 values, we report d/|a|, such that a value of 1 = complete dominance of the allele

678 associated with higher bacterial abundance, and a value of -1 = complete dominance of

679 the allele associated with lower bacterial abundance. Values above 1 or below -1 indicate

680 over/underdominance. We classified effects of significant regions the following

681 arbitrary d/|a| ranges to classify dominance of significant regions (Burke et al., 2002;

682 Miller et al., 2014): underdominant <-1.25, high abundance allele recessive between -1.25

683 and -0.75, partially recessive between -0.75 and -0.25, additive between -0.25 and 0.25,

684 partially dominant between 0.25 and 0.75, dominant 0.75 and 1.25, and

685 overdominant >1.25.

686 *Gene ontology and network analysis*

687 The nearest genes up- and downstream of the significant SNPs were identified

688 using the locateVariants() function from the VariantAnnotation R package (version

689 1.34.0; Valerie et al., 2014) using the default parameters. A maximum of two genes per

690 locus were included (one upstream, and one downstream of a given SNP).

691 To investigate functions and interactions of candidate genes, we calculated a a

692 protein-protein interaction (PPI) network with STRING version 11 (Szklarczyk et al.,

693 2019), on the basis of a list of the closest genes to all SNPs with significant trait

694 associations. We included network edges with an interaction score >0.9, based on

695 evidence from fusion, neighborhood, co-occurrence, experimental, text-mining,

696 database, and co-expression. We exported this network to Cytoscape v 3.8.2 (Shannon et

697 al., 2003) for identification of highly interconnected regions using the MCODE

698 Cytoscape plugin (Bader and Hogue, 2003), and functional annotation of clusters using

699 the stringApp Cytoscape plugin (Doncheva et al., 2019).

700 We identified overrepresented KEGG pathways and human diseases using the

701 clusterprofiler R package (version 3.16.1; Yu et al., 2012). *P* values were corrected for

702 multiple testing using the Benjamini-Hochberg method. Pathways and diseases with an

703 adjusted *P* value < .05 were considered over-represented.

704 *Calculating overlap with other studies and over-representation of IBD genes*

705   To test for significant overlap with loci identified in previous mapping studies and
706 for over-representation of IBD genes, we used the tool *poverlap* (Brent Pedersen, 2013) to
707 compare observed overlap to random expectations based on 10,000 permutations of
708 significant regions. We identified genes within overlapping regions using the
709 locateVariants() function from the VariantAnnotation R package (version 1.34.0; Valerie
710 et al., 2014).

711 *Combination of results*

712   Hub genes SNP network and their first neighbors, the hub genes from the
713 'differentially expressed in GF mice'-network and their respective first neighbors, genes
714 found in both Mills et al. (2020) and other mouse QTL studies, closest genes to a SNP
715 found in Mills et al. (2020), genes situated in the 20 smallest intervals, six genes in the
716 two intervals with the lowest *P* values, twenty genes in intervals found in most different
717 taxa, genes situated in the region with most overlap within our study, and finally the
718 genes situated in the intervals that most frequently overlapped with other studies were
719 combined into on gene set and analyzed with STRING. Genes situated in the same
720 genomic locus were curated according to the number of edges in the STRING network.

724 **Supplementary Materials: Suppl. Fig 1-14, Suppl. Table 1: Heritability estimates,**
725 **Suppl. Table 2: Genome-wide significant associations, Suppl. Table 3: Study-wide**
726 **significant associations, Suppl. Table 4: Intervals smaller than 1Mb, Suppl. Table 5:**
727 **Over-represented KEGG pathways, Suppl. Table 6: IBD genes, Suppl. Table 7:**
728 **Candidate genes.**

729   **Acknowledgments**: We thank Diethard Tautz for generous support of mouse
730 breeding and Camilo Medina and the MPI-Plön mouse team for performing mouse
731 husbandry, and Katja Cloppenborg-Schmidt and Dr. Sven Künzel for their excellent
732 technical assistance. We thank Mathieu Groussin for assistance with cospeciation rate
733 data. Research funding for this project was provided by the Deutsche
734 Forschungsgemeinschaft Collaborative Research Center 1182, 'Origin and Function of
735 Metaorganisms' (J.F.B. and A.F.) and TU 500/2-1 to L.M.T, and by the Max Planck
736 Society (to D. Tautz).

737   **Author contributions**: Conceptualization: L.M.T., S.I., A.F., and J.F.B; Methodology:
738 L.M.T, J.F.B., S.D., S.I., A.F., and A.K.; Software: S.D., M.R., and L.M.T; Validation: S.D.,
739 M.R., A.K., and L.M.T; Formal Analysis: S.D., M.R., A.K., and L.M.T; Investigation: S.D.
740 and L.M.T.; Resources: S.D., H.F., and C.C; Writing - Original Draft: S.D.; Writing -

741 Review & Editing: S.D., L.M.T, and J.F.B; Visualization: S.D. and L.M.T; Supervision:
742 L.M.T., A.F., and J.F.B.; Project Administration: J.F.B.; Funding Acquisition: A.F., L.M.T,
743 and J.F.B.

744 **Conflicts of Interest: The authors declare no conflict of interest.**

745   **Supplementary figures**

746   **Supplementary figure 1**: Selection of taxa for mGWAS analysis. A scatter plot showing the
747   association of average relative abundance of taxa with their prevalence in the G2 mapping
748   population. Taxa retained for analysis are colored according to the originating core. The size of
749   each dot represents the number of individuals that have a median abundance higher than 0.2% of
750   the taxon. The dashed lines represent the thresholds of the core (vertical: median abundance>0.2%
751   and horizontal prevalence of 25 %.

752 **Supplementary figure 2**: Relative abundances of core genera in G2 mapping population.
753 Each vertical line represents one individual. Subcross (see supplementary figure 14) is indicated at
754 the top.

755 **Supplementary figure 3**: Host genetic relatedness calculated from SNP data (x-axis) correlat-
756 ed with microbial composition-based relatedness (y-axis) calculated from ASV abundances.
757 The blue line represents a linear regression fit to the data.

Supplementary figure 4: Correlation of SNP-based heritability estimates based on DNA (x-axis) or RNA (y-axis). The blue line represents a linear regression fit to the data. Red dashed line represents the identity line with a slope of 1.

761   **Supplementary figure 5**: Manhattan plots for ASV184 (*Dorea*) of the complete model (A), the
762   additive effect (B) or the dominance effect (C). SNPs passing the study-wide significance
763   threshold (solid line) are shown in dark blue, while genome-wide significant SNPs (dashed
764   line) are shown in light blue. In panel A, the closest gene to the SNP is shown for a subset of
765   significant SNPs.

Supplementary figure 6: Number of significantly associated loci per bacterial taxon. Loci with significant additive effects (add.P), dominance effects (dom.P) or effects in full model (P) are indicated.

769

770 **Supplementary figure 7**: Top ten hub genes of the protein-protein interaction (PPI) network
771 with the closest genes to the host SNPs significantly associated with bacterial abundances.
772 The nodes are colored according to hub gene rank from 1 (red) to 10 (yellow). Blue nodes are
773 the first neighbors.

 Supplementary figure 8: Genes belonging to over-represented KEGG pathways within the
host genes closest to significant SNPs from association analysis.

776 **Supplementary figure 9**: Enriched KEGG pathways among closest genes to significant SNPs from
777 association analysis. Node color indicates FDR-adjusted *P* value of enrichment and node size indi-
778 cates number of candidate genes in pathway.

779 **Supplementary figure 10**: Enriched human diseases among genes closest to significant SNPs from
780 association analysis.

Supplementary figure 11: STRING (Szklarczyk et al., 2019) protein-protein interaction network of proteins that are differentially expressed in the intestine (small intestine and colon) of germ-free (GF) mice compared to conventionally raised mice, found in the present study. The color of the network nodes indicates whether the QTL hit was found using the DNA abundances (green), RNA abundances (purple) or was found in both (orange). The shape represents if the gene of the protein was the closest gene to the significant SNP (rectangle), if the gene was also found in QTLs of other studies (octagon), a combination of both (diamond), or only differentially expressed in GF mice vs. conventionally raised mice. The node size expresses the number of taxa where the gene was found in a QTL. The edges represent protein-protein interactions, where the line thickness indicates the strength of the data support from text mining, experiments, databases, co-expression, gene-fusion, and co-occurrence.

802   **Supplementary figure 13**: Original protein protein interaction (PPI) network of 304 candi-
803   date genes closest to SNPs significantly associated with bacterial abundances. Generated in
804   STRING (Szklarczyk et al., 2019) and Cytoscape (Shannon et al., 2003).

FS3   FS5   HO3   HO6   HA1   HA2   TU3   TU4

52.8 %   76.8 %   43.9 %   39.5 %   72.7 %   73.7 %   77.4 %   76.0 %

X

G1

A   B   E   F   C   D   G   H

G2

A/F   B/G   E/B   F/H   C/A   D/E   G/D   H/C

61.5 %   66.9 %   75.3 %   66.5 %   59.2 %   69.4 %   59.5 %   66.2 %

Supplementary figure 14: Overview of the intercross design. G0 mice are from eight partially
inbred lines derived from mice wild-caught in four hybrid zone sites. Hybrid index - the per-
centage of *musculus* alleles - is reported as the mean for the G0 mice from each line (top), or
mean of 40 G2s from each subcross (bottom). We performed eight G1 crosses with one line
with hybrid index ~50% (purple shades) and one line with hybrid index >50% (green
shades); color on the left side of mouse diagram indicates dam line and right side indicates
sire line. Next, G1 mice were crossed in eight combinations such that each G2 mouse had one
grandparent from each of the four breeding stocks, indicated by colors of mouse diagram,
and representative chromosomes below. Tail color indicates Y chromosome strain, and oval
indicates mitochondrial strain.

815 References

Abdi, Hervé (2007), 'The Bonferonni and Šidák Corrections for Multiple Comparisons', in Salkind, Neil J. (ed.), (Ency-
clopedia of Measurement and Statistics, SAGE), 9.

Alhasson, Firas, et al. (2017), 'Altered gut microbiome in a mouse model of Gulf War Illness causes neuroinflammation
and intestinal injury via leaky gut and TLR4 activation', *PLoS One*, 12 (3), e0172914.

Amato, Katherine R, et al. (2019), 'Evolutionary trends in host physiology outweigh dietary niche in structuring primate
gut microbiomes', *The ISME journal*, 13 (3), 576-87.

Backhed, F., et al. (2004), 'The gut microbiota as an environmental factor that regulates fat storage', *Proceedings of the*
National Academy of Sciences, 101 (44), 15718-23.

Bader, Gary D. and Christopher WV Hogue (2003), 'An automated method for finding molecular complexes in large pro-
tein interaction networks', *BMC Bioinformatics*, 4 (1), 2.

Barton, Nicholas H. and Peter D. Keightley (2002), 'Understanding quantitative genetic variation', *Nat. Rev. Genet.*, 3 (1),
11-21.

Beavis, WD (1994), 'The power and deceit of QTL experiments: lessons from comparative QTL studies', Proceedings of
the forty-ninth annual corn and sorghum industry research conference 250 266.

Belheouane, Meriem, et al. (2017), 'Improved detection of gene-microbe interactions in the mouse skin microbiota using
high-resolution QTL mapping of 16S rRNA transcripts', *Microbiome*, 5 (1), 1-17.

Benson, Andrew K., et al. (2010), 'Individuality in gut microbiota composition is a complex polygenic trait shaped by
multiple environmental and host genetic factors', *Proceedings of the National Academy of Sciences of the United States of*
America

Proc. Natl. Acad. Sci. U.S.A., 107 (44), 18933-38.

Bonder, Marc Jan, et al. (2016), 'The effect of host genetics on the gut microbiome', *Nat. Genet.*, 48 (11), 1407-12.

Brent Pedersen, Joe Brown (2013), 'poverlap: significance testing over interval overlaps',

Brooks, AW, et al. (2016), 'Phylosymbiosis: Relationships and Functional Effects of Microbial Communities across Host
Evolutionary History.', *PLoS Biol.*, 14 (11), e2000225.

Brucker, RM and SR Bordenstein (2012a), 'Speciation by symbiosis.', *Trends Ecol Evol*, 27 (8), 443-51.

Brucker, Robert M and Seth R Bordenstein (2012b), 'The roles of host evolutionary relationships (genus: Nasonia) and
development in structuring microbial communities', *Evolution: International Journal of Organic Evolution*, 66 (2),
349-62.

Burke, John M., et al. (2002), 'Genetic Analysis of Sunflower Domestication', *Genetics*, 161 (3), 1257-67.

Callahan, Benjamin J (2016), 'DADA2 pipeline', *DADA2*,

Callahan, Benjamin J, et al. (2016), 'DADA2: High resolution sample inference from Illumina amplicon data', *Nature*
methods

Nat Methods, 13 (7), 581-83.

Campbell, JH, et al. (2012), 'Host genetic and environmental effects on mouse intestinal microbiota.', *ISME J*, 6 (11),
2033-44.

Cani, Patrice D., et al. (2008), 'Changes in gut microbiota control metabolic endotoxemia-induced inflammation in high-
fat diet-induced obesity and diabetes in mice', *Diabetes*, 57 (6), 1470-81.

Carding, Simon, et al. (2015), 'Dysbiosis of the gut microbiota in disease', *Microb. Ecol. Health Dis.*, 26

Cardoso, JC, et al. (2012), 'Feeding and the rhodopsin family g-protein coupled receptors in nematodes and arthropods.',
Front Endocrinol (Lausanne), 3 157.

Carmody, RN, et al. (2015), 'Diet dominates host genotype in shaping the murine gut microbiota.', *Cell Host Microbe*, 17
(1), 72-84.

Castoldi, Angela, et al. (2015), 'They Must Hold Tight: Junction Proteins, Microbiota And Immunity In Intestinal Mu-
cosa', *Current Protein & Peptide Science*

Curr Protein Pept Sci, 16 (7), 655-71.

Chang, Christopher C, et al. (2015), 'Second-generation PLINK: rising to the challenge of larger and richer datasets',
GigaScience

GigaSci, 4 (1), 7.

Chen, Congying, et al. (2018), 'Contribution of Host Genetics to the Variation of Microbial Composition of Cecum
Lumen and Feces in Pigs', *Frontiers in Microbiology*

Front. Microbiol., 9

Chen, Haiwei, et al. (2019), 'A forward chemical genetic screen reveals gut microbiota metabolites that modulate host
physiology', *Cell*, 177 (5), 1217-1231.e18.

Chu, Hiutung and Sarkis K Mazmanian (2013), 'Innate immune recognition of the microbiota promotes host-microbial
symbiosis', *Nat. Immunol.*, 14 (7), 668-75.

Chung, HJ, et al. (2018), 'Gut Microbiota as a Missing Link Between Nutrients and Traits of Human.', *Front Microbiol*, 9
1510.

Clapp, M, et al. (2017), 'Gut microbiota's effect on mental health: The gut-brain axis.', *Clin Pract*, 7 (4), 987.

Cohen, Louis J., et al. (2017), 'Commensal bacteria make GPCR ligands that mimic human signalling molecules', *Nature*,
549 (7670), 48-53.

Cole, JR, et al. (2014), 'Ribosomal Database Project: data and tools for high throughput rRNA analysis.', *Nucleic Acids
Res.*, 42 (Database issue), D633-42.

Colosimo, Dominic A., et al. (2019), 'Mapping Interactions of Microbial Metabolites with Human G-Protein-Coupled Re-
ceptors', *Cell Host & Microbe*, 26 (2), 273-282.e7.

Cox, Laura M. and Howard L. Weiner (2018), 'Microbiota Signaling Pathways that Influence Neurologic Disease', *Neu-
rotherapeutics*, 15 (1), 135-45.

Daniel, Noëmie, Emelyne Lécuyer, and Benoit Chassaing (2021), 'Host/microbiota interactions in health and diseases—
Time for mucosal microbiology', *Mucosal Immunology*, 1-11.

Davenport, Emily R., et al. (2015), 'Genome-Wide Association Studies of the Human Gut Microbiota', *PLoS One*, 10
(11), e0140301.

Davenport, Emily R. (2020), 'Genetic Variation Shapes Murine Gut Microbiota via Immunity', *Trends in Immunology*, 41
(1), 1-3.

Deaver, Jessica A., Sung Y. Eum, and Michal Toborek (2018), 'Circadian Disruption Changes Gut Microbiome Taxa and
Functional Gene Composition', *Frontiers in Microbiology*

Front Microbiol, 9 737.

Delzenne, Nathalie M., et al. (2011), 'Targeting gut microbiota in obesity: effects of prebiotics and probiotics', *Nature Re-
views. Endocrinology*

Nat Rev Endocrinol, 7 (11), 639-46.

Doncheva, Nadezhda T., et al. (2019), 'Cytoscape StringApp: Network Analysis and Visualization of Proteomics Data',
Journal of Proteome Research

J Proteome Res, 18 (2), 623-32.

Erdman, S.E. and T. Poutahidis (2016), 'Microbes and Oxytocin', *131* (Int. Rev. Neurobiol., Elsevier), 91-126.

Fabian, Scheipl, Greven Sonja, and Kuechenhoff Helmut (2008), 'Size and power of tests for a zero random effect vari-
ance or polynomial regression in additive and linear mixed models.', *Computational Statistics & Data Analysis*, 52 (7),
3283-99.

Falconer, D. S (1996), *Introduction to quantitative genetics*, (Harlow, England: Prentice Hall).

Flux, M. C. and Christopher A. Lowry (2020), 'Finding intestinal fortitude: Integrating the microbiome into a holistic
view of depression mechanisms, treatment, and resilience', *Neurobiology of Disease*

Microbiome in neurological and psychiatric disease

Neurobiology of Disease, 135 104578.

Fonken, Laura K., et al. (2010), 'Light at night increases body mass by shifting the time of food intake', *Proceedings of
the National Academy of Sciences*

PNAS, 107 (43), 18664-69.

Foster, Jane A., Linda Rinaman, and John F. Cryan (2017), 'Stress & the gut-brain axis: Regulation by the microbiome',
Neurobiology of Stress, 7 124-36.

Fukata, Masayuki and Moshe Arditi (2013), 'The role of pattern recognition receptors in intestinal inflammation', *Mucos-
al immunology*, 6 (3), 451-63.

Gastelum, C, et al. (2021), 'Adaptive Changes in the Central Control of Energy Homeostasis Occur in Response to Varia-
tions in Energy Status.', *Int J Mol Sci*, 22 (5), 2728.

Gautam, D, et al. (2006), 'A critical role for beta cell M3 muscarinic acetylcholine receptors in regulating insulin release
and blood glucose homeostasis in vivo.', *Cell Metab*, 3 (6), 449-61.

Geraldes, A, et al. (2008), 'Inferring the history of speciation in house mice from autosomal, X-linked, Y-linked and mito-
chondrial genes.', *Mol Ecol*, 17 (24), 5349-63.

Gevers, Dirk, et al. (2014), 'The treatment-naive microbiome in new-onset Crohn's disease', *Cell host & microbe*, 15 (3),
382-92.

Gogarten, Jan F, et al. (2021), 'Primate phageomes are structured by superhost phylogeny and environment', *Proceedings*
of the National Academy of Sciences, 118 (15),

Goodrich, Julia K., et al. (2014), 'Human genetics shape the gut microbiome', *Cell*, 159 (4), 789-99.

Goodrich, Julia K., et al. (2016), 'Genetic Determinants of the Gut Microbiome in UK Twins', *Cell host & microbe*,

Gould, AL, et al. (2018), 'Microbiome interactions shape host fitness.', *Proc. Natl. Acad. Sci. U S A*, 115 (51), E11951-60.

Gregory R. Warnes, Ben Bolker and Thomas Lumley (2020), 'gtools: Various R Programming Tools',

Grieneisen, L, et al. (2021), 'Gut microbiome heritability is nearly universal but environmentally contingent.', *Science*,
373 (6551), 181-86.

Groussin, Mathieu, et al. (2017), 'Unraveling the processes shaping mammalian gut microbiomes over evolutionary time',
Nature Comm., 8 (1), 14319.

Hehemann, Jan-Hendrik, et al. (2010), 'Transfer of carbohydrate-active enzymes from marine bacteria to Japanese gut mi-
crobiota', *Nature*, 464 (7290), 908-12.

Hollander, Daniel and Jonathan D. Kaunitz (2020), 'The "Leaky Gut": Tight Junctions but Loose Associations', *Digestive*
Diseases and Sciences

Dig Dis Sci, 65 (5), 1277-87.

Hua, Yan, et al. (2020), 'Gut microbiota and fecal metabolites in captive and wild North China leopard (Panthera pardus
japonensis) by comparsion using 16 s rRNA gene sequencing and LC/MS-based metabolomics', *BMC Veterinary Re-*
search, 16 (1),

Hughes, David A., et al. (2020), 'Genome-wide associations of human gut microbiome variation and implications for
causal inference analyses', *Nature Microbiology*, 5 (9), 1079-87.

Ishida, Sachiko, et al. (2020), 'Genome-wide association studies and heritability analysis reveal the involvement of host
genetics in the Japanese gut microbiota', *Communications Biology*

Commun Biol, 3

Kelly, John R., et al. (2015), 'Breaking Down the Barriers: The Gut Microbiome, Intestinal Permeability and Stress-relat-
ed Psychiatric Disorders', *Frontiers in Cellular Neuroscience*

Front. Cell. Neurosci., 9

Kemis, Julia H., et al. (2019), 'Genetic determinants of gut microbiota composition and bile acid profiles in mice', *PLoS*
Genet., 15 (8), e1008073.

Khan, Farhat, et al. (2021), 'IBDDB: a manually curated and text-mining-enhanced database of genes involved in inflam-
matory bowel disease', *Database*, 2021

Klug-Micu, GM, et al. (2013), 'CD40 ligand and interferon-γ induce an antimicrobial response against Mycobacterium
tuberculosis in human monocytes.', *Immunology*, 139 (1), 121-28.

Kohl, KD and MD Dearing (2014), 'Wild-caught rodents retain a majority of their natural gut microbiota upon entrance
into captivity.', *Environ Microbiol Rep*, 6 (2), 191-95.

Korach-Rechtman, H, et al. (2019), 'Murine Genetic Background Has a Stronger Impact on the Composition of the Gut
Microbiota than Maternal Inoculation or Exposure to Unlike Exogenous Microbiota.', *Appl. Environ. Microbiol.*, 85 (18),
e00826-19.

Kovacs, Amir, et al. (2011), 'Genotype is a stronger determinant than sex of the mouse gut microbiota', *Microbial ecolo-*
gy, 61 (2), 423-28.

Kurilshikov, Alexander, et al. (2021), 'Large-scale association analyses identify host factors influencing human gut micro-
biome composition', *Nat. Genet.*, 53 (2), 156-65.

Leamy, Larry J, et al. (2014), 'Host genetics and diet, but not immunoglobulin A expression, converge to shape composi-
tional features of the gut microbiome in an advanced intercross population of mice', *Genome Biology*

Genome Biol, 15 (12),

Ley, RE, et al. (2006), 'Microbial ecology: human gut microbes associated with obesity.', *Nature*, 444 (7122), 1022-23.

Li, J. and L. Ji (2005), 'Adjusting multiple testing in multilocus analyses using the eigenvalues of a correlation matrix',
Heredity, 95 (3), 221-27.

Lim, SJ and SR Bordenstein (2020), 'An introduction to phylosymbiosis.', *Proc Biol Sci*, 287 (1922), 20192900.

Linnenbrink, Miriam, et al. (2013), 'The role of biogeography in shaping diversity of the intestinal microbiota in house
mice', *Molecular Ecology*, 22 (7), 1904-16.

Lynch, SV and O Pedersen (2016), 'The Human Intestinal Microbiome in Health and Disease.', *N. Engl. J. Med.*, 375

972   (24), 2369-79.

973 Malaguarnera, L (2020), 'Vitamin D and microbiota: Two sides of the same coin in the immunomodulatory aspects.', *Int*
974 *Immunopharmacol*, 79 106112.

975 McKnite, Autumn M., et al. (2012), 'Murine Gut Microbiota Is Defined by Host Genetics and Modulates Variation of
976 Metabolic Traits', *PLoS One*, 7 (6),

977 McMurdie, Paul J and Susan Holmes (2013), 'phyloseq: an R package for reproducible interactive analysis and graphics
978 of microbiome census data', *PLoS One*, 8 (4), e61217.

979 Metwaly, Amira, et al. (2020), 'Integrated microbiota and metabolite profiles link Crohn's disease to sulfur metabolism',
980 *Nature Comm.*, 11 (1),

981 Walsh, Michael Lynch and Bruce (1998), *Genetics and Analysis of Quantitative Traits*, (Sunderland, MA: Sinauer).

982 Miller, Craig T., et al. (2014), 'Modular Skeletal Evolution in Sticklebacks Is Controlled by Additive and Clustered
983 Quantitative Trait Loci', *Genetics*, 197 (1), 405-20.

984 Mills, Robert H., et al. (2020), 'Organ-level protein networks as a reference for the host effects of the microbiome',
985 *Genome Research*

986 *Genome Res.*, 30 (2), 276-86.

987 Moeller, Andrew H., et al. (2016), 'Cospeciation of gut microbiota with hominids', *Science (New York, N.Y.)*

988 *Science*, 353 (6297), 380-82.

989 Moeller, Andrew H., et al. (2019), 'Experimental Evidence for Adaptation to Species-Specific Gut Microbiota in House
990 Mice', *mSphere*, 4

991 Moran, Nancy A. and Daniel B. Sloan (2015), 'The Hologenome Concept: Helpful or Hollow', *PLoS Biol.*, 13 (12),
992 e1002311.

993 Morgan, Andrew P., et al. (2015), 'The Mouse Universal Genotyping Array: From Substrains to Subspecies', *G3:*
994 *GenesGenomesGenetics*

995 *G3 (Bethesda)*, 6 (2), 263-79.

996 Moya, Andrés and Manuel Ferrer (2016), 'Functional Redundancy-Induced Stability of Gut Microbiota Subjected to Dis-
997 turbance', *Trends Microbiol.*, 24 (5), 402-13.

998 Nagpal, Ravinder, et al. (2020), 'Role of TRP Channels in Shaping the Gut Microbiome', *Pathogens*, 9

999 Neumann, Philipp-Alexander, et al. (2014), 'Gut Commensal Bacteria and Regional Wnt Gene Expression in the Proxi-
1000 mal Versus Distal Colon', *The American Journal of Pathology*

1001 *Am J Pathol*, 184 (3), 592-99.

1002 Nicholson, Jeremy K., et al. (2012), 'Host-gut microbiota metabolic interactions', *Science (New York, N.Y.)*

1003 *Science*, 336 (6086), 1262-67.

1004 Nyholt, Dale R. (2019), 'matSpD local version - Statistical and Genomic Epidemiology Laboratory (SGEL)',

1005 O'Connor, Annalouise, et al. (2014), 'Responsiveness of cardiometabolic-related microbiota to diet is influenced by host
1006 genetics', *Mammalian Genome*, 25 (11), 583-99.

1007 Ochman, Howard, et al. (2010), 'Evolutionary relationships of wild hominids recapitulated by gut microbial communi-
1008 ties', *PLoS Biol.*, 8 (11), e1000546.

1009 Org, Elin, et al. (2015), 'Genetic and environmental control of host-gut microbiota interactions', *Genome Research*

1010 *Genome Res.*, 25 (10), 1558-69.

1011 Org, Elin and Aldons J. Lusis (2018), 'Using the natural variation of mouse populations to understand host-gut microbio-
1012 me interactions', *Drug discovery today. Disease models*

1013 *Drug Discov Today Dis Models*, 28 61-71.

1014 Ott, SJ, et al. (2004), 'Reduction in diversity of the colonic mucosa associated bacterial microflora in patients with active
1015 inflammatory bowel disease', *Gut*, 53 (5), 685-93.

1016 Pallares, LF, et al. (2014), 'Use of a natural hybrid zone for genomewide association mapping of craniofacial traits in the
1017 house mouse.', *Mol Ecol*, 23 5756-70.

1018 Pandey, Shubhi, Jagannath Maharana, and Arun K. Shukla (2019), 'The Gut Feeling: GPCRs Enlighten the Way', *Cell*
1019 *Host & Microbe*, 26 (2), 160-62.

1020 Papa, Eliseo, et al. (2012), 'Non-invasive mapping of the gastrointestinal microbiota identifies children with inflammatory
1021 bowel disease', *PLoS One*, 7 (6), e39242.

1022 Parker, Bianca J., et al. (2020), 'The Genus Alistipes: Gut Bacteria With Emerging Implications to Inflammation, Cancer,
1023 and Mental Health', *Frontiers in Immunology*

Front. Immunol., 11

Parker, CC, et al. (2014), 'High-resolution genetic mapping of complex traits from a combined analysis of F2 and ad-
vanced intercross mice.', *Genetics*, 198 (1), 103-16.

Peier, Andrea, et al. (2009), 'The Antiobesity Effects of Centrally Administered Neuromedin U and Neuromedin S Are
Mediated Predominantly by the Neuromedin U Receptor 2 (NMUR2)', *Endocrinology*, 150 (7), 3101-9.

Peng, Zhi, et al. (2020), 'The Gut Microbiome Is Associated with Clinical Response to Anti–PD-1/PD-L1 Immunotherapy
in Gastrointestinal Cancer', *Cancer Immunology Research*

Cancer Immunol Res, 8 (10), 1251-61.

Qin, Junjie, et al. (2012), 'A metagenome-wide association study of gut microbiota in type 2 diabetes', *Nature*, 490
(7418), 55-60.

Qin, Youwen, et al. (2020), 'Combined effects of host genetics and diet on human gut microbiota and incident disease in a
single population cohort',

Rapp, K (1972), 'HAN-rotation, a new system for rigorous outbreeding', *Z. Versuchstierk.*, 14 133-42.

Rausch, P, et al. (2016), 'Analysis of factors contributing to variation in the C57BL/6J fecal microbiota across German an-
imal facilities.', *Int. J. Med. Microbiol.*, 306 (5), 343-55.

Rehman, A, et al. (2016), 'Geographical patterns of the standing and active human gut microbiome in health and IBD.',
Gut, 65 (2), 238-48.

Reichardt, Nicole, et al. (2018), 'Specific substrate-driven changes in human faecal microbiota composition contrast with
functional redundancy in short-chain fatty acid production', *The ISME Journal*, 12 (2), 610-22.

Ricklin, D, et al. (2016), 'Complement component C3 - The "Swiss Army Knife" of innate immunity and host defense.',
Immunol. Rev., 274 (1), 33-58.

Rieseberg, Loren H, Margaret A Archer, and Robert K Wayne (1999), 'Transgressive segregation, adaptation and specia-
tion', *Heredity*, 83 (4), 363-72.

Rolig, AS, et al. (2015), 'Individual Members of the Microbiota Disproportionately Modulate Host Innate Immune Re-
sponses.', *Cell Host Microbe*, 18 (5), 613-20.

Rosshart, Stephan P., et al. (2017), 'Wild Mouse Gut Microbiota Promotes Host Fitness and Improves Disease Resis-
tance', *Cell*, 171 (5), 1015-1028.e13.

Roth, TL, et al. (2019), 'Reduced Gut Microbiome Diversity and Metabolome Differences in Rhinoceros Species at Risk
for Iron Overload Disorder.', *Front Microbiol*, 10 2291.

Rowland, Ian, et al. (2018), 'Gut microbiota functions: metabolism of nutrients and other food components', *European
Journal of Nutrition*

Eur J Nutr, 57 (1), 1-24.

Rühlemann, Malte Christoph, et al. (2021), 'Genome-wide association study in 8,956 German individuals identifies influ-
ence of ABO histo-blood groups on gut microbiome', *Nat. Genet.*, 1-9.

Saito, Yumiko, et al. (1999), 'Molecular characterization of the melanin-concentrating-hormone receptor', *Nature*, 400
(6741), 265-69.

Sarkar, Amar, et al. (2020), 'The role of the microbiome in the neurobiology of social behaviour', *Biol. Rev.*, 95 (5),
1131-66.

Sethi, JK and AJ Vidal-Puig (2008), 'Wnt signalling at the crossroads of nutritional regulation.', *Biochem. J.*, 416 (2),
e11-3.

Shannon, Paul, et al. (2003), 'Cytoscape: a software environment for integrated models of biomolecular interaction net-
works', *Genome Research*

Genome Res, 13 (11), 2498-504.

Shi, L and BP Tu (2015), 'Acetyl-CoA and the regulation of metabolism: mechanisms and consequences.', *Curr. Opin.
Cell Biol.*, 33 125-31.

Singh, Parul, et al. (2020), 'The potential role of vitamin D supplementation as a gut microbiota modifier in healthy indi-
viduals', *Scientific Reports*, 10 (1), 21641.

Škrabar, N, et al. (2018), 'Using the Mus musculus hybrid zone to assess covariation and genetic architecture of limb
bone lengths.', *Mol Ecol Resour*, 18 (4), 908-21.

Smith, Ashley E., et al. (2019), 'Binge-Type Eating in Rats is Facilitated by Neuromedin U Receptor 2 in the Nucleus Ac-
cumbens and Ventral Tegmental Area', *Nutrients*, 11 (2), 327.

Snijders, Antoine M., et al. (2016), 'Influence of early life exposure, host genetics and diet on the mouse gut microbiome
and metabolome', *Nature Microbiology*, 2 16221.

Spor, Aymé, Omry Koren, and Ruth Ley (2011), 'Unravelling the effects of the environment and host genotype on the gut

microbiome', *Nature Reviews Microbiology*, 9 (4), 279-90.

Sriram, K and PA Insel (2018), 'G Protein-Coupled Receptors as Targets for Approved Drugs: How Many Targets and
How Many Drugs', *Mol. Pharmacol.*, 93 (4), 251-58.

Steffen Durinck, Paul T. Spellman, Ewan

Birney and Wolfgang Huber (2009), 'Mapping identifiers for the integration of genomic datasets with the

R/Bioconductor package biomaRt.', *Nature Protocols*, 4 1184-91.

Suzuki, TA (2017), 'Links between Natural Variation in the Microbiome and Host Fitness in Wild Mammals.', *Integr
Comp Biol*, 57 (4), 756-69.

Suzuki, TA, et al. (2020), 'The gut microbiota and Bergmann's rule in wild house mice.', *Mol Ecol*, 29 (12), 2300-11.

Suzuki, Taichi A and Ruth E Ley (2020), 'The role of the microbiota in human genetic adaptation', *Science*, 370 (6521),

Suzuki, Taichi A., et al. (2019), 'Host genetic determinants of the gut microbiota of wild mice', *Molecular Ecology*, 28
(13), 3197-207.

Szklarczyk, Damian, et al. (2019), 'STRING v11: protein-protein association networks with increased coverage, support-
ing functional discovery in genome-wide experimental datasets', *Nucleic Acids Research*

Nucleic Acids Res, 47 (D1), D607-13.

Tanahashi, Yasuyuki, et al. (2009), 'Multiple muscarinic pathways mediate the suppression of voltage-gated Ca2+
channels in mouse intestinal smooth muscle cells', *Br. J. Pharmacol.*, 158 (8), 1874-83.

Taras, David, et al. (2002), 'Reclassification of Eubacterium formicigenerans Holdeman and Moore 1974 as Dorea formi-
cigenerans gen. nov., comb. nov., and description of Dorea longicatena sp. nov., isolated from human faeces.', *Internatio-
nal Journal of Systematic and Evolutionary Microbiology*, 52 (2), 423-28.

Thaiss, Christoph A., et al. (2014), 'Transkingdom control of microbiota diurnal oscillations promotes metabolic home-
ostasis', *Cell*, 159 (3), 514-29.

Thaiss, Christoph A., Maayan Levy, and Eran Elinav (2015a), 'Chronobiomics: The Biological Clock as a New Principle
in Host–Microbial Interactions', *PLoS Pathog.*, 11 (10), e1005113.

Thaiss, Christoph A., et al. (2015b), 'A day in the life of the meta-organism: diurnal rhythms of the intestinal microbiome
and its host', *Gut Microbes*, 6 (2), 137-42.

Thaiss, Christoph A., et al. (2016), 'Microbiota Diurnal Rhythmicity Programs Host Transcriptome Oscillations', *Cell*,
167 (6), 1495-1510.e12.

Tian, Liang, et al. (2020), 'Deciphering functional redundancy in the human microbiome', *Nature Comm.*, 11 (1),

Toderici, M, et al. (2016), 'Identification of Regulatory Mutations in SERPINC1 Affecting Vitamin D Response Elements
Associated with Antithrombin Deficiency.', *PLoS One*, 11 (3), e0152159.

Townsend, KL, et al. (2012), 'Bone morphogenetic protein 7 (BMP7) reverses obesity and regulates appetite through a
central mTOR pathway.', *FASEB J.*, 26 (5), 2187-96.

Turnbaugh, Peter J., et al. (2009), 'A core gut microbiome in obese and lean twins', *Nature*, 457 (7228), 480-84.

Turnbaugh, PJ, et al. (2008), 'Diet-induced obesity is linked to marked but reversible alterations in the mouse distal gut
microbiome.', *Cell Host Microbe*, 3 (4), 213-23.

Turner, Leslie M., Denise J. Schwahn, and Bettina Harr (2012), 'Reduced Male Fertility Is Common but Highly Variable
in Form and Severity in a Natural House Mouse Hybrid Zone', *Evolution*, 66 (2), 443-58.

Turner, Leslie M. and Bettina Harr (2014), 'Genome-wide mapping in a house mouse hybrid zone reveals hybrid sterility
loci and Dobzhansky-Muller interactions', *eLife*, 3 e02504.

Turpin, W., et al. (2016), 'Association of host genome with intestinal microbial composition in a large healthy cohort',
Nat. Genet., 48 (11), 1413-17.

Vaga, Stefania, et al. (2020), 'Compositional and functional differences of the mucosal microbiota along the intestine of
healthy individuals', *Scientific Reports*, 10 (1),

Valerie, Obenchain, et al. (2014), 'VariantAnnotation: a Bioconductor package for exploration and annotation of genetic
variants', *Bioinformatics*, 30 (14), 2076-78.

Velloso, Licio A., Franco Folli, and Mario J. Saad (2015), 'TLR4 at the Crossroads of Nutrients, Gut Microbiota, and
Metabolic Inflammation', *Endocrine Reviews*

Endocr Rev, 36 (3), 245-71.

Wang, Jun, et al. (2015), 'Analysis of intestinal microbiota in hybrid house mice reveals evolutionary divergence in a ver-
tebrate hologenome', *Nature Communications*

Nat Commun, 6

Wang, Jun, et al. (2016), 'Genome-wide association analysis identifies variation in vitamin D receptor and other host fac-

1131 tors influencing the gut microbiota', *Nat. Genet.*, 48 (11), 1396-406.

1132 Wang, Y, et al. (2010), 'Regional mucosa-associated microbiota determine physiological expression of TLR2 and TLR4 in
1133 murine colon.', *PLoS One*, 5 (10), e13607.

1134 Weldon, L, et al. (2015), 'The Gut Microbiota of Wild Mice.', *PLoS One*, 10 (8), e0134643.

1135 Wu, Guangyan, et al. (2018), 'Light exposure influences the diurnal oscillation of gut microbiota in mice', *Biochemical*
1136 *and Biophysical Research Communications*

1137 *Biochem Biophys Res Commun*, 501 (1), 16-23.

1138 Yang, M, et al. (2020a), 'Mucosal-Associated Microbiota Other Than Luminal Microbiota Has a Close Relationship With
1139 Diarrhea-Predominant Irritable Bowel Syndrome.', *Front. Cell. Infect. Microbiol.*, 10 515614.

1140 Yang, Q, et al. (2020b), 'Role of Dietary Nutrients in the Modulation of Gut Microbiota: A Narrative Review.', *Nutrients*,
1141 12 (2), E381.

1142 Yasuda, K, et al. (2021), 'Elucidation of metabolic pathways of 25-hydroxyvitamin D3 mediated by CYP24A1 and
1143 CYP3A using Cyp24a1 knockout rats generated by CRISPR/Cas9 system.', *J. Biol. Chem.*, 296 100668.

1144 Yatsunenko, Tanya, et al. (2012), 'Human gut microbiome viewed across age and geography', *Nature*, 486 (7402),
1145 222-27.

1146 Yi, Z and GA Bishop (2015), 'Regulatory role of CD40 in obesity-induced insulin resistance.', *Adipocyte*, 4 (1), 65-69.

1147 Yu, Guangchuang, et al. (2012), 'clusterProfiler: an R package for comparing biological themes among gene clusters',
1148 *Omics: a journal of integrative biology*, 16 (5), 284-87.

1149 Zhou, Xiang and Matthew Stephens (2012), 'Genome-wide efficient mixed-model analysis for association studies', *Nat.*
1150 *Genet.*, 44 (7), 821-24.

1151 Ziyatdinov, Andrey, et al. (2018), 'lme4qtl: linear mixed models with flexible covariance structure for genetic studies of
1152 related individuals', *BMC Bioinformatics*, 19 (1), 1-5.