Article

Rotational Activity Around an Obstacle in 2D Cardiac Tissue in Presence of Cellular Heterogeneity

Pavel Konovalov [1], **Daria Mangileva**[1], **Arsenii Dokuchaev** [1], **Olga Solovyova** [1,2] **and Alexander V. Panfilov** [2,3]*

1   Institute of Immunology and Physiology UB RAS, 620049 Ekaterinburg, Russia
2   Ural Federal University, 620075 Ekaterinburg, Russia
3   Department of Physics and Astronomy, Ghent University, 9000 Ghent, Belgium
*   Correspondence: Alexander.Panfilov@ugent.be

Abstract: Waves of electrical excitation rotating around an obstacle is one of the important
mechanisms of dangerous cardiac arrhythmias occurring in the heart damaged by post-infarction
scar. Such a scar also has a border zone around it, which has electrophysiological properties
different from the rest of normal myocardial tissue. Spatial patterns of wave rotation in the
presence of such tissue heterogeneity are poorly studied. In this paper we perform a comprehensive
numerical study of various regimes of rotation of a wave in a plane layer of the ventricular tissue
around an obstacle surrounded by a gray zone. We use a TP06 cellular ionic model which
reproduces the electrophysiological properties of cardiomyocytes in the left ventricle of human
heart. We vary the extent of obstacle and gray zone and study the pattern of wave rotation and its
period. We observed different regimes of wave rotation that can be subdivided into several classes:
(1) functional rotation and (2) scar rotation regimes, which were identified in the previous studies,
and new (3) gray zone rotation regime: where the wave instead of rotation around the obstacle,
rotates around the gray zone (an area of tissue heterogeneity) itself. For each class, the period of
rotation is determined by different factors, which we discuss and quantify. We also found that
due to regional pathological remodeling of myocardial tissue, we can obtain additional regimes
associated with dynamical instabilities of two types which may affect or not affect the period of
rotation.

Keywords: re-entry; myocardial heterogeneity; cardiac modeling; myocardial infarction; infarct
border zone; gray zone

1. Introduction

Propagating non-linear waves occur in various types of excitable media of physical
[1], chemical [2,3] and biological nature[4,5]. These waves can also form vortices which
organize spatial excitation patterns in the medium and their onset can have important
consequences. For example, rotating waves in the heart is the main mechanism of
dangerous cardiac arrhythmias [6,7] and study such regimes in the heart is important
area of research in cardiac electrophysiology. In some patients, arrhythmias may arise
from myocardial infarction, i.e. condition when due to poor blood supply a region of the
heart is damaged. As a result, electrical waves propagating through the heart break and
form vortices rotating in/around the damaged area [8,9].
From mathematical point such regimes can be viewed as rotation of non-linear
waves around an obstacle. This is one of the classical regimes studied in the theory of
excitable media. Such regimes were described in cellular automata models as early as in
1946 [10]. Later they were studied in reaction-diffusion models of excitable media [11].
More recently interest in such regimes, especially on rotation around obstacle of small
size was initiated by the concept of anchoring of vortices proposed by Davidenko and
co-authors [12]. Such processes were also studied using theoretical approaches [13,14].

Unfortunately, direct application of such studies to arrhythmias which occur around
infarcted area is difficult, due to the following problem. Post-infarction scar area in the
myocardial tissue has a complex geometry and functional heterogeneity [15]. It contains
a compact scar region, which is similar to unexcitable obstacle. Such compact scar is
surrounded by a border zone, which has both cellular and tissue properties different from
the rest of normal cardiac tissue [16]. This zone is also called gray zone (GZ) because of
its intermediate signal intensity in clinical MRI [17]. In mathematical sense this problem
can be viewed as wave rotation around an obstacle surrounded by a heterogeneous
media. There is a study by Bernus and co-authors [18] on the heterogeneity in the
gray zone which triggers such rotating wave. But this study focused on initiation of
arrhythmia, and it does not describe regimes of wave rotation after initiation. Such
rotation regimes are poorly understood.
Main aim of this paper is to perform a generic numerical study of this problem. In
this way, we use a simplified 2D representation of ventricular tissue with an obstacle and
gray zone of simple circular geometry, which gives us only two parameters (obstacle and
gray zone radii) to vary the extent of injury. In particular, we study rotation of waves in
such 2D models with unexcitable obstacles surrounded by a border region where the
ionic properties of cardiomyocytes are different form those in the rest of the tissue. We
vary the size of the obstacle and gray zone to reveal their effects on the pattern of wave
rotation and its period, which is an important characteristic of cardiac arrhythmia. We
show that depending on the parameters, the wave can rotate either around the compact
scar, or other regimes occur where the wave rotates within the gray zone or on its border
with the normal tissue. The switching between these regimes can be understood from
the principle of minimal period described in this paper. The dependency of the period
on the size of the scar in the physiologically relevant range is determined by the slope of
the conduction velocity restitution curve and can be estimated from a simple equation
derived in the paper. We also show that for small scars the regional tissue heterogeneity
can induce dynamical instability. Overall we provide classification of excitation patterns
and determine main factors which can affect the period of the wave rotation.

2. Materials and Methods

67   *2.1. Baseline ventricular myocardial tissue model. Numerical approach and software*

To describe propagation of the excitation wave in the ventricular myocardial tissue
from human heart, we used a 3-dimensional monodomain approach [19] and isotropic
medium:

$$C_m \frac{\partial V}{\partial t} = D\nabla^2 V - I_{ion}, \tag{1}$$

68   where V is the transmembrane potential; D is electro-diffusion matrix. Here, we used a
69   uniform D value throughout the tissue simulating isotropic media. C_m is the capacitance
70   of the cell membrane. I_{ion} is sum of all transmembrane ionic currents, described with
71   biophysically detailed cellular ionic TP06 model of human ventricular cardiomyocytes
72   [20]:

$$I_{ion} = I_{Na} + I_{K1} + I_{to} + I_{Kr} + I_{Ks} + I_{CaL} + I_{NaCa} + I_{NaK} + I_{pCa} + I_{pK} + I_{bNa} + I_{bCa}, \tag{2}$$

where I_{Na} is the Na^+ current, I_{K1} is the inward rectifier K^+ current, I_{to} is the
transient outward current, I_{Kr} is the delayed rectifier current, I_{Ks} is the slow delayed
rectifier current, I_{CaL} is the L-type Ca^{2+} current, I_{NaK} is the Na^+/Ca^{2+} exchanger current,
I_{NaK} is the Na^+/K^+ ATPhase current, I_{pCa} and I_{pK} are plateau Ca^{2+} and K^+ currents,
and I_{bNa} and I_{bCa} are background Na^+ and Ca^{2+} currents. This cellular model provides
a detailed description of voltage, ionic currents, and intracellular ion concentrations
and is based on a wide range of human electrophysiological data. The properties of all
these currents are fitted to their experimentally measured values and their dynamics is
fitted using additional differential equations. Each of the currents typically depends on

a maximal constant conductivity G_{max}, and on current values of time-dependent V and
certain gating variables.

In this study we used a thin single layer of excitable tissue simulating 2D myocardial
slab. Simulations were performed on grids $500 \times 500 \times 1$ elements ($140 \times 140 \times 0, 28$
mm^3) or $750 \times 750 \times 1$ elements ($210mm \times 210 \times 0, 28$ mm^3). Each tissue element can
represent either excitable myocardial tissue or inexcitable post-infaction scar. Larger
grids were used for modelling post-infarction scars with radius $r_{IS} > 16.8$ mm.

   Boundary conditions were formulated as the no flux through the boundaries:

$$\vec{n}\nabla V = 0 \tag{3}$$

where $\vec{n}$ is the normal to the boundary. Infarction scar elements were simulated as
non-conducting inexcitable obstacles and considered as internal boundaries (no flux) for
the myocardial elements.

2.2. Simulation of post-infarction scar and gray zone

Post-infarction scar (IS) was modelled as a non-conducting inexcitable circular
obstacle at the center of the tissue slab (black area in Fig. 1, left). It is surrounded by
a gray zone (gray area in Fig. 1, left). Here, we used a TP06 cellular ionic model [20]
with referent parameters for every cell in the healthy myocardial tissue, and a model
with modified parameters (Table 1) for every cell in the gray zone reflecting cellular
remodeling in myocardial tissue around the scar. Modification of TP06 ionic model for
description of cells in the gray zone was taken from [21] and is similar to the properties
of gray zone cells from [22]. In particular, the maximal conductances of several ionic
currents are changed against the reference values as specified in Table 1. The difference
in model parameters result in the difference in the action potential shape and duration
between the cells from gray zone and healthy myocardium (Fig. 1(right)). The action
potential duration is longer in the gray zone. The difference in the cellular activity
between the border zone and the rest of the healthy tissue provides the functional
heterogeneity in myocardial tissue around the scar.

Table 1: Maximal conductivity (G_*) of ionic currents in cellular models from the normal tissue (referent values) and gray zone (modified values).

Current description	Normal tissue	Gray zone
Fast Na+ current (I_{Na})	14.838	12.6123 ($0.85 \cdot G_{Na}$)
L-type Ca^{2+} current (I_{CaL})	$3.98 \cdot 10^{-5}$	$3.03674 \cdot 10^{-5}$ ($0.763 \cdot G_{CaL}$)
Rapid delayed rectifier current (I_{Kr})	0.134	0.0938 ($0.7 \cdot G_{Kr}$)
Slow delayed rectifier current (I_{Ks})	0.27	0.054 ($0.2 \cdot G_{Ks}$)
Inward Rectifier K^+ Current (I_{K1})	5.405	1.621 ($0.3 \cdot G_{K1}$)
Transient Outward Current (I_{to})	0.29372972	0.029372972 ($0.1 \cdot G_{to}$)

Figure 1. Left: Schematic representation of the geometry of cardiac tissue with the post-infarction scar: Normal tissue (pink), post-infarction scar (black) and gray zone around it (gray). Right: the shape of the action potential vs time in cells of healthy myocardium (pink) and gray zone (gray).

2.3. Protocol of numerical experiments in 2D tissue

A rotating wave in a 2D myocardial tissue slab with a post-infarction scar and gray zone was initiated counterclockwise using a standard protocol S1S2 (see Fig. S1 in the Supplement for graphical description of the protocol). Period of the stabilized wave was calculated in an arbitrary point in the healthy tissue as a time interval between the subsequent moments of the membrane depolarisation (zero V value at $V' > 0$).

We varied the size of scar from $r_{IS} = 0$ to $r_{IS} = 53.2$ mm with a step of 2.8 mm, and width of the gray zone from $w_{GZ} = 0$ to $w_{GZ} = 42$ mm also with a step of 2.8 mm and analysed the wave rotation pattern and evaluated the rotation period.

Note, in this study, we did not simulate the variability in the cellular population within the tissue. Here, we assumed two uniform areas of the myocardial tissue around the scar with the reference cellular model used throughout the healthy myocardium and the modified cellular model used throughout the gray zone. Thus, a single model run per each fixed pair (r_{IS}, w_{GZ}) of varied infarction parameters was performed.

Numerical methods

To solve the problem (1)-(3) we used a finite-difference method with 9-point stencil discretization scheme as described in [23] with 0.28 mm for the spatial step and 0.02 ms for the time step. Rush-Larsen formalism [24] was used for TP06 gating variables integration.

All calculations were performed on a C program on clusters "URAN" (IMM, Ural Branch of RAS) and "IIP" of Institute of immunology and Physiology (Ekaterinburg). The program uses CUDA for GPU parallelization and was compiled with a Nvidia C Compiler "nvcc". Computational nodes have graphical cards Tesla K40m0.

3. Results

3.1. Wave rotation regimes

Varying the radius of post-infarction scar (r_{IS}) and gray zone width (w_{GZ}) in our 2D tissue models, we observed several qualitatively different spatial patterns of the wave rotation. Representative types of the wave patterns are shown in Figure 2.

Figure 2. Examples of waves rotating around post-infarction scars of different sizes (up to down: $r_{IS} = 2.8mm; 19.6mm; 42mm$) with a gray zone of different sizes (left to right: $w_{GZ} = 0mm; 14mm; 42mm$). Four rotation regimes are shown: *scar rotation* (SR), *functional rotation* (FR, blue shading shows the effective obstacle around which the wave rotates), *gray zone rotation* (GZR), and *scar rotation 2* (SR2). See text for the description of the rotation regimes. Left column: typical view of the wave rotating around the scar in the absence of gray zone ($w_{GZ} = 0$ mm). Waves rotate in two regimes: *FR* (A1) and *SR* (B1, C1). Middle column: waves rotate around the gray zone in a *GZR* (A2, B2) and *SR2* (C2). Right column: ($w_{GZ} = 42$ mm) *SR2* regime in all cases (A3-C3). Arrows show the direction of the wave rotation. Dashed arrows in A2 ($r_{IS} = 19, 6mm$, $w_{GZ} = 14mm$) show wave direction inside the gray zone. Dashed circles show the border of the gray zones. Movies of rotation of spiral waves in *GZR* and *SR2* regimes can be found in the Supplement (movies S1 and S2).

In the absence of gray zone (Fig. 2 B1, C1, $w_{GZ} = 0$ mm) the rotation occurs with
the wavefront orthogonal to the boundary of the scar. Let us call this regime as a *scar*
rotation, as here the radius or rotation is determined by the radius of the scar. However,
for a small scar in Figure 2.A1 the situation is different. It looks as if the wave is rotating
around an effective obstacle of a larger radius: the point where displacement in the
normal direction coincides with the visible rotation around a center is located on a circle
of a larger radius than the scar (blue shaded circle). Let us call such regime as a *functional*
rotation. That means that the radius of rotation here is determined by the wave itself and
not by the obstacle.

For the gray zone of $w_{GZ} = 14$ mm (Fig.2 A2, B2) the rotation occurs with the
wavefront orthogonal to the boundary of the gray zone, as if it is non-excitable. Inside
the gray zone we see an area in red-orange colors indicating that spatial duration of
the action potential here in prolonged. Let us call this regime as a *gray zone rotation* as
here the radius or rotation is determined by the radius of the gray zone. For the largest
scar in Fig.2(C2) the wavefront is attached and almost orthogonal to the scar and we
observe scar rotation, however, as the wave here moves inside the gray zone let us call
this regime as a *scar rotation 2*.

For the largest gray zones of $w_{GZ} = 42$ mm (Fig.2 B3, C3) we see the *scar rotation 2*
regime, except that in Fig.2(A3), where we see some features of *functional rotation* inside
the gray zone.

Now let us analyse the period of rotation, which is one of the main characteristics
of cardiac arrhythmia.

3.2. Rotation period depending on geometry of the scar

158   We plotted dependencies of the rotation period on the radius of the scar for 4 values
159 of the gray zone width (Fig. 3) and dependencies of the rotation period on the width of
160 the gray zone for 4 values of the scar radius (Fig. 4).

Figure 3. Dependence of the wave rotation period on the radius of the scar with the gray zone
of different width. The black line shows cases in absence of gray zone. The gray dashed line
represents periods of waves rotating around a scar in tissue with properties of the gray zone
($w_{GZ} = \infty$). Filled dots (●) represent cases of the *scar rotation* regime. Open dots (○) represent cases
of the *gray zone rotation* regime. Marked dots (1)-(4) on this figure and Fig. 4 represent the same
cases.

Let us consider Figure 3 first. The black line shows the rotation period vs r_{IS} in the
absence of the gray zone ($w_{GZ} = 0$). We see that for small r_{IS} ($r_{IS} < 14$ mm) the period
is almost constant and does not depend on the size of the scar, and then dependence
becomes linear with a slope of 6.95. Similarly, in the absence of normal tissue, the dashed
line, ($w_{GZ} = \infty$), we have a dependency with the region of almost constant period
and then monotonically increasing dependency approaching linear with a slope of 5.45.
(Here entire tissue has the properties of the gray zone) We will use this line later to
explain the mechanisms of transitions between different wave rotation regimes. For the
gray zone of width $w_{GZ} = 8.4$ mm (the blue line) we have a similar dependency, with
a the linear slope=6.32, however the transition to the linear part here starts for smaller
infarct radius values of $r_{IS} > 2.8$ mm. For the gray zone with $w_{GZ} = 30.8$ and 36.4 mm
(green and yellow lines, respectively), both lines approach the asymptotic dashed line at

large scars. For smaller scar radius, the dependencies are linear before the green and
yellow lines approach the asymptotic dashed line. Their slopes here are 6.35 and 6.5,
respectively.

Figure 4. Dependence of the wave rotation period on the width of the gray zone (w_{GZ}) in experiments with different radius of infarction scars. Red line shows case with smaller radius of infarction scar ($r_{IS} = 8.4$ mm), purple line - case with bigger infarction scar ($r_{IS} = 36.4$ mm). Open symbols represent cases of the *gray zone rotation* regime. Filled symbols represent cases of *scar rotation* regime. Dots (1)-(4) on this figure and on Fig. 3 represent the same cases.

Now let us consider the dependence of the wave rotation period on the width of
the gray zone at constant r_{IS} (Fig. 4). For all r_{IS}, we have similar dependencies, which at
small w_{GZ} are almost linearly increasing functions, that for larger w_{GZ} instantly switch
to the almost horizontal straight lines. We also see that for larger scars the switch to
the plateau occurs at lower w_{GZ}. In particular, for a small scar of $r_{IS} = 8.4$ mm plateau
phase starts around $w_{GZ} = 36.0$ mm, while for the largest scar of $r_{IS} = 36.4$ mm, the
plateau is approached at around $w_{GZ} = 16.8$ mm.

3.3. Period and regimes of rotation

Now let us explain the observed dependencies and relate them to the rotation
regimes defined in section 3.1. Let us first consider the orange line in Fig. 4. Here the line
starts at $w_{GZ} = 0$ with the *scar rotation* regime and period of 276 ms. We illustrate this
wave pattern in Fig. 2 (B1). When w_{GZ} increases the wave detaches from the obstacle
and starts rotating around the gray zone in the *gray zone rotation* regime. We illustrate
this wave pattern with the period of rotation of 379 ms in Fig. 2 (B2). The reason for
onset of such regime is the following. It turns out that for wave it is faster to rotate
around the gray zone in the normal tissue, than around a scar in the gray zone tissue, as
the velocity of the wave there is smaller due to longer action potential in the gray zone.
Indeed, we found that the period of rotation inside the gray zone for scar of this size
is 460 ms. We can see it from the dashed line in Fig.3 for $r_{IS} = 19.6$ mm. This period
is longer than the period of 379 ms for rotation around the gray zone, and thus wave
chooses to rotate around the gray zone.

We can also explain the transition from the gray zone to scar rotation regime if we
further increase the size of the gray zone. Now the period of rotation around the gray
zone increases with its size, and for the gray zone of $w_{GZ} = 30.8$ mm it reaches a value
of 460 ms (see dot marked *SR2* in Fig. 4). For further increase in w_{GZ}, it will be faster for
the wave to rotate around the scar than around the gray zone, and we observe the *scar*
rotation2 regime with period of 460 ms (Fig. 2, B3). Thus further increase in w_{GZ} will

not change the wave rotation regime and the period will remain $T = 460$ ms, as we see
in Fig. 4.
Note, that as both the gray zone and scar rotations occur in the normal tissue, these
regimes are closely related to each other. For example, in our case of $r_{IS} = 19.6$ mm, and
$w_{GZ} = 14$ mm the radius of the gray zone is $r_{IS} + w_{GZ} = 33.6$ mm. From Fig. 3 we find
that for scar of $r_{IS} = 33.6$ mm the period of rotation is 362 ms, which is indeed close to
observed value of 379 ms.
Now let consider dependencies in Fig. 3. For $w_{GZ} = 0$ (the black line), the linear
part of the curve is associated with the *scar rotation* regime and increasing of period of the
wave is caused by increasing scar radius r_{IS}. However, the flat part of the dependency
for small r_{IS} is associated with the *functional rotation* regime. Indeed, for small r_{IS} the
rotation effectively occurs around a circle with the radius larger than the size of the
scar and thus size of the scar does not affect the rotation period An example of such
effective rotation circle is shown in Fig.2 A1 by shaded blue. Similar wave patterns of the
functional and scar rotation regimes are associated with the dashed curve ($w_{GZ} = \infty$).
However, here we obviously have the *scar rotation2* regime for large scars and the
functional rotation regimes for small scars as the wave here propagates in the tissue with
the properties of the gray zone.
For the gray zone with a large width of $w_{GZ} = 30.8$ mm (see green line in Fig. 3)
and large scars $r_{IS} >= 14$, the curve coincides with the dashed line corresponding to the
scar rotation2 regime. However for $r_{IS} < 14$ mm we have the *gray zone rotation* regime as
the rotation around gray zone is faster than *scar rotation 2*. For the gray zone with the
largest width of $w_{GZ} = 36.4$ (yellow line in Fig. 3) we have *scar rotation 2* except smaller
scars of $r_{IS} < 8.4$ mm. Here we have again the *gray zone rotation*. Compared to the green
curve the *scar rotation 2* here occurs at the smaller scar radius. This is also logical, as for
larger w_{GZ} the corresponding curve is closer to its asymptotic dashed curve.
Finally, let us consider the factors which determine the slopes of the dependencies
of period of rotation on the scar geometry parameters.

3.4. Factors affecting period of rotation

   To determine the factors responsible for the observed dependency of the period
on the size of the scar and gray zone let us consider the following simple equation. If
we assume that the velocity of wave (v) at the boundary of the obstacle is constant and
the perimeter of the obstacle is L, then the period T is given by: $T = L/v$, or $L = T * v$.
In turn, the velocity of the wave depends on the period of excitation T (conduction
velocity (CV) restitution) and the curvature of the wave front [25,26]. If we neglect the
dependency of the velocity on the curvature we get: $L = T * v(T)$. Thus the slope of the
dependency of the period on the obstacle size (shown in Fig.3) can be found from the
following simple calculation:

$$\frac{dL}{dT} = \frac{d(T * v(T))}{dT} = v(T) + T * \frac{dv}{dT}$$

$$\frac{dT}{dL} = \frac{1}{v(T) + T * dv/dT} \tag{4}$$

232 where dv/dT is the slope of the CV restitution curve.
233   Thus we see that the slope of the dependency of wave period T on the obstacle
234 perimeter L is determined not only by the velocity v itself but also by the slope of the CV
235 restitution curve.

Figure 5. Dependence of the wave speed on the activation period of the wave (restitution curve). Wave speed in normal tissue (black) and tissue with properties of the gray zone (gray) are shown. Black and gray dashed lines show maximum speed of waves in normal tissue ($v_{NT} = 0.71$ m/s) and in gray zone tissue ($v_{GZ} = 0.66$ m/s) respectively.

We computed CV restitution curves for the normal tissue and tissue with properties of the gray zone (Fig. 5). We see that both curves are monotonically increasing functions $v(T)$ approaching some saturated values (v_{NT} for healthy tissue, and v_{GZ} for gray zone tissue). Thus for large pacing periods, dv/dT approaches zero value, and according to the formulae (4) the slope dT/dL would approach the value $1/v_{NT}$ or $1/v_{GZ}$, respectively. However, such saturation occurs for rather long pacing periods, which are mostly outside of the physiologically relevant values. In our case, the rotation periods are in the range of $250 < T < 600$, where v strongly depends on T for the both CV restitution curves. Let us estimate, which terms in Eq. (4) primarily determine the slope of the dependencies in Fig.3, and explain why the curves with larger gray zone have smaller slopes than the curve without the gray zone. Let us use period $T = 500$, which is inside the region of linear dependency in Fig.3 for both curves: with and without the gray zone. For the case of a scar without the gray zone (the black curve), at $T = 500$ ms we find that $v = 0.678$, and $dv/dT = 0.0003$ derived from the black curve in Fig.5. We use it as in this case the wave rotates around the scar surrounded by the normal tissue. Thus in Eq. (4) the second term $T * dv/dT = 0.15$ is substantially smaller than the first term v. For a large gray zone (green curve in Fig.3) the rotation at $T = 500$ ms occurs inside the gray zone, thus we need to use the CV restitution curve for the border zone tissue (the gray curve in Fig.5). Similarly we find that for $T = 500$ ms $v = 0.46$, and $dv/dT = 0.00195$, thus in Eq.(4) $T * dv/dT = 0.979$ is higher than v. Thus in this case we see that now the second term $T * dv/dT$ determines the slope of the dependency. Thus we can conclude that the lower slope of the dependency of T on L in the presence of the gray zone is a result of CV restitution slope (dv/dT), where larger derivative occurs due to 'shift' of CV restitution curve to the longer periods because of tissue remodelling in the gray zone (Fig.5). Note however, that linear dependencies which we see in Fig.3 for large radii actually just a coincidence for specific values of parameters. The slopes will further change and for very large values of r they approach the values of $1/v_{NT}$ and $1/v_{GZ}$ correspondingly. At that stage the slope of the dashed line in Fig.3 will be larger than that of the black line. However, it occurs far beyond the physiologically relevant values for spiral waves.

265 *3.5. Additional rotation regimes*

Figure 6. Examples of waves with dynamical instabilities. Border of the gray zone is depicted by the dashed line. Arrows show the direction of the wave rotation. Dashed arrows show wave direction inside the gray zone. A: a passive regime. The breakthrough waves marked by dashed arrows do not exit the gray zone ($r_{IS} = 11.2$ mm, $w_{GZ} = 25.2$ mm). B: an active regime. The breakthrough waves exit the gray zone and force the wave rotate at some distance from the border of the gray zone ($r_{IS} = 2.8$ mm, $w_{GZ} = 5.6$ mm.) An illustration of such dynamics can be found in the Supplement movies (movies S3 and S4).

In addition to the regimes discussed above, for few parameter values we also
observed more complex rotation regimes. All these regimes occur for small scars and
can be distinguished into two types: active and passive. Fig.6 A shows an example of
passive regime, where we periodically see breakthrough of excitation into the gray zone,
however, they do not disturb the leading edge wave rotation. Fig.6 B shows an example
of active regime. Here, the wave comes outside the gray zone, actively interacts with the
leading edge of excitation and changes the trajectory of the wave rotation. As a result we
observe a non-monotonical dependency of the period on the size of the scar. However,
the active regime was not frequently observed, we saw it only in 3 cases (out of 360) with
perimeter of gray zone 52.7 mm ($r_{IS} = 0$ mm, $w_{GZ} = 8.4$ mm; $r_{IS} = 2.8$ mm, $w_{GZ} = 5.6$
mm; $r_{IS} = 5.6$ mm, $w_{GZ} = 2.8$ mm), thus they may have limited importance for overall
dynamics.

4. Discussion

In this paper we perform a detailed study of wave dynamics rotating around a
region, which generically represents a post-infarction scar and includes a compact inex-
citable obstacle surrounded by a gray zone with modified cellular activity as compared
with the rest of normal tissue. We found that all observed dynamics can be subdivided
into several main classes. These classes naturally occur if we combine properties of
wave rotating around inexcitable obstacle and effects of tissue heterogeneity on the wave
propagation.

In the absence of tissue heterogeneity, the period of wave rotating around an
obstacle is mainly determined by the velocity of the wave and the perimeter of the
obstacle [10]. This is a *scar rotation* regime, which was observed in previous studies.
However, compared to earlier studies [10] we show that one needs to take into account
the dependency of the velocity on the period (CV restitution), which substantially affects
the period of wave rotation. Another observed regime of *functional rotation*, was found in
early studies in the reaction diffusion models [11]. It was shown that for small obstacles
the period of rotation does not decrease anymore, wave detouches from the obstacle and
rotates around a circle which exceeds the obstacle size. In [11] it was also shown that
detachment from the obstacle is associated with a large increase in the period of rotation.
We did not observe such period increase in our study. This probably occurs because in
[11] a two variable model with lower excitability was used. In our study we used an
ionic model with high excitability, and thus boundary effects of the obstacle on the wave
propagation velocity in our case are less pronounced.

In the presence of the gray zone, the situation is more complex and, as far as we
aware, was not yet analyzed in details. Our study shows that here we can observe
additional rotation regimes, which we call *gray zone rotation* and *scar rotation 2* regimes.
We show that dependencies of period on the size of the scar and gray zone are bounded
by two asymptotic solutions for homogeneous tissue: rotation of wave around a scar in
normal tissue and rotation of wave around a scar in tissue with properties of the gray
zone. As we show in Fig. 3 all curves for different sizes of the gray zone are located
between these two asymptotic lines and the slopes of these lines bound from above and
below the slopes of all other curves. In addition, we found a new regime, *gray zone*
rotation. In that case, the wave rotates as if the compact scar coincides with the gray zone.
In the *gray zone rotation* regime the period of rotation depends on the perimeter of the
gray zone, and not on the size of the scar. We illustrate it in Fig. 7, where we show the
dependence of the period on the perimeter of the gray zone for various sizes of the scar.
We see that all curves in certain range of parameters almost coincide, which one more
time confirms that here the perimeter of the gray zone, and not the scar size determines
the period of rotation. However, we also see that this curves with the gray zone are little
above the dashed line, representing the case of pure scar rotation. It indicates that the
leading edge of rotation for the *gray zone rotation* regime is a little outsize the boundary
of the gray zone, which maybe a result of the electrotonic effects [27].

Figure 7. Dependence of the wave rotation period on the perimeter of the gray zone for different radii of infarction scar. Red line shows case with smallest radius of scar core, purple line - case with biggest scar core. The dashed line connects points where the gray zone is absent ($w_{GZ} = 0$ mm).

Important rule for transition between the gray zone and scar rotation regimes is the minimal period principle. In fact, if rotation in normal tissue along the gray zone is faster than the rotation around the scar in the gray zone, then the *gray zone* regime will occur. Such transition is clearly seen in Fig.3 and Fig. 4 where we see change from one type of dependency to another one. Note, this also implies that transitions between the regimes strongly depend on the relation of the velocity of the wave inside and outside the gray zone. Here we considered only one type of cellular remodeling based on the data reported in [21,22], where action potentials are longer in the gray zone. However in [28] it was shows that remodelling in the gray zone can also result in different changes in the properties of cardiac tissue and opposite effects on action potential duration. Thus it would be interesting to study similar regimes of wave rotation for different properties of cardiac tissue in the gray zone. In addition, the gray zone can also include fibrosis and reduced number of gap junctions. It also can decrease the propagation velocity and thus make *gray zone rotation* regime more essential.

The dynamics of rotating waves around an obstacle was studied in [14], where a general theory for the dynamics and instability of pinned spirals was developed. In particular, it was shown that the destabilization is better understood by the implementation of a mapping rule and dimension reduction.

Also transition between scar and functional rotation when the radius of the obstacle decreases were studied in [29]. There a sequence of transitions, from periodic motion to a modulated period-2 rhythm, and then to spiral wave breakup [30,31] were observed. It would be interesting to study how such dynamical instabilities will manifest itself in the presence of tissue heterogeneity around a scar, as considered in our paper.

It would be also interesting to study such rotational activity in anatomical models of the heart [22,32] and also include additional types of regional cellular heterogenety in the normal tissue as considered in [33,34]. Note that cellular heterogeneity, especially involving pathological elongation of the action potential, can lead to early afterdepolarizations and new types of spatio-temporal dynamics [35] .

In conclusion, we performed the first detailed studies of regimes of wave rotating around an obstacle surrounded by heterogeneous ventricular myocardial tissue, which is a generic model of the infarction scar with the gray zone. We found that the main regimes here include rotation around the compact scar and also rotation around the gray zone. We explained dependency of the rotation period on the size of the scar and showed

that it is strongly affected by the conduction velocity restitution. We also show that the transition between the regimes can be understood and predicted from the minimal period principle.

Supplementary Materials: The following are available at https://www.mdpi.com/2227-739 0/1/1/0/s1, Video S1: Example of wave rotating in *gray zone rotation* regime. Infarction scar size:$r_{IS} = 19.6$ mm. Gray zone width $w_{GZ} = 14$ mm. Video S2: Example of wave rotating in *scar rotation 2* regime. Infarction scar size:$r_{IS} = 19.6$ mm. Gray zone width $w_{GZ} = 42$ mm. Video S3: Example of the wave with passive dynamic instability in gray zone. Infarction scar size:$r_{IS} = 11.2$ mm. Gray zone width $w_{GZ} = 22.4$ mm. Video S4: Example of the wave with active dynamic instability in gray zone. Infarction scar size:$r_{IS} = 2.8$ mm. Gray zone width $w_{GZ} = 5.6$ mm.

Author Contributions: Conceptualization, A.P., D.M. and O.S; formal analysis, P.K., D.M.;Investigation, P.K.; methodology, A.P. and P.K.; software P.K., A.D. and D.M.; supervision, A.P. and O.S.; visualization, P.K. and A.D.; writing–original draft preparation, P.K., A.P.and O.S.; writing–review and editing, P.K., A.P. and O.S.; All authors have read and agreed to the published version of the manuscript.

Funding: A.P., P.K., D.M., A.D. and O.S. was funded by the Russian Foundation for Basic Research (#18-29-13008). A.P. and O.S. was funded by RF Government Act #211 of March 16, 2013 (agreement 02.A03.21.0006). P.K., D.M., A.D., and O.S. work was carried out within the framework of the IIF UrB RAS theme No AAAA-A21-121012090093-0. A.P. was partially funded by BOF Ghent University.

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

Abbreviations

The following abbreviations are used in this manuscript:

CV	Conduction velocity
FR	Functional rotation
GZ	Gray zone
GZR	Gray zone rotation
IS	Post-infarction scar
NT	Normal tissue
SR	Scar zone rotation
SR2	Scar rotation 2

References

1. Cross, M.C.; Hohenberg, P.C. Pattern formation outside of equilibrium. *Reviews of modern physics* **1993**, *65*, 851.
2. Winfree, A.T.; Strogatz, S.H. Organizing centres for three-dimensional chemical waves. *Nature* **1984**, *311*, 611–615.
3. Pertsov, A.M.; Aliev, R.R.; Krinsky, V.I. Three-dimensional twisted vortices in an excitable chemical medium. *Nature* **1990**, *345*, 419–421.
4. Gorelova, N.A.; Bureš, J. Spiral waves of spreading depression in the isolated chicken retina. *Journal of neurobiology* **1983**, *14*, 353–363.
5. Siegert, F.; Weijer, C.J. Three-dimensional scroll waves organize Dictyostelium slugs. *Proceedings of the National Academy of Sciences* **1992**, *89*, 6433–6437.
6. Jalife, J. Ventricular fibrillation: mechanisms of initiation and maintenance. *Annual review of physiology* **2000**, *62*, 25–50.
7. Karma, A. Physics of cardiac arrhythmogenesis. *Annu. Rev. Condens. Matter Phys.* **2013**, *4*, 313–337.
8. Nishida, K.; Qi, X.Y.; Wakili, R.; Comtois, P.; Chartier, D.; Harada, M.; Iwasaki, Y.k.; Romeo, P.; Maguy, A.; Dobrev, D. Mechanisms of atrial tachyarrhythmias associated with coronary artery occlusion in a chronic canine model. *Circulation* **2011**, *123*, 137–146.
9. Numata, A.; Miyauchi, Y.; Ono, N.; Fishbein, M.C.; Mandel, W.J.; SHIEN-FONG, L.I.N.; Weiss, J.N.; PENG-SHENG, C.H.E.N.; Karagueuzian, H.S. Spontaneous atrial fibrillation initiated by tyramine in canine atria with increased sympathetic nerve sprouting. *Journal of cardiovascular electrophysiology* **2012**, *23*, 415–422.
10. Weiner, N.; Rosenblunth, A. The mathematical formulation of the problem of conduction of impulses in a network of connected excitable elements specifically in cardiac muscle. *Arch. Inst. Cardiol. Mex.* **1946**, *16*, 205–265.
11. Pertsov, A.M.; Ermakova, E.A.; Panfilov, A.V. Rotating spiral waves in a modified Fitz-Hugh-Nagumo model. *Physica D: Nonlinear Phenomena* **1984**, *14*, 117–124.

12. Davidenko, J.M.; Pertsov, A.V.; Salomonsz, R.; Baxter, W.; Jalife, J. Stationary and drifting spiral waves of excitation in isolated cardiac muscle. *Nature* **1992**, *355*, 349–351.
13. Vinson, M.; Pertsov, A.; Jalife, J. Anchoring of vortex filaments in 3D excitable media. *Physica D: Nonlinear Phenomena* **1994**, *72*, 119–134.
14. Gao, X.; Feng, X.; Li, T.C.; Qu, S.; Wang, X.; Zhang, H. Dynamics of spiral waves rotating around an obstacle and the existence of a minimal obstacle. *Physical Review E* **2017**, *95*, 052218.
15. Balaban, G.; Halliday, B.P.; Bai, W.; Porter, B.; Malvuccio, C.; Lamata, P.; Rinaldi, C.A.; Plank, G.; Rueckert, D.; Prasad, S.K. Scar shape analysis and simulated electrical instabilities in a non-ischemic dilated cardiomyopathy patient cohort. *PLoS computational biology* **2019**, *15*, e1007421.
16. Baba, S.; Dun, W.; Cabo, C.; Boyden, P.A. Remodeling in cells from different regions of the reentrant circuit during ventricular tachycardia. *Circulation* **2005**, *112*, 2386–2396.
17. Schmidt, A.; Azevedo, C.F.; Cheng, A.; Gupta, S.N.; Bluemke, D.A.; Foo, T.K.; Gerstenblith, G.; Weiss, R.G.; Marbán, E.; Tomaselli, G.F. Infarct tissue heterogeneity by magnetic resonance imaging identifies enhanced cardiac arrhythmia susceptibility in patients with left ventricular dysfunction. *Circulation* **2007**, *115*, 2006–2014.
18. Bernus, O.; Zemlin, C.W.; Zaritsky, R.M.; Mironov, S.F.; Pertsov, A.M. Alternating conduction in the ischaemic border zone as precursor of reentrant arrhythmias: a simulation study. *EP Europace* **2005**, *7*, S93–S104.
19. Panfilov, A.; Holden, A. Computer simulation of re-entry sources in myocardium in two and three dimensions. *Journal of Theoretical Biology* **1993**, *161*, 271–285.
20. ten Tusscher, K.; Panfilov, A.V. Alternans and spiral breakup in a human ventricular tissue model. *American Journal of Physiology-Heart and Circulatory Physiology* **2006**, *291*, H1088–H1100. doi:https://doi.org/10.1152/ajpheart.00109.2006.
21. Decker, K.F.; Rudy, Y. Ionic mechanisms of electrophysiological heterogeneity and conduction block in the infarct border zone. *American Journal of Physiology-Heart and Circulatory Physiology* **2010**, *299*, H1588–H1597.
22. Arevalo, H.J.; Vadakkumpadan, F.; Guallar, E.; Jebb, A.; Malamas, P.; Wu, K.C.; Trayanova, N.A. Arrhythmia risk stratification of patients after myocardial infarction using personalized heart models. *Nature communications* **2016**, *7*, 1–8.
23. Ten Tusscher, K.H.W.J.; Panfilov, A.V. Modelling of the ventricular conduction system. *Progress in biophysics and molecular biology* **2008**, *96*, 152–170.
24. Rush, S.; Larsen, H. A practical algorithm for solving dynamic membrane equations. *IEEE Transactions on Biomedical Engineering* **1978**, *4*, 389–392.
25. Zykov, V.S. Analytic evaluation of the relationship between the speed of a wave of excitation in a two-dimensional excitable medium and the curvature of its front. *Biofizika* **1980**, *25*, 888–892.
26. Cabo, C.; Pertsov, A.M.; Baxter, W.T.; Davidenko, J.M.; Gray, R.A.; Jalife, J. Wave-front curvature as a cause of slow conduction and block in isolated cardiac muscle. *Circulation research* **1994**, *75*, 1014–1028.
27. Defauw, A.; Kazbanov, I.V.; Dierckx, H.; Dawyndt, P.; Panfilov, A.V. Action potential duration heterogeneity of cardiac tissue can be evaluated from cell properties using Gaussian Green's function approach. *PLoS one* **2013**, *8*, e79607.
28. Mendonca Costa, C.; Plank, G.; Rinaldi, C.A.; Niederer, S.A.; Bishop, M.J. Modeling the electrophysiological properties of the infarct border zone. *Frontiers in physiology* **2018**, *9*, 356.
29. Xie, F.; Qu, Z.; Garfinkel, A. Dynamics of reentry around a circular obstacle in cardiac tissue. *Physical Review E* **1998**, *58*, 6355.
30. Panfilov, A.V.; Holden, A.V. Self-generation of turbulent vortices in a two-dimensional model of cardiac tissue. *Physics Letters A* **1990**, *151*, 23–26.
31. Panfilov, A.V.; Holden, A.V. Spatiotemporal irregularity in a two-dimensional model of cardiac tissue. *International Journal of Bifurcation and Chaos* **1991**, *1*, 219–225.
32. Panfilov, A.; Keener, J.P. Re-entry in an anatomical model of the heart. *Chaos, Solitons & Fractals* **1995**, *5*, 681–689.
33. Keldermann, R.H.; ten Tusscher, K.H.; Nash, M.P.; Hren, R.; Taggart, P.; Panfilov, A.V. Effect of heterogeneous APD restitution on VF organization in a model of the human ventricles. *American Journal of Physiology-Heart and Circulatory Physiology* **2008**, *294*, H764–H774.
34. Keldermann, R.H.; Ten Tusscher, K.H.W.J.; Nash, M.P.; Bradley, C.P.; Hren, R.; Taggart, P.; Panfilov, A.V. A computational study of mother rotor VF in the human ventricles. *American Journal of Physiology-Heart and Circulatory Physiology* **2009**, *296*, H370–H379.
35. Vandersickel, N.; Kazbanov, I.V.; Nuitermans, A.; Weise, L.D.; Pandit, R.; Panfilov, A.V. A study of early afterdepolarizations in a model for human ventricular tissue. *PLoS one* **2014**, *9*, e84595.