Bursts of recent transposable element activity and signatures of

element removal in the monarch genome, *Danaus plexippus*

Tobias Baril[1,2]* and Alexander Hayward[1,3]*

[1] Centre for Ecology and Conservation, University of Exeter, Penryn Campus,

Cornwall, TR10 9FE, UK

[2] ORCID: 0000-0002-5936-7531

[3] ORCID: 0000-0001-7413-718X

* Corresponding Author information:

Tobias Baril: tb529@exeter.ac.uk

Dr Alexander Hayward: alex.hayward@exeter.ac.uk

Abstract

Background

Lepidoptera (butterflies and moths) are an important model system in ecology and evolution. A high-quality chromosomal genome assembly of the monarch butterfly (*Danaus plexippus*), famous for its North American migration, is available but lacks an in-depth transposable element (TE) annotation. This provides an opportunity to explore host-TE interactions, and the impact TEs have in shaping the monarch genome.

Results

6.47% of the monarch genome is comprised of TEs, a reduction of 6.59% compared to the original TE annotation performed on the draft genome assembly. TE content is low compared to two closely related species, *Danaus chrysippus* (26.70%) and *Danaus melanippus* (11.87%). The biggest contributors to genome size in the monarch are LINEs and *Penelope*-like elements, and 37.7% of TE content is contributed by five newly identified TE families (two LINE, two *Penelope*-like, and one SINE). Some young DNA TE families show similar activity profiles to these LINEs, with their success putatively due to horizontal transposon transfer from species sharing the same environment. There are several recent peaks of TE activity in the monarch, with little evidence for peaks of activity more anciently. LINE fragments demonstrate signatures of genomic deletions as reported by studies on *Heliconius* butterflies, indicating a high rate of TE turnover. Given previous associations in other species, we investigated the association of TEs with wing colouration and immune genes. We find a single unclassified element 7kb upstream of the myosin gene locus, associated with wing colouration, and 49 immune genes with TEs within 5kb upstream of the transcription start site, presenting the potential for the involvement of TEs in regulatory functions.

Conclusions

We provide an in-depth TE annotation and analysis of TE diversity and evolution for the

monarch genome. We identify highly successful novel DNA TE families, mirroring the activity

profile of the most successful LINEs. We also find evidence of ongoing TE expansion and

removal in the monarch, highlighting the dynamic nature of repeat content in genomes over

time. Further in-depth comparative studies across closely related species will be beneficial to

our understanding of the evolutionary dynamics of TEs and the processes leading to their

contrasting distributions.

55 Keywords

56 Transposon; TE Annotation; Lepidoptera; *Danaus plexippus*; Butterfly; Genomic Deletion;

57 TE-Host Dynamics; Genome Evolution

59 Background

Transposable elements (TEs) are autonomous DNA sequences that can move within the

genome. TEs are present in nearly all eukaryotes, and are important in shaping their

genomes [1–8]. TEs also represent a large reservoir of sequences with the potential to

contribute to host genomic novelty, and they have been implicated in the evolution of

regulatory networks [3,9–13], chromosomal rearrangements [3,14–19], exon shuffling [20–

22], and donation of coding sequence [20,22–27].

Due to the great diversity and dynamic nature of TEs, species-specific TE libraries

and accompanying TE annotations are required to provide an understanding of their

evolution and impact. By examining differences in the diversity and abundance of TEs within

a genome, as well as their integration histories, proliferation frequencies, and rate of

removal, we can begin to unravel the factors that govern host-TE evolutionary dynamics.

Curating TEs in a wide range of species is also an important step more generally, to expand

understanding of the interplay between TEs and host genomes, and consequences for host

evolution [28]. Meanwhile, from a practical perspective, accurate and reliable repeat

annotation is essential to provide quality genome annotations, ensuring that repeat

sequences are not incorrectly annotated as host genes and vice versa, especially given the

vast numbers of genomes currently being sequenced across the tree of life.

Lepidoptera (butterflies and moths) are an important model system in ecology and

evolution. Lepidopterans have captivated researchers for decades, proving a fascinating and

practical system for research into topics such as development, physiology, host-plant

interactions, coevolution, phylogenetics, mimicry, and speciation (e.g [29–33]). Over recent

decades, Lepidoptera has become an important model system in genomics especially. For

example, Lepidoptera genomes have been pivotal in investigations into the genomic basis of

migration [34–36], warning colouration [35], adaptive radiations [37], and evolutionary

studies into TEs, including TE horizontal transfer [38–40], the influence of TEs on genome

size [41], and the role of a TE in the evolution of industrial melanism in the peppered moth

[42].

The monarch butterfly, *Danaus plexippus*, is famous for its incredible North American

migration, global dispersal, and orange warning colouration [35]. The monarch has been the

focus of much research and broad public interest for decades, with a draft genome released

in 2011, and work during the last decade revealing the genetic bases of several key traits

[35,36]. Most recently, a high-quality chromosomal assembly of the *D. plexippus* genome

has been generated [43], providing a powerful opportunity for a detailed analysis of TEs in

the monarch genome. Despite the monarch being an important research focus, we lack a

detailed TE annotation for the species, and there is little understanding of host-TE

interactions, or the impact that TEs have had on shaping the monarch genome more

generally.

Here, we provide a detailed annotation of TEs within the monarch genome, including

analyses of TE diversity and abundance, interactions with host genes, genomic distribution

including hot and coldspots, and TE evolution. We also compare the efficacy of a manual

and automated *de novo* TE library curation method for repeat annotation. We find evidence

of considerable recent TE activity in the monarch, with two novel Tc1-Mariner element

families proliferating at a high rate over recent timescales (*danPle_TcM-1* and *danPle_TcM-*

2). In contrast, we find a pattern of high proliferation rates of LINE and *Penelope* elements

was historically dominant, and these elements still characterise the monarch TE landscape.

We also reveal the role of large genomic deletions in protecting the host genome from costly

LINE TEs, which show signatures of removal. Overall, we provide an overview of the

broadscale dynamics that shape the TE landscape of the monarch genome, and provide a

high-quality, curated, and species-specific TE annotation library for use by the scientific

community.

Results

Performance of automated and manual curation

113   We compared a basic manual annotation method with an iterative automated

114   curation method to assess their relative performance. Repetitive element quantifications

115   using both the manual and automatically curated libraries are presented in Table 1.

Repeat Class	Automated *de novo* Library (+ RepBase and Dfam)			Manual *de novo* Library (+ RepBase and Dfam)			Delta (Automated - Manual)		
	No. elements	Total Length (Mb)	Percentage sequence (%)	No. elements	Total Length (Mb)	Percentage sequence (%)	No. elements	Total Length (Mb)	Percentage sequence (%)
Retroelement	31716	9.90	3.98%	21673	7.93	3.19%	10043	1.97	0.79%
SINE	3977	0.90	0.36%	3338	0.85	0.34%	639	0.05	0.02%
LINE	12165	5.20	2.09%	10011	4.77	1.92	2154	0.43	0.17%

Penelope	14532	3.14	1.26%	7770	1.86	0.75%	6762	1.27	0.51%
LTR element	1042	0.66	0.27%	554	0.45	0.18%	488	0.22	0.09%
DNA transposon	4497	2.37	0.95%	3443	2.11	0.85%	1054	0.25	0.10%
Rolling-circle	5454	1.79	0.72%	2928	0.90	0.36%	2526	0.88	0.36%
Unclassified	6658	1.86	0.75%	11564	3.34	1.34%	-4906	-1.47	-0.59%
Other	555	0.17	0.07%	359	0.15	0.06%	196	0.02	0.01%
Total repeats	48880	16.08	6.47%	39967	14.43	5.80%	8913	1.65	0.66%

Table 1. Repeat content in the monarch using the manual and automated *de novo* library

curation methods. "+ RepBase and Dfam" indicates annotations performed with the *de novo*

library combined with the *Arthropoda* RepBase library and Dfam.

The automated *de novo* curated library contained 115 consensus sequences (59%

classified), whilst the manual *de novo* curated library contained 75 consensus sequences

(35% classified) (Datasets S1 and S2; Additional Files 1 and 2). A similar number of

identified TE families remained unclassified during the RepeatMasker step using both

methods, with 47 (41%) and 49 (65%) unclassified putative TE families being curated under

the automated and manual methods, respectively.

In general, consensus sequences in the manually-curated library were shorter than

their counterparts in the automated library (mean TE consensus length in automated library

= 3,534bp; mean TE consensus length in manual library = 1,236bp). For sequences initially

identified as SINEs by RepeatModeler, an automated version of the BLAST, Extract, Extend

process [28] resulted in reclassification of these as LINEs (Table S1; Additional File 3), whilst

they remained classified as SINEs following manual curation. These sequences were further

interrogated, and it was revealed that these novel families belonged to the HaSE2 SINE

family, which were misclassified due to their significant homology with RTE LINEs [44]. The

misclassification of these elements by RepeatMasker highlights one of the main issues with

homology-based TE detection methods, which often miss, or misclassify, SINEs due to high

sequence variation and lack of terminal repeats [45]. We find that further issues remain with

all-by-all BLAST methods, as used by RepeatMasker and RepeatModeler, for the

identification of putative TE families. When comparing the manual and automated *de novo*

libraries, certain consensus sequences shared extremely high levels of similarity. For

example, in one case, the "*rnd-1_family-48#LINE/Penelope*" family from the automated

library matched both the "*rnd-1_family-39#Unknown"* and the "*rnd-3_family-30#Unknown*"

families in the manual library with the same percentage identity and match length (Table S2;

Additional File 3). The key difference between these two matches was that one matched in

the forward direction and the other in the reverse. This situation was uncovered for two

cases where a single sequence from the automated library matched two sequences in the

manually curated library (Table S2; Additional File 3). To resolve this issue, cd-hit-est [46,47]

was used to cluster matching sequences from opposite strands (For discussion, see

Additional File 18).

When using the different *de novo* libraries, there was no significant difference in TE

coverage (One-Way ANOVA, F = 0.067, p = 0.799). (Figure S1; Additional File 4). In total,

1.65Mb more sequence was annotated as TE sequence using the automated library than the

manual library (Table 1). 807.40kb (5.02% of total TE content) of TE loci annotated with the

automated library lacked annotations using the manual library (Tables S3 and S4; Additional

File 3), with the largest contribution from rolling circle elements (Tables S3 and S4;

Additional File 3). A total of 176.16kb (1.22% of total TE content) of TE loci annotated with

the manual library lacked annotations using the automated library, with the largest

contribution from unclassified elements (Table S3 and S4; Additional File 3). Whilst the

difference in TE annotation coverage was not significant, the increase in TE coverage when

using the automated library is of a scale that could be consequential when assessing the

impact of TE insertions on host genomes. Given the similarity in annotation results between

the two libraries, and the generally longer consensus sequences achieved using the

automated *de novo* library (Figure S2; Additional File 5), the annotation results from the

automated *de novo* curation pipeline were used for all downstream analyses in this study.

Transposable Element Landscape

Repeats are a major determinant of genome size [48–53]. However, in the monarch,

we find that repeats comprise just 6.47% of total genome size (Assembly Scaffold N50 =

9,210kb, 97.5% Complete Busco orthologs) (Figure 1, Table 1). This differs considerably in

comparison to the TE annotation performed on the original monarch genome assembly,

where repeats were estimated to cover 13.06% of the genome (Table S5; Additional File 3)

[36]. When examining the source of this variation, the major difference in coverage can be

attributed to unclassified elements, where we find that coverage of unclassified elements

reduced by 16.47Mb (Table S5; Additional File 3). This is likely due to improvements in TE

databases and annotation tools over the last decade, particularly given that the original

annotation used similar tools and databases to those applied in this study [36], in addition to

improved repeat detection due to increased genome assembly quality. Consistent with this,

a standard 'no frills' RepeatMasker run using the *Arthropoda* database and Dfam identified a

repeat content of 5.60% (Table S6; Additional File 3). Thus, the sequences annotated as

unclassified in the original annotation were most likely not TE sequences and are no longer

annotated as such by current annotation tools. Nevertheless, we are unable to confirm this

for certain, as the original *de novo* repeat library generated for the draft genome is

unfortunately unavailable.

We annotated the two additional genomes available for species in the *Danaus* genus

using the same automated TE annotation method that we applied to the monarch. In

comparison, the monarch exhibits a lower TE content, with repetitive elements covering

26.70% of the *D. chrysippus* genome (Scaffold N50 = 1,465kb, 93.5% Complete Busco

orthologs), and 11.87% of the *D. melanippus* genome (Scaffold N50 = 890kb, 86.0%

Complete Busco orthologs) (Figure S3, Table S5; Additional Files 6 and 3). We suggest that

the reduced TE load observed in the monarch may have arisen through one of two

mechanisms: (i) loss of repeat content in the monarch, or, (ii) gains in repeat content in both

D. chrysippus and *D. melanippus* (or in their ancestral lineages). Relatively recent TE activity

is observed in all three *Danaus* species, indicated by repeats exhibiting low genetic distance

from consensus (Figure 2 and Figure S3; Additional File 6). The genome sizes of both *D.*

chrysippus and *D. melanippus* are also very similar, at 322Mb and 354Mb, whilst the

monarch is only 248Mb. Increases in genome coverage are observed for all TE

classifications for both *D. chrysippus* and *D. melanippus* in comparison to the monarch,

except for a decrease in *Penelope* coverage in *D. melanippus* (Table S5; Additional File 3).

Therefore, we suggest that alternative (ii) is more likely, whereby there have been

independent increases in TE content which have led to increased genome sizes for both *D.*

chrysippus and *D. melanippus.* The main TE contributors to increases in genome size are

LINEs in both *D. chrysippus* (7.7% of genome, 24.7Mb) and *D. melanippus* (4.6% of

genome, 16.2Mb) (Table S5; Additional File 3).

TEs are unevenly distributed across the monarch genome and between different

chromosomes, indicated by a high standard deviation in TE base pairs per 100kb (mean =

6284bp/100kb, SD = 4183bp/100kb) (Figure 1, Figure 3, Figure 4, Table S14; Additional File

3). A larger standard deviation indicates a less even spread of TEs in a genome, where a

standard deviation of 0 indicates TEs are evenly distributed across the genome. The largest

density of TEs is found on chromosome 28 (10,128bp/100kb), whilst chromosome 9 exhibits

the lowest TE load (4604bp/100kb) (Table S14; Additional File 3). Across all regions, the

biggest contributors to genome size are LINEs and *Penelope* elements (Figure 1).

TEs are known to be unevenly distributed between different genomic compartments,

with TE densities varying between genic and intergenic space [1]. LINEs make up 2.38% of

the genome, which comprises: 2.80% of gene flanking regions, 1.50% of intronic sequence,

0.93% of exonic sequence, and 3.52% of intergenic sequence. Meanwhile *Penelope*

elements make up 1.18% of the genome as a whole: 1.43% of gene flanking regions ,1.13%

of intronic sequence, 0.17% of exonic sequence, and 1.88% of intergenic sequence.

Collectively, the greatest proportion of TEs is found in intergenic regions (total 3.86%:

intergenic space 1.16%, and gene flanking regions 2.70%), in comparison to gene bodies

(total 2.49%: introns 2.29%, and exons 0.20%) (Figure 1). As expected, exons are largely

devoid of TE sequence, with only 1.72% of exonic sequence consisting of TEs [1,54] (Figure

1). Conversely, intronic regions have accumulated a considerably higher proportion of TE

sequences, with 5.13% of total intronic sequence derived from TEs (Figure 1).

Overall, we observed a weak negative correlation between gene and TE density in

the monarch genome (Spearman's rank, rho = -0.369, p < 0.01), suggesting regions that

contain a high density of host genes are less likely to contain TEs. We identified 23 hotspots

of TE accumulation in the monarch genome, defined as 100kb windows with a TE content

above a 99% AUC cutoff (see Methods). This equates to windows where TE density is at

least 3.09 times higher than expected (expected = 6.25kb, lowest hotspot density = 19.34kb)

(Figure 4). These windows represent regions of the genome able to tolerate large numbers

of TE insertions. Conversely, the presence of 38 cold spots, defined as the 100kb windows

with TE content below the 1% AUC cutoff, where TE density is at least 12.56 times lower

than expected (expected = 6.25kb, highest coldspot density = 498bp), highlights regions of

the genome where TE insertions are likely not tolerated [54]. Of note, we detected 6

windows labelled as TE coldspots but gene hotspots (Figure 4). Genes present in hotspots

that are TE coldspots are involved in RNA binding, splicing, polyadenylation, mRNA stability,

and translation initiation [55]; cell polarisation [56]; neuronal development; antenna and eye

development [57–61]; and olfactory processes, which are hypothesised to be involved in

social behaviour [36] (Table S12; Additional File 3). Given the importance of these functions,

it is likely that TE insertion in these hotspot regions would be severely detrimental to host

fitness, thus individuals where TEs insert into these regions may be non-viable and purged

early in development.

Transposable Element Activity

246   The genome of *D. plexippus* is dominated by five relatively young and very

247   successful novel TE families, which collectively comprise 37.7% of total TE content (Table 2,

248   Tables S7 and S10, Figure S4; Additional Files 3 and 7).

TE Family	TE Classification	Weighted Average Age (Mya)	Copy Number
danPle_penelope-1	Penelope-like Element	10.25	7814
danPle_penelope-2	Penelope-like Element	12.84	2033
danPle_penelope-3	Penelope-like Element	10.17	2504
danPle_HaSE2-1	SINE/HaSE2	6.67	1829
danPle_r2hero-1	LINE/R2-Hero	9.91	5405

251   Table 2. Age and Copy number of the five TE families responsible for 37.7% of total TE

252   content in the monarch.

254   In total, four families are estimated to have inserted <1Mya (Figure S4, Table S9;

Additional Files 8 and 3). The low estimated age of these insertions is a sign of recent host

colonisation, potentially via horizontal transposon transfer (HTT) from another host genome

[1,62–65]. Peaks of low genetic distance to TE consensus sequence indicate an abundance

of closely related elements and thus recent TE proliferation (Figure 2, Table S8; Additional

File 3). There has been a recent burst of activity in two Tc1-Mariner elements: *danPle_TcM-*

1 (499 new copies, from ~0.68Mya), and *danPle_TcM-2* (275 new copies, from ~0.34Mya),

with insertion rates of 733 insertions/My and 798 insertions/My, respectively (Tables S7 and

S9; Additional File 3). The identified Tc1-Mariner elements are closely related and located in

Clade 130 in a recent phylogenetic study of Tc1/mariner diversity [66], which contains

elements from diverse insect species, including butterflies, aphids and ants (Figure S5;

Additional File 8). Ants and aphids are known to share environments with monarch larvae on

milkweed plants [67], increasing the potential for such HTT events [64,68]. The insertion

rates of these Tc1-mariner elements are remarkably similar to the estimated insertion rate of

the most successful *Penelope* family, at 759 insertions/My, highlighting their great success

within the monarch genome when compared to the average TE insertion rate of just 8

insertions/My (Figure S6; Additional File 9). This recent expansion of DNA TEs is interesting

given the historic dominance of LINEs in the monarch genome (2.38% of the genome, 38%

of total TE content). Whilst there are more DNA TE families than LINE families in the

monarch genome (257 vs 207), LINEs have been more successful in proliferating and

colonising the genome than their DNA TE counterparts to date, although the reasons for this

are unknown. However, it appears that these new DNA TE families are challenging this

historic trend (Figure 5, Figure S4, Figure S7; Additional Files 7 and 10).

The oldest TE insertions were estimated to be ~86 Mya, predating the emergence of

the monarch ~32Mya [69], with the first invaders being rolling circle, SINE and LTR elements

(Figure 2, Figure 5, Figure S4, Table S8; Additional Files 7 and 3). These TEs likely

proliferated around the time that the Nymphalidae family emerged, ~91Mya (CI: 71-112Mya)

[69]. However, the majority of TE families are estimated to have inserted less than ~45Mya

(mean TE family age = 25.41Mya), after the speciation of the monarch (~32Mya [69]) (Figure

2). In total, 246 families (35.4%) are estimated to pre-date the emergence of the monarch,

whilst 448 families (64.6%) have emerged on the monarch branch (Figure 5). TEs estimated

to have inserted <25Mya are predominantly DNA, Rolling Circle, and LINE elements (Figure

2).

Turnover of LINE TEs

289   LINEs are reverse transcribed from their 3' end during mobilisation, and are often

290   incomplete at the 5' end as a consequence of the premature dissociation of reverse

291   transcriptase (RT) or the activity of cellular RNases [70,71]. In contrast, the absence of a

292   complete 3' region is unlikely to arise due to either process, and is evidence of a genomic

293   deletion, presumably as a consequence of ectopic recombination between homologous LINE

294   sequences [70,71].

295   Consistent with observations in *H. melpomene* [71], we find LINE fragments across

all families exhibiting patterns suggestive of genomic deletions acting to remove these

elements from the genome (Table 3, Figure 6, Figure S8, Figure S9; Additional Files 11 and

12). In total, 68.90% of LINE fragments contain an intact 3' region, while 31.10% of LINE

fragments are truncated at the 3' end. For the *CR1*, *I-Jockey*, *L2*, and *RTE-BovB* families,

most LINE fragments display patterns suggestive of genomic deletions (i.e. truncated at the

3' end), whilst the majority of fragments of the *R1*, *R2-Hero*, and *RTE-RTE* families display

patterns indicative of processes associated with degradation of LINEs, such as premature

disassociation of RT, or breakdown via cellular RNAse activity (Table 3).

304   Swathes of LINE fragments across all LINE families (mean LINE length = 411bp),

suggest a high rate of turnover of LINEs within the monarch genome, in contrast to patterns

observed in mammals, where low rates of DNA loss are observed [71,72]. This turnover

could occur through either: (i) some form of host mechanism resulting in LINE removal,

particularly if LINEs are associated with a high fitness cost, or (ii) removal and degradation of

LINEs due to their particularly repetitive nature, which provides loci for ectopic recombination

resulting in their partial removal from the host genome [1,19].

LINE Classification	Putative Removal Process	Fragment Types	Count	Proportion of Classification Type
CR1	Premature RT Dissociation/Activity of Cellular RNAses	Lone 3', Internal + 3'	16	0.36
	Putative Genomic Deletion	Lone 5', 5' + Internal, Internal Lacking 3'	29	0.64
I-Jockey	Premature RT Dissociation/Activity of Cellular RNAses	Lone 3', Internal + 3'	42	0.20
	Putative Genomic Deletion	Lone 5', 5' + Internal, Internal Lacking 3'	164	0.80
L2	Premature RT Dissociation/Activity of Cellular RNAses	Lone 3', Internal + 3'	61	0.20
	Putative Genomic Deletion	Lone 5', 5' + Internal, Internal Lacking 3'	244	0.80
R1	Premature RT Dissociation/Activity of Cellular RNAses	Lone 3', Internal + 3'	223	0.55
	Putative Genomic Deletion	Lone 5', 5' + Internal, Internal Lacking 3'	181	0.45
R2-Hero	Premature RT Dissociation/Activity of Cellular RNAses	Lone 3', Internal + 3'	2034	0.88
	Putative Genomic Deletion	Lone 5', 5' + Internal, Internal Lacking 3'	282	0.12
RTE-BovB	Premature RT Dissociation/Activity of Cellular RNAses	Lone 3', Internal + 3'	225	0.46
	Putative Genomic Deletion	Lone 5', 5' + Internal, Internal Lacking 3'	268	0.54
RTE-RTE	Premature RT Dissociation/Activity of Cellular RNAses	Lone 3', Internal + 3'	137	0.67
	Putative Genomic Deletion	Lone 5', 5' + Internal, Internal Lacking 3'	68	0.33

Table 3. Number of LINEs classified by fragment type, and putative removal process.

Despite evidence of removal, LINE sequence continues to dominate the TE

landscape of the monarch. To be active, an intact LINE will contain start and stop codons,

and an RT domain [73]. In total, we identified five intact LINEs in the monarch (Table S13,

Additional File 3), including *danPle_r2hero-1*, which is the second most populous TE in the

monarch genome (Table 2). We also observe a significant accumulation of certain LINE

families, with ~20% maintaining a length above 1kb (Table S11; Additional File 3). Short

elements are more likely to persist in the genome than their longer counterparts, as they are

less prone to recombination [19]. This, combined with the high activity level of LINEs before

and throughout the evolution of the monarch (Figure 2), explains the dominance of LINE

fragments across the genomic landscape, despite their high turnover [19,71].

Association of TEs with Monarch Colouration and Immune Genes

Given a previous role for TEs in wing colouration in the peppered moth (*Biston*

betularia) [42] and the clouded yellow (*Colias croceus*) [74], we investigated the possibility of

TE associations with wing colouration in the monarch. The myosin gene is associated with

wing colouration in the monarch [75]. We identified a single unclassified element 7kb

upstream of this gene (Figure S10; Additional File 12), contrasting the location of TEs

implicated in colouration in the peppered moth and clouded yellow, where the TE insertions

were found within an intron and 6kb downstream of the gene body, respectively [42,74]. This

is not entirely unexpected, as the colouration differences seen in the peppered moth and

clouded yellow are due to altered gene expression, whilst the white *nivosus* form of the

monarch is due to altered pigment transport [35,42,74].

Immune genes are some of the fastest evolving genes across metazoans [76–79],

and so are strong candidates for exploring the role of TEs in host evolution. Across the

monarch genome, TEs of all major types were found within immune gene flanks (20kb).

When normalised to compare TE enrichment around different immune gene types, we find

no significant difference in TE base coverage (Kruskal-Wallis χ^2 = 47.17, p = 0.1221), or in

TE count (Kruskal-Wallis χ^2 = 40.351, p = 0.3244) between different types of immune

genes, suggesting no enrichment around specific classes of immune genes (Figure S11;

Additional File 13). We also find no significant difference in coverage or count of TE

classifications around immune genes (TE count by classification: Kruskal-Wallis χ^2 = 14.46,

p = 0.8493; TE coverage by classification: Kruskal-Wallis χ^2 = 125.9, p = 0.5854). However,

there are TEs within 5kb of the 5' end in 49/103 identified immune genes. These proximal

TEs are of interest given the short distance between them and the immune gene

transcription start sites (TSS) , making regulatory functions a possibility [80].

Discussion

Here, we provide a detailed annotation of TEs in the monarch genome and an

analysis of their diversity and evolution. We found a large reduction in TE sequence

compared to the original annotation performed on the draft monarch genome. We suggest

that this is due to unclassified TE sequences in the original annotation no longer being

recognised as TEs with updated TE databases and annotation tools. However, we are

unable to confirm this, due to the original *de novo* TE library and annotation results being

unavailable. This raises an important issue in genomics currently. Given TE annotation is a

key step in genome assembly and annotation, we suggest that it should be required to

include TE libraries alongside the publication of genome projects, ideally in a specific

repository, to: (i) reduce duplication of efforts required for reannotation; (ii) improve

reproducibility in annotation; and, (iii) improve TE consensus libraries to reduce cases of a

single TE family being given different names in multiple species. As the number sequenced

genomes increases, the wider availability of TE libraries will benefit a range of studies,

where gene annotations can be improved by accurate TE annotation, and the availability of

large numbers of well-curated TE sequences can facilitate large-scale studies to study TE

evolutionary dynamics.

We identified an uneven distribution of TEs across the genome in line with

expectations of TE distribution in host genomes [1] (Figures 3 and 4). TE insertions in non-

coding regions are less likely to have a detrimental impact on host genome function than TE

insertions into coding regions [1,21,54,81–83], and so are less stringently removed by

selection, enabling TE accumulation in non-coding regions. Indeed, some TEs actively avoid

coding regions by targeting regions of the genome unlikely to impact host fitness [54,84],

with this process potentially occurring in the monarch, where we find overlapping gene

hotspots and TE coldspots.

The most populous regions of the monarch genome for TEs are intergenic regions

(Figure 1). The presence of large numbers of TEs in host gene flanks is interesting, given

the implication of TEs in the evolution of gene regulatory networks [10–13,85], although this

could also be due to the relatively compact nature of the monarch genome leaving TEs with

few other loci to integrate within. TEs are implicated in rapidly generating extensive intron

gain in eukaryotic genomes, where short non-autonomous DNA transposons independently

generated hundreds to thousands of introns across the genomes of eukaryotes [1,54,86]. It

is also hypothesised that transposon insertion-associated introns are responsible for the

prevalence of nucleosome-sized exons observed in eukaryotes [86]. The presence of TEs in

intronic sequence of the monarch could therefore indicate genomic refugia, where TE

insertions are less likely to be detrimental to the host, but they could also indicate ongoing

intron generation, where new insertions might lead to the generation of new, or longer,

introns over evolutionary time.

The presence of young and highly populous TE families in the monarch genome

demonstrates the success of TEs in naïve host genomes [84,87–90]. The putative recent

arrival of several young DNA TEs, namely two novel Tc1-Mariner families (*danPle_TcM-1*

and *danPle_TcM-2*), has led to another wave of TE proliferation. Over time, it is

hypothesised that the activity of these families will decrease as the host develops

mechanisms to suppress activity, either through capture of TE sequences in piRNA regions,

or through other defensive mechanisms such as methylation [90–92]. A reduction in

transposition activity of DDE transposase-containing DNA transposons can also be

attributed to increases in non-autonomous copies [66]. In the meantime, however, these TEs

are proliferating and dynamically altering the genomic landscape of *D. plexippus*, with

potential connotations for ongoing genome structural and functional evolution [1–3,19].

While factors leading to the relatively low TE content observed in the monarch

compared to other *Danaus* species are currently unknown, it is clear TEs have been unable

to accumulate to the same level as other members of the genus. The maintenance of a small

genome in the monarch can likely be partly attributed to high turnover of TEs, evidenced by

the lack of ancient peaks in repeat landscapes and the presence of few intact LINEs, many

of which show hallmarks of genomic deletions. To remove LINEs, ectopic recombination

between similar elements is likely to play a role, and it has been postulated that an increased

rate of ectopic recombination could be responsible for the differences in accumulation of TEs

in insects and mammals [93]. A lower ectopic recombination rate in mammals, despite

frequent large TE loads, enables older elements to accumulate, whereas high ectopic

recombination rates in insects can lead to removal of TEs before older copies accumulate in

large numbers [93]. Given our results, it is likely that ectopic recombination is occurring in

the monarch to remove LINEs. Indeed, the only families with an abundance of 3' fragments

are, *R1*, *R2-Hero*, and *RTE-RTE* (mean insertion times ~16.02Mya, ~9.9Mya, and

~27.38Mya). It is tempting to suggest that *R1*, *R2-Hero*, and *RTE-RTE* insertions have a

lower impact on host fitness, and so are "allowed" to degrade over time, rather than being

removed *en masse* through genomic mechanisms. However, it must also be noted that

ectopic recombination is a passive process. Whilst it is more likely that LINEs are being

fragmented and removed through ectopic recombination between copies, this does not

account for inter-family differences in the ratio of fragments undergoing degradation through

ectopic recombination versus degradation via the activity of RT or cellular RNAses. In the

case of the most populous family, *R2-Hero*, it is likely that these elements have so far

avoided ectopic recombination, given these elements are young compared to other LINEs

(insertion ~9.9Mya) (Figure 5, Table S8; Additional File 3)). Our results are consistent with

those of previous findings in *Heliconius* species [94], where a high TE turnover and lack of

ancient TEs was observed, suggesting these characteristics might be widespread among

lepidopterans, although further comparative studies will be required to confirm this.

In addition to TE turnover, genome size is likely also maintained through

sequestering TE activity via epigenetic mechanisms and piRNAs [1,91,95,96]. Further

research to characterise piRNA clusters in the monarch, and interrogation of its

transcriptome, will further improve our understanding of how these processes have shaped

the TE landscape within the monarch genome.

The lack of closer or more associations of TEs with the myosin gene associated with

warning colouration in the monarch is perhaps not unsurprising, as it has been postulated

that the white *nivosus* form of the monarch arises from altered pigment transport [35], rather

than altered gene expression, as is the case with the peppered moth and clouded yellow

[42,74]. Whilst TEs were not significantly enriched around immune genes, the prominence of

TEs around immune genes is nevertheless intriguing, as is the presence of TEs <5kb from

the TSS of 49 immune genes, which opens up the possibility for TEs to alter the regulation of

these genes. To further understand the impact of TEs in these regions, TE knockout studies

in combination with transcriptomic analyses may be able to elucidate any impacts TEs might

be having on immune genes in the monarch.

Conclusions

In-depth studies into TE landscapes and dynamics within host species of broad

scientific and public interest remain relatively few, even with widespread acknowledgement

of the key role that TEs have played in eukaryotic evolution [1,48,97]. We provide a detailed

annotation of TEs in the monarch genome and an analysis of TE diversity and evolution,

including the identification of highly successful young DNA TE families, which mirror the

activity profile of the most successful LINEs found in the monarch, and evidence of ongoing

TE expansion and removal in *D. plexippus*, highlighting the dynamic nature of repeat content

within genomes over time. Further large-scale comparative studies will be greatly beneficial

to further our understanding of the evolutionary dynamics of TEs, and the drivers leading to

diverse TE landscapes across closely related species.

Materials and Methods

Transposable Element Annotation

The monarch chromosomal assembly, Dplex_v4, was downloaded from NCBI (Refseq GCF_009731565.1), along with corresponding genome annotations in GFF format [43,98]. The monarch genome assembly is 248.676Mb, with a GC content of 32.1%, and comprises 4,115 scaffolds with a scaffold N50 of 9.21Mb.

The *D. plexippus* genome was annotated using a bespoke automated in-house TE annotation pipeline, as previously described [99–104]. For this, known repeats were first identified and masked using RepeatMasker (version 4.1.0) [105] using the *Arthropoda* library from RepBase (version 23.08) and Dfam (release 3.3) [106,107]. Low complexity repeats and RNA were not masked (*-nolow and -norna*) and a sensitive search was performed (*-s*). Following this, a *de novo* repeat library was constructed using RepeatModeler (version 1.0.11), with RECON (version 1.08) and RepeatScout (version 1.0.5) [108–110]. To generate maximum-length consensus sequences for the *de novo* repeat library, novel repeats identified by RepeatModeler were curated using an automated version of the 'BLAST, Extract, Extend' process [28]. A BLASTn search was performed on the *D. plexippus* genome to obtain up to the top 40 hits for each TE family identified by RepeatModeler [111]. Sequences were extracted with 1,000 base pairs of flanking sequence at the 5' and 3' ends. Each set of family sequences were aligned using MAFFT (version 7.453) [112]. Alignments were trimmed with trimAl (version 1.4.rev22) to retain high-quality positions in the alignment (*-gt* 0.6 *-cons* 60) [113]. Updated consensus sequences were computed with EMBOSS cons (*-plurality* 3) to generate a new TE library featuring consensus sequences with extended flanks [114]. This process was repeated through 5 iterations to obtain maximum-length consensus sequences. The resulting *de novo* repeat library was combined with the RepBase *Arthropoda* library and utilised to annotate repetitive elements using RepeatMasker with a score threshold at the relatively conservative level of 400 (*-cutoff* 400), to exclude poor

matches unlikely to be true TE sequences. Following this, the *D. plexippus* genome was

analysed with LTR_Finder (version 1.07), using the LTR_Finder parallel wrapper [115,116]

to identify full-length LTR elements. Final TE annotations were defragmented and refined

using the loose merge (*-loose*) command in RepeatCraft [117]. Overlapping annotations

were removed using MGKit (version 0.4.1) filter-gff, with the longest element of overlapping

annotations retained (*-c* length *-a* length) [118]. Finally, all repeats less than 100bp in length

were removed before the final TE quantification to decrease spurious hits. Final repeat

annotation file is provided in dataset S3 (Additional File 17).

To assess the performance of the automated annotation method, a manual curation

of the *de novo* library was also performed. Briefly, for this, the 'BLAST, Extract, Extend'

process was performed manually, rather than using the automated pipeline. For each

iteration, up to the top 40 hits for each TE family were extracted from the *D. plexippus*

genome using BLASTn [111]. Sequences were extracted with 1,000 base pairs of flanking

sequence at the 5' and 3' ends, before each set of family sequences was aligned using

MAFFT [112]. Alignments were then visualised using AliView, and poorly-aligned positions

and ends were trimmed where single-copy DNA was present [119]. If either the 5' or 3' end

of the alignment terminated within the repetitive region, the BLASTn and extension process

was repeated until single-copy DNA was observed at both the 5' and 3' ends of the

alignment for each family, signalling the end of the repeat and the transition to host genome

sequence. New consensus sequences were generated using EMBOSS cons to create a

new TE library of full-length elements. The presence of duplicate consensus sequences was

checked with *cd-hit-est* using stringent parameters (*-n* 10 *-c* 0.95) and any duplicates were

removed, retaining the longest consensus sequence only [46,47]. This manually curated *de*

novo library was combined with the RepBase *Arthropoda* library and utilised to provide a

separate TE annotation for the monarch genome. Full-length LTR elements were again

annotated using LTRFinder in the LTRFinder parallel wrapper and the final TE annotations

were defragmented and refined using RepeatCraft loose merge. The same post-processing

was applied to the annotations as with the automated curation.

Comparison of results from automated and manually curated libraries

The automated and manually curated libraries were compared to assess the

performance of the automated curation process. Final annotation quantifications were

compared using a One-way ANOVA, as observations were normally distributed for both

groups (Automated annotation: Shapiro-Wilk, $W = 0.88401$, $p > 0.05$; Manual annotation:

Shapiro-Wilk, $W = 0.90517$, $p > 0.05$). Comparisons were performed in R (version 3.6.3)

using the Rstudio IDE [120,121]. As differences in quantification between the two methods

were not statistically significant (see Results), the annotations from the automated process

were used for downstream analyses to maximise the reproducibility of this study. For

discussion, see [Additional File 18].

Identifying Transposable Element Association with Genomic Compartments

The *D. plexippus* gene annotation file was processed to generate a GFF file

containing coordinates of exons, introns, and 5' and 3' flanking regions. Gene flanking

regions are defined here as 20kb directly upstream or downstream of the gene body. These

flanking regions are determined to identify TEs at a distance that could be described as the

proximate promoter region, rather than just accounting for the core promoter region, to

include both promoter and more distal enhancer regions for genes [80]. Bedtools intersect

(version 2.27.1) was used to identify the number of bases of overlap (*-wao*) between TEs

and different genomic compartments [122]. Following this, quantification was performed in R.

TE Hotspots and Coldspots

To identify TE and gene hotspots and coldspots, chromosomal scaffolds were split

into 100kb windows, which were defined using the frequency distribution of observations. A

polynomial model was fitted to the frequency distribution, to generate a smooth curve

through the data points using the lm function in R. Using these models, the total area under

the curve (AUC) was calculated using the area_under_curve function of the bayestestR

package [123]. From this, the 1% and 99% AUC cutoff values were determined. Starting with

the first data point and iteratively adding subsequent datapoints with each round, the AUC

for each collection of points was determined. The maximum AUC value not exceeding 99%

of the total AUC was taken as the cutoff where windows containing higher TE or gene

coverage were considered to be hotspots. The minimum AUC value not exceeding 1% of the

total AUC was taken as the cutoff where windows containing lower TE or gene coverage

were considered to be coldspots (Figure S12; Additional File 15). Karyoplots of the *D.*

plexippus genome, illustrating gene and TE locations, highlighting TE hotspots and

coldspots were generated using the KaryoploteR package in R [124].

Estimating TE Age and Activity

TE activity dates were estimated using Kimura2-parameter genetic distances

(including CpG sites) calculated between individual insertions and the consensus sequence

of each TE family [125]. A neutral mutation rate of 0.0116 substitutions per site per million

years was used for *D. plexippus*, as previously described for monarch [126].

Turnover of LINEs

To interrogate LINE decay patterns, each LINE sequence was extracted from the

genome using bedtools getfasta [122]. LINE sequences were used to query the repeat

library using BLASTn with tabular output specified (*-outfmt* 6). In the event of multiple

matches, only the top match was retained (defined by percent identity). Coordinates of each

match were scaled to normalise the length of each LINE consensus sequence. Mapped

LINE fragments were plotted using ggplot2 of the tidyverse package in R [127,128].

Identification of Intact LINEs

To identify full-length LINEs, all elements longer than 2,700bp were extracted from

the monarch genome using bedtools getFasta [122]. Open reading frames (ORFs) were

detected using EMBOSS getORF [114]. ORF sequences were filtered to retain all

sequences > 600aa. RPSBLAST+ (v2.9.0+) [111] was used to query the ORF sequences

against the CDD database [129] to identify conserved protein domains. Sequences were

classified as intact LINEs based on the presence of relevant protein domains as previously

described [130] (Table S13, Additional File 3).

Classification of Novel DNA TEs

To further classify the novel DNA elements in the monarch beyond homology to

known elements, as used by RepeatModeler. DDE transposases were compared with

curated alignments for all DDE TE superfamilies [66]. Specifically, dashes in curated

alignments of all known DDE TE sequences for each of the 19 families were removed, and

the resultant fasta file was used as the subject in a blastx search against the *de novo*

monarch TE library, with an e-value threshold of 1×10^{-20} and tabular output specified (-

evalue 1e-20 *-outfmt* 6) [111]. The best match for each *de novo* query sequence was

retained, with a minimum 50% identity over at least 50% of the total length of the curated

DDE subject sequence. Following this, novel DNA elements were only detected for the Tc1-

Mariner and Sola1 families (Figures S5 and S13; Additional Files 8 and 16). Transposase

sequences for the *de novo* monarch sequences were extracted and aligned to the original

DDE alignments for their respective families using the profile alignment command in MAFFT

[112]. Sequences from related transposon superfamilies were used as outgroups, as

described [66]. To estimate phylogenies for the two DDE superfamilies, we performed 1,000

ultra-fast bootstrap replicates in IQ-TREE (v1.6.12) [131] specifying the best fit amino acid

model for each family as indicated by ModelFinder [132]. Phylogenetic trees were visualised

in figtree (v1.4.4) (http://tree.bio.ed.ac.uk/software/figtree/).

Association of TEs with Monarch Colouration and Immune Genes

We obtained the transcript sequence in fasta format for the myosin gene responsible

for wing colouration from MonarchBase, under accession DPOGS206617 [34]. To identify

the matching region in the v4 monarch assembly, a blastn search was performed to query

the monarch assembly [111]. The coordinates of the myosin gene were obtained and used in

a bedtools window search to identify TEs within 20kb (-*w* 20000)[122]. A karyoplot of the

myosin gene region including 20kb flanks was generated in R using the karyoploteR

package [124].

To investigate associations between monarch immunity genes and TEs, nucleotide

and protein sequences of monarch immune genes were obtained in fasta format from

MonarchBase [34]. To identify corresponding regions in the v4 monarch assembly, blastn

and tblastn searches were used to query the monarch genome assembly with the monarch

immune gene nucleotide and protein sequences [111]. Matches were filtered by percentage

identity and query match coordinates to identify the locus of each immune gene. A bed file

was generated, containing the name and location of each immune gene, and a bedtools

window search was performed to identify TEs within 20kb of each immune gene(-*w*

20000)[122]. Karyoplots of each immune gene region including 20kb flanks were generated

in R using the karyoploteR package [124]. TE bases surrounding each immune gene were

quantified in R [120], and normalised to account for variation in the number of each gene

type. As TE coverage and count were not normally distributed, a Kruskal-Wallis test was

used to compare TE coverage and count between different immune gene types. Unless

otherwise stated, all plots were generated using the ggplot2 package of the tidyverse suite in

R [127,128].

Figure Legends

Figure 1. TE content and location in the monarch. Major TE types are represented by

different colours indicated in the key (A) Pie chart illustrating proportions of the monarch

genome comprised of the main TE classifications. (B) Bars illustrating the number of base

pairs of the monarch genome found in different genomic compartments. (C) Bars illustrating

the proportion of each genomic compartment derived from bases annotated as TE

sequence.

Figure 2. Repeat landscape of the monarch. The x axis indicates the level of Kimura2-

parameter genetic distance observed between TE insertions and their respective consensus

sequences in percent. More recent elements are located to the right of the x axis. The y axis

indicates the percentage of the genome occupied by TE insertions of each genetic distance

class.

Figure 3. Violin plot showing kilobases of TE sequence per 100kb window across the

genome for each of the main TE classifications. Red points show the mean kilobases of TE

sequence per 100kb.

Figure 4. Karyoplots illustrating monarch chromosomes, with genes represented as black

bars. (A) Above each chromosome, TE density is shown for 100kb windows, with major TE

types represented by different colours indicated in the key. (B) Above each chromosome, TE

hotspots and coldspots are shown, with severity represented by the different colours

indicated in the key,

Figure 5. Violin plot illustrating the estimated insertion time (Mya) for each of the major TE

types, represented by different colours indicated in the key. The black bar shows the

estimated divergence time of the monarch, approximately 32Mya [69].

Figure 6. Plot illustrating LINE fragments annotated in the monarch genome. The x axis

indicates normalised start and end coordinates for each LINE element. Elements are

organised from longest to shortest, with short elements found towards the top of the plot.

LINE families are represented by different colours indicated by the key.

646   Additional Files

Additional File 1: Dataset S1. *De novo* transposable element consensus sequences

identified using the automated curation method, in FASTA format.

Additional File 2: Dataset S2. *De novo* transposable element consensus sequences

identified using the manual curation method, in FASTA format.

Additional File 3: Supplementary Tables, including contents page with table legends. Tables

S1-S13.

Additional File 4: Figure S1. Set of plots comparing the final TE quantifications between the

manual and automated *de novo* libraries, both in conjunction with RepBase *Arthropoda*

library and Dfam. Major TE types represented by different colours, indicated in the key. Top

row: Stacked bar showing number of elements identified with each library. Scatter plot

showing number of elements identified with each library, lines indicate matching TE

classifications. Box plot showing number of elements identified with each library. Middle row:

Stacked bar showing base pair coverage for TEs annotated with each library. Scatter plot

showing base pair coverage for TEs annotated with each library, lines indicate matching TE

classifications. Box plot showing base pair coverage for TEs annotated with each library.

Bottom row: Stacked bar showing proportion of monarch genome comprised of TEs

annotated with each library. Scatter plot showing proportion of monarch genome comprised

of TEs annotated with each library, lines indicate matching TE classifications. Box plot

showing proportion of monarch genome comprised of TEs annotated with each library.

Additional File 5: Figure S2. Scatter plots showing the length of consensus sequences

identified in the manual and automated curation methods. Lines join consensus sequences

identical prior to BLAST, Extract, Extend process. Major TE types are represented by

different colours indicated in the key. Orphaned points represent consensus sequences only

found in one of the two libraries at the end of the BLAST, Extract, Extend process.

Additional File 6: Figure S3. Repeat landscape for *Danaus chrysippus* and *Danaus*

melanippus. The x axis indicates the level of Kimura2-parameter genetic distance observed

between TE insertions and their respective consensus sequences in percent. More recent

elements are located to the right of the x axis. The y axis indicates the percentage of the

genome occupied by TE insertions of each genetic distance class.

Additional File 7: Figure S4. Scatter plot illustrating estimated TE age for each TE family.

Major TE types are represented by different colours indicated in the key. Elements are

ordered from oldest to youngest, within each major type.

Additional File 8: Figure S5. Phylogenetic tree of Tc1-Mariner DDE transposases, with novel

monarch families added (highlighted in red). Branch support from 1000 ultra-fast bootstrap

repetitions. Novel Tc1-Mariner families highlighted in teal are the young TEs found with

extremely high copy number.

Additional File 9: Figure S6. Scatter plots illustrating the number of insertions per million

years for each TE family. Major TE types are represented by different colours indicated in

the key.

Additional File 10: Figure S7. Scatter plots illustrating TE age against copy number. Major

TE types are represented by different colours indicated in the key. X axis shows estimated

age of each TE family, Y axis shows TE copy number identified in the monarch.

Additional File 11: Figure S8. Plots illustrating LINE fragments annotated in the monarch

genome, separated by LINE family. The x axis indicates normalised start and end

coordinates for each LINE element. Elements are organised from longest to shortest, with

short elements found towards the top of the plot. LINE families are represented by different

colours indicated by the key.

Additional File 12: Figure S9. Bar plots illustrating the abundance of LINE fragments

identified in the monarch. LINE classifications are represented by different colours as

indicated in the key.

Additional File 13: Figure S10. Karyoplot of the monarch gene association with colouration,

myosin, obtained from MonarchBase under accession DPOGS206617. Main TE types within

20kb are represented by different colours indicated in the key.

Additional File 14: Figure S11. Boxplots illustrating: (A) Normalised TE coverage in 20kb

regions surrounding each immune gene type. (B) Normalised TE count in 20kb regions

surrounding each immune gene type. Main TE types are represented by different colours

indicated in the key.

Additional File 15: Figure S12. Plots illustrating: (A) Distribution of genome windows with TE

coverage, with lines showing 1% and 99% AUC cutoffs used to define hotspots and

coldspots. (B) Distribution of genome windows with host gene coverage, with lines showing

1% and 99% AUC cutoffs used to define hotspots and coldspots.

Additional File 16: Figure S13. Phylogenetic tree of Sola1 DDE transposases, with novel

monarch families added (highlighted in red). Branch support from 1000 ultra-fast bootstrap

repetitions.

Additional File 17: Dataset S3. Coordinates and classification of all transposable element

sequences identified in the monarch genome using the automated *de novo* library in

conjunction with RepBase *Arthropoda* library and Dfam, in bed format.

Additional File 18: Discussion on the comparison between our automated annotation method

and a basic manual curation.

Abbreviations

TE: Transposable element

Non-LTR: Non-long terminal repeat

LTR: Long terminal repeat

LINEs: Long INterspersed Elements

SINEs: Short INterspersed Elements

TSS: Transcription Start Site

References

[1]   Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, et al. Ten

749   things you should know about transposable elements. Genome Biol 2018;19:199.

750   https://doi.org/10.1186/s13059-018-1577-z.

751   [2]   Klein SJ, O'Neill RJ. Transposable elements: genome innovation, chromosome

752   diversity, and centromere conflict. Chromosom Res 2018;26:5–23.

753   https://doi.org/10.1007/s10577-017-9569-5.

754   [3]   Feschotte C, Pritham EJ. DNA Transposons and the Evolution of Eukaryotic

755   Genomes. Annu Rev Genet 2007;41:331–68.

756   https://doi.org/10.1146/annurev.genet.40.110405.090448.

757   [4]   Belyayev A. Bursts of transposable elements as an evolutionary driving force. J Evol

758   Biol 2014;27:2573–84. https://doi.org/10.1111/jeb.12513.

759   [5]   Schrader L, Schmitz J. The impact of transposable elements in adaptive evolution.

760   Mol Ecol 2018. https://doi.org/10.1111/mec.14794.

761   [6]   Casola C, Lawing AM, Betrán E, Feschotte C. PIF-like transposons are common in

762   Drosophila and have been repeatedly domesticated to generate new host genes. Mol

763   Biol Evol 2007;24:1872–88. https://doi.org/10.1093/molbev/msm116.

764   [7]   Jangam D, Feschotte C, Betrán E. Transposable Element Domestication As an

765   Adaptation to Evolutionary Conflicts. Trends Genet 2017;33:817–31.

766   https://doi.org/10.1016/j.tig.2017.07.011.

767   [8]   Volff JN. Turning junk into gold: Domestication of transposable elements and the

768   creation of new genes in eukaryotes. BioEssays 2006.

769   https://doi.org/10.1002/bies.20452.

770   [9]   Mita P, Boeke JD. How retrotransposons shape genome regulation. Curr Opin Genet

771   Dev 2016;37:90–100. https://doi.org/10.1016/j.gde.2016.01.001.

[10]   Bourque G. Transposable elements in gene regulation and in the evolution of

vertebrate genomes. Curr Opin Genet Dev 2009;19:607–12.

https://doi.org/10.1016/j.gde.2009.10.013.

[11] Feschotte C. Transposable elements and the evolution of regulatory networks. Nat

Rev Genet 2008;9:397–405. https://doi.org/10.1038/nrg2337.

[12] Rebollo R, Romanish MT, Mager DL. Transposable Elements: An Abundant and

Natural Source of Regulatory Sequences for Host Genes. Annu Rev Genet

2012;46:21–42. https://doi.org/10.1146/annurev-genet-110711-155621.

[13] Chuong EB, Elde NC, Feschotte C. Regulatory activities of transposable elements:

From conflicts to benefits. Nat Rev Genet 2017;18:71–86.

https://doi.org/10.1038/nrg.2016.139.

[14] Bennetzen JL. Transposable element contributions to plant genome evolution. Plant

Mol Biol 2000;42:251–69. https://doi.org/10.1023/A:1006344508454.

[15] Casals F, Cáceres M, Ruiz A. The Foldback-like transposon Galileo is involved in the

generation of two different natural chromosomal inversions of Drosophila buzzatii. Mol

Biol Evol 2003;20:674–85. https://doi.org/10.1093/molbev/msg070.

[16] Campos-Sánchez R, Cremona MA, Pini A, Chiaromonte F, Makova KD. Integration

and Fixation Preferences of Human and Mouse Endogenous Retroviruses Uncovered

with Functional Data Analysis. PLoS Comput Biol 2016;12:1–41.

https://doi.org/10.1371/journal.pcbi.1004956.

[17] Oliver KR, Greene WK. Transposable elements: Powerful facilitators of evolution.

BioEssays 2009;31:703–14. https://doi.org/10.1002/bies.200800219.

[18] Hughes JF, Coffin JM. Human endogenous retroviral elements as indicators of

ectopic recombination events in the primate genome. Genetics 2005;171:1183–94.

https://doi.org/10.1534/genetics.105.043976.

[19] Kent T V., Uzunović J, Wright SI. Coevolution between transposable elements and

recombination. Philos Trans R Soc B Biol Sci 2017;372.

https://doi.org/10.1098/rstb.2016.0458.

[20] Coates BS, Hellmich RL, Grant DM, Abel CA. Mobilizing the genome of lepidoptera

through novel sequence gains and end creation by non-autonomous Lep1 Helitrons.

DNA Res 2012;19:11–21. https://doi.org/10.1093/dnares/dsr038.

[21] Biémont C, Vieira C. What transposable elements tell us about genome organization

and evolution: The case of Drosophila. Cytogenet Genome Res 2005;110:25–34.

https://doi.org/10.1159/000084935.

[22] Morgante M, Brunner S, Pea G, Fengler K, Zuccolo A, Rafalski A. Gene duplication

and exon shuffling by helitron-like transposons generate intraspecies diversity in

maize. Nat Genet 2005;37:997–1002. https://doi.org/10.1038/ng1615.

[23] Philippsen GS, Avaca-Crusca JS, Araujo APU, DeMarco R. Distribution patterns and

impact of transposable elements in genes of green algae. Gene 2016;594:151–9.

https://doi.org/10.1016/j.gene.2016.09.012.

[24] Wu M, Li L, Sun Z. Transposable element fragments in protein-coding regions and

their contributions to human functional proteins. Gene 2007;401:165–71.

https://doi.org/10.1016/j.gene.2007.07.012.

[25] Daniel C, Behm M, Öhman M. The role of Alu elements in the cis-regulation of RNA

processing. Cell Mol Life Sci 2015;72:4063–76. https://doi.org/10.1007/s00018-015-

1990-3.

[26] Vorechovsky I. Transposable elements in disease-associated cryptic exons. Hum

Genet 2010;127:135–54. https://doi.org/10.1007/s00439-009-0752-4.

[27] Bejerano G, Lowe CB, Ahituv N, King B, Siepel A, Salama SR, et al. A distal

enhancer and an ultraconserved exon are derived from a novel retroposon. Nature

2006;441:87–90. https://doi.org/10.1038/nature04696.

[28]   Platt RN, Blanco-Berdugo L, Ray DA. Accurate transposable element annotation is

vital when analyzing new genome assemblies. Genome Biol Evol 2016;8:403–10.

https://doi.org/10.1093/gbe/evw009.

[29]   Bates HW. Contributions to an Insect Fauna of the Amazon Valley.—Lepidoptera:—

Heliconinæ. J Proc Linn Soc London Zool 1862;6:73–7.

https://doi.org/10.1111/J.1096-3642.1862.TB00932.X.

[30]   Müller F. Ituna and Thyridia: a remarkable case of mimicry in butterflies. Trans

Entomol Soc L 1879;1879:20–9.

[31]   Ehrlich PR, Raven PH. Butterflies and Plants: A Study in Coevolution. Evolution (N Y)

1964;18:586. https://doi.org/10.2307/2406212.

[32]   Kettlewell, B. Evolution of melanism: the study of a recurring necessity 1973.

[33]   Dasmahapatra KK, Walters JR, Briscoe AD, Davey JW, Whibley A, Nadeau NJ, et al.

Butterfly genome reveals promiscuous exchange of mimicry adaptations among

species. Nature 2012;487:94–8. https://doi.org/10.1038/nature11041.

[34]   Zhan S, Reppert SM. MonarchBase: the monarch butterfly genome database. Nucleic

Acids Res 2013;41:D758–63. https://doi.org/10.1093/NAR/GKS1057.

[35]   Zhan S, Zhang W, Niitepõld K, Hsu J, Haeger JF, Zalucki MP, et al. The genetics of

monarch butterfly migration and warning colouration. Nature 2014;514:317–21.

https://doi.org/10.1038/nature13812.

[36]   Zhan S, Merlin C, Boore JL, Reppert SM. The monarch butterfly genome yields

insights into long-distance migration. Cell 2011;147:1171–85.

https://doi.org/10.1016/j.cell.2011.09.052.

[37]   Dasmahapatra KK, Walters JR, Briscoe AD, Davey JW, Whibley A, Nadeau NJ, et al.

Butterfly genome reveals promiscuous exchange of mimicry adaptations among

species. Nature 2012;487:94–8. https://doi.org/10.1038/nature11041.

[38]   Wu M, Sun Z, Luo G, Hu C, Zhang W, Han Z. Cloning and characterization of

849   piggyBac-like elements in lepidopteran insects. Genetica 2011;139:149–54.

850   https://doi.org/10.1007/s10709-010-9542-0.

851   [39]   Sormacheva I, Smyshlyaev G, Mayorov V, Blinov A, Novikov A, Novikova O. Vertical

852   evolution and horizontal transfer of CR1 non-LTR retrotransposons and Tc1/mariner

853   DNA transposons in Lepidoptera species. Mol Biol Evol 2012;29:3685–702.

854   https://doi.org/10.1093/molbev/mss181.

855   [40]   Reiss D, Mialdea G, Miele V, de Vienne D, Peccoud J, Gilbert C, et al. Global survey

856   of mobile DNA horizontal transfer in arthropods reveals Lepidoptera as a prime

857   hotspot. PLOS Genet 2019;15:e1007965.

858   https://doi.org/10.1371/journal.pgen.1007965.

859   [41]   Talla V, Suh A, Kalsoom F, Dinca V, Vila R, Friberg M, et al. Rapid increase in

860   genome size as a consequence of transposable element hyperactivity in wood-white

861   (leptidea) butterflies. Genome Biol Evol 2017;9:2491–505.

862   https://doi.org/10.1093/gbe/evx163.

863   [42]   Hof AEV t., Campagne P, Rigden DJ, Yung CJ, Lingley J, Quail MA, et al. The

864   industrial melanism mutation in British peppered moths is a transposable element.

865   Nature 2016;534:102–5. https://doi.org/10.1038/nature17951.

866   [43]   Gu L, Reilly PF, Lewis JJ, Reed RD, Andolfatto P, Walters JR. Dichotomy of dosage

867   compensation along the neo Z chromosome of the Monarch butterfly. Curr Biol

868   2019;29:4071–7.

869   [44]   Wang J, Wang A, Han Z, Zhang Z, Li F, Li X. Characterization of Three Novel SINE

870   Families with Unusual Features in Helicoverpa armigera. PLoS One 2012;7:e31355.

871   https://doi.org/10.1371/JOURNAL.PONE.0031355.

[45]   Ou S, Su W, Liao Y, Chougule K, Agda JRA, Hellinga AJ, et al. Benchmarking

transposable element annotation methods for creation of a streamlined,

comprehensive pipeline. Genome Biol 2019;20:1–45. https://doi.org/10.1186/s13059-

019-1905-y.

[46]   Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of

protein or nucleotide sequences. Bioinformatics 2006;22:1658–9.

[47]   Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-

generation sequencing data. Bioinformatics 2012;28:3150–2.

[48]   Chénais B, Caruso A, Hiard S, Casse N. The impact of transposable elements on

eukaryotic genomes: From genome size increase to genetic adaptation to stressful

environments. Gene 2012;509:7–15. https://doi.org/10.1016/j.gene.2012.07.042.

[49]   Osanai-Futahashi M, Suetsugu Y, Mita K, Fujiwara H. Genome-wide screening and

characterization of transposable elements and their distribution analysis in the

silkworm, Bombyx mori. Insect Biochem Mol Biol 2008;38:1046–57.

https://doi.org/10.1016/j.ibmb.2008.05.012.

[50]   Kapusta A, Suh A, Feschotte C. Dynamics of genome size evolution in birds and

mammals. Proc Natl Acad Sci U S A 2017;114:E1460–9.

https://doi.org/10.1073/pnas.1616702114.

[51]   Talla V, Suh A, Kalsoom F, Dincă V, Vila R, Friberg M, et al. Rapid increase in

genome size as a consequence of transposable element hyperactivity in wood-white

(leptidea) butterflies. Genome Biol Evol 2017;9:2491–505.

https://doi.org/10.1093/gbe/evx163.

[52]   Kidwell MG. Transposable elements and the evolution of genome size in eukaryotes.

Genetica 2002;115:49–63. https://doi.org/10.1023/A:1016072014259.

[53]   Stritt C, Gordon SP, Wicker T, Vogel JP, Roulin AC. Recent activity in expanding

populations and purifying selection have shaped transposable element landscapes

across natural accessions of themediterranean grass brachypodium distachyon.

Genome Biol Evol 2018;10:304–18. https://doi.org/10.1093/gbe/evx276.

[54]   Sultana T, Zamborlini A, Cristofari G, Lesage P. Integration site selection by

retroviruses and transposable elements in eukaryotes. Nat Rev Genet 2017;18:292–

308. https://doi.org/10.1038/nrg.2017.7.

[55]   Sawicka K, Bushell M, Spriggs KA, Willis AE. Polypyrimidine-tract-binding protein: A

multifunctional RNA-binding protein. Biochem Soc Trans 2008;36:641–7.

https://doi.org/10.1042/BST0360641.

[56]   Tree DRP, Shulman JM, Rousset R, Scott MP, Gubb D, Axelrod JD. Prickle mediates

feedback amplification to generate asymmetric planar cell polarity signaling. Cell

2002;109:371–81. https://doi.org/10.1016/S0092-8674(02)00715-8.

[57]   E L, Z Y, L F, S K, C V. A critical role of teashirt for patterning the ventral epidermis is

masked by ectopic expression of tiptop, a paralog of teashirt in Drosophila. Dev Biol

2005;283:446–58. https://doi.org/10.1016/J.YDBIO.2005.05.005.

[58]   RR D, JM L, JP K. Restriction of ectopic eye formation by Drosophila teashirt and

tiptop to the developing antenna. Dev Dyn 2009;238:2202–10.

https://doi.org/10.1002/DVDY.21927.

[59]   Contreras EG, Palominos T, Glavic Á, Brand AH, Sierralta J, Oliva C. The

transcription factor SoxD controls neuronal guidance in the Drosophila visual system.

Sci Reports 2018 81 2018;8:1–12. https://doi.org/10.1038/s41598-018-31654-5.

[60]   Matsubara D, Horiuchi SY, Shimono K, Usui T, Uemura T. The seven-pass

transmembrane cadherin Flamingo controls dendritic self-avoidance via its binding to

a LIM domain protein, Espinas, in Drosophila sensory neurons. Genes Dev

2011;25:1982–96. https://doi.org/10.1101/GAD.16531611.

[61]   Bessa J, Gebelein B, Pichaud F, Casares F, Mann RS. Combinatorial control of

Drosophila eye development by Eyeless, Homothorax, and Teashirt. Genes Dev

2002;16:2415–27. https://doi.org/10.1101/GAD.1009002.

[62] Oliveira SG, Bao W, Martins C, Jurka J. Horizontal transfers of Mariner transposons

between mammals and insects. Mob DNA 2012;3:1–6. https://doi.org/10.1186/1759-

8753-3-14.

[63] Ivancevic AM, Kortschak RD, Bertozzi T, Adelson DL. Horizontal transfer of BovB and

L1 retrotransposons in eukaryotes. Genome Biol 2018;19:1–13.

https://doi.org/10.1186/s13059-018-1456-7.

[64] Gilbert C, Schaack S, Pace JK, Brindley PJ, Feschotte C. A role for host-parasite

interactions in the horizontal transfer of transposons across phyla. Nature

2010;464:1347–50. https://doi.org/10.1038/nature08939.

[65] Wallau GL, Ortiz MF, Loreto ELS. Horizontal transposon transfer in eukarya:

Detection, bias, and perspectives. Genome Biol Evol 2012;4:689–99.

https://doi.org/10.1093/gbe/evs055.

[66] Dupeyron M, Baril T, Hayward A. Broadscale evolutionary analysis of eukaryotic DDE

transposons. BioRxiv 2021:2021.09.26.461848.

https://doi.org/10.1101/2021.09.26.461848.

[67] Prysby MD, Oberhauser KS. Temporal and Geographic Variation in Monarch

Densities: Citizen. Monarch Butterfly Biol Conserv 2004:9.

[68] Tang Z, Zhang HH, Huang K, Zhang XG, Han MJ, Zhang Z. Repeated horizontal

transfers of four DNA transposons in invertebrates and bats. Mob DNA 2015;6:1–10.

https://doi.org/10.1186/s13100-014-0033-1.

[69] Espeland M, Breinholt J, Willmott KR, Warren AD, Vila R, Toussaint EFA, et al. A

Comprehensive and Dated Phylogenomic Analysis of Butterflies. Curr Biol

2018;28:770-778.e5. https://doi.org/10.1016/J.CUB.2018.01.061.

[70] Ustyugova S V, Amosova AL, Lebedev YB, Sverdlov ED. Cell line fingerprinting using

retroelement insertion polymorphism. Biotechniques 2005;38:561–5.

[71] Lavoie CA, II RNP, Novick PA, Counterman BA, Ray DA. Transposable element

evolution in Heliconius suggests genome diversity within Lepidoptera. Mob DNA

2013;21:1–10.

[72] Blass E, Bell M, Boissinot S. Accumulation and rapid decay of non-LTR

retrotransposons in the genome of the three-spine stickleback. Genome Biol Evol

2012;4:687–702.

[73] Ivancevic AM, Kortschak RD, Bertozzi T, Adelson DL. LINEs between Species:

Evolutionary Dynamics of LINE-1 Retrotransposons across the Eukaryotic Tree of

Life. Genome Biol Evol 2016;8:3301–22. https://doi.org/10.1093/GBE/EVW243.

[74] Woronik A, Tunström K, Perry MW, Neethiraj R, Stefanescu C, Celorio-Mancera M de

la P, et al. A transposable element insertion is associated with an alternative life

history strategy. Nat Commun 2019 101 2019;10:1–11.

https://doi.org/10.1038/s41467-019-13596-2.

[75] Zhan S, Zhang W, Niitepõld K, Hsu J, Haeger JF, Zalucki MP, et al. The genetics of

monarch butterfly migration and warning colouration. Nat 2014 5147522

2014;514:317–21. https://doi.org/10.1038/nature13812.

[76] Hill T, Koseva BS, Unckless RL. The Genome of Drosophila innubila Reveals

Lineage-Specific Patterns of Selection in Immune Genes. Mol Biol Evol

2019;36:1405–17. https://doi.org/10.1093/MOLBEV/MSZ059.

[77] Sackton TB, Lazzaro BP, Schlenke TA, Evans JD, Hultmark D, Clark AG. Dynamic

evolution of the innate immune system in Drosophila. Nat Genet 2007 3912

2007;39:1461–8. https://doi.org/10.1038/ng.2007.60.

[78] Shultz AJ, Sackton TB. Immune genes are hotspots of shared positive selection

across birds and mammals. Elife 2019;8. https://doi.org/10.7554/ELIFE.41815.

[79]   Kimbrell DA, Beutler B. The evolution and genetics of innate immunity. Nat Rev Genet

2001 24 2001;2:256–67. https://doi.org/10.1038/35066006.

[80]   Lenhard B, Sandelin A, Carninci P. Metazoan promoters: Emerging characteristics

and insights into transcriptional regulation. Nat Rev Genet 2012;13:233–45.

https://doi.org/10.1038/nrg3163.

[81]   Jurka J, Kapitonov V V., Kohany O, Jurka M V. Repetitive Sequences in Complex

Genomes: Structure and Evolution. Annu Rev Genomics Hum Genet 2007;8:241–59.

https://doi.org/10.1146/annurev.genom.8.080706.092416.

[82]   Kidwell MG, Lisch DR. Transposable elements and host genome evolution. Trends

Ecol Evol 2000;15:95–9. https://doi.org/10.1016/S0169-5347(99)01817-0.

[83]   Biémont C, Vieira C. Junk DNA as an evolutionary force. Nature 2006;443:521–4.

[84]   Cosby RL, Chang N-C, Feschotte C. Host–transposon interactions: conflict,

cooperation, and cooption. Genes Dev 2019;33:1098–116.

https://doi.org/10.1101/GAD.327312.119.

[85]   Casacuberta E, González J. The impact of transposable elements in environmental

adaptation. Mol Ecol 2013;22:1503–17. https://doi.org/10.1111/mec.12170.

[86]   Huff JT, Zilberman D, Roy SW. Mechanism for DNA transposons to generate introns

on genomic scales. Nature 2016;538:533–6.

[87]   Gilbert C, Feschotte C. Horizontal acquisition of transposable elements and viral

sequences: patterns and consequences. Curr Opin Genet Dev 2018;49:15–24.

https://doi.org/10.1016/j.gde.2018.02.007.

[88]   Galbraith JD, Ludington AJ, Sanders KL, Suh A, Adelson DL. Horizontal transfer and

subsequent explosive expansion of a DNA transposon in sea kraits (Laticauda). Biol

Lett 2021;17:20210342. https://doi.org/10.1098/RSBL.2021.0342.

[89] Le Rouzic A, Capy P. The First Steps of Transposable Elements InvasionParasitic

Strategy vs. Genetic Drift. Genetics 2005;169:1033–43.

https://doi.org/10.1534/GENETICS.104.031211.

[90] Kofler R, Senti KA, Nolte V, Tobler R, Schlötterer C. Molecular dissection of a natural

transposable element invasion. Genome Res 2018;28:824–35.

https://doi.org/10.1101/gr.228627.117.

[91] Kelleher ES, Azevedo RBR, Zheng Y. The evolution of small-RNA-mediated silencing

of an invading transposable element. Genome Biol Evol 2018;10:3038–57.

https://doi.org/10.1093/gbe/evy218.

[92] Goodier JL. Restricting retrotransposons: A review. Mob DNA 2016;7.

https://doi.org/10.1186/s13100-016-0070-z.

[93] Eickbush TH, Furano A V. Fruit flies and humans respond differently to

retrotransposons. Curr Opin Genet Dev 2002. https://doi.org/10.1016/S0959-

437X(02)00359-3.

[94] Ray DA, Grimshaw JR, Halsey MK, Korstian JM, Osmanski AB, Sullivan KAM, et al.

Simultaneous TE Analysis of 19 Heliconiine Butterflies Yields Novel Insights into

Rapid TE-Based Genome Diversification and Multiple SINE Births and Deaths.

Genome Biol Evol 2019;11:2162–77. https://doi.org/10.1093/gbe/evz125.

[95] Levin HL, Moran J V. Dynamic interactions between transposable elements and their

hosts. Nat Rev Genet 2011;12:615–27. https://doi.org/10.1038/nrg3030.

[96] Fontanillas P, Hartl DL, Reuter M. Genome organization and gene expression shape

the transposable element distribution in the Drosophila melanogaster euchromatin.

PLoS Genet 2007;3:2256–67. https://doi.org/10.1371/journal.pgen.0030210.

[97] Parfrey LW, Lahr DJG, Katz LA. The dynamic nature of eukaryotic genomes. Mol Biol

Evol 2008;25:787–94. https://doi.org/10.1093/molbev/msn032.

[98] Sayers EW, Agarwala R, Bolton EE, Brister JR, Canese K, Clark K, et al. Database resources of the National Center for Biotechnology Information. Nucleic Acids Res 2014;42:D7-17. https://doi.org/10.1093/nar/gky1069.

[99] Qu Z, Nong W, Yu Y, Baril T, Yip HY, Hayward A, et al. Genome of the four-finger threadfin Eleutheronema tetradactylum (Perciforms: Polynemidae). BMC Genomics 2020;21:1–10.

[100] Qu Z, Nong W, So WL, Barton-Owen T, Li Y, Leung TCN, et al. Millipede genomes reveal unique adaptations during myriapod evolution. PLoS Biol 2020;18:e3000636.

[101] Li Y, Nong W, Baril T, Yip HY, Swale T, Hayward A, et al. Reconstruction of ancient homeobox gene linkages inferred from a new high-quality assembly of the Hong Kong oyster (Magallana hongkongensis) genome. BMC Genomics 2020;21:1–17.

[102] Nong W, Law STS, Wong AYP, Baril T, Swale T, Chu LM, et al. Chromosomal-level reference genome of the incense tree Aquilaria sinensis. Mol Ecol Resour 2020;20:971.

[103] Nong W, Qu Z, Li Y, Barton-Owen T, Wong AYP, Yip HY, et al. Horseshoe crab genomes reveal the evolution of genes and microRNAs after three rounds of whole genome duplication. Commun Biol 2021;4:1–11.

[104] da Fonseca RR, Couto A, Machado AM, Brejova B, Albertin CB, Silva F, et al. A draft genome sequence of the elusive giant squid, Architeuthis dux. Gigascience 2020;9:1–12. https://doi.org/10.1093/gigascience/giz152.

[105] Smit AFA, Hubley RR, Green PR. RepeatMasker Open-4.0. Http://RepeatmaskerOrg 2013.

[106] Jurka J, Kapitonov V V., Pavlicek A, Klonowski P, Kohany O, Walichiewicz J. Repbase Update, a database of eukaryotic repetitive elements. Cytogenet Genome

Res 2005;110:462–7. https://doi.org/10.1159/000084979.

[107] Hubley R, Finn RD, Clements J, Eddy SR, Jones TA, Bao W, et al. The Dfam

database of repetitive DNA families. Nucleic Acids Res 2016;44:D81–9.

https://doi.org/10.1093/nar/gkv1272.

[108] Smit A, Hubley R. RepeatModeler Open-1.0. Http://RepeatmaskerOrg 2015.

[109] Bao Z, Eddy SR. Automated De Novo Identification of Repeat Sequence Families in

Sequenced Genomes. Genome Res 2002;12:1269–76.

https://doi.org/10.1101/gr.88502.

[110] Price AL, Jones NC, Pevzner PA. De novo identification of repeat families in large

genomes. Bioinformatics 2005;21:i351–8.

[111] Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al.

BLAST+: Architecture and applications. BMC Bioinformatics 2009;10:1–9.

https://doi.org/10.1186/1471-2105-10-421.

[112] Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:

Improvements in performance and usability. Mol Biol Evol 2013;30:772–80.

https://doi.org/10.1093/molbev/mst010.

[113] Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: A tool for automated

alignment trimming in large-scale phylogenetic analyses. Bioinformatics

2009;25:1972–3. https://doi.org/10.1093/bioinformatics/btp348.

[114] Rice P, Longden L, Bleasby A. EMBOSS: The European Molecular Biology Open

Software Suite. Trends Genet 2000;16:276–7. https://doi.org/10.1016/S0168-

9525(00)02024-2.

[115] Xu Z, Wang H. LTR_FINDER: an efficient tool for the prediction of full-length LTR

retrotransposons. Nucleic Acids Res 2007;35:W265–8.

https://doi.org/10.1093/nar/gkm286.

[116]   Ou S, Jiang N. LTR_FINDER_parallel: parallelization of LTR_FINDER enabling rapid

1073   identification of long terminal repeat retrotransposons. BioRxiv 2019:2–6.

1074   [117]   Wong WY, Simakov O. RepeatCraft: a meta-pipeline for repetitive element de-

1075   fragmentation and annotation. Bioinformatics 2018;35:1051–2.

1076   https://doi.org/10.1093/bioinformatics/bty745.

1077   [118]   Rubino F, Creevey CJ. MGkit: Metagenomic framework for the study of microbial

1078   communities 2014.

1079   [119]   Larsson A. AliView: a fast and lightweight alignment viewer and editor for large

1080   datasets. Bioinformatics 2014;30:3276–8.

1081   [120]   Team RC. R: A language and environment for statistical computing 2013.

1082   [121]   Team Rs. RStudio: integrated development for R. RStudio, Inc, Boston, MA URL

1083   Http//Www Rstudio Com 2015.

1084   [122]   Quinlan AR, Hall IM. BEDTools: A flexible suite of utilities for comparing genomic

1085   features. Bioinformatics 2010;26:841–2. https://doi.org/10.1093/bioinformatics/btq033.

1086   [123]   Makowski D, Ben-Shachar M, Lüdecke D. bayestestR: Describing Effects and their

1087   Uncertainty, Existence and Significance within the Bayesian Framework. J Open

1088   Source Softw 2019;4:1541. https://doi.org/10.21105/joss.01541.

1089   [124]   Gel B, Serra E. karyoploteR: an R/Bioconductor package to plot customizable

1090   genomes displaying arbitrary data. Bioinformatics 2017;33:3088–90.

1091   https://doi.org/10.1093/bioinformatics/btx346.

1092   [125]   Kimura M. A simple method for estimating evolutionary rates of base substitutions

1093   through comparative studies of nucleotide sequences. J Mol Evol 1980;16:111–20.

1094   [126]   Pfeiler E, Nazario-Yepiz NO, Pérez-Gálvez F, Chávez-Mora CA, Laclette MRL,

1095   Rendón-Salinas E, et al. Population Genetics of Overwintering Monarch Butterflies,

Danaus plexippus (Linnaeus), from Central Mexico Inferred from Mitochondrial DNA

and Microsatellite Markers. J Hered 2017;108:163–75.

https://doi.org/10.1093/jhered/esw071.

[127] Wickham H. ggplot2: elegant graphics for data analysis. Springer; 2016.

[128] Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, et al. Welcome

to the Tidyverse. J Open Source Softw 2019;4:1686.

[129] Marchler-Bauer A, Bo Y, Han L, He J, Lanczycki CJ, Lu S, et al. CDD/SPARCLE:

functional classification of proteins via subfamily domain architectures. Nucleic Acids

Res 2017;45:D200–3. https://doi.org/10.1093/NAR/GKW1129.

[130] Kapitonov V V., Tempel S, Jurka J. Simple and fast classification of non-LTR

retrotransposons based on phylogeny of their RT domain protein sequences. Gene

2009;448:207. https://doi.org/10.1016/J.GENE.2009.07.019.

[131] Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol Evol

2015;32:268–74. https://doi.org/10.1093/MOLBEV/MSU300.

[132] Kalyaanamoorthy S, Minh BQ, Wong TKF, Haeseler A von, Jermiin LS. ModelFinder:

fast model selection for accurate phylogenetic estimates. Nat Methods 2017 146

2017;14:587–9. https://doi.org/10.1038/nmeth.4285.

[133] Barreiro LB, Quintana-Murci L. From evolutionary genetics to human immunology:

How selection shapes host defence genes. Nat Rev Genet 2010;11:17–30.

https://doi.org/10.1038/NRG2698.

[134] Brunham RC, Plummer FA, Stephens3 RS. Bacterial Antigenic Variation, Host

Immune Response, and Pathogen-Host Coevolution. INFECrION Immun 1993:2273–

6.

[135] Vilcinskas A. Coevolution between pathogen-derived proteinases and proteinase

inhibitors of host insects. Virulence 2010;1:206–14.

https://doi.org/10.4161/viru.1.3.12072.

[136] Yu P. Some host-parasite genetic interaction models. Theor Popul Biol 1972;3:347.

https://doi.org/10.1016/0040-5809(72)90008-1.

[137] Anderson RM, May RM. Coevolution of hosts and parasites. Parasitology

1982;85:411–26. https://doi.org/10.1017/S0031182000055360.

1127

Declarations

1129   Ethics approval and consent to participate: Not applicable

1130

1131   Consent for Publication: Not applicable

1132

1133   Availability of Data and Materials: All data generated or analysed during this study are

1134   included in this article and its supplementary information files.

1135

1136   Competing Interests: The authors declare that they have no competing interests.

1137

1138   Funding: TB was supported by a studentship from the Biotechnology and Biological

1139   Sciences Research Council-funded South West Biosciences Doctoral Training Partnership

1140   (BB/M009122/1). AH was supported by a Biotechnology and Biological Sciences

1141   Research Council (BBSRC) David Phillips Fellowship (BB/N020146/1).

1142

1143   Authors' Contributions: TB developed the pipelines and scripts for automated TE annotation

1144   and downstream analyses, performed manual curation of TEs for the monarch, produced the

1145   figures, and drafted the manuscript. AH conceived and coordinated the study and

1146   participated in writing the manuscript. All authors read and approved the final manuscript.

1147

1148   Acknowledgements: We would like to thank Dr James Galbraith, Ryan Biscocho, and Ryan

1149   Imrie for constructive comments on early drafts of this manuscript.

1150

1151

1152

1153

1154

1155

4A

4B

CR1
I-Jockey
L2
R1
R2-Hero
RTE-BovB
RTE-RTE

Scaled LINE Fragment Coordinates Compared to Consensus