# Migrators Within Migrators: Exploring Transposable Element

# Dynamics in the Monarch Butterfly, *Danaus plexippus*

Tobias Baril[1,2]* and Alexander Hayward[1,3]*

[1] Centre for Ecology and Conservation, University of Exeter, Penryn Campus,

Cornwall, TR10 9FE, UK

[2] ORCID: 0000-0002-5936-7531

[3] ORCID: 0000-0001-7413-718X

* Corresponding Author information:

Tobias Baril: tb529@exeter.ac.uk

Dr Alexander Hayward: alex.hayward@exeter.ac.uk

## Abstract

Background

Lepidoptera (butterflies and moths) are an important model system in ecology and evolution.

A high-quality chromosomal genome assembly is available for the monarch butterfly

(*Danaus plexippus*), but it lacks an in-depth transposable element (TE) annotation,

presenting an opportunity to explore monarch TE dynamics and the impact of TEs on

shaping the monarch genome.

Results

We find 6.21% of the monarch genome is comprised of TEs, a reduction of 6.85% compared

to the original TE annotation performed on the draft genome assembly. Monarch TE content

is low compared to two closely related species with available genomes, *Danaus chrysippus*

(33.97% TE) and *Danaus melanippus* (11.87% TE). The biggest TE contributions to genome

size in the monarch are LINEs and *Penelope*-like elements, and three newly identified

families, *r2-hero_dPle* (LINE), *penelope-1_dPle* (*Penelope*-like), and *hase2-1_dPle* (SINE),

collectively contribute 34.92% of total TE content. We find evidence of recent TE activity,

with two novel Tc1 families rapidly expanding over recent timescales (*tc1-1_dPle*, *tc1-*

*2_dPle*). LINE fragments show signatures of genomic deletions indicating a high rate of TE

turnover. We investigate associations between TEs and wing colouration and immune genes

and identify a three-fold increase in TE content around immune genes compared to other

host genes.

Conclusions

We provide a detailed TE annotation and analysis for the monarch genome, revealing a

considerably smaller TE contribution to genome content compared to two closely related

*Danaus* species with available genome assemblies. We identify highly successful novel DNA

TE families rapidly expanding over recent timescales, and ongoing signatures of both TE

expansion and removal highlight the dynamic nature of repeat content in the monarch

genome. Our findings also suggest that insect immune genes are promising candidates for

future interrogation of TE-mediated host adaptation.

## Keywords

52    Transposon; TE Annotation; Lepidoptera; *Danaus plexippus*; Butterfly; Genomic Deletion;

53    Repeat; Genome Evolution

## Background

Transposable elements (TEs) are autonomous DNA sequences that can move within the

genome. TEs are present in nearly all eukaryotes and are important in shaping their

genomes[1–8]. TEs also represent a large reservoir of sequences with the potential to

contribute to host genomic novelty, and they have been implicated in the evolution of

regulatory networks [3,9–13], chromosomal rearrangements [3,14–19], exon shuffling [20–

22], and donation of coding sequence [20,22–27].

62           Due to the great diversity and dynamic nature of TEs, species-specific TE libraries

and accompanying TE annotations are required to provide an understanding of their

evolution and impact. By examining differences in the diversity and abundance of TEs within

a genome, as well as their integration histories, proliferation frequencies, and rate of

removal, we can begin to unravel the factors that govern TE evolutionary dynamics. Curating

TEs in a wide range of species is also an important step more generally, to expand

understanding of variation in the interplay between TEs and host genomes, and the

consequences of this for host evolution [28]. Meanwhile, from a practical perspective,

accurate and reliable repeat annotation is essential to provide high-quality genome

annotations, and to help avoid repeat sequences being incorrectly annotated as host genes

and vice versa, especially given the vast numbers of genomes currently being sequenced

across the tree of life.

Lepidoptera (butterflies and moths) are an important model system in ecology and

evolution. Lepidopterans have captivated researchers for decades, proving a fascinating and

practical system for research into topics such as development, physiology, host-plant

interactions, coevolution, phylogenetics, mimicry, and speciation (e.g. [29–33]). Over recent

decades, Lepidoptera has become an important model system in genomics especially. For

example, Lepidoptera genomes have been pivotal in investigations into the genomic bases

of migration [34–36], warning colouration [35], adaptive radiations [33], and evolutionary

studies into TEs, including TE horizontal transfer [37–39], the influence of TEs on genome

size [40], and the role of a TE in the evolution of industrial melanism in the peppered moth

[41].

The monarch butterfly, *Danaus plexippus*, is famous for its incredible North American

migration, global dispersal, and striking orange warning colouration [35], and has been the

focus of much research and broad public interest for decades. A draft monarch genome was

released in 2011, and work during the last decade has focussed on revealing the genetic

bases of several key traits [35,36]. Most recently, a high-quality chromosomal assembly of

the monarch genome was generated [42]. Additionally, genome assemblies for the black

veined tiger (*D. melanippus*) and African Queen (*D. chrysippus*) provide a comparative

context to examine monarch genomics. Yet, despite being an important research focus, we

lack a detailed TE annotation for the monarch, and an accompanying understanding of

monarch TE dynamics and host-TE interactions.

Here, we provide a thorough annotation of TEs within the monarch genome,

alongside analyses of TE diversity and abundance, interactions with host genes, genomic

distribution including hot and cold spots, and TE evolution. We find evidence of considerable

recent TE activity in the monarch, with two novel Tc1 element families rapidly expanding

over recent timescales (*tc1-1_dPle* and *tc1-2_dPle*). In contrast, we find that historically a

pattern of high proliferation rates of LINE and *Penelope-like* elements (PLEs) was dominant,

with these elements still dominating the monarch TE landscape. We also reveal the

considerable role of genomic deletions in removing LINE TEs from the genome. Overall, we

provide an overview of the broadscale dynamics that have shaped the TE landscape of the

monarch genome, and provide a high-quality, curated, and species-specific TE annotation

library for use by the scientific community.

## Results

Transposable Element Landscape

Repeats are a major determinant of genome size [40,43–47]. However, in the

monarch, we find that repeats comprise just 6.21% of the genome (total assembly size =

248.68Mb, assembly scaffold N50 = 9,210kb, 97.5% Complete Busco orthologs) (Figure 1,

Table 1). This differs considerably in comparison to the TE annotation performed on the

original monarch genome assembly, where repeats were estimated to cover 13.06% of the

genome (Table S1; Additional File 1) [36]. When examining the source of this variation, the

major difference in TE content can be attributed to unclassified elements, for which we find

coverage is reduced by 16.31Mb (6.6% of total genome size) (Table S1; Additional File 1).

This difference is likely due to two issues: (i) improvements in TE annotation tools over the

last decade, where improved knowledge of TE structure has led to exclusion of erroneous

sequences previously annotated as putatively TE, particularly given that the original

annotation used similar tools and databases to those applied in this study [36]; (ii) We apply

a conservative approach to TE annotation that excludes very short fragments which cannot

be confidently identified as TE sequence, and we remove overlapping TE annotations.

Improvements in genome assembly quality will also have improved repeat detection.

Consistent with this, a standard 'no frills' RepeatMasker run using the Arthropoda RepBase

database [48] and the Dfam database of repetitive DNA families [49] identified a repeat

content of 5.60%, which is much closer to our finding of 6.21% repeat content using the

earlGrey pipeline (https://github.com/TobyBaril/EarlGrey)[50], and much lower than that

identified in the original TE annotation (Table S2; Additional File 1). Thus, many of the

sequences annotated as unclassified in the original annotation were most likely not TE

sequences and are no longer annotated as such by current annotation tools. We are,

however, unable to confirm this for certain since the original *de novo* repeat library

generated for the draft genome was not provided alongside the genome. We also

acknowledge the considerable difference in TE content annotated in the monarch genome in

this study compared to a recent study focussing on *D. chrysippus* which also annotated TEs

in the monarch genome [51]. When examining the source of this variation, we find this can

be attributed to the more conservative and detailed approach taken here, where for example,

95,956 extremely short annotations identified in [51] did not reach the length and score

thresholds for annotation in this study, whilst we also used a monarch-specific repeat library

for annotation (See Additional File 2 for full discussion).

We annotated the two additional genomes available in the genus *Danaus* using the

same automated TE annotation method that we applied for the monarch. In comparison, the

monarch exhibits a much lower TE content than the other two species, with repetitive

elements covering 33.97% of the *D. chrysippus* genome (assembly size = 354.02Mb, contig

N50 = 11.45Mb, 98.2% complete Busco orthologs), and 11.87% of the *D. melanippus*

genome (assembly size = 354.18Mb, scaffold N50 = 890kb, 86.0% complete Busco

orthologs) (Figure 1, Table 1). Relatively recent TE activity is indicated in all three *Danaus*

species, with several TE classifications exhibiting low genetic distance to their respective

consensus sequences, but there is evidence of a much greater burst of recent TE activity in

*D. chrysippus* (Figure 1). The genome size of both *D. chrysippus* and *D. melanippus* is

354Mb, whilst the monarch genome is 30% smaller at only 248Mb. Increases in TE content

are apparent for all TE classifications for *D. chrysippus* and *D. melanippus* compared to the

monarch, except for a decrease in PLE and SINE coverage in *D. melanippus* relative to the

monarch (Table 1). The most populous TEs in *D. chrysippus* are rolling circle TEs (8.71% of

genome, 30.8Mb), whilst LINEs dominate in *D. melanippus* (4.58% of genome, 16.2Mb)

(Table 1).

| Repeat Class | *Danaus plexippus (GCA_009731565.1)* | | | *Danaus chrysippus (GCA_916720795.1)* | | | *Danaus melanippus (GCA_010014825.1)* | | |
|---|---|---|---|---|---|---|---|---|---|
| | No. elements | Total Length (Mb) | Percentage sequence (%) | No. elements | Total Length (Mb) | Percentage sequence (%) | No. elements | Total Length (Mb) | Percentage sequence (%) |
| **Retroelement** | **29181** | **9.55** | **3.84%** | **91778** | **48.09** | **13.58%** | **65710** | **23.35** | **6.59%** |
| *SINE* | 3421 | 0.85 | 0.34% | 4973 | 1.83 | 0.52% | 33 | 0.01 | 0.00% |
| *LINE* | 10227 | 4.83 | 1.94% | 55533 | 28.47 | 8.04% | 41769 | 16.21 | 4.58% |
| *Penelope* | 10478 | 2.55 | 1.03% | 23263 | 9.69 | 2.74% | 6211 | 1.67 | 0.47% |
| *LTR element* | 5055 | 1.32 | 0.53% | 8009 | 8.09 | 2.28% | 17697 | 5.46 | 1.54% |
| **DNA transposon** | 3723 | 2.16 | 0.87% | 37897 | 22.87 | 6.46% | 19187 | 7.32 | 2.07% |
| **Rolling-circle** | 3659 | 1.50 | 0.60% | 56659 | 30.83 | 8.71% | 7484 | 2.41 | 0.68% |
| **Unclassified** | 6949 | 2.08 | 0.84% | 26595 | 1.83 | 5.22% | 28069 | 8.51 | 2.40% |
| **Other** | 362 | 0.15 | 0.06% | 13 | 0.00 | 0.00% | 2181 | 0.46 | 0.13% |
| **Total repeats** | 43874 | 15.44 | 6.21% | 212942 | 103.62 | 33.97% | 122631 | 42.05 | 11.87% |

Table 1. Transposable element content for the three *Danaus* species with available genome

assemblies.

The lower TE content observed in the monarch genome presumably arose by either: (i) loss

of repeat content in the monarch; (ii) independent gains of repeat content in both *D.*

*chrysippus* and *D. melanippus* (or in their ancestral lineages); or (iii) a gain of repeat content

in the ancestral lineage leading to *D. chrysippus* and *D. melanippus* (Figure 2A). The

ancestor of the monarch and *D. chrysippus* is estimated to have diverged ~8.3Mya [52].

Thus, using this estimate we were able to calculate DNA loss rates for the two species, to

examine if the smaller genome of the monarch is a result of an increased rate of DNA loss

(Table S3; Additional File 1). The estimated rates of micro- (<30bp) and mid-size (30bp to

10,000bp) deletions in the monarch and *D. chrysippus* are comparable. However, the

monarch exhibits similar DNA loss rates for both micro- and mid-size deletions, whilst *D.*

*chrysippus* has a reduced rate of DNA loss from mid-size deletions offset by an increased

rate of DNA loss from micro-deletions (monarch $\mu$=21.27 del_nt/10kb/My, micro-

deletions=20.57 del_nt/10kb/My, mid-size deletions=21.96 del_nt/10kb/My; *D. chrysippus*

$\mu$=20.61 del_nt/10kb/My, micro-deletions=26.61 del_nt/10kb/My, mid-size deletions=14.62

del_nt/10kb/My). This suggests observed differences in the quantity of TE sequence

between the monarch and *D. chrysippus* are due to increases in TE content in *D.*

*chrysippus,* rather than higher rates of DNA loss in the monarch. We were unable to

calculate comparative DNA loss rates for *D. melanippus*, since a divergence estimate was

not available for this species. However, mean estimated DNA loss was reduced in *D.*

*melanippus* compared to the other two species (*D. melanippus*=158.19 del_nt/10kb,

monarch=176.50 del_nt/10kb, *D. chrysippus*=171.11 del_nt/10kb). *D. melanippus* exhibited

a similar level of DNA loss from micro-deletions to the monarch, but a reduction compared to

*D. chrysippus*, while *D. melanippus* and *D. chrysippus* exhibit a reduction in mid-size

deletions compared to the monarch (Table S3; Additional File 1).  Meanwhile, considerable

expansions in species-specific TE families are evident in *D. chrysippus* and *D. melanippus*

(Figure 2B & C). Therefore, we suggest that hypothesis (ii) is the most likely explanation for

observed differences in TE load among *Danaus* species, whereby *D. chrysippus* and *D.*

*melanippus* have both experienced independent expansions in TE content (Figure 2).

Factors that may have been responsible for these gains, particularly the large gain in TE

content in the genome of *D. chrysippus*, are unclear.

Transposable Element Landscape

TEs are unevenly distributed across the monarch genome, as indicated by a high

standard deviation in TE base pairs per 100kb genomic window (mean = 6.1kb/100kb, SD =

4.1kb/100kb), whereby a standard deviation of 0 indicates TEs are evenly distributed across

the genome (Figure 3, Figure 4, Table S4; Additional File 1). TEs were also unevenly

distributed among different chromosomes (Kruskal-Wallis, $\chi2_{29}$=286.43, p<0.01). As

expected, a strong positive correlation between TE coverage and chromosome size was

observed (Pearson's Correlation, r = 0.76, $t_{28}$ = 6.2604, p < 0.01). The highest density of TEs

is present on chromosome 28 (10.1kb/100kb, chromosome size = 3.99Mb), whilst

chromosome 9 exhibits the lowest TE density, with less than half that observed for

chromosome 28 (4.5kb/100kb, chromosome size = 9.62Mb) (Table S4; Additional File 1).

Similarly, TEs are unevenly distributed across the genome of *D. chrysippus* (mean =

34.0kb/100kb, SD = 28.5kb/100kb) (Table S4; Additional File 1), which also has a

chromosomal-level assembly. Unlike the monarch, a weaker correlation between TE

coverage and chromosome size was observed (Pearson's Correlation, r = 0.43, $t_{28}$ = 2.4974,

p < 0.05). The highest density of TEs is found on chromosome 29 (69.8kb/100kb,

chromosome size = 10.47Mb), whilst chromosome 1 exhibits the lowest TE density, with a

nearly 6-fold decrease compared to chromosome 29 (12.6kb/100kb, chromosome size =

16.60Mb. In both the monarch and *D. chrysippus,* there is a significant negative correlation

between chromosome size and TE density, indicating that a larger proportion of small

chromosomes is composed of TE sequence (Pearson's Correlation, monarch: r = -0.70, $t_{28}$ =

5.1335, p < 0.01, *D. chrysippus*: r = -0.43, $t_{28}$ = 2.5395, p < 0.05). We were unable to

212 compare TE density in *D. melanippus* as a chromosomal-level genome assembly is not

213 available for this species.

214  TEs are known to be unevenly distributed between different genomic compartments,

215 with TE densities varying between genic and intergenic space [1]. In the monarch genome,

216 we find that the greatest amount of TE sequence is found in gene flanking regions (6.59Mb),

217 followed by introns (5.56MB), intergenic regions (2.82Mb) and, finally, exons (0.47Mb)

218 (Figure 3A). While when considered as a proportion of genomic compartment size,

219 intergenic regions have the greatest proportion of TE content (9.67%), followed by gene

220 flanking regions (8.22%), introns (5.00%), and exons (1.64%) (Figure 3B). The relative lack

221 of TEs in exonic regions is expected due to the potential for deleterious effects arising from

222 transposition into coding sequence [1,53]. LINEs and PLEs are the most populous TEs

223 present in the monarch genome. LINEs comprise 1.94% of the monarch genome,

224 accounting for 3.36% of intergenic sequences, 2.67% of gene flanking regions, 1.41% of

225 intronic sequences, and 0.76% of exonic sequences (Figure 3B). Meanwhile, PLEs make up

226 1.03% of the monarch genome, accounting for  1.63% of intergenic sequences, 1.26% of

227 gene flanking regions, 0.94% of intronic sequences, and 0.16% of exonic sequences (Figure

228 3B).

229  Overall, we observed a weak negative correlation between gene density and TE

230 density in the monarch genome (Spearman's rank, rho = -0.373, p < 0.01), suggesting that

231 regions that contain a high density of host genes are less likely to contain TEs. We identified

232 25 hotspots of TE accumulation in the monarch genome, defined as 100kb windows with a

233 TE content above a 99% area under the curve (AUC) cut-off (see Methods). This equates to

234 windows where TE density is at least 3.03 times higher than expected (i.e., expected TE

235 density assuming an even spread of TEs across 100kb windows = 6.15kb, lowest hotspot

236 density = 18.64kb, highest hotspot density = 28.02kb) (Figure 5). Hotspot windows represent

237 regions of the genome able to tolerate high densities of TE insertions. Conversely, the

238 presence of 36 cold spots, defined as 100kb windows with TE content below a 1% AUC cut-

off, where TE density is at least 12.42 times lower than expected (lowest cold spot density =

0.11kb, highest cold spot density = 0.5kb), highlights regions of the genome where very few

TE insertions are tolerated. Whilst numerous TE coldspots are found on the Z chromosome,

no TE hotspots are found there (Figure 5). Of note, we detected 6 separate 100kb windows

labelled as TE coldspots and gene hotspots (Figure 5). Gene ontology analyses suggest that

genes present in gene hotspots that are TE coldspots are involved in RNA binding, splicing,

polyadenylation, mRNA stability, and translation initiation [54]; cell polarisation [55]; neuronal

development; antenna and eye development [56–60]; and olfactory processes, which are

hypothesised to be involved in social behaviour [36] (Table S5; Additional File 1). Given the

importance of these functions, it is likely that TE insertions in these regions would be

severely detrimental to host fitness, thus individuals where TEs insert into these regions may

be non-viable and purged early in development.

Transposable Element Activity

The genome of *D. plexippus* is dominated by three highly successful novel TE

families, which collectively comprise 34.9% of total TE content (Table 2, Table S6; Additional

File 1). Of these families, the consensus sequence of the PLE, *penelope-1_dple*, is intact

and putatively transposition competent, containing both a GIY-YIG endonuclease domain

and a reverse transcriptase-like (RT-like) domain, whilst the LINE element, *r2-hero-1_dple*,

is putatively transposition-competent, with an intact RT-like domain (Table S7, Additional File

1). SINEs are nonautonomous elements containing an internal polymerase III promoter to

facilitate their transposition [61]. As such, there are no internal coding regions to determine

the transposition competency of this element, however *hase2-1_dple* is putatively full-length

at 306bp.

| TE Family | TE Classification | Weighted Average Age (Mya) | Copy Number |
|---|---|---|---|
| penelope-1_dple | Penelope-like Element | 10.74 | 9901 |

| hase2-1_dple | SINE/HaSE2 | 6.39 | 2865 |
| r2-hero-1_dple | LINE/R2-Hero | 8.58 | 5432 |

Table 2. Weighted average age (TE age accounting for copy number at each divergence)

and copy number of the three dominant TE families responsible for 34.9% of total TE content

in the monarch.

Only three TE families are estimated to have inserted <1Mya in the monarch genome

(Table S8, Figure S1; Additional Files 1 and 3). The low estimated age of these insertions is

a sign of recent host colonisation, potentially via horizontal transposon transfer (HTT) from

another host genome [1,62–65]. Peaks of low genetic distance to TE consensus sequence

indicate an abundance of closely related elements and thus recent TE proliferation (Figure 1,

Table S9 and Figure S2; Additional Files 1 and 4). There has been a recent burst of activity

in two Tc1 elements: *tc1-1_dple* (418 copies, from ~0.43Mya), and *tc1-2_dple* (216 copies,

from ~0.35Mya) (Table S8; Additional File 1). The rapid expansion of these Tc1 elements

over such a short time period highlights their great success within the monarch genome

when compared to other relatively young TEs, which are found at much lower copy numbers

(Figure 6 and Figure 7, Table S8 and Table S9; Additional File 1). This recent expansion of

DNA TEs is interesting given the historic dominance of LINEs in the monarch genome

(1.94% of the genome, 31% of total TE content). Whilst there are more DNA TE families

than LINE families in the monarch genome (319 vs 213), LINEs have been more successful

in proliferating and colonising the genome than their DNA TE counterparts to date, although

the reasons for this are unknown, it appears that these new DNA TE families may challenge

the historic trend (Figure 6 and Figure 7).

The oldest TE insertions present in the monarch genome were estimated to have

occurred ~43 Mya, predating the emergence of the monarch, which diverged ~32 Mya [66],

with the first invaders being DNA and LTR elements (Figure 6, Figure 7, Table S9, Figure

S1; Additional Files 1 and 3). Thus, these TEs appear to have proliferated in an ancestral

lineage after the Nymphalidae family emerged, ~91Mya (CI: 71-112Mya) [66]. However,

most TE families (79%) are estimated to have inserted after the emergence of the monarch

(mean TE family age = 25.41Mya) (Figure 6, Figure 7)

Turnover of LINEs and Penelope-like Elements

LINEs and PLEs are reverse transcribed from their 3' end during insertion of a new

copy and are often incomplete at the 5' end due to the premature dissociation of reverse

transcriptase (RT), or the activity of cellular RNases [67,68]. In contrast, the absence of a

complete 3' region is unlikely to arise due to either process and is evidence of a genomic

deletion, presumably as a consequence of ectopic recombination between homologous LINE

or PLE sequences [67,68].

Consistent with observations in the postman butterfly *Heliconius melpomene* [68]

also from the family Nymphalidae, we find LINE and PLE fragments across all LINE and PLE

families exhibit patterns suggestive of genomic deletions acting to remove these elements

from the genome (Table 3, Figure S3; Additional File 5). In total, just under half of all

LINE/PLE insertions (49.37%) are truncated at the 3' end. For the *CR1*, *I-Jockey*, *RTE-*

*BovB*, and *Penelope* families, most fragments display patterns suggestive of removal via

genomic deletions (i.e. truncated at the 3' end), whilst the majority of fragments of the *L2,*

*R1*, *R2-Hero*, and *RTE-RTE* families display patterns indicative of processes associated with

degradation of LINEs during mobilisation, such as premature disassociation of RT or

breakdown via cellular RNAse activity (i.e. truncated at the 5' end) (Table 3). The range of

insertion times between LINEs exhibiting an abundance of 3' fragments and those exhibiting

an abundance of 5' fragments overlap, suggesting that insertion time does not explain the

inter-family differences in fragmentation (Table S9, Additional File 1). Swathes of small

fragments across all LINE and PLE families (mean length = 356bp, n = 20,706) suggest a

high rate of turnover of these non-LTR elements within the monarch genome, in contrast to

patterns observed in mammals whereby a low rate of turnover leads to an accumulation of

ageing TEs [68,69].

| LINE Classification | Fragment Types | Count | Proportion of Total Elements |
|---|---|---|---|
| CR1 | 3' | 7 | 0.13 |
| | 5' | 48 | 0.87 |
| I-Jockey | 3' | 64 | 0.41 |
| | 5' | 92 | 0.59 |
| L2 | 3' | 138 | 0.54 |
| | 5' | 119 | 0.46 |
| R1 | 3' | 266 | 0.76 |
| | 5' | 85 | 0.24 |
| R2-Hero | 3' | 2971 | 0.85 |
| | 5' | 521 | 0.15 |
| RTE-BovB | 3' | 201 | 0.47 |
| | 5' | 220 | 0.53 |
| RTE-RTE | 3' | 138 | 0.67 |
| | 5' | 68 | 0.33 |
| LINE (Unknown Family) | 3' | 2 | 0.06 |
| | 5' | 34 | 0.94 |
| Penelope | Lone 3', Internal + 3' | 537 | 0.15 |
| | Lone 5', 5' + Internal, Internal Lacking 3' | 3029 | 0.85 |

Table 3. Number of LINEs and PLEs classified by fragment type, and putative removal

process. 3' encompasses element fragments containing a lone 3' end, or those with 3' ends

and internal regions, and indicates premature RT dissociation or activity of cellular RNAses.

5' encompasses element fragments containing a lone 5' end, or those with 5' ends and

internal regions, and indicates genomic deletions.

Despite evidence of loss, LINE and PLE sequences still dominate the TE landscape

of the monarch. To be autonomous, an intact LINE must contain start and stop codons and

an RT domain [70], whilst an intact PLE must contain an RT domain and a GIY-YIG

endonuclease domain [71]. In total, we identified five intact LINEs and two intact PLEs in the

monarch genome (Table S7, Additional File 1), including one copy of *r2-Hero-1_dPle* and 2

copies of *penelope-1_dPle*, which are the two most populous TE families in the monarch

genome, suggesting continued activity (Table 2). We identified 4 intact LINEs in *D.*

*melanippus* and 9 in *D. chrysippus*, however in both species intact PLEs were absent. We

also observed accumulations of certain LINE families in the monarch, however only 11.38%

of LINEs and 3.51% of PLEs have maintained a length above 1kb, falling short of the

expected length given an intact LINE ORF encoding RT is ~3kb [72] (Table S10; Additional

File 1). Short elements are more likely to persist in the genome than their longer

counterparts, as they are less prone to recombination [19]. This, combined with the high

activity level of LINEs and PLEs before and throughout the evolution of the monarch (Figure

3), explains the dominance of LINE and PLE fragments across the genomic landscape,

despite their high rates of turnover [19,68].

Association of TEs with Monarch Colouration and Immune Genes

Given a previous role for TEs in wing colouration in the peppered moth (*Biston*

*betularia*) [41] and the clouded yellow butterfly (*Colias croceus*) [73], we investigated the

possibility of TE associations with wing colouration in the monarch. The myosin gene is

associated with wing colouration in the monarch [35], and we identified a single unclassified

element 7kb upstream of this gene (Figure S4; Additional File 6). This location contrasts to

that of TEs implicated in colouration in the peppered moth and clouded yellow, where the TE

insertions were found within an intron, and 6kb downstream of the gene body, respectively

[41,73]. When compared to all monarch genes, the association of a single TE with the

myosin gene is within the expected range to find within 20kb of a genic locus (See Methods),

and therefore is not evidence of TE enrichment (Figure S5, Additional File 7).

Immune genes are among the fastest evolving metazoan genes given strong

selection arising from exposure to harmful pathogens and parasites [74–77], and so are key

candidates for exploring the role of TEs in host evolution. Across the monarch genome, TEs

of all major types were found within immune gene flanks (20kb either side of gene bodies).

Regions surrounding immune genes contain a three-fold increase in TE content when

compared to genes involved in other functions, although this level of enrichment is just

above a 5% level of significance (Welch's T-Test, $T_{9.0915} = 2.1351$, $p = 0.061$). However, a

significant difference in TE abundance between different types of immune genes was

apparent (Kruskal-Wallis $\chi^2_{38} = 104.88$, $p < 0.01$), suggesting an enrichment for TEs around

Class II-associated invariant chain peptide (CLIP)-like modulation genes when compared to

immune genes involved in IKK gamma, Tab2, SOCS, and IMD signalling. (Figure S6;

Additional File 8).

When comparing associations between major TE types around immune genes, we

find that LINEs and PLEs are significantly more abundant than rolling circles (Kruskal-Wallis,

$\chi^2_7 = 29.88$, $p < 0.01$), but there were no significant pairwise differences between the

remaining major TE types. However, LINEs and PLEs are not enriched around immune

genes compared to genome-wide levels (LINEs: genome-wide = 1.94% of genome size,

immune gene regions = 2.04% of immune gene compartment size; PLEs: genome-wide =

1.03% of genome size, immune gene regions = 0.96% of immune gene compartment size).

## Discussion

We provide a detailed annotation of TEs in the monarch genome, an analysis of their

diversity and evolution, an investigation of their influence in shaping the monarch genome,

and a comparative overview of TE load for the monarch and two congeneric species. We

find a large reduction in TE sequence compared to the original annotation performed on the

draft monarch genome. We suggest that this is due to many sequences in the original

annotation no longer being recognised as TEs when using updated TE databases and

current annotation tools. However, we are unable to confirm this, due to the original *de novo*

TE library and annotation results being unavailable. This raises an important issue in

genomics currently. Given TE annotation is a key step in genome assembly and annotation,

we suggest that it should be required to include TE libraries alongside the publication of

genome projects, ideally in a specific repository, to: (i) reduce duplication of efforts required

for reannotation; (ii) improve reproducibility in annotation; and (iii) improve TE consensus

libraries to reduce cases of a single TE family being given different names in multiple

species. As the number of sequenced genomes increases, the wider availability of TE

libraries will benefit a wide range of studies, where gene annotations can be improved by

accurate TE annotation. The availability of large numbers of well-curated TE libraries will

also facilitate large-scale studies of TE evolutionary dynamics, and the impacts of TEs on

specific traits of interest, and host evolution more generally.

We identify an uneven distribution of TEs across the genome. TE insertions in non-

coding regions are less likely to have a detrimental impact on host genome function than TE

insertions into coding regions [1,21,53,78–80], and so are less stringently removed by

selection, enabling TE accumulation in non-coding regions. Indeed, some TEs actively avoid

coding regions by targeting regions of the genome unlikely to impact host fitness [53,81],

with this process potentially occurring in the monarch, where we find overlapping gene

hotspots and TE cold spots. TE abundance is highest in intergenic regions. Of note, the

highest TE coverage is found in host gene flanking regions rather than intergenic regions

distal from host genes, where the risk of detrimental impacts of TE insertion are thought to

be reduced [53,81]. The finding is interesting given the potential involvement of TEs in the

evolution of gene regulatory networks [10–13,82] and the uneven distribution of TEs across

the monarch genome. Although this could also be due to the relatively compact nature of the

monarch genome, where genic regions account for 56% of the genome and host gene flanks

account for a further 32%, leaving only 12% of the genome distal from host genes (i.e

intergenic). TEs are implicated in rapid increases in intron size in eukaryotic genomes,

where short non-autonomous DNA TEs independently generate hundreds to thousands of

introns [1,53,83]. It is also hypothesised that TE-derived introns are responsible for the

prevalence of nucleosome-sized exons (DNA sequences ~150bp that can wrap around a

single histone core) observed in eukaryotes [83]. Thus, the presence of TEs in intronic

sequences in the monarch genome could reflect the status of introns as refugia where TE

insertions are not as likely to be purged, or alternatively it could indicate ongoing intron

generation, with TE insertions leading to the generation of new, or longer, introns over

evolutionary time.

The presence of several young high copy number TE families in the monarch

genome demonstrates the success of TEs in naïve host genomes [81,84–87].  The putative

recent arrival of two novel Tc1 families (*tc1-1_dple* and *tc1-2_dple*), has led to a wave of TE

proliferation in the monarch genome. These elements are closely related and located in

Clade 130 of Tc1/mariner phylogeny, which contains elements from diverse insect species,

including butterflies, aphids and ants, in a recent phylogenetic study of DDE TE diversity [88]

(Figure S7; Additional File 9). Ants and aphids are known to share environments with

monarch larvae on milkweed plants [89], raising the potential for HTT events between these

groups [64,90]. Elucidating the origin of recent Tc1 insertions in the monarch genome will

require a fuller comparison across related butterfly species and other host plants and insects

that are known to share environments with the monarch, as more genomes become

available. Over time, it is hypothesised that the activity of the novel families identified will

decrease as the host develops mechanisms to suppress activity, either through the capture

of TE sequences in piRNA regions, or through other defensive mechanisms such as

methylation [87,91,92]. A reduction in transposition of DDE transposase-containing DNA

transposons can also be attributed to increases in non-autonomous copies [88]. In the

meantime, however, these TEs are proliferating and dynamically altering the genomic

landscape of *D. plexippus*, with potential consequences for ongoing genome structure and

functional evolution [1–3,19].

Factors leading to the relatively low TE content observed in the monarch compared

to other *Danaus* species are currently unknown. The maintenance of a small genome in the

monarch may be partly attributed to the high turnover of TEs, evidenced by a lack of ancient

peaks in repeat landscapes and the presence of few intact LINEs and PLEs, many of which

show hallmarks of genomic deletions. We did not observe large differences in the number of

intact LINEs between the three *Danaus* species considered here, however both *D.*

*chrysippus* and *D. melanippus* lack transposition competent PLEs. In *D. melanippus*, there

are only small numbers of intact LINEs and a relative lack of recent LINE activity, despite

historical LINE activity having made the largest TE contribution to genome size (Figure 1 and

Figure 2). In contrast, there is evidence of very recent LINE activity in *D. chrysippus*,

although much very recent TE activity is attributed to DNA and rolling circle elements.

(Figure 1 and Figure 2).

Ectopic recombination between similar elements is likely to play a significant role in

the removal of LINEs and PLEs. The only LINE or PLE families with an abundance of 3'

fragments are: *L2*, *R1*, *R2-Hero*, and *RTE-RTE*. While it is tempting to suggest that L2, *R1*,

*R2-Hero*, and *RTE-RTE* insertions may have a lower impact on host fitness than other LINE

insertions, and so are "allowed" to degrade over time, rather than being removed *en masse*

through genomic mechanisms, it should be noted that ectopic recombination is a passive

process. However, removal via ectopic recombination does not account for inter-family

differences in the ratio of fragments undergoing degradation through ectopic recombination

versus degradation via the activity of RT or cellular RNAses. In the case of the most

abundant family, *R2-Hero*, these elements may have so far avoided ectopic recombination,

given these elements are relatively young compared to others in the same classification

(insertion ~8.58Mya, mean LINE insertion time 28.51Mya ) (Figure 6 and Figure 7, Table S9;

Additional File 1)). Our results are consistent with those of previous findings in *Heliconius*

species [93], in which a high TE turnover and a relative lack of more ancient TEs were

observed, suggesting these characteristics might be widespread among nymphalids or

lepidopterans more generally, although further comparative studies will be required to

confirm this. In addition to TE turnover, TE-driven increases in genome size are likely also

limited through inhibition of TE activity via epigenetic mechanisms and piRNAs [1,91,94,95].

Further research to characterise piRNA clusters in the monarch, and interrogation of its

transcriptome, will further improve our understanding of how these processes have shaped

its TE landscape.

The lack of TE hotspots on the Z chromosome is perhaps unexpected given sex

chromosomes are thought to accumulate TEs faster than autosomes due to suppressed

homologous recombination [96–98]. If this was the case in the monarch, one might expect to

observe TEs accumulating due to the inability of the host genome to purge insertions. The Z

chromosome of the monarch is the product of a fusion between an ancestral-Z and neo-Z

genome segment, with distinct modes of dosage compensation acting on either section [42].

The unique chromatin remodelling dynamics on the Z chromosome in the monarch might

moderate TE access to regions of the Z chromosome, although further research is required

to understand the interplay between host processes and TE dynamics on this unique sex

chromosome. Meanwhile, the three-fold increase in TE abundance around immune genes is

intriguing and warrants future research to further understand the impact of TEs in these

regions. The association with CLIP modulation genes is particularly interesting, as CLIP

peptides block class II MHC binding grooves to modulate the immune response by

influencing antigen presentation to CD4 T cells by class II MHC molecules [99]. Class II

MHC genes are diverse and evolve rapidly, maintained by strong positive selection and

frequent gene conversion [100,101]. Given the strong selection these loci are under, these

are ideal candidates for further study to investigate a potential role for TEs in the adaptive

immune system. The use of TE knockout studies in combination with transcriptomic

analyses may elucidate the potential impacts TEs have on immune gene evolution in the

monarch.

## 488 Conclusions

In-depth studies into TE landscapes and dynamics within host species of broad

scientific interest remain relatively few, even with widespread acknowledgement of the

important contributions that TEs have made to eukaryotic evolution [1,43,102]. We provide a

detailed annotation of TEs in the monarch genome and analysis of TE diversity and

evolution, including the identification of highly successful young DNA TE families, which

mirror the activity profile of the most successful LINEs present in the genome. We provide a

comparative context with two other *Danaus* species, demonstrating the considerable

differences in TE content that can occur even among congeneric species. We present

evidence of ongoing TE expansion and removal, highlighting the dynamic nature of repeat

content within genomes over time. We also identify an increase in TE abundance around

genes of immune function, providing potential candidates for future studies to elucidate the

impact of TEs on immune function in the monarch. Further large-scale comparative studies

will be greatly beneficial to further our understanding of the evolutionary dynamics of TEs,

and the drivers leading to diverse TE landscapes across lineages.

## Materials and Methods

### Transposable Element Annotation

The monarch chromosomal assembly, Dplex_v4, was downloaded from NCBI (Refseq GCF_009731565.1), along with corresponding genome annotations in GFF format [42,103]. The monarch genome assembly is 248.676Mb, with a GC content of 32.1%, and comprises 4,115 scaffolds with a scaffold N50 of 9.21Mb.

The *D. plexippus* genome was annotated using Earl Grey (github.com/TobyBaril/EarlGrey/) [104]. Known repeats were first identified and masked using RepeatMasker (version 4.1.0) [105] using the *Arthropoda* library from RepBase (version 23.08) and Dfam (release 3.3) [48,49]. Low complexity repeats and RNA were not masked (*-nolow and -norna*) and a sensitive search was performed (*-s*). Following this, a *de novo* repeat library was constructed using RepeatModeler (version 1.0.11), with RECON (version 1.08) and RepeatScout (version 1.0.5) [106–108]. To generate maximum-length consensus sequences for the *de novo* repeat library, novel repeats identified by RepeatModeler were curated using an automated version of the 'BLAST, Extract, Extend' process [28]. A BLASTn search was performed on the *D. plexippus* genome to obtain up to the top 40 hits for each TE family identified by RepeatModeler [109]. Sequences were extracted with 1,000 base pairs of flanking sequences at the 5' and 3' ends. Each set of family sequences were aligned using MAFFT (version 7.453) [110]. Alignments were trimmed with trimAl (version 1.4.rev22) to retain high-quality positions in the alignment (*-gt* 0.6 *-cons* 60) [111]. Updated consensus sequences were computed with EMBOSS cons (*-plurality* 3) to generate a new TE library featuring consensus sequences with extended flanks [112]. This process was repeated through 5 iterations to obtain maximum-length consensus sequences.

Following automated curation, *de novo* family sequences were subject to a manual curation following protocols described by Goubert et al. (2022)[113]. Here, the final

alignments generated by Earl Grey were visualised using AliView [114], and poorly-aligned

positions and ends were trimmed where single-copy DNA was present. Consensus

sequences were generated using EMBOSS cons [112] to generate a library of manually

curated elements, which was clustered using cd-hit-est [115] with the options recommended

by Goubert et al. (2022) [113] to reduce redundancy. To classify TEs, the TE library was

analysed with the diagnostic tool TE-Aid (https://github.com/clemgoub/TE-Aid) to identify TE-

associated protein domains in sequence ORFs and to look for the presence of TIRs and

LTRs [105,109,112]. Following this, blastn and nhmmer searches were performed against

the Dfam database (version 3.3) to identify homology to previously described elements

[49,109,116].

The resulting *de novo* repeat library was combined with the RepBase *Arthropoda*

library and utilised to annotate repetitive elements using RepeatMasker with a score

threshold at the relatively conservative level of 400 (*-cutoff* 400), to exclude poor matches

unlikely to be true TE sequences. Following this, the *D. plexippus* genome was analysed

with LTR_Finder (version 1.07), using the LTR_Finder parallel wrapper [117,118] to identify

full-length LTR elements. Final TE annotations were defragmented and refined using the

loose merge (*-loose*) command in RepeatCraft [119]. Overlapping annotations were

removed using MGKit (version 0.4.1) filter-gff, with the longest element of overlapping

annotations retained (*-c* length *-a* length) [120]. Finally, all repeats less than 100bp in length

were removed before the final TE quantification to decrease spurious hits. Final repeat

annotation is provided in datasets S1 and S2 (Additional Files 10 and 11), and *de novo* TE

library is provided in dataset S3 (Additional File 12).

Quantifying DNA Loss Rates Between the Monarch and *D. chrysippus*

To estimate DNA loss rates between the monarch and *D. chrysippus*, the DelGet.pl

script (https://github.com/4ureliek/DelGet) was used to estimate DNA loss from small (<30nt)

and mid-size (30<nt<10,000) deletions, as described previously [45]. Briefly, the script

identifies orthologous regions of 10,000bp between three input species (an outgroup and

two species with known branch length in Mya) using BLAT

(https://genome.ucsc.edu/FAQ/FAQblat.html). These regions are extracted and aligned with

MUSCLE [121] before deletions are quantified. Species-specific gaps are identified and

quantified. Deletion rates are then calculated between the two species of interest by dividing

the normalised deleted nucleotide count (del_nt/10kb) by the corresponding Mya since

divergence.

Identifying Transposable Element Association with Genomic Compartments

The *D. plexippus* gene annotation file was processed to generate a GFF file

containing coordinates of exons, introns, and 5' and 3' flanking regions. Gene flanking

regions are defined here as 20kb directly upstream or downstream of the gene body. These

flanking regions are determined to identify TEs at a distance that could be described as the

proximate promoter region, rather than just accounting for the core promoter region, to

include both promoter and more distal enhancer regions for genes [122]. Bedtools intersect

(version 2.27.1) was used to identify the number of bases of overlap (-*wao*) between TEs

and different genomic compartments [123]. Following this, quantification was performed in R.

TE Hotspots and Coldspots

To identify TE and gene hotspots and coldspots, chromosomal scaffolds were split

into 100kb windows, and the TE density in each window was calculated and used to produce

the frequency distribution of observations. A polynomial model was fitted to the frequency

distribution, to generate a smooth curve through the data points using the lm function in R.

Using these models, the total area under the curve (AUC) was calculated using the

area_under_curve function of the bayestestR package [124]. From this, the 1% and 99%

AUC cutoff values were determined. Starting with the first data point and iteratively adding

subsequent datapoints with each round, the AUC for each collection of points was

determined. The maximum AUC value not exceeding 99% of the total AUC was taken as the

cutoff where windows containing higher TE or gene coverage were considered to be

hotspots. The minimum AUC value not exceeding 1% of the total AUC was taken as the

cutoff where windows containing lower TE or gene coverage were considered to be

coldspots (Figure S8; Additional File 13). Karyoplots of the *D. plexippus* genome, illustrating

gene and TE locations, highlighting TE hotspots and coldspots were generated using the

KaryoploteR package in R [125].

Estimating TE Age and Activity

TE activity dates were estimated using Kimura 2-parameter genetic distances

(including CpG sites) calculated between individual insertions and the consensus sequence

of each TE family [126]. A neutral mutation rate of 0.0116 substitutions per site per million

years was used for *D. plexippus*, as previously described for the monarch [127].

Turnover of LINEs and PLEs

To interrogate LINE and PLE decay patterns, each LINE and PLE sequence was

extracted from the genome using bedtools getfasta [123]. Extracted sequences were used to

query the repeat library using BLASTn with tabular output specified (*-outfmt* 6). In the event

of multiple matches, only the top match was retained (defined by percent identity).

Coordinates of each match were scaled to normalise the length of each consensus

sequence. Mapped fragments were plotted using ggplot2 of the tidyverse package in R

[128,129].

## Identification of Intact LINEs

To identify full-length LINEs, all elements longer than 2,700bp were extracted from

the monarch genome using bedtools getFasta [123]. Open reading frames (ORFs) were

detected using EMBOSS getORF [112]. ORF sequences were filtered to retain all

sequences > 600aa. RPSBLAST+ (v2.9.0+) [109] was used to query the ORF sequences

against the CDD database [130] to identify conserved protein domains. Sequences were

classified as intact LINEs based on the presence of relevant protein domains as previously

described [72] (Table S7, Additional File 1).

## Identification of Intact PLEs

To identify intact PLEs, all elements longer than 2,400bp were extracted from the

monarch genome using bedtools getfasta [123] as full-length PLEs are ~2,500bp long [131].

Open reading frames (ORFs) were detected using EMBOSS getORF [112]. ORF sequences

were filtered to retain all sequences >200aa. RPSBLAST+ (v2.9.0+) [109]  was used to

query the ORF sequences against the CDD database [130]  to identify conserved  protein

domains. Sequences were classified as intact PLEs based on the presence of reverse

transcriptase (RT) and a GIY-YIG endonuclease.

## Classification of Novel DNA TEs

To further classify novel DNA elements in the monarch beyond homology to known

elements, as used by RepeatModeler, DDE transposases were compared with curated

alignments for all DDE TE superfamilies [88]. Specifically, dashes in curated alignments of

all known DDE TE sequences for each of the 19 families were removed, and the resultant

fasta file was used as the subject in a blastx search against the *de novo* monarch TE library,

with an e-value threshold of $1 \times 10^{-20}$ and tabular output specified (*-evalue* 1e-20 *-outfmt* 6)

[109]. The best match for each *de novo* query sequence was retained, with a minimum 50%

identity over at least 50% of the total length of the curated DDE subject sequence. Following

this, novel DNA elements were only detected for the Tc1-Mariner family (Figure S7;

Additional File 9). Transposase sequences for the *de novo* monarch sequences were

extracted and aligned to the original DDE alignments for their respective families using the

profile alignment command in MAFFT [110]. Sequences from related transposon

superfamilies were used as outgroups, as described [88]. To estimate phylogenies for the

two DDE superfamilies, we performed 1,000 ultra-fast bootstrap replicates in IQ-TREE

(v1.6.12) [132] specifying the best-fit amino acid model for each family as indicated by

ModelFinder [133]. Phylogenetic trees were visualised in figtree (v1.4.4)

(http://tree.bio.ed.ac.uk/software/figtree/).

Association of TEs with Monarch Colouration and Immune Genes

We obtained the transcript sequence in fasta format for the myosin gene responsible for

wing colouration from MonarchBase, under accession DPOGS206617 [34]. To identify the

matching region in the v4 monarch assembly, a blastn search was performed to query the

monarch assembly [109]. The coordinates of the myosin gene were obtained and used in a

bedtools window search to identify TEs within 20kb (-*w* 20000)[123]. TE copy number within

20kb of gene bodies was quantified for each monarch gene and a polynomial model was

fitted to the frequency distribution to generate a smooth curve through the data points using

the lm function in R. Using these models, the total area under the curve (AUC) was

calculated using the area_under_curve function of the bayestestR package [124]. From this,

the 2.5% and 97.5% AUC cutoff values were determined, equivalent to selecting the central

95% of the data distribution. Starting with the first data point and iteratively adding

subsequent datapoints with each round, the AUC for each collection of points was

determined. The maximum AUC value not exceeding 97.5% of the total AUC was taken as

the cutoff where genes containing higher TE copy number were considered significantly

enriched for TEs. The minimum AUC value not exceeding 2.5% of the total AUC was taken

as the cutoff where genes containing lower TE copy number were considered significantly

depleted for TEs (Figure S5; Additional File 7). A karyoplot of the myosin gene region

including 20kb flanks was generated in R using the karyoploteR package [125].

To investigate associations between monarch immunity genes and TEs, nucleotide

and protein sequences of monarch immune genes were obtained in fasta format from

MonarchBase [34]. To identify corresponding regions in the v4 monarch assembly, blastn

and tblastn searches were used to query the monarch genome assembly with the monarch

immune gene nucleotide and protein sequences [109]. Matches were filtered by percentage

identity and query match coordinates to identify the locus of each immune gene. A bed file

was generated, containing the name and location of each immune gene, and a bedtools

window search was performed to identify TEs within 20kb of each immune gene(-*w*

20000)[123]. The same method was applied to identify TEs within 20kb of all other genes.

TE bases surrounding each immune gene were quantified in R [134]. As TE count was not

normally distributed, a Kruskal-Wallis test was used to compare TE abundance between

immune genes and other genes, between different TE classifications around immune genes,

and between different immune gene types. Unless otherwise stated, all plots were generated

using the ggplot2 package of the tidyverse suite in R [128,129].

## Figure Legends

Figure 1. TE content in the monarch, *D. chrysippus* and *D. melanippus.* Major TE types are

represented by different colours indicated in the key (A) Pie charts illustrating proportions of

each *Danaus* genome comprised of the main TE classifications. (B) Repeat landscapes for

each *Danaus* species. The x axis indicates the level of Kimura 2-parameter genetic distance

observed between TE insertions and their respective consensus sequences in percent. More

recent elements are located to the right of the x axis. The y axis indicates the percentage of

the genome occupied by TE insertions of each genetic distance class.

Figure 2. Processes leading to genome size differences between the monarch, *D.*

*chrysippus*, and *D. melanippus*. (A) Three hypotheses leading to observed differences in

genome size between *Danaus* species. Hypotheses indicated by colours in the key. (B)

Shared and unique TE families across the three *Danaus* species. Numbers indicate distinct

TE families. Percentages indicate proportion of total TE families found in each section of the

Venn diagram. (C) TE coverage in each *Danaus* species split by TE shared status. Major TE

types are represented by different colours indicated in the key.

Figure 3. TE location in different genomic compartments in the monarch. Major TE types are

represented by different colours indicated in the key. (A) Quantity of TE base pairs found in

different genomic compartments. (B) Proportion of each genomic compartment contributed

by TEs.

Figure 4. Violin plot showing kilobases of TE sequence per 100kb window across the

genome for each of the main TE classifications. Red points show the mean kilobases of TE

sequence per 100kb.

Figure 5. Karyoplots illustrating monarch chromosomes. Grey boxes represent

chromosomes, with black regions representing genic regions of each chromosome. (A)

Above each chromosome, TE density is shown for 100kb windows, with major TE types

represented by different colours indicated in the key. (B) Above each chromosome, TE

hotspots and coldspots are shown, with severity represented by the different colours

indicated in the key,

Figure 6. Violin plot illustrating the estimated insertion time (Mya) for each of the major TE

types, represented by different colours indicated in the key. The black bar shows the

estimated divergence time of the monarch, approximately 32Mya [66].

Figure 7. TE age against copy number. Major TE types are represented by different colours

indicated in the key. X axis shows estimated age of each TE family, Y axis shows log10

transformed TE copy number identified in the monarch.

## Additional Files

Additional File 1: Supplementary Tables, including contents page with table legends. Tables

S1-S10.

Additional File 2: Detailed discussion on the discrepancies between TE annotations of the

monarch presented in this study, and those presented in another recent study focussing on

*D. chrysippus* [51].

Additional File 3: Figure S1. Scatter plot illustrating estimated TE age for each TE family.

Major TE types are represented by different colours indicated in the key. Elements are

ordered from oldest to youngest, within each major type.

Additional File 4: Figure S2. Repeat landscapes for the monarch. The x axis indicates the

level of Kimura 2-parameter genetic distance observed between TE insertions and their

respective consensus sequences in percent. More recent elements are located to the right of

the x axis. The y axis indicates the percentage of the genome occupied by TE insertions of

each genetic distance class.

Additional File 5: Figure S3. Plot illustrating LINE and Penelope fragments annotated in the

monarch genome. The x axis indicates normalised start and end coordinates for each

element relative to its consensus. Elements are organised from longest to shortest, with

short elements found towards the top of the plot. Families are represented by different

colours indicated by the key.

Additional File 6: Figure S4. Karyoplot of the monarch gene association with colouration,

myosin, obtained from MonarchBase under accession DPOGS206617. Main TE types within

20kb are represented by different colours indicated in the key.

Additional File 7: Figure S5. Plot illustrating the distribution of TE copy number found with

20kb of all genes, showing the 2.5% and 97.5% AUC cutoffs used to determine the TE copy

number within 20kb of a given gene for it to be significantly depleted or enriched in

comparison to other genes.

Additional File 8: Figure S6. Boxplots illustrating: (A) Normalised TE coverage in 20kb

regions surrounding each immune gene type. (B) Normalised TE count in 20kb regions

surrounding each immune gene type. Main TE types are represented by different colours

indicated in the key.

Additional File 9: Figure S7. Phylogenetic tree of Tc1-Mariner DDE transposases, with novel

monarch families added (highlighted in red). Branch support from 1000 ultra-fast bootstrap

repetitions. Novel Tc1-Mariner families highlighted in teal are the young TEs found with

extremely high copy number.

Additional File 10: Dataset S1.  Coordinates and classification of all transposable element

sequences identified in the monarch genome using the *de novo* library in conjunction with

RepBase *Arthropoda* library and Dfam, in bed format.

Additional File 11: Dataset S2.  Coordinates and classification of all transposable element

sequences identified in the monarch genome using the *de novo* library in conjunction with

RepBase *Arthropoda* library and Dfam, in GFF format.

Additional File 12: Dataset S3.  *De novo* transposable element consensus sequences

identified in the monarch in FASTA format.

Additional File 13: Figure S8. Plots illustrating: (A) Distribution of genome windows with TE

coverage, with lines showing 1% and 99% AUC cutoffs used to define hotspots and

coldspots. (B) Distribution of genome windows with host gene coverage, with lines showing

1% and 99% AUC cutoffs used to define hotspots and coldspots.

Abbreviations

TE: Transposable element

Non-LTR: Non-long terminal repeat

LTR: Long terminal repeat

LINEs: Long INterspersed Elements

SINEs: Short INterspersed Elements

PLEs: *Penelope*-like Elements

TSS: Transcription Start Site

## References

788    [1]    Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, et al. Ten

789            things you should know about transposable elements. Genome Biol 2018;19:199.

790            https://doi.org/10.1186/s13059-018-1577-z.

791    [2]    Klein SJ, O'Neill RJ. Transposable elements: genome innovation, chromosome

792            diversity, and centromere conflict. Chromosom Res 2018;26:5–23.

793            https://doi.org/10.1007/s10577-017-9569-5.

794    [3]    Feschotte C, Pritham EJ. DNA Transposons and the Evolution of Eukaryotic

795            Genomes. Annu Rev Genet 2007;41:331–68.

796            https://doi.org/10.1146/annurev.genet.40.110405.090448.

797    [4]    Belyayev A. Bursts of transposable elements as an evolutionary driving force. J Evol

798            Biol 2014;27:2573–84. https://doi.org/10.1111/jeb.12513.

799    [5]    Schrader L, Schmitz J. The impact of transposable elements in adaptive evolution.

800            Mol Ecol 2018. https://doi.org/10.1111/mec.14794.

[6]    Casola C, Lawing AM, Betrán E, Feschotte C. PIF-like transposons are common in

Drosophila and have been repeatedly domesticated to generate new host genes. Mol

Biol Evol 2007;24:1872–88. https://doi.org/10.1093/molbev/msm116.

[7]    Jangam D, Feschotte C, Betrán E. Transposable Element Domestication As an

Adaptation to Evolutionary Conflicts. Trends Genet 2017;33:817–31.

https://doi.org/10.1016/j.tig.2017.07.011.

[8]    Volff JN. Turning junk into gold: Domestication of transposable elements and the

creation of new genes in eukaryotes. BioEssays 2006.

https://doi.org/10.1002/bies.20452.

[9]    Mita P, Boeke JD. How retrotransposons shape genome regulation. Curr Opin Genet

Dev 2016;37:90–100. https://doi.org/10.1016/j.gde.2016.01.001.

[10]   Bourque G. Transposable elements in gene regulation and in the evolution of

vertebrate genomes. Curr Opin Genet Dev 2009;19:607–12.

https://doi.org/10.1016/j.gde.2009.10.013.

[11]   Feschotte C. Transposable elements and the evolution of regulatory networks. Nat

Rev Genet 2008;9:397–405. https://doi.org/10.1038/nrg2337.

[12]   Rebollo R, Romanish MT, Mager DL. Transposable Elements: An Abundant and

Natural Source of Regulatory Sequences for Host Genes. Annu Rev Genet

2012;46:21–42. https://doi.org/10.1146/annurev-genet-110711-155621.

[13]   Chuong EB, Elde NC, Feschotte C. Regulatory activities of transposable elements:

From conflicts to benefits. Nat Rev Genet 2017;18:71–86.

https://doi.org/10.1038/nrg.2016.139.

[14]   Bennetzen JL. Transposable element contributions to plant genome evolution. Plant

Mol Biol 2000;42:251–69. https://doi.org/10.1023/A:1006344508454.

[15]   Casals F, Cáceres M, Ruiz A. The Foldback-like transposon Galileo is involved in the

generation of two different natural chromosomal inversions of Drosophila buzzatii. Mol

Biol Evol 2003;20:674–85. https://doi.org/10.1093/molbev/msg070.

[16]  Campos-Sánchez R, Cremona MA, Pini A, Chiaromonte F, Makova KD. Integration

and Fixation Preferences of Human and Mouse Endogenous Retroviruses Uncovered

with Functional Data Analysis. PLoS Comput Biol 2016;12:1–41.

https://doi.org/10.1371/journal.pcbi.1004956.

[17]  Oliver KR, Greene WK. Transposable elements: Powerful facilitators of evolution.

BioEssays 2009;31:703–14. https://doi.org/10.1002/bies.200800219.

[18]  Hughes JF, Coffin JM. Human endogenous retroviral elements as indicators of

ectopic recombination events in the primate genome. Genetics 2005;171:1183–94.

https://doi.org/10.1534/genetics.105.043976.

[19]  Kent T V., Uzunović J, Wright SI. Coevolution between transposable elements and

recombination. Philos Trans R Soc B Biol Sci 2017;372.

https://doi.org/10.1098/rstb.2016.0458.

[20]  Coates BS, Hellmich RL, Grant DM, Abel CA. Mobilizing the genome of lepidoptera

through novel sequence gains and end creation by non-autonomous Lep1 Helitrons.

DNA Res 2012;19:11–21. https://doi.org/10.1093/dnares/dsr038.

[21]  Biémont C, Vieira C. What transposable elements tell us about genome organization

and evolution: The case of Drosophila. Cytogenet Genome Res 2005;110:25–34.

https://doi.org/10.1159/000084935.

[22]  Morgante M, Brunner S, Pea G, Fengler K, Zuccolo A, Rafalski A. Gene duplication

and exon shuffling by helitron-like transposons generate intraspecies diversity in

maize. Nat Genet 2005;37:997–1002. https://doi.org/10.1038/ng1615.

[23]  Philippsen GS, Avaca-Crusca JS, Araujo APU, DeMarco R. Distribution patterns and

impact of transposable elements in genes of green algae. Gene 2016;594:151–9.

https://doi.org/10.1016/j.gene.2016.09.012.

[24]    Wu M, Li L, Sun Z. Transposable element fragments in protein-coding regions and

853            their contributions to human functional proteins. Gene 2007;401:165–71.

854            https://doi.org/10.1016/j.gene.2007.07.012.

855    [25]    Daniel C, Behm M, Öhman M. The role of Alu elements in the cis-regulation of RNA

856            processing. Cell Mol Life Sci 2015;72:4063–76. https://doi.org/10.1007/s00018-015-

857            1990-3.

858    [26]    Vorechovsky I. Transposable elements in disease-associated cryptic exons. Hum

859            Genet 2010;127:135–54. https://doi.org/10.1007/s00439-009-0752-4.

860    [27]    Bejerano G, Lowe CB, Ahituv N, King B, Siepel A, Salama SR, et al. A distal

861            enhancer and an ultraconserved exon are derived from a novel retroposon. Nature

862            2006;441:87–90. https://doi.org/10.1038/nature04696.

863    [28]    Platt RN, Blanco-Berdugo L, Ray DA. Accurate transposable element annotation is

864            vital when analyzing new genome assemblies. Genome Biol Evol 2016;8:403–10.

865            https://doi.org/10.1093/gbe/evw009.

866    [29]    Bates HW. Contributions to an Insect Fauna of the Amazon Valley.—Lepidoptera:—

867            Heliconinæ. J Proc Linn Soc London Zool 1862;6:73–7.

868            https://doi.org/10.1111/J.1096-3642.1862.TB00932.X.

869    [30]    Müller F. Ituna and Thyridia: a remarkable case of mimicry in butterflies. Trans

870            Entomol Soc L 1879;1879:20–9.

871    [31]    Ehrlich PR, Raven PH. Butterflies and Plants: A Study in Coevolution. Evolution (N Y)

872            1964;18:586. https://doi.org/10.2307/2406212.

873    [32]    Kettlewell, B. Evolution of melanism: the study of a recurring necessity 1973.

874    [33]    Dasmahapatra KK, Walters JR, Briscoe AD, Davey JW, Whibley A, Nadeau NJ, et al.

875            Butterfly genome reveals promiscuous exchange of mimicry adaptations among

876            species. Nature 2012;487:94–8. https://doi.org/10.1038/nature11041.

[34]    Zhan S, Reppert SM. MonarchBase: the monarch butterfly genome database. Nucleic

Acids Res 2013;41:D758–63. https://doi.org/10.1093/NAR/GKS1057.

[35]    Zhan S, Zhang W, Niitepõld K, Hsu J, Haeger JF, Zalucki MP, et al. The genetics of

monarch butterfly migration and warning colouration. Nat 2014 5147522

2014;514:317–21. https://doi.org/10.1038/nature13812.

[36]    Zhan S, Merlin C, Boore JL, Reppert SM. The monarch butterfly genome yields

insights into long-distance migration. Cell 2011;147:1171–85.

https://doi.org/10.1016/j.cell.2011.09.052.

[37]    Wu M, Sun Z, Luo G, Hu C, Zhang W, Han Z. Cloning and characterization of

piggyBac-like elements in lepidopteran insects. Genetica 2011;139:149–54.

https://doi.org/10.1007/s10709-010-9542-0.

[38]    Sormacheva I, Smyshlyaev G, Mayorov V, Blinov A, Novikov A, Novikova O. Vertical

evolution and horizontal transfer of CR1 non-LTR retrotransposons and Tc1/mariner

DNA transposons in Lepidoptera species. Mol Biol Evol 2012;29:3685–702.

https://doi.org/10.1093/molbev/mss181.

[39]    Reiss D, Mialdea G, Miele V, de Vienne D, Peccoud J, Gilbert C, et al. Global survey

of mobile DNA horizontal transfer in arthropods reveals Lepidoptera as a prime

hotspot. PLOS Genet 2019;15:e1007965.

https://doi.org/10.1371/journal.pgen.1007965.

[40]    Talla V, Suh A, Kalsoom F, Dincă V, Vila R, Friberg M, et al. Rapid increase in

genome size as a consequence of transposable element hyperactivity in wood-white

(leptidea) butterflies. Genome Biol Evol 2017;9:2491–505.

https://doi.org/10.1093/gbe/evx163.

[41]    Hof AEV t., Campagne P, Rigden DJ, Yung CJ, Lingley J, Quail MA, et al. The

industrial melanism mutation in British peppered moths is a transposable element.

Nature 2016;534:102–5. https://doi.org/10.1038/nature17951.

[42]   Gu L, Reilly PF, Lewis JJ, Reed RD, Andolfatto P, Walters JR. Dichotomy of dosage
compensation along the neo Z chromosome of the Monarch butterfly. Curr Biol
2019;29:4071–7.

[43]   Chénais B, Caruso A, Hiard S, Casse N. The impact of transposable elements on
eukaryotic genomes: From genome size increase to genetic adaptation to stressful
environments. Gene 2012;509:7–15. https://doi.org/10.1016/j.gene.2012.07.042.

[44]   Osanai-Futahashi M, Suetsugu Y, Mita K, Fujiwara H. Genome-wide screening and
characterization of transposable elements and their distribution analysis in the
silkworm, Bombyx mori. Insect Biochem Mol Biol 2008;38:1046–57.
https://doi.org/10.1016/j.ibmb.2008.05.012.

[45]   Kapusta A, Suh A, Feschotte C. Dynamics of genome size evolution in birds and
mammals. Proc Natl Acad Sci U S A 2017;114:E1460–9.
https://doi.org/10.1073/pnas.1616702114.

[46]   Kidwell MG. Transposable elements and the evolution of genome size in eukaryotes.
Genetica 2002;115:49–63. https://doi.org/10.1023/A:1016072014259.

[47]   Stritt C, Gordon SP, Wicker T, Vogel JP, Roulin AC. Recent activity in expanding
populations and purifying selection have shaped transposable element landscapes
across natural accessions of the mediterranean grass brachypodium distachyon.
Genome Biol Evol 2018;10:304–18. https://doi.org/10.1093/gbe/evx276.

[48]   Jurka J, Kapitonov V V., Pavlicek A, Klonowski P, Kohany O, Walichiewicz J.
Repbase Update, a database of eukaryotic repetitive elements. Cytogenet Genome
Res 2005;110:462–7. https://doi.org/10.1159/000084979.

[49]   Hubley R, Finn RD, Clements J, Eddy SR, Jones TA, Bao W, et al. The Dfam
database of repetitive DNA families. Nucleic Acids Res 2016;44:D81–9.

927    https://doi.org/10.1093/nar/gkv1272.

928 [50] Baril T, Imrie R, Hayward A. TobyBaril/EarlGrey: Earl Grey v1.1 2021.

929    https://doi.org/10.5281/ZENODO.5675911.

930 [51] Singh KS, De-Kayne R, Omufwoko KS, Martins DJ, Bass C, ffrench-Constant R, et al.

931    Genome assembly of Danaus chrysippus and comparison with the Monarch Danaus

932    plexippus. G3 Genes|Genomes|Genetics 2021.

933    https://doi.org/10.1093/G3JOURNAL/JKAB449.

934 [52] Wiemers M, Chazot N, Wheat CW, Schweiger O, Wahlberg N. A complete time-

935    calibrated multi-gene phylogeny of the European butterflies. Zookeys 2020;938:97.

936    https://doi.org/10.3897/ZOOKEYS.938.50878.

937 [53] Sultana T, Zamborlini A, Cristofari G, Lesage P. Integration site selection by

938    retroviruses and transposable elements in eukaryotes. Nat Rev Genet 2017;18:292–

939    308. https://doi.org/10.1038/nrg.2017.7.

940 [54] Sawicka K, Bushell M, Spriggs KA, Willis AE. Polypyrimidine-tract-binding protein: A

941    multifunctional RNA-binding protein. Biochem Soc Trans 2008;36:641–7.

942    https://doi.org/10.1042/BST0360641.

943 [55] Tree DRP, Shulman JM, Rousset R, Scott MP, Gubb D, Axelrod JD. Prickle mediates

944    feedback amplification to generate asymmetric planar cell polarity signaling. Cell

945    2002;109:371–81. https://doi.org/10.1016/S0092-8674(02)00715-8.

946 [56] Laugier E, Yang Z, Fasano L, Kerridge S, Vola C. A critical role of teashirt for

947    patterning the ventral epidermis is masked by ectopic expression of tiptop, a paralog

948    of teashirt in Drosophila. Dev Biol 2005;283:446–58.

949    https://doi.org/10.1016/J.YDBIO.2005.05.005.

950 [57] Datta RR, Lurye JM, Kumar JP. Restriction of ectopic eye formation by Drosophila

951    teashirt and tiptop to the developing antenna. Dev Dyn 2009;238:2202–10.

https://doi.org/10.1002/DVDY.21927.

[58]  Contreras EG, Palominos T, Glavic Á, Brand AH, Sierralta J, Oliva C. The

transcription factor SoxD controls neuronal guidance in the Drosophila visual system.

Sci Reports 2018 81 2018;8:1–12. https://doi.org/10.1038/s41598-018-31654-5.

[59]  Matsubara D, Horiuchi SY, Shimono K, Usui T, Uemura T. The seven-pass

transmembrane cadherin Flamingo controls dendritic self-avoidance via its binding to

a LIM domain protein, Espinas, in Drosophila sensory neurons. Genes Dev

2011;25:1982–96. https://doi.org/10.1101/GAD.16531611.

[60]  Bessa J, Gebelein B, Pichaud F, Casares F, Mann RS. Combinatorial control of

Drosophila eye development by Eyeless, Homothorax, and Teashirt. Genes Dev

2002;16:2415–27. https://doi.org/10.1101/GAD.1009002.

[61]  Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, et al. A unified

classification system for eukaryotic transposable elements. Nat Rev Genet

2007;8:973–82. https://doi.org/10.1038/nrg2165.

[62]  Oliveira SG, Bao W, Martins C, Jurka J. Horizontal transfers of Mariner transposons

between mammals and insects. Mob DNA 2012;3:1–6. https://doi.org/10.1186/1759-

8753-3-14.

[63]  Ivancevic AM, Kortschak RD, Bertozzi T, Adelson DL. Horizontal transfer of BovB and

L1 retrotransposons in eukaryotes. Genome Biol 2018;19:1–13.

https://doi.org/10.1186/s13059-018-1456-7.

[64]  Gilbert C, Schaack S, Pace JK, Brindley PJ, Feschotte C. A role for host-parasite

interactions in the horizontal transfer of transposons across phyla. Nature

2010;464:1347–50. https://doi.org/10.1038/nature08939.

[65]  Wallau GL, Ortiz MF, Loreto ELS. Horizontal transposon transfer in eukarya:

Detection, bias, and perspectives. Genome Biol Evol 2012;4:689–99.

https://doi.org/10.1093/gbe/evs055.

[66]    Espeland M, Breinholt J, Willmott KR, Warren AD, Vila R, Toussaint EFA, et al. A

Comprehensive and Dated Phylogenomic Analysis of Butterflies. Curr Biol

2018;28:770-778.e5. https://doi.org/10.1016/J.CUB.2018.01.061.

[67]    Ustyugova S V, Amosova AL, Lebedev YB, Sverdlov ED. Cell line fingerprinting using

retroelement insertion polymorphism. Biotechniques 2005;38:561–5.

[68]    Lavoie CA, II RNP, Novick PA, Counterman BA, Ray DA. Transposable element

evolution in Heliconius suggests genome diversity within Lepidoptera. Mob DNA

2013;21:1–10.

[69]    Blass E, Bell M, Boissinot S. Accumulation and rapid decay of non-LTR

retrotransposons in the genome of the three-spine stickleback. Genome Biol Evol

2012;4:687–702.

[70]    Ivancevic AM, Kortschak RD, Bertozzi T, Adelson DL. LINEs between Species:

Evolutionary Dynamics of LINE-1 Retrotransposons across the Eukaryotic Tree of

Life. Genome Biol Evol 2016;8:3301–22. https://doi.org/10.1093/GBE/EVW243.

[71]    Evgen'ev MB, Arkhipova IR. Penelope-like elements – a new class of retroelements:

distribution, function and possible evolutionary significance. Cytogenet Genome Res

2005;110:510–21. https://doi.org/10.1159/000084984.

[72]    Kapitonov V V., Tempel S, Jurka J. Simple and fast classification of non-LTR

retrotransposons based on phylogeny of their RT domain protein sequences. Gene

2009;448:207. https://doi.org/10.1016/J.GENE.2009.07.019.

[73]    Woronik A, Tunström K, Perry MW, Neethiraj R, Stefanescu C, Celorio-Mancera M de

la P, et al. A transposable element insertion is associated with an alternative life

history strategy. Nat Commun 2019 101 2019;10:1–11.

https://doi.org/10.1038/s41467-019-13596-2.

[74]    Hill T, Koseva BS, Unckless RL. The Genome of Drosophila innubila Reveals

1003            Lineage-Specific Patterns of Selection in Immune Genes. Mol Biol Evol

1004            2019;36:1405–17. https://doi.org/10.1093/MOLBEV/MSZ059.

1005    [75]    Sackton TB, Lazzaro BP, Schlenke TA, Evans JD, Hultmark D, Clark AG. Dynamic

1006            evolution of the innate immune system in Drosophila. Nat Genet 2007 3912

1007            2007;39:1461–8. https://doi.org/10.1038/ng.2007.60.

1008    [76]    Shultz AJ, Sackton TB. Immune genes are hotspots of shared positive selection

1009            across birds and mammals. Elife 2019;8. https://doi.org/10.7554/ELIFE.41815.

1010    [77]    Kimbrell DA, Beutler B. The evolution and genetics of innate immunity. Nat Rev Genet

1011            2001 24 2001;2:256–67. https://doi.org/10.1038/35066006.

1012    [78]    Jurka J, Kapitonov V V., Kohany O, Jurka M V. Repetitive Sequences in Complex

1013            Genomes: Structure and Evolution. Annu Rev Genomics Hum Genet 2007;8:241–59.

1014            https://doi.org/10.1146/annurev.genom.8.080706.092416.

1015    [79]    Kidwell MG, Lisch DR. Transposable elements and host genome evolution. Trends

1016            Ecol Evol 2000;15:95–9. https://doi.org/10.1016/S0169-5347(99)01817-0.

1017    [80]    Biémont C, Vieira C. Junk DNA as an evolutionary force. Nature 2006;443:521–4.

1018    [81]    Cosby RL, Chang N-C, Feschotte C. Host–transposon interactions: conflict,

1019            cooperation, and cooption. Genes Dev 2019;33:1098–116.

1020            https://doi.org/10.1101/GAD.327312.119.

1021    [82]    Casacuberta E, González J. The impact of transposable elements in environmental

1022            adaptation. Mol Ecol 2013;22:1503–17. https://doi.org/10.1111/mec.12170.

1023    [83]    Huff JT, Zilberman D, Roy SW. Mechanism for DNA transposons to generate introns

1024            on genomic scales. Nature 2016;538:533–6.

[84]    Gilbert C, Feschotte C. Horizontal acquisition of transposable elements and viral

sequences: patterns and consequences. Curr Opin Genet Dev 2018;49:15–24.

https://doi.org/10.1016/j.gde.2018.02.007.

[85]   Galbraith JD, Ludington AJ, Sanders KL, Suh A, Adelson DL. Horizontal transfer and

subsequent explosive expansion of a DNA transposon in sea kraits (Laticauda). Biol

Lett 2021;17:20210342. https://doi.org/10.1098/RSBL.2021.0342.

[86]   Le Rouzic A, Capy P. The first steps of transposable elements invasion: Parasitic

strategy vs. genetic drift. Genetics 2005;169:1033–43.

https://doi.org/10.1534/genetics.104.031211.

[87]   Kofler R, Senti KA, Nolte V, Tobler R, Schlötterer C. Molecular dissection of a natural

transposable element invasion. Genome Res 2018;28:824–35.

https://doi.org/10.1101/gr.228627.117.

[88]   Dupeyron M, Baril T, Hayward A. Broadscale evolutionary analysis of eukaryotic DDE

transposons. BioRxiv 2021:2021.09.26.461848.

https://doi.org/10.1101/2021.09.26.461848.

[89]   Prysby MD, Oberhauser KS. Temporal and Geographic Variation in Monarch

Densities: Citizen Scientists Document Monarch Population Patterns. Monarch

Butterfly Biol Conserv 2004:9.

[90]   Tang Z, Zhang HH, Huang K, Zhang XG, Han MJ, Zhang Z. Repeated horizontal

transfers of four DNA transposons in invertebrates and bats. Mob DNA 2015;6:1–10.

https://doi.org/10.1186/s13100-014-0033-1.

[91]   Kelleher ES, Azevedo RBR, Zheng Y. The evolution of small-RNA-mediated silencing

of an invading transposable element. Genome Biol Evol 2018;10:3038–57.

https://doi.org/10.1093/gbe/evy218.

[92]   Goodier JL. Restricting retrotransposons: A review. Mob DNA 2016;7.

https://doi.org/10.1186/s13100-016-0070-z.

[93] Ray DA, Grimshaw JR, Halsey MK, Korstian JM, Osmanski AB, Sullivan KAM, et al.

Simultaneous TE Analysis of 19 Heliconiine Butterflies Yields Novel Insights into

Rapid TE-Based Genome Diversification and Multiple SINE Births and Deaths.

Genome Biol Evol 2019;11:2162–77. https://doi.org/10.1093/gbe/evz125.

[94] Levin HL, Moran J V. Dynamic interactions between transposable elements and their

hosts. Nat Rev Genet 2011;12:615–27. https://doi.org/10.1038/nrg3030.

[95] Fontanillas P, Hartl DL, Reuter M. Genome organization and gene expression shape

the transposable element distribution in the Drosophila melanogaster euchromatin.

PLoS Genet 2007;3:2256–67. https://doi.org/10.1371/journal.pgen.0030210.

[96] Mawaribuchi S, Takahashi S, Wada M, Uno Y, Matsuda Y, Kondo M, et al. Sex

chromosome differentiation and the W- and Z-specific loci in Xenopus laevis. Dev Biol

2017;426:393–400. https://doi.org/10.1016/J.YDBIO.2016.06.015.

[97] Dechaud C, Volff JN, Schartl M, Naville M. Sex and the TEs: transposable elements

in sexual development and function in animals. Mob DNA 2019 101 2019;10:1–15.

https://doi.org/10.1186/S13100-019-0185-0.

[98] Erlandsson R, Wilson JF, Pääd☐bo S. Sex Chromosomal Transposable Element

Accumulation and Male-Driven Substitutional Evolution in Humans. Mol Biol Evol

2000;17:804–12. https://doi.org/10.1093/OXFORDJOURNALS.MOLBEV.A026359.

[99] Chaturvedi P, Hengeveld R, Zechel MA, Lee-Chan E, Singh B. The functional role of

class II-associated invariant chain peptide (CLIP) in its ability to variably modulate

immune responses. Int Immunol 2000;12:757–65.

https://doi.org/10.1093/INTIMM/12.6.757.

[100] Minias P, Bateson ZW, Whittingham LA, Johnson JA, Oyler-Mccance S, Dunn PO.

Contrasting evolutionary histories of MHC class I and class II loci in grouse—effects

of selection and gene conversion. Heredity (Edinb) 2016;116:466.

1076    https://doi.org/10.1038/HDY.2016.6.

1077 [101] Eizaguirre C, Lenz TL, Kalbe M, Milinski M. Rapid and adaptive evolution of MHC

1078    genes under parasite selection in experimental vertebrate populations. Nat Commun

1079    2012 31 2012;3:1–6. https://doi.org/10.1038/ncomms1632.

1080 [102] Parfrey LW, Lahr DJG, Katz LA. The dynamic nature of eukaryotic genomes. Mol Biol

1081    Evol 2008;25:787–94. https://doi.org/10.1093/molbev/msn032.

1082 [103] Sayers EW, Agarwala R, Bolton EE, Brister JR, Canese K, Clark K, et al. Database

1083    resources of the National Center for Biotechnology Information. Nucleic Acids Res

1084    2014;42:D7-17. https://doi.org/10.1093/nar/gky1069.

1085 [104] Baril T, Imrie R, Hayward A. Earl Grey v1.2 2021.

1086    https://doi.org/10.5281/ZENODO.5718734.

1087 [105] Smit AFA, Hubley RR, Green PR. RepeatMasker Open-4.0. Http://RepeatmaskerOrg

1088    2013.

1089 [106] Smit A, Hubley R. RepeatModeler Open-1.0. Http://RepeatmaskerOrg 2015.

1090 [107] Bao Z, Eddy SR. Automated De Novo Identification of Repeat Sequence Families in

1091    Sequenced Genomes. Genome Res 2002;12:1269–76.

1092    https://doi.org/10.1101/gr.88502.

1093 [108] Price AL, Jones NC, Pevzner PA. De novo identification of repeat families in large

1094    genomes. Bioinformatics 2005;21:i351–8.

1095 [109] Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al.

1096    BLAST+: Architecture and applications. BMC Bioinformatics 2009;10:1–9.

1097    https://doi.org/10.1186/1471-2105-10-421.

1098 [110] Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:

1099    Improvements in performance and usability. Mol Biol Evol 2013;30:772–80.

https://doi.org/10.1093/molbev/mst010.

[111] Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: A tool for automated

alignment trimming in large-scale phylogenetic analyses. Bioinformatics

2009;25:1972–3. https://doi.org/10.1093/bioinformatics/btp348.

[112] Rice P, Longden L, Bleasby A. EMBOSS: The European Molecular Biology Open

Software Suite. Trends Genet 2000;16:276–7. https://doi.org/10.1016/S0168-

9525(00)02024-2.

[113] Goubert C, Craig RJ, Bilat AF, Peona V, Vogan AA, Protasio A V. A beginner's guide

to manual curation of transposable elements. Mob DNA (In Press 2022:1–98.

[114] Larsson A. AliView: a fast and lightweight alignment viewer and editor for large

datasets. Bioinformatics 2014;30:3276–8.

[115] Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-

generation sequencing data. Bioinformatics 2012;28:3150–2.

[116] Wheeler TJ, Eddy SR. nhmmer: DNA homology search with profile HMMs.

Bioinformatics 2013;29:2487–9. https://doi.org/10.1093/BIOINFORMATICS/BTT403.

[117] Xu Z, Wang H. LTR_FINDER: an efficient tool for the prediction of full-length LTR

retrotransposons. Nucleic Acids Res 2007;35:W265–8.

https://doi.org/10.1093/nar/gkm286.

[118] Ou S, Jiang N. LTR_FINDER_parallel: parallelization of LTR_FINDER enabling rapid

identification of long terminal repeat retrotransposons. BioRxiv 2019:2–6.

[119] Wong WY, Simakov O. RepeatCraft: a meta-pipeline for repetitive element de-

fragmentation and annotation. Bioinformatics 2018;35:1051–2.

https://doi.org/10.1093/bioinformatics/bty745.

[120] Rubino F, Creevey CJ. MGkit: Metagenomic framework for the study of microbial

communities 2014.

[121]   Edgar RC. MUSCLE: Multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res 2004;32:1792–7. https://doi.org/10.1093/nar/gkh340.

[122]   Lenhard B, Sandelin A, Carninci P. Metazoan promoters: Emerging characteristics

and insights into transcriptional regulation. Nat Rev Genet 2012;13:233–45.

https://doi.org/10.1038/nrg3163.

[123]   Quinlan AR, Hall IM. BEDTools: A flexible suite of utilities for comparing genomic

features. Bioinformatics 2010;26:841–2. https://doi.org/10.1093/bioinformatics/btq033.

[124]   Makowski D, Ben-Shachar M, Lüdecke D. bayestestR: Describing Effects and their

Uncertainty, Existence and Significance within the Bayesian Framework. J Open

Source Softw 2019;4:1541. https://doi.org/10.21105/joss.01541.

[125]   Gel B, Serra E. karyoploteR: an R/Bioconductor package to plot customizable

genomes displaying arbitrary data. Bioinformatics 2017;33:3088–90.

https://doi.org/10.1093/bioinformatics/btx346.

[126]   Kimura M. A simple method for estimating evolutionary rates of base substitutions

through comparative studies of nucleotide sequences. J Mol Evol 1980;16:111–20.

[127]   Pfeiler E, Nazario-Yepiz NO, Pérez-Gálvez F, Chávez-Mora CA, Laclette MRL,

Rendón-Salinas E, et al. Population Genetics of Overwintering Monarch Butterflies,

Danaus plexippus (Linnaeus), from Central Mexico Inferred from Mitochondrial DNA

and Microsatellite Markers. J Hered 2017;108:163–75.

https://doi.org/10.1093/jhered/esw071.

[128]   Wickham H. ggplot2: elegant graphics for data analysis. Springer; 2016.

[129]   Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, et al. Welcome

to the Tidyverse. J Open Source Softw 2019;4:1686.

[130]   Marchler-Bauer A, Bo Y, Han L, He J, Lanczycki CJ, Lu S, et al. CDD/SPARCLE:

1149           functional classification of proteins via subfamily domain architectures. Nucleic Acids

1150           Res 2017;45:D200–3. https://doi.org/10.1093/NAR/GKW1129.

1151     [131]   Evgen'ev MB, Zelentsova H, Shostak N, Kozitsina M, Barskyi V, Lankenau DH, et al.

1152           Penelope, a new family of transposable elements and its possible role in hybrid

1153           dysgenesis in Drosophila□virilis. Proc Natl Acad Sci 1997;94:196–201.

1154           https://doi.org/10.1073/PNAS.94.1.196.

1155     [132]   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective

1156           Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol Evol

1157           2015;32:268–74. https://doi.org/10.1093/MOLBEV/MSU300.

1158     [133]   Kalyaanamoorthy S, Minh BQ, Wong TKF, Haeseler A von, Jermiin LS. ModelFinder:

1159           fast model selection for accurate phylogenetic estimates. Nat Methods 2017 146

1160           2017;14:587–9. https://doi.org/10.1038/nmeth.4285.

1161     [134]   Team RC. R: A language and environment for statistical computing 2013.

1162

## 1163   Declarations

1164   Ethics approval and consent to participate: Not applicable

1165

1166   Consent for Publication: Not applicable

1167

1168   Availability of Data and Materials: All data generated or analysed during this study are

1169   included in this article and its supplementary information files.

1170

1171    Competing Interests: The authors declare that they have no competing interests.

1172

1173    Funding: TB was supported by a studentship from the Biotechnology and Biological

1174    Sciences Research Council-funded South West Biosciences Doctoral Training Partnership

1175    (BB/M009122/1). AH was supported  by  a  Biotechnology  and  Biological  Sciences

1176    Research  Council  (BBSRC)  David  Phillips  Fellowship  (BB/N020146/1).

1177

1178    Authors' Contributions: TB developed the pipelines and scripts for automated TE annotation

1179    and downstream analyses, produced the figures, and drafted the manuscript. AH conceived

1180    and coordinated the study and participated in writing the manuscript. All authors read and

1181    approved the final manuscript.

1182

1183    Acknowledgements: We would like to thank Dr James Galbraith, Ryan Biscocho, and Ryan

1184    Imrie for constructive comments on early drafts of this manuscript, and Dr Aurélie Kapusta

1185    for support in running the DNA deletions analysis.