Hippocampal-prefrontal theta coupling develops as mice become proficient in

associative odorant discrimination learning

Daniel Ramirez-Gordillo[1], Andrew A. Parra[1], K. Ulrich Bayer[2,3] and Diego Restrepo[1,2,#]

[1]Department of Cell and Developmental Biology, University of Colorado Anschutz Medical

Campus, Aurora, CO 80045, USA

[2]Program in Neuroscience, University of Colorado Anschutz Medical Campus, Aurora, CO

80045, USA

[3]Department of Pharmacology, University of Colorado Anschutz Medical Campus, Aurora, CO

80045, USA

[#]Corresponding author: Diego Restrepo, diego.restrepo@cuanschutz.edu

Abstract

Learning and memory requires coordinated activity between different regions of the brain. Here we studied the interaction between medial prefrontal cortex (mPFC) and hippocampal dorsal CA1 during associative odorant discrimination learning in the mouse. We found that as the animal learns to discriminate odorants in a go-no go task the coupling of high frequency neural oscillations to the phase of theta oscillations (phase-amplitude coupling or PAC) changes in a manner that results in divergence between rewarded and unrewarded odorant-elicited changes in the theta-phase referenced power (tPRP) for beta and gamma oscillations. In addition, in the proficient animal there was a decrease in the coordinated oscillatory activity between CA1 and mPFC in the presence of the unrewarded odorant. Furthermore, the changes in PAC resulted in a marked increase in the accuracy for decoding odorant identity from tPRP when the animal became proficient. Finally, we studied the role of Ca^{2+}/calmodulin-dependent protein kinase II α (CaMKIIα), a protein involved in learning and memory, in oscillatory neural processing in this task. We find that the accuracy for decoding the odorant identity from tPRP decreases in CaMKIIα knockout mice and that this accuracy correlates with behavioral performance. These results implicate a role for PAC and CaMKIIα in olfactory go-no go associative learning in the hippocampal-prefrontal circuit.

Introduction

Learning and memory are two crucial abilities important for survival of organisms (Klein et al., 2002). Our lives are enhanced, and our personalities are shaped due to the ability to learn and form memories. Therefore, it is not surprising that often diseases that affect these abilities are devastating and frequently the individual affected becomes dependent on others. Learning and memory requires coordinate activity between different regions of the brain (Colgin, 2011; Gordon, 2011; Headley and Paré, 2017; Lisman et al., 2017). Local field potential oscillations reflect activity of temporally coordinated neuronal groups providing a reference for spike timing-based codes and gating information transfer between distant brain regions. In phase amplitude coupling (PAC), the amplitude of the bursts of a faster oscillation is larger within a phase window of a slower carrier wave (Tort et al., 2010). In a previous publication, we characterized PAC in the olfactory bulb of mice learning to discriminate odorants in a go-no go associative learning task and we introduced a measure of the magnitude of phase-locked high gamma and beta bursts as a function of time: the theta phase-referenced power (tPRP). We found that PAC changes throughout learning and that odorant-elicited changes in tPRP increase for rewarded and decrease for unrewarded odorants in the proficient mouse (Losacco et al., 2020). Furthermore, we showed that contextual odorant identity (is the odorant rewarded?) can be decoded from peak tPRP in animals proficient in odorant discrimination, but not in naïve mice. As the animal learns to discriminate the odorants, the dimensionality of PRP decreased. These findings in the olfactory bulb, the first relay station in the olfactory system, raised the question whether

downstream areas of the brain receiving input from the olfactory bulb experience a

similar phenomenon.

In the current study, we assessed decoding of odorant identity from oscillatory neural

activity in animals learning to discriminate odorants in the go-no go task in two

downstream brain areas of the brain that receive input from the olfactory bulb: the

dorsal cornu ammonis area 1 (CA1) area of the hippocampus and medial prefrontal

cortex (mPFC). These brain areas are involved in learning of odorant valence and

attention to odorants (Cansler et al., 2021; Gourevitch et al., 2010; Li et al., 2017; Martin

et al., 2007) and exhibit oscillatory coupling disrupted in animal models of schizophrenia

(Colgin, 2011). Additionally, we chose to study decoding in Ca^{2+}/calmodulin-dependent

protein kinase II α (CaMKIIα) knockout mice (CaMKIIα KO) because of the essential

role of this protein in learning and memory in the hippocampus (Bayer and Schulman,

2019; Bear et al., 2018; Lisman et al., 2012). CaMKIIα plays a role in long term

potentiation (LTP) and long-term depression (LTD) and dentate gyrus neurogenesis,

essential processes for learning and memory (Cook et al., 2021; Coultrap et al., 2014;

Malinow et al., 1989; Silva et al., 1992b; Suarez-Pereira et al., 2015). Deficiencies in the

function of CaMKIIα have been implicated in a range of diseases including

schizophrenia, addiction, depression, epilepsy, and neurodevelopmental disorders

(Chia et al., 2018; Robison, 2014). CaMKIIα KO are viable and display impaired spatial

memory and reduced LTP in the hippocampus (Silva et al., 1992a; Silva et al., 1992b).

Mice heterozygous for CaMKIIα knockout (CaMKIIα Hets) have problems with working

memory and increased anxiety, and aggressiveness, characteristic of schizophrenia

(Chen et al., 1994; Hasegawa et al., 2009; Matsuo et al., 2009; Yamasaki et al., 2008).

Large genome screens have found significant heterozygous mutations in CaMKIIα in

schizophrenia patients (Fromer et al., 2014; Purcell et al., 2014). Furthermore, because

of the role of CaMKIIα in development and neurogenesis, CaMKIIα Het mice have an

immature dentate gyrus (Yamasaki et al., 2008). The undeveloped brain leads to

dysfunctions in the neuronal circuit leading to an altered perception of reality and

cognitive dysfunctions (Walton et al., 2012).

We recorded neural activity from both hippocampal dorsal CA1 and mPFC in mice

engaged in a go-no go task where they learn to respond to a rewarded odorant and

refrain from responding to the unrewarded odorant (Losacco et al., 2020). We surveyed

PAC changes throughout learning by calculating the modulation index (MI), a measure

of PAC strength, and the variance of peak angle, a measure of the variance of the theta

phase for gamma oscillations (Losacco et al., 2020; Tort et al., 2010). We show that the

MI and peak angle variance change as the animal becomes proficient and we find a

difference in these measures of PAC between wild type mice (WT) and CaMKIIα

knockout (KO) and heterozygous (Het) mice. In addition, the odorant-elicited changes in

tPRP decrease for unrewarded odorants as the animal learns and tPRP differs between

CaMKIIα KO and WT. Furthermore, the accuracy for decoding contextual odorant

identity from tPRP increased as the animal became proficient and this increase was

smaller for CaMKIIα KO mice. Finally, we show an increase in the odorant-induced

change in coordinated neural activity between CA1 and mPFC as the animal becomes

proficient at discriminating between odorants and the odorant-induced change in

coordinated neural activity is smaller in CaMKIIα KO mice. These results demonstrate

increased decoding accuracy from phase-referenced oscillatory power in CA1 and

mPFC as the animal learns to differentiate odorants and a role of CaMKIIα in learning-

induced changes in PAC and coordinated oscillatory neural activity between

hippocampus and mPFC.

Results

Dual CA1-mPFC tetrode recording in mice undergoing the go-no go olfactory discriminating task.

The goal of this study was to determine whether changes in coupled oscillations that occur in the olfactory bulb as mice learn to discriminate odorants in a go-no go task are observed in downstream areas of the brain (mPFC and hippocampus). These downstream areas receive input originating from the olfactory bulb and oscillations in the bulb are known to be coupled to the hippocampus and throughout the brain (Martin et al., 2007; Nguyen et al., 2016; Tort et al., 2018). We also wanted to determine whether there was a difference between the CaMKIIα KO, CaMKIIα Hets and the WT mice in PAC and whether the peak or trough of the oscillation carries different information allowing the mouse to learn to differentiate between a pair of odorants. Lastly, we determined whether coordinated neural activity changed as the animal learned to discriminate odorants and we asked whether there was a difference in the relationship of oscillations between CA1 and mPFC between the different CaMKIIα genotypes. Thirsty mice learned to lick on a spout in the presence of a rewarded odorant (S+) to obtain a water reward and refrained from licking in the presence of unrewarded odorant (S-) in a go-no go associative learning task (Losacco et al., 2020). The odorants were presented in pseudorandomized order in the go no-go task (Figure 1A–D). The odorant pairs tested were different volatile compounds (Table 1) and we found no difference in any measurements between odorant pairs.

In the go-no-go odorant discrimination task, mice start the trial at will by licking on the

lick port. The odorant is delivered at a random time 1–1.5 s after nose poke (the time

course for the trial is shown in Figure 1C). The mice learn to either lick a waterspout at

least once during each 0.5 s segment in the 2 s response area to obtain a water reward

for the rewarded odorant or refrain from licking for the unrewarded odorant. Mice refrain

from licking during presentation of the unrewarded odorant because of the effort it takes

to lick. Behavioral performance was termed naïve or proficient when their performance

estimated in a 20-trial window was below 65% for naïve and above 80% for proficient.

Mice were trained in the task for sessions of up to 200 trails. In the last session training

ended after the animal achieved two 20-trial blocks of proficient performance; valence

was reversed the next day or another odorant was tested. We recorded the LFP using 2

tetrodes (4 electrodes per tetrode) implanted in the CA1 of the hippocampus and 2

tetrodes in ipsilateral mPFC, and we analyzed the data to determine whether the

different genotypes differ in cross-frequency coupling for naïve or proficient mice (Figure

1E). The dataset is comprised of 747 recording sessions in 18 mice. The odorant pairs

tested were different volatile compounds (or mixtures) whose nomenclature addresses

the odorant names and the experimental set (e.g., APEB, see Table 1 for the

nomenclature). Table 2 enumerates the total number of sessions per odorant pair,

mouse, and experiment.

We performed PAC analysis of the LFP recorded in the go no-go behavioral task. PAC

analysis is a cross-frequency coupling analysis to determine whether high frequency

oscillation bursts take place at specific phases of low frequency theta oscillations (Tort

et al., 2010). PAC has been reported in brain areas such as the olfactory bulb,

hippocampus and prefrontal cortex (Belluscio et al., 2012; Colgin, 2015; Kaplan et al.,

2014; Losacco et al., 2020; Rojas-Libano et al., 2014; Scheffer-Teixeira and Tort, 2016).

Figures 1E i–iv show an example of high gamma/theta PAC for the LFP recorded in

CA1. Figure 1E i shows the extracellular LFP sampled for hippocampus at 20 kHz and

filtered between 1–750 Hz. The raw signal (Figure 1 E i) was filtered with a 20th order

Butterworth filter into different LFP frequency bands (Figures 1E ii, iii, iv) (theta, 6–14

Hz, adapted from (Nguyen Chi et al., 2016) high gamma, 65–95 Hz). We used PAC

analysis (Tort et al., 2010) to evaluate the degree of coupling of the amplitude of the

envelope of the beta or high gamma LFP on the phase of the theta LFP. Figure 1E iii

shows the theta phase and in Figure 1 E iv shows the envelope for the amplitude of the

high gamma LFP, both calculated with the Hilbert transform as detailed in Tort et al.

(2010). Figures 1 E ii and iv show that the filtered high gamma LFP changes amplitude

in manner that appears coordinated with the theta phase.

The peak angle variance of PAC increased for the unrewarded odorant as the

animals became proficient differentiating between odorants

We proceeded to ask whether the strength of PAC, quantified by the modulation index

(MI), changes as the animal learns to differentiate odorants in the go-no go task. MI is a

measure of how localized high frequency firing is within the phase of theta oscillations

(Tort et al 2010). Figures 2A–C illustrate an example of high gamma tPAC for S+ and S-

odorant trials in two sessions for naïve and proficient mice. The phase amplitude plots

for the trials are shown in pseudocolor in Figure 2A, for a WT mouse during the naïve

(2Ai-iii) and proficient (2Aii-iv) stages and the percent correct as a function of trial

number is shown for the two sessions in Figure 2B. In this example there appears to be

an increase in the strength of PAC when the animal becomes proficient differentiating

between the odorants. Furthermore, there was an increase in peak angle variance for

the unrewarded odorant as shown by the peak angle polar histograms in Figures 2C i, ii.

Figure 2C i shows that for the unrewarded odorant the peak angle was near 90 degrees

during the naïve stage while Figure 2C ii shows the peak angle for the unrewarded

odorant fluctuated widely during the proficient stage. This is in contrast with the

rewarded odorant for which the peak angle remained near 90 degrees during naïve

(Figure 2C iii) and proficient stages (Figure 2C iv).

Figure 2D shows the differences in MI computed per mouse, per odorant pair between

S+ and S- for naïve and proficient mice for tPAC for hippocampus (i) beta (ii) high

gamma and mPFC (iii) beta (iv) high gamma. For the beta tPAC MI for both the

hippocampus and mPFC a generalized linear model (GLM) analysis finds statistically

significant differences between S+ vs. S- and the interaction between S+ vs. S- and

naïve vs. proficient (p<0.001, 188 observations, 184 d.f., F-statistic=51,67.7, p<0.001, 6

mice, 8 odor pairs, Supplementary Table 1). GLM does not find significant differences

for MI for high gamma tPAC (p>0.05). Figure 2E shows differences for peak angle

variance between S+ and S- for naïve and proficient mice for hippocampus tPAC (i) for

beta and (ii) high gamma and for mPFC tPAC for (iii) beta and (iv) high gamma. For

both beta and gamma tPAC and for both the hippocampus and mPFC GLM finds

statistically significant differences between S+ vs. S- and the interaction between S+ vs.

S- and naïve vs. proficient (p<0.001, 188 observations, 184 d.f., F-statistic=19.5-26.4,

p<0.001, 6 mice, 8 odor pairs, Supplementary Table 2) and for naïve vs. proficient

(p<0.05). Asterisks in all figures show pairwise statistical significance (t test or ranksum

test, p< pFDR, p value for significance corrected for multiple comparisons using the

false discovery rate (Curran-Everett, 2000)).

Overall, we found an increase in peak angle variance for the unrewarded odorant as the

animal became proficient. We also found small, but significant changes in the strength

of beta tPAC that differed between the S+ and S- odorants as the animal learned to

discriminate odorants.

The odorant-elicited change in the theta phase-referenced beta and gamma power

became negative for the unrewarded odorant as the mice became proficient

discriminating between odorants

Wavelet power referenced to theta phase (peak or trough) was determined to evaluate

whether it changes as the animal learns. This analysis is referred to as theta phase-

referenced power (tPRP) (Losacco et al., 2020). Figure 3A shows examples of the time

course during the trial for the average wavelet broadband LFP spectrograms for 30

trials during naïve S+ (i), 27 trials during naïve S- (ii) 84 trials during proficient S+ (iii)

and 84 trials during proficient S- (iv). After odorant onset there was an increase in

broadband power for the S+ odorant and a decrease in power for the S- odorant as the

animal became proficient.

Figure 3B shows the time course during the trial for the average high gamma tPRP

referenced to the peak (red) or trough (blue) of theta for the proficient trials in Figure 3A.

For the rewarded (S+) odorant the power increased for the peak and decreased for

trough after addition of the odorant while for the unrewarded odorant both peak and

trough tPRPs decreased. Figure 3C shows that, as expected, there was an increase in

the lick rate for the rewarded odorant and a decrease for the unrewarded odorant.

Finally, Figures 3D,E show a per mouse per odorant pair analysis that indicated that

tPRP became negative for S- as the mice became proficient. E: peak, F: through, (i)

beta and (ii) gamma hippocampus (iii) beta and (iv) gamma mPFC. GLM finds

statistically significant differences for tPRP between S+ vs. S- and the interaction

between S+ vs. S- and naïve vs. proficient (p<0.001, 376 observations, 368 d.f., F-

statistic=72.7-103, p<0.001, 6 mice, 8 odor pairs, Supplementary Table 3).

In addition, we asked whether odorant-induced changes in peak tPRP changed when

odorant valence was reversed. Panels A i and ii of Figure 3 - figure supplement 1 show

an example of the time course for the peak (red) and trough (blue) high gamma tPRP

when the mouse was proficient for a forward session where the rewarded odorant (S+)

was acetophenone (AP) and the unrewarded odorant (S-) was ethyl benzoate (EB).

Panels A iii and iv of Figure 3 figure supplement 1 show that when the valence of the

odorant was reversed (AP was S- and EB was S+) the response to EB resembled the

response to AP in the forward sessions (compare panels A ii and iv of Figure 3 figure

supplement 1) indicating that the response is a response to the contextual identity of the

odorant (is it rewarded?) as opposed to the chemical identity of the odorant. Figure 3

figure supplement 1 (B,C) shows a summary bar graph/violin plot analysis per mouse

per odorant pair of all reversal experiments indicating that for the proficient mouse the

average peak and trough tPRP decreases for the unrewarded S- odorant regardless of

the identity of the odorant. For high gamma tPRP GLM finds statistically significant

differences for S+ vs. S- and peak vs. trough (p<0.001, 188 observations, 180 d.f., F-

statistic=61.8, p<0.001, 6 mice, 8 odor pairs, Supplementary Table 4) and does not find

a difference between forward and reverse sessions indicating that indeed the high

gamma tPRP responds to the contextual odorant identity. For beta tPRP GLM finds a

statistically significant difference for S+ vs. S- (p<0.001, 188 observations, 180 d.f., F-

statistic=87, p<0.001, 6 mice, 8 odor pairs, Supplementary Table 4) and interestingly,

GLM does find a small statistically significant differences between forward vs. reverse

(p<0.05) for beta tPRP. Although this is a relatively small difference, this indicates that

beta tPRP is not exclusively responsive to contextual odorant identity and that these

brain regions may also encode information on the chemical identity of the odorant.

The accuracy for decoding the contextual identity of the odorants from theta

phase referenced power increased when the mice became proficient

We proceeded to determine whether we could decode contextual odorant identity from

tPRP. Decoding was performed using a linear discriminant analysis (LDA) to set a

decision boundary hyper-plane between binary stimulus classes (S+ versus S-) (Vizcay

et al., 2015). LDA was trained with tPRP from each electrode for each mouse (8

electrodes in the hippocampus and 8 electrodes in the mPFC per mouse) for all trials

except one (the training dataset) and then the tPRP from the missing trial (test data)

was classified as S+ or S-. This was done at both naïve and proficient learning stages.

As a control we shuffled the identity of trials in the training set.

Figure 4A shows an example of the time course during the trial for the decoding

accuracy for one mouse for the LDA trained using CA1 tPRP for the EAPA odor pair for

(i) naïve stage beta, (ii) proficient stage beta, (iii) naïve stage gamma, (iv) proficient

stage gamma. For the naïve animal decoding accuracy increased slowly after the

addition of the odorant, and increased further after the animal received the reward

(Figures 4Ai,iii). When the animal became proficient decoding accuracy increased

rapidly beyond 80% after addition of the odorant (Figures 4Aii,iv). For high gamma tPRP

the accuracy was higher for the proficient animal for peak-referenced tPRP compared to

trough-referenced tPRP (Figure 4Aiv). Figures 4 B and C show mean bar graphs and

violin plots for the mean accuracy for decoding contextual odorant identity calculated for

the last second of the odorant epoch (1.5 to 2.5 sec after diverting the odorant towards

the mouse) for shuffled, naïve, and proficient. Figure 4B shows the accuracy for the

peak tPRP and Figure 4C shows the accuracy for trough tPRP. For all conditions the

accuracy is significantly higher for the proficient stage compared to naïve or shuffled.

GLM analysis finds for both CA1 and mPFC statistically significant differences between

naïve vs. proficient and shuffled vs. proficient for both beta and gamma tPRP (p<0.001,

756 observations, 744 d.f., F-statistic=25.1, p<0.001, 6 mice, 8 odor pairs,

Supplementary Table 5) and for gamma tPRP GLM finds statistically significant

differences between peak and trough (p<0.001, 756 observations, 744 d.f., F-

statistic=25.2, p<0.001, 6 mice, 8 odor pairs, Supplementary Table 5).

Coordinated hippocampal-prefrontal neural activity decreased for the unrewarded

odorant as the animal became proficient.

Coordinated hippocampal-prefrontal neural activity supports the organization of brain

rhythms and is present during a range of cognitive functions presumably underlying

transfer of information between these two brain regions (Colgin, 2011; Gordon, 2011;

Headley and Paré, 2017; Lisman et al., 2017). We proceeded to determine whether

there were changes in coordinated neural activity between dorsal CA1 and mPFC as

the animal learned to discriminate the odorants. We used two complementary methods

to quantify coordinated neural activity by calculating coherence and phase locking value

(PLV) (Borjigin et al., 2013; Namburi, 2021).

Figure 5A shows an example of a pseudocolor plot for the time course for coherence for

a session where a mouse was engaged in the go-no go task. The pseudocolor plots

show the average coherence time course per trial for (i) S+ naïve (ii) S- naïve (iii) S+

proficient (iv) S- proficient. For S- the pseudocolor plot shows that a decrease in

coherence elicited after the odor onset develops as the mouse becomes proficient

(compare Figures 5Aii and iv). Figure 5Bi shows the change of coherence (Δ

coherence) elicited by the odorants as a function of frequency for naïve and proficient

mice calculated as the mean per odorant pair per mouse. Δ coherence is defined as the

mean coherence in the window from 1.5 to 2.5 sec after the odorant onset minus the

mean coherence for -1.5 to 0 sec. We found that the Δ coherence became negative for

the unrewarded odorant as the animal became proficient discriminating between

odorants. Figures 5Aii to iv show bar graphs/violin plots for per mouse per odorant pair

for Δ coherence for the different bandwidths: (ii) theta, (iii) beta, (iv) gamma. A GLM

analysis finds for all bandwidths statistically significant differences for Δ coherence for

S+ vs. S- and the interaction between S+ vs. S- and naïve vs. proficient ($p < 0.001$, 190

observations, 186 d.f., F-statistic=21.6-32.4, $p < 0.001$, 6 mice, 8 odor pairs,

Supplementary Table 6) and for theta the GLM analysis finds additionally a significant

difference between naïve and proficient ($p < 0.05$).

Figure 5C shows the odor-elicited change in average PLV (Δ PLV) per mouse per odor

pair for (i) theta, (ii) beta, (iii) gamma. Here we also found a negative Δ PLV for the

unrewarded odorant for the proficient mice. For Δ PLV GLM finds statistically significant

differences for Δ PLV between naïve vs. proficient ($p < 0.05$), S+ vs. S- and the

interaction between S+ vs. S- and naïve vs. proficient ($p < 0.001$, 192 observations, 188

d.f., F-statistic=29.5-46.5, $p < 0.001$, 6 mice, 8 odor pairs, Supplementary Table 7).

Finally, Figure 5D shows a strong correlation between Δ coherence and Δ PLV for (i)

theta, (ii) beta, (iii) gamma. In conclusion, both the Δ coherence and Δ PLV measures

indicate that as the mouse becomes proficient there was a decrease in coordinated

hippocampal-prefrontal neural activity for the unrewarded odorant.

The strength of PAC and peak angle variance differed between the CaMKIIα

genotypes.

Since CaMKIIα is a protein involved in LTP, while PAC plays a role in learning and

memory in hippocampal space tasks we asked whether PAC and tPRP differed

between the different CaMKIIα genotypes (WT, CaMKIIα Het and CaMKIIα KO). We

trained mice from the three genotypes in the go-no go task. There was no difference in

the number of sessions to criterion (Figure 6 - Figure supplement 1 A, GLM p>0.05, 24

observations, 21 d.f., F-statistic=0.26, p>0.05, 8 odor pairs, Supplementary Table 8).

However, for proficient mice the percent correct was higher for WT compared to both

CaMKIIα Het and CaMKIIα KO (Figure 6 - figure supplement 1 B, GLM p<0.05, 137

observations, 134 d.f., F-statistic=3.7, p<0.05, 6 mice, 8 odor pairs, Supplementary

Table 8) and the intertrial interval was larger for CaMKIIα Het (Figure 6 - figure

supplement 1 B, p<0.001, 137 observations, 134 d.f., F-statistic=13.5, p<0.001, 6 mice,

8 odor pairs, Supplementary Table 8).

We then asked whether tPAC in the hippocampus and mPFC differs between the

different genotypes. When we compared strength of tPAC, measured by the modulation

index (MI), and the peak angle variance between genotypes (WT, CaMKIIα Het and

CaMKIIα KO), we found significant differences. Figure 6 - figure supplement 2 A, B

shows the average MI per mouse per odorant pair for beta (Ai), high gamma (Bi) for

hippocampus and for beta (Aii) and gamma (Bii) for prefrontal cortex. For beta tPAC

GLM finds a statistically significant difference for MI for the interaction between WT vs.

CaMKIIα KO and S+ vs. S- in both the hippocampus and mPFC (p<0.001, 544

observations, 532 d.f., F-statistic=10.3-34.3, p<0.001, 6 mice, 8 odor pairs,

Supplementary Table 9). For high gamma tPAC there was an increase in MI for

CaMKIIα Het and a decrease in MI for CaMKIIα KO compared to WT and GLM analysis

finds a statistically significant difference between WT vs. CaMKIIα KO and WT vs.

CaMKIIα Het for both hippocampus and mPFC (p<0.001, 544 observations, 532 d.f., F-

statistic=10.3-34.3, p<0.001, 6 mice, 8 odor pairs, Supplementary Table 9).

Figure supplement 2C, D shows average peak angle variance per mouse per odorant

pair for beta (Ci), high gamma (Ei) in the hippocampus and (Cii) beta, (Eii) high gamma

in prefrontal cortex. For proficient mice the peak angle variance for the rewarded

odorant decreased for CaMKα Het and increased for CaMKα KO. GLM analysis found a

statistically significant difference for WT vs. CaMKIIα Het, WT vs. CaMKIIα KO, naïve

vs. proficient and S+ vs. S- for theta/beta and theta/gamma for both the hippocampus

and mPFC (p<0.05, 544 observations, 532 d.f., F-statistic=20.6, p<0.001, 6 mice, 8 odor

pairs, Supplementary Table 10).

The accuracy for decoding the contextual identity of the odorant decreased in the

CaMKIIα knockout mouse and was correlated with percent correct discrimination.

The differences in PAC between CaMKIIα genotypes raises the question whether

decoding of contextual identity from tPRP is altered in the CaMKIIα KO and the

CaMKIIα Het mice. Figure 6 shows the results of our comparison of decoding accuracy

between genotypes. Figures 6A and B. show examples for proficient mice of the time

course for the accuracy of decoding of odorant contextual identity by LDA trained using

tPRP for the EAPA odor pair (Figure 6A theta/beta tPRP, Figure 6B theta/gamma tPRP,

i. WT, ii. Het, iii. KO). For the WT and CaMKIIα Het mice the accuracy increases

monotonously during the odorant epoch (Figures 6Ai,ii and 6Bi,ii) while for the CaMKIIα

KO mouse the accuracy reaches a maximum value after one second and then

decreases slightly for the rest of the odorant epoch (Figures 6Aiii and 6Biii). Figures 6C

and D show the differences in decoding accuracy between the different genotypes

assessed in the window from 1.5 to 2.5 sec after addition of the odorant. For both beta

and gamma peak or trough tPAC decoding for both the hippocampus and the mPFC

decoding accuracy was lowest for CaMKIIα KO. In addition, for gamma trough tPRP

decoding for mPFC the accuracy was highest for WT, and decreased for both CaMKIIα

Het and CaMKIIα KO mice (Figure 6Div). For beta tPRP LDA GLM found statistically

significant differences for decoding accuracy between WT and KO for all conditions

(p<0.001) and between WT and Het for peak gamma tPRP in the hippocampus, trough

gamma tPRP in mPFC (p<0.05, 139 observations, 136 d.f., F-statistic=9.9-34.3,

p<0.001, 6 mice, 8 odor pairs, Supplementary Table 11).

We then asked whether there was a relationship between contextual odorant identity

decoding accuracy and percent correct performance for proficient mice in the go-no go

task. Figures 6E and 6F show that there were statistically significant correlations

between decoding accuracy and percent correct performance for all the different

conditions. The correlation coefficients were as follows: 0.3 for hippocampal peak beta

tPRP, 0.24 for hippocampal peak gamma tPRP, 0.29 for mPFC peak beta tPRP, 0.23

for mPFC peak gamma tPRP, 0.3 for hippocampal trough beta tPRP, 0.36 for

hippocampal trough gamma tPRP, 0.29 for mPFC trough beta tPRP and 0.40 for mPFC

trough gamma tPRP, and the p value for significance of the correlation coefficient was

$p < 0.01$. This indicates that the decoding accuracy obtained with tPRP is related to

behavioral performance.

Coherent hippocampal-prefrontal neural activity differed between CaMKIIα

genotypes

Finally, we asked whether there were differences for coordinated hippocampal-

prefrontal neural activity for the different CaMKIIα genotypes. Figure 7A shows that for

the proficient mice the absolute value of the change of coherence (Δ coherence) elicited

by the odorants tended to be smaller for the CaMKIIα KO. For Δ coherence GLM

analysis found statistically significant differences for WT vs. KO (theta $p < 0.05$, beta and

gamma $p < 0.001$, 546 observations, 534 d.f., F-statistic=21.2-33.8, $p < 0.001$, 6 mice, 8

odor pairs, Supplementary Table 12). Δ PLV showed the same trends (Figure 7B), but

the GLM found no statistically significant differences between WT and KO ($p > 0.05$, 512

observations, 500 d.f., F-statistic=25.5-39.9, $p < 0.001$, 6 mice, 8 odor pairs,

Supplementary Table 13).

Discussion

Santiago Ramon y Cajal described the hippocampus as a quaternary region of the olfactory system with profuse olfactory input through entorhinal cortex (Cajal, 1904), but subsequent studies showed that the hippocampus is involved in neural processing for tasks independent of olfaction such as spatial learning and memory in the Morris water maze (Nakazawa et al., 2004). However, the hippocampus does play a role in olfactory learning. Indeed, recent studies implicate a role for olfactory input through the lateral entorhinal cortex to calbindin-expressing pyramidal cells in the dorsal hippocampal CA1 for neural processing of decision-making in the go-no go olfactory discrimination task and these investigators show that these pyramidal cells develop more selective spiking responses to odorants as the animal becomes proficient (Li et al., 2017). Furthermore, studies of oscillatory neural activity implicate directional coupling from the olfactory bulb to the hippocampus in go-no go olfactory discrimination learning (Gourevitch et al., 2010; Martin et al., 2007). In addition, Granger directionality analysis finds that distinct low frequency oscillation bandwidths link the olfactory bulb and hippocampus in a bi-directional manner (Nguyen Chi et al., 2016). Interestingly, mPFC has also been proposed as a downstream information processing brain area coupled with the olfactory bulb. Coupling of low frequency oscillations between the olfactory bulb and mPFC increases during freezing periods in auditory conditioned fear learning (Moberly et al., 2018) and at rest there is strong coupling of low frequency oscillations for both olfactory bulb-hippocampus and olfactory bulb-mPFC (Mofleh and Kocsis, 2021). Furthermore, a recent study found that beta and theta synchrony between mPFC and olfactory regions

was elevated as rats switched their attention to odors (Cansler et al., 2021). These

findings indicate that the hippocampus and mPFC are coupled to the olfactory bulb

through neural oscillatory activity and that these circuits play a role in learning and

attention. Taken together with the fact that in a previous publication we showed that

when mice become proficient in discriminating odorants in the go-no go task contextual

odorant identity can be decoded from beta and gamma tPRP (Losacco et al., 2020)

these findings raise the question whether coupled oscillations in mPFC and the

hippocampus play a role in olfactory discrimination in go-no go associative learning.

In this study we found that for both the dorsal CA1 and mPFC as animals became

proficient in the go-no go task there was an increase in the variance of the peak angle

for beta and high gamma tPAC for the unrewarded odorant (Figure 2) accompanied by

a decrease in tPRP for this odorant (Figure 3). This decrease in tPRP for S- was

accompanied by a small increase in tPRP for the rewarded odorant resulting in a sharp

increase in accuracy for decoding of the contextual odorant identity as the animal

became proficient differentiating between odorants (Figure 4). Likewise, decreases in

coordinated prefrontal-hippocampal activity elicited by the unrewarded odorant became

larger suggesting uncoupling of oscillations in these two brain areas for the unrewarded

odorant. When we tested CaMKIIα KO mice, known to be deficient in spatial learning

and memory, we found a decrease in the accuracy of the decoding of contextual

odorant identity (Figure 6), and a decrease in the magnitude of the decrease in odorant-

elicited changes in coherence (Figure 8). Finally, the behavioral performance for

CaMKIIα KO and CaMKIIα Het mice was lower than the performance of WT mice

(Figure 6 – figure supplement 1) and the accuracy for decoding of contextual odorant

identity from tPAC correlated with behavioral performance (Figure 6). These findings

indicate that coordinated oscillatory neuronal activity in the hippocampal-mPFC axis

plays a role in learning in the go-no go task for olfactory discrimination.

Theta frequency stimulation, eliciting intrinsic gamma frequency oscillations, is known to

be essential for long-term potentiation a process that is thought to be crucial for learning

(Bliss and Lomo, 1973; Butler et al., 2016; Larson and Munkácsy, 2015) and theta PAC

of higher frequency bursts is thought to be important for information transfer within and

between brain regions which is thought to be essential for learning (Fries, 2005). In the

hippocampus high frequency oscillations at different phases of theta carry different

information. Indeed, Siegle and Wilson used closed loop optogenetics to stimulate

parvalbumin (PV) interneurons showing that when PV interneurons are stimulated at the

peak of theta mice increased performance in the encoding epoch for a spatial

navigation task whereas when they stimulated at the trough of theta mice increased

performance in the retrieval epoch (Siegle and Wilson, 2014). Interestingly, here we find

that as the animal becomes proficient the variance of the peak angle of PAC increases

substantially for the unrewarded odorant (Figure 2). As a result, if the gamma or beta

frequencies were being read by a downstream observer at a fixed angle, the information

conveyed by oscillations elicited by the rewarded odorant would be missed because of

the constantly changing theta phase angle. In contrast, the peak angle variance for the

rewarded odorant is small, and presumably this would result in more faithful reading of

this information. This is likely what underlies the increased accuracy for decoding the

contextual odorant identity from beta and gamma tPRP we found in the proficient

mouse (Figure 4). Thus, we found that in the go-no go task phase-referenced power

carries information on contextual odorant identity.

Full genetic knockout of CaMKIIα results in impairments in hippocampus-dependent

cognitive tasks and impaired LTP and LTD in adult mice (Coultrap et al., 2014; Silva et

al., 1992b). In contrast, although CaMKIIα Het mice share some behavioral deficits with

the full CaMKIIα KO including deficient learning and memory in the Morris water maze

and deficient retention of contextual fear conditioning (Frankland et al., 2001; Silva et

al., 1996), adult CaMKIIα Het mice have additional working memory deficits such as

repeat entry errors in the radial arms version of the Morris water maze (Matsuo et al.,

2009; Yamasaki et al., 2008). In addition, while young (p 12-16) CaMKIIα Het show LTP

of the same magnitude as WT mice in CA1, young adult (9-11 week old) mice display

impaired basal synaptic transmission, but do not have a deficit in LTP (Goodell et al.,

2016) suggesting the development of compensatory mechanisms for LTP in the Het.

PAC plays a role in hippocampal learning and memory (Siegle and Wilson, 2014)

raising the question whether the CaMKIIα Het and CaMKIIα KO have deficits in cross

frequency coupling. Here we performed to our knowledge the first study to determine

whether CaMKIIα-deficient mice display altered PAC. We find that the accuracy for

decoding of contextual odorant identity from beta and gamma tPRP is decreased in

CaMKIIα KO mice, but not in CaMKIIα Hets (with the exception of a decrease accuracy

for mPFC gamma trough tPRP decoding, Figure 6). Interestingly, the strength of PAC,

measured as the MI, was highest for gamma tPAC CaMKIIα Hets compared to both WT

and CaMKIIα KO (Figure 6 -Figure Supplement 1B). For CaMKIIα Hets the peak angle

variance decreased significantly for the rewarded odorant when the animal became

proficient (Figure 6 -Figure Supplement 1C,D). If a downstream neural observer is

evaluating contextual odorant identity by observing beta or gamma frequency bursts in

phase with theta oscillations these changes in PAC for CaMKIIα Hets would tend to

increase the ability to discriminate. Thus, the differences found for tPAC for CaMKIIα

Hets may be compensatory leading to no difference in decoding accuracy between WT

and CaMKIIα Hets (Figure 6). This would agree with the interpretation by Goodell et al.

who indicated that LTP deficits in CaMKIIα Hets are restored by compensatory changes

during development (Goodell et al., 2016). Finally, given that CaMKIIα Hets have a

phenotype reminiscent of schizophrenia it is interesting that resting-state gamma tPAC

has been found to be increased in patients with schizophrenia compared to that in

healthy controls (Won et al., 2018). These authors speculate that increased tPAC may

be related to the compensatory hyper-arousal patterns of the dysfunctional default-

mode network in schizophrenia.

Whether the PAC/tPRP changes found in CaMKIIα Het and CaMKIIα KO mice are due

to the decreased expression of CaMKIIα protein or to developmental changes in circuits

known to change in these mice such as in the dentate gyrus is an open question that

will require future studies with temporally and spatially restricted changes in CaMKIIα

activity. CaMKIIα may alter plasticity of pyramidal neurons in CA1 or mPFC.

Furthermore, a decrease in CaMKIIα expression may alter the the postsynaptic

regulation by CaMKIIα of inhibitory synapse transmission onto the pyramidal neurons

(Cook et al., 2021; Udakis et al., 2020). Finally, a subpopulation of granule cells that

play a role in odorant discrimination in the go/no go task are found in the olfactory bulb

and these cells are involved in generating the gamma frequency oscillatory activity in

the bulb (Malvaut et al., 2017). The olfactory bulb granule cells may be involved in

changing coordinated oscillations between the olfactory bulb and hippocampus in the

CaMKIIα KO/Het mice.

In conclusion we found that as the mouse learns to differentiate odorants in the go-no

go associative learning task there are changes in PAC that result in an increase of the

accuracy of decoding of the contextual odorant identity from tPRP. We found that as the

animal becomes proficient the coordinated neural activity between CA1 and mPFC

decreases for the unrewarded odorant. Finally, the accuracy of decoding the contextual

odorant identity from tPRP decreased in the CaMKIIα KO, but did not decrease in the

CaMKIIα Het and this decoding accuracy correlated with behavioral performance across

genotypes.

Materials and methods

Key Resources Table

REAGENT TYPE	REAGENT or RESOURCE	SOURCE	IDENTIFIER	ADDITIONAL INFORMATION
Chemical compound, drug	Isoamyl acetate	Sigma-Aldrich	Cat#123-92-2	Odorant
Chemical compound, drug	Ethyl acetate	Sigma-Aldrich	Cat# 141-78-6	Odorant
Chemical compound, drug	Propyl acetate	Sigma-Aldrich	Cat# 109-60-4	Odorant
Chemical compound, drug	Mineral oil	Sigma-Aldrich	Cat# 8042-47-5	Odorant
Chemical compound, drug	Acetophenone	Sigma-Aldrich	Cat# 98-86-2	Odorant
Chemical compound, drug	Ethyl benzoate	Sigma-Aldrich	Cat# 93-89-0	Odorant
Other	Nickel-chrome wire	Sandvik	Cat#PX000002	Tetrode fabrication
Other	16 Channel Electrode Interface Board	Neuralynx	EIB-16	Tetrode fabrication
Software, algorithm	Analysis code	This paper	Available on github and by request	https://github.com/restrepd/drgMaster
Strain, strain background	Mouse: CaMKIIα WT	Bayer Lab	Coultrap et al., 2014	Male mice
Strain, strain background	Mouse: CaMKIIα HET	Bayer Lab	Coultrap et al., 2014	Male mice
Strain, strain background	Mouse: CaMKIIα KO	Bayer Lab	Coultrap et al., 2014	Male mice
Software, algorithm	MATLAB_R2018a	Mathworks	RRID: SCR_001622	
Software, algorithm	Illustrator	Adobe	RRID: SCR_010279	
Software, algorithm	Photoshop	Adobe	RRID: SCR_014199	

Animals

Littermate mice with genotypes of CaMKIIα KO, Het and WT were obtained from

heterozygous breeding (Coultrap et al., 2014). Mice were housed in a vivarium with a

reversed light cycle of 14/10 h light/dark periods with lights on at 10:00 p.m. Food

(Teklad Global Rodent Diet no. 2918; Harlan) was available ad libitum. Access to water

was restricted during the behavioral sessions. However, if mice did not obtain ~1ml of

water during the behavioral session, additional water was provided in a dish in the cage

(Slotnick and Restrepo, 2005). All mice were weighed daily and received sufficient

water to maintain >80% of the weight before water restriction. All experiments were

performed according to protocols approved by the University of Colorado Anschutz

Medical Campus Institutional Animal Care and Use Committee.

Surgery and double tetrode implantation

Male mice 2-4 months of age were anesthetized by brief exposure to isoflurane (2.5%)

and subsequently anesthesia was maintained with an intraperitoneal injection of

ketamine (100 mg/kg) and xylazine (10 mg/kg). The tetrode drive included one optical

fiber ferrule for support of an EIB 8 board with 2 tetrodes with four polyamide-coated

nichrome wires (diameter 12.5 μm; Sandvik, gold plated to an impedance of 0.2–

0.4MΩ). Tetrodes were connected, and the optic fiber ferrule was glued through an EIB-

8 interface board (Neuralynx). Mice were implanted with two tetrode drives aimed at

mPFC (+1.94 mm anterior, +0.25 mm lateral, and -3.12 mm below bregma) (Eleore et

al., 2011) and the second tetrode drive at the right CA1 layer of the dorsal hippocampus

(-3.16 mm posterior, +3.2 mm lateral, and -2 mm below bregma) (Eleore et al.,

2011)(Fig. 1A). One ground screw was inserted 1 mm posterior from bregma and 1 mm

lateral to the midline and sealed to the bone with dental acrylic. Mice were allowed to

recover for 1 week before the initiation of the behavioral studies. All behavioral and LFP

recording experiments were performed with 2.5-8 month old mice that had undergone

double tetrode drive implantation.

Go-No-Go Behavioral Task

We used the methods previously described in (Losacco et al., 2020). Briefly, water-

restricted mice were required to enter an odor port and lick at the water spout to initiate

the release of the odorants 1–1.5 s after the first lick (Figs. 1B,C). Mice were required to

lick at least once in four 0.5 s intervals during reinforced odorant delivery (S+) to obtain

10 μl of water. When exposed to the unreinforced odorant (S–), mice refrain from licking

for 2 sec because of the effort it takes to lick for this period. Licking was detected by

closing a circuit between the licking spout and the grounded floor of the cage (Slotnick

and Restrepo, 2005). The performance of the mice was assessed by calculating correct

response to the S+ and S– odorants in 20 trial blocks where 10 S+ and 10 S– odorants

were presented at random. Mice were first trained to discriminate between 1% isoamyl

acetate (S+) and mineral oil (S-). Once the mice learned to discriminate between

isoamyl acetate and mineral oil (percent correct > 80% in two blocks) the odorant pair

was switched to the 1% Acetophenone S+ and Ethyl Benzoate S- (APEB) odorant pair.

Once the mice learned to discriminate between acetophenone and ethyl benzoate

acetate the odor pair was reversed. Ethyl Benzoate became the S+ and Acetophenone

the S- (EBAP). When the mice became proficient at the reversal, the odor pair was

switched. The same pattern was followed for the rest of the odor pairs (see Table1 for

the list of all odorants used). All odorants were obtained from Sigma-Aldrich and were

diluted in mineral oil at room temperature. Mice were trained until they performed at 80

percent correct or better in the last two blocks of 20 trials. Fig. 2B shows an example of

the percent correct odorant discrimination performance per trial. We did not find

significant differences in the analysis between odorant pairs.

Double tetrode recordings

We followed procedures as previously described (Li et al., 2015). The mouse was

recorded within the olfactometer chamber with dimensions of 11.6 × 9.7 × 9.4 cm. The

EIB-8 boards that recorded signals from the tetrodes were connected to an INTAN

RHD2132 16 channel amplifier/A/D converter that interfaced with an RHD2000 USB

interface board. Extracellular potentials from the tetrodes were captured, filtered with

either a 500 Hz or a 5 kHz low pass filter, and digitized at 20 kHz. Meta data for the

behavioral events such as valve opening/closing times and odor identity were recorded

through a digital output from the olfactometer. Licks detected by the olfactometer were

recorded as an analog signal by the INTAN board.

PAC Analysis

As previously described in Losacco et al. (2020), phase amplitude coupling (PAC) data

were processed using the Hilbert transform using a method described by Tort et al.

(2010). Briefly, the signal was bandpass filtered with a 20^{th} order Butterworth filter using

Matlab's filtfilt function with zero phase shift to extract LFP in the low frequency

oscillation used for phase estimation and the high frequency oscillation used for

estimation of the amplitude of the envelope (Figure 1Ei,ii). The Hilbert transform

established the theta (6-14 Hz) phase and the envelope for beta (15-30 Hz) and high

gamma (65-95 Hz, referred to as gamma) (Figures 1Eiii,iv). To quantify the strength of

PAC, we calculated the modulation index (MI). If PAC is non-existent, MI = 0, meaning

the mean amplitude is distributed uniformly over theta phases, and if PAC is a delta

function MI = 1. MI for signals measured in brain areas such as the hippocampus

typically fall between 0–0.03 (Tort et al., 2010).

PRP Analysis

As previously described in Losacco et al. (2020), we developed the PRP approach

using custom MATLAB code. PAC was calculated following the approach used by Tort

et al. (2010), as described in PAC analysis and summarized in Figure 1. Peak and

trough theta phases are defined as the phase for maxima and minima of the PAC

distribution measured for the S+ trial. A continuous Morlet wavelet transform was used

to estimate the power for the high frequency oscillations (Buonviso et al., 2003). PRP

was estimated as the power of the high frequency oscillations measured at the peak or

trough of PAC. The MATLAB code used for data analysis has been deposited to

https://github.com/ restrepd/drgMaster.

Coherence and Phase Locking Value (PLV) Analysis

In order to quantify the interaction of LFP oscillations from electrodes impaled into

different brain regions we computed the complementary values of coherence and phase

locking value (PLV). We measured coherence following the method detailed in the

supplemental information of (Borjigin et al., 2013). Briefly, coherence between LFP

channels is measured by amplitude squared coherence $C_{xy}(f)$ where f is frequency

(mscohere.m in the MATLAB signal toolbox; MathWorks, Inc.). This is a coherence

estimate of the input signals x and y using Welch's averaged, modified periodogram

method. The magnitude squared coherence estimate $C_{xy}(f)$ is a function of frequency

with values between 0 and 1 that indicates how well x corresponds to y at each

frequency. Furthermore, we quantified phase-locking between the two LFPs by

computing PLV following the procedure detailed by Lachaux et al. (1999) using

MATLAB code generated by Praneeth Namburi (Namburi, 2021). Briefly, we compute

the convolution of each LFP with a complex Gabor wavelet centered at frequency f and

then we compute the PLV as the normalized absolute value of the sum of the

exponential of the difference in phase multiplied by the imaginary number i. If the phase

difference between the two LFPs varies little across the trials, PLV is close to 1; it is

close to zero when the relationship between the phases varies randomly across trials.

Statistical analysis

The statistical analysis was done as described in Losacco et al. (2020) using MATLAB

code. Both PAC and PRP parameters were calculated separately per electrode (16

electrodes per mouse, 8 hippocampus, 8 mPFC) for all electrodes per mouse. Statistical

significance for changes in measured parameters for multivariate factors such as the

mouse genotype, naïve vs. proficient and S+ vs. S-, and the interactions of these factors

was estimated using generalized linear model (GLM) analysis, with post-hoc tests for all

data pairs corrected for multiple comparisons using false discovery rate (Curran-Everett,

2000). The post hoc comparisons between sets of data were performed either with a t-

test, or a ranksum test, depending on the result of an Anderson-Darling test of

normality. Degrees of freedom and statistical significance have the same meaning in

GLM as in analysis of variance and regression (Agresti, 2015). In addition, as a

complementary assessment of significant differences (Halsey et al., 2015) we display

95% confidence intervals (CIs) shown in the figures as vertical black lines or shadow

boundaries that was estimated by bootstrap analysis of the mean by sampling with

replacement 1000 times.

Linear discriminant analysis

Decoding of contextual odorant identity from tPRP values was performed using linear

discriminant analysis (LDA) using MATLAB code as described in Losacco et al. (2020).

LDA was trained by a leave one out procedure where the algorithm was trained with all

trials but one, and the accuracy was assessed by predicting the odorant identity for the

trial that was left out. This was repeated for all trials and was performed separately for

peak and trough PRP, and for analysis where the identity of the odorants was shuffled.

LDA was performed on a per-mouse basis where the input was the PRP recorded from

16 electrodes.

Acknowledgements

We thank Ms. Nicole Arevalo for animal husbandry and Ms. Dnate' Baxter for laboratory

support. We thank Dr. Stephen Coultrap for helpful discussion and animal support and

Mrs. Lauraine Mediavillo for assistance in collecting data. This research was supported

by an Administrative Supplement S1 to NIH UF1 NS116241 (to DRG), NIH R01

NS081248 (to KUB), NIH R01 DC000566 (to DR) and by a pilot grant from the Center

for NeuroScience (CNS) of the University of Colorado School of Medicine (to KUB and

DR).

Competing interests. DR declares no competing interests.

717 **Table 1. Odorant pairs**

Name	Rewarded odorant (S+)	unrewarded odorant (S-)
APEB	1% Acetophenone	1% Ethyl benzoate
EBAP	1% Ethyl benzoate	1% Acetophenone
EAPA1	0.1% Ethyl acetate	0.05% Ethyl acetate + 0.05% Propyl acetate
EAPA1	0.05% Ethyl acetate + 0.05% Propyl acetate	0.1% Ethyl acetate
EAPA2	0.01% Ethyl acetate	0.005% Ethyl acetate + 0.005%Propyl acetate
PAEA2	0.005% Ethyl acetate + 0.005% Propyl acetate	0.01% Ethyl acetate
EAPA3	0.001% Ethyl acetate	0.0005% Ethyl acetate + 0.0005% Propyl acetate
PAEA3	0.0005% Ethyl acetate + 0.0005%Propyl acetate	0.001% Ethyl acetate

719 The odorants were diluted in odorless mineral oil.

Table 2. Number of sessions and number of mice per odorant pair

Odorant pair	# of mice	Total sessions
1% Acetophenone vs 1% Ethyl benzoate	6 WT	22
	7 Het	22
	5 KO	11
1% Ethyl benzoate vs 1% Acetophenone	6 WT	22
	7 Het	55
	5 KO	29
0.1% Ethyl acetate vs 0.05% ethyl acetate + 0.05% propyl acetate	6 WT	15
	7 Het	23
	5 KO	17
0.05% ethyl acetate + 0.05% propyl acetate vs 0.1% Ethyl acetate	6 WT	17
	7 Het	35
	5 KO	37
0.01% Ethyl acetate vs 0.005% ethyl acetate + 0.005% propyl acetate	6 WT	23
	7 Het	52
	4 KO	24
0.005% ethyl acetate + 0.005% propyl acetate vs 0.01% Ethyl acetate	6 WT	31
	7 Het	49
	4 KO	21
0.001% Ethyl acetate vs 0.0005% ethyl acetate + 0.0005% propyl acetate	6 WT	60
	7 Het	64
	3 KO	16
0.0005% ethyl acetate + 0.0005% propyl acetate vs 0.001% Ethyl acetate	6 WT	34
	7 Het	57
	3 KO	13

Figure 1. Tetrode location, behavioral task and PAC analysis.

(A) Mouse brain showing location of tetrodes in CA1 and mPFC. (B) Mouse undergoing

odor discrimination behavioral task. Mouse self-initiates trials by licking the lixit closing

the lick detection circuit. Odorants and water were delivered according to the timeline in

C and the decision tree in D. (C) Timeline for a single trial. When the animal entered the

port and licked at the lixit, it started the trial. TStart (1-1.5 sec) is the time from initiation

of the trial to odor delivery. At time 0 the final valve air flow was turned towards the odor

port for 2.5 sec resulting in odor onset ~100 ms after the valve was activated (Losacco

et al., 2020). To obtain the water reward for the rewarded odorant (S+) the mouse must

lick at least once during each 0.5 s block for four blocks in the response area. If the

stimulus is the rewarded odorant and the animal licked, a water reward was delivered

(Hit). If the animal did not lick in each 0.5 sec block (Miss), water was not delivered.

When the odorant was the unrewarded odorant (S-) the mouse was not rewarded with

water regardless of whether it licked in all 0.5 sec blocks (False Alarm, FA) or not

(Correct Rejection, CR). (D) Decision tree for each trial. Green check mark indicates the

correct decision was made. Red "X" mark indicates an incorrect decision was made.

Water reward is represented by the water droplet symbol. (E) PAC data analysis for the

LFP. For each electrode, the raw signal collected at 20 kHz (i. rewarded odorant [left],

unrewarded odorant right) was filtered into different frequency bands (ii. theta 6-14 Hz).

Hilbert transform was used to calculate the theta phase (iii) and the amplitude envelope

of higher oscillations such as high gamma (65-95 Hz) (red line in v; blue line is the

gamma filtered LFP).

Figure 2. The peak angle variance of PAC increased for the unrewarded odorant as the animal learned to discriminate odorants.

(A) Pseudocolor plots showing the phase amplitude relationship for the unrewarded odorant S- (left) and rewarded odorant S+ (right) for two example sessions (naïve and proficient) of the go-no-go task. As the animal became proficient differentiating between the rewarded and unrewarded odorants, PAC peak angle appeared to become more variable for the unrewarded odorant (S-) compared to the rewarded odorant (S+).

(B) Behavioral performance for the two sessions. (i) During the naïve stage, the animal licked for the presentation of the rewarded and unrewarded odorants (performance ~50%). (ii) During the proficient stage (red points), the animal licked more frequently during the presentation of the rewarded odorant and refrained from licking during the presentation of the unrewarded odorant. Blue = percent correct < 65%, red percent correct >80%. (C) Peak angle polar histograms for (i) naïve S- (ii) proficient S- (iii) naïve S+ and (iv) proficient S+. The variance of the peak angle appears to increase for the S- in comparison to the S+ during the proficient stage. (D) Bar graphs and violin plots showing the differences in MI between S+ and S- for hippocampus (i)theta/beta (ii) theta/high gamma and mPFC (iii) theta/beta (iv) theta/high gamma. For the beta tPAC for both the hippocampus and mPFC GLM finds statistically significant differences for S+ vs. S- and for the interaction between S+ vs. S- and naïve vs. proficient ($p<0.001$, 188 observations, 184 d.f., F-statistic=51,67.7, $p<0.001$, 6 mice, 8 odor pairs, Supplementary Table 1). (E) Bar graphs showing differences for peak angle variance between S+ and S- for hippocampus (i) for theta/beta (ii) theta/high gamma and mPFC (iii) theta/beta (iv) theta/high gamma. For both beta and gamma tPAC and for both the

776   hippocampus and mPFC GLM finds statistically significant differences for S+ vs. S- and

777   for the interaction between S+ vs. S- and naïve vs. proficient (p<0.001, 188

778   observations, 184 d.f., F-statistic=19.5-26.4, p<0.001, 6 mice, 8 odor pairs,

779   Supplementary Table 2). Asterisks show significant p values (p<pFDR) for post-hoc

780   pairwise tests.

Figure 3. Changes in phase referenced power as the animal became proficient

(A) Examples of the time course per trial for the average wavelet broadband LFP

spectrogram for (i) S+ naïve (30 trials) (ii) S- naïve (27 trials) (iii) S+ proficient (84 trials)

(iv) S- proficient (84 trials). Pseudocolor scale for LFP power is shown in dB. (B)

Average gamma wavelet power referenced to theta peak and through for the same trials

as shown in A (i) S- proficient (ii) S+ proficient, red: peak, blue: through, shadow:

confidence interval. (C) Lick rate for the same trials (i) S- proficient (ii) S+ proficient.

(D,E) Summary bar graphs/violin plots showing that peak tPRP decreases for S- as the

mice become proficient. D: peak, E: through, (i) theta/beta and (ii) theta/gamma

hippocampus (iii) theta/beta and (iv) theta/gamma prefrontal cortex. GLM finds

statistically significant differences for S+ vs. S- and for the interaction between S+ vs. S-

and naïve vs. proficient (p<0.001, 376 observations, 368 d.f., F-statistic=72.7-103,

p<0.001, 6 mice, 8 odor pairs, Supplementary Table 3). Asterisks show significant p

values (p<pFDR) for post-hoc pairwise tests.

Figure 4. Linear discriminant analysis for decoding the contextual odorant

identity from tPRP

A. Example for one mouse for the time course for the accuracy of odorant identity

decoding by a linear discriminant analysis algorithm trained using tPRP for the EAPA

odor pair (i) naïve theta/beta, (ii) proficient theta/beta, (iii) naïve theta/gamma, (iv)

proficient theta/gamma red: peak, blue: through, black: shuffled, shadow: confidence

interval, black bar: odorant application. B and C. Bar graphs/violin plots showing the

differences in decoding accuracy between shuffled, naïve, and proficient. B: Accuracy

for peak tPRP for (i) theta/beta in the hippocampus, (ii) theta/gamma in the

hippocampus, (iii) theta/beta in mPFC, (iv) theta/gamma in mPFC. C: Accuracy for

through for (i) theta/beta in the hippocampus, (ii) theta/gamma in the hippocampus, (iii)

theta/beta in mPFC, (iv) theta/gamma mPFC. The bars show the average accuracy, and

the points are the accuracy per mouse per odor pair. The vertical bars show the

confidence interval. For beta and gamma tPRP LDA GLM finds statistically significant

differences for naïve vs. proficient and shuffled vs. proficient (p<0.001, 756

observations, 744 d.f., F-statistic=25.1, p<0.001, 6 mice, 8 odor pairs, Supplementary

Table 5). For gamma tPRP LDA GLM finds statistically significant differences between

peak and trough (p<0.001, 756 observations, 744 d.f., F-statistic=25.2, p<0.001, 6 mice,

8 odor pairs, Table S4). Asterisks show significant p values (p<pFDR) for post-hoc

pairwise tests.

Figure 5. Odorant-elicited change in coherence and phase locking value decreased for the unrewarded odorant as the animal became proficient.

A. Example of pseudocolor plots of the average coherence time course per trial for (i) S+ naïve (30 trials) (ii) S- naïve (27 trials) (iii) S+ proficient (84 trials) (iv) S- proficient (84 trials). B. Summary for Δ coherence as the animal learns. (i) Frequency dependence for the average Δ coherence per mouse per odor pair elicited by the odorant for S+ naïve, S- naïve, S+ proficient and S- proficient (6 mice, 8 odor pairs, the bounded line

is the 95% CI). (ii to v) Average Δ coherence elicited by the odorant for the different

bandwidths (per mouse per odorant pair): (ii) theta, (iii) beta, (iv) gamma (6 mice, 8 odor

pairs, the vertical line is the 95% CI). Δ coherence decreases for S- as the animal

becomes proficient. GLM finds statistically significant differences for S+ vs. S- and the

interaction between S+ vs. S- and naïve vs. proficient (p<0.001, 190 observations, 186

d.f., F-statistic=21.6-32.4, p<0.001, 6 mice, 8 odor pairs, Supplementary Table 6). C.

Bar graphs showing odor-elicited change in average odorant-elicited change in Δ PLV

per mouse per odor pair for (i) theta, (ii) beta, (iii) gamma. (6 mice, 8 odor pairs, the

vertical line is the 95% CI). GLM finds statistically significant differences between naïve

vs. proficient (p<0.05) and for S+ vs. S- and the interaction between S+ vs. S- and

naïve vs. proficient (p<0.001, 192 observations, 188 d.f., F-statistic=29.5-46.5, p<0.001,

6 mice, 8 odor pairs, Supplementary Table 7). D. Correlation between the odor-elicited

change in coherence and the change in Δ PLV for (i) theta, (ii) beta, (iii) gamma. This

shows a relationship between Δ coherence and Δ PLV (correlation coefficient > 0.91,

p<0.001).

Figure 6. The accuracy for decoding the contextual identity of the odorant from

tPRP decreased in the CaMKIIα knockout mouse and was correlated with percent

correct discrimination. A and B. Examples for one mouse of the time course for the

accuracy of odorant identification by linear discriminant analysis trained using tPRP for

the EAPA odor pair. A. theta/beta PRP. B. theta/gamma PRP. (i) WT, (ii) Het, (iii) KO.

Shadow: confidence interval, black bar: odorant application. C and D. Bar graphs

showing the differences in discriminant accuracy between the different genotypes. C:

Accuracy for peak tPRP for (i) theta/beta in the hippocampus, (ii) theta/gamma in the

hippocampus, (iii) theta/beta in mPFC, (iv) theta/gamma in mPFC. D: Accuracy for

through for (i) theta/beta in the hippocampus, (ii) theta/gamma in the hippocampus, (iii)

theta/beta in mPFC, (iv) theta/gamma mPFC. The bars show the average accuracy, and

the points are the accuracy per mouse per odor pair. The vertical bars show the

confidence interval. For beta tPRP LDA GLM finds statistically significant differences

between WT and KO for all conditions (p<0.001) and between WT and Het for

theta/gamma trough PRP (p<0.05, 756 observations, 744 d.f., F-statistic=9.9-34.3,

p<0.001, 6 mice, 8 odor pairs, Supplementary Table 11). Asterisks show significant p

values (p<pFDR) for post-hoc pairwise tests. E. Relationship for proficient mice between

percent correct in the go-no go behavior and accuracy of odor identification by the LDA

decoding algorithm shown per mouse per odor pair (6 mice, 8 odor pairs). The

correlation coefficients were: E (i) 0.3, E (ii) 0.24, E (iii) 0.29, E (iv) 0.23, F (i) 0.3, F (ii)

0.36, F (iii) 0.29, F (iv) 0.40, and the p value for significance was p<0.01. Lines are best

fit lines.

Figure 7. The odorant-elicited change in coherence was smaller for CaMKIIα

knockout mice.

A. Summary bar graphs comparing the odorant-elicited change in coherence (Δ

coherence) for the different genotypes. (i) theta, (ii) beta, (iii) gamma. The bars show

the average Δ coherence, and the points are Δ coherence per mouse per odor pair. The

vertical bars show the confidence interval. For Δ coherence GLM finds statistically

significant differences between WT and KO, and for the interaction of WT vs. KO and

S+ vs. S- (p<0.05, 546 observations, 534 d.f., F-statistic=21.2-33.8, p<0.001, 6 mice, 8

odor pairs, Supplementary Table 12). Asterisks show significant p values (p<pFDR) for

post-hoc pairwise tests. B. Summary bar graphs comparing change in Δ PLV for the

different genotypes. (i) beta, (ii) theta, (iii) gamma. For Δ PLV GLM finds no statistically

significant differences between WT and KO, and for the interaction of WT vs. KO and

S+ vs. S- (p>0.05, 512 observations, 500 d.f., F-statistic=25.5-39.9, p<0.001, 6 mice, 8

odor pairs, Supplementary Table 13).

Figure 3 – Figure Supplement 1. Effect of reversal of odorant valence on the phase referenced power. (A) Average gamma wavelet power referenced to theta peak and through for the same trials when the mouse was proficient shown in Figure 3A. The gamma tPRP is normalized by subtracting the power before odorant application. (i,ii) Forward: (i) S- (ethyl benzoate, EB) (ii) S+ (acetophenone, AP); (iii,iv) Reversed: (iii) S- (acetophenone, AP) (iv) S+ (ethyl benzoate, EB), red: peak, blue: through, shadow; confidence interval. (B,C) Summary bar graphs showing that the average peak tPRP decreases for S- upon odorant application regardless of the identity of the odorant. B:

peak, C: through, (i) theta/beta and (ii) theta/gamma hippocampus (iii) theta/beta and

(iv) theta/gamma prefrontal cortex. For beta tPRP GLM finds statistically significant

differences for forward vs. reverse (p<0.05) and S+ vs. S- (p<0.001, 188 observations,

180 d.f., F-statistic=87, p<0.001, 6 mice, 8 odor pairs, Table S12). For gamma tPRP

GLM finds statistically significant differences for S+ vs. S- and peak vs. trough (p<0.001,

188 observations, 180 d.f., F-statistic=61.8, p<0.001, 6 mice, 8 odor pairs, Table S12).

Asterisks show significant p values (p<pFDR) for post-hoc pairwise tests.

Figure 6 – Figure Supplement 1. Analysis of the behavioral performance for mice

of the three genotypes. (A). Number of sessions to criterion per odorant pair (2 blocks

of 20 trials with performance >80%). There was no difference between genotypes in the

number of sessions to criterion (GLM p>0.05, 24 observations, 21 d.f., F-statistic=0.26,

p>0.05, 8 odor pairs). (B) Percent correct for proficient mice per odorant per mouse.

GLM finds a statistical difference for WT compared to both CaMKIIα Het and CaMKIIα

KO (GLM p<0.05, 137 observations, 134 d.f., F-statistic=3.7, p<0.05, 6 mice, 8 odor

pairs). (C). Intertrial interval (ITI) for proficient mice. GLM analysis finds that the ITI for

CaMKIIα Het differed from both CaMKIIα Het and CaMKIIα KO (p<0.001, 137

observations, 134 d.f., F-statistic=13.5, p<0.001, 6 mice, 8 odor pairs). Asterisks denote

statistically significant differences evaluated with either t test or ranksum corrected for

multiple comparisons (p<pFDR).

Figure 6 – Figure Supplement 2. PAC and peak angle variance per genotype

(A,B) Bar graphs showing average MI per mouse per odor pair for each genotype, S+

and S- in naïve and proficient mice for (A) theta/beta and (B) theta/gamma for (i)

hippocampus and (ii) mPFC. The bars show the average MI, and the points are the MI

per mouse per odor pair. For theta/beta PAC MI GLM finds a statistically significant

difference for S+ vs. S-, for the interaction of WT vs. KO and S+ vs. S- and the

interaction of WT vs. KO and naïve vs. proficient ($p < 0.001$, 544 observations, 532 d.f.,

F-statistic=10.3-34.3, $p < 0.001$, 6 mice, 8 odor pairs, Table S10). For theta/gamma PAC

MI GLM finds a statistically significant difference between WT vs. KO and WT vs. Het

($p < 0.01$) and for the interaction of WT vs. Het and S+ vs. S- ($p < 0.05$, 544 observations,

532 d.f., F-statistic=10.3-34.3, $p < 0.001$, 6 mice, 8 odor pairs, Table S10). (C,E) Bar

graphs showing average peak angle variance per mouse per odor pair for each

genotype, S+ and S- in naïve and proficient mice for (A) theta/beta and (B)

theta/gamma for (i) hippocampus and (ii) mPFC. For theta/beta PAC MI GLM finds a

statistically significant difference for WT vs. Het ($p < 0.001$) and WT vs. KO ($p < 0.05$), S+

vs. S-, ($p < 0.001$) and naïve vs. proficient ($p < 0.05$, 544 observations, 532 d.f., F-

statistic=20.6, $p < 0.001$, 6 mice, 8 odor pairs, Table S11). For theta/gamma PAC MI

GLM finds a statistically significant difference for WT vs. Het ($p < 0.05$) and WT vs. KO

($p < 0.05$), S+ vs. S-, ($p < 0.001$) and naïve vs. proficient ($p < 0.05$, 544 observations, 532

d.f., F-statistic=18.8, $p < 0.001$, 6 mice, 8 odor pairs, Table S11).

Note: Supplementary tables show the output of the MATLAB code that performs the

GLM analysis and the pairwise ranksum/t test. They will be included in our

resubmission.

References

Agresti, A. (2015). Foundations of linear and generalized linear models. In Wiley series in probability and statistics (Hoboken, New Jersey: John Wiley & Sons Inc.,), pp. 1 online resource.

Bayer, K.U., and Schulman, H. (2019). CaM Kinase: Still Inspiring at 40. Neuron *103*, 380-394.

Bear, M.F., Cooke, S.F., Giese, K.P., Kaang, B.K., Kennedy, M.B., Kim, J.I., Morris, R.G.M., and Park, P. (2018). In memoriam: John Lisman - commentaries on CaMKII as a memory molecule. Mol Brain *11*, 76.

Belluscio, M.A., Mizuseki, K., Schmidt, R., Kempter, R., and Buzsáki, G. (2012). Cross-Frequency Phase–Phase Coupling between Theta and Gamma Oscillations in the Hippocampus. The Journal of Neuroscience *32*, 423.

Bliss, T.V., and Lomo, T. (1973). Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path. J Physiol *232*, 331-356.

Borjigin, J., Lee, U., Liu, T., Pal, D., Huff, S., Klarr, D., Sloboda, J., Hernandez, J., Wang, M.M., and Mashour, G.A. (2013). Surge of neurophysiological coherence and connectivity in the dying brain. Proceedings of the National Academy of Sciences *110*, 14432.

Buonviso, N., Amat, C., Litaudon, P., Roux, S., Royet, J.P., Farget, V., and Sicard, G.

(2003). Rhythm sequence through the olfactory bulb layers during the time window of a

respiratory cycle. EurJNeurosci *17*, 1811-1819.

Butler, J.L., Mendonça, P.R.F., Robinson, H.P.C., and Paulsen, O. (2016). Intrinsic

Cornu Ammonis Area 1 Theta-Nested Gamma Oscillations Induced by Optogenetic

Theta Frequency Stimulation. The Journal of Neuroscience *36*, 4155.

Cajal, S. (1904). Textura del sistema nervioso del hombre y de los vertebrados (Madrid:

Moya).

Cansler, H.L., in 't Zandt, E.E., Carlson, K.S., Khan, W.T., Ma, M., and Wesson, D.W.

(2021). Organization and engagement of a prefrontal-olfactory network during olfactory

selective attention. bioRxiv, 2021.2009.2004.458996.

Chen, C., Rainnie, D.G., Greene, R.W., and Tonegawa, S. (1994). Abnormal fear

response and aggressive behavior in mutant mice deficient for alpha-calcium-

calmodulin kinase II. Science *266*, 291-294.

Chia, P.H., Zhong, F.L., Niwa, S., Bonnard, C., Utami, K.H., Zeng, R., Lee, H., Eskin,

A., Nelson, S.F., Xie, W.H.*, et al.* (2018). A homozygous loss-of-function CAMK2A

mutation causes growth delay, frequent seizures and severe intellectual disability. eLife

7, e32451.

Colgin, L.L. (2011). Oscillations and hippocampal–prefrontal synchrony. Current

Opinion in Neurobiology *21*, 467-474.

Colgin, L.L. (2015). Theta-gamma coupling in the entorhinal-hippocampal system. Curr

Opin Neurobiol *31*, 45-50.

Cook, S.G., Buonarati, O.R., Coultrap, S.J., and Bayer, K.U. (2021). CaMKII

holoenzyme mechanisms that govern the LTP versus LTD decision. Sci Adv *7*.

Coultrap, S.J., Freund, R.K., O'Leary, H., Sanderson, J.L., Roche, K.W., Dell'Acqua,

M.L., and Bayer, K.U. (2014). Autonomous CaMKII mediates both LTP and LTD using a

mechanism for differential substrate site selection. Cell Rep *6*, 431-437.

Curran-Everett, D. (2000). Multiple comparisons: philosophies and illustrations.

AmJPhysiol RegulIntegrComp Physiol *279*, R1-R8.

Eleore, L., Lopez-Ramos, J.C., Guerra-Narbona, R., and Delgado-Garcia, J.M. (2011).

Role of reuniens nucleus projections to the medial prefrontal cortex and to the

hippocampal pyramidal CA1 area in associative learning. PLoS One *6*, e23538.

Frankland, P.W., O'Brien, C., Ohno, M., Kirkwood, A., and Silva, A.J. (2001). Alpha-

CaMKII-dependent plasticity in the cortex is required for permanent memory. Nature

411, 309-313.

Fries, P. (2005). A mechanism for cognitive dynamics: neuronal communication through

neuronal coherence. Trends Cogn Sci *9*, 474-480.

Fromer, M., Pocklington, A.J., Kavanagh, D.H., Williams, H.J., Dwyer, S., Gormley, P.,

Georgieva, L., Rees, E., Palta, P., Ruderfer, D.M.*, et al.* (2014). De novo mutations in

schizophrenia implicate synaptic networks. Nature *506*, 179-184.

Goodell, D.J., Benke, T.A., and Bayer, K.U. (2016). Developmental restoration of LTP

deficits in heterozygous CaMKIIα KO mice. Journal of Neurophysiology *116*, 2140-2151.

Gordon, J.A. (2011). Oscillations and hippocampal–prefrontal synchrony. Current

Opinion in Neurobiology *21*, 486-491.

Gourevitch, B., Kay, L.M., and Martin, C. (2010). Directional coupling from the olfactory

bulb to the hippocampus during a go/no-go odor discrimination task. J Neurophysiol

103, 2633-2641.

Halsey, L.G., Curran-Everett, D., Vowler, S.L., and Drummond, G.B. (2015). The fickle

P value generates irreproducible results. Nature methods *12*, 179-185.

Hasegawa, S., Furuichi, T., Yoshida, T., Endoh, K., Kato, K., Sado, M., Maeda, R.,

Kitamoto, A., Miyao, T., Suzuki, R.*, et al.* (2009). Transgenic up-regulation of alpha-

CaMKII in forebrain leads to increased anxiety-like behaviors and aggression. Mol Brain

2, 6.

Headley, D.B., and Paré, D. (2017). Common oscillatory mechanisms across multiple

memory systems. npj Science of Learning *2*, 1.

Kaplan, R., Bush, D., Bonnefond, M., Bandettini, P.A., Barnes, G.R., Doeller, C.F., and

Burgess, N. (2014). Medial prefrontal theta phase coupling during spatial memory

retrieval. Hippocampus *24*, 656-665.

Klein, S.B., Cosmides, L., Tooby, J., and Chance, S. (2002). Decisions and the

evolution of memory: multiple systems, multiple functions. Psychol Rev *109*, 306-329.

Lachaux, J.P., Rodriguez, E., Martinerie, J., and Varela, F.J. (1999). Measuring phase

synchrony in brain signals. Hum Brain Mapp *8*, 194-208.

Larson, J., and Munkácsy, E. (2015). Theta-burst LTP. Brain Research *1621*, 38-50.

Li, A., Gire, D.H., and Restrepo, D. (2015). Y spike-field coherence in a population of

olfactory bulb neurons differentiates between odors irrespective of associated outcome.

J Neurosci *35*, 5808-5822.

Li, Y., Xu, J., Liu, Y., Zhu, J., Liu, N., Zeng, W., Huang, N., Rasch, M.J., Jiang, H., Gu,

X*., et al.* (2017). A distinct entorhinal cortex to hippocampal CA1 direct circuit for

olfactory associative learning. Nature Neuroscience *20*, 559-570.

Lisman, J., Buzsaki, G., Eichenbaum, H., Nadel, L., Ranganath, C., and Redish, A.D.

(2017). Viewpoints: how the hippocampus contributes to memory, navigation and

cognition. Nat Neurosci *20*, 1434-1447.

Lisman, J., Yasuda, R., and Raghavachari, S. (2012). Mechanisms of CaMKII action in

long-term potentiation. Nat Rev Neurosci *13*, 169-182.

Losacco, J., Ramirez-Gordillo, D., Gilmer, J., and Restrepo, D. (2020). Learning

improves decoding of odor identity with phase-referenced oscillations in the olfactory

bulb. Elife *9*, e52583.

Malinow, R., Schulman, H., and Tsien Richard, W. (1989). Inhibition of Postsynaptic

PKC or CaMKII Blocks Induction But Not Expression of LTP. Science *245*, 862-866.

Malvaut, S., Gribaudo, S., Hardy, D., David, L.S., Daroles, L., Labrecque, S., Lebel-

Cormier, M.-A., Chaker, Z., Coté, D., De Koninck, P.*, et al.* (2017). CaMKIIalpha

Expression Defines Two Functionally Distinct Populations of Granule Cells Involved in

Different Types of Odor Behavior. Current Biology *27*, 3315-3329.e3316.

Martin, C., Beshel, J., and Kay, L.M. (2007). An olfacto-hippocampal network is

dynamically involved in odor-discrimination learning. JNeurophysiol *98*, 2196-2205.

Matsuo, N., Yamasaki, N., Ohira, K., Takao, K., Toyama, K., Eguchi, M., Yamaguchi, S.,

and Miyakawa, T. (2009). Neural activity changes underlying the working memory deficit

in alpha-CaMKII heterozygous knockout mice. Front Behav Neurosci *3*, 20.

Moberly, A.H., Schreck, M., Bhattarai, J.P., Zweifel, L.S., Luo, W., and Ma, M. (2018).

Olfactory inputs modulate respiration-related rhythmic activity in the prefrontal cortex

and freezing behavior. Nature Communications *9*, 1528.

Mofleh, R., and Kocsis, B. (2021). Delta-range coupling between prefrontal cortex and

hippocampus supported by respiratory rhythmic input from the olfactory bulb in freely

behaving rats. Scientific Reports *11*, 8100.

Nakazawa, K., McHugh, T.J., Wilson, M.A., and Tonegawa, S. (2004). NMDA receptors,

place cells and hippocampal spatial memory. Nature Reviews Neuroscience *5*, 361-372.

Namburi, P. (2021). Phase locking value. https://praneethnamburi.com/2011/08/10/plv/.

Nguyen Chi, V., Muller, C., Wolfenstetter, T., Yanovsky, Y., Draguhn, A., Tort, A.B., and

Brankack, J. (2016). Hippocampal Respiration-Driven Rhythm Distinct from Theta

Oscillations in Awake Mice. J Neurosci *36*, 162-177.

Purcell, S.M., Moran, J.L., Fromer, M., Ruderfer, D., Solovieff, N., Roussos, P.,

O'Dushlaine, C., Chambert, K., Bergen, S.E., Kahler, A.*, et al.* (2014). A polygenic

burden of rare disruptive mutations in schizophrenia. Nature *506*, 185-190.

Robison, A.J. (2014). Emerging role of CaMKII in neuropsychiatric disease. Trends in

Neurosciences *37*, 653-662.

Rojas-Libano, D., Frederick, D.E., Egana, J.I., and Kay, L.M. (2014). The olfactory bulb

theta rhythm follows all frequencies of diaphragmatic respiration in the freely behaving

rat. Front Behav Neurosci *8*, 214.

Scheffer-Teixeira, R., and Tort, A.B. (2016). On cross-frequency phase-phase coupling

between theta and gamma oscillations in the hippocampus. eLife *5*, e20515.

Siegle, J.H., and Wilson, M.A. (2014). Enhancement of encoding and retrieval functions

through theta phase-specific manipulation of hippocampus. Elife *3*, e03061.

Silva, A.J., Paylor, R., Wehner, J.M., and Tonegawa, S. (1992a). Impaired spatial

learning in alpha-calcium-calmodulin kinase II mutant mice. Science *257*, 206-211.

Silva, A.J., Rosahl, T.W., Chapman, P.F., Marowitz, Z., Friedman, E., Frankland, P.W.,

Cestari, V., Cioffi, D., Südhof, T.C., and Bourtchuladze, R. (1996). Impaired learning in

mice with abnormal short-lived plasticity. Current Biology *6*, 1509-1518.

Silva, A.J., Stevens, C.F., Tonegawa, S., and Wang, Y. (1992b). Deficient hippocampal

long-term potentiation in alpha-calcium-calmodulin kinase II mutant mice. Science *257*,

201-206.

Slotnick, B.M., and Restrepo, D. (2005). Olfactometry with mice. In Current Protocols in

Neuroscience, J.N. Crawley, C.R. Gerefen, M.A. Rogawski, D.R. Sibley, P. Skolnick,

and S. Wray, eds. (New York: John Wiley and Sons, Inc), pp. 1-24

1095

Suarez-Pereira, I., Canals, S., and Carrion, A.M. (2015). Adult newborn neurons are

involved in learning acquisition and long-term memory formation: the distinct demands

on temporal neurogenesis of different cognitive tasks. Hippocampus *25*, 51-61.

Tort, A.B., Komorowski, R., Eichenbaum, H., and Kopell, N. (2010). Measuring phase-

amplitude coupling between neuronal oscillations of different frequencies. J

Neurophysiol *104*, 1195-1210.

Udakis, M., Pedrosa, V., Chamberlain, S.E.L., Clopath, C., and Mellor, J.R. (2020).

Interneuron-specific plasticity at parvalbumin and somatostatin inhibitory synapses onto

CA1 pyramidal neurons shapes hippocampal output. Nature Communications *11*, 4395.

Vizcay, M.A., Duarte-Mermoud, M.A., and Aylwin Mde, L. (2015). Odorant recognition

using biological responses recorded in olfactory bulb of rats. Comput Biol Med *56*, 192-

199.

Walton, N.M., Zhou, Y., Kogan, J.H., Shin, R., Webster, M., Gross, A.K., Heusner, C.L.,

Chen, Q., Miyake, S., Tajinda, K.*, et al.* (2012). Detection of an immature dentate gyrus

feature in human schizophrenia/bipolar patients. Transl Psychiatry *2*, e135.

Won, G.H., Kim, J.W., Choi, T.Y., Lee, Y.S., Min, K.J., and Seol, K.H. (2018). Theta-

phase gamma-amplitude coupling as a neurophysiological marker in neuroleptic-naïve

schizophrenia. Psychiatry Research *260*, 406-411.

Yamasaki, N., Maekawa, M., Kobayashi, K., Kajii, Y., Maeda, J., Soma, M., Takao, K.,

Tanda, K., Ohira, K., Toyama, K*., et al.* (2008). Alpha-CaMKII deficiency causes

immature dentate gyrus, a novel candidate endophenotype of psychiatric disorders.

Molecular Brain *1*, 6.

1118