Genotype-specific benefits and risks of a mutualist association during natural epidemics

Jenalle L. Eck,[1*] Minna-Maarit Kytöviita,[2] & Anna-Liisa Laine [1,3]

[1] Department of Evolutionary Biology and Environmental Studies, University of Zurich, 8057

Zurich, Switzerland

[2] Department of Biological and Environmental Science, University of Jyväskylä, 40014

Jyväskylä, Finland

[3] Organismal and Evolutionary Biology Research Program, University of Helsinki, 00790

Helsinki, Finland

* Correspondence to: jenalle.eck@ieu.uzh.ch

Keywords: mutualism, mycorrhizal fungi, plant disease, plant pathogen, *Plantago lanceolata*,

Podosphaera plantaginis, powdery mildew, protective symbiont

Summary

- While both plant-associated pathogenic and mutualistic microbes are ubiquitous across ecosystems, how they interact to determine disease risk in natural, genetically diverse populations is not known.

- To test whether mycorrhizal fungi provide protection against infection, and whether this functions in a genotype-specific manner, we conducted a field experiment in three naturally-occurring epidemics of a fungal pathogen, *Podosphaera plantaginis*, infecting a host plant, *Plantago lanceolata*, in the Åland Islands, Finland. In each population, we collected field epidemiological data on mycorrhizal-inoculated and non-mycorrhizal experimental plants originating from six allopatric populations.

- Mycorrhizal association caused host genotype-specific changes in growth and infection rates in the host populations, but reduced infection severity in hosts from every genetic origin. Protection occurred via changes in the relationship between growth and infection in the host individuals: mycorrhizal individuals grew larger without increasing their infection risk or load. More susceptible host genotypes received stronger protective effects from mycorrhizae.

- Our results show that mycorrhizal fungi produce host genotype-specific growth and defensive benefits and alter infection risks in wild host populations. Understanding how mutualism-derived protection alters host susceptibility to disease will be important for predicting infection outcomes in ecological communities and in agriculture.

Introduction

Protective symbionts – species that provide defensive benefits to their hosts – help

determine the outcome of species interactions and, thus, shape ecological and evolutionary

dynamics between hosts and parasites (Brownlie & Johnson, 2009; May & Nelson, 2014; King *et*

al., 2016; Sochard *et al*., 2020). Despite their importance, ecological studies examining the role

of protective symbionts in influencing host-parasite interactions in natural populations and

communities are rare (Oliver *et al*., 2014; Hafer-Hahmann & Vorburger, 2021). Protection

against pathogenic disease by mutualistic microbes, such as mycorrhizal fungi, has been

demonstrated under controlled laboratory conditions in several economically important

agricultural plant species (Norman *et al*., 1996; Pozo *et al*., 2002; Li *et al*., 2010; Song *et al*.,

2015; Berdeni *et al*., 2018). However, it has not been verified whether protective symbionts

mediate infection risk under natural epidemics, which are characterized by repeated pathogen

encounters, as well as by environmental and genotypic diversity in both hosts and pathogens

(Newsham *et al*., 1995). Mycorrhizal associations are widespread among terrestrial plants (Öpik

et al., 2006; van der Heijden *et al*., 2008), and have important impacts on plant fitness and

population dynamics (Barea *et al*., 2002; Koide & Dickie, 2002), community composition

(Hartnett & Wilson, 2002) and ecosystem functioning (Rillig, 2004). Understanding how

mycorrhizal and other mutualisms influence patterns of plant disease is essential given that

disease is a major factor shaping the abundance, diversity, and distribution of species in plant

communities (Bever *et al*., 2015), and affecting food production (Johansson *et al*., 2004; Gosling

et al., 2006; Pretty *et al*., 2011; Hohmann & Messmer, 2017). While both plant-associated

pathogenic and mutualistic microbes are ubiquitous across ecosystems, how they interact to

determine disease risk in natural, genetically diverse populations is not known.

Mycorrhizal fungi produce a suite of growth, nutritional, and/or defensive benefits that

may help protect plants from co-occurring antagonists, such as pathogenic microbes and

herbivores (Delavaux *et al*., 2017). Association with mycorrhizal fungi often improves plant

nutrient and water uptake (Smith & Read, 2008), though there is both intra- and interspecific

variation among plants in their affinity to form and benefit from mycorrhizal associations (Thrall

et al., 2011; Rasmussen *et al*., 2019). Increases in host size and nutritional status due to

mycorrhizal association can improve host tolerance against antagonists and abiotic stress

(Azcón-Aguilar & Barea, 1996). Mycorrhizal fungi can also influence host defenses directly, by

upregulating defense gene expression in their host plants (Azcón-Aguilar & Barea, 1996; Pozo &

Azcón-Aguilar, 2007; Jung *et al*., 2012). This form of protection, known as defense priming,

allows a more efficient activation of defense mechanisms in response to attack by potential

enemies, and has been shown to reduce the negative effects of interactions with a wide range of

antagonist species (Jung *et al*., 2012; Delavaux *et al*., 2017). Though mycorrhizal associations

occur belowground (at the root-soil interface), the induced resistance response in the host is

systemic (Cameron *et al*., 2013), meaning that even strictly aboveground antagonists may also be

affected (Koricheva *et al*., 2009). It is unclear how often mycorrhizal growth and defensive

benefits are conferred together in hosts, and how they operate simultaneously to determine the

incidence and outcome of interactions between hosts and parasites.

Association with mycorrhizal fungi or other mutualists can also present risks to hosts that

may influence infection dynamics (Polin *et al*., 2014). Ecological costs can occur when

combinations of host and mutualist species or genotypes are mismatched (Klironomos, 2003;

Hoeksema *et al*., 2010), resulting in inefficient mutualisms that fail to convert host resources in

to necessary benefits (Johnson *et al*., 1997; Jones & Smith, 2004; Grman, 2012). Furthermore,

unfavorable abiotic conditions could reduce or negate potential mutualist benefits (Hoeksema *et*

al., 2010; Qu *et al*., 2021). In addition, defensive benefits from mutualists may not be effective

against all antagonist species (e.g., depending on the lifestyle and infection strategy of the

pathogen) (Pozo & Azcón-Aguilar, 2007), or be durable to changes in pathogen traits or

composition in the environment. Hence, the potential ecological risks and/or benefits of

mutualist associations for disease protection may depend on genotype, and could vary among or

within host populations and environments; however, this has remained largely unexplored.

In addition, it is unclear how mutualism-derived protection acts alongside innate host

resistance to determine the outcome of host-pathogen interactions. Host genetic resistance can

vary widely among and within natural populations (Salvaudon *et al*., 2008, Laine *et al*., 2011) -

potentially due to costs associated with its maintenance (Brown, 2003; Susi & Laine, 2015) - and

may be under a different set of selection pressures than mutualism (Thompson, 1994). Whether

mutualism presents benefits or costs to particular host individuals or populations could depend

on their resistance status. In resistant hosts, resources provided to mutualists in return for

defensive benefits could represent an unnecessary metabolic cost. However, in susceptible hosts,

mutualist-derived protection could compensate for lack of genetic resistance, presenting a viable

alternate strategy for coping with pathogens. Mutualism-derived protection could be especially

beneficial when genetic resistance is costly to maintain, or when disease is ephemeral, as it often

is in natural populations (Burdon & Thrall, 2014). How much of host resistance is derived from

innate genetic defenses versus defenses derived from protective symbionts (e.g., via defense

priming, or improved tolerance to pathogens), and the relative importance of each type of

defense in different species combinations and environmental contexts, remains to be seen.

Despite general recognition for the potential impact of mycorrhizal fungi on plant fitness,

how mycorrhizal association may impact host infection risk – and to what extent this varies

among plant genotypes and populations - is poorly understood in natural populations. To

examine this, we conducted a field experiment to test whether association with arbuscular

mycorrhizal fungi affects infection dynamics by a fungal pathogen in a shared host under natural

epidemic conditions. Specifically, we ask: 1) Do growth effects due to mycorrhizal association

vary among host populations and maternal genotypes? 2) Does association with mycorrhizal

fungi influence host infection risk? 3) Upon infection, does association with mycorrhizal fungi

affect disease severity? 4) Are growth and defensive effects from mycorrhizal association linked

in host genotypes? and 5) Are host disease susceptibility and mycorrhizal-derived defensive

effects linked in host genotypes? To answer these questions within an ecologically relevant

context, we placed mycorrhiza-inoculated and non-mycorrhizal plants in wild populations during

a natural pathogen epidemic. The experiment was conducted in the long-term Åland Islands

study site (Finland), where infection by a fungal pathogen (powdery mildew) has been surveyed

on a large host population network of *Plantago lanceolata* L. (Plantaginaceae) since 2001

(Jousimo *et al*., 2014). From prior studies in this pathosystem, we know that several factors, such

as spatial context (Laine, 2006; Soubeyrand *et al*., 2009; Jousimo *et al*., 2014), pathogen genetic

diversity (Eck *et al*., accepted manuscript), local adaptation between host and pathogen

genotypes (Laine, 2004; Laine, 2005; Jousimo *et al*., 2014), and abiotic conditions (Laine, 2007;

Laine, 2008) are all critical in determining infection, but the impact of mutualistic interactions on

infection dynamics has not been determined. To our knowledge, this is the first study of the

impact of mycorrhizal fungi on infection risk by a plant pathogen under natural epidemics and

across different host populations and/or genotypes.

Methods

 Study system

Our study is focused on a fungal pathogen, *Podosphaera plantaginis* (Castagne) U. Braun

& S. Takam., infecting *Plantago lanceolata* L. (Plantaginaceae) in the Åland Islands (60°08'53"

N, 19°47'18" E), Finland. We carried out an experiment in three natural *Pl. lanceolata*

populations that are part of a larger network of > 4000 mapped populations in Åland (Hanski,

1999). These populations have been surveyed for infection by *Po. plantaginis* since 2001 (Laine

& Hanski, 2006). *Plantago lanceolata* (ribwort plantain) is native to Åland and much of Eurasia,

and occurs mainly in small meadows and disturbed areas in Åland. It is monoecious, self-

incompatible, and reproduces either sexually (via wind-dispersed pollen and seeds; Bos, 1992) or

asexually (via clonally-produced side-rosettes) (Sagar & Harper, 1964). *Plantago lanceolata*

associates commonly with mycorrhizal fungi, and has been found in association with a wide

variety of mycorrhizal species in Åland (Rasmussen *et al*., 2018; Eck *et al*., unpublished data).

Podosphaera plantaginis, a powdery mildew fungus (order Erysiphales), is an obligate

biotroph of foliar tissue, and is host-specific in Åland, infecting only *Pl. lanceolata* (Laine,

2004). Fungal hyphae grow on the surface of *Pl. lanceolata* leaves, producing localized

infections that mitigate host growth and reproduction (Bushnell, 2002), and may lead to

mortality in the presence of abiotic stress, such as drought (Laine, 2004; Susi & Laine, 2015).

Infections are transmitted via asexually-produced, wind-dispersed spores (conidia) that are

produced cyclically (~ every two weeks) throughout an epidemic season (~ June to September in

Åland) (Ovaskainen & Laine, 2006). Resting structures (chasmothecia), produced via haploid

selfing or outcrossing between strains (Tollenaere & Laine, 2013), allow the pathogen to

overwinter and initiate new infections in the next growing season (Tack & Laine, 2014).

In Åland, *Po. plantaginis* persists as a metapopulation, with frequent colonization and

extinction events (~ 3 – 17 % of the *Pl. lanceolata* populations are infected in a given year)

(Jousimo *et al*., 2014). Resistance in *Pl. lanceolata* against *Po. plantaginis* functions in the two-

step manner typical of most plants (Jones & Dangl, 2006). The first step is strain specific, such

that a host genotype expresses resistance against some strains (*i.e.* recognition) of the pathogen,

while being susceptible to others (*i.e.* non-recognition) (Laine, 2004; Laine, 2007). Once a strain

has successfully established, its development is affected by both pathogen and host genotype

(Laine, 2007). Previous studies have demonstrated high levels of intraspecific variation in

resistance within and among populations of the host (Laine, 2004; Laine 2007, Jousimo et al.

2014). Pathogen populations also contain variable levels of genetic diversity (Eck *et al*., accepted

manuscript). Infected hosts individuals have reduced survival during drought periods compared

to healthy hosts (Laine, 2004), and infection has a negative effect on host population growth rate

(Penczykowski *et al*., 2015).

Mycorrhizal inoculation of experimental plants

To measure the effects of association with mycorrhizal fungi in genetically diverse host

plants, seeds were collected from six geographically-variable populations of *Pl. lanceolata* in the

Åland Islands (a map of these populations is displayed in Fig. S1). In each population, seeds

were collected from five haphazardly chosen maternal plants in August 2007. Seeds from each

maternal plant were kept separate (seeds from one maternal plant are half or full siblings) and

were stored in paper envelopes. In April 2008, seeds from each of the 30 maternal plants were

germinated in separate 6 x 6 x 7 cm pots (one maternal genotype per pot) in a greenhouse at the

University of Oulu (Oulu, Finland) in sterilized sand. At two weeks of age, ten seedlings from

each maternal plant were transferred to identical, individual pots (one seeding per pot) filled with

experimental substrate: a 5:4:1 mixture of heat-sterilized sand : heat –sterilized garden soil :

perlite, combined with 1 g of bone meal and 3 g of dolomite per liter substrate. Five of the

seedlings in each maternal cohort were inoculated with mycorrhizal fungi, while the other five

received a non-mycorrhizal control treatment. The mycorrhizal inoculum consisted of a mixture

of spores of three arbuscular mycorrhizal fungal species native to Finland: *Glomus hoi*,

Claroideoglomus clareoideum, and *Gl. mosseae (BEG 29).* The mycorrhizal inoculum consisted

of allopatric isolates of each species (originating from central Finland, 61°10 N, 24°40 E, rather

than Åland) so as not to confound the experiment with potential local adaptation between

mycorrhizal fungi and their host populations (Hoeksema *et al*., 2010). A mixture of species was

used as inoculum because *Pl. lanceolata* is colonized naturally by several mycorrhizal symbionts

(Johnson *et al*., 2003). Spore inocula were produced by growing the mycorrhizal species with

non-experimental *Pl. lanceolata* in a soil substrate identical to the experimental one. Spores were

washed out of the non-experimental substrate with water and 15 spores of each species (45

spores in total) were pipetted on to the roots of each seedling in the mycorrhizal treatment in 2

ml of water (6 mL in total); seedlings in the non-mycorrhizal control treatment received 2 ml of

filtered spore washing water from each fungal species (6 mL in total). All experimental seedlings

were also inoculated with bacteria filtered from a soil mix collected from the six seed source

Åland populations to restore the natural, non-mycorrhizal soil microbial community. Seedlings

were fertilized with a dilute nitrogen-based solution, and natural light was supplemented with

Osram HQI lamps to provide a photoperiod of 18 h day / 6 h night. At six weeks of age, the

plants were moved to an outdoor area (at the University of Oulu) and grown under a transparent

plastic roof for an additional six weeks, to acclimatize to field conditions (as *Pl. lanceolata* and

Po. plantaginis do not inhabit this region, infection at this stage was highly unlikely).

Natural epidemic field experiment

Ten weeks post-inoculation, healthy experimental plants were placed in to three

naturally-occurring populations of *Pl. lanceolata*, infected by populations of *Po. plantaginis*, in

Åland, to gain infections naturally from the surrounding epidemic. The three field epidemic

populations were allopatric to the six seed origin populations (see Fig. S1 for a map). Thus, they

represent common garden field sites, in which the host genotypes are not locally adapted to the

local environmental conditions or to the local pathogen population. Presence of *Po. plantaginis*

at the sites was confirmed by surveying the populations for visible signs of infection prior to the

placement of the experimental plants. Each field site received 96 experimental plants: 16 from

each of the six seed origin populations (eight of which had received mycorrhizal inoculum and

eight of which were non-mycorrhizal). Within these subgroups, the five maternal genotypes

within each seed origin population were allocated as evenly as possible to the three field sites;

the particular individuals placed in each site were selected at random (full, pairwise replication

of every maternal genotype × mycorrhizal treatment combination within every field site was not

possible due to lack of available plants). At each site, experimental plants were randomly fitted

in to one of 96 plastic containers (14 x 10.5 x 4.5 cm) that were placed on the ground in a

random array in the proximity of naturally-infected *Pl. lanceolata* individuals inhabiting the field

population. The containers prevented contact between the bottom of the pots and the soil,

preventing the experimental plants from acquiring soil microbes from the field sites.

At the time of placement in the field epidemic populations, we measured the following

for each plant: the number of total leaves, the number of leaves infected by *Po. plantaginis*, as

well as the length and width (in cm) of the longest leaf. All experimental plants were uninfected

(i.e., had zero infected leaves) at the beginning of the experiment. Infection surveys were then

conducted every three days at each field site, on a rotating survey schedule (1st day field

population 9061, 2nd day field population 3484, and 3rd day field population 9066). During each

infection survey, the number of healthy leaves and the number of leaves infected by *Po.*

plantaginis were re-measured (leaves that withered during the experiment were not counted).

Plants were also watered if necessary, and were re-randomized in to a new position in the

experimental array (this was done to minimize the effect of spatial positioning, with respect to

distance to infected individuals and prevailing wind direction, on infection risk and severity).

Seven infection surveys were conducted in each field population. At the end of the experiment, a

randomly-selected subset of 21 mycorrhizal and 20 non-mycorrhizal individuals were harvested,

and their oven-dried total biomass was measured. To compare the size of the experimental plants

at the beginning of the experiment (prior to pathogen exposure), we created a metric

approximating the total leaf area of each plant. We did this by first applying the length and width

of the plant's longest leaf to the equation yielding the area of an oval (*Pl. lanceolata* leaves

resemble long, narrow, pointed ovals), then multiplying the area of the longest leaf by the total

number of leaves on the plant. Heavy rains during the 4th and 5th surveys resulted in missing

infection data for some field populations (identifying symptoms caused by *Po. plantaginis* on

wet leaves is challenging) and likely strongly influenced infection in the 6th and 7th surveys in all

populations (as heavy rain washes spores away from infected leaf tissue and damages spore

viability; Sivapalan, 1993). Because of this, we focus here on infection data from the 3rd survey,

which coincides with the completion of one 14-day initial infection and reproduction cycle of the

pathogen. Using data from this survey, we calculated the infection status (0 / 1, infected or

uninfected) and infection severity (i.e., the proportion of infected leaves) of each experimental

plant. Plants were considered infected if one or more leaves showed signs of infection.

Statistical methods

Do growth effects due to mycorrhizal association vary among host populations and maternal

genotypes?

To test whether the growth of healthy individuals (prior to pathogen exposure) was

explained by association with mycorrhizal fungi, host genetic origin, or an interaction between

these factors, we built two multiple linear regression models. To estimate the effect of genetic

origin on plant growth, seed origin population was included as an explanatory factor in one

model, while maternal genotype (without respect to population) was included in the second

model. In each model, mycorrhizal treatment (mycorrhizal-inoculated or non-mycorrhizal

control) was also included as an explanatory factor, as was an interaction term between

mycorrhizal treatment and plant genetic origin. We used a log-transformed metric of plant size

(i.e., an estimate of total leaf area prior to pathogen exposure) as the response variable. We also

tested whether mycorrhizal and non-mycorrhizal individuals varied in size (i.e., in oven-dried

total biomass) at the end of the experiment (i.e., post-pathogen exposure) using a separate linear

regression model (this model was applied to the subset of plants for which these data were

available; plant genetic origin was not included in this model due to low replication). All

statistical tests in our study were conducted in the R statistical environment (R Core Team,

2021).

Does association with mycorrhizal fungi influence host infection risk?

To test whether host infection risk upon exposure to natural epidemics is explained by

association with mycorrhizal fungi, plant genetic origin, or an interaction between these factors,

we built a generalized linear mixed-effects model. In this model, we used plant infection risk (0 /

1) at the end of the initial infection cycle (i.e., ~14 days post-pathogen exposure) as the binomial

response variable. We also included mycorrhizal treatment, seed origin population, and their

interaction as fixed explanatory variables in the model (maternal genotype could not be

examined in a similar manner due to low replication of each genotype × mycorrhizal treatment

combination in each infection category). In addition, field epidemic population was included as a

random effect in this model. All linear (and generalized linear) mixed-effects models in our

study were performed using the lme4 package (Bates *et al*., 2015) in conjunction with the

lmerTest package (Kuznetsova *et al*., 2017) in R.

We then tested the effect of host size (prior to pathogen exposure) on infection risk

within each mycorrhizal treatment group independently, using separate generalized logistic

regression models (to overcome strong multicollinearity between host size and mycorrhizal

treatment). Two models were used to examine the interacting effects of host size and seed origin

population on infection risks: first in mycorrhizal-inoculated individuals, and then separately in

non-mycorrhizal individuals. In each model, plant infection status at the end of the initial

infection cycle was used as the binomial response variable, while the log-transformed metric of

plant size (prior to pathogen exposure) was included as a fixed effect, as well as seed origin

population, and an interaction term between these two factors. Field epidemic population was not

included as a random effect in these models for reasons of model fit.

Upon infection, does association with mycorrhizal fungi affect disease severity?

To test whether disease severity in infected hosts is influenced by association with

mycorrhizal fungi, plant genetic origin, or an interaction between these factors, we built a

generalized linear mixed-effects model. In this model, we used the proportion of infected leaves

on an infected individual as the response variable (with weights set to the total number of leaves

on the individual, and with family set to binomial). We included mycorrhizal treatment, seed

origin population, and their interaction as fixed explanatory variables (the effect of maternal

genotype could not be examined in a similar model due to insufficient power). In addition, field

epidemic population was included as a random effect in this model. Non-significant interaction

terms were removed from the final model.

We then tested for the effect of host size on the disease severity experienced by infected

individuals using two generalized linear mixed-effects models (to overcome strong

multicollinearity between host size and mycorrhizal treatment). Two models were used to

examine the interacting effects of host size and seed origin population on disease severity (i.e.,

the proportion of infected leaves), first in infected mycorrhizal-inoculated individuals and then in

infected non-mycorrhizal individuals (to overcome strong multicollinearity between host size

and mycorrhizal treatment). In each model, the log-transformed metric of plant size (prior to

pathogen exposure) was included as a fixed effect, and field epidemic population was included

as a random effect whenever possible (i.e., without compromising model fit). Non-significant

interaction terms were removed from the final models. In the model examining infected

mycorrhizal individuals, seed origin population was not a significant predictor and did not

interact with size, so it was included instead as a random effect.

Are growth and defensive effects from mycorrhizal association linked in host genotypes?

To test for a relationship between the magnitude of the growth and defensive effects

received in the host genotypes due to association with mycorrhizal fungi, we built a linear

regression model. First, we quantified the magnitude of the growth effect due to association with

mycorrhizal fungi (in healthy plants, prior to pathogen exposure) for each maternal genotype. To

do this, we extracted the estimated coefficients of the difference in host size (i.e., estimated log-

leaf area, in cm^2) among the mycorrhizal and non-mycorrhizal treatments for each maternal

genotype from the multiple regression model used to test for differences in host size among

mycorrhizal treatments, maternal genotypes, and their interaction (i.e., the same model specified

in the first section of the statistical methods). Second, we calculated the magnitude of the defense

benefit due to association with mycorrhizal fungi for each genotype (when exposed to the

pathogen in the field epidemic experiment). To do this, we extracted the estimated coefficients

representing the difference in the proportion of infected leaves (in infected individuals only)

among the mycorrhizal and non-mycorrhizal treatments for each maternal genotype from the

generalized linear mixed-effects model used to test for differences among these factors (i.e., the

model specified in the third section of the statistical methods; a similar model representing the

difference in infection rates in the mycorrhizal and non-mycorrhizal treatments for each maternal

genotype could not be fit due to convergence issues). Finally, we built a linear regression model

that linked the coefficients for the difference in infection severity when the genotypes were

associated with mycorrhizal fungi (as the response variable) with the coefficients for the

difference in host size when the genotypes were associated with mycorrhizal fungi (as the

explanatory variable). Here and in the section below, coefficient estimates, standard errors, and

p-values for contrasts between mycorrhizal treatment groups within each genotype were obtained

using the lsmeans package (Lenth, 2016).

Are host disease susceptibility and mycorrhizal-derived defensive effects linked in host

genotypes?

To test for a relationship between the magnitude of baseline disease susceptibility in the

host genotypes (in the absence of mycorrhizal fungi) and the magnitude of the defensive effects

received in the host genotypes (due to association with mycorrhizal fungi), we built a linear

regression model. First, we quantified the magnitude of baseline disease susceptibility as the

estimated coefficient representing the proportion of infected leaves in infected individuals (in log

odds ratio scale) when the genotype was not associated with mycorrhizal fungi. Next, defensive

effects were quantified as the change in the proportion of infected leaves (in infected individuals)

when the genotype was associated with mycorrhizal fungi (as in the section above). Thus,

baseline disease susceptibility and defensive effects due to mycorrhizal association were

quantified using the effect size coefficients resulting from models contrasting the genotypes

(with epidemic field site included as a random effect in both models). Finally, we built a linear

regression model that linked the coefficients for the difference in infection severity when the

genotypes were associated with mycorrhizal fungi (as the response variable) with the coefficients

for the difference in baseline disease susceptibility when the genotypes were associated with

mycorrhizal fungi (as the explanatory variable).

Results

Do growth effects due to mycorrhizal association vary among host populations and

maternal genotypes?

We found that mycorrhizal association often provided growth benefits to hosts, but that

these benefits varied in magnitude among host populations and genotypes. Prior to pathogen

exposure, mycorrhizal individuals grew larger (i.e., had more total leaf area) than non-

mycorrhizal individuals in all six seed origin populations (Fig. 1A, Table S1; $p < 0.001$, $n = 282$

plants), and in 29 of the 30 maternal genotypes (Fig. 1B, Table S1; $p < 0.001$, $n = 282$ plants).

Yet, the strength of mycorrhizal growth benefits received varied among seed origin populations

and maternal genotypes: there were significant interactions between mycorrhizal status and host

genetic origin in determining host growth, at both the population (Fig. 1C & Table S1; $F = 3.31$,

$p = 0.006$, $n = 282$ plants) and genotype (Fig. 1D & Table S1; $F = 3.06$, $p < 0.001$, $n = 282$

plants) level. Growth effects in the genotypes ranged from 8 – 67.9 % (among-genotype mean:

32.9 ± 12.9 %) increases in host size (i.e., estimated total leaf area) when associated with

mycorrhizal fungi. In addition, host populations and maternal genotypes varied in their baseline

levels of growth, regardless of mycorrhizal status (Table S1; $p < 0.001$ (seed origin population),

$p < 0.001$ (maternal genotype); $n = 282$ plants). After pathogen exposure in natural field

conditions, host size continued to be linked tightly to mycorrhizal association: the harvest

biomass of mycorrhizal individuals was increased relative to non-mycorrhizal individuals (Fig.

S2 & Table S2; $p < 0.001$, $R^2 = 0.66$, $F = 79.65$ $_{(d.f.: 1, 39)}$, $n = 41$ plants).

Does association with mycorrhizal fungi influence host infection risk?

Upon exposure to naturally-occurring field epidemics, we found that infection risk was

influenced by an interaction between the mycorrhizal status and genetic origin of the host. The

effect of mycorrhizal association on infection risk varied among the six seed origin populations:

mycorrhizal association greatly increased infection risk in one seed origin population (2719),

while infection risk was similar among mycorrhizal and non-mycorrhizal individuals in the five

remaining seed origin populations (Fig. 2A & Table S3; myc. assoc. × seed origin pop.: $p = 0.01$,

$F = 1.33$, $n = 282$ plants). In addition, we found that host size (Fig. 2B) and seed origin

population interacted to influence the likelihood of infection within groups of mycorrhizal and

non-mycorrhizal individuals (Table S4). In non-mycorrhizal individuals, host size was linked to

increases in infection risk in two out of the six seed origin populations (Pop. 877 and Pop. 1553)

(Table S4 & Fig. S3A). In mycorrhizal individuals, host size was linked to increased infection

risk in only one seed origin population (Pop. 370) (Table S4 & Fig. S3B; $p = 0.04$, z = 2.09, $n =$

141 plants).

Upon infection, does association with mycorrhizal fungi affect disease severity?

Among infected individuals, we found that the severity of disease experienced during the

peak of the epidemic was influenced by both the mycorrhizal status and the genetic origin of the

host. Mycorrhizal individuals had less severe infections (i.e., had reduced proportions of infected

leaves) than non-mycorrhizal individuals (Fig. 3A & Table S5; $p < 0.001$, $F = 104.65$, $n = 205$

plants). In addition, seed origin population influenced the proportion of leaves infected on the

host (Fig. 3A & Table S5; $p = 0.004$, $F = 3.59$, $n = 205$ plants). We also found that host size (just

prior to pathogen exposure) affected disease severity differently in mycorrhizal versus non-

mycorrhizal plants (Table S6). Larger mycorrhizal individuals had lower proportions of their

leaves infected than smaller mycorrhizal individuals (Fig. 3B & Table S6; $p = 0.01$, F = 7.52, $n =$

109 plants), and this relationship was consistent across mycorrhizal plants from different genetic

origins. In contrast, non-mycorrhizal individuals from most genetic origins had similar

proportions of infected leaves regardless of their size (Fig. 3B & Table S6; $p = 0.06$, $F = 0.45$, n

$= 96$ plants), though there was a positive relationship between size and disease severity in one

seed origin population (Pop. 1915) (Fig. S4 & Table S6; $p = 0.04$, $F = 1.17$, $n = 96$ plants).

Are growth and defensive effects from mycorrhizal association linked in host genotypes?

We did not find a clear relationship between the growth and defensive benefits conferred

by mycorrhizal fungi across the maternal genotypes (Table S7; $p = 0.39$, $F = 0.76$, $n = 30$

maternal genotypes). Growth effects due to mycorrhizal fungi (quantified in healthy individuals,

prior to pathogen exposure; and which were positive in 29 out of the 30 genotypes) varied in

magnitude independently of defensive effects (quantified from infected individuals, post-

pathogen exposure); defensive effects varied in both strength and directionality among the

genotypes (Table S7). Lack of a relationship between the two types of potential mycorrhizal

effects (growth and/or defensive) was evident when examining changes in the proportion of

infected leaves when the genotypes were associated with mycorrhizae (Table S7).

Are host disease susceptibility and mycorrhizal-derived defensive effects linked in host

genotypes?

We found a strong, positive relationship between baseline host susceptibility to infection

(measured in non-mycorrhizal individuals) and the magnitude of the defensive effect received

from mycorrhizal association in the maternal genotypes (Fig 4 & Table S8; $p < 0.001$, $F = 22.65$,

$n = 30$ maternal genotypes). In general, maternal genotypes that suffered more severe infections

without mycorrhizal fungi experienced greater relief from infections with mycorrhizal fungi

(Table S8). This pattern was evident when considering the change in the proportion of infected

leaves when the genotypes were associated with mycorrhizae (Table S8).

<u>Discussion</u>

While both plant-associated pathogenic and mutualistic microbes are ubiquitous across

ecosystems, how they interact to determine disease risk in natural, genetically diverse

populations is not known. In a natural epidemic field experiment, we found that association with

a root microbial mutualist influenced infection by an aboveground pathogen in wild host

populations. Mycorrhizal symbiosis provided an array of genotype-specific growth and defensive

benefits to host plants, altering patterns of host growth and infection with a foliar fungal

pathogen. Mycorrhizal fungi have been shown to protect agricultural plants against belowground

(Azcón-Aguilar & Barea, 1996) and foliar (Fiorilli *et al*., 2018, Pozo de la Hoz *et al*., 2021)

pathogens in controlled laboratory conditions, but this is the first evidence of mycorrhizal effects

on pathogen dynamics in a genetically diverse wild host, and under natural ecological and

epidemic conditions. In addition to benefits, we found that association with mycorrhizal fungi

introduced occasional risks to host plants, as we observed an increased likelihood of infection in

some host populations. Association with mycorrhizal fungi altered the underlying relationships

between growth and pathogen susceptibility in the host individuals, allowing them to grow larger

without increasing infection risk or load. Moreover, the strength and directionality of

mycorrhizal effects depended on host population and/or genotype, with more susceptible

genotypes receiving stronger protective effects. Hence, this is some of the first evidence of

genotype-specificity of host infection risk mediation by a third interacting species. These results

underscore how genotype-specific interactions between hosts, microbial mutualists, and

pathogens combine to determine host fitness in genetically diverse natural populations.

Though mycorrhizal fungi provided growth benefits to hosts of almost every genetic

origin in our study, the magnitude of these benefits varied widely among host populations and

maternal genotypes. Due to the positive relationship between plant size and survival, growth is

often intrinsically linked to host fitness (Harper & White, 1974). More specifically, our results

point to the importance of mycorrhizal fungi in determining host fitness (Rasmussen *et al*., 2017,

Qin *et al*., 2021) within and among genetically variable plant populations. In addition, pathogen

susceptibility varied among hosts of different genetic origins in our study, consistent with other

studies in this pathosystem (Laine, 2005, Laine, 2007, Tack *et al*., 2014, Susi & Laine, 2015) and

in the broader wild plant disease literature (Carlsson-Granér, 1997, Price *et al*., 2004). Together,

these results provide evidence for the importance of genotype in mediating host-microbe

interactions (Eck *et al*. 2019, Sallinen *et al*., 2020), through more generalist interactions can also

certainly occur (Gilbert & Webb, 2007, Halbritter *et al*., 2012, Hersh *et al*., 2012).

Building upon such studies, we also show evidence of changes in host-parasite

interactions due to association with a microbial mutualist. Protective symbionts are known from

the plant and animal literature, and are thought to play an important role in determining host

fitness (Brownlie & Johnson, 2009; May & Nelson, 2014; King *et al*., 2016; Sochard *et al*.,

2020). Studies from agricultural plant species have shown that association with mycorrhizal

fungi can reduce the negative effects of pathogens (Norman *et al*., 1996; Pozo *et al*., 2002; Li *et*

al., 2010; Song *et al*., 2015; Berdeni *et al*., 2018), and theoretical studies speculate as to how

mycorrhizal benefits might counteract reductions in host fitness due to pathogens and affect

diversity in natural communities (Bachelot *et al*., 2015). However, symbiotic relationships in

cultivars may differ from those of wild plants (Xing *et al*., 2012); thus, it is not straightforward to

predict responses in wild populations from controlled agricultural trials. In this study,

mycorrhizal association mainly protected wild hosts against further development of established

infections, rather than preventing infection by the pathogen entirely. Mycorrhizal association had

little influence on infection risk in most host populations in our study, and even increased

infection risk in one seed origin population (highlighting the genotype-dependency of such

interactions). Instead, protective effects from mycorrhizal fungi, which resulted in reductions in

disease severity in mycorrhizal hosts from all genetic origins, occurred by altering the underlying

relationships between host growth and pathogen susceptibility.

In many free-living organisms, an evolutionary trade-off exists between growth and

defense (Herms & Mattson, 1992). In our experiment, insights in to how microbial mutualists

may affect this trade-off can be gained by examining how growth influences infection patterns in

hosts with and without mutualist association. In the absence of mycorrhizal mutualists, larger

hosts were more likely to become severely infected by the wind-dispersed pathogen (as increased

surface area increases the likelihood that hosts encounter pathogen spores), and the strength of

the relationship between growth and pathogen susceptibility varied among host populations. In

contrast, though mutualist association caused plants to grow larger, mycorrhizal plants from most

host populations grew larger without increasing their infection risk or load. Thus, mycorrhizal

plants appear to be benefitting from a more quantitative than qualitative mechanism of pathogen

resistance (Lo Iacono *et al*., 2012), which allows them to grow large while being relieved of the

increased infection risks that normally accompany growth. Molecular studies have shown that

mycorrhizal fungi can elicit induction of defence mechanisms (priming) in the absence of

pathogens, followed by a faster and stronger activation of pathogenesis-related genes upon

pathogen infection (Pozo & Azcón-Aguilar, 2007). Mycorrhizal protective effects in this wild

pathosystem best correspond to the latter, rather than the former, mechanism: mycorrhizal hosts

were infected, but not repeatedly (i.e., the pathogen did not spread to new leaves as readily as in

non-mycorrhizal plants). Other potential mechanisms include improvements in host tolerance to

pathogens due to mycorrhizal association, such as could be expected if mycorrhizal fungi, e.g.,

improve host nutritional status or increase leaf toughness, making foliar pathogen spread more

difficult (Meier & Hunter, 2018).

In addition to offsetting the costs of increased infection risk from growing large, our

findings indicate that mycorrhizal association could help compensate for lack of innate host

resistance. We found that the magnitude of mycorrhizal protective effects was linked to pathogen

susceptibility in the host genotypes: host genotypes that were more susceptible to severe

infections by the pathogen (in the absence of the mutualist) received stronger disease protection

effects from association with the mutualist. This could indicate that host genotypes experience a

trade-off in investment in genetic resistance vs. mycorrhizal-mediated resistance. Thus,

mycorrhizal association may have been selected for and maintained in host populations partially

because it increases the fitness of susceptible host genotypes. In this way, mycorrhizal protective

effects could contribute to the maintenance of diversity in pathogen genetic resistance within and

among host populations, by helping to maintain susceptible genotypes, which have not invested

heavily in defense. In contrast, the magnitude of the growth-enhancing and pathogen-protective

benefits conferred by mycorrhizal fungi were not linked in the host genotypes, suggesting that

the delivery of these two types of benefits does not necessarily create a resource allocation

conflict. These results highlight the range and variety of potential mycorrhizal effects on hosts,

and suggest that different selection processes may act to determine the magnitude and direction

of mycorrhizal growth vs. defensive effects.

Together, our results show that mutualism with mycorrhizal fungi produces host

genotype-specific benefits and alters infection risks from a pathogen during natural epidemics in

host populations. Altered growth and infection risk due to mutualist species may cascade to

affect larger ecological patterns of abundance and distribution of the associated organisms, as

well as ecosystem processes (Brown *et al*., 2001). The presence of genotype-specificity in host-

pathogen-mutualist interactions also highlights the importance of maintaining genetic diversity

within and among wild plant populations. Characterizing the potential for and frequency of

specialized interactions between microbial pathogens and mutualists and their host organisms

under natural conditions, and using the natural temporal sequence at which the various

associations are established, will also allow for more precise modelling of ecological patterns in

natural communities. In addition, the generality of growth and defensive effects due to

mutualism should also affect the co-evolutionary trajectories of the associated organisms,

because we expect organisms to be adapted to events that are common over evolutionary time

spans (van Dam & Heil, 2011). Moreover, we are also beginning to acknowledge the importance

of direct and indirect interactions among microbes within hosts in determining host fitness and

parasite population dynamics (Kemen, E, 2014; Kemen, A, *et al*., 2015; Kroll *et al*., 2017).

Belowground soil and rhizosphere processes may affect aboveground interactions with

pathogens and herbivores, and vice versa (van der Putten *et al*., 2001; Wardle *et al*., 2004; Frew,

2021), and mutualists and antagonists may act simultaneously to determine host fitness (Bezemer

& van Dam, 2005). Understanding how mutualism-derived protection alters host susceptibility

and parasite interactions will be important for understanding, predicting, and managing disease

in both natural communities and in agricultural settings.

Acknowledgements

We would like to thank Oulu University Botanic Gardens for providing greenhouse facilities, Riitta Ovaska for assistance during the field experiment, and Dr. Mauritz Vestberg for the fungal isolates. This work was funded by grants from the European Research Council (Consolidator Grant RESISTANCE: LS8 & ERC-2016-COG) and the Academy of Finland (MULTA #327222 & #296686) to A-LL, and the University of Zurich (Forschungskredit #K-74333-03) to JLE.

Author Contributions

A-LL and M-MK designed and conducted the experiment. JLE and A-LL analyzed the data. JLE wrote the first draft of the manuscript. All authors contributed to and approved the final version.

Data Availability

The data that support the findings of this study are openly available in the Dryad Digital Repository at http://doi.org/[doi], reference number [reference number].

References

- **Azcón-Aguilar, C, Barea, JM. 1996.** Arbuscular mycorrhizas and biological control of soil-borne plant pathogens – an overview of the mechanisms involved. *Mycorrhiza* **6**: 457-464.

- **Bachelot, B, Uriarte, M, McGuire, K. 2015.** Interactions among mutualism, competition, and predation foster species coexistence in diverse communities. *Theoretical Ecology* **8**: 297-312.

- **Barea, JM, Azcón, R, Azcón-Aguilar, C. 2002.** Mycorrhizosphere interactions to improve plant fitness and soil quality. *Antonie van Leeuwenhoek* **81**: 343-351.

- **Bates, D, Maechler, M, Bolker, B, Walker, S. 2015**. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* **67(1)**: 1-48.

- **Berdeni, D, Cotton, TEA, Daniell, TJ, Bidartondo, MI, Cameron, DD, Evans, KL. 2018.** The effects of arbuscular mycorrhizal fungal colonization on nutrient status, growth, productivity, and canker resistance of apple (*Malus pumila*). *Frontiers in Microbiology* **9:1461**. doi: 10.3389/fmicb.2018.01461

- **Bever, JD, Mangan, SA, Alexander, HM. 2015.** Maintenance of plant species diversity by pathogens. *Annual Review of Ecology, Evolution, and Systematics* **46**: 305-325.

- **Bezemer, TM, van Dam, NM. 2005**. Linking aboveground and belowground interactions via induced plant defenses. *Trends in Ecology and Evolution* **20(11)**: 617-624.

- **Bos, M. 1992**. Gene flow characters and population structure in *Plantago lanceolata*. In: Kuiper, PJC, Bos, M eds. *Plantago: A Multidisciplinary Study.* Berlin, DE: Springer-Verlag, 222-231.

- **Brown, JH, Whitham, TG, Ernest, SKM, Gehring, CA**. **2001.** Complex species interactions and the dynamics of ecological systems: long-term experiments. *Science* **293**: 643-650.

- **Brown, JKM**. **2003.** A cost of disease resistance: paradigm or peculiarity? *Trends in Genetics* **19(12)**: 667-671.

- **Brownlie, JC, Johnson, KN. 2009.** Symbiont-mediated protection in insect hosts. *Trends in Microbiology* **17(8)**: 348-354.

- **Burdon, JJ, Thrall, PH. 2014.** What have we learned from studies of wild plant-pathogen associations?—the dynamic interplay of time, space and life-history. *European Journal of Plant Pathology* **138,** 417–429.

• **Bushnell, WR. 2002.** The role of powdery mildew research in understanding host-pathogen interaction: past, present, and future. In: Bélanger, RR, Bushnell, WR, Dik, AJ, Carver, LW eds. *The Powdery Mildews – a Comprehensive Treatise*. St. Paul, USA: The American Phytopathology Society, 1-12.

• **Cameron, DD, Neal, AL, van Wees, SCM, Ton, J. 2013.** Mycorrhiza-induced resistance: more than the sum of its parts? *Trends in Plant Science* **18(10)**: 539-545.

• **Carlsson-Granér, U. 1997.** Anther-smut disease in *Silene dioica*: variation in susceptibility among genotypes and populations, and patterns of disease within populations. *Evolution* **51(5)**: 1416-1426.

• **van Dam, NM, Heil, M. 2010.** Multitrophic interactions below and above ground: *en route* to the next level. *Journal of Ecology* **99(1)**: 77-88.

• **Delavaux, CS, Smith-Ramesh, LM, Kuebbing, SE. 2017.** Beyond nutrients: a meta-analysis of the diverse effects of arbuscular mycorrhizal fungi on plants and soils. *Ecology* **98(8)**: 2111-2119.

• **Eck, JL, Stump, SM, Delavaux, CS, Mangan, SA, Comita, LS. 2019.** Evidence of within-species specialization by soil microbes and the implications for plant community diversity. *PNAS*. doi: 10.1073/pnas.1810767116

• **Eck, JL, Barrès, B, Soubeyrand, S, Siren, J, Numminen, E, Laine, A-L.** Strain diversity and spatial distribution are linked to epidemic dynamics in host populations. Accepted manuscript at *The American Naturalist*.

• **Fiorilli, V, Vannini, C, Ortolani, F, Garcia-Seco, D, Chiapello, M, Novero, M, Domingo, G, Terzi, V, Morcia, C, Bagnaresi, P, *et al*. 2018.** Omics approaches revealed how arbuscular mycorrhizal symbiosis enhances yield and resistance to leaf pathogen in wheat. *Scientific Reports* **8:** 9625. doi: 10.1038/s41598-018-27622-8

• **Frew, A, 2021.** Aboveground herbivory suppresses the arbuscular mycorrhizal symbiosis, reducing phosphorus uptake. *Applied Soil Ecology* **168**: 104133. doi: 10.1016/j.apsoil.2021.104133

• **Gilbert, GS, Webb, CO. 2007.** Phylogenetic signal in plant pathogen-host range. *PNAS* **104(12)**: 4979-4983.

• **Gosling, P, Hodge, A, Goodlass, G, Bending, GD. 2006.** Arbuscular mycorrhizal fungi and
organic farming. *Agriculture, Ecosystems, and Environment* **113**: 17-35.

• **Grman, E. 2012**. Plant species differ in their ability to reduce allocation to non-beneficial
arbuscular mycorrhizal fungi. *Ecology* **93(4):** 711-718.

• **Hafer-Hahmann, NC, Vorburger, C. 2021.** Positive association between the diversity of
symbionts and parasitoids of aphids in field populations. *Ecosphere* **12(1)**. doi:
10.1002/ecs2.3355

• **Halbritter, AH, Carroll, GC, Güsewell, S, Roy, BA. 2012.** Testing assumptions of the
enemy release hypothesis: generalist versus specialist enemies of the grass *Brachypodium*
sylvaticum. *Mycologia* **104(1)**: 34-44.

• **Hanski, I. 1999.** *Metapopulation Ecology*. Oxford University Press, Oxford, UK.

• **Harper JL, White J. 1974.** The demography of plants. *Annual Review of Ecology and*
Systematics **5:** 419–463.

• **Hartnett, DC, Wilson, GWT. 2002.** The role of mycorrhizas in plant community structure
and dynamics: lessons from grasslands. In: Smith SE, Smith, FA eds. *Diversity and*
integration in mycorrhizas. Developments in plant and soil sciences **94**. Dordrecht, NL:
Springer. doi: 10.1007/978-94-017-1284-2_30

• **van der Heijden, MGA, Bardgett, RD, van Straalen, NM. 2008.** The unseen majority: soil
microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecology*
Letters **11**: 296-310.

• **Herms DA, Mattson, WJ. 1992.** The dilemma of plants: to grow or defend. *The Quarterly*
Review of Biology **67(3)**: 283–335.

• **Hersh, MH, Vilgalys, R, Clark, JS. 2012.** Evaluating the impacts of multiple generalist
pathogens on temperate tree seedling survival. *Ecology* **93(3)**: 511-520.

• **Hoeksema, JD, Chaudhary, VB, Gehring, CA, Johnson, NC, Karst, J, Koide, RT,**
Pringle, A, Zabinski, C, Bever, JD, Moore, JC et al. 2010. A meta-analysis of context-
dependency in plant response to inoculation with mycorrhizal fungi. *Ecology Letters* **13**: 394-
407.

• **Hohmann, P, Messmer, MM. 2017.** Breeding for mycorrhizal symbiosis: focus on disease
resistance. *Euphytica* **213(113)**. doi: 10.1007/s10681-017-1900-x

• **Johansson, JF, Paul, LR, Finlay, RD. 2004.** Microbial interactions in the mycorrhizosphere
and their significance for sustainable agriculture. *FEMS Microbiology Ecology* **48**: 1-13.

• **Johnson, NC, Graham, JH, Smith, FA**. **1997**. Functioning of mycorrhizal associations
along the mutualism-parasitism continuum. *New Phytologist* **135**: 575-585.

• **Johnson, D, Vandenkoornhuyse, PJ, Leake, JR, Gilbert, L, Booth, RE, Grime, JP,**
Young, JPW, Read, DJ. 2003. Plant communities affect arbuscular mycorrhizal fungal
diversity and community composition in grassland microcosms. *New Phytologist* **161**: 503-
515.

• **Jones, JDG, Dangl, JL. 2006.** The plant immune system. *Nature* 444(16). doi:
10.1038/nature05286

• **Jones, MD, Smith, SE. 2004.** Exploring functional definitions of mycorrhizas: are
mycorrhizas always mutualisms? *Canadian Journal of Botany* **82**: 1089-1109.

• **Jousimo, J, Tack, AJM, Ovaskainen, O, Mononen, T, Susi, H, Tollenaere, C, Laine, A-**
L. 2014. Ecological and evolutionary effects of fragmentation and disease dynamics. *Science*
344(6189): 1289-1293.

• **Jung, SC, Martinez-Medina, A, Lopez-Raez, JA, Pozo, MJ. 2012.** Mycorrhiza-induced
resistance and priming of plant defenses. *Journal of Chemical Ecology* **38**: 651-664.

• **Kemen, E. 2014.** Microbe-microbe interactions determine oomycete and fungal host
colonization. *Current Opinion in Plant Biology* **20**: 75-81.

• **Kemen, AC, Agler, MT, Kemen, E. 2015.** Host-microbe and microbe-microbe interactions
in the evolution of obligate plant parasitism. *New Phytologist* **206**: 1207-1228.

• **King, KC, Brockhurst, MA, Vasieva, O, Paterson, S, Betts, A, Ford, SA, Frost, CL,**
Horsburgh, MJ, Haldenby, S, Hurst, GDD. 2016. Rapid evolution of microbe-mediated
protection against pathogens in a worm host. *The ISME Journal* **10**: 1915-1924.

• **Klironomos, J. 2003.** Variation in plant response to native and exotic arbuscular mycorrhizal
fungi. *Ecology* **84(9)**: 2292-2301.

• **Koide, RT, Dickie, IA. 2002.** Effects of mycorrhizal fungi on plant populations. In: Smith
SE, Smith, FA eds. *Diversity and integration in mycorrhizas. Developments in plant and soil*
sciences **94**. Dordrecht, NL: Springer. doi: 10.1007/978-94-017-1284-2_30

• **Koricheva, J, Gange, AC, Jones, T. 2009.** Effects of mycorrhizal fungi on insect
herbivores: a meta-analysis. *Ecology* **90(8)**: 2088-2097.

• **Kroll, S, Agler, MT, Kemen, E. 2017.** Genomic dissection of host-microbe and microbe-
microbe interactions for advanced plant breeding. *Current Opinion in Plant Biology* **36**: 71-
78.

• **Kuznetsova, A, Brockhoff, PB, Christensen, RHB. 2017.** lmerTest package: tests in linear
mixed effects models. *Journal of Statistical Software* **82(13)**. doi: 10.18673/jss.v082.i13.

• **Laine, A-L. 2004.** Resistance variation within and among host populations in a plant
pathogen metapopulation: implications for regional pathogen dynamics. *Journal of Ecology*
92(6): 990-1000.

• **Laine, A-L. 2005**. Spatial scale of local adaptation in a plant-pathogen metapopulation.
Journal of Evolutionary Biology **18**: 930-938.

• **Laine, A-L. 2006.** Evolution of host resistance: looking for coevolutionary hotspots at small
spatial scales. *Proceedings: Biological Sciences* **273(1584)**: 267-273.

• **Laine, A-L. 2007.** Pathogen fitness components and genotypes differ in their sensitivity to
nutrient and temperature variation in a wild plant-pathogen association. *Journal of*
Evolutionary Biology **20**: 2371-2378.

• **Laine, A-L. 2008**. Temperature-mediated patterns of local adaptation in a natural plant-
pathogen metapopulation. *Ecology Letters* **11**: 327-337.

• **Laine, A-L, Hanski, I. 2006.** Large-scale dynamics of a specialist plant pathogen in a
fragmented landscape. *Journal of Ecology* **94(1)**: 217-226.

• **Laine, A-L, Tellier, A. 2008**. Heterogeneous selection promotes polymorphism maintenance
in host-parasite interactions. *Oikos* **117**: 1281-1288.

• **Laine, A-L, Burdon, JJ, Dodds, PN, Thrall, PH. 2011.** Spatial variation in disease
resistance: from molecules to metapopulations. *Journal of Ecology* **99**: 96-112.

• **Lenth, RV. 2016**. Least-squares means: the R package lsmeans. *Journal of Statistical*
Software **69(1)**: 1-33.

• **Li, Y, Yanagi, A, Miyawaki, Y, Okada, T, Matsubara, Y. 2010.** Disease tolerance and
changes in antioxidative abilities in mycorrhizal strawberry plants. *Journal of the Japanese*
Society for Horticultural Science **79(2)**: 174-178.

• **Lo Iacono, G, van den Bosch, F, Paveley, N. 2012.** The evolution of plant pathogens in
response to host resistance: factors affecting the gain from deployment of qualitative and
quantitative resistance. *Journal of Theoretical Biology* **304**: 152-163.

• **May, G, Nelson, P. 2014.** Defensive mutualisms: do microbial interactions within hosts
drive the evolution of defensive traits? *Functional Ecology* **28**: 356-363.

• **Meier, AR, Hunter, MD. 2018.** Arbuscular mycorrhizal fungi mediate herbivore-induction
of plant defenses differently above and belowground. *Oikos* **127**: 1759-1775.

• **Newsham, KK, Fitter, AH, Watkinson, AR. 1995.** Arbuscular mycorrhiza protect an
annual grass from root pathogenic fungi in the field. *Journal of Ecology* **83(6)**: 991-1000.

• **Norman, JR, Atkinson, D, Hooker, JE. 1996.** Arbuscular mycorrhizal fungal-induced
alteration to root architecture in strawberry and induced resistance to the root pathogen
Phytophthora fragariae. *Plant and Soil* **185**: 191-198.

• **Oliver, KM, Smith, AH, Russell, JA. 2014.** Defensive symbiosis in the real world –
advancing ecological studies of heritable, protective bacteria in aphids and beyond.
Functional Ecology **28**: 341-355.

• **Öpik, M, Moora, M, Liira, J, Zobel, M. 2006.** Composition of root-colonizing arbuscular
mycorrhizal fungal communities in different ecosystems around the globe. *Journal of*
Ecology **94**: 778-790.

• **Ovaskainen, O, Laine, A-L. 2006**. Inferring evolutionary signals from ecological data in a
plant-pathogen metapopulation. *Ecology* **87(4)**: 880-891.

• **Penczykowski, R, Walker, E, Soubeyrand, S**, **Laine, A-L**. **2015**. Linking winter conditions
to large-scale disease dynamics in a wild plant-pathogen metapopulation. *New Phytologist*,
205: 1142-1152.

• **Polin, S., Simon, J-C, Outreman, Y. 2014.** An ecological cost associated with protective
symbionts of aphids. *Ecology and Evolution* **4(6)**: 836-840.

• **Pozo, MJ, Azcón-Aguilar, C. 2007.** Unraveling mycorrhiza-induced resistance. *Current*
Opinion in Plant Biology **10**: 393-398.

• **Pozo, MJ, Cordier, C, Dumas-Gaudot, E, Gianinazzi, S, Barea, JM, Azcón-Aguilar, C.**
2002. Localized versus systemic effect of arbuscular mycorrhizal fungi on defence responses
to *Phytophthora* infection in tomato plants. *Journal of Experimental Botany* **53(368)**: 525-
534.

• **Pozo de la Hoz, J, Rivero, J, Azcón-Aguilar, C, Urrestarazu, M, Pozo, MJ. 2021.**
Mycorrhiza-induced resistance against foliar pathogens is uncoupled of nutritional effects
under different light intensities. *Journal of Fungi* **7**: 402. doi: 10.3390/jof7060402

• **Pretty, J, Sutherland, WJ, Ashby, J, Auburn, J, Baulcombe, D, Bell, M, Bentley, J,**
Bickersteth, S, Brown, K, Burke, J et al. 2011. The top 100 questions of importance to the
future of global agriculture. *International Journal of Agricultural Sustainability* **8(4)**: 219-
236.

• **Price, JS, Bever, JD, Clay, K. 2004.** Genotype, environment, and genotype by environment
interactions determine quantitative resistance to leaf rust (*Coleosporium asterum*) in
Euthamia graminifolia (Asteraceae). *New Phytologist* **162**: 729-743.

• **van der Putten, WH, Vet, LEM, Harvey, JA, Wäckers, FL. 2001.** Linking above- and
belowground multitrophic interactions of plants, herbivores, pathogens, and their antagonists.
Trends in Ecology and Evolution **16(10)**: 547-554.

• **Qin, J, Geng, Y, Li, X, Zhang, C, Zhao, X, von Gadow, K. 2021.** Mycorrhizal type and
soil pathogenic fungi mediate tree survival and density dependence in a temperate forest.
Forest Ecology and Management 496: 119459. doi: 10.1016/j.foreco.2021.119459

• **Qu, L, Wang, M, Biere, A. 2021**. Interactive effects of mycorrhizae, soil phosphorus, and
light on growth and induction and priming of defense in *Plantago lanceolata*. *Frontiers in*
Plant Science **12**: 647372. doi: 10.3389/fpls.2021.647372

• **R Core Team. 2021**. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. URL: https://www.R-project.org

• **Rasmussen, PU, Amin, T, Bennett, AE, Karlssongreen, K, Timonen, S, van Nouhuys, S,**
Tack, AJM. 2017. Plant and insect genetic variation mediate the impact of arbuscular
mycorrhizal fungi on a natural plant-herbivore interaction. *Ecological Entomology* **42**: 793-
802.

• **Rasmussen, PU, Hugerth, LW, Blanchet, FG, Andersson, AF, Lindahl, BD, Tack, AJM.**
2018. Multiscale patterns and drivers of arbuscular mycorrhizal fungal communities in the
roots and root-associated soil of a wild perennial herb. *New Phytologist.* doi:
10.111/nph.15088

• **Rasmussen, PU, Chareesri, A, Neilson, R, Bennett, AE, Tack, AJM. 2019**. The impact of
dispersal, plant genotype and nematodes on arbuscular mycorrhizal fungal colonization. *Soil*
Biology and Biochemistry **132**: 28-35.

• **Rillig, MC. 2004.** Arbuscular mycorrhizae and terrestrial ecosystem processes. *Ecology*
Letters **7**: 740-754.

• **Sagar, GR, Harper, JL. 1964**. *Plantago major* L., *P. media* L. and *P. lanceolata* L. *Journal*
of Ecology **52(1)**: 189-221.

• **Sallinen, S, Norberg, A, Susi, H, Laine, A-L. 2020.** Intraspecific host variation plays a key
role in virus community assembly. *Nature Communications* **11**: 5610. doi: 10.1038/s41467-
020-19273-z

• **Salvaudon, L, Giraud, T, Shykoff, JA. 2008**. Genetic diversity in natural populations: a
fundamental component of plant-microbe interactions. *Current Opinion in Plant Biology*
11(2): 135-143.

• **Sivapalan, A. 1993.** Effects of water on germination of powdery mildew conidia.
Mycological Research **97(1)**: 71-76.

• **Smith, SE, Read, DJ. 2008.** *Mycorrhizal Symbiosis*. Cambridge, UK: Academic Press.

• **Sochard, C, Bellec, L, Simon, J-C, Outreman, Y. 2020.** Influence of "protective"
symbionts throughout the different steps of an aphid-parasitoid interaction. *Current Zoology*.
doi: 10.1093/cz/zoaa053

• **Song, Y, Chen, D, Lu, K, Sun, Z, Zeng, R. 2015**. Enhanced tomato disease resistance
primed by arbuscular mycorrhizal fungus. *Frontiers in Plant Science* **6(786)**: 1-13.

• **Soubeyrand, S, Laine, A-L, Hanski, I, Penttinen, A. 2009.** Spatiotemporal structure of
host-pathogen interactions in a metapopulation. *The American Naturalist* **174(3)**: 308-320.

• **Susi, H, Laine, A-L. 2015.** The effectiveness and costs of pathogen resistance strategies in a
perennial plant. *Journal of Ecology* 103: 303-315.

• **Tack, AJM, Laine, A-L. 2014.** Ecological and evolutionary implications of spatial
heterogeneity during the off-season for a wild plant pathogen. *New Phytologist* **202**: 297-308.

• **Tack, AJM, Hakala, J, Petäjä, T, Kulmala, M, Laine, A-L. 2014.** Genotype and spatial
structure shape pathogen dispersal and disease dynamics at small spatial scales. *Ecology*
95(3): 703-714.

• **Thompson, JN. 1994**. *The Coevolutionary Process*. Chicago, USA: The University of
Chicago Press.

• **Thrall, PH, Laine, A-L, Broadhurst, LM, Bagnall, DJ, Brockwell, J. 2011.** Symbiotic
effectiveness of rhizobial mutualists varies in interactions with native Australian legume
genera. *PLoS ONE* **6**: ge23545. doi: 10.1371/journal.pone.0023545

• **Tollenaere, C, Laine, A-L. 2013.** Investigating the production of sexual resting structures in
a plant pathogen reveals unexpected self-fertility and genotype-by-environment effects.
Journal of Evolutionary Biology **26**: 1716–1726.
• **Wardle, DA, Bardgett, RD, Klironomos, JN, Setälä, H, van der Putten, WH, Wall, DH.**
2004. Ecological linkages between aboveground and belowground biota. *Science* **304**: 1629-
1633.
• **Xing, X, Koch, AM, Jones, AMP, Ragone, D, Murch, S, Hart, MM. 2012.** Mutualism
breakdown in breadfruit domestication. *Proceedings of the Royal Society B* **279**: 1122–1130.

<u>Figures & Tables</u>

Fig. 1: Mycorrhizal growth benefits varied in magnitude among host populations and
maternal genotypes. Association with mycorrhizal fungi conferred growth benefits to host
individuals originating from six field populations (*Panel A*) and to 29 out of 30 maternal
genotypes (*Panel B*) (Table S1). Growth effects were measured as aboveground size prior to
pathogen exposure (i.e., a log-transformed estimate of total leaf area). Though host growth
almost always increased in association with mycorrhizal fungi, the magnitude of the growth
benefit received varied among the seed origin populations (*Panel C*) and among the maternal
genotypes (*Panel D*), due to an interaction between mycorrhizal association and host genetic
origin at both levels (Table S1).

Fig. 2: Infection risk varied among mycorrhizal treatments and host populations, and depended on plant size. Host infection risk upon exposure to field epidemics were influenced by an interaction between mycorrhizal association and seed origin population: mycorrhizal association greatly increased infection risk in one population, 2719 (*Panel A*; Table S3). In addition, host size influenced infection risk differently in mycorrhizal versus non-mycorrhizal plants: while the infection risk of non-mycorrhizal plants increased with size, mycorrhizal plants experienced similar infection risk, regardless of size (*Panel B*; Table S4).

824
825
Fig. 3: Disease severity was reduced in mycorrhizal plants, while host size affected infection
severity differently in mycorrhizal versus non-mycorrhizal plants. *Panel A*: In a natural
epidemic field experiment, the disease severity experienced by infected plants was influenced by
both mycorrhizal treatment and host genetic origin (though the two factors did not interact)
(Table S5; $n = 205$ plants). Mycorrhizal plants in all six seed origin populations had reduced
proportions of infected leaves, relative to non-mycorrhizal plants. Infection severity also differed
among seed origin populations, suggesting baseline differences in disease susceptibility among
populations. *Panel B*: Host size at the time of pathogen exposure was important in determining
infection severity, but different dynamics were found in mycorrhizal versus non-mycorrhizal
individuals (Table S6). Larger mycorrhizal individuals had reduced proportions of infected
leaves relative to smaller mycorrhizal individuals (Table S6; $p = 0.01$, $F = 7.52$, $n = 109$ plants),
while non-mycorrhizal individuals of all sizes had similar proportions of infected leaves (Table
S6; $p = 0.20$, $F = 0.45$, $n = 96$ plants).

840
841
Fig. 4: Host susceptibility to infection and the defensive effects of mycorrhizal fungi are
linked in the host genotypes. In a natural epidemic field experiment with plants from thirty
maternal genotypes, we found that host genotypes that were more susceptible to infection (when
not associated with mycorrhizae) received stronger disease protection effects from association
with mycorrhizal fungi (Table S8). Here, each point represents one maternal genotype,
originating from one of six seed origin populations (point colors indicate the seed origin
population). Infection severity is examined as the effect size representing the proportion of
infected leaves, calculated as a log odds ratio. On the x-axis, more positive values indicate more
severe infections and thus, higher disease susceptibility. On the y-axis, more positive values
indicate stronger mycorrhizal defensive effects (i.e., greater reductions in the proportion of
infected leaves when the genotype was associated with mycorrhizae).