Article

Ovipin: a new antimicrobial peptide from chicken eggs *Gallus gallus*

Sandra Regina dos **SANTOS**[1,2,3,*]; Antonio **MIRANDA**[1,2]; Pedro Ismael da **SILVA JUNIOR**[1,3,*]

[1] Biological Sciences (Molecular Biology) Post-Graduation Program - UNIFESP

[2] Department of Biophysics, UNIFESP, São Paulo CEP 04023-062, Brazil; amiranda@unifesp.br

[3] Laboratory for Applied Toxinology - Center of Toxins, Immune-Response and Cell Signaling

– CeTICS/CEPID, Butantan Institute, São Paulo CEP 05503-900, SP, Brazil

*Correspondence: sandrasantos111@gmail.com; Tel.: +55-15-98167-9581;

pisjr@butantan.gov.br; Tel.:+55-11-2627-9731

Abstract: The intensive and indiscriminate use of antibiotic has increased cases of microorganisms resistance and becoming a worldwide public health problem. In the last years, from natural sources such as hen's egg, have received special attention in the discovery of new bioactive compounds. This study aims to identify and characterize a new peptide from chicken egg of *Gallus gallus domesticus*. The chicken egg was subjected the extraction acid, the supernatant was prefractionated in Sep-Pak column and fractionated peptide by reverse-phase high-performance liquid chromatography (RP-HPLC). The antimicrobial activity of the fractions were evaluated through liquid growth inhibition assays. The molecular weight and amino acid sequence was determined by mass spectrometry (MS/MS), the characacterization performed by bioinformatics analysis with Peaks® tool and comparision with the NCBI and UniProt-SwisProt database. The physicochemical parameters of the samples were evaluated with online programs. One fraction named Ovipin peptide, showed antimicrobial activity against Gram-positive bacteria (*Micrococcus luteus* – MIC 1.94 μM) and Filamentous fungus (*Aspergillus niger* - MIC 31.01 μM). The minimum inhibitory concentration against *Cryptococcus neoformans* VNI (WM 148) Yeast was 22.75 μM, this microorganism an opportunistic yeast and mainly binds in immunosuppressed or immunocompromissed people. Ovipin is a hydrophoic peptide and not cause significant hemolytic effect against human erythrocytes. Ovipin primary sequence is YVSPVAIVKGLNIPL and a molecular weight of 1,581.94 Da. Ovipin shows 100% and 93.3%, respectively, sequence similarity with the fragments Apolipoprotein B of *Gallus gallus* and Apolipoprotein B of five others species of Aves. Our data suggest that Ovipin peptide could be a potential therapeutic candidate to be used an alternative against infections developed by resistant microorganisms, mainly in the fight against *Cryptococcus neoformans* opportunistic yeast.

Keywords: 1: Ovipin; 2: chicken eggs; 3: antimicrobial; 4: peptide; 5: AMPs.

1. Introduction

Immunocompromised patients presents a high risk of developing infection because in this group there is a decrease or lack the basic mechanisms of cellular defense. These patients are more susceptible to attack by microorganisms, such as bacteria or fungal infection are mainly genus *Aspergillus*, and the genera as *Mucor*, *Fusarium*, *Rhizopus*, *Petriellidium*, *Cryptococcus*, *Histoplasma*, *Coccidioides* and *Candida* (**FRARE e SILVA, 2010**).

Cryptococcosis or cryptococcal infection is a relevant problem global because affect mainly immunocompromised and immunocompetent hosts with high incidence of morbidity and mortality. *Cryptococcus* species are encapsulated yeasts and as an opportunistic microorganism, it can trigger in patients with human immunodeficiency virus (HIV) infection, solid organ transplantation, corticosteroid use, diabetes and cirrhosis. Pulmonary and central nervous system (CNS) are the well-known clinical manifestations of cryptococcosis followed by cryptococcal meningoencephalitis (**GUSHIKEN et al, 2021; ZAVALA, BADDLEY, 2020**). *Cryptococcus spp.* (approximately 30 species) has wide geographical distribution – *Cryptococcus neoformans* and *Cryptococcus gattii* are two species relationship with human disease. (**GUSHIKEN et al, 2021; HENAO-MARTÍNEZ et al., 2018**). The polysaccharide capsule, melanin production, grow at body temperature and capacity to produce extracellular enzymes are the factors which associated with the health state host, confer the success and survival of these microorganisms against the complex responses of the host's immune system (**ZAVALA, BADDLEY, 2020; ALMEIDA et al., 2017**).

The resistance antibacterial or antifungals arising from selective pressure of clinical and agricultural antifungal agent use is a serious concern because limit the options treatment, research into new antibacterial or antifungal agents or new combinations is necessary to control the advancement of cryptococcal disease or others infections disease (**BEN-AMI, KONTOYIANNIS, 2021**).

In this regard, antimicrobial peptides (AMPs) are evolutionarily conserved innate immune defense system and are produced by many organisms (**Peters et al., 2010**), such as Prokaryotes (e.g., bacteria) and Eukaryotes (e.g., protozoan, fungi, plants, insects and animals) (**Izadpanah & Gallo, 2005; Bahar & Ren, 2013**). They are widely studied by presenting a fast and broad spectrum of action against Gram-positive and Gram-negative bacteria, fungi, protozoa and viruses (**Kang et al., 2014; Andreu & Rivas, 1998; Cruz et al., 2014**). Thus presenting the ability to inhibit growth or kill multi-resistant microorganism (**Peters et al., 2010**). Antimicrobial peptides (AMPs) are small (weight <10 kDa) an in general amphipathic molecules, positively or negatively charged with amino acids sequences diversity (**Bahar & Ren, 2013; Cruz et al., 2014**). They are selective for prokaryotic membranes and act on the cell membrane-disruptive of microorganisms through electrostatic interactions and show low natural resistance. Some peptides have no action on

the cell membrane, but rather they cross this membrane and act on intracellular targets, which can interfere in crucial

processes for the growth and development of microorganisms (**Brogden, 2005**).

On the other hand, in the last years has also growth the interest in the search for bioactive molecules in food sources.

The functional foods contain amounts and types of bioactive compounds different, which provide health benefits and

may decrease the development of chronic diseases (**Gil-Chávez et al., 2013; Rutherfurd-Markwick, 2012**). As a

natural source, the chicken eggs with their diversity of structure and variety of components are increasingly becoming

targets of research in health or biotechnology (**Hancock & Sahl, 2006**). Chicken eggs is an important dietary source

because it is rich in proteins, lipids, vitamins and minerals, but which are associated with other functions such as

defense to inhibit the invasion and proliferation of pathogens (**Kovacs-Nolan et al., 2005**).

The aim of this work is to performed the bioprospecting on chicken eggs of specie *Gallus gallus domesticus*, with focus

in the discovery of new antimicrobial peptides with great potential to act against microorganisms, especially

microorganisms that have already developed some resistance mechanism.

2. Materials and Methods

2.1. Extraction acid and Pre-fractionated of Chicken eggs

In this study chicken eggs without the shell of the species *Gallus gallus domesticus* were utilized. They were purchased

from the commercial market network in the city of São Paulo/SP.

The total chicken eggs crude were submitted to acid extraction in the presence of Acetic Acid 2 mol.L^{-1}, with constant

agitation for 60 min at 4° C. After the stirring time, the insoluble material was separated by centrifugation at 3113 x g

for 30 min and the supernatant obtained was injected into coupled Sep-Pack$_{18}$ cartridges (Waters Associates – 20 cc

vac cartridge, 5 g, and particle weight 55-105 μm) equilibrated in trifluoroacetic acid (0,1% TFA). The supernatant was

pre-fractionated in three different acetonitrile (ACN) concentrations (5%, 40%, and 80% of ACN), using vacuum

manifold system equipment (SPE 12 positions aho 6023 - Phenomenex) and then were lyophilized.

2.2. Reverse-Phase High-Performance Liquid Chromatography (RP-HPLC)

The fractions, 5%, 40% and 80% of ACN with 2 g each, were reconstituted in 5 mL 0.1% TFA and subjected by

reverse-phase high-performance liquid chromatography (RP-HPLC) was carried out at room temperature on a

Shimadzu LC-8A system. The column was a Shim-Pack Preparative ODS C-18 (20 mm x 250 mm; particle weight 10

μm; pore weight 300 Å). The elution was performed with a linear ACN gradient, equilibrated in 0.1% TFA for 60 min

at a flow rate of 8.0 mL/min (0 to 20%) for the fraction eluted in 5%, 2 to 60% for the fraction in 40%, and 20 to 80%

fraction eluted in 80%. The ultraviolet (UV) absorbance of the effluent was monitored at 225 nm. The fractions were

manually collected, were vacuum-dried and suspended in 1 mL in ultrapure water for antimicrobial activity assays.

The molecule described in this work, fraction 21, was subjected to a second step of fractionation using linear gradient

of ACN (39 to 49%) with 0,05% TFA at a flow rate of 2 mL/min for 60 min on the same column Shim-Pack Preparative

ODS C-18 (20 mm x 250 mm; particle weight 10 µm; pore weight 300 Å). The fractions were hand-collected, were

vacuum-dried, suspended in 500 µL in ultrapure water and it was quantified based on absorbance 205 nm using a

NanoDrop 2000 spectrophotometer (Thermo Fischer Scientific, Waltham, MA, USA).

2.3. Microorganisms

Fungal and bacterial strains were obtained and stored in the freezer at -80°C in the Laboratory for Applied Toxicology

(LETA), of the Butantan Institute (São Paulo, Brazil). *Staphylococcus aureus* ATCC 29213 (Gram-positive bacteria),

Escherichia coli D31 (Gram–negative bacteria); *Aspergilus niger* (Filamentous fungus – isolated from bread); and

Cryptococcus neoformans yeast strain A serotype (CFP registration 55 (WM 148 = VNI) were used. The *Cryptococcus*

neoformans strain was granted by Dra. Marcia Melhem from the Adolfo Lutz Institute, Brazil - São Paulo.

2.4. Antimicrobial Assays and Minimal Inhibitory Concentration (**MIC**)

Antimicrobial activities of the fractions were evaluated by a liquid growth inhibition assay (**Bulet et al., 1993**). A

suspension of the microorganisms collected in the mid-logarithmic growth phase was used. Bacteria strains were

cultured in poor broth nutrient medium (PB: 1.0 g peptone in 100 mL of water containing 86 mM NaCl at pH 7.4; 217

mOsM); for the fungi and yeast strains were cultured in potato dextrose broth (1/2-strength PDB) (1.2 g potato

dextrose in 100 mL of H_2O at pH 5.0; 79 mOsM). Determination of antimicrobial activity was performed using 5-fold

microtiter broth dilution assay in 96-well sterile plates at a final volume of 100 µL. A mid-log phase culture were

diluted to a final concentration of 5 x 10^4 CFU/mL for bacteria and 5 x 10^5 CFU/mL for fungus and yeast (**Bulet, 2008;**

Hetru and Bulet, 1997). Sterile water and PB or PDB were used as growth control, and Streptomycin antibiotic (10.0

mg/mL) was used as growth inhibition control. Microtiter plate were incubated for 18 and 24 h for bacteria and

fungi/yeast, respectively, at 30°C; growth inhibition was determined by measuring absorbance at 595 nm. The

minimum inhibitory concentration (MIC) are expressed as the interval of the concentration [a]–[b], where [a] is the

highest tested concentration at which the microorganism grow, and [b] is the lowest concentration which causes a

growth inhibition of 100% (**Bulet et al., 1993**). The assay was performed using a serial dilution and duplicate in

96-well sterile plate; at 20 µL stock solution was used in each microtiter plate well and added to 80 µL of the

microorganism dilution (**Silva et al., 2000; Lorenzini et al., 2003; Riciluca et al., 2012**).

2.5. Determination of Hemolytic Activity

The hemolytic activity of the fractions with antimicrobial activity was assessed against human erythrocytes from a

health adult donor. The Ethics Committee was the University of São Paulo School of Medicine (USP - FMUSP), n°

18794419.7.0000.0065. The erythrocytes were collected in 0.15 M citrate buffer, pH 7.4, and washed three times by

centrifugation (700 x g, 10 min, 4°C) with 0.15 M phosphate buffer saline (PBS – 35 mM phosphate buffer, 0.15 M NaCl,

pH 7.4), and ressuspended in PBS to a final 3% (v/v) concentration.

Aliquots of 50 µL of each fraction were added to 50 µL of 3% (v/v) suspension erythrocytes in the well of U – shaped

bottom plates and incubated for 1h at 37° C (with a final volume of 100 µL). The supernatant was first collected and

hemolysis was measuring the absorbance at 405 nm of each well in a Victor³ (1420 Multilabel Counter/ Victor³, Perkin

Elmer). The hemolysis percentage was expressed in relation to a 100% lysis control (erythrocytes incubated with 0.1%

triton X-100) (Sigma-Aldrich, St. Louis, MO, USA); PBS was used as negative control. The fractions was tested serial

dilution and duplicate, and calculation was made according to the fallowing equation: %hemolysis = (Abs sample –

Abs negative)/(Abs positive – Abs negative) (**Riciluca et al., 2012; Díaz-Roa et al., 2018; Segura and Silva**

Junior,2018).

2.6. Sodium Dodecyl Sulfate-Polyacrylamide Gel (SDS-PAGE)

Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-Page) was carried out. This experiment was

performed, with modifications, based on the protocol described by **Laemmli** (**1970**). The fraction Ovipin (455 µg) were

analyzed under non-reducing conditions (without the presence β-mercaptoetanol) and diluted in four times

concentrated buffer (250 mmol.L^{-1} Tris-HCl pH 6.8; 300 mmol.L^{-1} SDS; 1 mmol.L^{-1} bromophenol blue; 40% glycerol),

and heated at 90° C for about 2 minutes. The stacking gel and the running gel were prepared in the concentrations

4.5% and 12% of the polyacrylamide, respectively. The constant voltage and temperature utilized was 120 V at 4° C,

and the total running time was 3 h. As a molecular weight marker was used See Blue (Thermo Fisher Scientific Inc.).

And the proteins were stained with Coomassie Blue R-250 (**Laemmli,1970**).

2.7. Mass Spectrometry

The fractions that showed antimicrobial activity were analyzed by positive mode mass spectrometry LC-MS/MS on a

LTQ XL (Thermo Scientific). The fractions (10 μL of which) were concentrated and ressuspended in 10 μL Formic

Acid 0,1%. The liquid chromatography was perfomed in a C18 column (13 cm; 75 μm, i. d.x 100 mm – 3 μm from

Waters), using a 5 to 95%B in 25 minutes with solvents A: 15% in acidified water (0,1% TFA) and B: 95% in acidified

water (0.1% TFA) for 2 min; with flow of 300 nL/min, voltage of 2.5 kV at 200°C. The full scan was performed with

m/z of 100 a 2000, occurring MS/MS–CID, and parameters of five most intense ions, with isolation width of 2 m/z at

oCID (collision energy at 35).

2.8. Database Search

The resulting spectra in "*.RAW*", format were collected and processed using MSconvert software (**Chambers et al.,**

2012). Thus, they were converted to the mascot generic format ("*.MGF*") and thus analyzed into database searches

using the Mascot tool (**Perkins et al., 1999**). The MS/MS peak list files were submitted to an in – house version of the

Mascot server (Matrix Science, United States) and screened against the Uniprot database. PEAKS Studio Software

(version 8.5, Bioinformatics Solutions Inc., Waterloo, Ontario, Canada) *de novo* sequencing/database search used for

establishing sequence. Analysis involved 10 ppm error tolerance for precursor ions and 0.6 Da for fragment ions.

Oxidation was considered a variable modification.

The physicochemical parameters of which the fraction (such as the total number of positively and negatively charged

residues, molecular weight, and theoretical pI) were calculated using the ProtParam tool, available through the portal

ExPASy of the Swiss Institute of Bioinformatics website (https://web.expasy.org/protparam/) (**Gasteiger et al., 2005**).

3. Results

3.1. Chicken Eggs Fractionation and Antimicrobial Activity

After acid extraction of the crude extract from the chicken eggs *Gallus gallus domesticus*. The acid extract was applied

to Sep-Pak® C18 cartridges for pre-fractionation. The supernatant were eluted by three successive concentrations of

acetonitrile (ACN) (5%, 40% and 80%) in acidified water (0.1% TFA). All the material at 80% ACN was subjected and

analyzed by RP-HPLC (**Figure 1A**), and the fractions obtained was lyophilized, suspended in ultrapure water (1 mL)

and tested against *Staphylococcus aureus*, *Escherichia coli*, *Aspergillus niger*, and *Cryptococcus neoformans* in a liquid

growth inhibitory assay. The searched for the eluted 80% ACN showed the presence of six fractions (9, 11, 20, 21, 22

and 24) with antimicrobial activity (**Table 1**).

185   **Table 1.** Antimicrobial activity fractions from elution 80% ACN Chicken eggs.

Microorganisms				
	Fungus		Bacteria	
	Filamentous	Yeast	Gram-positive	Gram-negative
Fractions	*A. niger*	*C. neoformans*	*S. aureus*	*E. coli*
9	-	+	-	-
11	+	+	-	-
20	+	+	+	+
21	+	+	+	+
22	+	+	+	+
24	+	+	-	-

188   Symbol (+) fraction showed antimicrobial activity; Symbol (-) fraction do not showed antimicrobial activity.

FIGURE 1 | Reversed-phase high-performance liquid chromatograph (RP-HPLC) of the Chicken eggs *Gallus gallus domesticus* extract eluted with 80% acetonitrile (ACN). After the pre-fractionation in a Sep-Pak C18 column, the fraction was eluted with 80% ACN were subjected to a Shim-Pack ODS C18 (20 x 250 mm, 300 Å, 10 µm); Solvents System: Solvent A: 0.1% TFA/H₂O and Solvent B: 0.1%TFA/ACN with a linear gradient 20% to 80% ACN, 60 min at 8 mL/min flow rate. (**A**)The fraction 21 indicated by an asterisk (*) had antimicrobial activity against microorganisms tested. (**B**) Ovipin, indicated with an arrow, using same column and solvents system: Solvent A: 0.05% TFA/H₂O and Solvent B: 0.05%TFA/ACN. Ovipin was eluted from linear gradient 39% to 49% ACN, for 60 min at 2 mL/min flow rate. Both runs the absorbance was monitored at 225 nm.

3.2. Minimum Inhibitory Concentration (MIC) of Ovipin peptide

The antimicrobial activity occurred in the fraction 21 eluted with retention time range of 32.7-34.1 min (**Figure 1A**).

This fraction was submitted to a second step of fractionation and showed only one a pronounced antimicrobial

fraction: named Ovipin (**Figure 1B**). The name to choice for which fraction referring the chicken eggs species *Galus*

gallus domesticus. The concentration of Ovipin peptide was determined based on the Lambert-Beer Law using the

molar extinction coefficient at 205 nm absorption (**Anthis and Clore, 2013**), obtained using the Protein Parameter

Calculator tool available at the Nick Anthis website (http://nickanthis.com/tools/a205.html; accessed on 15 September

2021). The antimicroial activity was performed by liquid growth inibition in liquid medium. The native Ovipin

peptide was capable of inhibited the growth of *M. luteus* A270 1.94 µM (2.84 µg/mL), *A. niger* 31.01 µM (45.5 µg/mL),

C. neoformans VNI (WM 148) 22.75 µM (15.51 µg/mL). And the *E. coli* D31 was the least sensitive to this peptide (**Table**

2).

Table 2. Antimicrobial activity spectrum of Ovipin peptide.

Microorganisms	MIC (µM (µg/mL))[1]
Gram-positive bacteria	
Micrococcus luteus A270	1.94 (2.84)
Gram-negative bacteria	
Escherichia coli D31	ND
Fungi	
Aspergillus niger	31.01 (45.5)
Yeast	
Cryptococcus neoformans VNI (WM 148)	22.75 (15.51)

MIC, minimum inhibitory concentration (µg/mL): µM. ND, not detectable (antimicrobial activity not detected in the concentrations assayed). [1]The MIC refers to the concentration necessary to achieve a growth inhibition of 100%.

3.3. Hemolytic Activity

To determine the effect of the Ovipin peptide on human erythrocytes at the antimicrobial concentrations, its hemolytic activity was tested. After incubating red blood cells from a healthy donor with Ovipin (124.05 µM up to 0.24 µM) concentrations for 1 h at 37 °C. No hemolytic activity was observed, demonstrated that Ovipin does not cause significant hemolytic effect against of human erythrocytes within these concentrations (**Figure 2**) (**Yacoub et al., 2014**). And the electrophoretic profile (data not show) of this peptide (up to 455 µg) could not observe bands in this gel concentration (12% polyacrilamide).

FIGURE 2 | Hemolytic Activity of the Ovipin peptide incubated with human erythrocytes in different concentrations of the peptides. Serial dilution and duplicated was performed, for 1 h at 37°C. The hemolysis percentage was expressed

in relation to 0% hemolysis with PBS and 100% lysis control (0.1% triton X-100)

3.4. Mass Spectrometry and Characterization of Ovipin Peptide

After the eletrophoresis, where it was not possible observe bands in the gel concentration (12% polyacrilamide), the

Ovipin peptide was analyzed using the MASCOT® software to search the *Gallus gallus* database. This tool revealed a

sequence with 15 amino acid – YVSPVAIVKGLNIPL- and the molecular weight of 1,581.9494 Da with high identity

with a fragment from the Apolipoprotein B (Apo-B) of *Gallus gallus* (**Figure 3**). By the deconvolution of the peptide

and analysis of the data in Peaks software, the PEAKS *de novo* sequencing showed, as well as the Mascot tool, 15

amino acid sequence do not having post-translational modification (PTM) (**Figure 4**). Collision-induced dissociation

(CID) spectrum from mass/charge (m/z) of its double charged ion give $[M + 2H]^{2+}$, m/z 792.3484 (**Figure 5**). Both tools

confirmed the amino acid sequence of Ovipin peptide.

Figure 3: Result obtained in searches databases using the tool Mascot® (Matrix Science). The Ovipin peptide is
similar with a fragment from the *Gallus gallus* Apolipoprotein B (APO B)

Figure 4: Peptide spectrum match indicated by a blue line below the sequence of the Apolipoprotein B of *Gallus gallus*.
Ovipin covered 2% of the whole protein sequence.

Figure 5: Collision-induced dissociation (CID) spectrum of *de novo* sequencing antimicrobial peptide, Ovipin. The ions belonging to –y (red) and –b (blue) series indicated in the spectrum correspond to the amino acid sequence of the peptide: YVSPVAIVKGLNIPL. The fragments of the sequenced peptide are represented by standard amino acid code letters.

Knowing the sequence, some tools were used predicting Ovipin significant physicochemical characteristics. ExPASY's (SIB Bioinformatics Resource Portal) PepDraw and Pep-Calc.com sequence analysis yielded a potential peptide isoelectric point (pI), molar extinction coefficient and net (**Table 3**).

The peptide was predicted to have one Lys positively-charged aa residue and six of the 15 aa were predicted to be hydrophobic (2 Ile, 2 Leu and 3 Val), showed hydrophobic ratio of 53% and suggesting poor water solubility for Ovipin. ExPASy's ProtParam tool predicted that the peptide would intact for up 2.8 hours in mammalian reticulocytes (*in vitro*), 10 min in yeast (*in vivo*) and 2 min in *Escherichia coli* (*in vivo*). This was likely due to the presence of a Tyr (Y) residue at the N-terminus.

Table 3: Ovipin theoretical physicochemical properties.

Peptide Properties	
Sequence:	YVSPVAIVKGLNIPL
Lenght:	15
Mass:	1581.9494
Isoelectric point (pI):	9.49
Net charge at pH:	1
Hydrophobicity:	+ 7.11 Kcal*mol^{-1}
Estimated solubility:	Poor water solubility
Extinction coefficient:	1280 M^{-1} cm^{-1}

Physicochemical parameters were calculated using ExPASy PepDraw and Pep-Calc.com

A BLAST (basic local alignment search tool) was used to identify a multiple alignment analysis of the peptide sequence Ovipin using Clustal Omega program. The BLAST searches and the alignments showed 100% sequence similarity with the fragment Apolipoprotein B of *Gallus gallus*, and 93.3% of similarity with fragments Apolipoprotein B with others five species of Aves, being *Patagioenas fasciata monilis*, *Amazona aestiva*, *Tinamus guttatus*, *Aptenodytes forsteri* and *Egretta garzetta* (**Figure 6**).

Figure 6: Alignment of the amino acid sequence of Ovipin. Using the Clustal Omega alignment, the Ovipin sequence was aligned with fragments of the Apolipoproteins of the *Gallus gallus* and others Aves species. Regions that show identical amino acid among all species are identified in dark red or (*****), and minor modifications are show in light red (**:**). Clustal Omega (http://www.ebi.ac.uk/Tools/msa/clustalo/; accessed on 24 January 2020) modified manually.

Table 4. Known antimicrobial peptides having similarity with Ovipin, as identified in the Antimicrobial Peptide Database (APD2) (**Wang et al., 2009**).

Peptide Name	Sequence Alignment	Source Organism	APD Identifier	Percentage Similarity
Temporin-1SKa (17 aa)	F L P V I L P V + I G K L L N G I + L + + Y V + S P V A I V K G L N + I P L	Frog *Rana sakuraii*	AP01924	47,36
Temporin-1GY (13 aa)	+ V I P + + I V S G L L S S L L Y V S P V A I V K G + L N I P L	Frogs *Pelophylax nigromaculatus*	AP02223	43,75
Temporin-1TGc (17 aa)	F L P V I L P V + I G K L L S G I + L + + Y V + S P V A I V K G L + N I P L	*Rana sakuraii*	AP01268	42,1
Peptide AN5-1 (12 aa)	Y S K S L P + L S V + + L N + P + Y + V S + P V A I V K G L N I P L	*Paenibacillus alvei* AN5	AP02193	41,17

The Antimicrobial Peptide Database (APD) prediction tool was used to align Ovipin

4. Discussion

The abundance of multidrug-resistance of the microorganisms has emerged, increasing the spectrum of untreatable

infections and triggering a worldwide public health problem (**Gaglione et al., 2011; Blair et al., 2015**). The accelerated

growth of antimicrobial resistance has created a great interest in the discovery for new compounds more effective,

mainly with antimicrobial action, to be used as tools for antimicrobial therapies (**Diaz-Roa et al., 2018; Chernysh et al.,**

2015)

During the hatched time, eggs are constantly attacked by a variety of microorganisms that decrease their chances of

survival and viability. (**Packard & Packard, 1980**). The compounds isolated from eggs showed a broad spectrum of

antimicrobial activity, mainly against bacteria's (**D'Alba & Shawkey, 2015**). In the present work we isolated and

characterize a peptide, denoted Ovipin, from chicken eggs *Gallus gallus domesticus*, its antimicrobial activity,

biochemical and structural properties were evaluated.

The antimicrobial activity Ovipin (MIC = 22.75 µM (15.51 µg/mL)) against *C. neoformans* was very interesting, because

this yeast has a strong relationship with human infections with highly incidence in immunocompromised and

immunocompetent individuals. *Cryptococcus* virulence factors protect it from host immune responses. According to

the host's immune response, this microorganism can make changes in the metabolism itself as well as induce changes

in the structure of its capsule, decreasing its sensitivity to conventional antifungals. The production of melanin

protects the microorganism from oxidative stress, the urease enzyme is able to neutralize the acidic pH of

phagolysosomes, as well as being able to carry out non-lytic exocytosis. *Cryptococcus* fuses its phagosomal membranes

with plasma membranes to escape the phagocytosis process. Since it can survive inside the phagolysosome without

being destroyed, it can more easily cross the blood-brain barrier. And finally, cryptococci can manipulate the adaptive

immune system response and remain in a latent state of infection (**GUSHIKEN et al, 2021**). Thus, as Ovipin was able

to inhibit the growth of *Cryptococcus neoformans*, could this peptide be interfering in any way with the virulence factors

of this microorganism?

In the work **Candido-Ferreira *et al.*** (**2017**) was identified and characterized the Oligoventin peptide (8 amino acid

residues and MW 1,061.4 Da) from the eggs of the Brazilian armed spider *Phoneutria nigriventer* (Ctenidae,

Araneomorphae). Oligoventin showed antimicrobial activity (MIC range at 47 to 188.9 µM) against Gram-positive and

Gram- negative bacteria, as also yeast (*Candida albicans* MDM8). The molecule shows 4,3% of hemolysis against

human erythrocytes at 188.9 µM and 2% at 94.5 µM . Therefore, it is worth pointing out that both Ovipin and

Oligoventin, although islolated from eggs of different animals, have antimicrobial action against opportunistic

pathogens of great medical importance.

Ovipin presented antimicrobial activity (MIC = 1.94 μM (2.84 μg/mL)) against *Micrococcus luteus* (Gram-positive

bacteria). In the work of **Memarpoor-Yazdia and Chamania (2012**), was identified from hydrolyzed lysozyme, the F2

peptide (MW 1753.98 Da) that was active against *Leuconostoc mesenteroides* (Gram-positive bacteria) and *Escherichia coli*

(Gram-negative bacteria). **Mine et al. (2004**) also isolated and identified from hydrolyzed lysozyme, the peptide F1d

(MW 1185 Da), with antimicrobial activity against *Escherichia coli* K-12, and the peptide F3b (MW 875 Da) that was

active against *Staphylococcus aureus* 23-394 (Gram-positive bacteria).

Through searches carried out in online databases, Ovipin demonstrated a high percentage of similarity with frog

peptides. Temporin-1SKa (47%), Temporin-1GY (43%) and Temporin-1TGc (42%), peptides from *Rana sakuraii* and

Pelophylax nigromaculatus, respectively, both peptides with antimicrobial activity against gram-positive and

gram-negative bacteria (**SONG et al, 2013; SUZUKI et al, 2007**). Amphibians, like chicken eggs, are frequently

exposed to many pathogens, and studies have shown several isolated and identified antimicrobial peptides or

pharmacological peptides from frogs with diverse function, mainly against multi-resistant microorganisms

(**HAYASHIDA, SILVA JUNIOR, 2021; XIAO et al., 2011**)

Ovipin showed to have 1,581.9494 Da and similarity of 93.3 to 100% with Apoliproteins B of the five groups of Aves.

Very low-density lipoproteins (VLDL) are precursors low-density lipoproteins (LDL) of eggs yolk. VLDL II and apo B

are mainly proteins that constituted VLDL. The molecular weight of apo B is 500 kDa, showed only one structural unit

and 64% of the similarity with human apo B-100. When transferred into the yolk, the apo B undergoes enzymatic

cleavages, forming news fragments (**Jolivet *et al.,* 2006**), which may have new functions in the yolk.

The apolipoproteins are recognized as source bioactive-peptides showed broad antimicrobial or antiviral activity

(**Gaglione *et al.* (2017**). In the work **Zambrovicz et al.,** (**2015**) was identified four peptides with molecular weight

ranging from 1,210.62 to 1,677.88 Da obtained from Apoliproteins B, Vitellogenin-2 and Apovitelenin-1, all proteins

localized in chicken egg yolk. In relation the peptides identified from Apolipoprotein B, one peptide showed amino

acid sequence YINQMPQKSRE (MW: 1,393.774 Da) and the other peptide showed amino acid sequence

YINQMPQKSREA (MW: 1,464.813). In this study not was evaluated the antimicrobial activity this peptides, but was

demonstrated the antioxidant activity, ACE inhibitory and/or antidiabetic activities. The peptide YINQMPQKSRE

demonstrated the strongest antioxidant, and the DPPH free radical scavenging and iron chelating activities reached

2.3 μM Trolox eq/mg and 37.4 μg Fe^{2+}/mg, respectively. The peptide showed YINQMPQKSREA the highest ferric

reducing activity (76.0 μg Fe^{2+}/mg), an activity very strong ACE (IC_{50} = 10.1 μg/mL) and DPP-IV (IC_{50} = 222.8 μg/mL)

inhibitory activity.

5. Conclusions

Whereas antimicrobial resistance has become a serious concern and a public health issue because it limits treatment

options, the identification and characterization of new antibiotics agents can lead to the development of new

compounds that act against resistant pathogenic microorganisms. In this work we reported one peptide that was

extracted from chicken eggs *Gallus gallus domesticus*, purified by RP-HPLC and characterized by amino acid

sequencing. The Ovipin, a small antimicrobial peptide that is a fragment of Apolipoprotein B of *Gallus gallus*; it

contains 15 amino acid sequence and is active against Gram-positive (*Micrococcus luteus*), Filamentous fungi

(*Aspergillus niger*) and Yeast (*Cryptococcus neoformans*), and which in the future as possible multifunctional candidates.

The results of this work show that chicken eggs, over time, have become increasingly important not only as a food to

meet nutritional needs, but also the chicken egg has become a great source for the discovery of new peptides, as the

literature has showed, and as has been effective against microorganisms that developed resistance to conventional

antibiotics. In searches carried out in the literature, it was not yet possible to find fragments of Apolipoprotein B,

whether from birds or humans, that have antimicrobial action, mainly against fungi. Perhaps the Ovipin peptide

could be the first fragment of Apolipoprotein B, was also able to inhibit the growth of *C. neoformans*, an important

yeast for medicine because act mainly imunossupressed patient. Further studies aimed at evaluating the activity of the

Ovipin peptide against others multidrug-resistance bacteria, fungi, viruses, and parasites are needed, as well as

evaluating its mechanism of action against *C. neoformans*, to understand in which possible virulence factor of this yeast

the Ovipin peptide may be acting and that inhibit its growth, and it can be used as a new peptide in the biotechnology

or pharmaceutical industry.

Author Contributions: Conceptualization, SRS, AM and PISJ; Data curation, SRS; Formal analysis, AM and PISJ; Funding

acquisition, AM and PISJ; Investigation, SRS; Methodology, AM and PISJ.; Project administration, AM and PISJ; Resources, PISJ;

Supervision, PISJ and AM; Validation, SRS, AM and PISJ; Writing-original draft, SRS; Writing-review and editing, SRS, AM and

PISJ.

Funding: This work was supported by the Research Support Foundation of the State of São Paulo (FAPESP/CeTICS) [Grant No. 2013/07467-1]; the Brazilian National Council for Scientific and Technological Development (CNPq) [Grant No. 472744/2012-7]; and the Coordination for the Improvement of Higher Education Personne (CAPES) [Grant No. 1646478].

Ethical Approval: This research was approved and performed in accordance with the Ethical Principles in Animal Research adopted by the Ethics Committee for the Use of Animals of UNIFESP - CEUA No 9183300118 and Plataforma Brasil CAAE N° 18794419.7.0000.0065 and was approved.

Data Availability Statement: The data that support the findings of this study are presented in the main manuscript or in the supplementary material of this article.

Acknowledgments: We would like to thank Caroline Correa, Marta Gomes, Rogerio Lauria and Sinval Gregorio laboratory technicians from the Biophysics Department at Unifesp for all their help with peptide synthesis. Professor Dr. Edgar Julian Paredes Gamero, and his students Heron and Wagner at Unifesp, for the opportunity to learn about the development of the antitumor study. *Cryptococcus neoformans* VNI (WM148) was kindly provided by Dra Márcia Melhem - Instituto Adolfo Lutz - São Paulo/SP, Brasil.

Conflicts of Interest: "The authors declare no conflict of interest".

References

Almeida, F., Wolf, J.M., da Silva, T.A. *et al.* Galectin-3 impacts *Cryptococcus neoformans* infection through direct antifungal effects. *Nat Commun* **8,** 1968 (2017). https://doi.org/10.1038/s41467-017-02126-7

Andreu, D. e Rivas, L. Animal antimicrobial peptides: an overview. *Biopolymers*, 47 (6) 415-433 (**1998**).

Anthis, N.J.; Clore, G.M. Sequence-specific determination of protein and peptide concentrations by absorbance at 205 nm. Protein Sci. **2013**, 22, 851–858.

Antimicrobial-peptide-database. Available online: http://aps.unmc.edu/AP/prediction/prediction_main.php (accessed on 12 September 2020).

Aspaugh, J.A. Virulence mechanisms and Cryptococcus neoformans pathogenesis. Fungal Genet Biol. 2015 May ; 78: 55–58. doi:10.1016/j.fgb.2014.09.004.

Bahar, A.A.; Ren, D. Antimicrobial peptides. *Pharmaceuticals (Basel)*, 6 (12) 1543-1575 (**2013**).

Ben-Ami, R., Kontoyiannis. Resistance to Antifungal Drugs. Infect Dis Clin N Am 35 (2021) 279–311 https://doi.org/10.1016/j.idc.2021.03.003

Brogden, K.A. Antimicrobial peptides: pore formers or metabolic inhibitors in bacteria? *Nat Rev Microbiol*, 3 (3) 238-250 (**2005**).

Bulet, P. Strategies for the discovery, isolation, and characterization of natural bioactive peptides from the immune system of invertebrates. Methods Mol. Biol. **2008**, 494, 9–29. [CrossRef] [PubMed]

Bulet, P., Dimarcq, J.L., Hetru, C., Lagueux, M., Charlet, M., Hegy, G., Van Dorsselaer, A. e Hoffmann, J.A. A novel inducible antibacterial peptide of Drosophila carries an O-glycosylated substitution. *J Biol Chem*, 268 (20) 14893-14897 (**1993**).

Candido-Ferreira, I.L.; Kronenberger, T.; Sayegh, R.S.R.; Batista, I.F.C.; Silva Jumior, P.I. Evidence of an antimicrobial peptide signature encrypted in HECT E3 uniquitin ligases. Front Immunol 2017 Jan 9;7:664. doi: 10.3389/fimmu.2016.00664. eCollection 2016.

Chambers, M.C., Maclean, B., Burke, R., Amodei, D., Ruderman, D.L., Neumann, S., Gatto, L., Fischer, B., Pratt, B., Egertson, J., Hoff, K., Kessner, D., Tasman, N., Shulman, N., Frewen, B., Baker, T.A., Brusniak, M.Y., Paulse, C., Creasy, D., Flashner, L., Kani, K., Moulding, C., Seymour, S.L., Nuwaysir, L.M., Lefebvre, B., Kuhlmann, F., Roark, J., Rainer, P., Detlev, S., Hemenway, T., Huhmer, A., Langridge, J., Connolly, B., Chadick, T., Holly, K., Eckels, J., Deutsch, E.W., Moritz, R.L., Katz, J.E., Agus, D.B., MacCoss, M., Tabb, D.L. e Mallick, P. A cross-platform toolkit for mass spectrometry and proteomics. *Nat Biotechnol*, 30 (10) 918-920 (**2012**).

Cruz, J., Ortiz, C., Guzman, F., Fernandez-Lafuente, R. e Torres, R. Antimicrobial peptides: promising compounds against pathogenic microorganisms. *Curr Med Chem*, 21 (20) 2299-2321 (**2014**).

Diaz-Roa, A., Patarroyo, M.A., Bello, F.J. e Da Silva, P.I., Jr. Sarconesin: Sarconesiopsis magellanica Blowfly Larval Excretions and Secretions With Antibacterial Properties. *Front Microbiol*, 9 2249 (**2018**).

Ercan, D. e Demirci, A. Recent advances for the production and recovery methods of lysozyme. *Crit Rev Biotechnol*, 36 (6) 1078-1088 (**2016**).

Frare e Silva, R. Fungal infections in immunocompromised patients. J Bras Pneumol. 2010;36(1):142-147

Gaglione R, Dell'Olmo E, Bosso A, Chino M, Pane K, Ascione F, Itri F, Caserta S, Amoresano A, Lombardi A, Haagsman HP, Piccoli R, Pizzo E, Veldhuizen EJA, Notomista E, Arciello A. Novel human bioactive peptides identified in Apolipoprotein B: Evaluation of their therapeutic potential. Biochem Pharmacol. 2017 Apr 15;130:34-50. doi: 10.1016/j.bcp.2017.01.009. Epub 2017 Jan 25. PMID: 28131846.

Garcia, F.; Villegas, E.; Espino-Solis, G.P.; Rodriguez, A.; Paniagua-Solis, J.F.; Sandoval-Lopez, G.; Possani, L.D.; Corzo, G. Antimicrobial peptides from arachnid venoms and their microbicidal activity in the presence of commercial antibiotics. J. Antibiot. **2013**, 66, 3–10. [CrossRef] [PubMed]

447 Gasteiger, E.; Hoogland, C.; Gattiker, A.; Duvaud, S.; Wilkins, M.R.; Appel, R.D.; Bairoch, A. Protein Identification and
448 Analysis Tools on the ExPASy Server. In The Proteomics Protocols Handbook; John, M.W., Ed.; Humans Press Inc.:
449 Totowa, NJ, USA, 2005; pp. 571–607.
450
451 Gil-Chávez, G.J.; Villa, J.A.; Ayala-Zavala, F.; Heredia, J.B.; Sepulveda, D.; Yahia, E.M.; González-Aguilar, G.A.
452 Technologies for Extraction and Production of Bioactive Compounds to be Used as Nutraceuticals and Food
453 Ingredients: An Overview. Comprehensive Reviews in Food Science and Food Safety, January (**2013**) Vol.12, Issue1,
454 Pages 5-23.
455 https://doi.org/10.1111/1541-4337.12005
456
457 Guimarães, D.O.; Momesso, L. S.; Pupo, M.T. Antibiotics: therapeutic importance and perspectives for the discovery
458 and   development   of   new   agents   Quim.   Nov.   Vol.   33,   N°   3,   São   Paulo,   (**2010**).
459 http://dx.doi.org/10.1590/S0100-40422010000300035
460
461 Gushiken, A.C.; Saharia, K.K.; Baddley, J.W. Cryptococcosis. Infect Dis Clin N Am 35 (2021) 493–514
462 https://doi.org/10.1016/j.idc.2021.03.012
463
464 Hancock, R.E. e Sahl, H.G. Antimicrobial and host-defense peptides as new anti-infective therapeutic strategies. Nat
465 Biotechnol, 24 (12) 1551-1557 (2006).
466
467 Hayashida, P.Y,; Silva Junior, P.I. Insights into Antimicrobial Peptides from *Limacus flavus* Mucus. Current
468 Microbiology (2021) 78:2970–2979 https://doi.org/10.1007/s00284-021-02552-3
469
470 Henao-Martínez, A.F.; Chastain, D.B.; Franco-Paredes, C. Treatment of cryptococcosis in non-HIV
471 immunocompromised patients. Curr Opin Infect Dis. 2018 Aug; 31(4):278-285. doi: 10.1097/QCO.0000000000000458.
472
473 Hetru, C.; Bulet, P. Strategies for the isolation and characterization of antimicrobial peptides of invertebrates. Methods
474 Mol. Biol. **1997**, 78, 35–49. [CrossRef]
475
476 Ibrahim, H.R., Sugimoto, Y. e Aoki, T. Ovotransferrin antimicrobial peptide (OTAP-92) kills bacteria through a
477 membrane damage mechanism. Biochim Biophys Acta, 1523 (2-3) 196-205 (2000).
478
479 Izadpanah A, Gallo RL. Antimicrobial   peptides.   J   Am   Acad   Dermatol.   2005;52(3   Pt   1):381-392.
480 doi:10.1016/j.jaad.2004.08.026
481
482 Jolivet, P.; Boulard, C.; Beaumal, V.; Chardot, T.; Anton, M. Protein components of low-density lipoproteins purified
483 from hen egg yolk. J. Agric. Food Chem., 2006, 54, 4424-4429.
484
485 Kang, S.J., Park, S.J., Mishig-Ochir, T. e Lee, B.J. Antimicrobial peptides: therapeutic potentials. Expert Rev Anti Infect
486 Ther, 12 (12) 1477-1486 (2014).
487

488   Kolaczkowska, A., Kolaczkowski, M., Sokolowska, A., Miecznikowska, H., Kubiak, A., Rolka, K. e Polanowski, A. The
489   antifungal properties of chicken egg cystatin against Candida yeast isolates showing different levels of azole resistance.
490   *Mycoses*, 53 (4) 314-320 (**2010**).
491
492   Kovacs-Nolan, J., Phillips, M. e Mine, Y. Advances in the value of eggs and egg components for human health. *J Agric*
493   *Food Chem*, 53 (22) 8421-8431 (**2005**).
494   Laemmli, U.K. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature (**1970**), 227,
495   680–685. [CrossRef] [PubMed]
496
497   Li J, Koh JJ, Liu S, Lakshminarayanan R, Verma CS, Beuerman RW. Membrane Active Antimicrobial Peptides:
498   Translating Mechanistic Insights to Design. *Front Neurosci*. 2017;11:73. Published 2017 Feb 14.
499   doi:10.3389/fnins.2017.00073
500
501   Lorenzini, D. M., Da Silva, P. I. Jr., Fogaca, A. C., Bulet, P., and Daffre, S. (2003). Acanthoscurrin: a novel glycine-rich
502   antimicrobial peptide constitutively expressed in the hemocytes of the spider Acanthoscurria gomesiana. Dev. Comp.
503   Immunol. 27, 781–791. doi: 10.1016/S0145-305X(03)00058-2
504
505   Memarpoor-Yazdia, M.A., A.; Chamania, J. A novel antioxidant and antimicrobial peptide from hen egg white lysozyme
506   hydrolysates. *Journal of Functional Foods*, 4 (1) 278-286 (**2012**).
507
508   Mine, Y., Ma, F. e Lauriau, S. Antimicrobial peptides released by enzymatic hydrolysis of hen egg white lysozyme. *J*
509   *Agric Food Chem*, 52 (5) 1088-1094 (**2004**).
510
511   Mookherjee, N.; Hancock, R.E.W. Cationic host defence peptides: Innate immune regulatory peptides as a novel
512   approach for treating infections. Cell. Mol. Life Sci. 64 (2007) 922 – 933. DOI 10.1007/s00018-007-6475-6
513
514   Perkins, D.N., Pappin, D.J., Creasy, D.M. e Cottrell, J.S. Probability-based protein identification by searching sequence
515   databases using mass spectrometry data. *Electrophoresis*, 20 (18) 3551-3567 (**1999**).
516
517   Peters, B.M.; Shirtli_, M.E.; Jabra-Rizk, M.A. Antimicrobial Peptides: Primeval molecules or future drogs? PLoS Pathog.
518   **2010**, 6, e1001067. [CrossRef] [PubMed]
519
520   Riciluca, K.C., Sayegh, R.S., Melo, R.L. e Silva, P.I., Jr. Rondonin an antifungal peptide from spider (Acanthoscurria
521   rondoniae) haemolymph. *Results Immunol*, 2 66-71 (**2012**).
522
523   Rutherfurd-Markwick KJ. Food proteins as a source of bioactive peptides with diverse functions. *Br J Nutr*. 2012;108
524   Suppl 2:S149-S157. doi:10.1017/S000711451200253X
525
526   Shai, Y. Mechanism of the binding, insertion and destabilization of phospholipid bilayer membranes by alpha-helical
527   antimicrobial and cell non-selective membrane-lytic peptides. *Biochim Biophys Acta*, 1462 (1-2) 55-70 (**1999**).
528

529 Segura-Ramírez PJ, Silva Júnior PI. *Loxosceles gaucho* Spider Venom: An Untapped Source of Antimicrobial
530 Agents. *Toxins (Basel)*. 2018;10(12):522. Published 2018 Dec 6. doi:10.3390/toxins10120522
531
532 Silva, P.I., Jr., Daffre, S. e Bulet, P. Isolation and characterization of gomesin, an 18-residue cysteine-rich defense
533 peptide from the spider Acanthoscurria gomesiana hemocytes with sequence similarities to horseshoe crab
534 antimicrobial peptides of the tachyplesin family. *J Biol Chem*, 275 (43) 33464-33470 (**2000**).
535
536 Song, Y.; Ji, S,; Liu, W.; Yu, X.; Meng, Q.; Lai, R. Different Expression Profiles of Bioactive Peptides in *Pelophylax*
537 *nigromaculatus* from Distinct Regions. Biosci. Biotechnol. Biochem., 77 (5), 1075–1079, 2013.
538 https://doi.org/10.1271/bbb.130044
539
540 Srikanta, D.; Santiago-Tirado, F.H.; Doering, T.L. *Cryptococcus neoformans*: Historical curiosity to modern pathogen.
541 *Yeast*. 2014 February ; 31(2): 47–60. doi:10.1002/yea.2997.
542
543 Suzuki, H.; Iwamuro, S.; Ohnuma, A.; Coquet, L.; Leprince, J.; Jouenne, T.; Vaudry, H.; Taylor, C.K.; Abel, P.W.;
544 Conlon, J.M. Expression of genes encoding antimicrobial and bradykinin-related peptides in skin of the stream brown
545 frog Rana sakuraii. Peptide 28 (2007) 505-514. doi:10.1016/j.peptides.2006.10.016
546
547 Wang, G.; Li, X.; Wang, Z. APD2: The updated antimicrobial peptide database and its application in peptide design.
548 Nucleic Acids Res. **2009**, 37, 933–937.
549
550 Xiao, Y.; Liu, C.; Lai, R. Antimicrobial peptides from amphibians. BioMol Concepts, Vol. 2 (2011), pp. 27–38. doi
551 10.1515/BMC.2011.006
552
553 Zambrowicz, A.; pokora, M.; Setner, B.; Dabrowska, A.; Szoltysik, M.; Babij, K.; Szewczuk, Z.; Trziszka, T.; Lubec, G.;
554 Chrzanowska, J. Multifunctional peptides derived from an egg yolk protein hydrolysate: isolation and characterization.
555 *Amino Acids* **47,** 369–380 (2015). https://doi.org/10.1007/s00726-014-1869-x
556
557 Zasloff, M. Antimicrobial peptides of multicellular organisms. *Nature*, 415 (6870) 389-395 (**2002**).