1   **Enhanced nociceptor sprouting in a rabbit model of cerebral palsy**

2   **Authors**
3   E. J. REEDICH[1,2], L. T. GENRY[1,2,3], M. A. SINGER[4], C. F. CAVARSAN[1,2], E. MENA AVILA[1,2], D. M.
4   BOUDREAU[1,2], M. C. BRENNAN[1,2], A. M. GARRETT [1,2,3], L. DOWALIBY[1,2], M. R. DETLOFF[4*], K. A. QUINLAN[1,2*]
5
1 George and Anne Ryan Institute for Neuroscience, University of Rhode Island, Kingston, RI, USA
2 Department of Biomedical and Pharmaceutical Sciences, College of Pharmacy, University of Rhode
Island, Kingston, RI, USA
3 Interdisciplinary Neuroscience Program, University of Rhode Island, Kingston, RI, USA
4 Department of Neurobiology and Anatomy, Marion Murray Spinal Cord Research Center, College of
Medicine, Drexel University, Philadelphia, PA, USA
12
*Corresponding Authors:

Katharina A. Quinlan
George and Anne Ryan Institute for Neuroscience
Department of Biomedical and Pharmaceutical Sciences
College of Pharmacy
University of Rhode Island
130 Flagg Rd
Kingston, RI 02881
kaquinlan@uri.edu

Megan R. Detloff
Department of Neurobiology & Anatomy
Marion Murray Spinal Cord Research Center
College of Medicine
Drexel University
2900 W. Queen Lane
Philadelphia, PA 19129
mrd64@drexel.edu
30

Abstract

Spastic cerebral palsy (CP) is a movement disorder marked by hypertonia and hyperreflexia, and
the most prevalent comorbidity is pain. Since spinal nociceptive afferents contribute to both the
sensation of painful stimuli as well as reflex circuits involved in movement, we investigated the
relationship between prenatal hypoxia-ischemia (HI) injury which can cause CP, and possible changes in
spinal nociceptive circuitry. To do this, we examined nociceptive afferents and mechanical and thermal
sensitivity of New Zealand White rabbit kits after prenatal HI or a sham surgical procedure. As described
previously, a range of motor deficits similar to spastic CP was observed in kits born naturally after HI (40
minutes at ~70-80% gestation). We found that HI caused an expansion of peptidergic afferents (marked
by expression of calcitonin gene-related peptide; CGRP) in both the superficial and deep dorsal horn at
postnatal day (P)5. Non-peptidergic nociceptive afferent arborization (labeled by isolectin B4; IB4) was
unaltered in HI kits but overlap of the two populations (peptidergic and non-peptidergic nociceptors)
was increased by HI. Density of glial fibrillary acidic protein (GFAP) was unchanged within spinal white
matter regions important in nociceptive transmission at P5. We found that mechanical and thermal
allodynia was present in HI kits even in the absence of motor deficits. These findings suggest that
prenatal HI injury impacts spinal sensory pathways in addition to the more well-established disruptions
to descending motor circuits. In conclusion, changes to spinal nociceptive circuitry could disrupt spinal
reflexes and contribute to pain experienced by individuals with CP.

Key words: cerebral palsy, hypoxia-ischemia, sensation, nociception, rabbit, primary afferent

Abbreviations: Serotonin (5HT); calcitonin gene-related peptide (CGRP); central nervous system (CNS);
cerebral palsy (CP); Diamidino-2-Phenylindole (DAPI); dorsal column (DC); dorsal root ganglion (DRG);
glial fibrillary acidic protein (GFAP); Gross Motor Function Classification System (GMFCS); hypoxia-
ischemia (HI); isolectin B4 (IB4); immunohistochemistry (IHC); lateral spinothalamic tract (LST); normal
horse serum (NHS); nonsteroidal anti-inflammatory drugs (NSAIDs); postnatal day (P); phosphate
buffered saline (PBS); paraformaldehyde (PFA); regions of interest (ROIs); superficial dorsal gray matter
(SDGM); ventral spinothalamic tract (VST)

Introduction:

Cerebral palsy (CP) is the most common motor disability in children, affecting 1:500 live births (Oskoui et al., 2013). Chronic pain is the most prevalent co-morbidity of CP (McDowell et al., 2017; Novak et al., 2012): up to 77% of children with CP experience pain (Mckinnon et al., 2018; McKinnon et al., 2020). Individuals with CP experience chronic pain at significantly higher rates than the rest of the population (Badia et al., 2014; Doralp and Bartlett, 2010; García Jalón et al., 2021; Westbom et al., 2017). Pain decreases quality of life (Findlay et al., 2016), yet the pathophysiology of pain is rarely researched in the context of CP. This is likely due to the pervasive view that pain in individuals with CP is a symptom arising from secondary effects of physical stressors, including spasticity, hip subluxation, and joint contractures. Individuals with severe CP (as measured by Gross Motor Function Classification System (GMFCS) V) do experience more pain (García Jalón et al., 2021; Mckinnon et al., 2018); however, CP-affected individuals across all GMFCS levels report pain (Flanigan et al., 2020; Sienko, 2018). Even individuals born very prematurely (<26 weeks gestion), who do not develop CP, report recurrent pain and altered thermal sensitivity as adults (Walker et al., 2018). Thus, we hypothesized that the same trauma that can impact developing corticospinal tracts in CP could impact development of nociceptive circuitry in the spinal cord.

There is a wealth of evidence regarding the interrelatedness of sensory and motor performance (Cooper et al., 1995; Gupta et al., 2017; Matusz et al., 2018; Zarkou et al., 2020). Somatosensory deficits are present in CP, and their presence could contribute to motor dysfunction. Typically, primary afferent input onto spinal circuitry is a key mediator of functional motor performance. Proprioceptive and tactile afferent feedback informs the central nervous system (CNS) of motor command outcomes and sculpts ongoing drive to motoneurons. In particular, the role of primary nociceptive afferent fibers in motor function is critically overlooked—especially after CNS damage. Importantly, there are a few exceptions, including the restriction of motor unit activation in response to excessive stretch, detection of dangerous pH and other metabolite levels during muscle fatigue, and the exercise pressor reflex and motor unit modulation during exercise-induced fatigue (Laurin et al., 2015; Zając et al., 2015). Cutaneous nociceptors (mechanosensitive C, thermosensitive C and Aδ neurons) are small, pseudounipolar neurons in the dorsal root ganglion (DRG) that are analogous to Type III (Aδ) and Type IV (C) primary afferent fibers from the muscle, though their axons terminate in different dorsal horn laminae (Kandel et al., 2013; Lallemend and Ernfors, 2012; Laurin et al., 2015). Importantly, nociceptive and pain information can be relayed from muscle and joints via the nociceptive primary afferent reflex arcs, modulating motor output by generating a withdrawal reflex (Kandel et al., 2013; Kimura et al., 2004). The interconnection between motor and sensory systems can make it challenging to parse out their relative contribution to functional impairments.

As in other CNS disorders, there is a strong correlation between the integrity of the somatosensory system (including primary afferent input) and motor performance in CP (Gupta et al., 2017; Zarkou et al., 2020). Tractography and physiology experiments in individuals with unilateral CP revealed that integrity of somatosensory connections has a relatively larger impact on hand function compared to the integrity of motor connections (Gupta et al., 2017), and stimulation of peripheral nerves, which provides additional sensory feedback, improved motor performance in individuals with diplegic CP (Bumin and Kayihan, 2001; Kaelin-Lang, 2008).

Physical insults during pregnancy have been linked to later development of autism, schizophrenia, and the motor impairments that are hallmarks of CP (Marriott et al., 2017), but it is not

known if these prenatal events impact nociceptive afferents. Neural development can be negatively
impacted by relatively common conditions in the perinatal period, including maternal infection and
inflammation, placental insufficiency, and difficult birth. These conditions can all result in varying
degrees of hypoxia-ischemia (HI) and injury to the developing CNS. Effects of prenatal HI include white
matter injury, cortical damage and motor deficits (Buser et al., 2010; Derrick et al., 2004; Drobyshevsky
and Quinlan, 2017; Rocha-Ferreira and Hristova, 2016), but the effect on nociceptors is completely
unstudied. Notably, cell death in the deep dorsal horn (where synaptic targets of nociceptors are
located) is present after HI in rabbits and in cases of infant death after hypoxia (Clancy et al., 1989;
Drobyshevsky and Quinlan, 2017; Sladky and Rorke, 1986). Dorsal root ganglia (DRG) neurons undergo
perinatal growth and refinement of projections during the period of susceptibility to injuries causing CP
(Marmigère and Ernfors, 2007; Olson and Luo, 2018), though afferent plasticity is retained in adulthood
(Jiang et al., 2016; Tan et al., 2012). Changes to nociceptive DRG neurons are entirely unexplored in the
context of CP.

In this study, we have chosen to focus on nociceptors at the spinal level because they are the
neuronal link between sensation of stimuli in the periphery (that is perceived as painful by higher brain
centers) and hyperreflexia that is the hallmark of spastic CP. Nociceptors impact movement by
modulating spinal reflex circuits via primary afferent depolarization (PAD)(Black, 2019; Lucas-Osma et
al., 2019). PAD profoundly alters reflex responses (Hari et al., 2021; Hochman et al., 2010; Rudomin and
Schmidt, 1999) and could directly contribute to the hyperreflexia and reflex irradiation that is observed
in people with CP (Achache et al., 2010; Leonard et al., 1991; Leonard and Hirschfeld, 1995).
Importantly, monosynaptic stretch reflexes can be potently suppressed via nociceptor inhibition (Black,
2019; Lucas-Osma et al., 2019). Thus, not only could expanded distribution of nociceptive afferents
contribute to the perception of pain by higher brain centers, but also to hyperreflexia and disruptions in
motor coordination that are the hallmarks of CP.

Thus, we examined nociceptors and altered nociception at the level of the spinal cord in rabbit
kits that were subjected to HI injury *in utero* (40 minutes at ~70-80% gestation) using both
immunohistochemistry to quantify molecular markers for nociceptors and *in vivo* behavioral testing at
an age roughly corresponding to human toddlers (postnatal day [P] 5). We found that rabbit kits
exposed to prenatal HI demonstrated increased nociceptive behavior in response to cutaneous
mechanical and thermal stimuli. This increased nociceptive response corresponded to aberrant
distribution and density of nociceptive primary afferent fibers but not astrogliosis at this age.
Interestingly, our data show that enhanced forepaw and hindpaw sensitivity occurs independent of
motor deficits produced by this model. Together, these results indicate that sensory circuits are
vulnerable to perturbation by perinatal HI injury, and suggest that neuropathic pain could accompany
motor deficits characteristic of CP.

Materials and Methods:

Animal model: Experiments were performed in accordance with the ARRIVE guidelines and the United
States National Institutes of Health's Guide for Care and Use of Laboratory Animals. Approval of the
University of Rhode Island's Institutional Animal Care and Use Committee was obtained for all
experiments performed in this study. 62 kits of both sexes from 11 different litters originating from 6
dams were used for this study; some dams provided more than one litter of rabbits. The inherent

variability in behavioral data warranted large sample sizes. All efforts were made to minimize animal
suffering and to reduce the number of animals used.

Hypoxia-ischemia surgery: Pregnant New Zealand White rabbits (bred in house or ordered from Charles
River Laboratories, Inc., Wilmington MA), underwent HI procedures as previously described (Derrick et
al., 2004). Briefly, at ~70-80% gestation (day 22 -26 of gestation), dams were premedicated with
ketamine and xylazine (25 mg/kg and 3.75 mg/kg, respectively), anesthetized with isoflurane (1-3%;
inhaled via V gel tube) and occasionally treated with buprenorphine (0.01 - 0.24 mg/kg, as needed). The
left or right femoral artery was isolated. A Fogarty balloon catheter (2-4 French) was inserted and
advanced to the level of the descending aorta, above the uterine arteries, and inflated for 40 minutes to
cause HI in the fetuses. Dams subjected to sham surgical procedures underwent the same anesthesia
protocols as the HI dams but without insertion and/or inflation of the catheter.

After the procedures, dams recovered and later gave birth to kits. Neonatal kits of both sexes were
characterized for motor deficits, nociceptive behavior, and used for anatomical investigations as
described below. Since we examined behavior in neonatal animals far from sexual maturity (NZW
rabbits typically reach sexual maturity at ~6 months of age), and whose external genitalia were not yet
visible, we grouped together male and female kits.

Behavior: There is a wide range in motor phenotypes after HI, with some kits showing motor deficits
while others appear unaffected. Categorization of motor deficits was performed using a modified
Ashworth scale, observations/tests for activity, locomotion, posture, righting reflex, and muscle tone as
previously described (Derrick et al., 2004). Based on these tests, kits were labeled as "motor-affected
HI" or "motor-unaffected HI."

von Frey test for mechanical nociception: Neonatal rabbit kits of both sexes were tested for mechanical
allodynia using von Frey filaments. Animals were allowed to acclimatize in a chamber on an elevated
gridded (8 mm x 8 mm) surface. von Frey filaments (Touch Test® Sensory Evaluators; range: 1-26 g) were
applied in ascending order to the plantar surface of the forepaws and hindpaws of the kit; if paw
withdrawal occurred in response to application of the 1 g von Frey filament, filaments (0.6-0.008 g) were
applied in descending order until no response was observed. Each filament was tested at least five
consecutive times per paw to ensure the observed reaction was in response to the filament and not
general movement of the animal. The force (in g) of the filament that elicited a consistent withdrawal
response was recorded as the paw withdrawal threshold. If no response was elicited by application of
the 26 g von Frey filament, that paw was recorded as non-responsive and given a score of 60 g (which is
the subsequent filament). Filament forces eliciting withdrawal of left and right forepaws and hindpaws
were averaged for statistical analyses.

Hargreaves' test for thermal nociception: Thermal sensitivity was tested using a Hargreaves apparatus.
Specifically, a model 336G Plantar/Tail Stimulator Analgesia Meter (IITC Life Sciences, Woodland Hills,
CA) was used to apply a beam of radiant heat (4x6 mm area; 80% intensity; 8 s max duration) to the
plantar surface of each hindpaw and forepaw. Rabbit kits were allowed to acclimatize in a chamber on a
glass surface, and then each paw was tested three times with at least 30 seconds between applications.
The beam of radiant heat increased temperature gradually from rest (20-30 °C) to a maximum of 83 °C
at 8 s. Latency to paw withdrawal was recorded; if the animal did not withdraw its paw, that trial was
recorded as a non-response and given a latency of 9 s. Withdrawal latencies of left and right forepaws
and hindpaws were averaged for statistical analyses.

Immunohistochemistry (IHC): Kits were deeply anesthetized by intraperitoneal injection of ketamine/xylazine (100 mg/kg and 20 mg/kg, respectively) or sodium pentobarbital before tissue harvest at P5. Transcardial perfusion was performed using chilled 0.01 M phosphate buffered saline (PBS; pH 7.4) until tissue blanching was observed, then 4% paraformaldehyde (PFA) (in PBS) was infused at a constant flow rate of approximately 5-8 mL/min (depending on rabbit kit size). Perfused spinal cords were dissected and post-fixed in 4% paraformaldehyde for ~24-72 hrs. Next, fixed spinal cords were washed with PBS and cryoprotected in 30% sucrose (in PBS) until sinking was observed. Cryoprotected spinal cords were embedded in optimal cutting temperature (OCT) compound and frozen at −80 °C until cryostat sectioning (Leica Lm 1850, Germany). Transverse lumbar spinal cord cryosections were collected in serial at 30-μm or 40-μm thickness; to avoid repeat regions of analysis, every 15th section was used for analysis.

IHC for glial fibrillary acidic protein (GFAP): Slides were thawed, dried in an oven at 37 °C to improve section adherence, then rehydrated in PBS. Sections were blocked in 0.2% Triton X-100 and 5% normal horse serum (NHS) in PBS for 30 minutes at room temperature. Following blocking, slices were incubated in primary antibody solution containing mouse-anti GFAP antibody (diluted 1:500 in 0.1% Triton X-100 and 1% NHS in PBS; EnCor Biotechnology, cat. #MCA-5C10) overnight at room temperature. Sections were washed in PBS and then were incubated in secondary antibody solution consisting of Atto 488-goat anti-mouse IgG (whole molecule) (diluted 1:500 in 0.1% Triton X-100 and 1% NHS in PBS, Sigma-Aldrich, cat. # 62197) for 2 h at room temperature. Nuclei were labeled with 4',6-Diamidino-2-Phenylindole (DAPI) (Invitrogen™, cat. # D1306) and slides were mounted using Fluoromount mounting medium (SouthernBioTech). Z-stack photomicrographs of dorsal and ventral lower lumbar spinal cord were acquired at 10x magnification using a Nikon Eclipse Ti2 inverted confocal microscope. Maximum intensity projections of the green channel were generated and stitched (if possible) using Image Composite Editor (Microsoft). Grayscale images were imported into ImageJ and blinded densitometric analysis of GFAP immunofluorescent signal (mean gray value) in regions of interest containing dorsal columns, superficial dorsal gray matter, lateral spinothalamic tract, and ventral spinothalamic tract (divided by background fluorescent signal/mean gray value) was performed using Image J (NIH). The grey or white matter in our regions of interest was targeted to encompass the entire region of interest but could also contain portions of other areas.

IHC for primary afferent fibers: Sections were rinsed in PBS, blocked in 10% normal goat serum and 0.2% Triton-X100 in PBS for 1 hour at room temperature followed by a 30-minute incubation with rabbit-to-rabbit blocking reagent (ScyTek Laboratories, Inc.). Sections were then incubated with a solution of biotinylated IB4 (1:2000, Sigma) and anti-calcitonin gene-regulated peptide (CGRP; 1:1500 Peninsula Laboratories) in 0.1% Triton X-100 in PBS at room temperature for 48 h. After three washes with PBS, sections were incubated in avidin-conjugated with Alexa Fluor 594 (Invitrogen, diluted at 1:400) and goat-anti-rabbit Alexa Fluor 488 (Invitrogen, diluted at 1:400) in 5% goat serum in PBS for 2 hours at room temperature. Following three washes, sections were rinsed in PBS and cover slipped with FluorSave Reagent (Calbiochem, Bedford, MA). Three images of the left and right lumbar dorsal horn were taken for each rabbit at the same magnification, lens aperture and exposure time on a Leica DM5500B epifluorescent microscope and captured with Slidebook software. Images were averaged to calculate values per animal. The proportional area of IB4, CGRP and colocalized immunoreactive tissue within the cap of the dorsal horn (laminae I-III) was quantified in ImageJ as previously described (Detloff

et al., 2016). The proportional area of CGRP+ fibers that expand in dorsal horn ventral to the IB4+ band
was measured in the same sections.

Statistical Analyses: All statistical analyses were performed using GraphPad Prism (GraphPad Software
Inc.; San Diego, CA, USA). For each dataset, normality of distributions was tested using D'Agostino and
Pearson tests. For normally distributed datasets with equal variances, unpaired t-tests were used to
identify statistically significant differences between groups. For datasets in which one or more
distributions was non-normally distributed, Mann-Whitney tests were used.

Results:

In this study, we investigated the link between prenatal HI injury and nociception in a rabbit model
of CP. Rabbit kits exposed to prenatal HI and age-matched sham animals underwent testing for
mechanical and thermal allodynia at P5 using von Frey and Hargreaves' testing, respectively. After
prenatal HI injury, neonatal rabbit kits displayed increased nociceptive behavior in response to
mechanical and thermal stimuli that manifested even in the absence of motor deficits. HI kits
demonstrated significantly greater sensitivity to von Frey filaments applied to the paws compared to
sham kits, withdrawing paws in response to application of lower forces (forepaws, P=0.0014; hindpaws,
P=0.0043) (**Figure 1A**). Hindpaw von Frey scores of motor-unaffected HI kits (mean: 8.2 g; light red
symbols in Figure 1A) were significantly lower than that of motor-affected HI kits (mean: 19.4 g; dark red
symbols; P=0.0416). There was a similar but non-significant trend in forelimb paw withdrawal thresholds
between motor-unaffected (mean: 5.6 g; light red symbols in Figure 1A) and motor-affected HI kits
(mean: 10.5 g; dark red symbols; P=0.9068).

Similar to von Frey, the Hargreaves' test revealed allodynia in HI kits. HI kits had reduced paw
withdrawal latencies compared to sham rabbits in response to targeted radiant heat (**Figure 1B**),
indicating increased thermal sensitivity (forepaws: P<0.0001; hindpaws: P=0.0011). Unlike with
mechanical stimulation, paw withdrawal latencies to noxious thermal stimulation were comparable
between motor-unaffected HI (forepaw latency: 3.6 s; hindpaw latency: 3.0 s) and motor-affected HI kits
(forepaw latency: 4.4 s; hindpaw latency: 3.8 s) (P=0.1637 and 0.2711, respectively).

Figure 1: HI kits exhibit mechanical and thermal allodynia. **A:** Average forepaw (left) and hindpaw (right) von Frey scores at P5. Sham-operated controls (blue) have higher scores, indicating less mechanosensitivity, while HI kits (red) display greater mechanosensitivity (lower scores). Lower von Frey scores indicate mechanical allodynia in HI kits. Data are represented as mean ± SEM; N=32 sham and N=30 HI rabbits (dark red, motor-affected). **P <0.01; Mann-Whitney tests. **B:** Average forepaw (left) and hindpaw (right) withdrawal latencies during Hargreaves testing at P5. Shorter latencies to paw withdrawal in response to radiant heat are observed in HI kits compared to sham animals, indicative of enhanced thermal sensitivity. Data are represented as mean ± SEM; N=12 sham and N=18 HI rabbits (dark red, motor-affected). **P <0.01, ****P <0.0001; unpaired t-tests.

We hypothesized that mechanical and thermal allodynia in HI rabbit kits may be caused by spinal astrogliosis and/or expansion of nociceptive afferents in the spinal cord dorsal horn. To determine whether prenatal HI leads to long-lasting postnatal astrogliosis in gray and white matter spinal cord regions involved in cutaneous sensation or nociception, the degree of glial fibrillary acidic protein (GFAP) expression in four lumbar spinal cord regions of interest (ROIs) was quantified. These ROIs included tracts/laminae involved in cutaneous sensation or nociceptive neurotransmission: the dorsal column, superficial dorsal gray matter, ventral spinothalamic tract, and lateral spinothalamic tract. Densitometric analysis revealed no significant differences in GFAP labeling in these ROIs at P5 (**Figure 2B**). Baseline GFAP immunoreactivity in the white matter of neonatal rabbits found here is consistent with the role of astrocytes in myelination during early postnatal development (Landry et al., 1990; Liedtke et al., 1996; Nash et al., 2011). No evidence was found to support a contribution of neuroinflammation to increased nociception at this age.

Figure 2: Astrogliosis is not present in HI rabbits at P5. **A**: Transverse lumbar spinal cord sections from sham (left) and HI (right) rabbits immunostained for GFAP (green). Scale bar applies to both images. **B:** Quantification of GFAP densitometry in spinal regions containing tracts/laminae of interest: dorsal column (DC), superficial dorsal gray matter (SDGM), ventral spinothalamic tract (VST), and lateral spinothalamic tract (LST). GFAP expression was unchanged in all ROIs in HI spinal cords (ns, not significant; unpaired t-tests). Data are represented as mean ± SEM; N=3-4 rabbits per group.

To determine whether prenatal HI affects nociceptive primary afferent distribution and density
within the dorsal horn of the spinal cord, lumbar spinal cords (which include hindpaw dermatomes)
from sham and HI kits were labeled for CGRP and IB4 to identify peptidergic and non-peptidergic
nociceptors, respectively. The distributions of these two nociceptor subpopulations had little overlap in
the dorsal horn of sham kits (**Figure 3 A-A''**), however, HI led to an increased overlap in the topographic
distribution of these two fiber types (Figure 3 B-B''). In addition, HI kits exhibited increased CGRP-
positive fibers in the deep dorsal horn (Figure C-C'') that were not present in sham kits. Quantification of
the proportional area of each afferent subpopulation independently revealed that CGRP+ fibers (Figure
3D, G) but not IB4+ (Figure 3E) fibers exhibited increased density in the superficial and deep dorsal horn.
Despite no increase in the density or distribution of IB4-positive fibers in the dorsal horn,

Figure 3: Increased density and overlap of nociceptive primary afferent fibers in HI kits at P5. Representative images of the lumbar dorsal horn of sham (A-A'') and HI (B-B'') kits immunolabeled for calcitonin gene-regulated peptide (CGRP; A, B, C) or labeled by isolectin B-4 (IB4; A', B', C'). Note the increased presence of CGRP-positive fibers in the deep dorsal horn in the HI but not the sham spinal cord (C-C''). The proportional area of CGRP-positive tissue was significantly increased in the superficial and deep dorsal horn of HI rabbits compared to sham (D, G), while the density and distribution of IB4-positive afferents was unchanged between groups (E). The degree of overlap of these two nociceptor populations can be viewed in the merged images (A'', B'') and was significantly increased in HI kits compared to sham kits (F). Data are represented as mean ± SEM; individual data points represent each animal (n=5 sham and n=6 HI kits) from averages of 3 separate images of left and right dorsal horns per animal. **P<0.01, ***P<0.001; unpaired t-tests were used for statistical analysis.

there was significant colocalization of CGRP and IB4 labeling in the dorsal horn of HI kits compared to
sham kits, suggesting that HI may induce plasticity or phenotype switching of these nociceptor
populations.

Conclusions:

Experimental animal models that mimic human disease with high fidelity are more likely to drive
meaningful discovery of therapeutic interventions to treat such diseases. Here, we use a preclinical
model that displays prominent motor deficits to show that nociceptive afferents are also disrupted, in
addition to the previously characterized damage to the cortex, thinning of the corticospinal tract, loss of
spinal motoneurons and altered motoneuron excitability (Buser et al., 2010; Drobyshevsky et al., 2007;
Drobyshevsky and Quinlan, 2017; Steele et al., 2020). Importantly, both rabbits after HI and human
infants after asphyxiation show cell death in the dorsal horn (Clancy et al., 1989; Drobyshevsky and
Quinlan, 2017; Sladky and Rorke, 1986). Moreover, this is one of few studies that utilize quantitative
assessments of nociception in the rabbit, despite the rabbit's extensive use in biomedical research (Lei
et al., 2017; Mans, 2020; Yoshioka et al., 1996). Our findings indicate that after prenatal HI, sprouting of
peptidergic nociceptive primary afferent fibers in the superficial and deep dorsal horn is present along
with behavior that is suggestive of allodynia.

This is the first study to show that nociceptors are altered in an animal model of CP. The
increased overlap and density of nociceptive afferent fibers in the dorsal horn of HI kits may indicate
intrinsic changes within nociceptive DRG neurons. Despite significantly lower von Frey scores in HI kits
indicating increased mechanosensitivity, non-peptidergic IB4-positive mechanical nociceptors, which are
activated by noxious mechanical stimuli (Marmigère and Ernfors, 2007), were not found to have
expanded afferent arborizations in the dorsal horn. It could be that these IB4-positive nociceptors are
hyperexcitable, and that altered activity, not anatomy, contributes to mechanical allodynia in HI kits. In
other models of nervous system injury, chronic pain is associated with dysfunctional nociceptors that
exhibit robust anatomical and electrophysiological plasticity (Bedi et al., 2010; Detloff et al., 2016, 2014;
Dougherty et al., 2004; Moy et al., 2018; Zhang et al., 2013); further, emerging evidence suggests that
nociceptive primary afferent input impedes motor output following CNS injury (Keller et al., 2017, 2019,
2018). In a series of studies, Keller *et al*. demonstrated that nociceptor depletion reduces muscle
contractures and spasticity of the hindlimbs and improves overall locomotor performance following
spinal cord injury. Nociceptor silencing via activation of inhibitory $5HT_{1D}$ receptors has also been found
to be helpful in reducing clonus and spasms after spinal cord injury (Lucas-Osma et al., 2019). Taken
together, these findings indicate that altered activity in nociceptors contributes to motor deficits in
cases of CNS injury.

In adults, physical trauma is known to alter neural circuits and have long lasting effects on
nociception (Costigan et al., 2009; Price and Dussor, 2014). The effect of neural injury on nociceptive
primary sensory neurons is a major contributor to the development of chronic pain (Hulsebosch and
Coggeshall, 1981a, 1981b; Koerber et al., 1994; Ramer et al., 2012; Rodin et al., 1983; Walters, 2012;
Woolf et al., 1992). After trauma, it is well-established that the afferent fibers of these nociceptors
exhibit robust arborizations into the superficial and deep dorsal horn of the spinal cord (Bedi et al.,
2010; Detloff et al., 2016, 2014; Krenz and Weaver, 1998; Ondarza et al., 2003; Weaver et al., 2001). In

addition, nociceptors increase spontaneous activity and become hyperexcitable following CNS trauma
(Bedi et al., 2010; Burchiel et al., 1985; Gold and Gebhart, 2010; Gracely et al., 1993; Li et al., 2018;
North et al., 2019; Pitcher and Henry, 2008). In neonates, effects of neural injury on nociception are less
studied. Peripheral inflammation in neonatal rats enhances collateral sprouting of primary nociceptive
afferents and enhances nociception in adulthood (Ruda et al., 2000). Astrocytes and microglia are
activated by neural injury and release a plethora of pro-inflammatory molecules into the spinal milieu,
and their activation corresponds to the development of neuropathic pain (Chhaya et al., 2018; Detloff et
al., 2008; Gao et al., 2009; Hulsebosch et al., 2009; Ishikura et al., 2021; Lu et al., 2021; Xie et al., 2009;
Zhang et al., 2016, 2007).

Neuroinflammation and pain are tightly intertwined (Grace et al., 2014) and astrogliosis has a
well-established role in other painful conditions. After complete spinal cord injury, increased GFAP
persists for at least 3 weeks in the adult rat spinal cord (Choi et al., 2021; Ducza et al., 2021; Morin-
Richaud et al., 1998; Nesic et al., 2006; Xu et al., 2021). However, the time course of the GFAP response
to injury is different in neonates. Unlike adults, neonatal rat pups respond to spared nerve injury with an
early increase in GFAP (one day after injury), along with the same elevation after 7 days that is observed
in adults (Vega-Avelaira et al., 2007). Density of GFAP fiber length and density of GFAP+ cell bodies are
increased in the lumbar gray matter of motor-affected (hypertonic) HI rabbit kits at P1, seven days after
HI (Drobyshevsky and Quinlan, 2017; Synowiec et al., 2019). The current study was performed in HI kits
that were 4 days older (P5) and examined specific laminae/tracts involved in cutaneous sensation or
nociception: (1) the dorsal column, the ascending pathway for cutaneous sensation; (2) the superficial
dorsal gray matter, containing second-order projection neurons relaying nociception; (3) the ventral
spinothalamic tract, carrying crude touch and pressure information; and (4) the lateral spinothalamic
tract, the ascending pathway for nociception. While astrogliosis is not present in the spinal regions of
interest we studied at P5 (11-14 days after HI), their reactivity closer to the time of injury could be a
causal link of enhanced nociception as has been demonstrated in other models (Milligan et al., 2000).
Astrocytes have been shown to exhibit different phenotypes or reactive states following similar CNS
injuries that can be either destructive or pro-reparative (Escartin et al., 2021; Liddelow and Barres,
2017). Moreover, it has been reported that ischemia can induce an "A2-like" astrocyte phenotype that
upregulates thrombospondins and neurotrophic factors that promote survival and repair of neurons
(Gao et al., 2005; Hayakawa et al., 2014; Zador et al., 2009). Future studies could examine more closely
the time course and specific regions of elevated astrocyte activation in the rabbit after prenatal HI and
its relationship to enhanced nociception observed here at P5.

The stereotypical aversive paw withdrawal response to nocifensive tactile or thermal stimuli is
mitigated by a segmental spinal reflex and is often inferred as a proxy for cortical perception of the
stimulus. It is unclear whether the results of the von Frey and Hargreaves tests performed here reflect
hyperreflexia of nociceptive segmental spinal cord circuits (Advokat, 2002; Kauppila et al., 1998; Murphy
and Zemlan, 1990; Schomburg and Steffens, 1998) or the complicated perception of neuropathic pain. It
is likely these stimuli were offensive to the rabbits because paw withdrawal was typically accompanied
by paw licking and/or a rapid jump away from the stimuli. Unlike in rodent experiments, there are
surprisingly few reports measuring pain or nociception in experimental rabbit models, even those
modeling knee osteoarthritis or intervertebral disc dislocation (Lei et al., 2017; Mans, 2020; Yoshioka et
al., 1996). Composite pain scales for rabbits in a clinical environment (Banchi et al., 2020) and the Bristol
Rabbit Pain Scale (Benato et al., 2021) examine and score rabbit facial expressions and other behaviors
to assess pain, such as the grimace scale (Evangelista et al., 2021; Keating et al., 2012). Yet, recent

reports suggest that the experimenter's presence can have a profound effect on the animal's behavior and subsequent score on these tests (Pinho et al., 2020). In the present study, unlike assessments of spontaneous pain using the rabbit grimace scale, these assessments identify the dermatomal location and relevant spinal cord or dorsal root ganglion level in which dysfunctional sensation occurs. This is especially useful in examining changes in primary afferent input and/or dorsal horn processing in specific spinal segments. Also, since our study examined an early postnatal age (P5) when the neonatal rabbits' eyes are not yet open, the tests we performed were more appropriate to assess nociceptive behavior than using the grimace scale, place-preference or aversion tests. However, future work to assess pain perception and an associated fear response at later postnatal ages and could include these assessments to further characterize the perception of pain and translate these findings clinically.

Individuals with CP have disruptions in somatosensation in addition to motor dysfunction. Disruptions in somatosensory cortex connectivity (Hoon et al., 2002) and increased latency of sensory evoked potentials (Cahan et al., 1987; Kundi et al., 1989) have been well documented in individuals with CP. Innocuous tactile sensitivity actually decreases in individuals with CP while pain sensitivity is enhanced (Riquelme and Montoya, 2010). The mechanism of this is not clear, but it is possible that this is due to afferent switching after injury: in a rat model of neuropathic pain, large diameter cutaneous/somatic Aβ afferents can undergo fiber-type switching after nerve ligation and begin expressing CGRP, like nociceptive fibers (Nitzan-Luques et al., 2011). This switch in identity corresponds to a period of tactile allodynia. Another possibility is that the impaired CST creates a permissive environment for afferent sprouting. When competition between CST and afferent input to the spinal cord is lost after transection of the pyramidal tracts in adult rats, afferent sprouting into the deep dorsal horn is observed (Jiang et al., 2016; Tan et al., 2012). Afferent sprouting after CST transection is accompanied by loss of rate dependent depression of the H reflex, a hallmark of CP (Tan et al., 2012). The common thread in both of these mechanisms is the remarkable plasticity of afferents, even in adulthood. This raises the possibility that aberrant nociceptive sprouting arising during perinatal development could be corrected later in life.

Movement and motor unit firing patterns in neurotypical adults are directly affected by pain (Hodges and Tucker, 2011; Martinez-Valdes et al., 2020), perhaps through neuromodulators including serotonin (5HT) (Mesquita et al., 2020). In our study, we found that HI kits were more sensitive to mechanical and thermal stimuli than sham control kits. When comparing HI kits that were motor-affected and those that were motor-unaffected, motor-affected HI kits appeared less sensitive to mechanical stimuli and their response to thermal stimuli trended in the same direction. However, we attribute this to the greater difficulty these kits had in making movements to respond to stimuli due to their hypertonia. No kits in this study were completely incapable of movement and unable to react physically to the stimuli, but their hypertonia could have impeded their reaction. Further investigation into differences between motor-affected and motor-unaffected HI kits warrants further study.

Each individual's pain perception includes complex psychosocial components that were not the focus of this study. In situations where both nociception and the affective experience of pain can be altered, such as after HI, it is critical to accurately identify and classify aberrant sensory behavior. Here, we utilized methods to assess and quantify shifts in sensitivity to mechanical and thermal stimuli that are sensitive, reliable, and validated in models of central and peripheral neuropathic pain across species. We found that prenatal HI heightened nociception at the level of spinal nociceptors and reflexive responses to mechanical and thermal stimuli. It could be that CP-causative perinatal injuries contribute to other aspects of the pain experience in individuals with CP, including changes to mood and cognition, in addition to altered nociception. Thus, treatment options for chronic pain in individuals with CP should

412 be explored with greater specificity, considering that the pathophysiology of their pain could be distinct
413 from that of chronic pain derived from other etiologies. Novel targets for fully alleviating CP-induced
414 chronic pain should be explored specifically within the context of treating individuals with CP or those
415 who have survived perinatal physical trauma.
416
417 The current study demonstrates enhanced nociceptive behavior and nociceptor sprouting in
418 neonatal rabbits after prenatal HI injury. Identification of how prenatal physical stressors impact
419 nociceptors could be the first step to reducing hyperreflexia and preventing aberrant nociception and
420 ultimately chronic pain in those at risk.

Acknowledgements

422 This project was supported by funding from NIH NINDS including NS097880 to MRD and
423 NS104436 to KQ. MCB and DMB received a grant from the University of Rhode Island Undergraduate
424 Research & Innovation. The funders of this project had no role in study design, the collection, analysis and
425 interpretation of data, in writing the report, or in the decision to publish these results.

Author contributions

EJ Reedich contributed to data curation, formal analysis, investigation, methodology, supervision,
validation and writing, editing and reviewing the manuscript. LT Genry contributed to data curation,
formal analysis, investigation, methodology, supervision, validation, and reviewing and editing the
manuscript. MA Singer contributed to data curation, formal analysis, investigation, methodology, and
validation. CF Cavarsan contributed to formal analysis, investigation, methodology, and validation. E
Mena Avila contributed to data curation, formal analysis, investigation, methodology, validation, and
reviewing the manuscript. DM Boudreau contributed to data curation, formal analysis, investigation,
methodology, validation, and reviewing the manuscript. MC Brennan contributed to data curation,
formal analysis, investigation, methodology, validation, and reviewing the manuscript. AM Garrett
contributed to data curation, formal analysis, investigation, methodology, validation, and reviewing the
manuscript. L Dowaliby contributed to investigation, methodology and supervision of this project. MR
Detloff contributed to contributed to conceptualization, data curation, funding acquisition, project
administration, resources, supervision, and writing, editing and review of the manuscript. KA Quinlan
contributed to conceptualization, data curation, funding acquisition, project administration, resources,
supervision, and writing, editing and review of the manuscript.
442

References

Achache, V., Roche, N., Lamy, J.C., Boakye, M., Lackmy, A., Gastal, A., Quentin, V., Katz, R., 2010.
Transmission within several spinal pathways in adults with cerebral palsy. Brain 133, 1470–1483.
https://doi.org/10.1093/brain/awq053

Advokat, C., 2002. Spinal transection increases the potency of clonidine on the tail-flick and hindlimb
flexion reflexes. Eur. J. Pharmacol. 437, 63–67. https://doi.org/10.1016/S0014-2999(02)01259-1

Badia, M., Riquelme, I., Orgaz, B., Acevedo, R., Longo, E., Montoya, P., 2014. Pain, motor function and
health-related quality of life in children with cerebral palsy as reported by their physiotherapists.
BMC Pediatr. 14, 2–7. https://doi.org/10.1186/1471-2431-14-192

Banchi, P., Quaranta, G., Ricci, A., Von Degerfeld, M.M., 2020. Reliability and construct validity of a composite pain scale for rabbit (CANCRS) in a clinical environment. PLoS One 15, 1–12. https://doi.org/10.1371/journal.pone.0221377

Bedi, S.S., Yang, Q., Crook, R.J., Du, J., Wu, Z., Fishman, H.M., Grill, R.J., Carlton, S.M., Walters, E.T., 2010. Chronic spontaneous activity generated in the somata of primary nociceptors is associated with pain-related behavior after spinal cord injury. J. Neurosci. 30, 14870–14882. https://doi.org/10.1523/JNEUROSCI.2428-10.2010

Benato, L., Murrell, J., Knowles, T.G., Rooney, N.J., 2021. Development of the Bristol Rabbit Pain Scale (BRPS): A multidimensional composite pain scale specific to rabbits (Oryctolagus cuniculus). PLoS One 16, 1–18. https://doi.org/10.1371/journal.pone.0252417

Black, S., 2019. C Fibres facilitate proprioceptive afferent transmission to motoneurons. Masters Thesis, Univ. Alberta.

Bumin, G., Kayihan, H., 2001. Effectiveness of two different sensory-integration programmes for children with spastic diplegic cerebral palsy. Disabil. Rehabil. 23, 394–399. https://doi.org/10.1080/09638280010008843

Burchiel, K.J., Russell, L.C., Lee, R.P., Sima, A.A.F., 1985. Spontaneous Activity of Primary Afferent Neurons in Diabetic BB/Wistar Rats: A possible mechanism of Diabetic Neuropathic Pain. Diabetes 34.

Buser, J.R., Segovia, K.N., Dean, J.M., Nelson, K., Beardsley, D., Gong, X., Luo, N.L., Ren, J., Wan, Y., Riddle, A., Mcclure, M.M., Ji, X., Derrick, M., Hohimer, A.R., Back, S.A., Tan, S., 2010. Timing of appearance of late oligodendrocyte progenitors coincides with enhanced susceptibility of preterm rabbit cerebral white matter to hypoxia-ischemia. J. Cereb. Blood Flow Metab. 30, 1053–1065. https://doi.org/10.1038/jcbfm.2009.286

Cahan, L.D., Kundi, M.S., McPherson, D., Starr, A., Peacock, W., 1987. Electrophysiologic Studies in Selective Dorsal Rhizotomy for Spasticity in Children with Cerebral Palsy. Appl. Neurophysiol. 50, 459–460.

Chhaya, S.J., Quiros-Molina, D., Tamashiro-Orrego, A.D., Houle, J.D., Detloff, M.R., 2018. Exercise-Induced Changes to the Macrophage Response in the Dorsal Root Ganglia Prevent Neuropathic Pain after Spinal Cord Injury. J. Neurotrauma 14, neu.2018.5819. https://doi.org/10.1089/neu.2018.5819

Choi, J.-G., Choi, S.-R., Kang, D.-W., Kim, J., Park, J.B., Lee, J.-H., Kim, H.-W., 2021. Sigma-1 receptor increases intracellular calcium in cultured astrocytes and contributes to mechanical allodynia in a model of neuropathic pain. Brain Res. Bull. 178, 69–81. https://doi.org/10.1016/j.brainresbull.2021.11.010

Clancy, R.R., Sladky, J.T., Rorke, L.B., 1989. Hypoxic-ischemic spinal cord injury following perinatal asphyxia. Ann. Neurol. 25, 185–189. https://doi.org/10.1002/ana.410250213

Cooper, J., Majnemer, A., Rosenblatt, B., 1995. The Determination of Sensory Deficits in Children With Hemiplegic Cerebral Palsy. J. Child Neurol. 10, 300–309. https://doi.org/10.1177/088307389501000412

Costigan, M., Scholz, J., Woolf, C.J., 2009. Neuropathic pain: A maladaptive response of the nervous system to damage. Annu. Rev. Neurosci. 32, 1–32.

493 https://doi.org/10.1146/annurev.neuro.051508.135531

494 Derrick, M., Luo, N.L., Bregman, J.C., Jilling, T., Ji, X., Fisher, K., Gladson, C.L., Beardsley, D., Murdoch, G.,
495  Back, S.A., Tan, S., 2004. Preterm Fetal Hypoxia-Ischemia Causes Hypertonia and Motor Deficits in
496  the Neonatal Rabbit: A Model for Human Cerebral Palsy? J. Neurosci. 24, 24–34.
497  https://doi.org/10.1523/JNEUROSCI.2816-03.2004

498 Detloff, M.R., Fisher, L.C., McGaughy, V., Longbrake, E.E., Popovich, P.G., Basso, D.M., 2008. Remote
499  activation of microglia and pro-inflammatory cytokines predict the onset and severity of below-
500  level neuropathic pain after spinal cord injury in rats. Exp. Neurol. 212, 337–347.
501  https://doi.org/10.1016/j.expneurol.2008.04.009

502 Detloff, M.R., Quiros-Molina, D., Javia, A.S., Daggubati, L., Nehlsen, A.D., Naqvi, A., Ninan, V., Vannix,
503  K.N., McMullen, M.K., Amin, S., Ganzer, P.D., Houlé, J.D., 2016. Delayed Exercise Is Ineffective at
504  Reversing Aberrant Nociceptive Afferent Plasticity or Neuropathic Pain after Spinal Cord Injury in
505  Rats. Neurorehabil. Neural Repair 30, 685–700. https://doi.org/10.1177/1545968315619698

506 Detloff, M.R., Smith, E.J., Quiros Molina, D., Ganzer, P.D., Houlé, J.D., 2014. Acute exercise prevents the
507  development of neuropathic pain and the sprouting of non-peptidergic (GDNF- and artemin-
508  responsive) c-fibers after spinal cord injury. Exp. Neurol. 255, 38–48.
509  https://doi.org/10.1016/j.expneurol.2014.02.013

510 Doralp, S., Bartlett, D.J., 2010. The prevalence, distribution, and effect of pain among adolescents with
511  cerebral palsy. Pediatr. Phys. Ther. 22, 26–33. https://doi.org/10.1097/PEP.0b013e3181ccbabb

512 Dougherty, P.M., Cata, J.P., Cordella, J. V., Burton, A., Weng, H.R., 2004. Taxol-induced sensory
513  disturbance is characterized by preferential impairment of myelinated fiber function in cancer
514  patients. Pain 109, 132–142. https://doi.org/10.1016/j.pain.2004.01.021

515 Drobyshevsky, A., Derrick, M., Wyrwicz, A.M., Ji, X., Englof, I., Ullman, L.M., Zelaya, M.E., Northington,
516  F.J., Tan, S., 2007. White matter injury correlates with hypertonia in an animal model of cerebral
517  palsy. J. Cereb. Blood Flow Metab. 27, 270–281. https://doi.org/10.1038/sj.jcbfm.9600333

518 Drobyshevsky, A., Quinlan, K.A., 2017. Spinal cord injury in hypertonic newborns after antenatal
519  hypoxia-ischemia in a rabbit model of cerebral palsy. Exp. Neurol. 293, 13–26.
520  https://doi.org/10.1016/j.expneurol.2017.03.017

521 Ducza, L., Szücs, P., Hegedűs, K., Bakk, E., Gajtkó, A., Wéber, I., Holló, K., 2021. Nlrp2 is overexpressed in
522  spinal astrocytes at the peak of mechanical pain sensitivity during complete freund adjuvant-
523  induced persistent pain. Int. J. Mol. Sci. 22. https://doi.org/10.3390/ijms222111408

524 Escartin, C., Galea, E., Lakatos, A., O'Callaghan, J.P., Petzold, G.C., Serrano-Pozo, A., Steinhäuser, C.,
525  Volterra, A., Carmignoto, G., Agarwal, A., Allen, N.J., Araque, A., Barbeito, L., Barzilai, A., Bergles,
526  D.E., Bonvento, G., Butt, A.M., Chen, W.T., Cohen-Salmon, M., Cunningham, C., Deneen, B., De
527  Strooper, B., Díaz-Castro, B., Farina, C., Freeman, M., Gallo, V., Goldman, J.E., Goldman, S.A., Götz,
528  M., Gutiérrez, A., Haydon, P.G., Heiland, D.H., Hol, E.M., Holt, M.G., Iino, M., Kastanenka, K. V.,
529  Kettenmann, H., Khakh, B.S., Koizumi, S., Lee, C.J., Liddelow, S.A., MacVicar, B.A., Magistretti, P.,
530  Messing, A., Mishra, A., Molofsky, A. V., Murai, K.K., Norris, C.M., Okada, S., Oliet, S.H.R., Oliveira,
531  J.F., Panatier, A., Parpura, V., Pekna, M., Pekny, M., Pellerin, L., Perea, G., Pérez-Nievas, B.G.,
532  Pfrieger, F.W., Poskanzer, K.E., Quintana, F.J., Ransohoff, R.M., Riquelme-Perez, M., Robel, S., Rose,
533  C.R., Rothstein, J.D., Rouach, N., Rowitch, D.H., Semyanov, A., Sirko, S., Sontheimer, H., Swanson,
534  R.A., Vitorica, J., Wanner, I.B., Wood, L.B., Wu, J., Zheng, B., Zimmer, E.R., Zorec, R., Sofroniew, M.

V., Verkhratsky, A., 2021. Reactive astrocyte nomenclature, definitions, and future directions. Nat.
Neurosci. 24, 312–325. https://doi.org/10.1038/s41593-020-00783-4

Evangelista, M.C., Monteiro, B.P., Steagall, P. V, 2021. Measurement properties of grimace scales for
pain assessment in non-human mammals, Pain. https://doi.org/10.1097/j.pain.0000000000002474

Findlay, B., Switzer, L., Narayanan, U., Chen, S., Fehlings, D., 2016. Investigating the impact of pain, age,
Gross Motor Function Classification System, and sex on health-related quality of life in children
with cerebral palsy. Dev. Med. Child Neurol. 58, 292–297. https://doi.org/10.1111/dmcn.12936

Flanigan, M., Gaebler-Spira, D., Kocherginsky, M., Garrett, A., Marciniak, C., 2020. Spasticity and pain in
adults with cerebral palsy. Dev. Med. Child Neurol. 62, 379–385.
https://doi.org/10.1111/dmcn.14368

Gao, Q., Li, Y., Chopp, M., 2005. Bone marrow stromal cells increase astrocyte survival via upregulation
of phosphoinositide 3-kinase/threonine protein kinase and mitogen-activated protein kinase
kinase/extracellular signal-regulated kinase pathways and stimulate astrocyte trophic factor .
Neuroscience 136, 123–134. https://doi.org/10.1016/j.neuroscience.2005.06.091

Gao, Y.J., Zhang, L., Samad, O.A., Suter, M.R., Yasuhiko, K., Xu, Z.Z., Park, J.Y., Lind, A.L., Ma, Q., Ji, R.R.,
2009. JNK-induced MCP-1 production in spinal cord astrocytes contributes to central sensitization
and neuropathic pain. J. Neurosci. 29, 4096–4108. https://doi.org/10.1523/JNEUROSCI.3623-
08.2009

García Jalón, E.G., Maguire, A., Perra, O., Gavin, A., O'Reilly, D., Thurston, A., 2021. Data linkage and pain
medication in people with cerebral palsy: a cross-sectional study. Dev. Med. Child Neurol. 1–8.
https://doi.org/10.1111/dmcn.14854

Gold, M.S., Gebhart, G.F., 2010. Nociceptor sensitization in pain pathogenesis. Nat. Med. 16, 1248–1257.
https://doi.org/10.1038/nm.2235

Grace, P.M., Hutchinson, M.R., Maier, S.F., Watkins, L.R., 2014. Pathological pain and the neuroimmune
interface. Nat. Rev. Immunol. 14, 217–231. https://doi.org/10.1038/nri3621

Gracely, R.H., Lynch, S.A., Bennett, G.J., 1993. Painful neuropathy: altered central processing maintained
dynamically by peripheral input. Pain 52, 251–253. https://doi.org/10.1016/0304-3959(93)90141-B

Gupta, D., Barachant, A., Gordon, A.M., Ferre, C., Kuo, H.C., Carmel, J.B., Friel, K.M., 2017. Effect of
sensory and motor connectivity on hand function in pediatric hemiplegia. Ann. Neurol. 82, 766–
780. https://doi.org/10.1002/ana.25080

Hari, K., Lucas-Osma, A.M., Metz, K., Lin, S., Pardell, N., Roszko, D., Black, S., Minarik, A., Singla, R.,
Stephens, M.J., Fouad, K., Jones, K.E., Gorassini, M.A., Fenrich, K.K., Li, Y., Bennett, D.J., 2021.
Nodal GABA facilitates axon spike transmission in the spinal cord. bioRxiv 2021.01.20.427494.

Hayakawa, K., Pham, L.D.D., Arai, K., Lo, E.H., 2014. Reactive astrocytes promote adhesive interactions
between brain endothelium and endothelial progenitor cells via HMGB1 and beta-2 integrin
signaling. Stem Cell Res. 12, 531–538. https://doi.org/10.1016/j.scr.2013.12.008

Hochman, S., Shreckengost, J., Kimura, H., Quevedo, J., 2010. Presynaptic inhibition of primary afferents
by depolarization : observations supporting nontraditional mechanisms 1198, 140–152.
https://doi.org/10.1111/j.1749-6632.2010.05436.x

Hodges, P.W., Tucker, K., 2011. Moving differently in pain: A new theory to explain the adaptation to pain. Pain 152, S90–S98. https://doi.org/10.1016/j.pain.2010.10.020

Hoon, A.H., Lawrie, W.T., Melhem, E.R., Reinhardt, E.M., Van Zijl, P.C.M., Solaiyappan, M., Jiang, H., Johnston, M. V., Mori, S., 2002. Diffusion tensor imaging of periventricular leukomalacia shows affected sensory cortex white matter pathways. Neurology 59, 752–756. https://doi.org/10.1212/WNL.59.5.752

Hulsebosch, C.E., Coggeshall, R.E., 1981a. Quantitation of Sprouting of Dorsal Root Axons. Science 213, 1020–1021.

Hulsebosch, C.E., Coggeshall, R.E., 1981b. Sprouting of dorsal root axons. Brain Res. 224, 170–174. https://doi.org/10.1016/0006-8993(81)91128-8

Hulsebosch, C.E., Hains, B.C., Crown, E.D., Carlton, S.M., 2009. Mechanisms of chronic central neuropathic pain after spinal cord injury. Brain Res. Rev. 60, 202–213. https://doi.org/10.1016/j.brainresrev.2008.12.010

Ishikura, T., Kinoshita, M., Shimizu, M., Yasumizu, Y., Motooka, D., Okuzaki, D., Yamashita, K., Murata, H., Beppu, S., Koda, T., Tada, S., Shiraishi, N., Sugiyama, Y., Miyamoto, K., Kusunoki, S., Sugimoto, T., Kumanogoh, A., Okuno, T., Mochizuki, H., 2021. Anti-AQP4 autoantibodies promote ATP release from astrocytes and induce mechanical pain in rats. J. Neuroinflammation 18, 1–12. https://doi.org/10.1186/s12974-021-02232-w

Jiang, Y.-Q., Zaaimi, B., Martin, J.H., 2016. Competition with Primary Sensory Afferents Drives Remodeling of Corticospinal Axons in Mature Spinal Motor Circuits. J. Neurosci. 36, 193–203. https://doi.org/10.1523/JNEUROSCI.3441-15.2016

Kaelin-Lang, A., 2008. Enhancing rehabilitation of motor deficits with peripheral nerve stimulation. NeuroRehabilitation 23, 89–93. https://doi.org/10.3233/nre-2008-23109

Kandel, E.R., Schwartz, J.H., Jessell, T.M., Siegelbaum, S.A., Hudspeth, A.J. (Eds.), 2013. Principles of Neural Science, Fifth. ed. McGraw Hill, United States of America.

Kauppila, T., Kontinen, V.K., Pertovaara, A., 1998. Influence of spinalization on spinal withdrawal reflex responses varies depending on the submodality of the test stimulus and the experimental pathophysiological condition in the rat. Brain Res. 797, 234–242. https://doi.org/10.1016/S0006-8993(98)00379-5

Keating, S.C.J., Thomas, A.A., Flecknell, P.A., Leach, M.C., 2012. Evaluation of EMLA Cream for Preventing Pain during Tattooing of Rabbits: Changes in Physiological, Behavioural and Facial Expression Responses. PLoS One 7, 1–11. https://doi.org/10.1371/journal.pone.0044437

Keller, A.V.P., Wainwright, G., Shum-Siu, A., Prince, D., Hoeper, A., Martin, E., Magnuson, D.S.K., 2017. Disruption of locomotion in response to Hindlimb muscle stretch at acute and chronic time points after a spinal cord injury in rats. J. Neurotrauma 34, 661–670. https://doi.org/10.1089/neu.2015.4227

Keller, A. V., Hainline, C., Rees, K., Krupp, S., Prince, D., Wood, B.D., Shum-Siu, A., Burke, D.A., Petruska, J.C., Magnuson, D.S.K., 2019. Nociceptor-dependent locomotor dysfunction after clinically-modeled hindlimb muscle stretching in adult rats with spinal cord injury. Exp. Neurol. 318, 267–276. https://doi.org/10.1016/j.expneurol.2019.03.006

Keller, A. V., Rees, K.M., Seibt, E.J., Wood, B.D., Wade, A.D., Morehouse, J., Shum-Siu, A., Magnuson,
D.S.K., 2018. Electromyographic patterns of the rat hindlimb in response to muscle stretch after
spinal cord injury. Spinal Cord 56, 560–568. https://doi.org/10.1038/s41393-018-0069-z

Kimura, S., Honda, M., Tanabe, M., Ono, H., 2004. Noxious stimuli evoke a biphasic flexor reflex
composed of Aδ-fiber-mediated short-latency and C-fiber-mediated long-latency withdrawal
movements in mice. J. Pharmacol. Sci. 95, 94–100. https://doi.org/10.1254/jphs.95.94

Koerber, H.R., Mirnics, K., Brown, P.B., Mendell, L.M., 1994. Central sprouting and functional plasticity of
regenerated primary afferents. J. Neurosci. 14, 3655–3671. https://doi.org/10.1523/jneurosci.14-
06-03655.1994

Krenz, N.., Weaver, L.., 1998. Sprouting of primary afferent fibers after spinal cord transection in the rat.
Neuroscience 85, 443–458. https://doi.org/10.1016/S0306-4522(97)00622-2

Kundi, M., Starr, A., Cahan, L., 1989. Somatosensory Evoked Potentials in Cerebral Palsy After Partial
Dorsal Root Rhizotomy. Arch. Neurol. 46, 524–527.
https://doi.org/10.1001/archneur.1989.00520410058024

Lallemend, F., Ernfors, P., 2012. Molecular interactions underlying the specification of sensory neurons.
Trends Neurosci. 35, 373–381. https://doi.org/10.1016/j.tins.2012.03.006

Landry, C.F., Ivy, G.O., Brown, I.R., 1990. Developmental expression of glial fibrillary acidic protein mRNA
in the rat brain analyzed by in situ hybridization. J. Neurosci. Res. 25, 194–203.
https://doi.org/10.1002/jnr.490250207

Laurin, J., Pertici, V., Dousset, E., Marqueste, T., Decherchi, P., 2015. Group III and IV muscle afferents:
Role on central motor drive and clinical implications. Neuroscience 290, 543–551.
https://doi.org/10.1016/j.neuroscience.2015.01.065

Lei, T., Zhang, Y., Zhou, Q., Luo, X., Tang, K., Chen, R., Yu, C., Quan, Z., 2017. A novel approach for the
annulus needle puncture model of intervertebral disc degeneration in rabbits. Am. J. Transl. Res. 9,
900–909.

Leonard, C.T., Hirschfeld, H., 1995. Myotatic Reflex Responses of Non-Disabled Children and Children
with Spastic Cerebral Palsy. Dev. Med. Child Neurol. 37, 783–799.

Leonard, C.T., Hirschfeld, H., Moritani, T., Forssberg, H., 1991. Myotatic reflex development in normal
children and children with cerebral palsy. Exp. Neurol. 111, 379–382.
https://doi.org/10.1016/0014-4886(91)90106-M

Li, Y., North, R.Y., Rhines, L.D., Tatsui, C.E., Rao, G., Edwards, D.D., Cassidy, R.M., Harrison, D.S.,
Johansson, C.A., Zhang, H., Dougherty, P.M., 2018. DRG voltage-gated sodium channel 1.7 is
upregulated in paclitaxel-induced neuropathy in rats and in humans with neuropathic pain. J.
Neurosci. 38, 1124–1136. https://doi.org/10.1523/JNEUROSCI.0899-17.2017

Liddelow, S.A., Barres, B.A., 2017. Reactive Astrocytes: Production, Function, and Therapeutic Potential.
Immunity 46, 957–967. https://doi.org/10.1016/j.immuni.2017.06.006

Liedtke, W., Edelmann, W., Bieri, P.L., Chiu, F.C., Cowan, N.J., Kucherlapati, R., Raine, C.S., 1996. GFAP is
necessary for the integrity of CNS white matter architecture and long-term maintenance of
myelination. Neuron 17, 607–615. https://doi.org/10.1016/S0896-6273(00)80194-4

Lu, R., Cui, S.S., Wang, X.X., Chen, L., Liu, F., Gao, J., Wang, W., 2021. Astrocytic c-Jun N-terminal kinase-
histone deacetylase-2 cascade contributes to glutamate transporter-1 decrease and mechanical
allodynia following peripheral nerve injury in rats. Brain Res. Bull. 175, 213–223.
https://doi.org/10.1016/j.brainresbull.2021.07.024

Lucas-Osma, A.M., Li, Y., Murray, K., Lin, S., Black, S., Stephens, M.J., Ahn, A.H., Heckman, C.J., Fenrich,
K.K., Fouad, K., Bennett, D.J., 2019. 5-HT1D receptors inhibit the monosynaptic stretch reflex by
modulating C-fiber activity. J. Neurophysiol. 121, 1591–1608.
https://doi.org/10.1152/jn.00805.2018

Mans, C., 2020. Insights into pain assessment and management in rabbits. Vet. Rec. 186, 600–602.
https://doi.org/10.1136/vr.m2265

Marmigère, F., Ernfors, P., 2007. Specification and connectivity of neuronal subtypes in the sensory
lineage. Nat. Rev. Neurosci. 8, 114–127. https://doi.org/10.1038/nrn2057

Marriott, A.L., Rojas-Mancilla, E., Morales, P., Herrera-Marschitz, M., Tasker, R.A., 2017. Models of
progressive neurological dysfunction originating early in life. Prog. Neurobiol. 155, 2–20.
https://doi.org/10.1016/j.pneurobio.2015.10.001

Martinez-Valdes, E., Negro, F., Farina, D., Falla, D., 2020. Divergent response of low- versus high-
threshold motor units to experimental muscle pain. J. Physiol. 598, 2093–2108.
https://doi.org/10.1113/JP279225

Matusz, P.J., Key, A.P., Gogliotti, S., Pearson, J., Auld, M.L., Murray, M.M., Maitre, N.L., 2018.
Somatosensory Plasticity in Pediatric Cerebral Palsy following Constraint-Induced Movement
Therapy. Neural Plast. 2018. https://doi.org/10.1155/2018/1891978

McDowell, B.C., Duffy, C., Lundy, C., 2017. Pain report and musculoskeletal impairment in young people
with severe forms of cerebral palsy: A population-based series. Res. Dev. Disabil. 60, 277–284.
https://doi.org/10.1016/j.ridd.2016.10.006

Mckinnon, C.T., Meehan, E.M., Harvey, A.R., Antolovich, G.C., Morgan, P.E., 2018. Prevalence and
characteristics of pain in children and young adults with cerebral palsy: a systematic review. Dev.
Med. Child Neurol. 61, 305–314. https://doi.org/10.1111/dmcn.14111

McKinnon, C.T., Morgan, P.E., Antolovich, G.C., Clancy, C.H., Fahey, M.C., Harvey, A.R., 2020. Pain in
children with dyskinetic and mixed dyskinetic/spastic cerebral palsy. Dev. Med. Child Neurol. 62,
1294–1301. https://doi.org/10.1111/dmcn.14615

Mesquita, R.N.O., Škarabot, J., Pearcey, G.E.P., 2020. Low-threshold motor units can be a pain during
experimental muscle pain. J. Physiol. 598, 2545–2547. https://doi.org/10.1113/JP279872

Milligan, E.D., Mehmert, K.K., Hinde, J.L., Harvey, L.O., Martin, D., Tracey, K.J., Maier, S.F., Watkins, L.R.,
2000. Thermal hyperalgesia and mechanical allodynia produced by intrathecal administration of
the human immunodeficiency virus-1 (HIV-1) envelope glycoprotein, gp120. Brain Res. 861, 105–
116. https://doi.org/10.1016/S0006-8993(00)02050-3

Morin-Richaud, C., Feldblum, S., Privat, A., 1998. Astrocytes and oligodendrocytes reactions after a total
section of the rat spinal cord. Brain Res. 783, 85–101. https://doi.org/10.1016/S0006-
8993(97)01282-1

Moy, J.K., Kuhn, J.L., Szabo-Pardi, T.A., Pradhan, G., Price, T.J., 2018. eIF4E phosphorylation regulates

ongoing pain, independently of inflammation, and hyperalgesic priming in the mouse CFA model.
Neurobiol. Pain 4, 45–50. https://doi.org/10.1016/j.ynpai.2018.03.001

Murphy, R.M., Zemlan, F.P., 1990. Selective serotonin1A/1B agonists differentially affect spinal
nociceptive reflexes. Neuropharmacology 29, 463–468. https://doi.org/10.1016/0028-
3908(90)90168-Q

Nash, B., Ioannidou, K., Barnett, S.C., 2011. Astrocyte phenotypes and their relationship to myelination.
J. Anat. 219, 44–52. https://doi.org/10.1111/j.1469-7580.2010.01330.x

Nesic, O., Lee, J., Ye, Z., Unabia, G.C., Rafati, D., Hulsebosch, C.E., Perez-Polo, J.R., 2006. Acute and
chronic changes in aquaporin 4 expression after spinal cord injury. Neuroscience 143, 779–792.
https://doi.org/10.1016/j.neuroscience.2006.08.079

Nitzan-Luques, A., Devor, M., Tal, M., 2011. Genotype-selective phenotypic switch in primary afferent
neurons contributes to neuropathic pain. Pain 152, 2413–2426.
https://doi.org/10.1016/j.pain.2011.07.012

North, R.Y., Li, Y., Ray, P., Rhines, L.D., Tatsui, C.E., Rao, G., Johansson, C.A., Zhang, H., Kim, Y.H., Zhang,
B., Dussor, G., Kim, T.H., Price, T.J., Dougherty, P.M., 2019. Electrophysiological and transcriptomic
correlates of neuropathic pain in human dorsal root ganglion neurons. Brain 142, 1215–1226.
https://doi.org/10.1093/brain/awz063

Novak, I., Hines, M., Goldsmith, S., Barclay, R., 2012. Clinical Prognostic Messages From a Systematic
Review on Cerebral Palsy. Pediatrics 130, e1285–e1312. https://doi.org/10.1542/peds.2012-0924

Olson, W., Luo, W., 2018. Somatotopic organization of central arbors from nociceptive afferents
develops independently of their intact peripheral target innervation. J. Comp. Neurol. 526, 3058–
3065. https://doi.org/10.1002/cne.24533

Ondarza, A.B., Ye, Z., Hulsebosch, C.E., 2003. Direct evidence of primary afferent sprouting in distant
segments following spinal cord injury in the rat: Colocalization of GAP-43 and CGRP. Exp. Neurol.
184, 373–380. https://doi.org/10.1016/j.expneurol.2003.07.002

Oskoui, M., Coutinho, F., Dykeman, J., Jetté, N., Pringsheim, T., 2013. An update on the prevalence of
cerebral palsy: A systematic review and meta-analysis. Dev. Med. Child Neurol. 55, 509–519.
https://doi.org/10.1111/dmcn.12080

Pinho, R.H., Leach, M.C., Minto, B.W., Del Lama Rocha, F., Luna, S.P.L., 2020. Postoperative pain
behaviours in rabbits following orthopaedic surgery and effect of observer presence. PLoS One 15,
1–21. https://doi.org/10.1371/journal.pone.0240605

Pitcher, G.M., Henry, J.L., 2008. Governing role of primary afferent drive in increased excitation of spinal
nociceptive neurons in a model of sciatic neuropathy. Exp. Neurol. 214, 219–228.
https://doi.org/10.1016/j.expneurol.2008.08.003

Price, T.J., Dussor, G., 2014. Evolution: The advantage of "maladaptive" pain plasticity. Curr. Biol. 24,
R384–R386. https://doi.org/10.1016/j.cub.2014.04.011

Ramer, L.M., Peter van Stolk, A., Inskip, J.A., Ramer, M.S., Krassioukov, A. V., 2012. Plasticity of TRPV1-
expressing sensory neurons mediating autonomic dysreflexia following spinal cord injury. Front.
Physiol. 3, 1–16. https://doi.org/10.3389/fphys.2012.00257

Riquelme, I., Montoya, P., 2010. Developmental changes in somatosensory processing in cerebral palsy
and healthy individuals. Clin. Neurophysiol. 121, 1314–1320.
https://doi.org/10.1016/j.clinph.2010.03.010

Rocha-Ferreira, E., Hristova, M., 2016. Plasticity in the neonatal brain following hypoxic-ischaemic injury.
Neural Plast. 2016. https://doi.org/10.1155/2016/4901014

Rodin, B., Sampogna, S.L., Kruger, L.W., 1983. An examination of intraspinal sprouting in dorsal root
axons with the tracer horseradish peroxidase. J. Comp. Neurol. 215, 187–198.
https://doi.org/10.1002/cne.902150206

Ruda, M.A., Ling, Q.D., Hohmann, A.G., Peng, Y.B., Tachibana, T., 2000. Altered nociceptive neuronal
circuits after neonatal peripheral inflammation. Science 289, 628–630.
https://doi.org/10.1126/science.289.5479.628

Rudomin, P., Schmidt, R., 1999. Presynaptic inhibition in the vertebrate spinale chord revisited. Exp.
brain Res. 129, 1–37.

Schomburg, E.D., Steffens, H., 1998. Comparative analysis of L-DOPA actions on nociceptive and non-
nociceptive spinal reflex pathways in the cat. Neurosci. Res. 31, 307–316.
https://doi.org/10.1016/S0168-0102(98)00058-3

Sienko, S.E., 2018. An exploratory study investigating the multidimensional factors impacting the health
and well-being of young adults with cerebral palsy. Disabil. Rehabil. 40, 660–666.
https://doi.org/10.1080/09638288.2016.1274340

Sladky, J.T., Rorke, L.B., 1986. Perinatal hypoxic/ischemic spinal cord injury. Fetal Pediatr. Pathol. 6, 87–
101. https://doi.org/10.3109/15513818609025927

Steele, P.R., Cavarsan, C.F., Dowaliby, L., Westefeld, M., Katenka, N., Drobyshevsky, A., Gorassini, M.A.,
Quinlan, K.A., 2020. Altered Motoneuron Properties Contribute to Motor Deficits in a Rabbit
Hypoxia-Ischemia Model of Cerebral Palsy. Front. Cell. Neurosci. 14, 1–12.
https://doi.org/10.3389/fncel.2020.00069

Synowiec, S., Lu, J., Yu, L., Goussakov, I., Lieber, R., Drobyshevsky, A., 2019. Spinal Hyper-Excitability and
Altered Muscle Structure Contribute to Muscle Hypertonia in Newborns After Antenatal Hypoxia-
Ischemia in a Rabbit Cerebral Palsy Model. Front. Neurol. 9, 1–18.
https://doi.org/10.3389/fneur.2018.01183

Tan, A.M., Chakrabarty, S., Kimura, H., Martin, J.H., 2012. Selective corticospinal tract injury in the rat
induces primary afferent fiber sprouting in the spinal cord and hyperreflexia. J. Neurosci. 32,
12896–12908. https://doi.org/10.1523/JNEUROSCI.6451-11.2012

Vega-Avelaira, D., Moss, A., Fitzgerald, M., 2007. Age-related changes in the spinal cord microglial and
astrocytic response profile to nerve injury. Brain. Behav. Immun. 21, 617–623.
https://doi.org/10.1016/j.bbi.2006.10.007

Walker, S.M., Melbourne, A., O'Reilly, H., Beckmann, J., Eaton-Rosen, Z., Ourselin, S., Marlow, N., 2018.
Somatosensory function and pain in extremely preterm young adults from the UK EPICure cohort:
sex-dependent differences and impact of neonatal surgery. Br. J. Anaesth. 121, 623–635.
https://doi.org/10.1016/j.bja.2018.03.035

Walters, E.T., 2012. Nociceptors as chronic drivers of pain and hyperreflexia after spinal cord injury: An

772   adaptive-maladaptive hyperfunctional state hypothesis. Front. Physiol. 3, 1–13.
773   https://doi.org/10.3389/fphys.2012.00309

774 Weaver, L.C., Verghese, P., Bruce, J.C., Fehlings, M.G., Krenz, N.R., Marsh, D.R., 2001. Autonomic
775   dysreflexia and primary afferent sprouting after clip-compression injury of the rat spinal cord. J.
776   Neurotrauma 18, 1107–1119. https://doi.org/10.1089/08977150152693782

777 Westbom, L., Rimstedt, A., Nordmark, E., 2017. Assessments of pain in children and adolescents with
778   cerebral palsy: a retrospective population-based registry study. Dev. Med. Child Neurol. 59, 858–
779   863. https://doi.org/10.1111/dmcn.13459

780 Woolf, C.J., Shortland, P., Coggeshall, R.E., 1992. Peripheral nerve injury triggers central sprouting of
781   myelinated afferents. Nature 355, 75–78. https://doi.org/10.1038/355075a0

782 Xie, W., Strong, J.A., Zhang, J.M., 2009. Early blockade of injured primary sensory afferents reduces glial
783   cell activation in two rat neuropathic pain models. Neuroscience 160, 847–857.
784   https://doi.org/10.1016/j.neuroscience.2009.03.016

785 Xu, Q., Ford, N.C., He, S., Huang, Q., Anderson, M., Chen, Z., Yang, F., Crawford, L.K., Caterina, M.J.,
786   Guan, Y., Dong, X., 2021. Astrocytes contribute to pain gating in the spinal cord. Sci. Adv. 7, 1–13.

787 Yoshioka, B.Y.M., Coutts, R.D., Amiel, D., Hacker, S.A., 1996. Characterization of a model of osteoarthritis
788   in the rabbit knee. Osteoarthr. Cartil. 4, 87–98.

789 Zador, Z., Stiver, S., Wang, V., Manley, G.T., 2009. Role of Aquaporin-4 in Cerebral Edema and Stroke.
790   Handb. Exp. Pharmacol. 190, 159–170. https://doi.org/10.1007/978-3-540-79885-9_7

791 Zając, A., Chalimoniuk, M., Gołaš, A., Lngfort, J., Maszczyk, A., 2015. Central and peripheral fatigue
792   during resistance exercise - A critical review. J. Hum. Kinet. 49, 159–169.
793   https://doi.org/10.1515/hukin-2015-0118

794 Zarkou, A., Lee, S.C.K., Prosser, L.A., Jeka, J.J., 2020. Foot and Ankle Somatosensory Deficits Affect
795   Balance and Motor Function in Children With Cerebral Palsy. Front. Hum. Neurosci. 14, 1–12.
796   https://doi.org/10.3389/fnhum.2020.00045

797 Zhang, H., Boyette-Davis, J.A., Kosturakis, A.K., Li, Y., Yoon, S.-Y., Walters, E.T., Dougherty, P.M., 2013.
798   Induction of monocyte chemoattractant protein-1 (MCP-1) and its receptor CCR2 in primary
799   sensory neurons contributes to paclitaxel-induced peripheral neuropathy. J. Pain 14, 1031–1044.
800   https://doi.org/10.1016/j.jpain.2013.03.012.Induction

801 Zhang, H., Li, Y., De Carvalho-Barbosa, M., Kavelaars, A., Heijnen, C.J., Albrecht, P.J., Dougherty, P.M.,
802   2016. Dorsal Root Ganglion Infiltration by Macrophages Contributes to Paclitaxel Chemotherapy-
803   Induced Peripheral Neuropathy. J. Pain 17, 775–786. https://doi.org/10.1016/j.jpain.2016.02.011

804 Zhang, J., Xiang, Q.S., Echeverry, S., Mogil, J.S., De Koninck, Y., Rivest, S., 2007. Expression of CCR2 in
805   both resident and bone marrow-derived microglia plays a critical role in neuropathic pain. J.
806   Neurosci. 27, 12396–12406. https://doi.org/10.1523/JNEUROSCI.3016-07.2007

807