Enhanced nociceptive behavior and expansion of associated primary afferents in a rabbit model of cerebral palsy

Authors

E. J. REEDICH[1,2], L. T. GENRY[1,2,3], M. A. SINGER[4], C. F. CAVARSAN[1,2], E. MENA AVILA[1,2], D. M. BOUDREAU[1,2], M. C. BRENNAN[1,2], A. M. GARRETT [1,2,3], L. DOWALIBY[1,2], M. R. DETLOFF[4*], K. A. QUINLAN[1,2*]

1 George and Anne Ryan Institute for Neuroscience, University of Rhode Island, Kingston, RI, USA

2 Department of Biomedical and Pharmaceutical Sciences, College of Pharmacy, University of Rhode Island, Kingston, RI, USA

3 Interdisciplinary Neuroscience Program, University of Rhode Island, Kingston, RI, USA

4 Department of Neurobiology and Anatomy, Marion Murray Spinal Cord Research Center, College of Medicine, Drexel University, Philadelphia, PA, USA

*Corresponding Authors:

Katharina A. Quinlan

George and Anne Ryan Institute for Neuroscience

Department of Biomedical and Pharmaceutical Sciences

College of Pharmacy

University of Rhode Island

130 Flagg Rd

Kingston, RI 02881

kaquinlan@uri.edu

Megan R. Detloff

Department of Neurobiology & Anatomy

Marion Murray Spinal Cord Research Center

College of Medicine

Drexel University

2900 W. Queen Lane

Philadelphia, PA 19129

mrd64@drexel.edu

Declaration of interest: None

Abstract

Spastic cerebral palsy (CP) is a movement disorder marked by hypertonia and hyperreflexia, and the most prevalent comorbidity is pain. Since spinal nociceptive afferents contribute to both the sensation of painful stimuli as well as reflex circuits involved in movement, we investigated the relationship between prenatal hypoxia-ischemia (HI) injury which can cause CP, and possible changes in spinal nociceptive circuitry. To do this, we examined nociceptive afferents and mechanical and thermal sensitivity of New Zealand White rabbit kits after prenatal HI or a sham surgical procedure. As described previously, a range of motor deficits similar to spastic CP was observed in kits born naturally after HI (40 minutes at ~70-80% gestation). We found that HI caused an expansion of peptidergic afferents (marked by expression of calcitonin gene-related peptide; CGRP) in both the superficial and deep dorsal horn at postnatal day (P)5. Non-peptidergic nociceptive afferent arborization (labeled by isolectin B4; IB4) was unaltered in HI kits but overlap of the two populations (peptidergic and non-peptidergic nociceptors) was increased by HI. Density of glial fibrillary acidic protein (GFAP) was unchanged within spinal white matter regions important in nociceptive transmission at P5. We found that mechanical and thermal nociception was enhanced in HI kits even in the absence of motor deficits. These findings suggest that prenatal HI injury impacts spinal sensory pathways in addition to the more well-established disruptions to descending motor circuits. In conclusion, changes to spinal nociceptive circuitry could disrupt spinal reflexes and contribute to pain experienced by individuals with CP.

Significance Statement:

Perinatal injuries that cause cerebral palsy (CP) typically involve global insults to the central nervous system and are capable of modulating development of both motor and sensory systems. Most individuals with CP experience pain, yet whether nociception is enhanced in this disorder is unexplored. Here, we demonstrate altered topographic distribution of nociceptive afferents in the spinal cord dorsal horn of neonatal rabbits that experienced hypoxic-ischemic (HI) injury *in utero*; these anatomical changes were associated with nocifensive behavior indicative of pain-like behaviors. Our findings suggest that CP-causative injuries alter spinal sensory pathways (not only descending motor circuits), contributing to increased pain in CP.

Key words: cerebral palsy, hypoxia-ischemia, sensation, nociception, rabbit, primary afferent

Abbreviations: Serotonin (5HT); calcitonin gene-related peptide (CGRP); central nervous system (CNS); cerebral palsy (CP); Diamidino-2-Phenylindole (DAPI); dorsal column (DC); dorsal root ganglion (DRG); glial fibrillary acidic protein (GFAP); Gross Motor Function Classification System (GMFCS); hypoxia-ischemia (HI); isolectin B4 (IB4); immunohistochemistry (IHC); lateral spinothalamic tract (LST); normal horse serum (NHS); nonsteroidal anti-inflammatory drugs (NSAIDs); postnatal day (P); phosphate buffered saline (PBS); paraformaldehyde (PFA); regions of interest (ROIs); superficial dorsal gray matter (SDGM); ventral spinothalamic tract (VST)

Introduction:

Cerebral palsy (CP) is the most common motor disability in children, affecting 1:500 live births (Oskoui et al., 2013). Chronic pain is the most prevalent co-morbidity of CP (McDowell et al., 2017; Novak et al., 2012): up to 77% of children with CP experience pain (McKinnon et al., 2020; Mckinnon et al., 2018). Individuals with CP experience chronic pain at significantly higher rates than the rest of the population (Badia et al., 2014; Doralp and Bartlett, 2010; García Jalón et al., 2021; Westbom et al., 2017). Pain decreases quality of life (Findlay et al., 2016), yet the pathophysiology of pain is rarely researched in the context of CP. This is likely due to the pervasive view that pain in individuals with CP is a symptom arising from secondary effects of physical stressors, including spasticity, hip subluxation, and joint contractures. Individuals with severe CP (as measured by Gross Motor Function Classification System (GMFCS) V) do experience more pain (García Jalón et al., 2021; Mckinnon et al., 2018); however, CP-affected individuals across all GMFCS levels report pain (Flanigan et al., 2020; Sienko, 2018). Even individuals born very prematurely (<26 weeks gestion), who do not develop CP, report recurrent pain and altered thermal sensitivity as adults (Walker et al., 2018). Thus, we hypothesized that the same trauma that can impact developing corticospinal tracts in CP could impact development of nociceptive circuitry in the spinal cord.

There is a wealth of evidence regarding the interrelatedness of sensory and motor performance (Cooper et al., 1995; Gupta et al., 2017; Matusz et al., 2018; Zarkou et al., 2020). Somatosensory deficits are present in CP, and their presence could contribute to motor dysfunction. Typically, primary afferent input onto spinal circuitry is a key mediator of functional motor performance. Proprioceptive and tactile afferent feedback informs the central nervous system (CNS) of motor command outcomes and sculpts ongoing drive to motoneurons. In particular, the role of primary nociceptive afferent fibers in motor function is critically overlooked—especially after CNS damage. Importantly, there are a few exceptions, including the restriction of motor unit activation in response to excessive stretch, detection of dangerous pH and other metabolite levels during muscle fatigue, and the exercise pressor reflex and motor unit modulation during exercise-induced fatigue (Laurin et al., 2015; Zając et al., 2015). Cutaneous nociceptors (mechanosensitive C, thermosensitive C and Aδ neurons) are small, pseudounipolar neurons in the dorsal root ganglion (DRG) that are analogous to Type III (Aδ) and Type IV (C) primary afferent fibers from the muscle, though their axons terminate in different dorsal horn laminae (Kandel et al., 2013; Lallemend and Ernfors, 2012; Laurin et al., 2015). Importantly, nociceptive and pain information can be relayed from muscle and joints via the nociceptive primary afferent reflex arcs, modulating motor output by generating a withdrawal reflex (Kandel et al., 2013; Kimura et al., 2004). The interconnection between motor and sensory systems can make it challenging to parse out their relative contribution to functional impairments.

As in other CNS disorders, there is a strong correlation between the integrity of the somatosensory system (including primary afferent input) and motor performance in CP (Gupta et al., 2017; Zarkou et al., 2020). Tractography and physiology experiments in individuals with unilateral CP revealed that integrity of somatosensory connections has a relatively larger impact on hand function compared to the integrity of motor connections (Gupta et al., 2017), and stimulation of peripheral nerves, which provides additional sensory feedback, improved motor performance in individuals with diplegic CP (Bumin and Kayihan, 2001; Kaelin-Lang, 2008).

Physical insults during pregnancy have been linked to later development of autism, schizophrenia, and the motor impairments that are hallmarks of CP (Marriott et al., 2017), but it is not

known if these prenatal events impact nociceptive afferents. Neural development can be negatively
impacted by relatively common conditions in the perinatal period, including maternal infection and
inflammation, placental insufficiency, and difficult birth. These conditions can all result in varying
degrees of hypoxia-ischemia (HI) and injury to the developing CNS. Effects of prenatal HI include white
matter injury, cortical damage and motor deficits (Buser et al., 2010; Derrick et al., 2004; Drobyshevsky
and Quinlan, 2017; Rocha-Ferreira and Hristova, 2016), but the effect on nociceptors is completely
unstudied. Notably, cell death in the deep dorsal horn (where synaptic targets of nociceptors are
located) is present after HI in rabbits and in cases of infant death after hypoxia (Clancy et al., 1989;
Drobyshevsky and Quinlan, 2017; Sladky and Rorke, 1986). Dorsal root ganglia (DRG) neurons undergo
perinatal growth and refinement of projections during the period of susceptibility to injuries causing CP
(Marmigère and Ernfors, 2007; Olson and Luo, 2018), though afferent plasticity is retained in adulthood
(Jiang et al., 2016; Tan et al., 2012). Changes to nociceptive DRG neurons are entirely unexplored in the
context of CP.

In this study, we have chosen to focus on nociceptors at the spinal level using the rabbit model
of CP. Investigating the relationship between motor and sensory dysfunction in CP requires using an
animal model that displays both phenotypes of motor dysfunction and enhanced nociception.
Unfortunately, rodents, the most commonly used models, can sustain very severe brain injuries yet
display mild or nonexistent motor deficits (Fan et al., 2005; Rice et al., 1981)HI kits show pronounced
motor dysfunction (Cavarsan et al., 2019; Derrick et al., 2007, 2004; Shi et al., 2021). Furthermore,
emerging evidence suggests that nociceptors may link afferents to hyperreflexia. Nociceptors impact
movement by modulating spinal reflex circuits (Black, 2019; Lucas-Osma et al., 2019), and nociceptor
input to reflex circuits is a feasible potential contributor to disrupted reflex responses (Hari et al., 2021)
particularly in those with CP who may suffer from chronic pain. Monosynaptic stretch reflexes can be
potently suppressed via nociceptor inhibition (Black, 2019; Lucas-Osma et al., 2019). In the case of
people with CP, rather than suppression of the monosynaptic stretch reflexes, over active nociceptors
may drive hyperreflexia and reflex irradiation (Achache et al., 2010; Leonard et al., 1991; Leonard and
Hirschfeld, 1995). Whether this mechanism is contributing to spasticity and hyperreflexia in CP is the
focus of ongoing study. Thus, not only could expanded distribution of nociceptive afferents contribute to
the perception of pain by higher brain centers, but also to hyperreflexia and disruptions in motor
coordination that are the hallmarks of CP.

Thus, we examined nociceptors and altered nociception at the level of the spinal cord in rabbit
kits that were subjected to HI injury *in utero* (40 minutes at ~70-80% gestation) using both
immunohistochemistry to quantify molecular markers for nociceptors and *in vivo* behavioral testing at
an age roughly corresponding to human toddlers (postnatal day [P] 5). We found that rabbit kits
exposed to prenatal HI demonstrated increased nociceptive behavior in response to cutaneous
mechanical and thermal stimuli. This increased nociceptive response corresponded to aberrant
distribution and density of nociceptive primary afferent fibers but not astrogliosis at this age.
Interestingly, our data show that enhanced forepaw and hindpaw sensitivity occurs independent of
motor deficits produced by this model. Together, these results indicate that sensory circuits are
vulnerable to perturbation by perinatal HI injury, and suggest that neuropathic pain could accompany
motor deficits characteristic of CP.

Materials and Methods:

Animal model: Experiments were performed in accordance with the ARRIVE guidelines and the United States National Institutes of Health's Guide for Care and Use of Laboratory Animals. Approval of the University of Rhode Island's Institutional Animal Care and Use Committee was obtained for all experiments performed in this study. 66 kits of both sexes from 15 different litters originating from 9 dams were used for this study; some dams provided more than one litter of rabbits. The inherent variability in behavioral data warranted large sample sizes. All efforts were made to minimize animal suffering and to reduce the number of animals used.

Hypoxia-ischemia surgery: Pregnant New Zealand White rabbits (bred in house or ordered from Charles River Laboratories, Inc., Wilmington MA), underwent HI procedures as previously described (Derrick et al., 2004). Briefly, at ~70-80% gestation (day 22 -26 of 31.5 days typical gestation period), dams were premedicated with ketamine and xylazine (25 mg/kg and 3.75 mg/kg, respectively), anesthetized with isoflurane (1-3%; inhaled via V gel tube) and occasionally treated with buprenorphine (0.01 - 0.24 mg/kg, as needed). Indications for treatment with buprenorphine include elevated heart rate (over 220BPM); prolonged rapid, shallow breathing / panting; elevated CO_2 levels (due to panting). The left or right femoral artery was isolated. A Fogarty balloon catheter (2-4 French) was inserted and advanced to the level of the bifurcation of the descending aorta, above the uterine arteries, and inflated to completely occlude blood flow for 40 minutes to cause HI in the fetuses. Placement of the catheter and confirmation that blood flow was occluded was performed with ultrasound. Dams subjected to sham surgical procedures underwent the same anesthesia protocols as the HI dams but without insertion and/or inflation of the catheter.

After the procedures, dams recovered and later gave birth to kits on the expected due date +/- two days. Sham and HI litters included in this study were all born at term (>92% of gestation). Neonatal kits of both sexes were characterized for motor deficits, nociceptive behavior, and used for anatomical investigations as described below. Since we examined behavior in neonatal animals far from sexual maturity (NZW rabbits typically reach sexual maturity at ~6 months of age), and whose external genitalia were not yet visible, we grouped together male and female kits.

Behavior: There is a wide range in motor phenotypes after HI, with some kits showing motor deficits while others appear unaffected. Categorization of motor deficits was performed using a modified Ashworth scale, observations/tests for activity, locomotion, posture, righting reflex, and muscle tone as previously described (Derrick et al., 2004). Based on these tests, kits were labeled as "motor-affected HI" or "motor-unaffected HI." Motor-affected HI kits had modified Ashworth muscle tone scores of ≥3 in one or more evaluated joints; Ashworth scores did not exceed 2 in any joints of motor-unaffected HI kits.

von Frey test for mechanical nociception: Neonatal rabbit kits of both sexes were tested for mechanical hypersensitivity using von Frey filaments. Animals were allowed to acclimatize in a chamber on an elevated gridded (8 mm x 8 mm) surface. von Frey filaments (Touch Test® Sensory Evaluators; range: 1-26 g) were applied in ascending order to the plantar surface of the forepaws and hindpaws, just proximal to the digits of the kit. See inset in Figure 1 which depicts location of stimulus application. If paw withdrawal occurred in response to application of the 1 g von Frey filament, filaments (0.6-0.008 g) were applied in descending order until no response was observed. Each filament was tested at least five consecutive times per paw to ensure the observed reaction was in response to the filament and not general movement of the animal. The force (in g) of the filament that elicited a consistent withdrawal

response was recorded as the paw withdrawal threshold. If no response was elicited by application of
the 26 g von Frey filament, that paw was recorded as non-responsive and given a score of 60 g (which is
the subsequent filament). Filament forces eliciting withdrawal of left and right forepaws and hindpaws
were entered independently for statistical analyses, since motor deficits are frequently disparate across
limbs and responsiveness to stimuli was therefore not assumed to be equal across limbs.

Hargreaves' test for thermal nociception: Thermal sensitivity was tested using a Hargreaves apparatus.
Specifically, a model 336G Plantar/Tail Stimulator Analgesia Meter (IITC Life Sciences, Woodland Hills,
CA) was used to apply a beam of radiant heat (4x6 mm area; 80% intensity; 8 s max duration) to the
plantar surface of each hindpaw and forepaw. Rabbit kits were allowed to acclimatize in a chamber on a
glass surface, and then the skin on the plantar surface of each paw proximal to the digits was tested
three times with at least 30 seconds between applications. See insets in Figure 2 for schematic depicting
the location of stimulus application to the forepaws and hindpaws. The beam of radiant heat increased
temperature gradually from rest (20-30 °C) to a maximum of 83 °C at 8 s. Latency to paw withdrawal
was recorded; if the animal did not withdraw its paw, that trial was recorded as a non-response and
given a latency of 9 s. For each paw, the stimulus was applied 3 times, and the withdrawal latencies
were averaged to generate a single score per paw. Since weight distribution and function are different
between forelimbs and hindpaws, withdrawal latencies from the forepaws were examined separately
from hindpaws. Each group contains individual data points generated from the left or right paws.

Immunohistochemistry (IHC): Kits were deeply anesthetized by intraperitoneal injection of
ketamine/xylazine (100 mg/kg and 20 mg/kg, respectively) or sodium pentobarbital before tissue
harvest at P5. Transcardial perfusion was performed using chilled 0.01 M phosphate buffered saline
(PBS; pH 7.4) until tissue blanching was observed, then 4% paraformaldehyde (PFA) (in PBS) was infused
at a constant flow rate of approximately 5-8 mL/min (depending on rabbit kit size). Perfused spinal cords
were dissected and post-fixed in 4% paraformaldehyde for ~24-72 hrs. Next, fixed spinal cords were
washed with PBS and cryoprotected in 30% sucrose (in PBS) until sinking was observed. Cryoprotected
spinal cords were embedded in optimal cutting temperature (OCT) compound and frozen at −80 °C until
cryostat sectioning (Leica Lm 1850, Germany). Transverse lumbar spinal cord cryosections were
collected in serial at 30-μm or 40-μm thickness; to avoid repeat regions of analysis, every 15th section
was used for analysis. Manufacturer Information, catalog number and lot information for all
immunohistochemical antibodies and lectins used are provided in Table 1.

IHC for glial fibrillary acidic protein (GFAP): Slides were thawed, dried in an oven at 37 °C to improve
section adherence, then rehydrated in PBS. Sections were blocked in 0.2% Triton X-100 and 5% normal
horse serum (NHS) in PBS for 30 minutes at room temperature. Following blocking, slices were
incubated in primary antibody solution containing mouse-anti GFAP antibody (diluted 1:500 in 0.1%
Triton X-100 and 1% NHS in PBS;) overnight at room temperature. Sections were washed in PBS and
then were incubated in secondary antibody solution consisting of Atto 488-goat anti-mouse IgG (whole
molecule) (diluted 1:500 in 0.1% Triton X-100 and 1% NHS in PBS,) for 2 h at room temperature. Nuclei
were labeled with 4',6-Diamidino-2-Phenylindole (DAPI) and slides were mounted using Fluoromount
mounting medium (SouthernBioTech). Z-stack photomicrographs of dorsal and ventral lower lumbar
spinal cord were acquired at 10x magnification using a Nikon Eclipse Ti2 inverted confocal microscope.
Maximum intensity projections of the green channel were generated and stitched (if possible) using
Image Composite Editor (Microsoft). Grayscale images were imported into ImageJ and blinded
densitometric analysis of GFAP immunofluorescent signal (mean gray value) in regions of interest

containing dorsal columns, superficial dorsal gray matter, lateral spinothalamic tract, and ventral
spinothalamic tract (divided by background fluorescent signal) was performed using Image J (NIH).
Background fluorescent signal was the mean gray value of adjacent unlabeled gray matter. The grey or
white matter in our regions of interest was targeted to encompass the entire region of interest but
could also contain portions of other areas.

IHC for primary afferent fibers: Sections were rinsed in PBS, blocked in 10% normal goat serum and 0.2%
Triton-X100 in PBS for 1 hour at room temperature followed by a 30-minute incubation with rabbit-to-
rabbit blocking reagent (ScyTek Laboratories, Inc., catalog #: RTR008). Sections were then incubated
with a solution of biotinylated IB4 (1:2000, Sigma) and anti-calcitonin gene-regulated peptide in 0.1%
Triton X-100 in PBS at room temperature for 48 h. After three washes with PBS, sections were incubated
in avidin-conjugated with Alexa Fluor 594 and goat-anti-rabbit Alexa Fluor 488 in 5% goat serum in PBS
for 2 hours at room temperature. Following three washes, sections were rinsed in PBS and cover slipped
with FluorSave Reagent (Calbiochem, catalog #: 345789, Bedford, MA). Three images of the left and
right lumbar dorsal horn were taken for each rabbit at
the same magnification, lens aperture and exposure
time on a Leica DM5500B epifluorescent microscope
and captured with Slidebook software. Images were
averaged to calculate values per animal. The
proportional area of IB4, CGRP and colocalized
immunoreactive tissue within the cap of the dorsal horn
(laminae I-III) was quantified in ImageJ as previously
described (Detloff et al., 2016). The proportional area of
CGRP+ fibers that expand in dorsal horn ventral to the
IB4+ band was measured in the same sections.

Statistical Analyses: All statistical analyses were
performed using GraphPad Prism (GraphPad
Software Inc.; San Diego, CA, USA). For each
dataset, normality of distributions was tested
using D'Agostino and Pearson tests. For normally
distributed datasets with equal variances,
unpaired t-tests were used to identify statistically
significant differences between groups. For

Figure 1: HI kits exhibit pain-like behavior in response to
mechanical stimuli . **A:** Forepaw (left) and hindpaw (right)
von Frey scores at P5. Sham-operated controls (blue) have
higher scores, indicating less mechanosensitivity, while HI
kits (red) display greater mechanosensitivity (lower scores).
B: HI kits from (A) were separated by whether they show
motor deficits (motor-affected) or no deficits (motor-
unaffected). There was no difference in forepaw withdrawal
threshold (left) but hindpaw withdrawal (right) scores were
higher in motor-affected HI kits, potentially indicating less
sensitivity or impaired ability to withdraw. Data are
represented as mean ± SEM; N=64 sham and N=60 HI paws
from N=32 sham and N=30 HI rabbits. **P<0.01, ****P
<0.0001; Mann-Whitney tests.

272 datasets in which one or more distributions was
273 non-normally distributed, Mann-Whitney tests
274 were used.

Results:

In this study, we investigated the link between
prenatal HI injury and nociception in a rabbit
model of CP. Rabbit kits exposed to prenatal HI
and age-matched sham animals underwent testing
for enhanced nociceptive reflexes in response to mechanical and thermal stimuli at P5 using von Frey
and Hargreaves' testing, respectively. After prenatal HI injury, neonatal rabbit kits displayed increased
nociceptive behavior in response to mechanical and thermal stimuli that manifested even in the absence
of motor deficits. HI kits demonstrated significantly greater sensitivity to von Frey filaments applied to
the paws compared to sham kits, withdrawing paws in response to application of lower forces
(forepaws, *P*<0.0001; hindpaws, *P*<0.0001) (**Figure 1**). Hindpaw von Frey scores of motor-unaffected HI
kits (mean: 8.2 g;) were significantly lower than that of motor-affected HI kits (mean: 19.4 g; *P*=0.0087)
while forepaw withdrawal thresholds were equivalent (*P*=0.9908).

Similar to von Frey, the Hargreaves' test revealed paw hypersensitivity in HI kits. HI kits had reduced
paw withdrawal latencies compared to sham rabbits in response to targeted radiant heat (**Figure 2**),
indicating increased thermal sensitivity (forepaws: $P<0.0001$; hindpaws: $P<0.0001$). Paw withdrawal
latencies to noxious thermal stimulation were
comparable between motor-unaffected HI
(forepaw latency: 3.6 s; hindpaw latency: 3.0 s)
and motor-affected HI kits (forepaw latency: 4.4 s;
hindpaw latency: 3.8 s) ($P=0.0740$ and 0.1279,
respectively).

Figure 2: HI kits display pain-like behavior in response to thermal stimuli **A:** Forepaw (left) and hindpaw (right) withdrawal latencies during Hargreaves' testing at P5. Shorter latencies to paw withdrawal in response to radiant heat are observed in HI kits compared to sham animals, indicative of enhanced thermal sensitivity. **B:** HI kits from (A) were separated by whether they show motor deficits (motor-affected) or no deficits (motor-unaffected). There was no difference in forepaw or hindpaw withdrawal scores in motor-affected HI kits. Data are represented as mean ± SEM; N= 24 sham and N=36 HI paws from N=12 sham and N=18 HI rabbits. ****$P <0.0001$; unpaired t-tests.

We hypothesized that mechanical and thermal hypersensitivity in HI rabbit kits may be caused by
spinal astrogliosis and/or expansion of nociceptive afferents in the spinal cord dorsal horn. To determine
whether prenatal HI leads to long-lasting postnatal astrogliosis in gray and white matter spinal cord
regions involved in cutaneous sensation or nociception, the degree of glial fibrillary acidic protein (GFAP)
expression in four lumbar spinal cord regions of interest (ROIs) was quantified. These ROIs included
tracts/laminae involved in cutaneous sensation or nociceptive neurotransmission: the dorsal column,
superficial dorsal gray matter, ventral spinothalamic tract, and lateral spinothalamic tract. Densitometric
analysis revealed no significant differences in GFAP labeling in these ROIs at P5 (**Figure 3**). Baseline GFAP
immunoreactivity in the white matter of neonatal rabbits found here is consistent with the nervous
system expression profile established in mice (Yoon et al., 2017) and may reflect the role of astrocytes in

Figure 3: Astrogliosis is not present in HI rabbits at P5. **A**: Transverse lumbar spinal cord sections from sham (left) and HI (right) rabbits immunostained for GFAP (green). Scale bar applies to both images. **B:** Quantification of GFAP densitometry in spinal regions containing tracts/laminae of interest: dorsal column (DC), superficial dorsal gray matter (SDGM), ventral spinothalamic tract (VST), and lateral spinothalamic tract (LST). GFAP expression was unchanged in all ROIs in HI spinal cords (ns, not significant; unpaired t-tests). Data are represented as mean ± SEM; N=7 rabbits per group.

myelination during early postnatal development (Landry et al., 1990; Liedtke et al., 1996; Nash et al.,
2011). No evidence was found to support a contribution of neuroinflammation to increased nociception
at this age.

To determine whether prenatal HI affects primary afferent distribution and density within the dorsal
horn of the spinal cord, lumbar spinal cords (which include hindpaw dermatomes) from sham and HI kits
were labeled for CGRP and IB4 to identify peptidergic and non-peptidergic afferents, respectively. In
other species, small diameter fibers emanating from the dorsal root ganglia that contain CGRP and bind
IB4 transmit nociceptive information from the periphery to the superficial dorsal horn of the spinal cord
(REF). Similar to the rodent, distributions of these two nociceptor subpopulations had little overlap in
the dorsal horn of sham kits (**Figure 4 A-A''**), however, HI led to an increased overlap in the topographic
distribution of these two fiber types (**Figure 4 B-B''**). In addition, HI kits exhibited increased CGRP-
positive fibers in the deep dorsal horn (**Figure 4 C-C''**) that were not present in sham kits. Quantification
of the proportional area of each afferent subpopulation independently revealed that CGRP+ fibers
(**Figure 3D, G**) but not IB4+ (**Figure 4E**) fibers exhibited increased density in the superficial and deep
dorsal horn. Despite no increase in the density or distribution of IB4-positive fibers in the dorsal horn,
there was significant colocalization of CGRP and IB4 labeling in the dorsal horn of HI kits compared to
sham kits, suggesting that HI may induce plasticity or phenotype switching of these nociceptor
populations.

Figure 4 Increased density and overlap of nociceptive primary afferent fibers in HI kits at P5.
Representative images of the lumbar dorsal horn of sham (A-A'') and HI (B-B'') kits immunolabeled for
calcitonin gene-regulated peptide (CGRP; A, B, C) or labeled by isolectin B-4 (IB4; A', B', C'). Note the
increased presence of CGRP-positive fibers in the deep dorsal horn in the HI but not the sham spinal cord
(C-C''). The proportional area of CGRP-positive tissue was significantly increased in the superficial and
deep dorsal horn of HI rabbits compared to sham (D, G), while the density and distribution of IB4-positive
afferents was unchanged between groups (E). The degree of overlap of these two nociceptor populations
can be viewed in the merged images (A'', B'') and was significantly increased in HI kits compared to sham
kits (F). Data are represented as mean ± SEM; individual data points represent each animal (n=5 sham and
n=6 HI kits) from averages of 3 separate images of left and right dorsal horns per animal. **$P<0.01$,

Discussion:

Experimental animal models that mimic human disease with high fidelity are more likely to drive
meaningful discovery of therapeutic interventions to treat such diseases. Here, we use a preclinical
model that displays prominent motor deficits to show that nociceptive afferents are also disrupted, in
addition to the previously characterized damage to the cortex, thinning of the corticospinal tract, loss of
spinal motoneurons and altered motoneuron excitability (Buser et al., 2010; Drobyshevsky et al., 2007;
Drobyshevsky and Quinlan, 2017; Steele et al., 2020). Importantly, both rabbits after HI and human
infants after asphyxiation show cell death in the dorsal horn (Clancy et al., 1989; Drobyshevsky and
Quinlan, 2017; Sladky and Rorke, 1986). Moreover, this is one of few studies that utilize quantitative
assessments of nociception in the rabbit, despite the rabbit's extensive use in biomedical research (Lei
et al., 2017; Mans, 2020; Yoshioka et al., 1996). Our findings indicate that after prenatal HI, increased
topographic distribution of peptidergic nociceptive primary afferent fibers in the dorsal horn is present
along with a hypersensitive withdrawal response that is suggestive of allodynia.

There is substantial information regarding the formation of sensory connections during
development in the rodent, cat and primate (Beggs et al., 2002; Brown and Fuchs, 1975; Fitzgerald and
Jennings, 1999; Fitzgerald and Swett, 1983; Jennings and Fitzgerald, 1996; Ralston and Ralston, 1979;
Scheibel and Scheibel, 1968; Swett and Woolf, 1985; Zhang et al., 1997), however, there is a paucity of
information regarding primary afferent distribution in the developing or adult rabbit. This is the first
study to show that nociceptors are altered in an animal model of CP and one of the first to examine
primary afferent fiber distribution in the developing rabbit. The increased overlap and density of
peptidergic nociceptive afferent fibers in the dorsal horn of HI kits may indicate intrinsic changes within
nociceptive DRG neurons. In normal rodents, CGRP and IB4 are rarely co-expressed in primary sensory
neurons of the DRG; these markers normally indicate distinctive subpopulations of nociceptors
(Marmigère and Ernfors, 2007; Orozco et al., 2001). In chronic pathological situations, the distribution of
the axonal arbors within the dorsal horn exhibits significant overlap, and there is some indication that
the degree of coexpression of these two nociceptive markers increases (Detloff et al., 2016; Price and
Flores, 2007; Sliwinski et al., 2018). Despite significantly lower von Frey scores in HI kits indicating
increased mechanosensitivity, non-peptidergic IB4-positive mechanical nociceptors, which are activated
by noxious mechanical stimuli, were not found to have expanded afferent arborizations in the dorsal
horn. It could be that these IB4-positive nociceptors are hyperexcitable, and that altered activity, not
anatomy, contributes to pain-like behavior in response to mechanical stimuli in HI kits. In other rodent
models of nervous system injury, chronic pain and paw hypersensitivity is associated with dysfunctional
nociceptors that exhibit robust anatomical and electrophysiological plasticity (Bedi et al., 2010; Detloff
et al., 2016, 2014; Dougherty et al., 2004; Moy et al., 2018; Zhang et al., 2013); further, emerging
evidence suggests that nociceptive primary afferent input impedes motor output following CNS injury
(Keller et al., 2017, 2019, 2018). In a series of studies, Keller *et al.* demonstrated that nociceptor
depletion reduces muscle contractures and spasticity of the hindlimbs and improves overall locomotor
performance following spinal cord injury. Nociceptor silencing via activation of inhibitory $5HT_{1D}$
receptors has also been found to be helpful in reducing clonus and spasms after spinal cord injury
(Lucas-Osma et al., 2019). Taken together, these findings indicate that altered activity in nociceptors
contributes to motor deficits in cases of CNS injury. While the relationship with motor deficits is
particularly important in CP, the methods used in this study are unable to distinguish CGRP and/or IB4
positive primary afferent fibers emanating from the skin, muscle or viscera, and behavioral assessments

here only assess cutaneous afferent responses. Moreover, as dorsal root ganglia, where nociceptive cell
bodies reside, were not systematically harvested in this study, we cannot determine if the altered
topographic distribution of these afferents may represent a sprouting response or it may be indicative of
a change in the population of DRG neurons which express these nociceptive markers. Populations of
primary sensory neurons have been identified in many mammalian species from rodents to humans, and
size, molecular markers and functions are mostly conserved (Shiers et al., 2020). While isolated
anatomical reports suggest that the primary sensory neurons of rabbits may be anatomically similar to
other mammals (Kusunoki et al., 1992; Mori, 1986), a systematic evaluation of the anatomical and
physiological function of DRG neurons, especially nociceptors, in the developing rabbit is warranted.

In adults, physical trauma is known to alter neural circuits and have long lasting effects on
nociception (Costigan et al., 2009; Price and Dussor, 2014). The effect of neural injury on nociceptive
primary sensory neurons is a major contributor to the development of tactile and thermal
hypersensitivity and chronic pain (Hulsebosch and Coggeshall, 1981a, 1981b; Koerber et al., 1994;
Ramer et al., 2012; Rodin et al., 1983; Walters, 2012; Woolf et al., 1992). After trauma, it is well-
established that the afferent fibers of these nociceptors exhibit robust arborizations into the superficial
and deep dorsal horn of the spinal cord (Bedi et al., 2010; Detloff et al., 2016, 2014; Krenz and Weaver,
1998; Ondarza et al., 2003; Weaver et al., 2001). Thus, it is possible that these primary afferents make
inappropriate synaptic connections on dorsal horn neurons subserving non-nociceptive functions such
as low threshold mechanosensation or touch that may not occur during normal development (Abraira et
al., 2017; Abraira and Ginty, 2013; Boyle et al., 2019; Peirs et al., 2021). Piers et al (Peirs et al., 2021)
demonstrate that spinal cord circuitry that transmits nociceptive information is dependent upon the
nature or etiology of hypersensitivity or pain behaviors. Physiologically, nociceptors increase
spontaneous activity and become hyperexcitable following CNS trauma (Bedi et al., 2010; Burchiel et al.,
1985; Gold and Gebhart, 2010; Gracely et al., 1993; Li et al., 2018; North et al., 2019; Pitcher and Henry,
2008).

In neonates, effects of CNS trauma (like CP or brain injury) on nociception are less studied,
however there is a wealth of information on the effects peripheral nerve injuries and inflammation
(Fitzgerald, 1985; Himes and Tessler, 1989; Pattinson and Fitzgerald, 2004; Shortland and Fitzgerald,
1994; Torsney and Fitzgerald, 2002; Walker et al., 2003). Peripheral inflammation enhances collateral
sprouting of primary nociceptive afferents and enhances nociception in adulthood (Ruda et al., 2000).
We would be remiss if we did not also postulate that perinatal HI may also induce anatomical and
physiological plasticity of other subtypes of primary afferent fibers or supraspinal centers involved in
pain perception or pain modulation. Anoxia in the neonatal rat led to paw hypersensitivity and
significantly fewer cells in the ventroposterolateral nucleus of the thalamus and somatosensory and
insular cortex (Helou et al., 2021; Kumar et al., 2020, 2017).

Neuroinflammation, nociception, and pain are tightly intertwined (Grace et al., 2014).
Astrocytes and microglia are activated by neural injury and release a plethora of pro-inflammatory
molecules into the spinal milieu, and their activation corresponds to the development of neuropathic
pain (Chhaya et al., 2018; Detloff et al., 2008; Gao et al., 2009; Hulsebosch et al., 2009; Ishikura et al.,
2021; Lu et al., 2021; Xie et al., 2009; Zhang et al., 2016, 2007). After complete spinal cord injury,
increased GFAP persists for at least 3 weeks in the adult rat spinal cord (Choi et al., 2021; Ducza et al.,
2021; Morin-Richaud et al., 1998; Nesic et al., 2006; Xu et al., 2021). However, the time course of the

GFAP response to injury is different in neonates. Unlike adults, neonatal rat pups respond to spared
nerve injury with an early increase in GFAP (one day after injury), along with the same elevation after 7
days that is observed in adults (Vega-Avelaira et al., 2007). Density of GFAP fiber length and density of
GFAP+ cell bodies are increased in the lumbar gray matter of motor-affected (hypertonic) HI rabbit kits
at P1, seven days after HI (Drobyshevsky and Quinlan, 2017; Synowiec et al., 2019). The current study
was performed in HI kits that were 4 days older (P5) and examined specific laminae/tracts involved in
cutaneous sensation or nociception: (1) the dorsal column, the ascending pathway for cutaneous
sensation; (2) the superficial dorsal gray matter, containing second-order projection neurons relaying
nociception; (3) the ventral spinothalamic tract, carrying crude touch and pressure information; and (4)
the lateral spinothalamic tract, the ascending pathway for nociception. While astrogliosis is not present
in the spinal regions of interest we studied at P5 (11-14 days after HI), their reactivity closer to the time
of injury could be a causal link of enhanced nociception as has been demonstrated in other models
(Milligan et al., 2000). Astrocytes have been shown to exhibit different phenotypes or reactive states
following similar CNS injuries that can be either destructive or pro-reparative (Escartin et al., 2021;
Liddelow and Barres, 2017). Moreover, it has been reported that ischemia can induce an "A2-like"
astrocyte phenotype that upregulates thrombospondins and neurotrophic factors that promote survival
and repair of neurons (Gao et al., 2005; Hayakawa et al., 2014; Zador et al., 2009). Future studies could
examine more closely the time course and specific regions of elevated astrocyte activation in the rabbit
after prenatal HI and its relationship to enhanced nociception observed here at P5.

The stereotypical aversive paw withdrawal response to nocifensive tactile or thermal stimuli is
mediated by a segmental spinal reflex and is often inferred as a proxy for cortical perception of the
stimulus. It is unclear whether the results of the von Frey and Hargreaves' tests performed here reflect
hyperreflexia of nociceptive segmental spinal cord circuits (Advokat, 2002; Kauppila et al., 1998; Murphy
and Zemlan, 1990; Schomburg and Steffens, 1998) or the complicated perception of neuropathic pain. It
is likely these stimuli were offensive to the rabbits because paw withdrawal was typically accompanied
by paw licking and/or a rapid jump away from the stimuli. Unlike in rodent experiments, there are
surprisingly few reports measuring pain or nociception in experimental rabbit models, even those
modeling knee osteoarthritis or intervertebral disc dislocation (Lei et al., 2017; Mans, 2020; Yoshioka et
al., 1996). Composite pain scales for rabbits in a clinical environment (Banchi et al., 2020) and the Bristol
Rabbit Pain Scale (Benato et al., 2021) examine and score rabbit facial expressions and other behaviors
to assess pain, such as the grimace scale (Evangelista et al., 2021; Keating et al., 2012). Yet, recent
reports suggest that the experimenter's presence can have a profound effect on the animal's behavior
and subsequent score on these tests (Pinho et al., 2020). In the present study, unlike assessments of
spontaneous pain using the rabbit grimace scale, these assessments identify the dermatomal location
and relevant spinal cord or dorsal root ganglion level in which dysfunctional sensation occurs. This is
especially useful in examining changes in primary afferent input and/or dorsal horn processing in
specific spinal segments. Also, since our study examined an early postnatal age (P5) when the neonatal
rabbits' eyes are not yet open, the tests we performed were more appropriate to assess nociceptive
behavior than using the grimace scale, place-preference or aversion tests. However, future work could
include assessments of pain perception and an associated fear response at later postnatal ages to
further characterize the perception of pain and translate these findings clinically.
452
Individuals with CP have disruptions in somatosensation in addition to motor dysfunction.
Disruptions in somatosensory cortex connectivity (Hoon et al., 2002) and increased latency of sensory
evoked potentials (Cahan et al., 1987; Kundi et al., 1989) have been well documented in individuals with
CP. Innocuous tactile sensitivity actually decreases in individuals with CP while pain sensitivity is

enhanced (Riquelme and Montoya, 2010). The mechanism of this is not clear, but it is possible that this
is due to afferent switching after injury: in a rat model of neuropathic pain, large diameter
cutaneous/somatic Aβ afferents can undergo fiber-type switching after nerve ligation and begin
expressing CGRP, like nociceptive fibers (Nitzan-Luques et al., 2011). This switch in identity corresponds
to a period of tactile hypersensitivity. Another possibility is that the impaired CST creates a permissive
environment for afferent sprouting. When competition between CST and afferent input to the spinal
cord is lost after transection of the pyramidal tracts in adult rats, afferent sprouting into the deep dorsal
horn is observed (Jiang et al., 2016; Tan et al., 2012). Afferent sprouting after CST transection is
accompanied by loss of rate dependent depression of the H reflex, a hallmark of CP (Tan et al., 2012).
The common thread in both of these mechanisms is the remarkable plasticity of afferents, even in
adulthood. This raises the possibility that aberrant nociceptive input either by an increased anatomical
distribution and/or functional /physiological alterations arising during perinatal development could be
corrected later in life.
470
Movement and motor unit firing patterns in neurotypical adults are directly affected by pain
(Hodges and Tucker, 2011; Martinez-Valdes et al., 2020), perhaps through neuromodulators including
serotonin (5HT) (Mesquita et al., 2020). In our study, we found that HI kits were more sensitive to
mechanical and thermal stimuli than sham control kits. When comparing HI kits that were motor-
affected and those that were motor-unaffected, motor-affected HI kits were less sensitive to mechanical
stimuli to their hindpaws while their responses in forepaws and to thermal stimuli was unchanged. We
attribute this to the greater difficulty these kits had in making movements to respond to stimuli due to
their hypertonia. No kits in this study were completely incapable of movement and unable to react
physically to the stimuli, but their hypertonia could have impeded their reaction. Differences between
motor-affected and motor-unaffected HI kits and the relationship between nociceptive afferent
feedback and motor dysfunction is the focus of ongoing studies. This study represents the first evidence
of altered nociceptive afferents in a model of CP.
483
Each individual's pain perception includes complex psychosocial components that were not the
focus of this study. In situations where both nociception and the affective experience of pain can be
altered, such as after HI, it is critical to accurately identify and classify aberrant sensory behavior. Here,
we utilized methods to assess and quantify shifts in sensitivity to mechanical and thermal stimuli that
are sensitive, reliable, and validated in models of central and peripheral neuropathic pain across species.
We found that prenatal HI heightened nociception at the level of spinal nociceptors and reflexive
responses to mechanical and thermal stimuli. It could be that CP-causative perinatal injuries contribute
to other aspects of the pain experience in individuals with CP, including changes to mood and cognition,
in addition to altered nociception. Thus, treatment options for chronic pain in individuals with CP should
be explored with greater specificity, considering that the pathophysiology of their pain could be distinct
from that of chronic pain derived from other etiologies. Novel targets for fully alleviating CP-induced
chronic pain should be explored specifically within the context of treating individuals with CP or those
who have survived perinatal physical trauma.
497
Conclusions
The current study demonstrates enhanced nociceptive behavior and nociceptive primary
afferent distribution in neonatal rabbits after prenatal HI injury. Identification of how prenatal physical
stressors impact nociceptors could be the first step to preventing aberrant nociception and ultimately
chronic pain in those at risk.

Conflict of Interest Statement

Authors have no conflicts of interests to disclose.

Author contributions

EJ Reedich contributed to data curation, formal analysis, investigation, methodology, supervision,
validation and writing, editing and reviewing the manuscript. LT Genry contributed to data curation,
formal analysis, investigation, methodology, supervision, validation, and reviewing and editing the
manuscript. MA Singer contributed to data curation, formal analysis, investigation, methodology, and
validation. CF Cavarsan contributed to formal analysis, investigation, methodology, and validation. E
Mena Avila contributed to data curation, formal analysis, investigation, methodology, validation, and
reviewing the manuscript. DM Boudreau contributed to data curation, formal analysis, investigation,
methodology, validation, and reviewing the manuscript. MC Brennan contributed to data curation,
formal analysis, investigation, methodology, validation, and reviewing the manuscript. AM Garrett
contributed to data curation, formal analysis, investigation, methodology, validation, and reviewing the
manuscript. L Dowaliby contributed to investigation, methodology and supervision of this project. MR
Detloff contributed to contributed to conceptualization, data curation, funding acquisition, project
administration, resources, supervision, and writing, editing and review of the manuscript. KA Quinlan
contributed to conceptualization, data curation, funding acquisition, project administration, resources,
supervision, and writing, editing and review of the manuscript.
521

Other Acknowledgements

This project was supported by funding from NIH NINDS including NS097880 to MRD and
NS104436 to KQ. MCB and DMB received a grant from the University of Rhode Island Undergraduate
Research & Innovation. The funders of this project had no role in study design, the collection, analysis
and interpretation of data, in writing the report, or in the decision to publish these results.

Data Accessibility

The data that support the findings of this study are available from the corresponding author upon
reasonable request.

Literature Cited

Abraira, V.E., Ginty, D.D., 2013. The sensory neurons of touch. Neuron 79, 618–639.
https://doi.org/10.1016/j.neuron.2013.07.051

Abraira, V.E., Kuehn, E.D., Chirila, A.M., Springel, M.W., Toliver, A.A., Zimmerman, A.L., Orefice,
L.L., Boyle, K.A., Bai, L., Song, B.J., Bashista, K.A., O'Neill, T.G., Zhuo, J., Tsan, C., Hoynoski, J.,
Rutlin, M., Kus, L., Niederkofler, V., Watanabe, M., Dymecki, S.M., Nelson, S.B., Heintz, N.,
Hughes, D.I., Ginty, D.D., 2017. The Cellular and Synaptic Architecture of the Mechanosensory
Dorsal Horn. Cell 168, 295-310.e19. https://doi.org/10.1016/j.cell.2016.12.010

Achache, V., Roche, N., Lamy, J.C., Boakye, M., Lackmy, A., Gastal, A., Quentin, V., Katz, R., 2010.
Transmission within several spinal pathways in adults with cerebral palsy. Brain 133, 1470–
1483. https://doi.org/10.1093/brain/awq053

Advokat, C., 2002. Spinal transection increases the potency of clonidine on the tail-flick and
hindlimb flexion reflexes. European Journal of Pharmacology 437, 63–67.
https://doi.org/10.1016/S0014-2999(02)01259-1

Badia, M., Riquelme, I., Orgaz, B., Acevedo, R., Longo, E., Montoya, P., 2014. Pain, motor function
and health-related quality of life in children with cerebral palsy as reported by their
physiotherapists. BMC Pediatrics 14, 2–7. https://doi.org/10.1186/1471-2431-14-192

Banchi, P., Quaranta, G., Ricci, A., Von Degerfeld, M.M., 2020. Reliability and construct validity of a
composite pain scale for rabbit (CANCRS) in a clinical environment. PLoS ONE 15, 1–12.
https://doi.org/10.1371/journal.pone.0221377

Bedi, S.S., Yang, Q., Crook, R.J., Du, J., Wu, Z., Fishman, H.M., Grill, R.J., Carlton, S.M., Walters, E.T.,
2010. Chronic spontaneous activity generated in the somata of primary nociceptors is
associated with pain-related behavior after spinal cord injury. Journal of Neuroscience 30,
14870–14882. https://doi.org/10.1523/JNEUROSCI.2428-10.2010

Beggs, S., Torsney, C., Drew, L.J., Fitzgerald, M., 2002. The postnatal reorganization of primary
afferent input and dorsal horn cell receptive fields in the rat spinal cord is an activity-
dependent process. European Journal of Neuroscience 16, 1249–1258.
https://doi.org/10.1046/j.1460-9568.2002.02185.x

Benato, L., Murrell, J., Knowles, T.G., Rooney, N.J., 2021. Development of the Bristol Rabbit Pain
Scale (BRPS): A multidimensional composite pain scale specific to rabbits (Oryctolagus
cuniculus). PLoS ONE 16, 1–18. https://doi.org/10.1371/journal.pone.0252417

Black, S., 2019. C Fibres facilitate proprioceptive afferent transmission to motoneurons. Masters
Thesis, University of Alberta.

Boyle, K.A., Gradwell, M.A., Yasaka, T., Dickie, A.C., Polgár, E., Ganley, R.P., Orr, D.P.H., Watanabe,
M., Abraira, V.E., Kuehn, E.D., Zimmerman, A.L., Ginty, D.D., Callister, R.J., Graham, B.A.,
Hughes, D.I., 2019. Defining a Spinal Microcircuit that Gates Myelinated Afferent Input:
Implications for Tactile Allodynia. Cell Reports 28, 526-540.e6.
https://doi.org/10.1016/j.celrep.2019.06.040

Brown, P.B., Fuchs, J.L., 1975. Somatotopic Representation of Hindlimb Skin in Cat Dorsal Horn.
Journal of Neurophysiology 38, 1–9.

Bumin, G., Kayihan, H., 2001. Effectiveness of two different sensory-integration programmes for
children with spastic diplegic cerebral palsy. Disability and Rehabilitation 23, 394–399.
https://doi.org/10.1080/09638280010008843

Burchiel, K.J., Russell, L.C., Lee, R.P., Sima, A.A.F., 1985. Spontaneous Activity of Primary Afferent
Neurons in Diabetic BB/Wistar Rats: A possible mechanism of Diabetic Neuropathic Pain.
Diabetes 34.

Buser, J.R., Segovia, K.N., Dean, J.M., Nelson, K., Beardsley, D., Gong, X., Luo, N.L., Ren, J., Wan, Y.,
Riddle, A., Mcclure, M.M., Ji, X., Derrick, M., Hohimer, A.R., Back, S.A., Tan, S., 2010. Timing of
appearance of late oligodendrocyte progenitors coincides with enhanced susceptibility of
preterm rabbit cerebral white matter to hypoxia-ischemia. Journal of Cerebral Blood Flow and
Metabolism 30, 1053–1065. https://doi.org/10.1038/jcbfm.2009.286

Cahan, L.D., Kundi, M.S., McPherson, D., Starr, A., Peacock, W., 1987. Electrophysiologic Studies in
Selective Dorsal Rhizotomy for Spasticity in Children with Cerebral Palsy. Applied
Neurophysiology 50, 459–460.

Cavarsan, C.F., Gorassini, M.A., Quinlan, K.A., 2019. Animal models of developmental motor
disorders: parallels to human motor dysfunction in cerebral palsy. Journal of Neurophysiology
122, 1238–1253. https://doi.org/10.1152/jn.00233.2019

Chhaya, S.J., Quiros-Molina, D., Tamashiro-Orrego, A.D., Houle, J.D., Detloff, M.R., 2018. Exercise-
Induced Changes to the Macrophage Response in the Dorsal Root Ganglia Prevent
Neuropathic Pain after Spinal Cord Injury. Journal of Neurotrauma 14, neu.2018.5819.
https://doi.org/10.1089/neu.2018.5819

Choi, J.-G., Choi, S.-R., Kang, D.-W., Kim, J., Park, J.B., Lee, J.-H., Kim, H.-W., 2021. Sigma-1 receptor
increases intracellular calcium in cultured astrocytes and contributes to mechanical allodynia
in a model of neuropathic pain. Brain Research Bulletin 178, 69–81.
https://doi.org/10.1016/j.brainresbull.2021.11.010

Clancy, R.R., Sladky, J.T., Rorke, L.B., 1989. Hypoxic-ischemic spinal cord injury following perinatal
asphyxia. Annals of Neurology 25, 185–189. https://doi.org/10.1002/ana.410250213

Cooper, J., Majnemer, A., Rosenblatt, B., 1995. The Determination of Sensory Deficits in Children
With Hemiplegic Cerebral Palsy. Journal of Child Neurology 10, 300–309.
https://doi.org/10.1177/088307389501000412

Costigan, M., Scholz, J., Woolf, C.J., 2009. Neuropathic pain: A maladaptive response of the
nervous system to damage. Annual Review of Neuroscience 32, 1–32.
https://doi.org/10.1146/annurev.neuro.051508.135531

Derrick, M., Drobyshevsky, A., Ji, X., Tan, S., 2007. A model of cerebral palsy from fetal hypoxia-
ischemia. Stroke 38, 731–735. https://doi.org/10.1161/01.STR.0000251445.94697.64

Derrick, M., Luo, N.L., Bregman, J.C., Jilling, T., Ji, X., Fisher, K., Gladson, C.L., Beardsley, D.,
Murdoch, G., Back, S.A., Tan, S., 2004. Preterm Fetal Hypoxia-Ischemia Causes Hypertonia and
Motor Deficits in the Neonatal Rabbit: A Model for Human Cerebral Palsy? Journal of
Neuroscience 24, 24–34. https://doi.org/10.1523/JNEUROSCI.2816-03.2004

Detloff, M.R., Fisher, L.C., McGaughy, V., Longbrake, E.E., Popovich, P.G., Basso, D.M., 2008.
Remote activation of microglia and pro-inflammatory cytokines predict the onset and severity
of below-level neuropathic pain after spinal cord injury in rats. Experimental Neurology 212,
337–347. https://doi.org/10.1016/j.expneurol.2008.04.009

Detloff, M.R., Quiros-Molina, D., Javia, A.S., Daggubati, L., Nehlsen, A.D., Naqvi, A., Ninan, V.,
Vannix, K.N., McMullen, M.K., Amin, S., Ganzer, P.D., Houlé, J.D., 2016. Delayed Exercise Is
Ineffective at Reversing Aberrant Nociceptive Afferent Plasticity or Neuropathic Pain after
Spinal Cord Injury in Rats. Neurorehabilitation and Neural Repair 30, 685–700.
https://doi.org/10.1177/1545968315619698

Detloff, M.R., Smith, E.J., Quiros Molina, D., Ganzer, P.D., Houlé, J.D., 2014. Acute exercise prevents
the development of neuropathic pain and the sprouting of non-peptidergic (GDNF- and
artemin-responsive) c-fibers after spinal cord injury. Experimental Neurology 255, 38–48.
https://doi.org/10.1016/j.expneurol.2014.02.013

Doralp, S., Bartlett, D.J., 2010. The prevalence, distribution, and effect of pain among adolescents
with cerebral palsy. Pediatric Physical Therapy 22, 26–33.
https://doi.org/10.1097/PEP.0b013e3181ccbabb

Dougherty, P.M., Cata, J.P., Cordella, J. V., Burton, A., Weng, H.R., 2004. Taxol-induced sensory
disturbance is characterized by preferential impairment of myelinated fiber function in cancer
patients. Pain 109, 132–142. https://doi.org/10.1016/j.pain.2004.01.021

Drobyshevsky, A., Derrick, M., Wyrwicz, A.M., Ji, X., Englof, I., Ullman, L.M., Zelaya, M.E.,
Northington, F.J., Tan, S., 2007. White matter injury correlates with hypertonia in an animal
model of cerebral palsy. Journal of Cerebral Blood Flow and Metabolism 27, 270–281.
https://doi.org/10.1038/sj.jcbfm.9600333

Drobyshevsky, A., Quinlan, K.A., 2017. Spinal cord injury in hypertonic newborns after antenatal
hypoxia-ischemia in a rabbit model of cerebral palsy. Experimental Neurology 293, 13–26.
https://doi.org/10.1016/j.expneurol.2017.03.017

Ducza, L., Szücs, P., Hegedűs, K., Bakk, E., Gajtkó, A., Wéber, I., Holló, K., 2021. Nlrp2 is
overexpressed in spinal astrocytes at the peak of mechanical pain sensitivity during complete
freund adjuvant-induced persistent pain. International Journal of Molecular Sciences 22.
https://doi.org/10.3390/ijms222111408

Escartin, C., Galea, E., Lakatos, A., O'Callaghan, J.P., Petzold, G.C., Serrano-Pozo, A., Steinhäuser, C.,
Volterra, A., Carmignoto, G., Agarwal, A., Allen, N.J., Araque, A., Barbeito, L., Barzilai, A.,
Bergles, D.E., Bonvento, G., Butt, A.M., Chen, W.T., Cohen-Salmon, M., Cunningham, C.,
Deneen, B., De Strooper, B., Díaz-Castro, B., Farina, C., Freeman, M., Gallo, V., Goldman, J.E.,
Goldman, S.A., Götz, M., Gutiérrez, A., Haydon, P.G., Heiland, D.H., Hol, E.M., Holt, M.G., Iino,
M., Kastanenka, K. V., Kettenmann, H., Khakh, B.S., Koizumi, S., Lee, C.J., Liddelow, S.A.,
MacVicar, B.A., Magistretti, P., Messing, A., Mishra, A., Molofsky, A. V., Murai, K.K., Norris,
C.M., Okada, S., Oliet, S.H.R., Oliveira, J.F., Panatier, A., Parpura, V., Pekna, M., Pekny, M.,
Pellerin, L., Perea, G., Pérez-Nievas, B.G., Pfrieger, F.W., Poskanzer, K.E., Quintana, F.J.,
Ransohoff, R.M., Riquelme-Perez, M., Robel, S., Rose, C.R., Rothstein, J.D., Rouach, N.,
Rowitch, D.H., Semyanov, A., Sirko, S., Sontheimer, H., Swanson, R.A., Vitorica, J., Wanner,
I.B., Wood, L.B., Wu, J., Zheng, B., Zimmer, E.R., Zorec, R., Sofroniew, M. V., Verkhratsky, A.,
2021. Reactive astrocyte nomenclature, definitions, and future directions. Nature
Neuroscience 24, 312–325. https://doi.org/10.1038/s41593-020-00783-4

Evangelista, M.C., Monteiro, B.P., Steagall, P. V, 2021. Measurement properties of grimace scales
for pain assessment in non-human mammals, Pain.
https://doi.org/10.1097/j.pain.0000000000002474

Fan, L.W., Lin, S., Pang, Y., Lei, M., Zhang, F., Rhodes, P.G., Cai, Z., 2005. Hypoxia-ischemia induced
neurological dysfunction and brain injury in the neonatal rat. Behavioural Brain Research 165,
80–90. https://doi.org/10.1016/j.bbr.2005.06.033

Findlay, B., Switzer, L., Narayanan, U., Chen, S., Fehlings, D., 2016. Investigating the impact of pain,
age, Gross Motor Function Classification System, and sex on health-related quality of life in
children with cerebral palsy. Developmental Medicine and Child Neurology 58, 292–297.
https://doi.org/10.1111/dmcn.12936

Fitzgerald, M., 1985. The sprouting of saphenous nerve terminals in the spinal cord following early
postnatal sciatic nerve section in the rat. Journal of Comparative Neurology 240, 407–413.
https://doi.org/10.1002/cne.902400408

Fitzgerald, M., Jennings, E., 1999. The postnatal development of spinal sensory processing. Proc
Natl Acad Sci U S A 96, 7719–7722. https://doi.org/10.1073/pnas.96.14.7719

Fitzgerald, M., Swett, J., 1983. THE TERMINATION PATTERN OF SCIATIC NERVE AFFERENTS IN THE
SUBSTANTIA GELATINOSA OF NEONATAL RATS. Neuroscience Letters 43, 149–154.

Flanigan, M., Gaebler-Spira, D., Kocherginsky, M., Garrett, A., Marciniak, C., 2020. Spasticity and
pain in adults with cerebral palsy. Developmental Medicine and Child Neurology 62, 379–385.
https://doi.org/10.1111/dmcn.14368

Gao, Q., Li, Y., Chopp, M., 2005. Bone marrow stromal cells increase astrocyte survival via
upregulation of phosphoinositide 3-kinase/threonine protein kinase and mitogen-activated
protein kinase kinase/extracellular signal-regulated kinase pathways and stimulate astrocyte
trophic factor . Neuroscience 136, 123–134.
https://doi.org/10.1016/j.neuroscience.2005.06.091

Gao, Y.J., Zhang, L., Samad, O.A., Suter, M.R., Yasuhiko, K., Xu, Z.Z., Park, J.Y., Lind, A.L., Ma, Q., Ji,
R.R., 2009. JNK-induced MCP-1 production in spinal cord astrocytes contributes to central
sensitization and neuropathic pain. Journal of Neuroscience 29, 4096–4108.
https://doi.org/10.1523/JNEUROSCI.3623-08.2009

García Jalón, E.G., Maguire, A., Perra, O., Gavin, A., O'Reilly, D., Thurston, A., 2021. Data linkage
and pain medication in people with cerebral palsy: a cross-sectional study. Developmental
Medicine and Child Neurology 1–8. https://doi.org/10.1111/dmcn.14854

Gold, M.S., Gebhart, G.F., 2010. Nociceptor sensitization in pain pathogenesis. Nature Medicine 16,
1248–1257. https://doi.org/10.1038/nm.2235

Grace, P.M., Hutchinson, M.R., Maier, S.F., Watkins, L.R., 2014. Pathological pain and the
neuroimmune interface. Nature Reviews Immunology 14, 217–231.
https://doi.org/10.1038/nri3621

Gracely, R.H., Lynch, S.A., Bennett, G.J., 1993. Painful neuropathy: altered central processing
maintained dynamically by peripheral input. Pain 52, 251–253. https://doi.org/10.1016/0304-
3959(93)90141-B

Gupta, D., Barachant, A., Gordon, A.M., Ferre, C., Kuo, H.C., Carmel, J.B., Friel, K.M., 2017. Effect of
sensory and motor connectivity on hand function in pediatric hemiplegia. Annals of
Neurology 82, 766–780. https://doi.org/10.1002/ana.25080

Hari, K., Lucas-Osma, A.M., Metz, K., Lin, S., Pardell, N., Roszko, D., Black, S., Minarik, A., Singla, R.,
Stephens, M.J., Fouad, K., Jones, K.E., Gorassini, M.A., Fenrich, K.K., Li, Y., Bennett, D.J., 2021.
Nodal GABA facilitates axon spike transmission in the spinal cord. bioRxiv 2021.01.20.427494.

Hayakawa, K., Pham, L.D.D., Arai, K., Lo, E.H., 2014. Reactive astrocytes promote adhesive
interactions between brain endothelium and endothelial progenitor cells via HMGB1 and
beta-2 integrin signaling. Stem Cell Research 12, 531–538.
https://doi.org/10.1016/j.scr.2013.12.008

Helou, A.Y., Martins, D.O., Arruda, B.P., de Souza, M.C., Cruz-Ochoa, N.A., Nogueira, M.I., Chacur,
M., 2021. Neonatal anoxia increases nociceptive response in rats: Sex differences and lumbar
spinal cord and insula alterations. International Journal of Developmental Neuroscience 81,
686–697. https://doi.org/10.1002/jdn.10145

Himes, B.T., Tessler, A., 1989. Death of some dorsal root ganglion neurons and plasticity of others
following sciatic nerve section in adult and neonatal rats. Journal of Comparative Neurology
284, 215–230. https://doi.org/10.1002/cne.902840206

Hodges, P.W., Tucker, K., 2011. Moving differently in pain: A new theory to explain the adaptation
to pain. Pain 152, S90–S98. https://doi.org/10.1016/j.pain.2010.10.020

Hoon, A.H., Lawrie, W.T., Melhem, E.R., Reinhardt, E.M., Van Zijl, P.C.M., Solaiyappan, M., Jiang, H.,
Johnston, M. V., Mori, S., 2002. Diffusion tensor imaging of periventricular leukomalacia
shows affected sensory cortex white matter pathways. Neurology 59, 752–756.
https://doi.org/10.1212/WNL.59.5.752

Hulsebosch, C.E., Coggeshall, R.E., 1981a. Quantitation of Sprouting of Dorsal Root Axons. Science
213, 1020–1021.

Hulsebosch, C.E., Coggeshall, R.E., 1981b. Sprouting of dorsal root axons. Brain Research 224, 170–
174. https://doi.org/10.1016/0006-8993(81)91128-8

Hulsebosch, C.E., Hains, B.C., Crown, E.D., Carlton, S.M., 2009. Mechanisms of chronic central
neuropathic pain after spinal cord injury. Brain Research Reviews 60, 202–213.
https://doi.org/10.1016/j.brainresrev.2008.12.010

Ishikura, T., Kinoshita, M., Shimizu, M., Yasumizu, Y., Motooka, D., Okuzaki, D., Yamashita, K.,
Murata, H., Beppu, S., Koda, T., Tada, S., Shiraishi, N., Sugiyama, Y., Miyamoto, K., Kusunoki,
S., Sugimoto, T., Kumanogoh, A., Okuno, T., Mochizuki, H., 2021. Anti-AQP4 autoantibodies
promote ATP release from astrocytes and induce mechanical pain in rats. Journal of
Neuroinflammation 18, 1–12. https://doi.org/10.1186/s12974-021-02232-w

Jennings, E., Fitzgerald, M., 1996. C-fos can be induced in the neonatal rat spinal cord by both
noxious and innocuous peripheral stimulation. Pain 68, 301–306.

Jiang, Y.-Q., Zaaimi, B., Martin, J.H., 2016. Competition with Primary Sensory Afferents Drives
Remodeling of Corticospinal Axons in Mature Spinal Motor Circuits. Journal of Neuroscience
36, 193–203. https://doi.org/10.1523/JNEUROSCI.3441-15.2016

Kaelin-Lang, A., 2008. Enhancing rehabilitation of motor deficits with peripheral nerve stimulation.
NeuroRehabilitation 23, 89–93. https://doi.org/10.3233/nre-2008-23109

Kandel, E.R., Schwartz, J.H., Jessell, T.M., Siegelbaum, S.A., Hudspeth, A.J. (Eds.), 2013. Principles of
Neural Science, Fifth. ed. McGraw Hill, United States of America.

Kauppila, T., Kontinen, V.K., Pertovaara, A., 1998. Influence of spinalization on spinal withdrawal
reflex responses varies depending on the submodality of the test stimulus and the
experimental pathophysiological condition in the rat. Brain Research 797, 234–242.
https://doi.org/10.1016/S0006-8993(98)00379-5

Keating, S.C.J., Thomas, A.A., Flecknell, P.A., Leach, M.C., 2012. Evaluation of EMLA Cream for
Preventing Pain during Tattooing of Rabbits: Changes in Physiological, Behavioural and Facial
Expression Responses. PLoS ONE 7, 1–11. https://doi.org/10.1371/journal.pone.0044437

Keller, A.V.P., Wainwright, G., Shum-Siu, A., Prince, D., Hoeper, A., Martin, E., Magnuson, D.S.K.,
2017. Disruption of locomotion in response to Hindlimb muscle stretch at acute and chronic
time points after a spinal cord injury in rats. Journal of Neurotrauma 34, 661–670.
https://doi.org/10.1089/neu.2015.4227

Keller, A. V., Hainline, C., Rees, K., Krupp, S., Prince, D., Wood, B.D., Shum-Siu, A., Burke, D.A.,
Petruska, J.C., Magnuson, D.S.K., 2019. Nociceptor-dependent locomotor dysfunction after
clinically-modeled hindlimb muscle stretching in adult rats with spinal cord injury.
Experimental Neurology 318, 267–276. https://doi.org/10.1016/j.expneurol.2019.03.006

Keller, A. V., Rees, K.M., Seibt, E.J., Wood, B.D., Wade, A.D., Morehouse, J., Shum-Siu, A.,
Magnuson, D.S.K., 2018. Electromyographic patterns of the rat hindlimb in response to
muscle stretch after spinal cord injury. Spinal Cord 56, 560–568.
https://doi.org/10.1038/s41393-018-0069-z

Kimura, S., Honda, M., Tanabe, M., Ono, H., 2004. Noxious stimuli evoke a biphasic flexor reflex
composed of Aδ-fiber-mediated short-latency and C-fiber-mediated long-latency withdrawal
movements in mice. Journal of Pharmacological Sciences 95, 94–100.
https://doi.org/10.1254/jphs.95.94

Koerber, H.R., Mirnics, K., Brown, P.B., Mendell, L.M., 1994. Central sprouting and functional
plasticity of regenerated primary afferents. Journal of Neuroscience 14, 3655–3671.
https://doi.org/10.1523/jneurosci.14-06-03655.1994

Krenz, N.R., Weaver, L.C., 1998. Sprouting of primary afferent fibers after spinal cord transection in
the rat. Neuroscience 85, 443–458. https://doi.org/10.1016/S0306-4522(97)00622-2

Kumar, A.J., Martins, D.O., Arruda, B.P., Lee, V.Y., Chacur, M., Nogueira, M.I., 2020. Impairment of
767   nociceptive responses after neonatal anoxia correlates with somatosensory thalamic damage:
768   A study in rats. Behavioural Brain Research 390. https://doi.org/10.1016/j.bbr.2020.112690

769   Kumar, A.J., Takada, S.H., Motta-Teixeira, L.C., Lee, V.Y., Xavier, G.F., Nogueira, M.I., 2017. Sex
770   differences in somatic and sensory motor development after neonatal anoxia in Wistar rats.
771   Behavioural Brain Research 333, 242–250. https://doi.org/10.1016/j.bbr.2017.07.009

772   Kundi, M., Starr, A., Cahan, L., 1989. Somatosensory Evoked Potentials in Cerebral Palsy After
773   Partial Dorsal Root Rhizotomy. Archives of Neurology 46, 524–527.
774   https://doi.org/10.1001/archneur.1989.00520410058024

775   Kusunoki, S., Inoue, K., Iwamori, M., Mannen, T., Kanazawa, I., 1992. Developmental changes of
776   fucosylated glycoconjugates in rabbit dorsal root ganglia, Neuroscienee Research.

777   Lallemend, F., Ernfors, P., 2012. Molecular interactions underlying the specification of sensory
778   neurons. Trends in Neurosciences 35, 373–381. https://doi.org/10.1016/j.tins.2012.03.006

779   Landry, C.F., Ivy, G.O., Brown, I.R., 1990. Developmental expression of glial fibrillary acidic protein
780   mRNA in the rat brain analyzed by in situ hybridization. Journal of Neuroscience Research 25,
781   194–203. https://doi.org/10.1002/jnr.490250207

782   Laurin, J., Pertici, V., Dousset, E., Marqueste, T., Decherchi, P., 2015. Group III and IV muscle
783   afferents: Role on central motor drive and clinical implications. Neuroscience 290, 543–551.
784   https://doi.org/10.1016/j.neuroscience.2015.01.065

785   Lei, T., Zhang, Y., Zhou, Q., Luo, X., Tang, K., Chen, R., Yu, C., Quan, Z., 2017. A novel approach for
786   the annulus needle puncture model of intervertebral disc degeneration in rabbits. American
787   Journal of Translational Research 9, 900–909.

788   Leonard, C.T., Hirschfeld, H., 1995. Myotatic Reflex Responses of Non-Disabled Children and
789   Children with Spastic Cerebral Palsy. Developmental Medicine & Child Neurology 37, 783–
790   799.

791   Leonard, C.T., Hirschfeld, H., Moritani, T., Forssberg, H., 1991. Myotatic reflex development in
792   normal children and children with cerebral palsy. Experimental Neurology 111, 379–382.
793   https://doi.org/10.1016/0014-4886(91)90106-M

794   Li, Y., North, R.Y., Rhines, L.D., Tatsui, C.E., Rao, G., Edwards, D.D., Cassidy, R.M., Harrison, D.S.,
795   Johansson, C.A., Zhang, H., Dougherty, P.M., 2018. DRG voltage-gated sodium channel 1.7 is
796   upregulated in paclitaxel-induced neuropathy in rats and in humans with neuropathic pain.
797   Journal of Neuroscience 38, 1124–1136. https://doi.org/10.1523/JNEUROSCI.0899-17.2017

798   Liddelow, S.A., Barres, B.A., 2017. Reactive Astrocytes: Production, Function, and Therapeutic
799   Potential. Immunity 46, 957–967. https://doi.org/10.1016/j.immuni.2017.06.006

800   Liedtke, W., Edelmann, W., Bieri, P.L., Chiu, F.C., Cowan, N.J., Kucherlapati, R., Raine, C.S., 1996.
801   GFAP is necessary for the integrity of CNS white matter architecture and long-term
802   maintenance of myelination. Neuron 17, 607–615. https://doi.org/10.1016/S0896-
803   6273(00)80194-4

Lu, R., Cui, S.S., Wang, X.X., Chen, L., Liu, F., Gao, J., Wang, W., 2021. Astrocytic c-Jun N-terminal
kinase-histone deacetylase-2 cascade contributes to glutamate transporter-1 decrease and
mechanical allodynia following peripheral nerve injury in rats. Brain Research Bulletin 175,
213–223. https://doi.org/10.1016/j.brainresbull.2021.07.024

Lucas-Osma, A.M., Li, Y., Murray, K., Lin, S., Black, S., Stephens, M.J., Ahn, A.H., Heckman, C.J.,
Fenrich, K.K., Fouad, K., Bennett, D.J., 2019. 5-HT1D receptors inhibit the monosynaptic
stretch reflex by modulating C-fiber activity. Journal of Neurophysiology 121, 1591–1608.
https://doi.org/10.1152/jn.00805.2018

Mans, C., 2020. Insights into pain assessment and management in rabbits. Veterinary Record 186,
600–602. https://doi.org/10.1136/vr.m2265

Marmigère, F., Ernfors, P., 2007. Specification and connectivity of neuronal subtypes in the sensory
lineage. Nature Reviews Neuroscience 8, 114–127. https://doi.org/10.1038/nrn2057

Marriott, A.L., Rojas-Mancilla, E., Morales, P., Herrera-Marschitz, M., Tasker, R.A., 2017. Models of
progressive neurological dysfunction originating early in life. Progress in Neurobiology 155, 2–
20. https://doi.org/10.1016/j.pneurobio.2015.10.001

Martinez-Valdes, E., Negro, F., Farina, D., Falla, D., 2020. Divergent response of low- versus high-
threshold motor units to experimental muscle pain. Journal of Physiology 598, 2093–2108.
https://doi.org/10.1113/JP279225

Matusz, P.J., Key, A.P., Gogliotti, S., Pearson, J., Auld, M.L., Murray, M.M., Maitre, N.L., 2018.
Somatosensory Plasticity in Pediatric Cerebral Palsy following Constraint-Induced Movement
Therapy. Neural Plasticity 2018. https://doi.org/10.1155/2018/1891978

McDowell, B.C., Duffy, C., Lundy, C., 2017. Pain report and musculoskeletal impairment in young
people with severe forms of cerebral palsy: A population-based series. Research in
Developmental Disabilities 60, 277–284. https://doi.org/10.1016/j.ridd.2016.10.006

Mckinnon, C.T., Meehan, E.M., Harvey, A.R., Antolovich, G.C., Morgan, P.E., 2018. Prevalence and
characteristics of pain in children and young adults with cerebral palsy: a systematic review.
Developmental Medicine & Child Neurology 61, 305–314.
https://doi.org/10.1111/dmcn.14111

McKinnon, C.T., Morgan, P.E., Antolovich, G.C., Clancy, C.H., Fahey, M.C., Harvey, A.R., 2020. Pain
in children with dyskinetic and mixed dyskinetic/spastic cerebral palsy. Developmental
Medicine and Child Neurology 62, 1294–1301. https://doi.org/10.1111/dmcn.14615

Mesquita, R.N.O., Škarabot, J., Pearcey, G.E.P., 2020. Low-threshold motor units can be a pain
during experimental muscle pain. Journal of Physiology 598, 2545–2547.
https://doi.org/10.1113/JP279872

Milligan, E.D., Mehmert, K.K., Hinde, J.L., Harvey, L.O., Martin, D., Tracey, K.J., Maier, S.F., Watkins,
L.R., 2000. Thermal hyperalgesia and mechanical allodynia produced by intrathecal
administration of the human immunodeficiency virus-1 (HIV-1) envelope glycoprotein, gp120.
Brain Research 861, 105–116. https://doi.org/10.1016/S0006-8993(00)02050-3

842 Mori, K., 1986. Lectin Ulex europaeus agglutinin I specifically labels a subset of primary afferent
843   fibers which project selectively to the superficial dorsal horn of the spinal cord. Brain
844   Research 365, 404–408.

845 Morin-Richaud, C., Feldblum, S., Privat, A., 1998. Astrocytes and oligodendrocytes reactions after a
846   total section of the rat spinal cord. Brain Research 783, 85–101.
847   https://doi.org/10.1016/S0006-8993(97)01282-1

848 Moy, J.K., Kuhn, J.L., Szabo-Pardi, T.A., Pradhan, G., Price, T.J., 2018. eIF4E phosphorylation
849   regulates ongoing pain, independently of inflammation, and hyperalgesic priming in the
850   mouse CFA model. Neurobiology of Pain 4, 45–50.
851   https://doi.org/10.1016/j.ynpai.2018.03.001

852 Murphy, R.M., Zemlan, F.P., 1990. Selective serotonin1A/1B agonists differentially affect spinal
853   nociceptive reflexes. Neuropharmacology 29, 463–468. https://doi.org/10.1016/0028-
854   3908(90)90168-Q

855 Nash, B., Ioannidou, K., Barnett, S.C., 2011. Astrocyte phenotypes and their relationship to
856   myelination. Journal of Anatomy 219, 44–52. https://doi.org/10.1111/j.1469-
857   7580.2010.01330.x

858 Nesic, O., Lee, J., Ye, Z., Unabia, G.C., Rafati, D., Hulsebosch, C.E., Perez-Polo, J.R., 2006. Acute and
859   chronic changes in aquaporin 4 expression after spinal cord injury. Neuroscience 143, 779–
860   792. https://doi.org/10.1016/j.neuroscience.2006.08.079

861 Nitzan-Luques, A., Devor, M., Tal, M., 2011. Genotype-selective phenotypic switch in primary
862   afferent neurons contributes to neuropathic pain. Pain 152, 2413–2426.
863   https://doi.org/10.1016/j.pain.2011.07.012

864 North, R.Y., Li, Y., Ray, P., Rhines, L.D., Tatsui, C.E., Rao, G., Johansson, C.A., Zhang, H., Kim, Y.H.,
865   Zhang, B., Dussor, G., Kim, T.H., Price, T.J., Dougherty, P.M., 2019. Electrophysiological and
866   transcriptomic correlates of neuropathic pain in human dorsal root ganglion neurons. Brain
867   142, 1215–1226. https://doi.org/10.1093/brain/awz063

868 Novak, I., Hines, M., Goldsmith, S., Barclay, R., 2012. Clinical Prognostic Messages From a
869   Systematic Review on Cerebral Palsy. Pediatrics 130, e1285–e1312.
870   https://doi.org/10.1542/peds.2012-0924

871 Olson, W., Luo, W., 2018. Somatotopic organization of central arbors from nociceptive afferents
872   develops independently of their intact peripheral target innervation. Journal of Comparative
873   Neurology 526, 3058–3065. https://doi.org/10.1002/cne.24533

874 Ondarza, A.B., Ye, Z., Hulsebosch, C.E., 2003. Direct evidence of primary afferent sprouting in
875   distant segments following spinal cord injury in the rat: Colocalization of GAP-43 and CGRP.
876   Experimental Neurology 184, 373–380. https://doi.org/10.1016/j.expneurol.2003.07.002

877 Orozco, O.E., Walus, L., Sah, D.W.Y., Pepinsky, R.B., Sanicola, M., 2001. GFRalpha3 is expressed
878   predominantly in nociceptive sensory neurons. European Journal of Neuroscience 13, 2177–
879   2182. https://doi.org/10.1046/j.0953-816X.2001.01596.x

Oskoui, M., Coutinho, F., Dykeman, J., Jetté, N., Pringsheim, T., 2013. An update on the prevalence
of cerebral palsy: A systematic review and meta-analysis. Developmental Medicine and Child
Neurology 55, 509–519. https://doi.org/10.1111/dmcn.12080

Pattinson, D., Fitzgerald, M., 2004. The Neurobiology of Infant Pain: Development of Excitatory and
Inhibitory Neurotransmission in the Spinal Dorsal Horn. Regional Anesthesia and Pain
Medicine 29, 36–44.

Peirs, C., Williams, S.P.G., Zhao, X., Arokiaraj, C.M., Ferreira, D.W., Noh, M. chul, Smith, K.M.,
Halder, P., Corrigan, K.A., Gedeon, J.Y., Lee, S.J., Gatto, G., Chi, D., Ross, S.E., Goulding, M.,
Seal, R.P., 2021. Mechanical Allodynia Circuitry in the Dorsal Horn Is Defined by the Nature of
the Injury. Neuron 109, 73-90.e7. https://doi.org/10.1016/j.neuron.2020.10.027

Pinho, R.H., Leach, M.C., Minto, B.W., Del Lama Rocha, F., Luna, S.P.L., 2020. Postoperative pain
behaviours in rabbits following orthopaedic surgery and effect of observer presence. PLoS
ONE 15, 1–21. https://doi.org/10.1371/journal.pone.0240605

Pitcher, G.M., Henry, J.L., 2008. Governing role of primary afferent drive in increased excitation of
spinal nociceptive neurons in a model of sciatic neuropathy. Experimental Neurology 214,
219–228. https://doi.org/10.1016/j.expneurol.2008.08.003

Price, T.J., Dussor, G., 2014. Evolution: The advantage of "maladaptive" pain plasticity. Current
Biology 24, R384–R386. https://doi.org/10.1016/j.cub.2014.04.011

Price, T.J., Flores, C.M., 2007. Critical evaluation of the colocalization between calcitonin gene-
related peptide, substance P, transient receptor potential vanilloid subfamily type 1
immunoreactivities and isolectin B 4 binding in primary afferent neurons of the rat and
mouse, J Pain.

Ralston, H.J., Ralston, D.D., 1979. The distribution of dorsal root axons in laminae I, II and III of the
macaque spinal cord: A quantitative electron microscope study. Journal of Comparative
Neurology 184, 643–683. https://doi.org/10.1002/cne.901840404

Ramer, L.M., Peter van Stolk, A., Inskip, J.A., Ramer, M.S., Krassioukov, A. V., 2012. Plasticity of
TRPV1-expressing sensory neurons mediating autonomic dysreflexia following spinal cord
injury. Frontiers in Physiology 3, 1–16. https://doi.org/10.3389/fphys.2012.00257

Rice, J.E., Vannucci, R.C., Brierley, J.B., 1981. The influence of immaturity on hypoxic-ischemic brain
damage in the rat. Annals of Neurology 9, 131–141. https://doi.org/10.1002/ana.410090206

Riquelme, I., Montoya, P., 2010. Developmental changes in somatosensory processing in cerebral
palsy and healthy individuals. Clinical Neurophysiology 121, 1314–1320.
https://doi.org/10.1016/j.clinph.2010.03.010

Rocha-Ferreira, E., Hristova, M., 2016. Plasticity in the neonatal brain following hypoxic-ischaemic
injury. Neural Plasticity 2016. https://doi.org/10.1155/2016/4901014

Rodin, B., Sampogna, S.L., Kruger, L.W., 1983. An examination of intraspinal sprouting in dorsal
root axons with the tracer horseradish peroxidase. Journal of Comparative Neurology 215,
187–198. https://doi.org/10.1002/cne.902150206

Ruda, M.A., Ling, Q.D., Hohmann, A.G., Peng, Y.B., Tachibana, T., 2000. Altered nociceptive
neuronal circuits after neonatal peripheral inflammation. Science 289, 628–630.
https://doi.org/10.1126/science.289.5479.628

Scheibel, M.E., Scheibel, A.B., 1968. TERMINAL AXONAL PATTERNS IN CAT SPINAL CORD. II. THE
DORSAL HORN. Brain Research 9, 32–58.

Schomburg, E.D., Steffens, H., 1998. Comparative analysis of L-DOPA actions on nociceptive and
non-nociceptive spinal reflex pathways in the cat. Neuroscience Research 31, 307–316.
https://doi.org/10.1016/S0168-0102(98)00058-3

Shi, Z., Luo, K., Jani, S., February, M., Fernandes, N., Venkatesh, N., Sharif, N., Tan, S., 2021.
Mimicking partial to total placental insufficiency in a rabbit model of cerebral palsy. Journal of
Neuroscience Research 1–16. https://doi.org/10.1002/jnr.24901

Shiers, S., Klein, R.M., Price, T.J., 2020. Quantitative differences in neuronal subpopulations
between mouse and human dorsal root ganglia demonstrated with RNAscope in situ
hybridization. Pain 161, 2410–2424. https://doi.org/10.1097/j.pain.0000000000001973

Shortland, P., Fitzgerald, M., 1994. Neonatal Sciatic Nerve Section Results in a Rearrangement of
the Central Terminals of Saphenous and Axotomized Sciatic Nerve Afferents in the Dorsal
Horn of the Spinal Cord of the Adult Rat. European Journal of Neuroscience 6, 75–86.

Sienko, S.E., 2018. An exploratory study investigating the multidimensional factors impacting the
health and well-being of young adults with cerebral palsy. Disability and Rehabilitation 40,
660–666. https://doi.org/10.1080/09638288.2016.1274340

Sladky, J.T., Rorke, L.B., 1986. Perinatal hypoxic/ischemic spinal cord injury. Fetal and Pediatric
Pathology 6, 87–101. https://doi.org/10.3109/15513818609025927

Sliwinski, C., Nees, T.A., Puttagunta, R., Weidner, N., Blesch, A., 2018. Sensorimotor activity
partially ameliorates pain and reduces nociceptive fiber density in the chronically injured
spinal cord. Journal of Neurotrauma 35, 2222–2238. https://doi.org/10.1089/neu.2017.5431

Steele, P.R., Cavarsan, C.F., Dowaliby, L., Westefeld, M., Katenka, N., Drobyshevsky, A., Gorassini,
M.A., Quinlan, K.A., 2020. Altered Motoneuron Properties Contribute to Motor Deficits in a
Rabbit Hypoxia-Ischemia Model of Cerebral Palsy. Frontiers in Cellular Neuroscience 14, 1–12.
https://doi.org/10.3389/fncel.2020.00069

Swett, J.E., Woolf, C.J., 1985. The somatotopic organization of primary afferent terminals in the
superficial laminae of the dorsal horn of the rat spinal cord. Journal of Comparative
Neurology 231, 66–77. https://doi.org/10.1002/cne.902310106

Synowiec, S., Lu, J., Yu, L., Goussakov, I., Lieber, R., Drobyshevsky, A., 2019. Spinal Hyper-
Excitability and Altered Muscle Structure Contribute to Muscle Hypertonia in Newborns After
Antenatal Hypoxia-Ischemia in a Rabbit Cerebral Palsy Model. Frontiers in Neurology 9, 1–18.
https://doi.org/10.3389/fneur.2018.01183

Tan, A.M., Chakrabarty, S., Kimura, H., Martin, J.H., 2012. Selective corticospinal tract injury in the
rat induces primary afferent fiber sprouting in the spinal cord and hyperreflexia. Journal of
Neuroscience 32, 12896–12908. https://doi.org/10.1523/JNEUROSCI.6451-11.2012

Torsney, C., Fitzgerald, M., 2002. Age-dependent effects of peripheral inflammation on the
electrophysiological properties of neonatal rat dorsal horn neurons. Journal of
Neurophysiology 87, 1311–1317. https://doi.org/10.1152/jn.00462.2001

Vega-Avelaira, D., Moss, A., Fitzgerald, M., 2007. Age-related changes in the spinal cord microglial
and astrocytic response profile to nerve injury. Brain, Behavior, and Immunity 21, 617–623.
https://doi.org/10.1016/j.bbi.2006.10.007

Walker, S.M., Melbourne, A., O'Reilly, H., Beckmann, J., Eaton-Rosen, Z., Ourselin, S., Marlow, N.,
2018. Somatosensory function and pain in extremely preterm young adults from the UK
EPICure cohort: sex-dependent differences and impact of neonatal surgery. British Journal of
Anaesthesia 121, 623–635. https://doi.org/10.1016/j.bja.2018.03.035

Walker, S.M., Meredith-Middleton, J., Cooke-Yarborough, C., Fitzgerald, M., 2003. Neonatal
inflammation and primary afferent terminal plasticity in the rat dorsal horn. Pain 105, 185–
195. https://doi.org/10.1016/S0304-3959(03)00201-X

Walters, E.T., 2012. Nociceptors as chronic drivers of pain and hyperreflexia after spinal cord
injury: An adaptive-maladaptive hyperfunctional state hypothesis. Frontiers in Physiology 3,
1–13. https://doi.org/10.3389/fphys.2012.00309

Weaver, L.C., Verghese, P., Bruce, J.C., Fehlings, M.G., Krenz, N.R., Marsh, D.R., 2001. Autonomic
dysreflexia and primary afferent sprouting after clip-compression injury of the rat spinal cord.
Journal of Neurotrauma 18, 1107–1119. https://doi.org/10.1089/08977150152693782

Westbom, L., Rimstedt, A., Nordmark, E., 2017. Assessments of pain in children and adolescents
with cerebral palsy: a retrospective population-based registry study. Developmental Medicine
and Child Neurology 59, 858–863. https://doi.org/10.1111/dmcn.13459

Woolf, C.J., Shortland, P., Coggeshall, R.E., 1992. Peripheral nerve injury triggers central sprouting
of myelinated afferents. Nature 355, 75–78. https://doi.org/10.1038/355075a0

Xie, W., Strong, J.A., Zhang, J.M., 2009. Early blockade of injured primary sensory afferents reduces
glial cell activation in two rat neuropathic pain models. Neuroscience 160, 847–857.
https://doi.org/10.1016/j.neuroscience.2009.03.016

Xu, Q., Ford, N.C., He, S., Huang, Q., Anderson, M., Chen, Z., Yang, F., Crawford, L.K., Caterina, M.J.,
Guan, Y., Dong, X., 2021. Astrocytes contribute to pain gating in the spinal cord. Science
Advances 7, 1–13.

Yoon, H., Walters G., Paulsen A.R., Scarisbrick, I.A., 2017. Astrocyte heterogeneity across the brain
and spinal cord occurs developmentally in adulthood and in response to demyelination. PLoS
ONE 12(7): e018697.

Yoshioka, B.Y.M., Coutts, R.D., Amiel, D., Hacker, S.A., 1996. Characterization of a model of
osteoarthritis in the rabbit knee. Osteoarthritis and Cartilage 4, 87–98.

Zador, Z., Stiver, S., Wang, V., Manley, G.T., 2009. Role of Aquaporin-4 in Cerebral Edema and
Stroke. Handbook of Experimental Pharmacology 190, 159–170. https://doi.org/10.1007/978-
3-540-79885-9_7

Zając, A., Chalimoniuk, M., Gołaš, A., Lngfort, J., Maszczyk, A., 2015. Central and peripheral fatigue
during resistance exercise - A critical review. Journal of Human Kinetics 49, 159–169.
https://doi.org/10.1515/hukin-2015-0118

Zarkou, A., Lee, S.C.K., Prosser, L.A., Jeka, J.J., 2020. Foot and Ankle Somatosensory Deficits Affect
Balance and Motor Function in Children With Cerebral Palsy. Frontiers in Human
Neuroscience 14, 1–12. https://doi.org/10.3389/fnhum.2020.00045

Zhang, H., Boyette-Davis, J.A., Kosturakis, A.K., Li, Y., Yoon, S.-Y., Walters, E.T., Dougherty, P.M.,
2013. Induction of monocyte chemoattractant protein-1 (MCP-1) and its receptor CCR2 in
primary sensory neurons contributes to paclitaxel-induced peripheral neuropathy. Journal of
Pain 14, 1031–1044. https://doi.org/10.1016/j.jpain.2013.03.012.Induction

Zhang, H., Li, Y., De Carvalho-Barbosa, M., Kavelaars, A., Heijnen, C.J., Albrecht, P.J., Dougherty,
P.M., 2016. Dorsal Root Ganglion Infiltration by Macrophages Contributes to Paclitaxel
Chemotherapy-Induced Peripheral Neuropathy. Journal of Pain 17, 775–786.
https://doi.org/10.1016/j.jpain.2016.02.011

Zhang, J., Xiang, Q.S., Echeverry, S., Mogil, J.S., De Koninck, Y., Rivest, S., 2007. Expression of CCR2
in both resident and bone marrow-derived microglia plays a critical role in neuropathic pain.
Journal of Neuroscience 27, 12396–12406. https://doi.org/10.1523/JNEUROSCI.3016-07.2007

Zhang, J.-M., Song, X.-J., Lamotte, R.H., 1997. An in vitro study of ectopic discharge generation and
adrenergic sensitivity in the intact, nerve-injured rat dorsal root ganglion. Pain 72, 51–57.

1014

1015 **Table 1. Immunohistochemical Reagents**

Antibody Name	Immunogen Structure	Manufacturer Information Catalog / Lot /RRID	Concentration
Primary Antibodies			
Monoclonal mouse-anti-GFAP	Purified porcine spinal cord GFAP	EnCor Biotechnology cat. #MCA-5C10	1:500
Polyclonal rabbit-anti-CGRP	Synthetic peptide	Peninsula Laboratories cat. #T4032; Lot #A17601	1:1500
Lectins			
Biotinylated-isolectin B_4	Not applicable	Sigma Aldrich cat. #L2140; Lot #058M101V	1:2000
DNA Stains			
4',6-Diamidino-2-Phenylindole (DAPI)	Not applicable	Invitrogen cat. #D1306	1:5000

Secondary Antibodies			
Atto 488-goat anti-mouse IgG	Purified mouse IgG	Sigma-Aldrich, cat. # 62197	1:500
Avidin-conjugated with Alexa Fluor 594	Not applicable	Jackson ImmunoResearch cat. #016-580-084; Lot #151793	1:400
Goat-anti-rabbit IgG (H+L) Cross-Adsorbed, Alexa Fluor 488	Purified rabbit IgG	Invitrogen cat. #A11008; Lot #2284594	1:400

1016

Figure Legends

Figure 1: HI kits exhibit pain-like behavior in response to mechanical stimuli . **A:** Forepaw (left) and
hindpaw (right) von Frey scores at P5. Sham-operated controls (blue) have higher scores, indicating less
mechanosensitivity, while HI kits (red) display greater mechanosensitivity (lower scores). **B:** HI kits from
(A) were separated by whether they show motor deficits (motor-affected) or no deficits (motor-
unaffected). There was no difference in forepaw withdrawal threshold (left) but hindpaw withdrawal
(right) scores were higher in motor-affected HI kits, potentially indicating less sensitivity or impaired
ability to withdraw. Data are represented as mean ± SEM; N=64 sham and N=60 HI paws from N=32
sham and N=30 HI rabbits. **P<0.01, ****P <0.0001; Mann-Whitney tests.

Figure 2: HI kits display pain-like behavior in response to thermal stimuli **A:** Forepaw (left) and hindpaw
(right) withdrawal latencies during Hargreaves' testing at P5. Shorter latencies to paw withdrawal in
response to radiant heat are observed in HI kits compared to sham animals, indicative of enhanced
thermal sensitivity. **B:** HI kits from (A) were separated by whether they show motor deficits (motor-
affected) or no deficits (motor-unaffected). There was no difference in forepaw or hindpaw withdrawal
scores in motor-affected HI kits. Data are represented as mean ± SEM; N= 24 sham and N=36 HI paws
from N=12 sham and N=18 HI rabbits. ****P <0.0001; unpaired t-tests.

Figure 3: Astrogliosis is not present in HI rabbits at P5. **A**: Transverse lumbar spinal cord sections from
sham (left) and HI (right) rabbits immunostained for GFAP (green). Scale bar applies to both images. **B:**
Quantification of GFAP densitometry in spinal regions containing tracts/laminae of interest: dorsal
column (DC), superficial dorsal gray matter (SDGM), ventral spinothalamic tract (VST), and lateral
spinothalamic tract (LST). GFAP expression was unchanged in all ROIs in HI spinal cords (ns, not
significant; unpaired t-tests). Data are represented as mean ± SEM; N=7 rabbits per group.

Figure 4 Increased density and overlap of nociceptive primary afferent fibers in HI kits at P5.
Representative images of the lumbar dorsal horn of sham (**A-A''**) and HI (**B-B''**) kits immunolabeled for
calcitonin gene-regulated peptide (CGRP**; A, B, C**) or labeled by isolectin B-4 (IB4; **A', B', C'**). Note the
increased presence of CGRP-positive fibers in the deep dorsal horn in the HI but not the sham spinal
cord (**C-C''**). The proportional area of CGRP-positive tissue was significantly increased in the superficial
and deep dorsal horn of HI rabbits compared to sham (**D, G**), while the density and distribution of IB4-
positive afferents was unchanged between groups (**E**). The degree of overlap of these two nociceptor
populations can be viewed in the merged images (**A'', B''**) and was significantly increased in HI kits
compared to sham kits (**F**). Data are represented as mean ± SEM; individual data points represent each

animal (n=5 sham and n=6 HI kits) from averages of 3 separate images of left and right dorsal horns per

animal. **$P<0.01$, ***$P<0.001$; unpaired t-tests were used for statistical analysis.

1050

	HI vs. Sham
Tactile Hypersensitivity	↑
Thermal Hypersensitivity	↑
Astrocyte Reactivity	↔
CGRP Distribution	↑
IB4 Distribution	↔

Mechanical Nociception

A

Forepaws

Hindpaws

B

Forepaws

Hindpaws

Thermal Nociception

A

Forepaws

Hindpaws

B

Forepaws

Hindpaws

A

Sham

DC
SDGM
LST
VST

HI

GFAP

248 μm

B

Dorsal Column

ns

Superficial Dorsal Gray Matter

ns

Ventral Spinothalamic Tract

ns

Lateral Spinothalamic Tract

ns

GFAP Densitometric Signal (Relative Mean Gray Value)

Sham   HI
Sham   HI
Sham   HI
Sham   HI