1 **Influenza A virus undergoes compartmentalized replication *in vivo***
2 **dominated by stochastic bottlenecks**

3
4

5 Katherine A. Amato[1], Luis A. Haddock III[2], Katarina M. Braun[2], Victoria Meliopoulos[3], Brandi Livingston[3],
6 Rebekah Honce[3], Grace A. Schaack[1], Emma Boehm[2], Christina A. Higgins[1], Gabrielle L. Barry[2], Katia
7 Koelle[4], Stacey Schultz-Cherry[3], Thomas C. Friedrich[2,5], Andrew Mehle[1]
8
[1]Department of Medical Microbiology & Immunology, University of Wisconsin-Madison, Madison, WI
53706, USA
[2]Department of Pathobiological Sciences, University of Wisconsin School of Veterinary Medicine,
Madison, WI 53706, USA
[3]Department of Infectious Diseases, St. Jude Children's Research Hospital, Memphis, TN 38105, USA
[4]Department of Biology, Emory University, Atlanta, GA 30322, USA
[5]Wisconsin National Primate Research Center, Madison, WI 53715, USA

Abstract

Transmission of influenza A viruses (IAV) between hosts is subject to numerous physical and biological
barriers that impose genetic bottlenecks, constraining viral diversity and adaptation. The presence of
bottlenecks within individual hosts and their potential impacts on evolutionary pathways taken during
infection and subsequent transmission are poorly understood. To address this knowledge gap, we
created highly diverse IAV libraries bearing molecular barcodes on two independent gene segments,
enabling high-resolution tracking and quantification of unique virus lineages within hosts. Here we show
that IAV infection in lungs is characterized by multiple within-host bottlenecks that result in "islands" of
infection in lung lobes, each with genetically distinct populations. We performed site-specific inoculation
of barcoded IAV in the upper respiratory tract of ferrets and tracked viral diversity as infection spread to
the trachea and lungs. We observed compartmentalized replication of discrete barcoded populations
within the lobes of the lung. Bottlenecks stochastically sampled individual viruses from the upper
respiratory tract or the trachea that became the dominant genotype in a particular lobe. These
populations are shaped strongly by founder effects, with no evidence for positive selection. The
segregated sites of replication highlight the jackpot-style events that contribute to within-host influenza
virus evolution and may account for low rates of intrahost adaptation.

INTRODUCTION

The constant evolution of influenza viruses results in recurring seasonal epidemics and has the potential to initiate pandemics in the human population. Viral evolution occurs on multiple scales. Influenza A virus evolves rapidly on the global scale where population-level immunity positively selects new antigenic variants, necessitating reformulation of the seasonal influenza vaccine[1–3]. Yet, on smaller scales, variants with a predicted fitness advantage rarely rise to high frequency within a host or transmit to a recipient, even in the face of vaccine-induced immunity[4–7]. Within-host variation is low and genetic drift plays a large role[8]. To reconcile these observations, it is currently thought that positive selection drives more deterministic processes at the global scale while genetic drift shapes stochastic processes during local transmission[9].

Genetic drift of influenza virus between hosts is dominated by bottlenecks, neutral processes in which a small number of viral particles found a new infection, often resulting in dramatic reductions in the effective population size. In natural human infections, bottlenecks during airborne transmission of influenza A virus are exceedingly tight, with as few as 1-2 genomes founding infections in the recipient[8]. Similar results have been detected in animal models, where a very small proportion of intrahost variants successfully found new infections[10]. Under these conditions, genetic drift is expected to have a large impact, where diversity in the transmitted population is dramatically decreased with potential loss of beneficial variants as the population transits the narrow bottleneck[11]. Repeated experimental bottlenecks severely restrict viral fitness, and it has been suggested that transmission bottlenecks restrain jumps across host species and the global rate of influenza virus evolution[8,12,13].

Influenza virus infection occurs in heterogeneous cell populations within complex anatomical structures in a host[14]. Anatomical structures may help establish local sites with high multiplicity of infection that impact reassortment and complementation. Initial sites of infection are influenced by the tissue-specific distribution of sialic acid receptors and their topology[15–17]. Influenza virus replication can occur throughout the respiratory tract, but recent evidence suggests discrete locations contribute virus that is transmitted. Viruses in the upper respiratory tract, specifically those replicating in the soft palate or nasal epithelial cells, contribute most to the population that is transmitted in animal models[10,18,19]. Thus, it is the diversity at the site of transmission that is likely the most important for onward evolution. How viral movement in the respiratory tract and potential compartmentalization affect within-host evolution and ultimately transmission is poorly understood.

A clearly defined intrahost population structure is required to accurately model and predict influenza virus evolution and onward transmission of variants. Although error-prone genome replication generates influenza A viruses with distinct genotypes, the relatively low levels of naturally occurring variation within hosts do not provide sufficient information for fine-grained analysis of population dynamics. We overcame this limitation by introducing a neutral barcode of 10 random nucleotides into two segments of the influenza virus genome and creating rich viral populations with $\sim$0.6-3x10^5 uniquely quantifiable members. Using these barcoded viral populations, we capture soft selective sweeps in cell culture and show adaptive changes arose independently multiple times, yet only one lineage became dominant. Infection in ferrets revealed highly compartmentalized replication as virus migrated from the upper respiratory tract to the lung. Bottlenecks between sites led to stochastic sampling of individual viruses from the upper respiratory tract or the trachea that became the dominant lineage in lung lobes, while there was no evidence of positive selection. Thus, viruses infecting the lung do not constitute a large homogeneously mixed population, but rather multiple isolated populations that each undergo

Figure 1. Creation of molecularly barcoded influenza A virus populations. A) Molecular barcodes containing 10 randomized nucleotides were encoded downstream of the open reading frame in the *PA* and *HA* genes, shown as cRNAs. A registration mark was also added onto *HA* to distinguish unique barcode libraries. Sequences were repeated downstream of the barcode to maintain contiguous packaging signals required for replication, and silent mutations were introduced into the open reading frame to avoid direct repeats. B) Experimental overview where randomized barcodes were cloned into reverse genetics vectors followed by large-scale, parallel virus rescues to ensure unbiased barcode distribution. C) Optimized rescue plasmids enhance viral yield. Rescue efficiency was determined by measuring viral titers at the indicated times post-transfection with the standard 8-plasmid system, a consolidated 3-plasmid system, or the 3-plasmid system plus a vector expressing TMPRSS2. (data presented as mean of n = 3 ± sd. ANOVA with Tukey's post-hoc, * = p<0.05 relative to 8-plasmid rescue, # = p<0.05 relative to 3-plasmid system.)

79 bottlenecking events that can severely constrain population diversity and the potential for selection of fit
80 variants.

RESULTS

Large scale incorporation of barcodes avoids artificial bottlenecking of variants

Naturally occurring genetic diversity in influenza virus populations is poorly suited for high-resolution
characterization of population-level dynamics and cannot accurately enumerate the full spectrum of
individuals present. Moreover, genetic mutations may affect the fitness of the virus, biasing
representation of specific variants in the population and precluding their use as a neutral marker. To
better resolve and quantify the dynamics of the IAV population, we introduced dual molecular barcodes
into the genome of the influenza virus isolate A/California/07/2009 (H1N1; CA07) to create individual
viruses that are uniquely recognizable and quantifiable via deep sequencing **(Fig 1A)**. Barcodes of 10
randomized nucleotides were introduced onto the *HA* and *PA* segments that were subsequently used for

high-efficiency virus rescue **(Fig 1B)**. HA is under selective pressure as the viral attachment protein and
principal target of neutralizing antibodies. We might therefore expect barcodes embedded in *HA* to
increase or decrease in frequency as a result of selection on this segment. Conversely, PA is subject to
less intense selection pressure, and barcodes on this segment may be expected to better represent total
population size. For both *HA* and *PA*, barcodes were encoded between the ORF and the UTR.
Packaging and bundling signals were duplicated downstream of the barcode to ensure proper gene
replication and virion formation[20,21]. The utility of the system was further increased by using the PA
reporter construct that co-transcriptionally expresses Nanoluciferase (PASTN)[20]. Finally, the *HA* segment
encodes an additional "registration mark," six nucleotides creating either NheI or PstI restriction sites,
that allows us to index and identify separate libraries of *HA* variants.
101
The limited efficiency of virus rescue can introduce artificial bottlenecks[22]. To increase efficiency and
recover a larger, more diverse barcode population, we reduced the number of individual plasmids
needed for virus rescue[23]. We combined the 6 non-barcoded segments of CA07 IAV onto a single
plasmid, reducing the entire reverse genetics system from 8 to 3 plasmids **(Fig 1C)**. Rescue titers of the
3-plasmid system increased > 200-fold compared to the 8-plasmid system. Titers were further increased
by co-expressing transmembrane protease serine 2 (TMPRSS2) during virus rescue, which cleaves and
activates HA. We performed 120 parallel virus rescues, pooled the resultant supernatants, and passaged
them at high representation to ensure unbiased barcode distribution (**Fig 1B**). NheI and PstI registration
marked libraries were prepared independently.
111

Viral barcodes reveal selective sweeps *in vitro*

Deep sequencing revealed ~180,000 unique *PA* barcodes and over 238,000 *HA* barcodes in each of our
original virus libraries **(Fig 2A)**. Given that each library has an invariant registration mark, we could
measure the fidelity of the quantification pipeline by assessing the number of reads that do not match the
predicted registration mark sequence. Over 99% of mapped reads perfectly matched the appropriate
registration marks on *HA*, with the majority of those that did not match differing by a single nucleotide
from the intended registration mark, indicating the neutrality of the registration mark sequence and
overall high fidelity of our sequencing pipeline. Viral populations were characterized by richness, the total
number of unique lineages present, Shannon's diversity, a measure that considers both the presence
and relative abundance of a lineage, and evenness, a parameter that compares the frequencies of all
lineages in the population to that of a theoretically evenly distributed population[24–26]. Barcode
enumeration for viruses recovered from the transfected cell supernatant (i.e. passage 0, P0) revealed
rich and highly diverse populations with evenly distributed lineages (**Fig 2A-B**). However, a single
passage to create amplified P1 stocks resulted in highly skewed populations in which individual lineages
dominated the population, evidenced by dramatic drops in Shannon's diversity and evenness (**Fig 2B**).
The dominant barcodes in each library represented 64-77% of the population. Neither of these barcodes
was dramatically over-represented in the plasmid or P0 stocks. In fact, the most abundant lineages in the
P0 rescue stocks did not become the most prevalent in our amplified stocks.
130
131 Changes in population composition can occur through drift or selection. We passaged virus at a large
132 effective population size, minimizing the propensity for drift and raising the possibility that the expanding
133 lineages became dominant because they had acquired a selective advantage. Whole genome
134 sequencing of the P1 stocks identified the single nucleotide variants (SNVs) A540G (numbering based
135 on cRNA) in almost 50% of reads from NheI registration mark viruses and A543G in almost 70% of reads

Figure 2. Single viruses seed selective sweeps in *HA*. A) Creation of two dual-barcode libraries with distinct registration marks and uniquely addressable members. B) Frequency of each lineage as a fraction of total population size. Colors indicate unique barcode identities. C) Whole genome sequencing identifies adaptive variants in *HA*. Individual single nucleotide variant (SNV) frequencies are indicated at each nucleotide position in a concatenated IAV genome for each library. D) Long-read sequencing reveals selective sweeps by linking adaptive mutants in HA to single dominant barcodes. The frequencies of mutations coding for the indicated change that are linked to the dominant barcode are shown for both libraries (left). The frequencies of all barcodes linked to the adaptive glutamic acid variant are indicated (right), with the dominant barcode in each library colored as in B.

from PstI marked viruses (**Fig 2C**), mirroring the abundance of the dominant barcode. These mutations
code for HA K153E and K154E (H1 numbering), respectively. Importantly, these changes in HA had
previously been identified as adaptations that provide a growth advantage to CA07 in cell culture[27]. Long-
read sequencing showed that the dominant barcode was linked to the adaptive mutations (**Fig 2D, left**).
62-75% of reads containing the dominant barcode also encoded the adaptive mutation. Similarly, the
adaptive mutation was primarily linked to the dominant barcode (**Fig 2D, right**). However, up to 27% of
reads encoding these adaptive variants were associated with different very low-frequency viral lineages.
These data are consistent with a soft selective sweep where *HA* A540G or A543G arose on multiple
genetic backgrounds, even though only one lineage ultimately became the most abundant, possibly
suggesting clonal interference. These observations demonstrate that our barcoded viruses capture
lineage dynamics and selective processes at extremely high resolution.

Pre-adaptation creates large and diverse viral libraries

To create diverse libraries without tissue-culture-induced skewing, we made new libraries on a "pre-adapted" HA K153E background. The libraries contained at least 57,000 unique members (**Fig 3A**). Multicycle growth curves of HA K153E NheI- or PstI-marked barcoded viruses were indistinguishable (**Fig 3B**). Insertion of the barcode cassette and the *PA* reporter gene slightly reduces titers when compared to HA K153E alone, consistent with prior results[20,28]. The new libraries were highly diverse (Shannon's Diversity Index (H') >8.8 for all barcodes) and evenly distributed; the vast majority of lineages were present at low frequency, and no single lineage in our amplified stocks was present at a frequency above 0.6% for HA, or above 1.2% for PA barcodes **(Fig 3C)**. We utilized the HA-NheI stock for extensive quality control of our barcode enumeration pipeline. Libraries were prepared from four independent RNA extractions and sequenced. Lineage frequencies from the four replicates showed strong correlation (**Supp Fig 1A**). Over 94% of all sequence reads were shared across the four replicates (**Supp Fig 1B**). However, when considering barcode identity, only ~37% of lineages were identified in all replicate runs (**Supp Fig 1C**). This apparent discrepancy is due to variable detection of extremely low-frequency lineages. The frequency of an individual lineage was correlated with the number of replicates in which it was detected (**Supp Fig 1D**). These data highlight the complexity of our viral populations and our ability to reliably detect individual members. Replicate sequencing of the *PA* barcode and the HA-PstI stock produced similarly well-correlated results (**Supp Fig 1E, 2A-B**). No additional SNVs were detected at high frequency in our libraries, while K153E remained fixed **(Fig 3D)**.

Figure 3. Generation of large and evenly distributed dual-barcoded virus libraries. A) Properties of the A/CA/07/2009 HA-K153E PASTN virus libraries. Data are from a single sequencing replicate; see Supplemental Figs 1-2 for additional analyses. A) Unique barcodes identified in each population. B) Multistep growth curves of dual-barcoded PASTN libraries compared to the parental strain A/CA/07/2009 HA-K153E. Viral titer was measured by plaque assay (n=3 ± sd, ANOVA with Tukey's post hoc, * = p < 0.05 and ** = p < 0.01 compared to the parental). C) Frequency distribution of unique barcodes on *HA* and *PA* binned in 0.1% increments. D) Whole genome sequencing was performed and SNV frequencies are indicated at each nucleotide position for each library.

Rich and diverse populations replicate in mice

The dual barcoded virus libraries provide a key opportunity to quantify population dynamics *in vivo*. Mice are frequent models for influenza virus replication, pathogenesis, and immune response[29]. Mice were inoculated with the CA07 PASTN-barcoded virus libraries containing either barcoded HA-K153E populations or a non-barcoded HA-K153E control. Weight loss was similar for all conditions **(Fig 4A),**

Figure 4. Replication of diverse virus populations in mice. A) Mice were inoculated with 10^5 TCIDGlo$_{50}$ of virus containing barcoded PASTN with either HA-K153E or barcoded variants and body weights were measured daily. Half of the mice were sacrificed at 3 dpi. Data presented as mean ± sd for n=6 1-3 dpi, and n=3 for 4-6 dpi. B) Viral titers in mouse lungs harvested at 3 and 6 dpi were determined by plaque assay (left) or TCIDGlo$_{50}$/mL (right). Data presented as mean of n=3 ± sd. C) Barcodes in the viral stock and mouse lungs were quantified and the frequency of clustered HA barcodes as a fraction of total population size is indicated. Each color in a series represents an individual barcode cluster. Total number of unique barcode clusters per sample and the most abundant barcode with its frequency are listed at right. D) Shannon's diversity index for viral populations in the stock and mouse lungs (mean of n=3 ± sd). E) Venn diagrams displaying the number of unique and shared lineages within each mouse for NheI and PstI libraries.

172 and viral titers in the lungs were not significantly different at either 3 or 6 dpi (**Fig 4B**). Thus, introduction
173 of barcodes onto *HA* does not compromise replication *in vivo*.
174
Deep sequencing of mouse lung homogenates revealed that the majority of mice harbored diverse viral
lineages (**Fig 4C-D**). Mouse infections were characterized by a high richness, where approximately one
third of lineages present in the stock were detected in the lungs at 3 dpi. The frequency of lineages in the
NheI-marked stock was moderately predictive of their abundance in the mouse at 3 dpi (Pearson's R >
0.57)(**Supp Fig 3A**). For example, the most abundant lineage in the HA-NheI stock was also the most
abundant lineage in two mice at 3 dpi (**Fig 4C, Supp Fig 3A**). However, there are notable exceptions in
which low-frequency lineages in the stock rose to high relative abundance in the mouse (**Supp Fig 3A**).
In addition, we cannot exclude the possibility that some of our barcodes are derived from residual viruses
from the inoculum that did not initiate a productive infection. Mice infected with PstI-marked libraries
showed less correlation between the inoculum and lungs at 3dpi (**Supp Fig 3A**). For all viral libraries, the
titers decreased from 3 to 6 dpi, as did richness and overall diversity (**Fig 4C-D**). In an extreme example,
a single lineage in mouse 11 rose to over 40% prevalence at 6 dpi (**Fig 4C, Supp Fig 3B**). The
correlation between lineage frequency in the stock and the mouse lung was greatly diminished at 6 dpi
(**Supp Fig 3**). The lineage identities are highly heterogeneous among mice, with a small fraction shared
across animals, suggesting that barcodes themselves are not under selection *in vivo* (**Fig 4E**). Together,
these data show virus populations replicating in mice approximate the diversity present in the inoculum,
as might be expected from a high-dose challenge. Within-host richness decreases as the infection is
resolved, with lineages lost as the overall population size decreases.
193

Within-host bottlenecks result in compartmentalized replication in the ferret lower
respiratory tract

Ferrets are often considered the "gold standard" infection model with lung physiology, sialic acid
distribution, pathogenesis, and transmission capacity that are all similar to humans[30,31]. We intranasally
inoculated 3 ferrets with the HA-K153E dual barcoded library containing the NheI registration mark. We
used a site-specific inoculation strategy in which the inoculum is retained in the upper respiratory tract
without unintentional introduction into the trachea or lower respiratory tract[32], allowing us to track the
natural movement of virus. Ferrets were monitored daily for signs of infection with nasal washes obtained
1, 3 and 5dpi. Ferrets exhibited slight weight loss over the course of infection (**Fig 5A**), consistent with
prior work[28]. Similarly, we detected high viral titers in nasal washes 1 dpi that declined over time (**Fig 5B**).

Sequencing revealed rich and complex viral populations in the nasal washes of ferrets, consistent with a
high dose inoculation in this compartment (**Fig 5C-E**). Frequency trajectory plots showed heterogeneous
and well mixed viral populations undergoing little, if any, selection or bottlenecks in the upper respiratory
tract (**Fig 5C**). This is consistent with a highly diverse population with generally even lineage distribution,
reflected by a high Shannon's diversity index and evenness (**Fig 5D**). Approximately 19,000-22,400 unique
lineages were detected in each nasal wash 1 dpi, and richness remained high over time, with at least
16,800 lineages at 3 dpi and 10,000 lineages at 5 dpi (**Fig 5E**). Sequencing of the barcode on *PA* revealed
remarkably similar lineage dynamics. The fact that frequencies of *HA* and *PA* barcodes moved in parallel
is perhaps surprising given that these are unlinked genes and that influenza viruses can undergoes
frequent reassortment[33].

Figure 5. Population diversity is reduced when influenza virus moves from the upper to the lower respiratory tract in ferrets. A) Ferrets were inoculated with a site-specific intranasal dose of 10^5 PFU of dual-barcoded A/CA/07/2009 HA-K153E PASTN virus containing the NheI registration mark. Ferret weight was monitored daily. B) Viral titer in ferret nasal washes were determined by TCIDGlo$_{50}$. C) Changes in frequency for HA (top) and PA (bottom) barcodes present in nasal wash samples over the course of infection. D) Shannon's diversity index and evenness for the indicated viral populations. (cont'd on next page)

Intranasally inoculated viruses spread throughout the respiratory tract by 5 dpi. Low levels of infectious
virus were present in the trachea of ferrets 35 and 36, whereas deep sequencing detected IAV genetic
material in all trachea samples (**Fig 5F-G**). Compared to the nasal wash, population richness dropped
significantly, with 1250 or fewer lineages present in the trachea. Lineage distribution differed in the trachea
of each animal, ranging from a more diverse population in ferret 34 to a largely homogenous population in
ferret 36, in which a single lineage accounted for 71% of the population (**Fig 5G**). Our site-specific
inoculation requires virus replication in the upper respiratory tract prior to movement into the trachea or
lower respiratory tract. We therefore used lineages present in nasal washes at 3 dpi as a comparator for
populations in the trachea and lungs at 5 dpi. Lineage frequency is poorly correlated between nasal washes
and the trachea (**Supp Fig 4A**). Migration into the trachea is associated with a drastic reduction in richness,
a poor correlation with the source, and skewed distribution of the resultant population, indicating that viral
population bottlenecks between compartments and founder effects may play a role during the seeding of
the trachea from the upper respiratory tract.

Virus also spread to the lungs of infected animals (**Fig 5H**). Moderate titers were detected in all five lung
lobes in ferret 34, even though infectious viral titers in the trachea were below the limit of detection for this
animal. Infectious virus was also detected in all lung lobes for ferret 36, but only the middle right lobe for
ferret 35. Thus, while virus must traverse the trachea to access the lungs, the presence of virus in the
trachea 5 dpi was not predictive of the extent of spread in the lungs. Moreover, populations in lung lobes
had higher lineage richness than that detected in the trachea at 5 dpi; the vast majority of lineages in the
lung were not detected in the trachea (**Supp Fig 4B**). This raises the possibility that population richness in
the trachea at 5 dpi shrank significantly from the populations at earlier time points that may have seeded
infection in the lung, or that virus can transit through the trachea to directly inoculate the lung.

Lineage analysis revealed heterogeneous populations of barcodes in each of the distinct lobes (**Fig 5I**).
Over 25,000 lineages were detected across the five lobes in ferret 34. However, each lung lobe of ferret
34 had a different dominant barcode sequence, and when this same barcodes was detected in other lung
lobes its frequency varied. The only infected lung lobe of ferret 35 had very low viral titers and was
dominated by a single lineage reaching ~95% abundance (**Fig 5H-I**). Ferret 36 yielded another outcome,
where the same lineage was dominant at a frequency of 35-98% in each lobe except for the upper left,
which maintained a richer and more diverse population. For all animals, individual lobes showed reduced
diversity and evenness compared to the virus population in the upper respiratory tract (**Fig 5D**). In the two
animals in which virus was detected in all lobes, the upper left lobe consistently had the highest richness,
diversity, and evenness. These data suggest that anatomical features associated with each lobe, such as
tracheal bifurcation patterns or bronchus size[32], may affect patterns of virus establishment and population
size. The differences in composition of the populations in each lung lobe was determined. Bray-Curtis
dissimilarity assessment revealed highly compartmentalized replication, in which each lung behaved as a
distinct anatomical "island" with a unique population composition (**Fig 6A**). Four lobes in ferret 36 showed
lower dissimilarity, as they were dominated by the same lineage, yet the non-dominant lineages still
contributed unique populations to each lobe. Only 167 lineages were common to all lobes of ferret 34, and
only 42 in ferret 36 (**Fig 6B**).

The drastic reduction in richness and diversity between the upper respiratory tract and the lung lobes suggested populations were subject to selection or bottlenecks. *HA* sequencing from select lung lobes for each animal revealed that no intrahost SNVs (iSNVs) surpassed the 3% threshold set for accurate estimate of iSNV frequency, excluding the possibility that barcode frequency changes were driven by selection of a linked adaptive variant (**Fig 6C**). We therefore further assessed the possibility of viral population bottlenecks in driving the observed reduction in richness and diversity. To this end, we developed a simple mutlinomial model to estimate bottleneck sizes (N_b) as virus transmits from the nasal wash at 3 dpi to lung lobes at 5 dpi (**Fig 6D**). Specifically, this model estimated N_b using data on the lineages and their frequencies in the 3 dpi nasal wash and on the number of lineages observed in a focal 5 dpi lung lobe. The model yielded maximum likelihood estimates of N_b of ~66,000 (ferret 36) and ~217,000 (ferret 34) virions for the upper left lobe, and lower estimates for the other lobes (**Fig 6D**), with maximum likelihood estimates spanning between ~200 and ~15,000 virions. The larger bottleneck size between the nasal wash and the upper left lobe compared to that for the other lung lobes was consistent with higher levels of viral lineage diversity detected in the upper left lobe compared to the others. Nonetheless, the bottleneck size estimates still appeared unexpectedly large to us, given the drastic reductions in richness and diversity observed in Fig 5D. We therefore forward simulated mock transmission events from the nasal wash using our

Figure 6. Lung lineage diversity and population bottlenecks are dominated by stochastic pressures. A) Pair-wise Bray-Curtis dissimilarity for lung lobes within single animals. B) The number of barcode lineages common and unique to lung lobes are illustrated. C) Whole-HA sequencing of virus in lung homogenates identified multiple iSNVs, but not rose above the 3% threshold set for accurate estimation of iSNV frequency. D) Transmission bottlenecks (N_b) from nasal wash 3 dpi to each lung lobe at 5 dpi were calculated by maximum likelihood estimation and plotted ± 95% confidence intervals.

estimated bottleneck sizes under the assumption that virions were transmitted from the nasal wash to a focal lung lobe at a single timepoint. These simulations predicted a high degree of similarity between lineage frequencies in the viral populations sampled in the nasal wash and lung lobes (**Supp Fig 5**). They further predicted levels of viral genetic diversity and richness in the lung lobes that were significantly higher than those observed (**Fig 5D,I**). These initially seemingly inconsistent results could be parsimoniously explained under a model where many virions are transmitted between the nasal wash and each of the lung lobes over the course of infection, but at multiple timepoints rather than at a single one.

Considering the lung as a whole for ferrets 34 and 36, a substantial number of lineages were shared between the nasal wash and lung (**Fig 7A**). But, this appeared to be largely driven by the rich and diverse population in the upper left lobe, as this overlap was largely lost when lobes were considered individually. Many of the dominant lineages in a lung lobe were poorly represented in the nasal wash, such as that in the upper right lobe of ferret 34 and the dominant lineage shared in 4 lobes of ferret 36 (**Fig 7B, C, Supp Fig 6A, C**). Lineage enrichment in the lung samples compared to nasal washes revealed many lineages that were unique to nasal washes or lung lobes (**Sup Fig 6A-C**). These data show that treating the lung

Figure 7. Compartmentalized infections establish replication islands within the lung. A) Overlap between barcodes in the nasal wash 3 dpi and those present in distinct lobes, or composite data for all barcodes in the lung. B-D) The frequency of barcodes within the nasal wash are compared to frequencies within individual lobes for B) ferret 34, C) ferret 35 and D) ferret 36. Red arrowheads highlight dominant barcodes with frequency > 30%, which for ferret 36 is the same barcode that dominated in multiple lobes of the lung. Note that barcodes unique to the nasal wash or lung are not plotted here and can be found in Supplemental Figure 6.

as a whole can lead to very different and misleading population structures, highlighting the importance of
assessing each lobe individually.

The same lineage dominated in four lobes of the lung for ferret 36, whereas underlying diversity and a lack
of overlap in each population still suggests each lobe is seeded by distinct transmission events (**Fig 5I, 6B
7D**). In this animal, enrichment for the dominating lineage appeared to occur in the trachea (**Fig 5G, Supp**

Fig 4A, 7A). This may be a general trend, as high-frequency lineages in the trachea were often over-
represented in the lobes (**Supp Fig 7A-B**). Combined, the large differences in lineage identity and
frequency show that each lobe is independently seeded, with little mixing between compartments.
Moreover, comparisons across compartments suggest each lobe presents a bottleneck and that genetic
drift repeatedly reshapes the population as virus traverses the respiratory tract.

DISCUSSION

313
A quantitative understanding of population dynamics is crucial for determining how evolutionary forces
shape viral populations within and between hosts. Error-prone replication by influenza virus generates
genetic diversity within an infected host. However, the full extent of that diversity does not survive
transmission to a new host. Transmission events between hosts involve stringent bottlenecks, with only
2-6 viral genomes founding the next infection during aerosol transmission[8,10,34]. Here we show that
influenza virus also faces multiple bottlenecks within a host as it seeds different compartments. Rich and
diverse populations in the upper respiratory tract were stochastically sampled as virus transited into the
lower respiratory tract, introducing strong founder effects that skewed the resultant population. Our data
suggest a scenario in which repeated within-host bottlenecks severely reduce diversity, richness, and
evenness, resulting in distinct, compartmentalized "island populations" in each lobe of the lung.
324
Our results show that influenza virus evolution within a host involves multiple bottlenecks, tempering the
impact of positive selection and likely creating additional barriers for host-adaption and onward
transmission. This could involve multiple physical bottlenecks as the virus sequentially infects distinct
tissues, repeated exposure to the same bottleneck if transmission to a new tissue occurs more than
once, or a combination of the two. And while we considered entire lung lobes, it remains possible that
focal infections within tissues may further subdivide viral populations. In all scenarios, bottlenecks
decrease the richness and overall population size and are expected to increase the strength of genetic
drift[35]. These processes combine to constrain the diversity and evolution during influenza virus infection,
which slows the fixation of adaptive variants within a host.
334
Immune pressure positively selects antigenically advanced influenza virus variants on the global scale[1].
Yet, these same variants rarely emerge in an acutely infected host, where infections are dominated by
purifying selection and low intra-host diversity[4,5,7,8,36]. The apparent disconnect between influenza virus
evolution on the global versus individual scale is incompletely understood. Our findings reveal that
intrahost bottlenecks are major contributors to the limited evolution detected in an infected individual.
This may partially explain why highly pathogenic avian H5N1 strains have not yet acquired the ability to
transmit within people. H5N1 strains infect the lower respiratory tract in humans, where the preferred
α2,3-sialic acid receptors are more abundant[15,16]. We saw little evidence for mixing between
compartments in the lower respiratory tract. Thus, while only a few mutations are needed to confer
airborne transmission to H5N1 strains in experimental settings, the spatial structure and
compartmentalization of infection in humans may impose bottlenecks that prevent fixation of variants and
migration to the upper respiratory tract, let alone transmission to a new host[37,38].
347
Intrahost bottlenecks are common features during viral infections. Physical barriers establish bottlenecks
as enteroviruses escape the gut in mouse models[39,40]. Poliovirus then faces another bottleneck related to
IFN responses that restrict diversity as virus invades the central nervous system[39,41,42]. Our data do not

reveal whether the bottlenecks we describe for influenza virus are due to physical barriers, innate
immune response, or other factors. Similar forces shape evolution in arthropod vectors. Repeated
bottlenecks winnow diversity as virus moves between different tissues within mosquitoes for West Nile
virus, Venezuelan equine encephalitis virus and Zika virus[35,43,44]. Error-prone replication then repopulates
diversity in the new sites of replication. Repeated bottlenecks, such as those we have identified during
intra-host dissemination, decrease diversity and may reduce fitness. Yet, they also act to purge
deleterious mutations and facilitate escape from local fitness maxima and exploration of additional
evolutionary space. The combined effects of these bottlenecks shape the kinetics of intra-host influenza
virus evolution.
360
In summary, we demonstrate the existence of multiple bottlenecks during dissemination of influenza virus
throughout the respiratory tract. We posit that these bottlenecks contribute to the limited impact of
positive selection on intrahost evolution. Moreover, they provide additional barriers to initial cross-species
transmissions and sustained transmission once a virus spills over into a new host. Coupled with the
stringent bottlenecks that occur during inter-host transmission, our results help explain the stochastic
nature of influenza virus evolution at the local scale.
367

METHODS

Cells

370 MDCK cells (ATCC), MDCK-SIAT1-TMPRSS2 cells[45] and 293T cells (ATCC) were maintained in
371 Dulbecco's modified Eagle medium (DMEM) supplemented with 10% heat-inactivated fetal bovine serum
372 (Atlanta Biologicals), 100 μg/mL streptomycin, and 100 U/mL penicillin at 37°C and 5% CO_2.

Generation of pHW2000-all-ΔPA-ΔHA A/California/07/2009

374 Bidirectional cassettes for CA07 PB2, PB1, NP, NA, M, and NS gene segments were sequentially
375 amplified and inserted into a pHW2000 vector via Gibson assembly. The resulting 17.3 kb plasmid was
376 sequence verified. Annotated plasmids sequences can be found at
377 https://github.com/mehlelab/barcoded_flu_analysis

Generation of barcoded PA

Barcodes were inserted into the previously described PASTN rescue construct that expresses a
polyprotein encoding PA and Nanoluciferase separated by the 2A cleavage site[20,46]. The barcode was
originally synthesized (IDT) as a single-stranded DNA oligo containing ten random nucleotides flanked by
sequence to enable cloning immediately downstream of the Nanoluciferase open reading frame
(Supplementary Table 1). The oligo was amplified in 25 individual low-cycle PCRs using Q5 polymerase
(NEB).The amplicons were pooled and gel purified prior to cloning into a modified PASTN plasmid using
50 individual ligation reactions. Ligations were transformed into Mach1 competent *E. coli* (Thermo
Fisher), plated on LB agar supplemented with 200 μg/mL ampicillin, and grown overnight at 37°C to yield
220,000 transformants, indicative of a theoretical maximum library size. Colonies were scraped off
plates, collected in 750 mL liquid LB with ampicillin, and grown for 3 hr at 37°C. DNA was then purified
using the Zymo Midiprep kit to create the plasmid library. Plasmid stocks were deep sequenced to
determine library size and diversity.

Generation of barcoded HA

HA from A/California/07/2009 was cloned into the pHW2000 rescue plasmid. Silent mutations were introduced into the final 25 amino acids of the open reading frame and 80 nt from the 3' end of the open reading frame were repeated downstream to recreate a contiguous packaging signal[21]. Initial libraries utilized the native HA sequence, whereas subsequent libraries included the tissue-culture adaptive K153E (nt A540G in cRNA) mutation[27]. A single-stranded oligo (IDT) was synthesized containing a randomized 10-nt barcode as well as a 6-nt registration mark, either GCTAGC (NheI) or CTGCAG (PstI) (Supplementary Table 1). This registration mark is a conserved and identifiable region for all the individuals within a given library. Barcodes were amplified in 50 low-cycle PCRs and cloned following the same strategy as for PA libraries. Approximately 60,000 transformants were obtained with the NheI registration mark, and 60,000 for the PstI registration mark. Plasmid stocks were deep sequenced to determine library size and diversity.

Rescue of dual-barcoded virus libraries

Barcoded CA07 virus libraries were rescued via reverse genetics using the pHW2000-all-ΔPA-ΔHA and PA and HA barcoded plasmids described above. Briefly, 293T were forward transfected with 2.7 µg pHW2000-all-ΔPA-ΔHA, 450 ng pHW2000-PASTN-barcode library, 450 ng pHW2000-HA-barcode library (NheI or PstI variant), and 400 ng pHAGE2–EF1αInt–TMPRSS2–IRES–mCherry-W[45] in a 6-well format. Plasmids were combined with 200 µL jetPRIME Buffer and 8 µL jetPRIME reagent (Polyplus) per well. 120 independent transfections were performed per viral library. 24 hr post-transfection, media was removed and cultures were overlaid with MDCK-SIAT1-TMPRSS2 cells in OptiVGM (OptiMEM supplemented with 0.3% bovine serum albumin, 100 µg/mL calcium chloride, 100 µg/mL streptomycin, and 100 U/mL penicillin). Rescue viruses were harvested 48-72 hr later and pooled based on their registration mark. Viruses were amplified on 20, 15 cm dishes of MDCK-SIAT1-TMPRSS2 cells in OptiVGM for 66 hr. Viruses were pooled based on their registration mark and cellular debris was removed by centrifugation. Viral titers were determined by plaque assay and TCID50Glo assays on MDCK and MDCK-SIAT-TMPRSS2 cells[47].

Library preparation for barcoded amplicons.

Viral RNA was extracted from all samples using the Maxwell RSC Viral Total Nucleic Acid Purification Kit (Promega) according to the manufacturer's instructions. RNA was subjected to DNAse treatment using the TURBO DNAse (Invitrogen), and reverse transcribed in 20µl using the SuperScript IV VILO master mix (Invitrogen) with PA and HA gene segment-specific primers (Supplementary Table 1). DNA amplicons for HA and PA gene segments with partial sequencing adapters were generated via PCR amplification of cDNA using the Phusion High-Fidelity DNA Polymerase (New England BioLabs) and gene segment specific primers (Supplementary Table 1). To minimize technical bottlenecks during library preparation, reverse transcription and the first PCR amplification for all samples were performed in triplicate and pooled together prior to DNA purification. PCR products from mouse samples were gel purified whereas PCR products from ferret samples were purified with paramagnetic beads using the AMPure XP for PCR Purification kit (Beckman Coulter). Purified PCR products were used in a second PCR reaction for incorporating sample-specific 5'-end indexes and additional Illumina sequencing adapters (Supplementary Table 2). Final PCR products were gel purified and individual DNA concentrations were determined with the Qubit dsDNA High Sensitivity Assay Kit on the Qubit Fluorometer (Invitrogen). Samples were quality controlled using the Bioanalyzer High Sensitivity DNA Analysis Kit and the Agilent 2100 Bioanalyzer (Agilent). All samples were prepared and sequenced in

434   technical replicate. Detailed protocols and code available at
435   https://github.com/mehlelab/barcoded_flu_analysis.

Library preparation for whole-genome sequencing.

Library preparation was similar to our prior approaches[48]. Briefly, viral RNA was extracted from all
samples using the Maxwell RSC Viral Total Nucleic Acid Purification Kit (Promega) according to the
manufacturer's instructions. RNA were subjected to DNAse treatment using the TURBO DNAse
(Invitrogen), and reverse transcribed in 20µl using the SuperScript IV VILO master mix (Invitrogen) with
the Uni12 primer (Supplementary Table 1) that targets conserved ends of all gene segments. Segments
were amplified by PCR with gene-specific primers (Supplementary Table 1), gel purified, and DNA
concentrations were determined using the Qubit dsDNA High Sensitivity Assay Kit on the Qubit
Fluorometer (Invitrogen). 1 ng of each segment was pooled and used as input for the Nextera DNA
Library Prep kit where samples were tagmented and indexed according to the manufacturer's
instructions. Tagmented and amplified products were purified with AMPure XP paramagnetic beads for
PCR Purification kit (Beckman Coulter) in two consecutive steps (0.5x and 0.7x) and were quantified
using Qubit dsDNA high-sensitivity kit (Invitrogen, USA). Sample quality control was performed using the
Bioanalyzer High Sensitivity DNA Analysis Kit in the Agilent 2100 Bioanalyzer (Agilent). All samples were
prepared and sequenced in technical replicate. Detailed protocols can be found in
https://github.com/haddocksoto/bcflu_protocols.

Deep sequencing

Amplicon and whole-genome libraries were sequenced on the Illumina MiSeq system using the MiSeq
Reagent Kit v2-500 and v3-600, respectively (Illumina). Amplicon and whole-genome samples that
passed quality control were pooled in a 4nM library with nuclease-free water. 5 µl of the 4nM library pool
was denatured with 5 µl 0.2N of NaOH and diluted using the HT1 Hybridization Buffer (Illumina) to a
concentration of 8 pM for amplicon samples and 10 pm for whole-genome samples. A PhiX library was
prepared similarly, and added at 30% of the input for amplicon sequencing and 1% for whole-genome
libraries. Samples were loaded on the respective MiSeq cartridge and paired-end sequencing reads were
generated (Illumina).

Sequencing data analysis for barcoded amplicons

We generated a custom bioinformatic pipeline to process raw FASTQ files and quantify barcode
frequencies (https://github.com/mehlelab/barcoded_flu_analysis). Briefly, raw FASTQ paired reads were
demultiplexed, merged and aligned to a custom amplicon reference containing barcode and registration
mark regions as strings of N's (BBMap Tools v38.87). Reads with lengths the same as the average insert
size were aligned, sorted and indexed with Samtools (v1.11, htslib v1.11). BAM files with aligned reads
were processed and trimmed to the region containing the registration mark and/or the barcode sequence
using command line tools (Seqtk v1.3, Bash v3.2.57).
469
Sequencing of the invariant registration mark was used to benchmark fidelity of our sequencing. Over
99.1% of reads were perfect matches, where most differences were the result of a single nucleotide
change from the expected sequence. Therefore, to correct for amplification and/or sequencing errors that
may inflate the number of unique barcodes, we used UMI-tools (v1.1.1) to generate consensus barcode
clusters with read counts via the adjacency network-based clusterer method[49]. Parental barcodes and
their apparent mutational offspring were clustered prior to enumeration and cluster frequencies were

generated and visualized via a custom Python pipeline (Python v3.8.5, Pandas v1.1.3, Matplotlib v3.3.2,
Numpy v1.19.2) or in Prism 9. Manual inspection of sequences in the raw FASTQ files confirmed these
as *bona fide* barcodes and not the product of misalignment.

Sequencing data analysis for whole gene segments

We generated a custom bioinformatic pipeline to process raw FASTQ files and determine single
nucleotide variants (SNV) from our barcoded viral samples
(https://github.com/mehlelab/barcoded_flu_analysis). Briefly, raw FASTQ paired reads were
demultiplexed and merged with a minimum overlap region of 30 nucleotides using bbmerge (BBMap
Tools v38.87). We mapped reads to the full IAV genome using a Burrows-Wheeler alignment (BWA
v.0.7.17). Using Samtools (v1.11, htslib v1.11), we sorted our aligned reads and called variants using
LoFreq (v2.1.5) with a minimum coverage of 500 reads, base call quality of at least 30 and a frequency
exceeding 0.03 (3%). Our reference IAV genome included barcode insertions in *PA* and *HA*, registration
marks on *HA* as a strings of 10 N's, NanoLuc inserted in *PA*, and repeated packaging signals. SNVs
were annotated using SnpEff (v5.0e) to determine the impact of each variant on the amino acid
sequence, and the resulting variant call format (VCF) files were manipulated using bcftools (v1.11) to
transform into user-defined formats. Plots were generated using custom bioinformatic pipelines in Python
language (Python v3.8.5, Pandas v1.1.3, Matplotlib v3.3.2, Numpy v1.19.2).

Long-read sequencing of *HA* and analysis

Viral RNA was made compatible for sequencing on an Oxford Nanopore Technologies instrument using
the 1D PCR Barcoding Kit. Briefly, viral RNA was extracted using the Maxwell RSC Viral Total Nucleic
Acid Purification Kit (Promega), subjected to DNAse treatment using TURBO DNAse (Invitrogen), and
reverse transcribed and amplified using the SuperScript IV One-Step RT-PCR system with HA-specific
primers. Amplified DNA was gel purified using the QIAquick Gel Extraction Kit (Qiagen) and quantified
using the Qubit dsDNA High Sensitivity Assay Kit on the Qubit Fluorometer (Invitrogen). Normalized
samples were made compatible for long-read sequencing using the 1D Native Barcoding ONT Kit (SQK-
LSK-109), following the manufacturer protocol. Libraries were loaded onto a flow cell and run on the ONT
GridION machine. Bases were called in real time using the ONT software package Guppy 3.2.6.
Minimap2 was used to map reads to the influenza virus *HA* segment and discard low-quality reads[50]. The
Sam2Tsv module from Jvarkit[51] was used to convert the sam to a tsv file so Pandas could be used to
extract the bases at position 540 (NheI library) or 543 (PstI library) and the barcode sequence. Results
were visualized with Prism 9.

Sequencing data availability

508   All sequencing files have been deposited as BioProjects PRJNA746307, PRJNA746319, and
509   PRJNA746307 with details and SRA accessions in Supplementary Table 3.

Growth kinetics

511   Triplicate dishes of confluent MDCK-SIAT1-TMPRSS2 cells were inoculated at an MOI of 0.01 with virus
512   diluted in OptiVGM. Virus was adsorbed for 1 hr at 37°C, removed from cells, and replaced with fresh
513   OptiVGM. Virus was sampled over time and titered by plaque assay on MDCK-SIAT1-TMPRSS2 cells.

Mouse infections

All mouse experiments were approved by the University of Wisconsin Madison Institutional Animal Care and Use Committee. 18 9-week old female BALB/c mice (Charles River Labs) were randomly divided into groups of 6 to receive either WT HA, HA-K153E-NheI-bc, or HA-K153E-PstI-bc virus. All viruses contained barcoded PASTN. Mice were inoculated intra-nasally with 10^5 TCIDGlo50 of virus in 35 µL media. Mice were weighed daily and monitored for clinical signs of infection. 3 animals from each group were sacrificed at 3 dpi and the remainder at 6 dpi. Lungs were removed, dounce homogenized in 1x DPBS, and clarified at 2000 x g for 5 min. The clarified homogenate was titered via plaque assay and TCIDGlo50 assay on MDCK-SIAT1-TMPRSS2 cells. Viral RNA was recovered from homogenate and sequenced as described above.

Ferret infections

All ferret experiments were approved by the St Jude Children's Research Hospital Animal Care and Use Committee. 12-week old male ferrets (Triple F Farms, Sayre, PA) confirmed to be seronegative for influenza virus were housed individually. Each animal was infected intranasally following previous approaches that ensure site-specific inoculation of the upper respiratory tract[32]. Animals were inoculated with 5x10[5] PFU HA-K153E-NheI-bc with barcoded PASTN diluted in PBS (Corning) containing 100 U/mL penicillin and 100 ug/mL streptomycin (Corning) in a total volume of 500 µl. Nasal washes were collected at 1, 3, and 5 dpi. Briefly, animals were anesthetized with 0.25 mL ketamine and nasal washes collected by administering 1 mL PBS/pen-strep dropwise onto the nostrils. Animals were sacrificed at 5 dpi and trachea and separated lung lobes were removed and frozen prior to processing. Tissue was homogenized in 10% (w/v) L-15 media and using an OMNI TH220-PCRD homogenizer and clarified at 500 x g for 10 min to remove cell debris. Viral titers in homogenates and nasal washes were determined by TCIDGlo50 assay. Viral RNA was recovered from homogenate and sequenced as described above.

Statistical analysis

Viral titers are presented as the mean of n=3 and significance was tested with a two-way ANOVA with Tukey's post hoc analysis. Replicate sequencing runs were analyzed with a Pearson's and Spearman's correlation coefficient (Prism 9). Population diversity and evenness were assessed by measuring Shannon's diversity (H')[26] as follows:

$$H' = -\sum_{i=1}^{S} p_i \ln p_i$$

Where S is the total number of barcodes detected (richness) and p_i is the frequency of the i-th barcode in that sample. Population evenness is bounded at 0 and 1 and defined as the actual barcode diversity divided by the maximum possible diverstiy (H'$_{max}$) for the sample[25]:

$$evenness = \frac{H'}{H_{max}}$$

Bray-Curtis dissimilarity (BC$_{ij}$) was used to assess compositions and compare lung lobes[52], defined as:

$$BC_{ij} = 1 - \frac{2C_{ij}}{S_i + S_j}$$

where C_{ij} is the sum of the lesser values for only barcodes found in both lobes. S_i and S_j are the total number of barcode reads detected in either lobe.

Multinomial model of bottleneck size

While several methods currently exist that estimate viral transmission bottleneck sizes between a donor sample and a recipient sample, none of these methods are appropriate for estimating Nb from the barcode lineage frequencies available from this study. The beta-binomial approach outlined in[53] assumes that that each locus is biallelic and unlinked to other loci. A more recent approach developed by[54] allows for linked loci by reconstructing haplotypes. However, the number of haplotypes that can be considered in their approach is very low relative to the number of barcode lineages observed in this study. We thus developed a simple statistical approach to estimate Nb that allows for a large number of observed haplotypes (here, barcode lineages). The approach is as follows:

1. Identify the barcode lineages observed in the donor sample and calculate their frequencies
2. Calculate the overall number of barcode lineages present in the recipient sample that are also found in the donor sample
3. Over a range of Nb values, for each Nb do as follows:
   a. For a given Nb value, draw n separate times from a multinomial distribution with the probability vector given by the barcode lineage frequencies in the donor sample and the number of trials being Nb. A given draw can be considered a random realization of barcode lineages in a recipient sample seeded by a bottleneck size of Nb. We let n = 500; higher values of n did not alter the results.
   b. For each of the n draws from the multinomial distribution, quantify the number of barcode lineages present in the recipient sample.
   c. From the n independent draws, calculate the mean and standard deviation of the number of barcode lineages present in the recipient sample.
   d. Calculate the probability of observing the observed number of barcodes in the recipient using a normal distribution with mean and standard deviations as calculated above. The log of this probability yields the log-likelihood of the bottleneck size being Nb.
4. Identify the maximum likelihood estimate of Nb as the Nb yielding the highest log-likelihood. Calculate the 95% confidence interval of Nb as the set of Nb values that yield log-likelihood values within 1.92 log-likelihood units of the maximum likelihood estimate of Nb.

ACKNOWLEDGEMENTS

We thank Jesse Bloom and Yoshihiro Kawaoka for providing key plasmids and reagents. We thank Christopher Brooke and Brigette Martin for assistance in virus rescue protocols. This work is funded by AI125392 to AM and TCF and by the National Institute of Allergy and Infectious Diseases under HHS contract HHSN27220140006C for the St. Jude Center of Excellence for Influenza Research and Surveillance, NIH grant R01AI140766, and ALSAC to SSC. KAA is supported by GM007215. KMB is supported by AI145182. LAH is supported by HG002760. GAS is supported by a Rath Foundation Wisconsin Distinguished Graduate Fellowship and AI007414. AM holds an Investigators in the Pathogenesis of Infectious Disease Award from the Burroughs Wellcome Fund.

References

1.   Smith, D. J. *et al.* Mapping the antigenic and genetic evolution of influenza virus. *Science* **305**,
371–376 (2004).

2.   Ghedin, E. *et al.* Large-scale sequencing of human influenza reveals the dynamic nature of viral
genome evolution. *Nature* **437**, 1162–1166 (2005).

3.   Rambaut, A. *et al.* The genomic and epidemiological dynamics of human influenza A virus. *Nature*
453, 615–619 (2008).

4.   Dinis, J. M. *et al.* Deep Sequencing Reveals Potential Antigenic Variants at Low Frequencies in
Influenza A Virus-Infected Humans. *J. Virol.* **90**, 3355–3365 (2016).

5.   Debbink, K. *et al.* Vaccination has minimal impact on the intrahost diversity of H3N2 influenza
viruses. *PLoS Pathog.* **13**, e1006194 (2017).

6.   Imai, H. *et al.* Diversity of Influenza A(H5N1) Viruses in Infected Humans, Northern Vietnam,
2004-2010. *Emerg. Infect. Dis.* **24**, 1128–1238 (2018).

7.   Moncla, L. H. *et al.* Quantifying within-host diversity of H5N1 influenza viruses in humans and
poultry in Cambodia. *PLoS Pathog.* **16**, e1008191 (2020).

8.   McCrone, J. T. *et al.* Stochastic processes constrain the within and between host evolution of
influenza virus. *Elife* **7**, e35962 (2018).

9.   Xue, K. S. & Bloom, J. D. Linking influenza virus evolution within and between human hosts. *Virus*
Evol. **6**, veaa010 (2020).

10.   Varble, A. *et al.* Influenza A Virus Transmission Bottlenecks Are Defined by Infection Route and
Recipient Host. *Cell Host Microbe* **16**, 691–700 (2014).

11.   Willi, Y., Van Buskirk, J. & Hoffmann, A. A. Limits to the Adaptive Potential of Small Populations.
Annu. Rev. Ecol. Evol. Syst. **37**, 433–458 (2006).

12.   Duarte, E., Clarke, D., Moya, A., Domingo, E. & Holland, J. Rapid fitness losses in mammalian
RNA virus clones due to Muller's ratchet. *Proc. Natl. Acad. Sci. U. S. A.* **89**, 6015–6019 (1992).

13.   Zaraket, H. *et al.* Mammalian adaptation of influenza A(H7N9) virus is limited by a narrow genetic
bottleneck. *Nat. Commun.* **6**, 6553 (2015).

14.   Jonsson, C. B. *et al.* Molecular Imaging Reveals a Progressive Pulmonary Inflammation in Lower
Airways in Ferrets Infected with 2009 H1N1 Pandemic Influenza Virus. *PLoS One* **7**, 1–12 (2012).

15.   Shinya, K. *et al.* Avian flu: Influenza virus receptors in the human airway. *Nature* **440**, 435–436
(2006).

16.   van Riel, D. *et al.* H5N1 Virus Attachment to Lower Respiratory Tract. *Science* **312**, 399 (2006).

17.   Chandrasekaran, A. *et al.* Glycan topology determines human adaptation of avian H5N1 virus
hemagglutinin. *Nat. Biotechnol.* **26**, 107–113 (2008).

18.   Lakdawala, S. S. *et al.* The soft palate is an important site of adaptation for transmissible influenza
viruses. *Nature* **526**, 122–5 (2015).

19.   Richard, M. *et al.* Influenza A viruses are transmitted via the air from the nasal respiratory
epithelium of ferrets. *Nat. Commun.* **11**, 766 (2020).

20.   Tran, V., Moser, L. A., Poole, D. S. & Mehle, A. Highly sensitive real-time in vivo imaging of an
influenza reporter virus reveals dynamics of replication and spread. *J. Virol.* **87**, 13321–9 (2013).

21.   Marsh, G. A., Hatami, R. & Palese, P. Specific residues of the influenza A virus hemagglutinin viral
RNA are important for efficient packaging into budding virions. *J. Virol.* **81**, 9727–9736 (2007).

22.   Doud, M. B. & Bloom, J. D. Accurate measurement of the effects of all amino-acid mutations on

influenza hemagglutinin. *Viruses* **8**, 155 (2016).

23.   Neumann, G., Fujii, K., Kino, Y. & Kawaoka, Y. An improved reverse genetics system for influenza
A virus generation and its implications for vaccine production. *Proc. Natl. Acad. Sci. U. S. A.* **102**,
16825–16829 (2005).

24.   Jost, L. Entropy and diversity. *Oikos* **113**, 363–375 (2006).

25.   Pielou, E. C. The measurement of diversity in different types of biological collections. *J. Theor.*
Biol. **13**, 131–144 (1966).

26.   Shannon, C. E. A Mathematical Theory of Communication. *Bell Syst. Tech. J.* **27**, 379–423 (1948).

27.   Chen, Z. *et al.* Generation of Live Attenuated Novel Influenza Virus A/California/7/09 (H1N1)
Vaccines with High Yield in Embryonated Chicken Eggs. *J. Virol.* **84**, 44–51 (2010).

28.   Karlsson, E. A. *et al.* Visualizing real-time influenza virus infection, transmission and protection in
ferrets. *Nat. Commun.* **6**, 6378 (2015).

29.   Bouvier, N. M. & Lowen, A. C. Animal Models for Influenza Virus Pathogenesis and Transmission.
Viruses **2**, 1530–1563 (2010).

30.   Belser, J. A., Pulit-Penaloza, J. A. & Maines, T. R. Ferreting Out Influenza Virus Pathogenicity and
Transmissibility: Past and Future Risk Assessments in the Ferret Model. *Cold Spring Harb.*
Perspect. Med. **10**, (2020).

31.   Honce, R., Wohlgemuth, N., Meliopoulos, V. A., Short, K. R. & Schultz-Cherry, S. Influenza in
High-Risk Hosts—Lessons Learned from Animal Models. *Cold Spring Harb. Perspect. Med.* **10**,
(2020).

32.   Moore, I. N. *et al.* Severity of Clinical Disease and Pathology in Ferrets Experimentally Infected
with Influenza Viruses Is Influenced by Inoculum Volume. *J. Virol.* **88**, 13879–13891 (2014).

33.   Marshall, N., Priyamvada, L., Ende, Z., Steel, J. & Lowen, A. C. Influenza virus reassortment
occurs with high frequency in the absence of segment mismatch. *PLoS Pathog.* **9**, e1003421
(2013).

34.   Lumby, C. K., Nene, N. R. & Illingworth, C. J. R. A novel framework for inferring parameters of
transmission from viral sequence data. *PLoS Genet.* **14**, e1007718 (2018).

35.   Grubaugh, N. D. *et al.* Genetic Drift during Systemic Arbovirus Infection of Mosquito Vectors
Leads to Decreased Relative Fitness during Host Switching. *Cell Host Microbe* **19**, 481–492
(2016).

36.   Valesano, A. L. *et al.* Influenza B Viruses Exhibit Lower Within-Host Diversity than Influenza A
Viruses in Human Hosts. *J. Virol.* **94**, e01710-19 (2020).

37.   Herfst, S. *et al.* Airborne transmission of influenza A/H5N1 virus between ferrets. *Science (80-.).*
336, 1534–1541 (2012).

38.   Imai, M. *et al.* Experimental adaptation of an influenza H5 HA confers respiratory droplet
transmission to a reassortant H5 HA/H1N1 virus in ferrets. *Nature* **486**, 420–428 (2012).

39.   Kuss, S. K., Etheredge, C. A. & Pfeiffer, J. K. Multiple host barriers restrict poliovirus trafficking in
mice. *PLoS Pathog.* **4**, e1000082 (2008).

40.   McCune, B. T., Lanahan, M. R., TenOever, B. R. & Pfeiffer, J. K. Rapid Dissemination and
Monopolization of Viral Populations in Mice Revealed Using a Panel of Barcoded Viruses. *J. Virol.*
94, e01590-19 (2020).

41.   Lauring, A. S. & Andino, R. Exploring the Fitness Landscape of an RNA Virus by Using a
Universal Barcode Microarray. *J. Virol.* **85**, 3780–3791 (2011).

42.   Pfeiffer, J. K. & Kirkegaard, K. Bottleneck-mediated quasispecies restriction during spread of an
RNA virus from inoculation site to brain. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 5520–5525 (2006).

43.   Forrester, N. L., Guerbois, M., Seymour, R. L., Spratt, H. & Weaver, S. C. Vector-borne
683   transmission imposes a severe bottleneck on an RNA virus population. *PLoS Pathog.* **8**,
684   e1002897 (2012).

685   44.   Weger-Lucarelli, J. *et al.* Using barcoded Zika virus to assess virus population structure in vitro
686   and in Aedes aegypti mosquitoes. *Virology* **521**, 138–148 (2018).

687   45.   Lee, J. M. *et al.* Deep mutational scanning of hemagglutinin helps predict evolutionary fates of
688   human H3N2 influenza variants. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E8276–E8285 (2018).

689   46.   Tran, V. *et al.* Multi-Modal Imaging with a Toolbox of Influenza A Reporter Viruses. *Viruses* **7**,
690   5319–5327 (2015).

691   47.   Karlsson, E. A. *et al.* Measuring Influenza Virus Infection Using Bioluminescent Reporter Viruses
692   for In Vivo Imaging and In Vitro Replication Assays. in *Methods in Molecular Biology* vol. 1836
693   431–459 (2018).

694   48.   Poole, D. S. *et al.* Influenza A virus polymerase is a site for adaptive changes during experimental
695   evolution in bat cells. *J. Virol.* **88**, 12572–12585 (2014).

696   49.   Smith, T., Heger, A. & Sudbery, I. UMI-tools: modeling sequencing errors in Unique Molecular
697   Identifiers to improve quantification accuracy. *Genome Res.* **27**, 491–499 (2017).

698   50.   Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–3100
699   (2018).

700   51.   Lindenbaum, P. JVarkit: java-based utilities for Bioinformatics. *figshare*
701   DOI:10.6084/m9.figshare.1425030.v1 (2015) doi:10.6084/m9.figshare.1425030.v1.

702   52.   Bray, J. R. & Curtis, J. T. An Ordination of the Upland Forest Communities of Southern Wisconsin.
703   *Ecol. Monogr.* **27**, 325–349 (1957).

704   53.   Sobel Leonard, A., Weissman, D. B., Greenbaum, B., Ghedin, E. & Koelle, K. Transmission
705   Bottleneck Size Estimation from Pathogen Deep-Sequencing Data, with an Application to Human
706   Influenza A Virus. *J. Virol.* **91**, e00171-17 (2017).

707   54.   Mahan, G., K., L. C., B., W. D., R., I. C. J. & R., P. C. Inferring Transmission Bottleneck Size from
708   Viral Sequence Data Using a Novel Haplotype Reconstruction Method. *J. Virol.* **94**, e00014-20
709   (2021).