**Bcr4 is a Chaperone for the Inner Rod Protein in the *Bordetella* Type III Secretion**

**System**

Masataka Goto[a], Akio Abe[a], Tomoko Hanawa[b], Masato Suzuki[c], and Asaomi Kuwae[a#]

[a]Laboratory of Bacterial Infection, Graduate School of Infection Control Sciences, Kitasato

University, Tokyo, Japan

[b]Department of Infectious Diseases, Kyorin University School of Medicine, Tokyo, Japan

[c]Antimicrobial Resistance Research Center, National Institute of Infectious Diseases, Tokyo,

Japan

**Running Head:** Bcr4 is a Chaperone for Inner Rod in *Bordetella* T3SS

[#]**Address Correspondence to:** Asaomi Kuwae, kuwae@lisci.kitasato-u.ac.jp

**Abstract**

*Bordetella bronchiseptica* injects virulence proteins called effectors into host cells via a

type III secretion system (T3SS) conserved among many Gram-negative bacteria. Small

proteins called chaperones are required for stabilizing some T3SS components or

localizing them to the T3SS machinery. In a previous study, we identified a chaperone-like

protein named Bcr4 that regulates T3SS activity in *B. bronchiseptica*. Bcr4 does not show

strong sequence similarity to well-studied T3SS proteins of other bacteria, and its function

remains to be elucidated. Here, we investigated the mechanism by which Bcr4 controls

T3SS activity. A pull-down assay revealed that Bcr4 interacts with BscI, based on its

homology to other bacterial proteins, to be an inner rod protein of the T3SS machinery. An

additional pull-down assay using truncated Bcr4 derivatives and secretion profiles of *B.*

*bronchiseptica* producing truncated Bcr4 derivatives showed that the Bcr4 C-terminal

region is necessary for the interaction with BscI and activation of the T3SS. Moreover, the

deletion of BscI abolished the secretion of type III secreted proteins from *B. bronchiseptica*

and the translocation of a cytotoxic effector into cultured mammalian cells. Finally, we

showed that BscI is unstable in the absence of Bcr4. These results suggest that Bcr4

supports the construction of the T3SS machinery by stabilizing BscI. This is the first

demonstration of a chaperone for the T3SS inner rod protein among the virulence bacteria

possessing the T3SS.

**Importance**

The type III secretion system (T3SS) is a needle-like complex that projects outward from

bacterial cells. *Bordetella bronchiseptica* uses the T3SS to inject virulence proteins into

host cells. Our previous study reported that a protein named Bcr4 is essential for the

secretion of virulence proteins from *B. bronchiseptica* bacterial cells and delivery through

the T3SS. Because other bacteria lack a Bcr4 homologue, the function of Bcr4 has not

been elucidated. In this study, we discovered that Bcr4 interacts with BscI, a component of

the T3SS machinery. We showed that a *B. bronchiseptica* BscI-deficient strain was unable

to secrete type III secreted proteins. Furthermore, in a *B. bronchiseptica* strain that

overproduces T3SS component proteins, Bcr4 is required to maintain BscI in bacterial cells.

These results suggest that Bcr4 stabilizes BscI to allow construction of the T3SS in *B.*

*bronchiseptica*.

61
62

## Introduction

The genus *Bordetella* consists of Gram-negative bacteria that infect the respiratory tracts of mammals including humans. *Bordetella pertussis* causes a severe coughing attack called whooping cough in humans (1, 2). *Bordetella bronchiseptica* causes atrophic rhinitis in pigs and kennel cough in dogs (1, 2). These *Bordetella* spp. harbor a virulence factor secretion apparatus called the type III secretion system (T3SS).

The T3SS consists of a basal body that penetrates the inner and outer membranes of the bacteria and a needle structure that protrudes outside the bacteria, and is conserved in many Gram-negative bacteria such as *Yersinia*, *Salmonella*, and *Pseudomonas*. *B. bronchiseptica* uses the T3SS to inject virulence factors, called effectors, into host cells to disrupt the physiological functions of host cells (3). Once the basal body and the export apparatus of the T3SS are completed, the type III secreted proteins are secreted in a fixed order. First, the components (Table S1) of the needle structure (SctF) and the inner rod (SctI) that ties the needle structure to the basal body are secreted. After the needle is completed, translocators (SctE and SctB) are secreted, which form small pores in the host cell membrane to create a pathway for effectors. Finally, the effectors translocate into the host cell via the T3SS (4, 5). In *B. bronchiseptica*, BopB (SctE) (6), BopD (SctB) (7), and Bsp22 (SctA) (8) function as translocators (Table S1), while BteA (also referred to as BopC) (9, 10), BopN (11, 12), and BspR (also referred to as BtrA) (13, 14) function as effectors. Bsp22 is located at the tip of the needle and bridges the needle to the pore-forming factors

embedded in the plasma membrane (15). BteA has been shown to cause

membrane-disrupting cytotoxicity in mammalian cells (10). BspR is a regulator that

represses transcription of the *bteA* gene and genes on the *bsc* locus (Fig. S1), where genes

encoding type III secreted proteins and components of T3SS are located (14, 16).

According to secondary structure prediction—e.g., the predicted positions of helix, and the

overall structure of the operon—the BscF and BscI of *B. bronchiseptica* correspond to

*Yersinia* needle YscF (SctF) and inner rod YscI (SctI), respectively (Table S1).

In addition, many type III secreted proteins have unique chaperones that are involved in

stabilizing the substrate and preventing premature polymerization of the substrate in the

bacterial cytosol, and then in efficiently transporting the substrate to the T3SS machinery

(17-20). For example, PscE and PscG function as chaperones of the needle PscF (SctF) in

*Pseudomonas* (Table S1). These chaperones stabilize PscF in the bacterial cytosol and are

involved in PscF secretion through the T3SS (21). In bacteria such as *Yersinia* and

*Pseudomonas*, an inner rod chaperone is thought to exist, because the inner rod is

secreted out of the bacterial cell and polymerized (22, 23). However, a chaperone for the

inner rod has not been reported.

So far, a chaperone-like protein called Bcr4 has been identified in *B. bronchiseptica* (24).

*Bordetella* Bcr4 is highly conserved among *B. pertussis*, *B. parapertussis*, and *B.*

*bronchiseptica* (Fig. S2). In this study, we attempted to identify factors that interact with

Bcr4 in order to investigate how Bcr4 is involved in the T3SS regulation.

## Results

**Bcr4 binds to BscI, an inner rod protein of the *Bordetella* type III secretion system.**

The results of a previous study suggested that Bcr4 is a chaperone for components of the type III secretion system (T3SS) (24). T3SS chaperones are known to be involved in substrate stability and efficient transport of substrates to the T3SS machinery (17, 19, 20). In addition, it is generally known that the genes of these chaperones are localized adjacent to genes encoding their substrates (25). On the *B. bronchiseptica* S798 chromosome, the genes encoding BcrH2, BscI, BscJ, and BscK are located in the vicinity of the *bcr4* gene (Fig. 1A), and these proteins are predicted to function as a translocator chaperone, inner rod (SctI), inner membrane ring (SctJ), and ATPase cofactor (SctK), respectively (Table S1) (5, 26). To test whether Bcr4 binds to BscI, BscJ or BscK, we added *E. coli* lysates containing the V5-tagged target proteins (BscI-V5, BscJ-V5 or BscK-V5) to Strep-Tactin beads loaded with the purified Strep-tagged Bcr4 (Bcr4-Strep), and then performed the pull-down assay. The supernatant fraction (Sup) and pellet fraction (Pellet) samples were prepared, separated by SDS-PAGE, and subjected to Western blotting with anti-V5 antibody (Fig. 1B). When the beads were washed with TBS, the V5 signal of the BscI-V5 pellet sample was detected in the beads loaded with Bcr4, but not in the unloaded beads (Fig. 1B). The V5 signals of the BscK-V5 and BscJ-V5 pellet samples were detected in both the Bcr4-loaded beads and the unloaded beads when washed with TBS, and not detected when washed with TBS containing 0.1% Triton X-100 (Fig. 1B). These results suggest that

Bcr4 binds to BscI, an inner rod protein of the *Bordetella* T3SS.

Next, to confirm that Bcr4 binds to BscI, a pull-down assay was performed by adding *E.*

*coli* lysate containing V5-tagged Bcr4 (Bcr4-V5) to Strep-Tactin beads with the purified

Strep-tagged BscI (BscI-Strep) or BteA N-terminal 1–312 amino acids region

(BteA-N-Strep). BteA is a protein secreted from the type III secretion system and interacts

with its cognate chaperone BtcA through the N-terminal (9). As a result, the V5 signal was

detected in the pellet sample of the BscI-Strep, but was hardly detected in that of the

BteA-N-Strep (Fig. 1C). These results strongly suggested that Bcr4 binds to BscI.

**The C-terminal region of Bcr4 is required for the binding of Bcr4 to BscI.**

Next, to investigate the Bcr4 region responsible for the binding to BscI, we

produced Strep-tagged full-length Bcr4 (Bcr4-FL-Strep), Bcr4 lacking amino acids region

58–109 (Bcr4Δ58-109-Strep), and Bcr4 lacking amino acids region 110–173

(Bcr4Δ110-173-Strep) in *E. coli* (Fig. 2A). A pull-down assay was performed by adding *E.*

*coli* lysate containing V5-tagged BscI (BscI-V5) to Strep-Tactin beads with the purified

Bcr4-FL-Strep, Bcr4Δ58-109-Strep, or Bcr4Δ110-173-Strep. The Sup and Pellet samples

were then prepared, separated by SDS-PAGE, and subjected to Western blotting using

anti-V5 antibody (Fig. 2B). As a result, the V5 signal was detected in the pellet sample of

Bcr4-FL-Strep, but not in those of Bcr4Δ58-109-Strep and Bcr4Δ110-173-Strep. Although it

is still unknown which region of Bcr4 directly interacts with BscI, our results strongly

suggest that both the Bcr4-58-109 and Bcr4-110-173 regions are required for the

interaction.

**The C-terminal region of Bcr4 is required for T3SS activity.**

The results in Figure 2 suggest that the C-terminal region of Bcr4 is required for binding

to BscI. Therefore, we introduced plasmids encoding the full-length FLAG-tagged Bcr4

(Bcr4-FL-FLAG), Bcr4 with the 5 C-terminal amino acids deleted (Bcr4Δ169-173-FLAG),

Bcr4 with the 10 C-terminal amino acids deleted (Bcr4Δ164-173-FLAG), and Bcr4 with the

15 C-terminal amino acids deleted (Bcr4Δ159-173-FLAG) (Fig. 3 A) into a Bcr4-deficient

strain         (Δ*bcr4*)         (Δ*bcr4+bcr4*-FL-FLAG,         Δ*bcr4+bcr4*Δ169-173-FLAG,

Δ*bcr4+bcr4*Δ164-173-FLAG and Δ*bcr4+bcr4*Δ159-173-FLAG, respectively). Whole cell

lysates (WCL) and culture supernatant fraction (CS) samples were prepared from these

strains and separated by SDS-PAGE. Western blottings were then carried out using

antibodies against Bcr4, FLAG, BteA (an effector and a type III secreted protein), BopD

(SctB, a translocator and a type III secreted protein) or RpoB (an internal control of WCL)

(Fig. 3B). Since the antibody against Bcr4 was generated using its 18 C-terminal amino

acids (amino acids (aa) 156–173) as the antigen peptide, we considered that it may not

recognize the Bcr4 partial deletion mutant proteins used in this experiment. Therefore, we

also performed Western blotting using anti-FLAG antibody (Fig. 3B). As a result, when

anti-Bcr4 antibody was used against the WCL samples, signals were detected in the

wild-type, Δ*bcr4*+*bcr4*-FL-FLAG and Δ*bcr4*+*bcr4* Δ169-173-FLAG, but almost no signals

were detected in Δ*bcr4*+*bcr4*Δ164-173-FLAG and Δ*bcr4*+*bcr4*Δ159-173-FLAG (Fig. 3B).

On the other hand, when anti-FLAG antibody was used, signals were detected in

Δ*bcr4*+*bcr4*Δ164-173-FLAG and Δ*bcr4*+*bcr4*Δ159-173-FLAG (Fig. 3B). These results

confirmed that the Bcr4 partial deletion mutant proteins were produced in the Δ*bcr4* strain.

When anti-BteA or anti-BopD antibodies were used against the CS samples, signals were

detected in the wild-type, Δ*bcr4*+*bcr4*-FL-FLAG, Δ*bcr4*+*bcr4*Δ169-173-FLAG and

Δ*bcr4*+*bcr4*Δ164-173-FLAG, respectively, but not in Δ*bcr4* or Δ*bcr4*+*bcr4*Δ159-173-FLAG

(Fig. 3B). We attempted to create *B. bronchiseptica* strains that produce shorter Bcr4, e.g.

amino acid regions 1-57, 58-109, and 110-173, however, those truncated Bcr4 were

produced at very low levels (data not shown). Therefore, we were unable to evaluate

whether or not these truncated proteins were functional in *B. bronchiseptica*. We then

infected L2 cells (a rat lung epithelial cell line) with the wild-type, Δ*bcr4*,

Δ*bcr4*+*bcr4*-FL-FLAG, Δ*bcr4*+*bcr4*Δ169-173-FLAG, Δ*bcr4*+*bcr4*Δ164-173-FLAG or

Δ*bcr4*+*bcr4*Δ159-173-FLAG strain at a multiplicity of infection (MOI) of 50 for 1 h and

measured the amounts of LDH released into the culture medium as an index of cytotoxicity.

As a result, LDH was detected in the medium of cells infected with Δ*bcr4*+*bcr4*-FL-FLAG,

Δ*bcr4*+*bcr4*Δ169-173-FLAG or Δ*bcr4*+*bcr4*Δ164-173-FLAG, but not in the medium of cells

infected with Δ*bcr4*+*bcr4*Δ159-173-FLAG (Fig. 3C). These results suggest that the region

required for the T3SS function is located in the 159–163 amino acids region of Bcr4.

**BscI is required for the function of the T3SS in *B. bronchiseptica*.**

BscI is a homologue of YscI, an inner rod (SctI) of the *Yersinia* T3SS, and YscI is

essential for the function of the T3SS (22). To confirm that BscI is required for the function

of the *B. bronchiseptica* T3SS, the *B. bronchiseptica* S798 wild-type, BscI-deficient strain

(Δ*bscI*), BscI-complemented strain (Δ*bscI+bscI*), and BscN-deficient strain (Δ*bscN*) were

cultured in SS medium. BscN is a protein predicted to function as an ATPase (SctN) and is

required for the T3SS function in *Bordetella bronchiseptica* (Table S1) (6, 27). We used the

Δ*bscN* as a T3SS-deficient strain. WCL and CS samples were prepared, and these

samples were separated by SDS-PAGE and analyzed by Western blotting using antibodies

against BscI, BteA, BopD, Bsp22 (SctA, a translocator and a component of the filament-like

structure; Fig. 7) or FHA (filamentous hemagglutinin, an adhesion factor) (Fig. 4A). As

expected, the FHA signals were detected in the WCL and CS samples of all strains used

here (Fig. 4A). In contrast, no signals of BscI were detected in the WCL samples of all

strains used here (Fig. 4A). The BscI signal was detected in the CS of Δ*bscI+bscI*, and a

faint BscI signal was detected in the CS of the wild-type (Fig. 4A). In order to examine

whether the signal we obtained around 20 kDa in the western blot using anti-BscI antibody

in Fig. 4A was specific or nonspecific, we prepared the supernatant fraction from Δ*bsp22*

strain. As a result, the signal disappeared in the supernatant fraction of the Δ*bsp22* strain

(Fig. S3), suggesting that the signal around 20 kDa obtained in the western blot was a

nonspecific interaction between anti-BscI antibody and an excess amount of Bsp22 on the

membrane. The BteA, BopD and Bsp22 signals were detected in the CS of the wild-type

and Δ*bscI*+*bscI*, but not in those of Δ*bscI* and Δ*bscN* (Fig. 4A). BspR is a negative regulator

that represses the transcription of genes on the *bsc* locus (14). In *Bordetella* strains that are

unable to secrete type III secreted proteins, such as Bcr4-deficient strains (Δ*bcr4*), the

repression of the *bsc* locus transcriptions by BspR is enhanced (24). Therefore, we

speculated that the production of BopD and Bsp22 was intensely suppressed by BspR in

Δ*bscI* due to the loss of T3SS activity. To test whether this hypothesis is correct, we

generated Δ*bspR*Δ*bscI* (a BspR- and BscI-deficient strain). WCL and CS samples were

prepared from the *B. bronchiseptica* S798 wild-type strain, Δ*bspR*, Δ*bspR*Δ*bscI* or

Δ*bspR*Δ*bscI*+*bscI* (BscI-complemented BspR- and BscI-deficient strain), and these

samples were separated by SDS-PAGE and analyzed by Western blotting using antibodies

against Bsp22 or BopB (SctE, a translocator and a type III secreted protein). As a result, the

Bsp22 and BopB signals were detected in the WCL samples of the wild-type, Δ*bspR*,

Δ*bspR*Δ*bscI* and Δ*bspR*Δ*bscI*+*bscI* strains (Fig. 4B). The Bsp22 and BopB signals were

detected in the CS samples of the wild-type, Δ*bspR* and Δ*bspR*Δ*bscI*+*bscI* strains, but not

in that of the Δ*bspR*Δ*bscI* strain (Fig. 4B). These results suggest that in *B. bronchiseptica*,

BscI is required for the function of the T3SS and is secreted out of the bacterial cell.

**BscI plays an important role in the function of the *B. bronchiseptica* T3SS during**

**infection of cultured mammalian cells.**

To further investigate whether BscI is required for the construction and function of the T3SS

in *B. bronchiseptica*, we infected L2 cells with the wild-type, Δ*bscI*, Δ*bscI+bscI* or Δ*bscN*

and measured the number of Bsp22 (SctA, a translocator, and a component of the

filament-like structure; Fig. 5) signals detected on L2 cells by immunofluorescence

microscopy. For the efficient detection of Bsp22 signals, we cultured the bacteria in

iron-depleted SS medium, which has been shown to increase the amount of type III

secreted protein secreted by *B. bronchiseptica* (28). L2 cells were infected with bacteria for

1 h at an MOI of 2000. Then, Bsp22, F-actin and bacteria were stained with anti-Bsp22

antibody, Rhodamine-Phalloidin and DAPI, respectively (Fig. 5A). The DAPI fluorescence

signals showing bacterial genomic DNA were detected to the same extent among the L2

cells infected with each bacterial strain, suggesting that the amounts of each mutant

adhered to the cells were not significantly reduced when compared to the wild-type strain

(Fig. 5A). The fluorescence signals of Bsp22 were detected on the L2 cells infected with the

wild-type or Δ*bscI+bscI*, but not on the L2 cells infected with Δ*bscI* or Δ*bscN* (Fig. 5 A, B).

We also infected L2 cells with the wild-type, Δ*bscI*, Δ*bscI+bscI* or Δ*bscN* for 1 h at an MOI of

200 and measured the amount of LDH released into the extracellular medium. The results

showed that LDH was detected in the medium of L2 cells infected with the wild-type or

Δ*bscI+bscI*, but not in the medium of L2 cells infected with Δ*bscI* or Δ*bscN* (Fig. 5 C). These

results suggest that BscI is required for the construction of the T3SS and induction of

T3SS-dependent death of mammalian cells in *B. bronchiseptica*.

**Bcr4 is required for the stability of BscI and is suggested to interact with BscI in *B.***

***bronchiseptica*.**

As mentioned above, our results suggested that Bcr4 interacts with BscI (Fig. 1). Because

Bcr4 has various properties like those of chaperones for the type III secreted proteins (25),

e.g., low molecular mass (18.1 kDa) and low isoelectric point (pI 4.41), we investigated

whether Bcr4 is involved in the stability of BscI. We cultured the *B. bronchiseptica* wild-type,

Bcr4-deficient strain (Δ*bcr4* , T3SS-inactive strain) (24), BspR-deficient strain (Δ*bspR* ,

T3SS-overproducing and -oversecreting strain) (14) or BspR- and Bcr4-deficient strain

(Δ*bspR* , a T3SS-overproducing but -inactive strain) (24) in SS medium. The prepared WCL

samples were separated by SDS-PAGE, and Western blotting was performed using

antibodies against BscI, BscJ (SctJ, a protein composing the inner membrane ring of the

T3SS), or Bcr4. As shown in Fig. S1, the negative regulatory function of BspR, which

represses the *bsc* locus transcription, is enhanced in the strain that loses the activity

of T3SS, e.g., Δ*bcr4* (24). Because of this property of BspR, the BscJ signal of the WCL

sample was weaker in Δ*bcr4* than in the wild-type (Fig. 6A). In Δ*bspR* and Δ*bspR*Δ*bcr4*,

BscJ signals of the WCL samples were more strongly detected when compared to the

wild-type (Fig. 6A) due to the BspR deficiency. On the other hand, the BscI signals of the

WCL and CS samples were detected in Δ*bspR*, but not in the Δ*bspR*Δ*bcr4* strain (Fig. 6A).

To investigate the presence of *bscI* mRNA in Δ*bspR*Δ*bcr4*, a quantitative RT-PCR was

performed. The results showed that the amount of *bscI* mRNA in Δ*bspR*Δ*bcr4* was 1.5-fold

higher than that of *bscI* in Δ*bspR* (Fig. S4), demonstrating that the *bscI* gene is transcribed

in Δ*bspR*Δ*bcr4*. These results suggest that Bcr4 is necessary for the stability of BscI in *B.*

*bronchiseptica*. Finally, we examined whether Bcr4 binds to BscI inside of the *B.*

*bronchiseptica* cells. As described in the Discussion section, we were unable to detect an

interaction of Bcr4 with BscI by immunoprecipitation because of the low level of BscI protein

in bacterial cells. Therefore, we attempted to reveal the interaction by using a cross linker.

We treated the wild-type, Δ*bspR* and Δ*bspR*Δ*bcr4* strains with disuccinimidyl suberate

(DSS) and then prepared the whole cell lysate samples. The samples were separated by

SDS-PAGE and then analyzed by Western blotting using anti-Bcr4 antibody. As a result, an

extra signal around 33 kDa was evident in Δ*bspR* but not in Δ*bspR*Δ*bscI* (Fig. 6B). The size

of the extra signal was almost equivalent to that of the Bcr4-BscI complex, suggesting the

possibility that Bcr4 binds to BscI in the *B. bronchiseptica* cells.

**Discussion**

In the present study, we found that deletion of BscI in *B. bronchiseptica* abolished the

extracellular secretion of type III secreted proteins (Fig. 4, 5). In addition, when Bcr4 was

deleted in the *B. bronchiseptica* BspR-deficient strain, BscI (SctI, a protein composing the

inner rod of the *Bordetella* T3SS) was not detected in the bacterial cells (Fig. 6A). These

results suggest that Bcr4 contributes to the establishment of the T3SS in *B. bronchiseptica*

by stabilizing BscI (Fig. 7).

To date, chaperones for needles, translocators and effectors have been reported in

bacteria that retain the T3SS, such as *Yersinia* (29-32). However, a chaperone for the inner

rod protein has not been reported, and the mechanism underlying stability of the inner rod

in the bacterial cell and the transport of inner rods to the T3SS remain unclear.

In the present study, it was suggested that both the 58–109 and 110–173 amino acid

regions of Bcr4 are required for the binding of Bcr4 to BscI (Fig. 2B). To investigate whether

the 1–57 amino acids region of Bcr4 is also required for the binding of Bcr4 to BscI, we

attempted to purify Bcr4 lacking the N-terminal 1–57 amino acids region (Bcr4Δ1-57-Strep).

However, the amount of Bcr4Δ1-57-Strep produced in *E. coli* was extremely low (data not

shown), so a pull-down assay could not be performed. Figure 3B suggests that the 159–

163 amino acids region of Bcr4 is required for the function of the T3SS. However, it is still

unclear whether the 159−163 amino acids region of Bcr4 is required for binding to BscI.

Further analysis is needed to clarify whether the binding of Bcr4 to BscI is required for the

function of T3SS.

In this study, the BopD signal was detected in the WCL of the *B. bronchiseptica*

wild-type strain, but the BscI signal was not detected (Fig. 4A). We detected *bscI* mRNA in

*B. bronchiseptica*, and the amount of *bscI* mRNA was not less than that of *bopD* mRNA (Fig.

S5). Therefore, the amount of BscI protein present in *B. bronchiseptica* was considered to

be much lower than that of BopD. The reason for this finding is unknown, but it could be

due to the low efficiency of translation of the *bscI* gene. In order to detect BscI in the whole

cell lysate, we prepared the samples at 0 hr to 18 hr after suspending the bacteria in liquid

broth. However, no signals were detected in the whole cell lysate samples prepared from

the wild-type strain or the Δ*bcr4* strain (Fig. S6). We also used the Δ*bspR* and Δ*bspR*Δ*bcr4*

strains, and detected BscI signals in the Δ*bspR* strain, but not in the Δ*bspR*Δ*bcr4* double

knockout strain (Fig. S6), strongly suggesting that Bcr4 is important for the stability of BscI.

In this study, we attempted to detect the BscI signal using a BspR-deficient strain

(Δ*bspR*), in which the gene transcriptions on the *bsc* locus are promoted (Fig. 6), because

the BscI signal was not detected in the WCL sample of the *B. bronchiseptica* wild-type (Fig.

4). The *bscJ* gene is located downstream of the *bscI* gene (Fig. 1A), and the BscJ protein is

predicted to function as an inner membrane ring (SctJ), a component of the T3SS. The

signal intensity of BscJ was not decreased in the WCL sample of the BspR/Bcr4

double-deficient strain (Δ*bspR*Δ*bcr4*) compared to that of Δ*bspR* (Fig. 6A), suggesting that

Bcr4 is not involved in the stability of BscJ. Therefore, it is suggested that Bcr4 specifically

stabilizes BscI. In order to detect the interaction between Bcr4 and BscI in the *B.*

*bronchiseptica* cells, immunoprecipitation was performed using anti-BscI antibody against

Δ*bspR* lysate. The immunoprecipitated fractions were analyzed by Western blotting using

anti-BscI antibody; however, an evident BscI signal was not detected (data not shown). The

results of our immunoprecipitation assay were thus unable to demonstrate a specific

interaction between Bcr4 and BscI.   We speculate that *B. bronchiseptica* did not maintain

a sufficient amount of BscI in whole cell lysate to detect the interaction by

immunoprecipitation. In Fig. 6B, we obtained an extra signal around 33 kDa in Western

blotting using anti-Bcr4 antibody when we analyzed the lysate of *B. bronchiseptica* treated

with a cross-linker, DSS. When we performed the Western blotting using anti-BscI antibody

for the cross-linked lysate, the extra signal was not detected (data not shown).

In order to examine whether Bcr4 has structural similarity to any chaperones for the type

III secreted proteins produced by other bacteria, we used AlphaFold2. As a result, we

detected significant structural similarities to the other chaperones, *Aeromonas* AcrH (34)

and *Pseudomonas* PscG (35) (Fig. S7). Although we obtained no plausible structural model

for the interaction between BscI and Bcr4, the results strongly suggest that Bcr4 is a

chaperone.

It is still unclear how Bcr4 antagonizes the BspR-negative regulation activity, and how

overexpression of Bcr4 promotes the production and secretion of type III secreted proteins

in *B. bronchiseptica*. Further analysis is needed to elucidate the detailed molecular

mechanism by which Bcr4 contributes to the regulation and establishment of the T3SS.

## Materials and Methods

### Bacterial strains and cell culture.

The strains used in this study are listed in Table 1. *B. bronchiceptica* S798 (6) was used as the wild-type strain. Δ*bscN* (a BscN-deficient strain), Δ*bcr4* (a Bcr4-deficient strain), Δ*bspR* (a BspR-deficient strain) and Δ*bspR*Δ*bcr4* (a BspR and Bcr4-deficient strain) have been described elsewhere (6, 14, 24). *Escherichia coli* DH10B, BL21 (36) and Sm10λ*pir* were used for DNA cloning, recombinant protein expression and conjugation, respectively (Table 1). *B. bronchiceptica* was grown on Bordet-Gengou agar plates at 37ºC for 2 days. Fresh colonies of *B. bronchiceptica* were cultured in Stainer-Scholt (SS) medium (37) with a starting $A_{600}$ of 0.23 under vigorous shaking conditions at 37ºC. L2 cells, a rat lung epithelial cell line (ATCC CCL-149), were maintained in F-12K (Invitrogen). The cell culture medium was supplemented with 10% fetal calf serum (FCS). L2 cells were grown at 37°C under a 5% $CO_2$ atmosphere.

### Preparation of proteins from culture supernatants and whole bacterial cell lysates.

Proteins secreted into bacterial culture supernatants and whole bacterial cell lysates were prepared as described previously (38). The loaded sample amounts were adjusted by the $A_{600}$ of each bacterial culture to load samples prepared from the same amounts of bacteria. The protein samples were separated by SDS-PAGE and analyzed by Western blotting.

**Antibodies.**

Anti-BteA, anti-BopD, anti-Bsp22, BspR, BopB, and anti-Bcr4 antibodies were purified

from rabbit serum in our previous study (6, 7, 10, 14, 24, 39). Mouse anti-V5 and anti-RNA

polymerase beta subunit (RpoB) monoclonal antibodies were purchased from Santa Cruz

Biotechnology and BioLegend, respectively. To detect filamentous hemagglutinin (FHA)

signals, we used mouse anti-FHA serum kindly gifted from K. Kamachi (National Institute of

Infectious Diseases). To prepare the anti-BscI and anti-BscJ antibodies, the peptides

corresponding to the C-terminus regions of BscI (CKAIGRATQNVDTLARMS) and BscJ

(CRGEGRGGAGAGATEGAGHD) were conjugated with hemocyanin from keyhole limpets

(Sigma), respectively, by using 3-maleimidobenzoic acid N-hydroxysuccinimide ester

(Sigma). These cross-linked peptides were used to immunize rabbits, and the resulting

anti-sera were incubated with the same peptides immobilized on epoxy-activated

sepharose 6B (Amersham) to obtain specific Ig-fractions.

**Immunofluorescence staining of L2 cells infected with *B. bronchiseptica.***

Immunofluorescent staining assay was performed as previously described (38), with

slight modifications. Briefly, L2 cells were seeded on coverslips in 6-well plates and

incubated overnight. In order to detect adequate amounts of Bsp22 signals, we infected the

L2 cells with *B. bronchiseptica* cultured in iron-depleted SS medium at an MOI of 2000. To

avoid excessive killing of L2 cells with bacteria, the plates were not centrifuged. After being

incubated for 1 h at 37°C under a 5% $CO_2$ atmosphere, the infected L2 cells were

immunostained. Average numbers of Bsp22 signals on single L2 cells were measured by a

fluorescence microscope.

**LDH Assays.**

LDH assays were performed as previously described (38). Briefly, to examine whether

lactate dehydrogenase (LDH) is released from *B. bronchiseptica*-infected cells, $5.0×10^4$

cells/well of L2 cells seeded in 24-well plates were infected with bacteria at an MOI of 50 or

200. The plates were centrifuged at $900×g$ for 5 min and incubated for 3 h at 37°C under a

5% $CO_2$ atmosphere. The amounts of LDH were measured spectrophotometrically using a

Cyto-Tox 96 Non-radioactive Cytotoxicity Assay kit (Promega). The LDH (%) level is shown

as a ratio, with the value obtained from the well treated with 0.1% Triton X-100 set as

100%.

**Cross-linking Assay**

The cross-linking assay was performed as previously described (40). A 100 μL culture of

*B. bronchiseptica* was centrifuged at $20,000×g$ and the supernatant was removed. The

pellet was washed with PBS and resuspended in 100 μL of PBS. Then, disuccinimidyl

suberate (DSS (Thermo)) dissolved in DMSO was added at a final concentration of 10 mM.

After incubation on ice for 1 h, Tris-HCl pH8.0 was added at a final concentration of 50 mM.

The solution was centrifuged at 20,000×$g$, and the supernatant was removed. The pellet

was dissolved in 2×SDS-PAGE sample buffer. The samples were prepared from the same

amounts of bacteria based on the $A_{600}$ values of the bacterial culture, then separated by

SDS-PAGE, and analyzed by Western blotting.

**Statistical analyses**

The statistical analyses were performed using the nonparametric test with a two-tailed

p-value with Prism ver. 5.0f software (Graphpad, La Jolla, CA). Values of $p < 0.05$ were

considered significant.

**Others**

The pull-down assay, the construction of *bscI* gene-disrupted or *bspR*/*bscI* double strains,

the construction of the plasmids used for producing Bcr4 derivatives and BscI

complementation, the quantitative reverse transcription-PCR, and Protein structure

prediction are described in Text S. Plasmids and Primers used in this study are listed in

Table S2 and S3, respectively.

**Acknowledgements**

This work was supported in part by grants (19K07542 to A.A.; 20K07485 to A.K.; 19K07561 to T.H.) from the Ministry of Education, Culture, Sports, Science and Technology and the Japan Society for the Promotion of Science (KAKENHI), and a grant (JP22gm1610003 to M.S. and A.K.) from the Japan Agency for Medical Research and Development (AMED).

The funders had no role in the study design, date collection or analysis, the decision to publish, or the preparation of manuscript.

**References**

1. Goodnow RA. 1980. Biology of *Bordetella bronchiseptica*. Microbiol Rev 44:722-38.

2. Mattoo S, Cherry JD. 2005. Molecular pathogenesis, epidemiology, and clinical manifestations of respiratory infections due to *Bordetella pertussis* and other *Bordetella* subspecies. Clin Microbiol Rev 18:326-82.

3. Cornelis GR, Van Gijsegem F. 2000. Assembly and function of type III secretory systems. Annu Rev Microbiol 54:735-74.

4. Büttner D. 2012. Protein export according to schedule: architecture, assembly, and regulation of type III secretion systems from plant- and animal-pathogenic bacteria. Microbiol Mol Biol Rev 76:262-310.

5. Deng W, Marshall NC, Rowland JL, McCoy JM, Worrall LJ, Santos AS, Strynadka NCJ, Finlay BB. 2017. Assembly, structure, function and regulation of type III secretion systems. Nat Rev Microbiol 15:323-337.

6. Kuwae A, Ohishi M, Watanabe M, Nagai M, Abe A. 2003. BopB is a type III secreted protein in *Bordetella bronchiseptica* and is required for cytotoxicity against cultured mammalian cells. Cell Microbiol 5:973-83.

7. Nogawa H, Kuwae A, Matsuzawa T, Abe A. 2004. The type III secreted protein BopD in *Bordetella bronchiseptica* is complexed with BopB for pore formation on the host plasma membrane. J Bacteriol 186:3806-13.

8. Yuk MH, Harvill ET, Cotter PA, Miller JF. 2000. Modulation of host immune

responses, induction of apoptosis and inhibition of NF-kappaB activation by the

*Bordetella* type III secretion system. Mol Microbiol 35:991-1004.

9. Panina EM, Mattoo S, Griffith N, Kozak NA, Yuk MH, Miller JF. 2005. A genome-wide

screen identifies a Bordetella type III secretion effector and candidate effectors in

other species. Mol Microbiol 58:267-79.

10. Kuwae A, Matsuzawa T, Ishikawa N, Abe H, Nonaka T, Fukuda H, Imajoh-Ohmi S,

Abe A. 2006. BopC is a novel type III effector secreted by *Bordetella bronchiseptica*

and has a critical role in type III-dependent necrotic cell death. J Biol Chem

281:6589-600.

11. Nagamatsu K, Kuwae A, Konaka T, Nagai S, Yoshida S, Eguchi M, Watanabe M,

Mimuro H, Koyasu S, Abe A. 2009. *Bordetella* evades the host immune system by

inducing IL-10 through a type III effector, BopN. J Exp Med 206:3073-88.

12. Abe A, Nishimura R, Kuwae A. 2017. *Bordetella* effector BopN is translocated into

host cells via its N-terminal residues. Microbiol Immunol 61:206-214.

13. Ahuja U, Liu M, Tomida S, Park J, Souda P, Whitelegge J, Li H, Harvill ET, Parkhill J,

Miller JF. 2012. Phenotypic and genomic analysis of hypervirulent human-associated

*Bordetella bronchiseptica*. BMC Microbiol 12:167.

14. Kurushima J, Kuwae A, Abe A. 2012. The type III secreted protein BspR regulates

the virulence genes in *Bordetella bronchiseptica*. PLoS One 7:e38925.

15. Medhekar B, Shrivastava R, Mattoo S, Gingery M, Miller JF. 2009. *Bordetella* Bsp22

forms a filamentous type III secretion system tip complex and is immunoprotective in

vitro and in vivo. Mol Microbiol 71:492-504.

16.   Ahuja U, Shokeen B, Cheng N, Cho Y, Blum C, Coppola G, Miller JF. 2016.

Differential regulation of type III secretion and virulence genes in *Bordetella*

*pertussis* and *Bordetella bronchiseptica* by a secreted anti-sigma factor. Proc Natl

Acad Sci U S A 113:2341-8.

17.   Feldman MF, Cornelis GR. 2003. The multitalented type III chaperones: all you can

do with 15 kDa. FEMS Microbiol Lett 219:151-8.

18.   Stebbins CE, Galán JE. 2001. Maintenance of an unfolded polypeptide by a cognate

chaperone in bacterial type III secretion. Nature 414:77-81.

19.   Lloyd SA, Norman M, Rosqvist R, Wolf-Watz H. 2001. *Yersinia* YopE is targeted for

type III secretion by N-terminal, not mRNA, signals. Mol Microbiol 39:520-31.

20.   Lee SH, Galán JE. 2004. *Salmonella* type III secretion-associated chaperones

confer secretion-pathway specificity. Mol Microbiol 51:483-95.

21.   Quinaud M, Chabert J, Faudry E, Neumann E, Lemaire D, Pastor A, Elsen S,

Dessen A, Attree I. 2005. The PscE-PscF-PscG complex controls type III secretion

needle biogenesis in *Pseudomonas aeruginosa*. J Biol Chem 280:36293-300.

22.   Wood SE, Jin J, Lloyd SA. 2008. YscP and YscU switch the substrate specificity of

the *Yersinia* type III secretion system by regulating export of the inner rod protein

YscI. J Bacteriol 190:4252-62.

23.  Monlezun L, Liebl D, Fenel D, Grandjean T, Berry A, Schoehn G, Dessein R, Faudry

E, Attree I. 2015. PscI is a type III secretion needle anchoring protein with in vitro

polymerization capacities. Mol Microbiol 96:419-36.

24.  Nishimura R, Abe A, Sakuma Y, Kuwae A. 2018. *Bordetella bronchiseptica* Bcr4

antagonizes the negative regulatory function of BspR via its role in type III secretion.

Microbiol Immunol 62:743-754.

25.  Wattiau P, Bernier B, Deslée P, Michiels T, Cornelis GR. 1994. Individual chaperones

required for Yop secretion by *Yersinia*. Proc Natl Acad Sci U S A 91:10493-7.

26.  Kamanova J. 2020. *Bordetella* Type III Secretion Injectosome and Effector Proteins.

Front Cell Infect Microbiol 10:466.

27.  Yuk MH, Harvill ET, Miller JF. 1998. The BvgAS virulence control system regulates

type III secretion in *Bordetella bronchiseptica*. Mol Microbiol 28:945-59.

28.  Kurushima J, Kuwae A, Abe A. 2012. Iron starvation regulates the type III secretion

system in *Bordetella bronchiseptica*. Microbiol Immunol 56:356-62.

29.  Sun P, Tropea JE, Austin BP, Cherry S, Waugh DS. 2008. Structural characterization

of the *Yersinia pestis* type III secretion system needle protein YscF in complex with

its heterodimeric chaperone YscE/YscG. J Mol Biol 377:819-30.

30.  Day JB, Guller I, Plano GV. 2000. *Yersinia pestis* YscG protein is a Syc-like

chaperone that directly binds yscE. Infect Immun 68:6466-71.

31.  Francis MS, Aili M, Wiklund ML, Wolf-Watz H. 2000. A study of the YopD-lcrH

interaction from *Yersinia pseudotuberculosis* reveals a role for hydrophobic residues

within the amphipathic domain of YopD. Mol Microbiol 38:85-102.

32.   Woestyn S, Sory MP, Boland A, Lequenne O, Cornelis GR. 1996. The cytosolic SycE

and SycH chaperones of *Yersinia* protect the region of YopE and YopH involved in

translocation across eukaryotic cell membranes. Mol Microbiol 20:1261-71.

33.   Yang H, Shan Z, Kim J, Wu W, Lian W, Zeng L, Xing L, Jin S. 2007. Regulatory role

of PopN and its interacting partners in type III secretion of *Pseudomonas aeruginosa*.

J Bacteriol 189:2599-609.

34.   Nguyen VS, Jobichen C, Tan KW, Tan YW, Chan SL, Ramesh K, Yuan Y, Hong Y,

Seetharaman J, Leung KY, Sivaraman J, Mok YK. 2015. Structure of AcrH-AopB

Chaperone-Translocator Complex Reveals a Role for Membrane Hairpins in Type III

Secretion System Translocon Assembly. Structure 23:2022-31.

35.   Quinaud M, Ple S, Job V, Contreras-Martel C, Simorre JP, Attree I, Dessen A. 2007.

Structure of the heterotrimeric complex that regulates type III secretion needle

formation. Proc Natl Acad Sci U S A 104:7803-8.

36.   Studier FW, Moffatt BA. 1986. Use of bacteriophage T7 RNA polymerase to direct

selective high-level expression of cloned genes. J Mol Biol 189:113-30.

37.   Stainer DW, Scholte MJ. 1970. A simple chemically defined medium for the

production of phase I *Bordetella pertussis*. J Gen Microbiol 63:211-20.

38.   Goto M, Hanawa T, Abe A, Kuwae A. 2020. Transcriptional Downregulation of a Type

III Secretion System under Reducing Conditions in *Bordetella pertussis*. J Bacteriol

202.

39.  Kurushima J, Kuwae A, Abe A. 2012. Btc22 chaperone is required for secretion and

stability of the type III secreted protein Bsp22 in *Bordetella bronchiseptica*. FEMS

Microbiol Lett 331:144-51.

40.  Rappl C, Deiwick J, Hensel M. 2003. Acidic pH is required for the functional

assembly of the type III secretion system encoded by *Salmonella* pathogenicity

island 2. FEMS Microbiol Lett 226:363-72.

 **Figure legends**

**Fig. 1. The interaction of Bcr4 with inner rod protein BscI.** (A) The *bcr4* gene and its

peripheral genes localized in the T3SS apparatus locus (*bsc* locus) on the *B.*

*bronchiseptica* S798 chromosome are depicted. (B) The purified Bcr4 (Bcr4-Strep) was

loaded on Strep-Tactin beads. Then, the Bcr4-Strep-loaded beads (Bcr4) and the beads

alone (beads) were mixed with the lysates prepared from *E. coli* BL21 producing BscI,

BscK or BscJ tagged with V5 (BscI-V5, BscK-V5 or BscJ-V5), respectively. After 3 h

incubation at 4□, each supernatant was prepared as the supernatant fraction (Sup) sample,

and each pellet was washed with TBS or TBS containing 0.1% Triton X-100 (Triton) and

prepared as the pellet fraction (Pellet) sample. The Sup and Pellet samples were separated

by SDS-PAGE and analyzed by Western blotting (WB) with anti-V5 antibody (top). Pellet

samples were also stained with Coomassie Brilliant Blue (CBB, bottom). (C) The purified

BscI (BscI-Strep) or N-terminal moiety of BteA (amino acids region 1–312, BteA-N-Strep)

were loaded on Strep-Tactin beads. Then, the beads were mixed with the lysate prepared

from *E. coli* BL21 to produce Bcr4 tagged with V5 (Bcr4-V5) at 4□ for 3 h. The prepared

Sup and Pellet samples were separated by SDS-PAGE and analyzed by WB with anti-V5

antibody (top). Pellet samples were also stained with CBB (bottom). Experiments were

performed at least three times, and representative data are shown.

**Fig. 2. Pull-down assays between BscI and truncated versions of Bcr4.** (A) Bcr4

derivatives used for pull-down assay are depicted.   (B) The purified Bcr4-FL, Bcr4Δ58-109

or Bcr4Δ110-173 (Bcr4-FL-Strep, Bcr4Δ58-109-Strep or Bcr4Δ110-17-Strep) was loaded on

Strep-Tactin beads, and then the beads were mixed with the lysate prepared from *E. coli*

BL21, producing BscI-V5, respectively. After 3 h incubation at 4□, the supernatant fraction

(Sup) and pellet fraction (Pellet) samples were prepared. The Sup and Pellet samples were

separated by SDS-PAGE and analyzed by Western blotting (WB) with anti-V5 antibody

(top), and stained with Coomassie Brilliant Blue (CBB, bottom). Experiments were

performed at least three times, and representative data are shown.

**Fig. 3. The Bcr4 domain required for T3SS activity** (A) Bcr4 derivatives used for the

analysis of T3SS activity are depicted.   (B) The *B. bronchiseptica* wild-type strain,

Bcr4-deficient strain (Δ*bcr4*), and Δ*bcr4* producing Bcr4-FL-FLAG (Δ*bcr4*+*bcr4*-FL-FLAG),

Bcr4Δ169-173-FLAG   (Δ*bcr4*+*bcr4*Δ169-173-FLAG),   Bcr4Δ164-173-FLAG   (Δ*bcr4*+*bcr4*

Δ164-173-FLAG), and Bcr4Δ159-173-FLAG (Δ*bcr4*+*bcr4*Δ159-173-FLAG) were cultured in

SS medium. Whole cell lysates (WCL) and culture supernatant fraction (CS) samples were

separated by SDS-PAGE and analyzed by Western blotting using antibodies against Bcr4,

FLAG, BteA, BopD or RpoB. RpoB was used as an internal loading control. AP indicates

the amino acids region of the antigen peptide used for anti-Bcr4 antibody generation. WCL

and CS samples were prepared from equal volumes of bacterial culture. When we

performed Western blotting using anti-BteA or BopD antibodies, we loaded a 100- or

50-fold smaller amount of Δ*bcr4*+*bcr4*-FL-FLAG, Δ*bcr4*+*bcr4*Δ169-173-FLAG and

Δ*bcr4*+*bcr4*Δ164-173-FLAG CS samples on the SDS-PAGE gel than on the wild-type,

Δ*bcr4* and Δ*bcr4*+*bcr4*Δ159-173-FLAG CS samples to avoid obtaining excess signal

intensities. The numbers at the bottom of the lower panel indicate the relative signal

intensity of RpoB measured using ImageJ software. (C) L2 cells were infected with each

strain at an moi of 50 for 1 h. The amounts of LDH released into the extracellular medium

from infected cells are shown, and the relative cytotoxicity (percent) was determined as

described in the Materials and Methods. Error bars indicate the SEMs from triplicate

experiments. *$P$<0.05. Experiments were performed at least three times, and

representative data are shown.

**Fig. 4. The effect of BscI deletion on T3SS activity in *B. bronchiseptica*.** (A) Whole cell

lysates (WCL) and culture supernatant fraction (CS) samples were prepared from the

wild-type strain, Δ*bscI* (BscI-deficient strain), Δ*bscI*+*bscI* (BscI-complemented strain) and

Δ*bscN* (T3SS-inactive strain) cultured in SS medium. WCL and CS samples were

separated by SDS-PAGE and analyzed by Western blotting using antibodies against BscI,

BteA, BopD, Bsp22, FHA or RpoB. BscI tagged with Strep (BscI-Strep) was loaded as a

control. *BscI-Strep signal; **NS (nonspecific signals); ***A faint signal of BscI. The

numbers at the bottom of the lower panel indicate the relative signal intensity of RpoB

measured using ImageJ software. (B) WCL and CS were prepared from the wild-type,

Δ*bspR* (BspR-deficient strain), Δ*bspR*Δ*bscI* (BspR- and BscI-deficient strain) and

Δ*bspR*Δ*bscI*+*bscI* (BscI-complemented BspR- and Bcr4-deficient strain) cultured in SS

medium. WCL and CS were separated by SDS-PAGE and analyzed by Western blotting

using antibodies against BscI, BspR, Bsp22 or BopB antibodies. Loaded WCL and CS

samples were prepared from equal volumes of bacterial culture. When we performed

Western blotting using antibodies against Bsp22 or BopB, we loaded a 25- or 5-fold smaller

amount of Δ*bspR*, Δ*bspR*Δ*bscI* and Δ*bspR*Δ*bscI*+*bscI* WCL samples, and 5- or 10-fold

smaller amount of Δ*bspR* and Δ*bspR*Δ*bscI*+*bscI* CS samples on the SDS-PAGE gel than

the wild-type to avoid obtaining excess signal intensities. Experiments were performed at

least three times, and representative data are shown.

**Fig. 5. Immunofluorescent staining of Bsp22 on L2 cells infected with *B.**

***bronchiseptica* and the results of the LDH assay.** (A) L2 cells were infected with the

wild-type strain, Δ*bscI* (BscI-deficient strain), Δ*bscI*+*bscI* (BscI-complemented strain) or

Δ*bscN* (T3SS-inactive strain) as described in the Materials and Methods. After fixation, cells

were stained with anti-Bsp22 antibody (green), rhodamine phalloidin (red) and DAPI (blue).

The inset shows higher magnification of the boxed area in the image. Arrowheads and

arrows indicate the signals of Bsp22 (green) and bacteria, respectively. (B) Bsp22 signals

per one cell were counted under a fluorescent microscope. At least 120 cells were

randomly chosen. (C) The results of LDH assay at an MOI of 200 are shown as a histogram.

Error bars indicate the SEMs from triplicate experiments. *P<0.05. Experiments were

performed at least three times, and representative data are shown.

**Fig. 6. The effect of Bcr4 on BscI stability and the suggestive interaction of Bcr4 with**

**BscI in *B. bronchiseptica*.** (A) The whole cell lysates (WCL) and culture supernatant

fraction (CS) were prepared from the wild-type strain, Δ*bcr4* (Bcr4-deficient strain), Δ*bspR*

(BspR-deficient strain) or Δ*bspR*Δ*bcr4* (BspR- and Bcr4-deficient strain) cultured in SS

medium. WCL and CS were separated by SDS-PAGE and analyzed by Western blotting

with antibodies against BscI (inner-rod protein), BscJ (inner-ring protein) and Bcr4. The

samples were prepared from equal volumes of bacterial culture. When we carried out

Western blotting using anti-BscJ or Bcr4 antibodies, we loaded a 10-fold-smaller amount of

Δ*bspR* or Δ*bspR*Δ*bcr4* WCL samples on the SDS-PAGE than WT or Δ*bcr4* WCL samples to

avoid obtaining excess signal intensities. (B) WCL were prepared from the wild-type, Δ*bspR*

or Δ*bspR*Δ*bcr4* treated with (+) or without (−) cross linker disuccinimidyl suberate (DSS).

WCL were separated by SDS-PAGE and analyzed by Western blotting with anti-Bcr4

antibody. The lower panel is a short exposure of the upper panel. Experiments were

performed at least three times, and representative data are shown.

**Fig. 7. Schematic depiction of the Bcr4 and BscI functions.** In the *B. bronchiseptica*

bacterial cytosol, Bcr4 (blue) stabilizes BscI (orange). Then, BscI is presumed to be

localized to the T3SS machinery (gray) and form the inner rod. BscI is essential for T3SS

activity and secreted from the bacterial cell. BscJ (pink) is predicted to function as an inner

membrane ring. Bsp22 (green) forms a filament-like structure. OM and IM indicate the outer

membrane (white) and inner membrane (white), respectively.

Table 1. Bacterial strains used in the study

| Name | Relevant genotype | Reference or source |
|---|---|---|
| *B. bronchiseptica* | | |
| S798 | wild-type strain | 6 |
| △*bscl* | Bscl-deficient strain | This study |
| △*bscN* | BscN-deficient strain | 6 |
| △*bcr4* | Bcr4-deficient strain | 24 |
| △*bspR* | BspR-deficient strain | 14 |
| △*bspR*△*bscl* | BspR and Bscl-deficient strain | This study |
| △*bspR*△*bcr4* | BspR- and Bcr4-deficient strain | 24 |
| *E. coli* | | |
| DH10B | Host strain for pDONR201, p99-*ccdB*-V5 and pRK415-R4-R3-F | Invitrogen |
| BL21 | Host strain for pColdII | 36 |
| Sm10λ*pir* | Host strain for pABB-CRS2 | 6 |

**Fig. 1. The interaction of Bcr4 with inner rod protein BscI.** (A) The *bcr4* gene and its peripheral genes localized in the T3SS apparatus locus (*bsc* locus) on the *B. bronchiseptica* S798 chromosome are depicted. (B) The purified Bcr4 (Bcr4-Strep) was loaded on Strep-Tactin beads. Then, the Bcr4-Strep-loaded beads (Bcr4) and the beads alone (beads) were mixed with the lysates prepared from *E. coli* BL21 producing BscI, BscK or BscJ tagged with V5 (BscI-V5, BscK-V5 or BscJ-V5), respectively. After 3 h incubation at 4°C, each supernatant was prepared as the supernatant fraction (Sup) sample, and each pellet was washed with TBS or TBS containing 0.1% Triton X-100 (Triton) and prepared as the pellet fraction (Pellet) sample. The Sup and Pellet samples were separated by SDS-PAGE and analyzed by Western blotting (WB) with anti-V5 antibody (top). Pellet samples were also stained with Coomassie Brilliant Blue (CBB, bottom). (C) The purified BscI (BscI-Strep) or N-terminal moiety of BteA (amino acids region 1–312, BteA-N-Strep) were loaded on Strep-Tactin beads. Then, the beads were mixed with the lysate prepared from *E. coli* BL21 to produce Bcr4 tagged with V5 (Bcr4-V5) at 4°C for 3 h. The prepared Sup and Pellet samples were separated by SDS-PAGE and analyzed by WB with anti-V5 antibody (top). Pellet samples were also stained with CBB (bottom). Experiments were performed at least three times, and representative data are shown.

Fig. 1

Fig. 2. Pull-down assays between BscI and truncated versions of Bcr4. (A) Bcr4 derivatives used for pull-down assay are depicted.  (B) The purified Bcr4-FL, Bcr4Δ58-109 or Bcr4Δ110-173 (Bcr4-FL-Strep, Bcr4Δ58-109-Strep or Bcr4Δ110-17-Strep) was loaded on Strep-Tactin beads, and then the beads were mixed with the lysate prepared from *E. coli* BL21, producing BscI-V5, respectively. After 3 h incubation at 4°C, the supernatant fraction (Sup) and pellet fraction (Pellet) samples were prepared. The Sup and Pellet samples were separated by SDS-PAGE and analyzed by Western blotting (WB) with anti-V5 antibody (top), and stained with Coomassie Brilliant Blue (CBB, bottom). Experiments were performed at least three times, and representative data are shown.

Fig. 2

A

Bcr4

Bcr4-FL    1 ——————————————— 173

Bcr4Δ169-173    1 —————————————— 168

Bcr4Δ164-173    1 ————————————— 163

Bcr4Δ159-173    1 ———————————— 158

B

|  | WCL |
|---|---|
| Bcr4 (AP: 156-173 aa) | |
| FLAG | |

|  | CS |
|---|---|
| BteA multimer | |
| BopD | |
| RpoB | |

1.00  1.10  1.05  1.02  1.20  1.25

C

**Fig. 3. The Bcr4 domain required for T3SS activity** (A) Bcr4 derivatives used for the analysis of T3SS activity are depicted. (B) The *B. bronchiseptica* wild-type strain, Bcr4-deficient strain (Δ*bcr4*), and Δ*bcr4* producing Bcr4-FL-FLAG (Δ*bcr4*+*bcr4*-FL-FLAG), Bcr4Δ169-173-FLAG (Δ*bcr4*+*bcr4*Δ169-173-FLAG), Bcr4Δ164-173-FLAG (Δ*bcr4*+*bcr4* Δ164-173-FLAG), and Bcr4Δ159-173-FLAG (Δ*bcr4*+*bcr4*Δ159-173-FLAG) were cultured in SS medium. Whole cell lysates (WCL) and culture supernatant fraction (CS) samples were separated by SDS-PAGE and analyzed by Western blotting using antibodies against Bcr4, FLAG, BteA, BopD or RpoB. RpoB was used as an internal loading control. AP indicates the amino acids region of the antigen peptide used for anti-Bcr4 antibody generation. WCL and CS samples were prepared from equal volumes of bacterial culture. When we performed Western blotting using anti-BteA or BopD antibodies, we loaded a 100- or 50-fold smaller amount of Δ*bcr4*+*bcr4*-FL-FLAG, Δ*bcr4*+*bcr4*Δ169-173-FLAG and Δ*bcr4*+*bcr4*Δ164-173-FLAG CS samples on the SDS-PAGE gel than on the wild-type, Δ*bcr4* and Δ*bcr4*+*bcr4*Δ159-173-FLAG CS samples to avoid obtaining excess signal intensities. The numbers at the bottom of the lower panel indicate the relative signal intensity of RpoB measured using ImageJ software. (C) L2 cells were infected with each strain at an moi of 50 for 1 h. The amounts of LDH released into the extracellular medium from infected cells are shown, and the relative cytotoxicity (percent) was determined as described in the Materials and Methods. Error bars indicate the SEMs from triplicate experiments. *$P<0.05$. Experiments were performed at least three times, and representative data are shown.

Fig. 3

**Fig. 4. The effect of BscI deletion on T3SS activity in *B. bronchiseptica*.** (A) Whole cell lysates (WCL) and culture supernatant fraction (CS) samples were prepared from the wild-type strain, Δ*bscI* (BscI-deficient strain), Δ*bscI*+*bscI* (BscI-complemented strain) and Δ*bscN* (T3SS-inactive strain) cultured in SS medium. WCL and CS samples were separated by SDS-PAGE and analyzed by Western blotting using antibodies against BscI, BteA, BopD, Bsp22, FHA or RpoB. BscI tagged with Strep (BscI-Strep) was loaded as a control. *BscI-Strep signal; **NS (nonspecific signals); ***A faint signal of BscI. The numbers at the bottom of the lower panel indicate the relative signal intensity of RpoB measured using ImageJ software. (B) WCL and CS were prepared from the wild-type, Δ*bspR* (BspR-deficient strain), Δ*bspR*Δ*bscI* (BspR- and BscI-deficient strain) and Δ*bspR*Δ*bscI*+*bscI* (BscI-complemented BspR- and Bcr4-deficient strain) cultured in SS medium. WCL and CS were separated by SDS-PAGE and analyzed by Western blotting using antibodies against BscI, BspR, Bsp22 or BopB antibodies. Loaded WCL and CS samples were prepared from equal volumes of bacterial culture. When we performed Western blotting using antibodies against Bsp22 or BopB, we loaded a 25- or 5-fold smaller amount of Δ*bspR*, Δ*bspR*Δ*bscI* and Δ*bspR*Δ*bscI*+*bscI* WCL samples, and 5- or 10-fold smaller amount of Δ*bspR* and Δ*bspR*Δ*bscI*+*bscI* CS samples on the SDS-PAGE gel than the wild-type to avoid obtaining excess signal intensities. Experiments were performed at least three times, and representative data are shown.

Fig. 4

**Fig. 5. Immunofluorescent staining of Bsp22 on L2 cells infected with *B. bronchiseptica* and the results of the LDH assay.** (A) L2 cells were infected with the wild-type strain, Δ*bscI* (BscI-deficient strain), Δ*bscI*+*bscI* (BscI-complemented strain) or Δ*bscN* (T3SS-inactive strain) as described in the Materials and Methods. After fixation, cells were stained with anti-Bsp22 antibody (green), rhodamine phalloidin (red) and DAPI (blue). The inset shows higher magnification of the boxed area in the image. Arrowheads and arrows indicate the signals of Bsp22 (green) and bacteria, respectively. (B) Bsp22 signals per one cell were counted under a fluorescent microscope. At least 120 cells were randomly chosen. (C) The results of LDH assay at an MOI of 200 are shown as a histogram. Error bars indicate the SEMs from triplicate experiments. *$P$<0.05. Experiments were performed at least three times, and representative data are shown.

Fig. 5

**Fig. 6. The effect of Bcr4 on BscI stability and the suggestive interaction of Bcr4 with BscI in *B. bronchiseptica*.** (A) The whole cell lysates (WCL) and culture supernatant fraction (CS) were prepared from the wild-type strain, Δ*bcr4* (Bcr4-deficient strain), Δ*bspR* (BspR-deficient strain) or Δ*bspR*Δ*bcr4* (BspR- and Bcr4-deficient strain) cultured in SS medium. WCL and CS were separated by SDS-PAGE and analyzed by Western blotting with antibodies against BscI (inner-rod protein), BscJ (inner-ring protein) and Bcr4. The samples were prepared from equal volumes of bacterial culture. When we carried out Western blotting using anti-BscJ or Bcr4 antibodies, we loaded a 10-fold-smaller amount of Δ*bspR* or Δ*bspR*Δ*bcr4* WCL samples on the SDS-PAGE than WT or Δ*bcr4* WCL samples to avoid obtaining excess signal intensities. (B) WCL were prepared from the wild-type, Δ*bspR* or Δ*bspR*Δ*bcr4* treated with (+) or without (–) cross linker disuccinimidyl suberate (DSS). WCL were separated by SDS-PAGE and analyzed by Western blotting with anti-Bcr4 antibody. The lower panel is a short exposure of the upper panel. Experiments were performed at least three times, and representative data are shown.

Fig. 6

**Fig. 7. Schematic depiction of the Bcr4 and BscI functions.** In the *B. bronchiseptica* bacterial cytosol, Bcr4 (blue) stabilizes BscI (orange). Then, BscI is presumed to be localized to the T3SS machinery (gray) and form the inner rod. BscI is essential for T3SS activity and secreted from the bacterial cell. BscJ (pink) is predicted to function as an inner membrane ring. Bsp22 (green) forms a filament-like structure. OM and IM indicate the outer membrane (white) and inner membrane (white), respectively.

Fig. 7

**Supplemental material**

**Text S. Supplementary Information**

**Fig. S1. Construction of the T3SS machinery in strains lacking Bcr4 and/or BspR.** In

the *B. bronchiseptica* wild-type (upper left), the BspR negative regulation level for the *bsc*

locus transcription is moderate, and the T3SS machinery is established. In the Bcr4-deficient

strain (upper right), BspR strongly represses the *bsc* locus transcription, and construction of

the T3SS machinery is incomplete. In the BspR-deficient strain (lower left), the negative

regulatory effect of BspR is cancelled, and the construction of the T3SS machinery is

promoted. In the BspR/Bcr4 double-deficient strain (lower right), while the *bsc* locus

transcription is promoted because of BspR deficiency, T3SS is not functional.

**Fig. S2. Alignment of Bcr4 amino acid sequences in representative *Bordetella* species.**

Bcr4 amino acid sequences of *B. bronchiseptica* S798 (Bb), *B. pertussis* Tohama I (Bp) and

*B. parapertussis* 12822 (Bp) were compared using ClustalW. The grey highlighted letters

represent amino acid residues that were different from those in Bb. Bcr4 of Bp and Bpp have

98.3% and 97.1% identities with those of Bp, respectively.

**Fig. S3. The nonspecific reaction of anti-BscI antibody to Bsp22.** The culture

supernatants (CS) were prepared from the wild-type strain, Δ*bsp22* (Bsp22-deficient strain),

Δ*bspR* (BspR-deficient strain) or Δ*bscI+bscI* (BscI-complemented strain) cultured in SS

medium. The CS samples were separated by SDS-PAGE and stained with Coomassie

Brilliant Blue (CBB, left panel) or analyzed by Western blotting (WB) with anti-BscI antibody

(right panel). NS indicates nonspecific signals. Experiments were performed at least three

times, and representative data are shown.

**Fig. S4. The results of the RT-PCR analysis for the mRNA level of *bscI* in *B.**

***bronchiseptica* strains.** Total RNA was prepared from the wild-type strain, Δ*bspR* (BspR-

deficient strain) or Δ*bspR*Δ*bcr4* (BspR- and Bcr4-deficient strain) cultured in SS medium and

subjected to a quantitative RT-PCR analysis. The histogram shows the relative amount of

*bscI* mRNA normalized by the housekeeping gene, *recA* mRNA. Experiments were

performed at least three times, and representative data are shown.

**Fig. S5. The results of the RT-PCR analysis for mRNA levels of *bopD* and *bscI* in the**

**wild-type *B. bronchiseptica*.** Total RNA was prepared from the wild-type strain cultured in

SS medium and subjected to a quantitative RT-PCR analysis. The histogram shows the

relative amount of *bopD* and *bscI* mRNA in the wild-type. The relative ratio of *bscI* mRNA is

shown when the *bopD* mRNA amount is set as 1. Experiments were performed at least three

times, and representative data are shown.

**Fig. S6. The time course of BscI production in _B. bronchiseptica_.** The whole cell lysates

(WCL) were prepared from the wild-type strain, Δ*bcr4* (Bcr4-deficient strain), Δ*bspR* (BspR-

deficient strain) or Δ*bspR*Δ*bcr4* (BspR- and Bcr4-deficient strain) cultured in SS medium for

0, 2, 5, 8 or 18 hr. The WCL were separated by SDS-PAGE and analyzed by Western blotting

with antibodies against BscI, BopB and RpoB. Experiments were performed at least three

times, and representative data are shown.

**Fig. S7. The predicted structural model of _B. bronchiseptica_ Bcr4**

(A) The AlphaFold2 (AF2)-predicted structural model of _B. bronchiseptica_ Bcr4 and

structural comparison with _Aeromonas hydrophila_ AcrH (PDB: 3WXX) and _Pseudomonas_

_aeruginosa_ PscG (PDB: 2UWJ). Z-score, root mean square deviation (RMSD), and amino

acid identity (AA %ID) of AcrH and PscG compared with Bcr4 are shown, respectively. (B)

The pairwise sequence alignment of Bcr4, AcrH, and PscG. The most frequent amino acid

type is colored. The secondary structure assignments (H/h: helix, E/e: strand, L/l: coil) are

also shown.

61      **Table S1. Nomencalture of the *Bordetella* T3SS component**

63      **Table S2. Plasmids used in the study**

65      **Table S3. Primers used in the study**

**Supplementary Information**

**SI Materials and Methods**

**Pull-down assay.**

The plasmids and primers used in this study are listed in Table S1 and S2, respectively. In order to express *bscI*, *bscK*, *bscJ* or *bcr4* tagged with a V5 sequence at the respective C-terminus (*bscI*-V5, *bscK*-V5, *bscJ*-V5 or *bcr4*-V5), we amplified DNA fragments encoding the *bscI*, *bscK*, *bscJ* or *bcr4* genes with the primer sets of B1-bscI-V5 and B2-bscI-V5, B1-bscK-V5 and B2-bscK-V5, B1-bscJ-V5 and B2-bscJ-V5 or B1-bcr4-comp and B2-bcr4-V5, respectively, using *B. bronchiseptica* S798 genomic DNA as the template. Each resulting PCR product was cloned into pDONR201 to obtain pMGKU404, pMGKU405, pMGKU406 or pMGKU407, respectively, by means of adapter PCR and site-specific recombination techniques using the Gateway cloning system (Invitrogen). Each plasmid was mixed with an expression vector such as p99*ccdB*-V5 (1) to obtain pMGKU408, pMGKU409, pMGKU410 or pMGKU411 using the Gateway cloning system.

BL21 cells carrying pMGKU408, pMGKU409 pMGKU410 or pMGKU411 were cultured overnight at 37°C with shaking, and then diluted 1:40 in LB liquid medium containing 50 $\mu$L/mL ampicillin and incubated for 2 h at 37°C with shaking. Each bacterial culture was further incubated for 5 h at 30°C in the presence of isopropyl-beta-thiogalactopyranoside (IPTG) at the final concentration of 1 mM. Bacteria were collected by centrifugation at 2,600×$g$ for 15 min, and suspended in cold TBS containing protease inhibitor cocktail,

cOmplete (Roche). Each bacterial suspension was sonicated, and each supernatant was

used for the pull-down assay.

Next, in order to purify Bcr4 or BscI tagged with six histidine residues (6×His) at the

respective N-terminus and Strep at the respective C-terminus, we amplified DNA fragments

encoding *bcr4* or *bscI* with the primer sets of 5-HindIII-bcr4 and 3-bcr4-Strep, or 5-HindIII-

bscI and 3-bscI-Strep using *B. bronchiseptica* S798 genomic DNA as the template. Each

amplified DNA fragment was used as a template for 2nd PCR with a primer set consisting

of the upper primer used in the 1st PCR and 3-Strep-HindIII to add a 24 bp sequence

encoding the Strep tag. Each resulting PCR product was cloned into the HindIII recognition

sites of pColdII to obtain pMGKU412 or pMGKU413 by the In-Fusion Cloning System

(Clontech), respectively. In order to purify Bcr4 lacking the amino acids region 58–109 or

110–173, we amplified the DNA fragment with the primer sets of 5-bcr4-58-109-Strep-IF and

3-IF-bcr4-58-109-Strep, or 5-bcr4-110-173-Strep-IF and 3-IF-bcr4-110-173-Strep using

pColdII-*bcr4*-Strep as the template. Each amplified fragment was self-ligated by the In-

Fusion Cloning System, respectively, and then designated pMGKU414 or pMGKU415.

BL21 cells carrying pMGKU412, pMGKU413, pColdII-*bteA*-N-Strep (2), pMGKU414 or

pMGKU415 were cultured overnight at 37°C with shaking, and then diluted 1:100 in LB liquid

medium containing 50 $\mu$L/mL ampicillin and incubated for 2 h at 37°C with shaking,

respectively. Each bacterial culture was further incubated overnight at 15°C in the presence

of IPTG at the final concentration of 0.05 mM. Bacteria were collected by centrifugation at

2,600×*g* for 20 min, and suspended in cold TBS containing cOmplete. Each bacterial

suspension was sonicated and each supernatant except for that of pMGKU413 was

subjected to purification using Ni-NTA agarose (Qiagen) according to the manufacturer's

instructions. The purified proteins were dialyzed with TBS. As for pMGKU413, the bacterial

suspension was sonicated and the pellet was suspended in Inclusion Body Solubilization

Reagent (Thermo) according to the manufacturer's instructions. The BscI-Strep in

solubilized solution was refolded as described previously (3). Briefly, the solubilized solution

was diluted with 5-fold larger refolding buffer (20 mM Tris-HCl, pH-8, 150 mM NaCl, 10%

glycerol) overnight at 4°C on a rotator, and then centrifugated at 20,000×*g* for 15 min. The

supernatant was dialyzed with dialysis buffer (20 mM Tris-HCl, pH-8, 150 mM NaCl, 10%

glycerol), and then centrifugated at 20,000×*g* for 15 min. The supernatant was used as

purified protein for the pull-down assay.

We mixed the dual-tagged protein (6×His and Strep) and 30 $\mu$l Strep-Tactin resin (IBA) in

an Eppendorf tube and rotated the tube at 4°C for 1 h. Then, we washed the beads with

TBS three times. Next, the V5-tagged protein-containing *E. coli* lysate was added to the tube

and rotated at 4°C for 3 h. We transferred 30 $\mu$L supernatant to new Eppendorf tube and

added 30 $\mu$L 2×SDS-PAGE sample buffer to prepare the Sup. Then we washed the beads

with TBS (Fig. 1B and 1C) or TBS containing 0.1% Triton X-100 (Fig. 1C) three times and

added 30 $\mu$L 2×SDS-PAGE sample buffer to prepare the Pellet samples.

**Construction of a *bscI* gene-disrupted or *bspR/bscI* double strains.**

To construct the *bscI* or *bspR/bscI* double mutants, a 2.4 kb DNA fragment encoding *bscI*

and its flanking regions was amplified by PCR with primers B1-bscI and B2-bscI using *B.*

*bronchiceptica* S798 genomic DNA as the template. The resulting PCR product was cloned

into pDONR201 to obtain pMGKU416 using the Gateway cloning system. An inverse PCR

was carried out with the primers R1-bscI and R2-bscI using circular pMGKU416 as the

template. The resulting PCR product was self-ligated using the In-Fusion Cloning System to

obtain pMGKU417. This plasmid contained a 369-bp in-frame deletion from 30 bp

downstream of the 5' end of the *bscI* gene to 30 bp upstream of the 3' end of the gene. This

plasmid, pMGKU417, was mixed with a positive suicide vector such as pABB-CRS2 (4) to

obtain pMGKU418 using the Gateway cloning system. The pMGKU418 or pABB-CRS2-

*bspR* (5) plasmids in turn were   introduced into *E. coli* Sm10λ*pir,* and transconjugated into

the S798 wild-type or Δ*bscI* as described previously (6). The resulting mutant strains were

designated Δ*bscI* or Δ*bspR*Δ*bscI,* respectively.

**Construction of plasmids used for producing Bcr4 derivatives and BscI**

**complementation**

In order to produce full-length Bcr4 tagged with a FLAG sequence at the C-terminus, we

performed an inverse PCR with primers of 5-bcr4-FLAG-IF and 3-IF-bcr4-FLAG using

pDONR-*bcr4* (7) as the template. The resulting PCR product was self-ligated to obtain

pMGKU419 by the In-Fusion Cloning System. To produce Bcr4 lacking the amino acid

regions 159–173, 164–173 or 169–173, inverse PCR was carried out with the primer sets of

5-*bcr4*-FLAG-IF and 3-IF-bcr4Δ169-173-FLAG, 5-*bcr4*-FLAG-IF and 3-IF-*bcr4*Δ164-173-

FLAG, or 5-*bcr4*-FLAG-IF and 3-IF-*bcr4*Δ159-173-FLAG using pMGKU419 as the template.

Each amplified fragment was self-ligated by the In-Fusion Cloning System to obtain

pMGKU420, pMGKU421 or pMGKU422, respectively. For BscI complementation, a PCR

was carried out with the primers of B1-bscI-V5 and 3-B2-bscI-comp using *B. bronchiceptica*

S798 genomic DNA as the template. The resulting PCR product was cloned into pDONR201

to obtain pMGKU423 using the Gateway cloning system. These plasmids were mixed with

pRK-R4-R3-F, pDONR-*fha*P and pDONR-*rrnB* (8) to obtain pMGKU424, pMGKU425,

pMGKU426, pMGKU427 and pMGKU428 using the Gateway cloning system.

**Quantitative reverse transcription-PCR**

Total RNA was prepared from the *B. bronchiseptica* culture using a Trizol Max Bacterial RNA

isolation Kit (Invitrogen), RNeasy Mini Kit (Qiagen), and RNase-free DNase Kit (Qiagen).

The reverse transcription reaction was carried out using Transcriptor Universal cDNA Master

(Roche) and T100 Thermal Cycler (Bio-Rad). The quantitative RT-PCR reaction was carried

out using FastStart Essential DNA Probes Master (Roche) and Light Cycler 96 (Roche). To

amplify the *bscI*, *bopD*, and *recA* genes, the primer sets 5-bscI and 3-bscI, 5-bopD and 3-

bopD, 5-recA and 3-recA were used, respectively. For the experiment to determine the

presence of *bscI* mRNA in the Bcr4 mutant (Fig. S4), *recA* was used as an internal control.

The values obtained from each strain were standardized with *recA*, and the relative amounts

to the wild-type strain were determined. For the experiment to compare the amounts of *bscI*

and *bopD* mRNA in the wild-type (Fig. S5), we followed the protocol provided by Roche. The

genome of *B. bronchiseptica* was used as a template for quantitative RT-PCR to generate a

calibration curve. The relative amount of *bopD* mRNA to *bscI* mRNA was then determined

using the calibration curve.

**Protein structure prediction**

Protein structure prediction using amino acid sequences by AlphaFold2 with MMseqs2

(ColabFold) was performed on the Google Colab server with default parameters (9). A

structure-based protein homology search using AlphaFold2-predicted structural models was

performed on the Dali server (10). In addition, primary, secondary, and tertiary structures of

query proteins were visualized and compared with those of neighbors on the Dali server.

**References**

202     1.     Ogino T, Ohno R, Sekiya K, Kuwae A, Matsuzawa T, Nonaka T, Fukuda H, Imajoh-

203            Ohmi S, Abe A. 2006. Assembly of the type III secretion apparatus of

204            enteropathogenic *Escherichia coli*. J Bacteriol 188:2801-11.

205     2.     Kuwae A, Momose F, Nagamatsu K, Suyama Y, Abe A. 2016. BteA Secreted from the

206            *Bordetella bronchiseptica* Type III Secetion System Induces Necrosis through an

207            Actin Cytoskeleton Signaling Pathway and Inhibits Phagocytosis by Macrophages.

208            PLoS One 11:e0148387.

209     3.     Halder PK, Roy C, Datta S. 2019. Structural and functional characterization of type

210            three secretion system ATPase PscN and its regulator PscL from *Pseudomonas*

211            *aeruginosa*. Proteins 87:276-288.

212     4.     Sekiya K, Ohishi M, Ogino T, Tamano K, Sasakawa C, Abe A. 2001. Supermolecular

213            structure of the enteropathogenic *Escherichia coli* type III secretion system and its

214            direct interaction with the EspA-sheath-like structure. Proc Natl Acad Sci U S A

215            98:11638-43.

216     5.     Kurushima J, Kuwae A, Abe A. 2012. The type III secreted protein BspR regulates

217            the virulence genes in *Bordetella bronchiseptica*. PLoS One 7:e38925.

218     6.     Donnenberg MS, Kaper JB. 1991. Construction of an eae deletion mutant of

219            enteropathogenic *Escherichia coli* by using a positive-selection suicide vector. Infect

220            Immun 59:4310-7.

7.    Nishimura R, Abe A, Sakuma Y, Kuwae A. 2018. *Bordetella bronchiseptica* Bcr4

antagonizes the negative regulatory function of BspR via its role in type III secretion.

Microbiol Immunol 62:743-754.

8.    Kuwae A, Matsuzawa T, Ishikawa N, Abe H, Nonaka T, Fukuda H, Imajoh-Ohmi S,

Abe A. 2006. BopC is a novel type III effector secreted by *Bordetella bronchiseptica*

and has a critical role in type III-dependent necrotic cell death. J Biol Chem 281:6589-

600.

9.    Mirdita M, Schutze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. 2022.

ColabFold: making protein folding accessible to all. Nat Methods 19:679-682.

10.   Holm L. 2020. Using Dali for Protein Structure Comparison. Methods Mol Biol

2112:29-42.

| Table S1. Nomencalture of the *Bordetella* T3SS | | | | |
|---|---|---|---|---|
| Unified name | *Bordetella* | *Yersinia* | *Pseudomonas* | Predicted function |
| SctF | BscF | YscF | PscF | Needle |
| SctI | BscI | YscI | PscI | Inner rod |
| SctE | BopB | YopB | PopB | Translocation pore |
| SctB | BopD | YopD | PopD | Translocation pore |
| SctA | Bsp22 | LcrV | PcrV | Needle tip or Filament |
| SctJ | BscJ | YscJ | PscJ | Inner membrane ring |
| SctK | BscK | YscK | PscK | ATPase cofactor |
| SctN | BscN | YscN | PscN | ATPase |

| Table S2. Plasmids used in the study | | |
|---|---|---|
| Name | Description | Reference or source |
| pDONR201 | DNA cloning vector, Km[r] | Invitrogen |
| pMGKU404 | pDONR201, *bscI* gene tagged with V5 sequence at the C terminus | This study |
| pMGKU405 | pDONR201, *bscK* gene tagged with V5 sequence at the C terminus | This study |
| pMGKU406 | pDONR201, *bscJ* gene tagged with V5 sequence at the C terminus | This study |
| pMGKU407 | pDONR201, *bcr4* gene tagged with V5 sequence at the C terminus | This study |
| p99*ccdB*-V5 | Expression vector for V5 tagged gene, Amp[r] | 1 |
| pMGKU408 | p99-*ccdB*-V5, *bscI* gene tagged with V5 sequence at the C terminus | This study |
| pMGKU409 | p99-*ccdB*-V5, *bscK* gene tagged with V5 sequence at the C terminus | This study |
| pMGKU410 | p99-*ccdB*-V5, *bscJ* gene tagged with V5 sequence at the C terminus | This study |
| pMGKU411 | p99-*ccdB*-V5, *bsc4* gene tagged with V5 sequence at the C terminus | This study |
| pColdII | Expression vector for His tagged gene, Amp[r] | TAKARA |
| pMGKU412 | pColdII, *bcr4* gene tagged with Strep sequence at the C terminus | This study |
| pMGKU413 | pColdII, *bscI* gene tagged with Strep sequence at the C terminus | This study |
| pMGKU414 | pColdII, *bcr4* gene lacking amino acids region 58-109 tagged with Strep sequence at the C terminus | This study |
| pMGKU415 | pColdII, *bcr4* gene lacking amino acids region 110-173 tagged with Strep sequence at the C terminus | This study |
| pColdII-*bteA*-N-Strep | pColdII, *bteA* gene coding amino acids region 1-312 tagged with Strep sequence at the C terminus | 2 |
| pMGKU416 | pDONR201, *bscI* gene | This study |
| pMGKU417 | pDONR201, *bscI* gene containing internal sequence-deletion and its flanking region | This study |
| pABB-CRS2 | Suicide vector for conjugation, Amp[r] | 4 |
| pMGKU418 | pABB-CRS2, *bscI* gene containing internal sequence-deletion and its flanking region | This study |
| pABB-CRS2-*bspR* | pABB-CRS2, *bspR* gene containing internal sequence-deletion and its flanking region | 5 |
| pDONR-*bcr4* | pDONR201, *bcr4* gene for complementation | 7 |
| pMGKU419 | pDONR201, *bcr4* gene tagged with FLAG sequence at the C terminus for complementation | This study |
| pMGKU420 | pDONR201, *bcr4* gene lacking amino acids region 169-173 tagged with FLAG sequence at the C terminus | This study |
| pMGKU421 | pDONR201, *bcr4* gene lacking amino acids region 164-173 tagged with FLAG sequence at the C terminus | This study |
| pMGKU422 | pDONR201, *bcr4* gene lacking amino acids region 159-173 tagged with FLAG sequence at the C terminus | This study |
| pMGKU423 | pDONR201, *bscI* gene for complementation | This study |
| pRK415-R4-R3-F | pRK415, recombination sites for MultiSite Gateway, Tet[r] | 8 |
| pDONR-*fhaP* | pDONR-P4-P1R, *fha* promoter | 8 |
| pDONR-*rrnB* | pDONR-P2R-P3, *rrnB* terminator | 8 |
| pMGKU424 | pRK415-R4-R3-F, *bcr4* gene tagged with FLAG sequence at the C terminus for complementation | This study |
| pMGKU425 | pRK415-R4-R3-F, *bcr4* gene lacking amino acids region 169-173 tagged with FLAG sequence at the C terminus | This study |
| pMGKU426 | pRK415-R4-R3-F, *bcr4* gene lacking amino acids region 164-173 tagged with FLAG sequence at the C terminus | This study |
| pMGKU427 | pRK415-R4-R3-F, *bcr4* gene lacking amino acids region 159-173 tagged with FLAG sequence at the C terminus | This study |
| pMGKU428 | pRK415-R4-R3-F, *bscI* gene | This study |

| Table S3. Primers used in the study | |
|---|---|
| Name | Sequence (5'→3') |
| B1-bscl-V5 | AAAAAGCAGGCTTGTTGGACAGCGGACCCGCC |
| B2-bscl-V5 | AGAAAGCTGGGTTTGACATTCTCGCCAGCGTGTC |
| B1-bscK-V5 | AAAAAGCAGGCTTGGCGCACGCGCTGGTCCCG |
| B2-bscK-V5 | AGAAAGCTGGGTTGGCTTGGGCGGGGAACGAGG |
| B1-BscJ-V5 | AAAAAGCAGGCTTGGAGTACGAGTTGGTGGGC |
| B2-BscJ-V5 | AGAAAGCTGGGTTGTCATGCCCGGCTCCTTCCG |
| B1-bcr4-comp | AAAAAGCAGGCTGCCAGGTCCGGTCTCGCACCG |
| B2-bcr4-V5 | AGAAAGCTGGGTTTCCAGGAGCTCCAGGTAATG |
| 5-HindIII-bcr4 | ATCCGAATTCAAGCTTCATTCAGACTCAGGTTCAGATTC |
| 3-bcr4-Strep | CGGGTGGCTCCATCCAGGAGCTCCAGGTAATG |
| 5-HindIII-bscl | ATCCGAATTCAAGCTTAATTTGGATCTGACGGCGATC |
| 3-bscl-Strep | CGGGTGGCTCCATGACATTCTCGCCAGCGTGTC |
| 3-Strep-HindIII | GCAGGTCGACAAGCTTTCATTTTTCGAACTGCGGGTGGCTCCA |
| 5-bcr4-58-109-Strep-IF | GCCGTCAGCGATCTGTCGGACTGGACGCCTGGTG |
| 3-IF-bcr4-58-109-Strep | CAGATCGCTGACGGCCGCCG |
| 5-bcr4-110-173-Strep-IF | TGGAGCCACCCGCAGTTCGA |
| 3-IF-bcr4-110-173-Strep | CTGCGGGTGGCTCCAGCGGTCCTCCGCCTGTGCCG |
| B1-bscl | AAAAAGCAGGCTTCATGACTGTTCACGACGAC |
| B2-bscl | AGAAAGCTGGGTGCACAAGATCCAGCGCGACC |
| R1-bscl | CGACGTTGGCGTTGATCGCCGTCAGATC |
| R2-bscl | TCAACGCCAACGTCGACACGCTGGCGAG |
| 5-bcr4-FLAG-IF | CCACCCGCAGTTCGAAAAATGAATTTGGATCTACCCAGCTTTCTTG |
| 3-IF-bcr4-FLAG | TCGAACTGCGGGTGGCTCCATCCAGGACCTCCAGGTAATGGCTCG |
| 3-IF-bcr4△169-173-FLAG | GTCATCCTTGTAGTCTAATGGCTCGGACTGCAACG |
| 3-IF-bcr4△164-173-FLAG | GTCATCCTTGTAGTCCAACGGCGCGGGCCGCATCATG |
| 3-IF-bcr4△159-173-FLAG | GTCATCCTTGTAGTCCATCATGGCGGGTCCGCTGTC |
| 3-B2-bscl-comp | AGAAAGCTGGGTTTGACATTCTCGCCAGCGTGTC |
| 5-bscl | CGATCTGCAGGCCAGGTT |
| 3-bscl | CTATTGCCTTGCCCACCAA |
| 5-bopD | CGGCTCGGTGAAGACATC |
| 3-bopD | CCTCCCGCATCTGTTGAC |
| 5-recA | ATGAAGATCGGCGTGATGT |
| 3-recA | TAGAACTTGAGCGCGTTGC |

**Fig. S1. Construction of the T3SS machinery in strains lacking Bcr4 and/or BspR.** In the *B. bronchiseptica* wild-type (upper left), the BspR negative regulation level for the *bsc* locus transcription is moderate, and the T3SS machinery is established. In the Bcr4-deficient strain (upper right), BspR strongly represses the *bsc* locus transcription, and construction of the T3SS machinery is incomplete. In the BspR-deficient strain (lower left), the negative regulatory effect of BspR is cancelled, and the construction of the T3SS machinery is promoted. In the BspR/Bcr4 double-deficient strain (lower right), while the *bsc* locus transcription is promoted because of BspR deficiency, T3SS is not functional.

Fig. S1

```
Bb     1 MHSDSGSDSGSDSGSGS--PMASSIHPSEPIQPMEHVLEEADARLLTEVGFLAAAVSDLT    58
Bp     1 MHSDSGSDSGSDSGSGS--PMVSSIHPSEPIQPMEHVLEEADARLLTEVGFLAAAVSDLT    58
Bpp    1 MHSDSGSDSGSGSGSGSGSPMASSIHPSEPIQPMEHVLEEADARLLTEVGFLAAAVSDLT    60

Bb    59 RADAIFNALQRVRPGRTYPCIGLAVARMNAGLPDEAAEILANFQPAQAEDRSELDAWCGF   118
Bp    59 RADAIFNALQRVRPGRTHPCIGLAVARMNAGLPDEAAEILANFQPAQPEDRSELDAWCGF   118
Bpp   61 RADAIFNALQRVRPGRTYPCIGLAVARMNAGLPDEAAEVLANFQPAQAEDRSELDAWCGF   120

Bb   119 ALLLAGRSDEARRMLQRAIDAGGEAARLAQVVLDSGPAMMRPAPLQSEPLPGAPG        173
Bp   119 ALLLAGRSDEARRMLQRAIDAGGEAARLAQVVLDSGPAMMRPAPLQSEPLPGAPG        173
Bpp  121 ALLLAGRSDEARRMLQRAIDAGGEAARLAQVVLDSGPAMMRPAPLQSEPLPGAPG        175
```

**Fig. S2. Alignment of Bcr4 amino acid sequences in representative *Bordetella* species.** Bcr4 amino acid sequences of *B. bronchiseptica* S798 (Bb), *B. pertussis* Tohama I (Bp) and *B. parapertussis* 12822 (Bp) were compared using ClustalW. The grey highlighted letters represent amino acid residues that were different from those in Bb. Bcr4 of Bp and Bpp have 98.3% and 97.1% identities with those of Bp, respectively.

**Fig. S3. The nonspecific reaction of anti-BscI antibody to Bsp22.** The culture supernatants (CS) were prepared from the wild-type strain, Δ*bsp22* (Bsp22-deficient strain), Δ*bspR* (BspR-deficient strain) or Δ*bscI+bscI* (BscI-complemented strain) cultured in SS medium. The CS samples were separated by SDS-PAGE and stained with Coomassie Brilliant Blue (CBB, left panel) or analyzed by Western blotting (WB) with anti-BscI antibody (right panel). NS indicates nonspecific signals. Experiments were performed at least three times, and representative data are shown.

**Fig. S4. The results of the RT-PCR analysis for the mRNA level of *bscI* in *B. bronchiseptica* strains.** Total RNA was prepared from the wild-type strain, Δ*bspR* (BspR-deficient strain) or Δ*bspR*Δ*bcr4* (BspR- and Bcr4-deficient strain) cultured in SS medium and subjected to a quantitative RT-PCR analysis. The histogram shows the relative amount of *bscI* mRNA normalized by the housekeeping gene, *recA* mRNA. Experiments were performed at least three times, and representative data are shown.

**Fig. S5. The results of the RT-PCR analysis for mRNA levels of *bopD* and *bscI* in the wild-type *B. bronchiseptica*.** Total RNA was prepared from the wild-type strain cultured in SS medium and subjected to a quantitative RT-PCR analysis. The histogram shows the relative amount of *bopD* and *bscI* mRNA in the wild-type. The relative ratio of *bscI* mRNA is shown when the *bopD* mRNA amount is set as 1. Experiments were performed at least three times, and representative data are shown.

**Fig. S6. The time course of BscI production in _B. bronchiseptica_.** The whole cell lysates (WCL) were prepared from the wild-type strain, Δ*bcr4* (Bcr4-deficient strain), Δ*bspR* (BspR-deficient strain) or Δ*bspR*Δ*bcr4* (BspR- and Bcr4-deficient strain) cultured in SS medium for 0, 2, 5, 8 or 18 hr. The WCL were separated by SDS-PAGE and analyzed by Western blotting with antibodies against BscI, BopB and RpoB. Experiments were performed at least three times, and representative data are shown.

Fig. S6

**Fig. S7. The predicted structural model of *B. bronchiseptica* Bcr4**

(A) The AlphaFold2 (AF2)-predicted structural model of *B. bronchiseptica* Bcr4 and structural comparison with *Aeromonas hydrophila* AcrH (PDB: 3WXX) and *Pseudomonas aeruginosa* PscG (PDB: 2UWJ). Z-score, root mean square deviation (RMSD), and amino acid identity (AA %ID) of AcrH and PscG compared with Bcr4 are shown, respectively. (B) The pairwise sequence alignment of Bcr4, AcrH, and PscG. The most frequent amino acid type is colored. The secondary structure assignments (H/h: helix, E/e: strand, L/l: coil) are also shown.

Fig. S7