Bcr4 is a Chaperone for the Inner Rod Protein in the *Bordetella* Type III Secretion

System

Masataka Goto[a], Akio Abe[a], Tomoko Hanawa[b], and Asaomi Kuwae[a#]

[a]Laboratory of Bacterial Infection, Graduate School of Infection Control Sciences, Kitasato

University, Tokyo, Japan

[b]Department of Infectious Diseases, Kyorin University School of Medicine, Tokyo, Japan

Running Head: Bcr4 is a Chaperone for Inner Rod in *Bordetella* T3SS

[#]**Address Correspondence to:** Asaomi Kuwae, kuwae@lisci.kitasato-u.ac.jp

Abstract

Bordetella bronchiseptica injects virulence proteins called effectors into host cells via a

type III secretion system (T3SS) conserved among many Gram-negative bacteria. Small

proteins called chaperones are required for stabilizing some T3SS components or

localizing them to the T3SS machinery. In a previous study, we identified a chaperone-like

protein named Bcr4 that regulates T3SS activity in *B. bronchiseptica*. Bcr4 does not show

strong sequence similarity to well-studied T3SS proteins of other bacteria, and its function

remains to be elucidated. Here, we investigated the mechanism by which Bcr4 controls

T3SS activity. A pull-down assay revealed that Bcr4 interacts with BscI, based on its

homology to other bacterial proteins, to be an inner rod protein of the T3SS machinery. A

pull-down assay using truncated Bcr4 derivatives and secretion profiles of *B.*

bronchiseptica producing truncated Bcr4 derivatives showed that the Bcr4 C-terminal

region is necessary to interact with BscI and to activate the T3SS. Moreover, the deletion of

BscI abolished the secretion of type III secreted proteins from *B. bronchiseptica* and the

translocation of a cytotoxic effector into cultured mammalian cells. Finally, we showed that

BscI is unstable in the absence of Bcr4. These results suggest that Bcr4 supports the

construction of the T3SS machinery by stabilizing BscI. This is the first demonstration of a

chaperone for the T3SS inner rod protein among the virulence bacteria possessing the

T3SS.

Importance

The type III secretion system (T3SS) is a needle-like complex that projects outward from bacterial cells. *Bordetella bronchiseptica* uses the T3SS to translocate virulence proteins into host cells. Our previous study reported that a protein named Bcr4 is essential for the secretion of virulence proteins from *B. bronchiseptica* bacterial cells through the T3SS. Because other bacteria lack a Bcr4 homologue, the function of Bcr4 has not been elucidated. In this study, we discovered that Bcr4 interacts with BscI, a component of the T3SS machinery. We showed that a *B. bronchiseptica* BscI-deficient strain was unable to secrete type III secreted proteins. Furthermore, the deletion of Bcr4 led to BscI instability in *B. bronchiseptica*. These results suggest that Bcr4 stabilizes BscI to allow construction of the T3SS in *B. bronchiseptica*.

Introduction

The genus *Bordetella* consists of Gram-negative bacteria that infect the respiratory tracts of humans and animals. *Bordetella pertussis* causes a severe coughing attack called whooping cough in humans (1, 2). *Bordetella bronchiseptica* causes atrophic rhinitis in pigs and kennel cough in dogs (1, 2). These *Bordetella* spp. harbor a virulence factor secretion apparatus called the type III secretion system (T3SS).

The T3SS consists of a basal body that penetrates the inner and outer membranes of the bacteria and a needle structure that protrudes outside the bacteria, and is conserved in many Gram-negative bacteria such as *Yersinia*, *Salmonella*, and *Pseudomonas*. *B. bronchiseptica* uses the T3SS to inject virulence factors, called effectors, into host cells to disrupt the physiological functions of host cells (3). Once the basal body and the export apparatus of the T3SS are completed, the type III secreted proteins are secreted in a fixed order. First, the components of the needle structure and the inner rod that ties the needle structure to the basal body are secreted. After the needle is completed, translocators are secreted, which form small pores in the host cell membrane to create a pathway for effectors. Finally, the effectors translocate into the host cell via the T3SS (4, 5). BopB (6), BopD (7), and Bsp22 (8) function as translocators, while BteA (also referred to as BopC) (9, 10), BopN (11, 12), and BspR (also referred to as BtrA) (13, 14) function as effectors in *B. bronchiseptica*. Bsp22 is located at the tip of the needle and bridges the needle to the pore-forming factors embedded in the plasma membrane (15). BteA has been shown to

cause membrane-disrupting cytotoxicity in mammalian cells (10). BspR is a regulator that

represses transcription of the *bteA* gene and genes on the *bsc* locus, where genes

encoding type III secreted proteins and components of T3SS are located (14, 16). The

BscF and BscI proteins of *B. bronchiseptica* have 18.4% and 18.3% identity with the

Yersinia needle YscF and inner rod YscI, respectively, suggesting that they are the needle

and inner rod of the *Bordetella* T3SS.

In addition, many type III secreted proteins have unique chaperones that are involved in

stabilizing the substrate and preventing premature polymerization of the substrate in the

bacterial cytosol, and then in efficiently transporting the substrate to the T3SS machinery

(17-20). For example, PscE and PscG function as chaperones of the needle PscF in

Pseudomonas. These chaperones stabilize PscF in the bacterial cytosol and are involved

in PscF secretion through the T3SS (21). In bacteria such as *Yersinia* and *Pseudomonas*,

an inner rod chaperone is thought to exist, because the inner rod has the properties of

being secreted out of the bacterial cell and polymerizing (22, 23). However, a chaperone for

the inner rod has not been reported.

So far, a chaperone-like protein called Bcr4 has been identified in *B. bronchiseptic*a (24).

In this study, we attempted to identify factors that interact with Bcr4 in order to investigate

how Bcr4 is involved in the T3SS regulation.

Results

Bcr4 binds to BscI, an inner rod protein of the *Bordetella* type III secretion system.

The results of a previous study suggested that Bcr4 is a chaperone for components of

the type III secretion system (T3SS) (24). T3SS chaperones are known to be involved in

substrate stability and efficient transport of substrates to the T3SS machinery (17, 19, 20).

In addition, it is generally known that the genes of these chaperones are localized adjacent

to genes encoding their substrates (25). On the *B. bronchiseptica* S798 chromosome, the

genes encoding BcrH2, BscI, BscJ, and BscK are located in the vicinity of the *bcr4* gene

(Fig. 1A), and these proteins are predicted to function as a translocator chaperone, inner

rod, inner membrane ring, and ATPase cofactor, respectively (5, 26). Based on the above,

we hypothesized that Bcr4 binds to either BscI, BscJ, or BscK. To test whether Bcr4 binds

to BscI or BscK, we added *E. coli* lysates containing the V5-tagged target proteins (BscI-V5

or BscK-V5) to Strep-Tactin beads with the purified Strep-tagged Bcr4 (Bcr4-Strep), and

then performed the pull-down assay. Supernatant fraction (Sup) and pellet fraction (Pellet)

samples were prepared, separated by SDS-PAGE, and subjected to Western blotting with

antibodies against V5 (Fig. 1B). The results showed that the V5 signal was detected in the

Pellet sample of the BscI-V5, but not in that of BscK-V5 (Fig. 1B). These results suggest

that Bcr4 binds to BscI, an inner rod protein of the *Bordetella* T3SS.

Next, to reconfirm that Bcr4 binds to BscI, a pull-down assay was performed by adding *E.*

coli lysate containing V5-tagged Bcr4 (Bcr4-V5) to Strep-Tactin beads with the purified

Strep-tagged BscI (BscI-Strep) or BteA N-terminal 1–312 amino acids region

(BteA-N-Strep). As a result, the V5 signal was detected in the Pellet sample of the

BscI-Strep, but was hardly detected in that of the BteA-N-Strep (Fig. 1C). These results

strongly suggested that Bcr4 binds to BscI.

The C-terminal region of Bcr4 is required for the binding of Bcr4 to BscI and T3SS

activity.

Next, to investigate the Bcr4 region responsible for the binding to BscI, we produced

Strep-tagged full-length Bcr4 (Bcr4-FL-Strep), Bcr4 lacking amino acids region 58–109

(Bcr4Δ58-109-Strep), and Bcr4 lacking amino acids region 110–173 (Bcr4Δ110-173-Strep)

in *E. coli* (Fig. 2A). A pull-down assay was performed by adding *E. coli* lysate containing

V5-tagged BscI (BscI-V5) to Strep-Tactin beads with the purified Bcr4-FL-Strep,

Bcr4Δ58-109-Strep, or Bcr4Δ110-173-Strep. The Sup and Pellet samples were then

prepared, separated by SDS-PAGE, and subjected to Western blotting using antibodies

against V5 (Fig. 2B). As a result, the V5 signal was detected in the Pellet sample of

Bcr4-FL-Strep, but not in those of Bcr4Δ58-109-Strep and Bcr4Δ110-173-Strep. These

results suggest that the BscI-binding domain is located in the 58–173 amino acids region of

Bcr4.

To determine the region of Bcr4 required for the function of T3SS, we produced

full-length FLAG-tagged Bcr4 (Bcr4-FL-FLAG), Bcr4 with the 5 C-terminal amino acids

deleted (Bcr4Δ169-173-FLAG), Bcr4 with the 10 C-terminal amino acids deleted

(Bcr4Δ164-173-FLAG), and Bcr4 with the 15 C-terminal amino acids deleted

(Bcr4Δ159-173-FLAG) (Fig. 2A) within a Bcr4-deficient strain (ΔBcr4)

(ΔBcr4+Bcr4-FL-FLAG, ΔBcr4+Bcr4Δ169-173-FLAG, ΔBcr4+Bcr4Δ164-173-FLAG and

ΔBcr4+Bcr4Δ159-173-FLAG, respectively). Whole cell lysates (WCL) and Sup samples

were prepared from these strains and separated by SDS-PAGE. Western blottings were

then carried out using antibodies against Bcr4, FLAG, BteA (an effector and a type III

secreted protein) or BopD (a translocator and a type III secreted protein) (Fig. 2C). Since

the antibody against Bcr4 was generated using its 18 C-terminal amino acids (amino acids

(aa) 156–173) as the antigen peptide, it may not recognize the Bcr4 partial deletion mutant

proteins used in this experiment. Therefore, we also performed Western blotting using

antibodies against FLAG (Fig. 2C). As a result, when anti-Bcr4 antibodies were used

against the WCL samples, signals were detected in the wild-type, ΔBcr4+Bcr4-FL-FLAG

and ΔBcr4+Bcr4Δ169-173-FLAG, but almost no signals were detected in

ΔBcr4+Bcr4Δ164-173-FLAG and ΔBcr4+Bcr4Δ159-173-FLAG (Fig. 2C). On the other hand,

when anti-FLAG antibody was used, signals were detected in ΔBcr4+Bcr4Δ164-173-FLAG

and ΔBcr4+Bcr4Δ159-173-FLAG (Fig. 2C). These results confirmed that the Bcr4 partial

deletion mutant proteins were produced in the ΔBcr4 strain. When anti-BteA or anti-BopD

antibodies were used against the Sup samples, signals were detected in the wild-type,

ΔBcr4+Bcr4-FL-FLAG, ΔBcr4+Bcr4Δ169-173-FLAG and ΔBcr4+Bcr4Δ164-173-FLAG,

respectively, but not in ΔBcr4 or Bcr4Δ159-173-FLAG (Fig. 2C). These results suggest that

the region required for the T3SS function is located in the 159–163 amino acids region of

Bcr4.

BscI is required for the function of the T3SS in *B. bronchiseptica*.

BscI is a homologue of YscI, an inner rod of the *Yersinia* T3SS, and YscI is essential for

the function of the T3SS (Wood, JB, 2008). To confirm that BscI is required for the function

of the *B. bronchiseptica* T3SS, the *B. bronchiseptica* S798 wild-type, BscI-deficient strain

(ΔBscI), BscI-complemented strain (ΔBscI+BscI), and BscN-deficient strain (ΔBscN) were

cultured in SS medium. WCL and Sup samples were prepared, and these samples were

separated by SDS-PAGE, followed by Western blotting using antibodies against BscI, BteA,

BopD or FHA (filamentous hemagglutinin, an adhesion factor) (Fig. 3). As a result, no

signals of BscI were detected in the WCL samples of all strains used here (Fig. 3). The BscI

signal was detected in the Sup of ΔBscI+BscI, and a faint BscI signal was detected in the

Sup of the wild-type (Fig. 3). The BteA and BopD signals were detected in the Sup of the

wild-type and ΔBscI+BscI, but not in those of ΔBscI and ΔBscN (Fig. 3). As expected, the

FHA signals were detected in the WCL and Sup samples of all strains used here (Fig. 3).

These results suggest that in *B. bronchiseptica*, BscI is required for the function of the

T3SS and is secreted out of the bacterial cell.

BscI plays an important role in the function of the *B. bronchiseptica* T3SS during infection of cultured mammalian cells.

To further investigate whether BscI is required for the construction and function of the T3SS in *B. bronchiseptica*, we infected L2 cells (a rat lung epithelial cell line) with the wild-type, ΔBscI, ΔBscI+BscI or ΔBscN and measured the number of Bsp22 (a translocater, and a component of the filament-like structure; Fig. 6) signals detected on L2 cells by immunofluorescence microscopy. For the efficient detection of Bsp22 signals, we cultured the bacteria in iron-depleted SS medium, which has been shown to increase the amount of type III secreted protein secreted by *B. bronchiseptica* (27). L2 cells were infected with bacteria for 1 h at a multiplicity of infection (MOI) of 2000. Then, Bsp22 and F-actin were stained with antibodies against Bsp22 and Rhodamine-Phalloidin, respectively (Fig. 4A). As a result, the fluorescence signals of Bsp22 were detected on the L2 cells infected with the wild-type or ΔBscI+BscI, but not on the L2 cells infected with ΔBscI or ΔBscN (Fig. 4A, B). We also infected L2 cells with the wild-type, ΔBscI, ΔBscI+BscI or ΔBscN for 1 h at an MOI of 200 and measured the amount of lactate dehydrogenase (LDH) released into the extracellular medium. The results showed that LDH was detected in the medium of L2 cells infected with the wild-type or ΔBscI+BscI, but not in the medium of L2 cells infected with ΔBscI or ΔBscN (Fig. 4C). These results suggest that BscI is required for the construction of the T3SS and induction of T3SS-dependent death of mammalian cells in *B. bronchiseptica*.

Bcr4 is required for the stability of BscI in *B. bronchiseptica*.

As mentioned above, it was suggested that the interaction between Bcr4 and BscI is necessary for the function of the T3SS (Fig. 2). Therefore, to investigate whether Bcr4 is involved in the stability of BscI, we cultured the *B. bronchiseptica* wild-type, Bcr4-deficient strain (ΔBcr4, T3SS-inactive strain) (24), BspR-deficient strain (ΔBspR, T3SS-overproducing and -oversecreting strain) (14) or BspR- and Bcr4-deficient strain (ΔBspRΔBcr4, a T3SS-overproducing but -inactive strain) (24) in SS medium. The prepared WCL samples were separated by SDS-PAGE, and Western blotting was performed using antibodies against BscI, BscJ (a protein composing the inner membrane ring of the T3SS), and Bcr4. As a result, the BscJ signals were detected in the WCL samples of ΔBspR and ΔBspRΔBcr4 (Fig. 5). On the other hand, the BscI signal was detected in ΔBspR but not in ΔBspRΔBcr4 (Fig. 5). These results suggest that Bcr4 is necessary for the stability of BscI in *B. bronchiseptica*.

Discussion

In the present study, we found that deletion of BscI in *B. bronchiseptica* abolished the

extracellular secretion of type III secreted proteins (Fig. 3, 4). In addition, when Bcr4 was

deleted in the *B. bronchiseptica* BspR-deficient strain, BscI (a protein composing the inner

rod of the *Bordetella* T3SS) was not detected in the bacterial cells (Fig. 5). These results

suggest that Bcr4 contributes to the establishment of the T3SS in *B. bronchiseptica* by

stabilizing BscI (Fig. 6).

To date, chaperones for needles, translocators and effectors have been reported in

bacteria that retain the T3SS, such as *Yersinia* (28-31). However, a chaperone for the inner

rod protein has not been reported, and the stability of the inner rod in the bacterial cell and

the transport of inner rods to the T3SS remain unclear.

In the present study, it was suggested that the 58–173 amino acids region of Bcr4 is

required for the binding of Bcr4 to BscI (Fig. 2B). To investigate whether the 1–57 amino

acids region of Bcr4 is also required for the binding of Bcr4 to BscI, we attempted to purify

Bcr4 lacking the N-terminal 1–57 amino acids region (Bcr4Δ1-57-Strep). However, the

amount of Bcr4Δ1-57-Strep produced in *E. coli* was extremely low (data not shown), so a

pull-down assay could not be performed. Figure 2C suggests that the 159–163 amino acids

region of Bcr4 is required for the function of the T3SS. These results suggest that the

binding of Bcr4 to BscI is necessary for the T3SS function, because the region of Bcr4

required for the T3SS function (aa 159–163) is contained in the region where Bcr4 binds to

BscI (aa 58–173) (Fig. 2).

Since the *bscI* and *bopD* genes are located in the *bsc* locus (Fig. 1A) and transcribed

from the same promoter, the mRNA levels of the *bscI* and *bopD* genes in the bacterial cells

are likely to be the same. In this study, the BopD signal was detected in the WCL of the *B.*

bronchiseptica wild-type strain, but the BscI signal was not detected (Fig. 3). To test the

anti-BscI antibody sensitivity, purified recombinant BscI and the Sup sample containing

BopD were subjected to SDS-PAGE. The amounts of BscI and BopD in the samples were

adjusted to the same weight when we loaded the samples on the gel. A Western blotting

analysis was then performed using antibodies against BscI or BopD. The results showed

that the BscI and BopD signals were detected with almost the same intensity, indicating that

the sensitivity of the anti-BscI antibody for antigen detection was not low. Therefore, the

amount of BscI present in *B. bronchiseptica* was considered to be lower than that of BopD.

The reason for this is unknown, but it could be due to the low efficiency of translation of the

bscI gene.

BspR is a negative regulator that represses the transcription of genes on the *bsc* locus

(14). In strains that are unable to secrete type III secreted proteins, such as Bcr4-deficient

strains (ΔBcr4), the repression of gene transcriptions on the *bsc* locus by BspR is enhanced

(24), and thus the BscJ signal in the WCL sample of ΔBcr4 was weaker than that of the

wild-type (Fig. 5). In this study, we attempted to detect the BscI signal using a

BspR-deficient strain (ΔBspR), in which the gene transcriptions on the *bsc* locus are

promoted (Fig. 5), because the BscI signal was not detected in the WCL sample of the *B.*

bronchiseptica wild-type (Fig. 3). The *bscJ* gene is located downstream of the *bscI* gene

(Fig. 1A), and the BscJ protein is predicted to function as an inner membrane ring, a

component of the T3SS. The signal intensity of BscJ was not decreased in the WCL sample

of the BspR/Bcr4 double-deficient strain (ΔBspRΔBcr4) compared to that of ΔBspR (Fig. 5),

suggesting that Bcr4 is not involved in the stability of BscJ. Therefore, it is suggested that

Bcr4 specifically stabilizes BscI. In addition, when BscI was overexpressed in ΔBcr4, a

small amount of extracellular secretion of type III secreted proteins was observed (data not

shown), suggesting that Bcr4 is not a component of the T3SS but a chaperone of BscI.

Previous studies have shown that overexpression of Bcr4 promotes the production and

secretion of type III secreted proteins in *B. bronchiseptica* (24). It is also known that Bcr4 is

not secreted out of the *B. bronchiseptica* bacterial cell (24). *Pseudomonas aeruginosa* Pcr4

is a homologue of Bcr4 (identity: 26.6%). Since there has been no report on overexpression

of Pcr4 in *Pseudomonas*, it is unclear whether Pcr4 overexpression enhances the

production and secretion of type III secreted proteins (32). In addition, since Pcr4 has been

reported to be an extracellularly secreted protein in *Pseudomonas*, it is quite possible that

Bcr4 functions differently from Pcr4. The hypothetical protein (NH44784_014631) of

Achromobacter xylosoxidans, which like *Bordetella*, belongs to the family Alcaligenaceae

has 49.1% identity with Bcr4. This identity is higher than the identity between Pcr4 and Bcr4.

Therefore, it is possible that NH44784_014631 and Bcr4 have the same function. However,

there has been no report describing how NH44784_014631 functions in *Achromobacter*.

Further functional analysis of Bcr4 homologues in other bacteria will provide more

information on the function of Bcr4.

Homologues of BscI include YscI of *Yersinia pseudotuberculosis*, but the identity

between BscI and YscI is low at 18.3%. In addition, no gene encoding a homologue of Bcr4

has been found in the genome of *Yersinia*. Therefore, in bacteria such as *Yersinia* and other

Enterobacteriaceae, for which there are numerous reports on the T3SS, the inner rod may

be stabilized by a system different from that in *Bordetella*. It is also unknown why inner rod

production is low in *B. bronchiseptica* (Fig. 3). Furthermore, it is still unclear how Bcr4

antagonizes the BspR negative regulation activity, and how overexpression of Bcr4

promotes the production and secretion of type III secreted proteins in *B. bronchiseptica*.

Further analysis is needed to elucidate the detailed molecular mechanism by which Bcr4

contributes to the establishment of the T3SS.

Materials and Methods

Bacterial strains and cell culture.

The strains used in this study are listed in Table 1. *B. bronchiceptica* S798 (6) was used as the wild-type strain. ΔBscN (a BscN-deficient strain), ΔBcr4 (a Bcr4-deficient strain), ΔBspR (a BspR-deficient strain) and ΔBspRΔBcr4 (a BspR and Bcr4-deficient strain) have been described elsewhere (6, 14, 24). *Escherichia coli* DH10B, BL21 (33) and Sm10λ*pir* were used for DNA cloning, recombinant protein expression and conjugation, respectively (Table 1). *B. bronchiceptica* was grown on Bordet-Gengou agar plates at $37^{\circ}C$ for 2 days. Fresh colonies of *B. bronchiceptica* were cultured in Stainer-Scholt (SS) medium (34) with a starting A_{600} of 0.23 under vigorous shaking conditions at $37^{\circ}C$. L2 cells, a rat lung epithelial cell line (ATCC CCL-149), were maintained in F-12K (Invitrogen). The cell culture medium was supplemented with 10% fetal calf serum (FCS). L2 cells were grown at $37^{\circ}C$ under a 5% CO_2 atmosphere.

Pull-down assay.

The primers used in this study are listed in Table 2. In order to express *bscI*, *bscK* or *bcr4* tagged with a V5 sequence at the respective C-terminus (*bscI*-V5, *bscK*-V5 or *bcr4*-V5), we amplified DNA fragments encoding the *bscI*, *bscK* or *bcr4* genes with the primer sets of B1-bscI-V5 and B2-bscI-V5, B1-bscK-V5 and B2-bscK-V5 or B1-bcr4-comp and B2-bcr4-V5, respectively, using *B. bronchiseptica* S798 genomic DNA as the template.

Each resulting PCR product was cloned into pDONR201 to obtain pMGKU404, pMGKU405

or pMGKU406, respectively, by means of adapter PCR and site-specific recombination

techniques using the Gateway cloning system (Invitrogen). Each plasmid was mixed with

an expression vector such as p99*ccdB*-V5 (35) to obtain pMGKU407, pMGKU408 or

pMGKU409 using the Gateway cloning system.

BL21 cells carrying pMGKU407, pMGKU408 or pMGKU409 were cultured overnight at

37°C with shaking, and then diluted 1:40 in LB liquid medium containing 50 μL/mL ampicillin

and incubated for 2 h at 37°C with shaking. Each bacterial culture was further incubated for

5 h at 30°C in the presence of isopropyl-beta-thiogalactopyranoside (IPTG) at the final

concentration of 1 mM. Bacteria were collected by centrifugation at 2,600×*g* for 15 min, and

suspended in cold TBS containing protease inhibitor cocktail, cOmplete (Roche). Each

bacterial suspension was sonicated, and each supernatant was used for the pull-down

assay.

Next, in order to purify Bcr4 or BscI tagged with six histidine residues (6×His) at the

respective N-terminus and Strep at the respective C-terminus, we amplified DNA fragments

encoding *bcr4* or *bscI* with the primer sets of 5-HindIII-bcr4 and 3-bcr4-Strep, or

5-HindIII-bscI and 3-bscI-Strep using *B. bronchiseptica* S798 genomic DNA as the template.

Each amplified DNA fragment was used as a template for 2nd PCR with a primer set

consisting of the upper primer used in the 1st PCR and 3-Strep-HindIII to add a 24 bp

sequence encoding the Strep tag. Each resulting PCR product was cloned into the HindIII

recognition sites of pColdII to obtain pMGKU410 or pMGKU411 by the In-Fusion Cloning

System (Clontech), respectively. In order to purify Bcr4 lacking the amino acid regions 58–

109 or 110–173, we amplified the DNA fragment with the primer sets of

5-bcr4-58-109-Strep-IF and 3-IF-bcr4-58-109-Strep, or 5-bcr4-110-173-Strep-IF and

3-IF-bcr4-110-173-Strep using pColdII-*bcr4*-Strep as the template. Each amplified fragment

was self-ligated by the In-Fusion Cloning System, respectively, and then designated

pMGKU412 or pMGKU413.

BL21 cells carrying pMGKU410, pMGKU411, pColdII-*bteA*-N-Strep (36), pMGKU412 or

pMGKU413 were cultured overnight at 37ºC with shaking, and then diluted 1:100 in LB

liquid medium containing 50 μL/mL ampicillin and incubated for 2 h at 37ºC with shaking,

respectively. Each bacterial culture was further incubated overnight at 15ºC in the presence

of IPTG at the final concentration of 0.05 mM. Bacteria were collected by centrifugation at

2,600×*g* for 20 min, and suspended in cold TBS containing cOmplete. Each bacterial

suspension was sonicated and each supernatant except for that of pMGKU411 was

subjected to purification using Ni-NTA agarose (Qiagen) according to the manufacturer's

instructions. The purified proteins were dialyzed with TBS. As for pMGKU411, the bacterial

suspension was sonicated and the pellet was suspended in Inclusion Body Solubilization

Reagent (Thermo) according to the manufacturer's instructions. The BscI-Strep in

solubilized solution was refolded as described previously (37). Briefly, the solubilized

solution was diluted with 5-fold larger refolding buffer (20 mM Tris-HCl, pH-8, 150 mM NaCl,

10% glycerol) overnight at 4°C on a rotator, and then centrifugated at 20,000×g for 15 min.

The supernatant was dialyzed with dialysis buffer (20 mM Tris-HCl, pH-8, 150 mM NaCl,

10% glycerol), and then centrifugated at 20,000×g for 15 min. The supernatant was used

as purified protein for the pull-down assay.

We mixed the dual-tagged protein (6×His and Strep) and 30 μl Strep-Tactin resin (IBA) in

an Eppendorf tube and rotated the tube at 4°C for 1 h. Then, we washed the beads with

TBS three times. Next, the V5-tagged protein-containing *E. coli* lysate was added to the

tube and rotated at 4°C for 3 h. We transferred 30 μL supernatant to new Eppendorf tube

and added 30 μL 2×SDS-PAGE sample buffer to prepare the Sup. Then we washed the

beads with TBS (Fig. 1B) or TBS containing 0.1% Triton X-100 (Fig. 1C) three times and

added 30 μL 2×SDS-PAGE sample buffer to prepare the Pellet samples.

Construction of a *bscI* gene-disrupted strain.

To construct the *bscI* mutant, a 2.4 kb DNA fragment encoding *bscI* and its flanking

regions was amplified by PCR with primers B1-bscI and B2-bscI using *B. bronchiceptica*

S798 genomic DNA as the template. The resulting PCR product was cloned into

pDONR201 to obtain pMGKU414 using the Gateway cloning system. An inverse PCR was

carried out with the primers R1-bscI and R2-bscI using circular pMGKU414 as the template.

The resulting PCR product was self-ligated using the In-Fusion Cloning System to obtain

pMGKU415. This plasmid contained a 369-bp in-frame deletion from 30 bp downstream of

the 5' end of the *bscI* gene to 30 bp upstream of the 3' end of the gene. This plasmid,

pMGKU415, was mixed with a positive suicide vector such as pABB-CRS2 (38) to obtain

pMGKU416 using the Gateway cloning system. The pMGKU416 plasmid in turn was

introduced into *E. coli* Sm10λ*pir*, and transconjugated into the S798 wild-type as described

previously (39). The resulting mutant strain was designated ΔBscI.

Construction of plasmids used for producing Bcr4 derivatives and BscI

complementation

In order to produce full-length Bcr4 tagged with a FLAG sequence at the C-terminus, we

performed an inverse PCR with primers of 5-bcr4-FLAG-IF and 3-IF-bcr4-FLAG using

pDONR-*bcr4* (24) as the template. The resulting PCR product was self-ligated to obtain

pMGKU417 by the In-Fusion Cloning System. To produce Bcr4 lacking the amino acid

regions 159–173, 164–173 or 169–173, inverse PCR was carried out with the primer sets of

5-*bcr4*-FLAG-IF   and   3-IF-bcr4Δ169-173-FLAG,   5-*bcr4*-FLAG-IF   and

3-IF-*bcr4*Δ164-173-FLAG,   or   5-*bcr4*-FLAG-IF   and   3-IF-*bcr4*Δ159-173-FLAG   using

pMGKU417 as the template. Each amplified fragment was self-ligated by the In-Fusion

Cloning System to obtain pMGKU418, pMGKU419 or pMGKU420, respectively. For BscI

complementation, a PCR was carried out with the primers of B1-bscI-V5 and

3-B2-bscI-comp using *B. bronchiceptica* S798 genomic DNA as the template. The resulting

PCR product was cloned into pDONR201 to obtain pMGKU421 using the Gateway cloning

system. These plasmids were mixed with pRK-R4-R3-F, pDONR-*fha*P and pDONR-*rrnB* (10) to obtain pMGKU422, pMGKU423, pMGKU424, pMGKU425 and pMGKU426 using the Gateway cloning system.

Preparation of proteins from culture supernatants and whole bacterial cell lysates.

Proteins secreted into bacterial culture supernatants and whole bacterial cell lysates were prepared as described previously (40). The loaded sample amounts were adjusted by the A_{600} of each bacterial culture to load samples prepared from the same amounts of bacteria. The protein samples were separated by SDS-PAGE and analyzed by Western blotting.

Antibodies.

Anti-BteA, anti-BopD, anti-Bsp22 and anti-Bcr4 antibodies were purified from rabbit serum in our previous study (7, 10, 24, 41). Mouse anti-V5 monoclonal antibody was purchased from Santa Cruz Biotechnology. To detect filamentous hemagglutinin (FHA) signals, we used mouse anti-FHA serum kindly gifted from K. Kamachi (National Institute of Infectious Diseases). To prepare the anti-BscI and anti-BscJ antibodies, the peptides corresponding to the C-terminus regions of BscI (CKAIGRATQNVDTLARMS) and BscJ (CRGEGRGGAGAGATEGAGHD) were conjugated with hemocyanin from keyhole limpets (Sigma), respectively, by using 3-maleimidobenzoic acid N-hydroxysuccinimide ester

(Sigma). These cross-linked peptides were used to immunize rabbits, and the resulting

anti-sera were incubated with the same peptides immobilized on epoxy-activated

sepharose 6B (Amersham) to obtain specific Ig-fractions.

Immunofluorescence staining of L2 cells infected with *B. bronchiseptica.*

Immunofluorescent staining assay was performed as previously described (40), with

slight modifications. Briefly, L2 cells were seeded on coverslips in 6-well plates and

incubated overnight. In order to detect adequate amounts of Bsp22 signals, we infected the

L2 cells with *B. bronchiseptica* cultured in iron-depleted SS medium at an MOI of 2000. To

avoid excessive killing of L2 cells with bacteria, the plates were not centrifuged. After being

incubated for 1 h at 37°C under a 5% CO_2 atmosphere, the infected L2 cells were

immunostained. Average numbers of Bsp22 signals on single L2 cells were measured by a

fluorescence microscope.

LDH Assays.

LDH assays were performed as previously described (40). Briefly, to examine whether

lactate dehydrogenase (LDH) is released from *B. bronchiseptica*-infected cells, 5.0×10^4

cells/well of L2 cells seeded in 24-well plates were infected with bacteria at an MOI of 200.

The plates were centrifuged at $900 \times g$ for 5 min and incubated for 3 h at 37°C under a 5%

CO_2 atmosphere. The amounts of LDH were measured spectrophotometrically using a

Cyto-Tox 96 Non-radioactive Cytotoxicity Assay kit (Promega). The LDH (%) level is shown

as a ratio, with the value obtained from the well treated with 0.1% Triton X-100 set as

100%.

Statistical analyses

The statistical analyses were performed using the nonparametric unpaired t-test with a

one-tailed p-value with Prism ver. 5.0f software (Graphpad, La Jolla, CA). Values of $p<0.05$

were considered significant.

Acknowledgements

This work was supported in part by grants from the Ministry of Education, Culture, Sports, Science and Technology and the Japan Society for the Promotion of Science (KAKENHI) nos. 19K07542 (to A.A.), 20K07485 (to A.K.) and 19K07561 (to T.H.).

The funders had no role in the study design, date collection or analysis, the decision to publish, or the preparation of manuscript.

References

1. Goodnow RA. 1980. Biology of *Bordetella bronchiseptica*. Microbiol Rev 44:722-738.

2. Mattoo S, Cherry JD. 2005. Molecular pathogenesis, epidemiology, and clinical manifestations of respiratory infections due to *Bordetella* pertussis and other *Bordetella* subspecies. Clin Microbiol Rev 18:326-382.

3. Cornelis GR, Van Gijsegem F. 2000. Assembly and function of type III secretory systems. Annu Rev Microbiol 54:735-774.

4. Büttner D. 2012. Protein export according to schedule: architecture, assembly, and regulation of type III secretion systems from plant- and animal-pathogenic bacteria. Microbiol Mol Biol Rev 76:262-310.

5. Deng W, Marshall NC, Rowland JL, McCoy JM, Worrall LJ, Santos AS, Strynadka NCJ, Finlay BB. 2017. Assembly, structure, function and regulation of type III secretion systems. Nat Rev Microbiol 15:323-337.

6. Kuwae A, Ohishi M, Watanabe M, Nagai M, Abe A. 2003. BopB is a type III secreted protein in *Bordetella bronchiseptica* and is required for cytotoxicity against cultured mammalian cells. Cell Microbiol 5:973-983.

7. Nogawa H, Kuwae A, Matsuzawa T, Abe A. 2004. The type III secreted protein BopD in *Bordetella bronchiseptica* is complexed with BopB for pore formation on the host plasma membrane. J Bacteriol 186:3806-3813.

8.   Yuk MH, Harvill ET, Cotter PA, Miller JF. 2000. Modulation of host immune

responses, induction of apoptosis and inhibition of NF-*kappa*B activation by the

Bordetella type III secretion system. Mol Microbiol 35:991-1004.

9.   Panina EM, Mattoo S, Griffith N, Kozak NA, Yuk MH, Miller JF. 2005. A genome-wide

screen identifies a *Bordetella* type III secretion effector and candidate effectors in

other species. Mol Microbiol 58:267-279.

10.   Kuwae A, Matsuzawa T, Ishikawa N, Abe H, Nonaka T, Fukuda H, Imajoh-Ohmi S,

Abe A. 2006. BopC is a novel type III effector secreted by *Bordetella bronchiseptica*

and has a critical role in type III-dependent necrotic cell death. J Biol Chem

281:6589-6600.

11.   Nagamatsu K, Kuwae A, Konaka T, Nagai S, Yoshida S, Eguchi M, Watanabe M,

Mimuro H, Koyasu S, Abe A. 2009. *Bordetella* evades the host immune system by

inducing IL-10 through a type III effector, BopN. J Exp Med 206:3073-3088.

12.   Abe A, Nishimura R, Kuwae A. 2017. *Bordetella* effector BopN is translocated into

host cells via its N-terminal residues. Microbiol Immunol 61:206-214.

13.   Ahuja U, Liu M, Tomida S, Park J, Souda P, Whitelegge J, Li H, Harvill ET, Parkhill J,

Miller JF. 2012. Phenotypic and genomic analysis of hypervirulent human-associated

Bordetella bronchiseptica. BMC Microbiol 12:167.

14.   Kurushima J, Kuwae A, Abe A. 2012. The type III secreted protein BspR regulates

the virulence genes in *Bordetella bronchiseptica*. PLoS One 7:e38925.

15.   Medhekar B, Shrivastava R, Mattoo S, Gingery M, Miller JF. 2009. *Bordetella* Bsp22

522   forms a filamentous type III secretion system tip complex and is immunoprotective in

523   vitro and in vivo. Mol Microbiol 71:492-504.

524   16.   Ahuja U, Shokeen B, Cheng N, Cho Y, Blum C, Coppola G, Miller JF. 2016.

525   Differential regulation of type III secretion and virulence genes in *Bordetella*

526   *pertussis* and *Bordetella bronchiseptica* by a secreted anti-sigma factor. Proc Natl

527   Acad Sci U S A 113:2341-2348.

528   17.   Feldman MF, Cornelis GR. 2003. The multitalented type III chaperones: all you can

529   do with 15 kDa. FEMS Microbiol Lett 219:151-158.

530   18.   Stebbins CE, Galán JE. 2001. Maintenance of an unfolded polypeptide by a cognate

531   chaperone in bacterial type III secretion. Nature 414:77-81.

532   19.   Lloyd SA, Norman M, Rosqvist R, Wolf-Watz H. 2001. *Yersinia* YopE is targeted for

533   type III secretion by N-terminal, not mRNA, signals. Mol Microbiol 39:520-531.

534   20.   Lee SH, Galán JE. 2004. *Salmonella* type III secretion-associated chaperones

535   confer secretion-pathway specificity. Mol Microbiol 51:483-495.

536   21.   Quinaud M, Chabert J, Faudry E, Neumann E, Lemaire D, Pastor A, Elsen S,

537   Dessen A, Attree I. 2005. The PscE-PscF-PscG complex controls type III secretion

538   needle biogenesis in *Pseudomonas aeruginosa*. J Biol Chem 280:36293-36300.

539   22.   Wood SE, Jin J, Lloyd SA. 2008. YscP and YscU switch the substrate specificity of

540   the *Yersinia* type III secretion system by regulating export of the inner rod protein

YscI. J Bacteriol 190:4252-4262.

23.   Monlezun L, Liebl D, Fenel D, Grandjean T, Berry A, Schoehn G, Dessein R, Faudry

E, Attree I. 2015. PscI is a type III secretion needle anchoring protein with in vitro

polymerization capacities. Mol Microbiol 96:419-436.

24.   Nishimura R, Abe A, Sakuma Y, Kuwae A. 2018. *Bordetella bronchiseptica* Bcr4

antagonizes the negative regulatory function of BspR via its role in type III secretion.

Microbiol Immunol 62:743-754.

25.   Wattiau P, Bernier B, Deslée P, Michiels T, Cornelis GR. 1994. Individual chaperones

required for Yop secretion by *Yersinia*. Proc Natl Acad Sci U S A 91:10493-10497.

26.   Kamanova J. 2020. *Bordetella* Type III Secretion Injectosome and Effector Proteins.

Front Cell Infect Microbiol 10:466.

27.   Kurushima J, Kuwae A, Abe A. 2012. Iron starvation regulates the type III secretion

system in *Bordetella bronchiseptica*. Microbiol Immunol 56:356-362.

28.   Sun P, Tropea JE, Austin BP, Cherry S, Waugh DS. 2008. Structural characterization

of the *Yersinia pestis* type III secretion system needle protein YscF in complex with

its heterodimeric chaperone YscE/YscG. J Mol Biol 377:819-830.

29.   Day JB, Guller I, Plano GV. 2000. *Yersinia pestis* YscG protein is a Syc-like

chaperone that directly binds yscE. Infect Immun 68:6466-6471.

30.   Francis MS, Aili M, Wiklund ML, Wolf-Watz H. 2000. A study of the YopD-lcrH

interaction from *Yersinia pseudotuberculosis* reveals a role for hydrophobic residues

within the amphipathic domain of YopD. Mol Microbiol 38:85-102.

31.   Woestyn S, Sory MP, Boland A, Lequenne O, Cornelis GR. 1996. The cytosolic SycE

and SycH chaperones of *Yersinia* protect the region of YopE and YopH involved in

translocation across eukaryotic cell membranes. Mol Microbiol 20:1261-1271.

32.   Yang H, Shan Z, Kim J, Wu W, Lian W, Zeng L, Xing L, Jin S. 2007. Regulatory role

of PopN and its interacting partners in type III secretion of *Pseudomonas aeruginosa*.

J Bacteriol 189:2599-2609.

33.   Studier FW, Moffatt BA. 1986. Use of bacteriophage T7 RNA polymerase to direct

selective high-level expression of cloned genes. J Mol Biol 189:113-130.

34.   Stainer DW, Scholte MJ. 1970. A simple chemically defined medium for the

production of phase I *Bordetella pertussis*. J Gen Microbiol 63:211-220.

35.   Ogino T, Ohno R, Sekiya K, Kuwae A, Matsuzawa T, Nonaka T, Fukuda H,

Imajoh-Ohmi S, Abe A. 2006. Assembly of the type III secretion apparatus of

enteropathogenic *Escherichia coli*. J Bacteriol 188:2801-2811.

36.   Kuwae A, Momose F, Nagamatsu K, Suyama Y, Abe A. 2016. BteA Secreted from

the *Bordetella bronchiseptica* Type III Secetion System Induces Necrosis through an

Actin Cytoskeleton Signaling Pathway and Inhibits Phagocytosis by Macrophages.

PLoS One 11:e0148387.

37.   Halder PK, Roy C, Datta S. 2019. Structural and functional characterization of type

three secretion system ATPase PscN and its regulator PscL from *Pseudomonas*

aeruginosa. Proteins 87:276-288.

38.   Sekiya K, Ohishi M, Ogino T, Tamano K, Sasakawa C, Abe A. 2001. Supermolecular

structure of the enteropathogenic *Escherichia coli* type III secretion system and its

direct interaction with the EspA-sheath-like structure. Proc Natl Acad Sci U S A

98:11638-11643.

39.   Donnenberg MS, Kaper JB. 1991. Construction of an *eae* deletion mutant of

enteropathogenic *Escherichia coli* by using a positive-selection suicide vector. Infect

Immun 59:4310-4317.

40.   Goto M, Hanawa T, Abe A, Kuwae A. 2020. Transcriptional Downregulation of a Type

III Secretion System under Reducing Conditions in *Bordetella pertussis*. J Bacteriol

202.

41.   Kurushima J, Kuwae A, Abe A. 2012. Btc22 chaperone is required for secretion and

stability of the type III secreted protein Bsp22 in *Bordetella bronchiseptica*. FEMS

Microbiol Lett 331:144-151.

  Figure legends

  Fig. 1. The interaction of Bcr4 with inner rod protein BscI. (A) The *bcr4* gene and its

peripheral genes localized in the T3SS apparatus locus (*bsc* locus) on the *B.*

bronchiseptica S798 chromosome are depicted. (B) The purified Bcr4 (Bcr4-Strep) was

loaded on Strep-Tactin beads. Then, the beads were mixed with the lysates prepared from

E. coli BL21 producing BscI or BscK tagged with V5 (BscI-V5 or BscK-V5), respectively.

After 3 h incubation at 4°C, the supernatant fraction (Sup) and pellet fraction (Pellet)

samples were prepared. The Sup and Pellet samples were separated by SDS-PAGE and

analyzed by Western blotting (WB) with anti-V5 antibody (top). Pellet samples were also

stained with Coomassie Brilliant Blue (CBB, bottom). (C) The purified BscI (BscI-Strep) or

N-terminal moiety of BteA (amino acid region 1–312, BteA-N-Strep) were loaded on

Strep-Tactin beads. Then, the beads were mixed with the lysate prepared from *E. coli* BL21

to produce Bcr4 tagged with V5 (Bcr4-V5) at 4°C for 3 h. The prepared Sup and Pellet

samples were separated by SDS-PAGE and analyzed by WB with anti-V5 antibody (top).

Pellet samples were also stained with CBB (bottom). Experiments were performed at least

three times, and representative data are shown.

Fig. 2. The Bcr4 domain is required for interaction with BscI and T3SS activity. (A)

Bcr4 derivatives used for pull-down assay in panel B and the analysis of T3SS activity in

panel C are depicted. The yellow part shows the Bcr4 domain required for T3SS activity. (B)

The purified Bcr4-FL, Bcr4Δ58-109 or Bcr4Δ110-173 (Bcr4-FL-Strep, Bcr4Δ58-109-Strep or

Bcr4Δ110-17-Strep) was loaded on Strep-Tactin beads, and then the beads were mixed

with the lysate prepared from *E. coli* BL21, producing BscI-V5, respectively. After 3 h

incubation at 4°C, the supernatant fraction (Sup) and pellet fraction (Pellet) samples were

prepared. The Sup and Pellet samples were separated by SDS-PAGE and analyzed by

Western blotting (WB) with anti-V5 antibody (top), and stained with Coomassie Brilliant

Blue (CBB, bottom). (C) The *B. bronchiseptica* wild-type strain, Bcr4-deficient strain (ΔBcr4),

and ΔBcr4 producing Bcr4-FL-FLAG (ΔBcr4+Bcr4-FL-FLAG), Bcr4Δ169-173-FLAG

(ΔBcr4+Bcr4Δ169-173-FLAG), Bcr4Δ164-173-FLAG (ΔBcr4+Bcr4Δ164-173-FLAG), and

Bcr4Δ159-173-FLAG (ΔBcr4+Bcr4Δ159-173-FLAG) were cultured in SS medium. Whole

cell lysates (WCL) and supernatant fraction (Sup) samples were separated by SDS-PAGE

and analyzed by Western blotting using antibodies against Bcr4, FLAG, BteA or BopD. AP

indicates the antigen peptide used for anti-Bcr4 antibody generation. WCL and Sup

samples were prepared from equal volumes of bacterial culture. When we performed

western blotting using anti-BteA or BopD antibodies, we loaded a 100- or 50-fold smaller

amount of ΔBcr4+Bcr4-FL-FLAG, ΔBcr4+Bcr4Δ169-173-FLAG and

ΔBcr4+Bcr4Δ164-173-FLAG Sup samples on the SDS-PAGE gel than on the wild type,

ΔBcr4 and ΔBcr4+Bcr4Δ159-173-FLAG Sup samples to avoid obtaining excess signal

intensities. Experiments were performed at least three times, and representative data are

shown.

Fig. 3. The effect of BscI deletion on T3SS activity in *B. bronchiseptica*. Whole cell lysates (WCL) and supernatant fraction (Sup) samples were prepared from the wild-type strain, ΔBscI (BscI-deficient strain), ΔBscI+BscI (BscI-complemented strain) and ΔBscN (T3SS-inactive strain) cultured in SS medium. WCL and Sup samples were separated by SDS-PAGE and analyzed by Western blotting using antibodies against BscI, BteA, BopD or FHA. BscI tagged with Strep (BscI-Strep) was loaded as a control. *BscI-Strep signal; **NS (nonspecific signals); ***A faint signal of BscI. Loaded WCL and Sup samples were prepared from equal volumes of bacterial culture. Experiments were performed at least three times, and representative data are shown.

Fig. 4. Immunofluorescent staining of Bsp22 on L2 cells infected with *B. bronchiseptica* and the results of the LDH assay. (A) L2 cells were infected with the wild-type strain, ΔBscI (BscI-deficient strain), ΔBscI+BscI (BscI-complemented strain) or ΔBscN (T3SS-inactive strain) as described in the Materials and Methods. After fixation, cells were stained with anti-Bsp22 antibody (green) and rhodamine phalloidin (red). Arrowheads indicate the Bsp22 (green) signals. (B) Bsp22 signals per one cell were counted under a fluorescent microscope. At least 120 cells were randomly chosen. (C) The results of LDH assay at an MOI of 200 are shown as a histogram. Error bars indicate the SEMs from triplicate experiments. *$P<0.05$. Experiments were performed at least three

times, and representative data are shown.

Fig. 5. The effect of Bcr4 on BscI stability in *B. bronchiseptica*. The whole cell lysates

(WCL) were prepared from wild-type strain, ΔBcr4 (Bcr4-deficient strain), ΔBspR

(BspR-deficient strain) or ΔBspRΔBcr4 (BspR- and Bcr4-deficient strain) cultured in SS

medium. WCL were separated by SDS-PAGE and analyzed by Western blotting with

antibodies against BscI (inner-rod protein), BscJ (inner-ring protein) and Bcr4. The samples

were prepared from equal volumes of bacterial culture. When we carried out western

blotting using anti-BscJ or Bcr4 antibodies, we loaded a 10-fold-smaller amount of ΔBspR

or ΔBspRΔBcr4 WCL samples on the SDS-PAGE than WT or ΔBcr4 WCL samples to avoid

obtaining excess signal intensities. The production of type III secreted proteins and the

secretion level of each strain are shown as native (+), high (+++) or null (–). Experiments

were performed at least three times, and representative data are shown.

Fig. 6. Schematic depiction of the Bcr4 and BscI functions. In the *B. bronchiseptica*

bacterial cytosol, Bcr4 (blue) stabilizes BscI (orange). Then, BscI is presumed to be

localized to the T3SS machinery (gray) and form the inner rod. BscI is essential for T3SS

activity and secreted from the bacterial cell. BscJ (pink) is predicted to function as an inner

membrane ring. Bsp22 (green) forms a filament-like structure. OM and IM indicate the outer

membrane (white) and inner membrane (white), respectively.

Fig. 1. The interaction of Bcr4 with inner rod protein BscI. (A) The *bcr4* gene and its peripheral genes localized in the T3SS apparatus locus (*bsc* locus) on the *B. bronchiseptica* S798 chromosome are depicted. (B) The purified Bcr4 (Bcr4-Strep) was loaded on Strep-Tactin beads. Then, the beads were mixed with the lysates prepared from *E. coli* BL21 producing BscI or BscK tagged with V5 (BscI-V5 or BscK-V5), respectively. After 3 h incubation at 4°C, the supernatant fraction (Sup) and pellet fraction (Pellet) samples were prepared. The Sup and Pellet samples were separated by SDS-PAGE and analyzed by Western blotting (WB) with anti-V5 antibody (top). Pellet samples were also stained with Coomassie Brilliant Blue (CBB, bottom). (C) The purified BscI (BscI-Strep) or N-terminal moiety of BteA (amino acid region 1–312, BteA-N-Strep) were loaded on Strep-Tactin beads. Then, the beads were mixed with the lysate prepared from *E. coli* BL21 to produce Bcr4 tagged with V5 (Bcr4-V5) at 4°C for 3 h. The prepared Sup and Pellet samples were separated by SDS-PAGE and analyzed by WB with anti-V5 antibody (top). Pellet samples were also stained with CBB (bottom). Experiments were performed at least three times, and representative data are shown.

Fig. 1

Fig. 2. The Bcr4 domain is required for interaction with BscI and T3SS activity. (A) Bcr4 derivatives used for pull-down assay in panel B and the analysis of T3SS activity in panel C are depicted. The yellow part shows the Bcr4 domain required for T3SS activity. (B) The purified Bcr4-FL, Bcr4Δ58-109 or Bcr4Δ110-173 (Bcr4-FL-Strep, Bcr4Δ58-109-Strep or Bcr4Δ110-17-Strep) was loaded on Strep-Tactin beads, and then the beads were mixed with the lysate prepared from *E. coli* BL21, producing BscI-V5, respectively. After 3 h incubation at 4°C, the supernatant fraction (Sup) and pellet fraction (Pellet) samples were prepared. The Sup and Pellet samples were separated by SDS-PAGE and analyzed by Western blotting (WB) with anti-V5 antibody (top), and stained with Coomassie Brilliant Blue (CBB, bottom). (C) The *B. bronchiseptica* wild-type strain, Bcr4-deficient strain (ΔBcr4), and ΔBcr4 producing Bcr4-FL-FLAG (ΔBcr4+Bcr4-FL-FLAG), Bcr4Δ169-173-FLAG (ΔBcr4+Bcr4Δ169-173-FLAG), Bcr4Δ164-173-FLAG (ΔBcr4+Bcr4Δ164-173-FLAG), and Bcr4Δ159-173-FLAG (ΔBcr4+Bcr4Δ159-173-FLAG) were cultured in SS medium. Whole cell lysates (WCL) and supernatant fraction (Sup) samples were separated by SDS-PAGE and analyzed by Western blotting using antibodies against Bcr4, FLAG, BteA or BopD. AP indicates the antigen peptide used for anti-Bcr4 antibody generation. WCL and Sup samples were prepared from equal volumes of bacterial culture. When we performed western blotting using anti-BteA or BopD antibodies, we loaded a 100- or 50-fold smaller amount of ΔBcr4+Bcr4-FL-FLAG, ΔBcr4+Bcr4Δ169-173-FLAG and ΔBcr4+Bcr4Δ164-173-FLAG Sup samples on the SDS-PAGE gel than on the wild type, ΔBcr4 and ΔBcr4+Bcr4Δ159-173-FLAG Sup samples to avoid obtaining excess signal intensities. Experiments were performed at least three times, and representative data are shown.

Fig. 2

Fig. 3. The effect of BscI deletion on T3SS activity in *B. bronchiseptica*. Whole cell lysates (WCL) and supernatant fraction (Sup) samples were prepared from the wild-type strain, ΔBscI (BscI-deficient strain), ΔBscI+BscI (BscI-complemented strain) and ΔBscN (T3SS-inactive strain) cultured in SS medium. WCL and Sup samples were separated by SDS-PAGE and analyzed by Western blotting using antibodies against BscI, BteA, BopD or FHA. BscI tagged with Strep (BscI-Strep) was loaded as a control. *BscI-Strep signal; **NS (nonspecific signals); ***A faint signal of BscI. Loaded WCL and Sup samples were prepared from equal volumes of bacterial culture. Experiments were performed at least three times, and representative data are shown.

Fig. 3

Fig. 4. Immunofluorescent staining of Bsp22 on L2 cells infected with *B. bronchiseptica* and the results of the LDH assay. (A) L2 cells were infected with the wild-type strain, ΔBscI (BscI-deficient strain), ΔBscI+BscI (BscI-complemented strain) or ΔBscN (T3SS-inactive strain) as described in the Materials and Methods. After fixation, cells were stained with anti-Bsp22 antibody (green) and rhodamine phalloidin (red). Arrowheads indicate the Bsp22 (green) signals. (B) Bsp22 signals per one cell were counted under a fluorescent microscope. At least 120 cells were randomly chosen. (C) The results of LDH assay at an MOI of 200 are shown as a histogram. Error bars indicate the SEMs from triplicate experiments. *$P<0.05$. Experiments were performed at least three times, and representative data are shown.

Fig. 4

Fig. 5. The effect of Bcr4 on BscI stability in *B. bronchiseptica*. The whole cell lysates (WCL) were prepared from wild-type strain, ΔBcr4 (Bcr4-deficient strain), ΔBspR (BspR-deficient strain) or ΔBspRΔBcr4 (BspR- and Bcr4-deficient strain) cultured in SS medium. WCL were separated by SDS-PAGE and analyzed by Western blotting with antibodies against BscI (inner-rod protein), BscJ (inner-ring protein) and Bcr4. The samples were prepared from equal volumes of bacterial culture. When we carried out western blotting using anti-BscJ or Bcr4 antibodies, we loaded a 10-fold-smaller amount of ΔBspR or ΔBspRΔBcr4 WCL samples on the SDS-PAGE than WT or ΔBcr4 WCL samples to avoid obtaining excess signal intensities. The production of type III secreted proteins and the secretion level of each strain are shown as native (+), high (+++) or null (–). Experiments were performed at least three times, and representative data are shown.

Fig. 5

BscI

Bsp22 (filament-like structure)

T3SS

OM

BscI (Inner rod)
is essential for T3SS activity

IM

BscJ (Inner membrane ring)

Bcr4   BscI

Bcr4 stabilizes BscI

Fig. 6. Schematic depiction of the Bcr4 and BscI functions. In the *B. bronchiseptica* bacterial cytosol, Bcr4 (blue) stabilizes BscI (orange). Then, BscI is presumed to be localized to the T3SS machinery (gray) and form the inner rod. BscI is essential for T3SS activity and secreted from the bacterial cell. BscJ (pink) is predicted to function as an inner membrane ring. Bsp22 (green) forms a filament-like structure. OM and IM indicate the outer membrane (white) and inner membrane (white), respectively.

Fig. 6

Table 1. Bacterial strains and plasmids used in the study

Strain or plasmid	Relevant genotype or description	Reference or source
Strains		
B. bronchiseptica		
S798	wild-type strain	6
ΔBscI	BscI-deficient strain	This study
△BscN	BscN-deficient strain	6
△Bcr4	Bcr4-deficient strain	24
△BspR	BspR-deficient strain	14
△BspR△Bcr4	BspR and Bcr4-deficient strain	24
E. coli		
DH10B	Host strain for pDONR201, p99-*ccdB*-V5 and pRK415-R4-R3-F	Invitrogen
BL21	Host strain for pColdII	33
Sm10λ*pir*	Host strain for pABB-CRS2	6
Plasmids		
pDONR201	DNA cloning vector, Km^r	Invitrogen
pMGKU404	pDONR201, *bscI* gene tagged with V5 sequence at the C terminus	This study
pMGKU405	pDONR201, *bscK* gene tagged with V5 sequence at the C terminus	This study
pMGKU406	pDONR201, *bsc4* gene tagged with V5 sequence at the C terminus	This study
p99*ccdB*-V5	Expression vector for V5 tagged gene, Amp^r	35
pMGKU407	p99-*ccdB*-V5, *bscI* gene tagged with V5 sequence at the C terminus	This study
pMGKU408	p99-*ccdB*-V5, *bscK* gene tagged with V5 sequence at the C terminus	This study
pMGKU409	p99-*ccdB*-V5, *bsc4* gene tagged with V5 sequence at the C terminus	This study
pColdII	Expression vector for His tagged gene, Amp^r	TAKARA
pMGKU410	pColdII, *bcr4* gene tagged with Strep sequence at the C terminus	This study
pMGKU411	pColdII, *bscI* gene tagged with Strep sequence at the C terminus	This study
pMGKU412	pColdII, *bcr4* gene lacking amino acids region 58-109 tagged with Strep sequence at the C terminus	This study
pMGKU413	pColdII, *bcr4* gene lacking amino acids region 110-173 tagged with Strep sequence at the C terminus	This study
pColdII-*bteA*-N-Strep	pColdII, *bteA* gene coding amino acids region 1-312 tagged with Strep sequence at the C terminus	36
pMGKU414	pDONR201, *bscI* gene	This study
pMGKU415	pDONR201, *bscI* gene containing internal sequence-deletion and its flanking region	This study
pABB-CRS2	Suicide vector for conjugation, Amp^r	38
pMGKU416	pABB-CRS2, bscI gene containing internal sequence-deletion and its flanking region	This study
pDONR-*bcr4*	pDONR201, *bcr4* gene for complementation	24
pMGKU417	pDONR201, *bcr4* gene tagged with FLAG sequence at the C terminus for complementation	This study
pMGKU418	pDONR201, *bcr4* gene lacking amino acids region 169-173 tagged with FLAG sequence at the C terminus	This study
pMGKU419	pDONR201, *bcr4* gene lacking amino acids region 164-173 tagged with FLAG sequence at the C terminus	This study
pMGKU420	pDONR201, *bcr4* gene lacking amino acids region 159-173 tagged with FLAG sequence at the C terminus	This study
pMGKU421	pDONR201, *bscI* gene for complementation	This study
pRK415-R4-R3-F	pRK415, recombination sites for MultiSite Gateway, Tet^r	10
pDONR-*fhaP*	pDONR-P4-P1R, *fha* promoter	10
pDONR-*rrnB*	pDONR-P2R-P3, *rrnB* terminator	10
pMGKU422	pRK415-R4-R3-F, *bcr4* gene tagged with FLAG sequence at the C terminus for complementation	This study
pMGKU423	pRK415-R4-R3-F, *bcr4* gene lacking amino acids region 169-173 tagged with FLAG sequence at the C terminus	This study
pMGKU424	pRK415-R4-R3-F, *bcr4* gene lacking amino acids region 164-173 tagged with FLAG sequence at the C terminus	This study
pMGKU425	pRK415-R4-R3-F, *bcr4* gene lacking amino acids region 159-173 tagged with FLAG sequence at the C terminus	This study
pMGKU426	pRK415-R4-R3-F, *bscI* gene	This study

Table 2. Primers used in the study

Name	Sequence (5'→3')
B1-bscl-V5	AAAAAGCAGGCTTGTTGGACAGCGGACCCGCC
B2-bscl-V5	AGAAAGCTGGGTTTGACATTCTCGCCAGCGTGTC
B1-bscK-V5	AAAAAGCAGGCTTGGCGCACGCGCTGGTCCCG
B2-bscK-V5	AGAAAGCTGGGTTGGCTTGGGCGGGGAACGAGG
B1-bcr4-comp	AAAAAGCAGGCTGCCAGGTCCGGTCTCGCACCG
B2-bcr4-V5	AGAAAGCTGGGTTTCCAGGAGCTCCAGGTAATG
5-HindIII-bcr4	ATCCGAATTCAAGCTTCATTCAGACTCAGGTTCAGATTC
3-bcr4-Strep	CGGGTGGCTCCATCCAGGAGCTCCAGGTAATG
5-HindIII-bscl	ATCCGAATTCAAGCTTAATTTGGATCTGACGGCGATC
3-bscl-Strep	CGGGTGGCTCCATGACATTCTCGCCAGCGTGTC
3-Strep-HindIII	GCAGGTCGACAAGCTTTCATTTTTCGAACTGCGGGTGGCTCCA
5-bcr4-58-109-Strep-IF	GCCGTCAGCGATCTGTCGGACTGGACGCCTGGTG
3-IF-bcr4-58-109-Strep	CAGATCGCTGACGGCCGCCG
5-bcr4-110-173-Strep-IF	TGGAGCCACCCGCAGTTCGA
3-IF-bcr4-110-173-Strep	CTGCGGGTGGCTCCAGCGGTCCTCCGCCTGTGCCG
B1-bscl	AAAAAGCAGGCTTCATGACTGTTCACGACGAC
B2-bscl	AGAAAGCTGGGTGCACAAGATCCAGCGCGACC
R1-bscl	CGACGTTGGCGTTGATCGCCGTCAGATC
R2-bscl	TCAACGCCAACGTCGACACGCTGGCGAG
5-bcr4-FLAG-IF	CCACCCGCAGTTCGAAAAATGAATTTGGATCTACCCAGCTTTCTTG
3-IF-bcr4-FLAG	TCGAACTGCGGGTGGCTCCATCCAGGACCTCCAGGTAATGGCTCG
3-IF-bcr4△169-173-FLAG	GTCATCCTTGTAGTCTAATGGCTCGGACTGCAACG
3-IF-bcr4△164-173-FLAG	GTCATCCTTGTAGTCCAACGGCGCGGGCCGCATCATG
3-IF-bcr4△159-173-FLAG	GTCATCCTTGTAGTCCATCATGGCGGGTCCGCTGTC
3-B2-bscl-comp	AGAAAGCTGGGTTTGACATTCTCGCCAGCGTGTC