Large-scale multi-omics analysis suggests specific roles for intragenic cohesin in transcriptional regulation

Jiankang Wang[1,2], Masashige Bando[1], Katsuhiko Shirahige[1,2], Ryuichiro Nakato[1,2,*]

[1]Institute for Quantitative Biosciences, The University of Tokyo; [2]Graduate School of Medicine, The University of Tokyo

*Corresponding author. Tokyo 1130023, Japan. Tel: +81-3-5841-1471; Fax: +81-3-5841-7308; E-mail: rnakato@iqb.u-tokyo.ac.jp

Abstract

Cohesin, an essential protein complex for chromosome segregation, regulates transcription through a variety of mechanisms. It is not a trivial task to genome-widely assign the diverse cohesin functions. Moreover, the context-specific roles of cohesin-mediated interactions, especially on intragenic regions, have not been thoroughly investigated. Here we performed a comprehensive characterization of cohesin binding sites in several human cell types. We integrated epigenomic, transcriptomic and chromatin interaction data with and without transcriptional stimulation, to explore context-specific functions of intragenic cohesin related to gene activation. We identified a new subset of cohesin binding sites, decreased intragenic cohesin sites (DICs), which have a different function from previously known ones. The intron-enriched DICs were negatively correlated with transcriptional regulation: a subgroup of DICs were related to enhancer markers and paused RNA polymerase II, whereas others contributed to chromatin architecture. We implemented machine learning and successfully isolated DICs with distinct genomic features. We observed DICs in various cell types, including cells from cohesinopathy patients. These results suggest a previously unidentified function of cohesin at intragenic regions for transcription regulation.

Keywords: Cohesin, Epigenomics, Transcription regulation, Chromosome structure, Machine learning

Introduction

Cohesin, a ring-shaped chromosome-bound protein complex, is required for holding sister chromatids together during certain phases of the cell cycle[1]. Recent studies suggest that cohesin also has a role in transcription regulation, maintenance of chromosome architecture[2] and DNA repair[3]. The context-specific functions of cohesin have been investigated using chromatin immunoprecipitation followed by sequencing (ChIP-seq) and high-throughput chromosome conformation capture (Hi-C). The early study reported that most cohesin-binding sites overlap with CTCF to function as an insulator[4]. Conversely, a group of cohesin is reported to be CTCF independent and co-bind with tissue-specific transcription factors (TFs) to contribute to transcriptional regulation[5, 6], possibly via the mediation of interactions between enhancers and promoters[7]. Other studies using Hi-C suggested that cohesin and CTCF are essential for forming topologically associated domains (TADs), the evolutionarily conserved chromatin domains that are hundreds of kilobases to several megabases in length[8, 9]. These studies focused on cohesin functions with respect to blocked interactions by insulation, or the formation of enhancer-promoter interactions which implicitly assume the positive regulation of gene expression. In contrast, a recent report showed that transcription elongation within gene bodies causes displacement of cohesin binding from chromatin, leading to disruption of cohesin-mediated chromatin loops[10]. Thus a subset of chromatin loops (either end of which may be located within the intragenic regions) mediated by cohesin is suggested to be negatively correlated with gene activation. Whereas modifications in intragenic regions affect transcription events[11, 12, 13], the function of intragenic cohesin has hardly been discussed.

Mutations in cohesin complex and its loader (NIPBL) are observed in the cohesinopathy Cornelia de Lange syndrome (CdLS), a multisystem developmental disorder[14], and in multiple types of cancers[15, 16]. Our previous study found that the diagnostic phenotype of CdLS is very similar to that of CHOPS syndrome[17], which is caused by missense mutations in AFF4, a core component of the super elongation complex. Given the diverse functions of cohesin with respect to gene expression and chromatin folding, the underlying molecular mechanism responsible for the similarity between CdLS and CHOPS is yet unknown. Noteworthily, the CHOPS-related mutations in super elongation complex are also associated with transcriptional regulation by cohesin, indicating a common pathogenetic mechanism of cohesin in both CHOPS and CdLS. It can be a feasible hypothesis that intragenic cohesin has a distinct role that links the phenotypic similarities between CdLS and CHOPS.

Here, we conducted a large-scale epigenomic analysis to clarify the context-specific functions of cohesin sites, especially those in intragenic regions. To investigate the perturbation of cohesin binding sites by gene activation, we generated RNA sequencing (RNA-seq) and ChIP-seq data for cohesin and

several TFs from MCF-7 cells with or without transcription stimulus. We also used many publicly available datasets including Hi-C, ChIP-seq, RNA-seq and chromatin interaction analysis by paired-end tag (ChIA-PET). First, we clarified that a subset of cohesin sites, which we refer to as 'decreased intragenic cohesin sites' (DICs), is distinct from the other groups of cohesin sites. Cohesin bindings on DICs are negatively correlated with transcriptional activation and locus compaction of chromatin. A part of DICs exhibit the high preference for enhancer marks and paused RNA polymerase II, whereas others contribute to chromatin architecture. Second, we performed ChIP-seq and RNA-seq with cohesin-depleted cells and confirmed that cohesin has an active function on DICs. Third, we applied machine learning and captured DICs with distinct epigenomic landscape, which is predictable across cell types. Finally, we conducted plenty of ChIP-seq in other cell types. Importantly, DICs can be observed across multiple cell types, including cells derived from patients with CdLS and CHOPS, in a cell-type-specific manner. The findings from our integrated analysis and machine learning approaches suggest an additional role for cohesin in the regulation of gene expression.

Results

Classification of DICs

MCF-7 cells, when treated with the transcriptional stimulator estradiol[18], is a widely used model for investigating the transcription-dependent perturbation[5]. We prepared ChIP-seq data of cohesin (Rad21), cohesin loader (Mau2), CTCF and several TFs (ER, CBP, P300, AFF4, TAF1) from MCF-7 cells treated with vehicle (control, or Ctrl) or estradiol (E2, 45min). ChIP-seq and RNA-seq data generated in this study and their statistics are summarized in Supplementary Table 1. Other omics data used in this study are listed in Supplementary Tables 2-3. In total, we obtained 76,668 and 89,111 peaks as cohesin binding sites in the E2 and control data, respectively. Next, we examined the stimulation-dependent cohesin sites (Fig. 1a). Although the total number of cohesin peaks decreased after E2 stimulation, the proportion of peaks that increased (9.3%) after stimulation (log-fold change of peak intensity Mvalue[19] > 0.5) was larger than the one that decreased (6.2%) (Mvalue < −0.5) (Fig. 1a, bottom). We also found that around 40% (36.3% for E2, 41.2% for control) of cohesin peaks did not overlap with CTCF peaks (Supplementary Fig. 1a). Such 'cohesin-non-CTCF sites' (hereafter, CNCs) overlapped with peaks of the enhancer markers P300 and CBP (Supplementary Fig. 1b), which is consistent with an earlier ChIP-seq study[5]. The cohesin loader Mau2 also preferred enhancer sites. In fact, 88.7% of CNCs with enhancer markers overlapped with Mau2, and Mau2 was localized at enhancer sites with and without cohesin binding (Supplementary Figs. 1c-d). This result implies the role of Mau2 for enhancer activity or for chromatin interaction, which can precede cohesin localization.

We classified cohesin sites based on gene annotation information (Supplementary Fig. 1e). We

defined 'intragenic cohesin sites' as those located within gene bodies, with the exception of transcription

start sites (TSSs), transcription end sites (TESs) and alternative promoters. As a result, 13.8% of cohesin

sites were identified as intragenic ones, 19.6% of which had enhancer activities. We did not observe a

difference in the proportion of up- or down-regulated cohesin peaks between intergenic and intragenic

sites (Fig. 1a, lower panel). To investigate the correlation of cohesin binding and transcription activation,

we conducted the ChIP-seq of RNA polymerase II (Pol2), and RNA pol II CTD serine-2 phosphorylation

(Pol2ser2) that represents transcription elongation activity[20]. We identified 499 E2-responsive genes for

which the Pol2ser2 signal increased after E2 stimulation (Method). We then validated these genes by

RNA-seq and confirmed that their expressions were mostly up-regulated by E2 stimulation

(Supplementary Fig. 1f). Within the 499 E2-responsive genes, we identified 4,346 intragenic cohesin

peaks, 976 (22.4%) of which were decreased after stimulation (Fig. 1b, Supplementary Fig. 1g). Because

our main interest is the negative correlation between active transcription and the signal intensity of

intragenic cohesin[10], we focused on the intragenic cohesin sites for which the peak intensity decreased

after E2 stimulation. Hereafter we refer to these sites as DICs. Among the E2-responsive genes, 53.5%

(267/499) contained one or more DICs. We found that almost all (82.8%) DICs were located in intronic

regions (Fig. 1c). This result implies a function of DICs on intron, whose mechanism remains

unrecognized[21].

Next, we investigated the correlation between decreased cohesin binding and the level of chromatin

interaction using Hi-C data (GSE99451). Aggregate peak analysis (APA)[22] showed that DICs-centered

chromatin interactions were weakened by E2 treatment, whereas no difference was observed for all

cohesin sites (Fig. 1d). These results suggested that at least some intragenic cohesin was required for

chromatin loop formation, which were disrupted due to the induction of transcription according to

previous study[10]. In contrast to the positive regulation of gene expression by CNCs[6], DICs possibly

function negatively with respect to gene expression. We then applied DLR (distal-to-local interaction ratio)

and ICF (inter-chromosomal fraction of interactions) metrics[10] to represent changes in locus-specific

intra- and inter-chromosomal interactions, respectively. The differences (Δ) for DLR or ICF between two

Hi-C samples represent chromatin compaction (negative value) or de-compaction (positive value). ΔDLR

showed a positive value at DICs (Fig. 1e). In contrast, ΔDLR had a negative peak at all cohesin sites,

whereas all enhancers showed no enrichment. Chromatin compaction at all cohesin sites could be

explained by more frequent *cis*-regulatory interactions after estrogen stimulation. Conversely, DICs did

not show a clear difference as compared with all cohesin sites for ΔICF (Supplementary Fig. 1h). These

results suggested that DICs were involved in intra-chromosome decompaction that created a more open

architecture around DICs.

Classification of LC-DICs and HC-DICs

We next investigated the binding pattern of cohesin and other TFs, including the estrogen receptor

(ER). We found that DICs could be clearly classified into two categories: HC-DICs (high CTCF binding)

and LC-DICs (low CTCF binding), which co-localized with strong and weak CTCF peaks, respectively

(Fig. 1f). LC-DICs had a higher probability of co-binding with many TFs as compared with HC-DICs (Figs.

1f-g). This tendency was similar, but not identical, to cohesin peaks in the other regions. For example,

cohesin localized with strong CTCF on promoters, where many TFs also bound[6, 23] (Supplementary Fig.

2a). A majority of intergenic cohesin sites (possibly insulator sites or TAD boundaries) did not show TF

enrichment (Supplementary Fig. 2b). Moreover, the TFs on LC-DICs (Fig. 1g, Supplementary Fig. 2c,

except Mau2 and P300), including 16 publicly available TFs (Supplementary Table 4), were increased

after E2 treatment. This suggested that enhancer markers Mau2 and P300 localized to LC-DICs even

before stimulation, whereas other TFs, including another enhancer marker CBP, were recruited by E2

stimulation. In addition, the weakened interactions were observed in both LC- and HC-DICs

(Supplementary Fig. 2d).

Fig. 1h showed examples of two E2-responsive genes (*MREG* and *PAK4*; see Supplementary Fig.

2e for publicly available TFs). For instance, at the *MREG* locus, there were both HC-DICs, which

co-localized with strong CTCF signals but with scarcely any TFs, and LC-DICs, which corresponded with

the frequent binding of many TFs yet without strong CTCF signals. Overall, at LC-DICs, the peak

intensity of cohesin decreased after E2 stimulation, whereas that of many TFs increased. Consistently,

for the E2-activated gene *MREG* (Supplementary Fig. 3a), we could also clearly observe weakened

interactions (Fig. 1i) and chromatin decompaction (Supplementary Fig. 3b).

More genomic characteristics were detected by motif analysis (Supplementary Fig. 3c). Not

surprisingly, all types of cohesin showed the motifs of CTCF and CTCFL (BORIS). Specifically, LC-DICs

were highly enriched for motifs of the forkhead box (FOX) protein family, which is responsible for

remodeling chromatin structure[24] and controlling transcription[25]. Of note, FOXA1 is a pioneer factor

before ER activation in MCF-7 cells[26]. Meanwhile, HC-DICs showed motifs for transcription repressors

including the tumor suppressor gene *HIC1*, implying a possible role for HC-DICs in transcription

repression. Taken together, these results highlighted the unique features of LC-DICs and HC-DICs

relative to other cohesin sites.

Characterization of LC-DICs as enhancers

The bindings of the enhancer markers CBP and P300 were frequently observed at LC-DICs (Figs.

1f-h). We confirmed that a significantly higher percentage of LC-DICs overlapped with CBP binding as

compared with other cohesin sites (Fig. 2a, Fisher's exact test). In addition, LC-DICs were also enriched

for enhancer markers H3K27ac and H3K4me1 as well as FANTOM5 enhancers[27] (Fig. 2b,

Supplementary Fig. 4a; publicly available data). In contrast, HC-DICs were not annotated as enhancers.

Moreover, although intergenic cohesin also (including both CTCF-dependent and -independent) in

conjunction with many TFs, they were not highly enriched for enhancer markers (Fig. 2a, Supplementary

Fig. 4b). This is consistent with the finding that only 18% of intergenic cohesin was co-bound with CBP,

which is reasonable because only a subset of intergenic cohesin sites serves as enhancers.

Characterization of chromatin loops on DICs

To explore DIC-mediated loops, we investigated what kind of chromatin loci interacted with

LC-DICs and HC-DICs. Remarkably, when Pol2-mediated chromatin loops identified by ChIA-PET

(GSE33664) were analyzed, the LC-DICs contained multiple Pol2 loops that interacted with the TSS of

the host gene, whereas HC-DICs at the *MREG* locus did not have Pol2 loops (Fig. 2b, red arcs). To

further investigate this tendency, we also analyzed DIC-anchored loops detected by Hi-C data

(GSE99451). Interestingly, this result was in direct contrast to the Hi-C and ChIA-PET Pol2 loops (Fig. 2c,

Fisher's exact test). HC-DICs had a significantly lower occurrence probability with respect to

Pol2-mediated chromatin loops as compared with LC-DICs and all cohesin sites. In contrast, HC-DICs

exhibited a significantly higher occurrence with respect to Hi-C loops. We also compared loops with

CTCF ChIA-PET data (GSE39495) and found that over 81% of HC-DICs overlapped with CTCF loops

(27% for LC-DICs). This result suggested that LC-DICs were anchored by chromatin loops with Pol2 and

other TFs to function as an enhancer in a CTCF-independent manner, whereas HC-DICs were more

likely to interact with CTCF to form chromatin loops to participate in chromatin architecture independently

of the Pol2 machinery.

We then investigated the other anchor sites of the DIC-mediated loops. The other anchor sites of

DIC-mediated Hi-C loops also overlapped with cohesin, which also showed decreasing tendency

(Supplementary Fig. 5a). As shown in Figs 2d-e, LC-DIC loops (ChIA-PET and Hi-C) mainly interacted

with enhancers (40.8%) or promoters (51.2%), which were confirmed by highly enriched active histone

markers (Supplementary Fig. 5b). We also observed that only a subset of LC-DIC loops (19.2%)

interacted with the promoter of their host genes, suggesting that LC-DICs also contribute to the

regulation of distant non-host genes, possibly as intragenic enhancer sites. In contrast, most of the

HC-DIC loops did not interact with promoter or enhancer sites (Figs. 2d-e). Instead, over half of these sites interacted with intronic regions (Fig. 2e; example loci are shown in Supplementary Fig. 5c). In summary, these results suggested that LC-DICs participated in transcriptional regulation, whereas HC-DICs were more likely to connect the intronic regions of two genes.

We also examined the insulation score (IS) from Hi-C data, for which a lower value indicates more insulated regions, e.g., TAD boundaries. Although the IS profile showed clear valley at all, intergenic and intragenic cohesin sites (Fig. 2f, Supplementary Fig. 5d), it peaked at DICs (Fig. 2f, top left). Interestingly, the IS profile for HC-DICs showed bimodal peaks around small valley, whereas there was neither a peak nor valley at LC-DICs (Fig. 2f, lower right). These results suggested that LC-DICs possibly act as enhancers within TADs and that HC-DICs participate in the formation of boundaries, which is consistent with our loop analysis described above.

Assessment of Pol2 stalling on DICs

Pol2 is released from promoter-proximal pausing to transcribe the entire gene body, although it may be temporarily paused by roadblocks within gene bodies[12]. To test whether DICs can function as roadblocks, we investigated the Pol2 enrichment at DICs using our Pol2 and Pol2ser2 ChIP-seq data (Supplementary Fig. 6a). Our Pol2ser2 and public global nuclear run-on sequencing data (GRO-seq, GSE99508) showed that transcription elongation was activated by E2 (Fig. 3a). Moreover, we found that Pol2 peaked at LC-DICs, and its intensity decreased after E2 stimulation (Fig. 3a-b), possibly due to the release of paused Pol2. This tendency toward a decrease in Pol2 binding was statistically significant as compared with the other cohesin sites (Fig. 3c). Importantly, ChIP-seq data treated with E2 for 30min, and public ChIP-seq datasets also showed the decreased Pol2 (Supplementary Figs. 6b-c). Given that bindings of most TFs were increased by E2 stimulation at LC-DICs (Figs. 1g-h), cohesin binding that decreased at LC-DICs was more likely to be accordant with Pol2, rather than TF bindings. In contrast, Pol2ser2 was increased on all DICs due to transcription activation (Fig. 3b, right panel). Pol2ser2 also exhibited peak-like enrichment at LC-DICs, which was increased by E2 stimulation (Figs. 3a-b). This is remarkable given that Pol2 enrichment at LC-DICs decreased significantly after E2 stimulation (Figs. 3b-c). In addition, whereas Pol2 binding on TSSs of DIC-host genes did not show any difference after E2 stimulation, the intensity of Pol2ser2 on TSSs increased (Supplementary Fig. 6d). These results support the model that Pol2 temporarily stalls within DICs, which function as roadblocks, and then is released by the loss of cohesin.

Knockdown analysis of cohesin revealed that cohesin had a function at DICs

Although we observed cohesin binding at DICs that was synchronized with Pol2 binding and was

negatively correlated with gene expression, there is still a possibility that cohesin is "passively" localized

to DICs and therefore does not have any active role with respect to gene expression. To determine

whether cohesin functions in the pausing of Pol2 at LC-DICs, we prepared Pol2 and Pol2ser2 ChIP-seq

data in the absence and presence of Rad21-specific siRNA (siRad21) to generate Rad21 knockdown

(KD; Supplementary Fig. 6e) and investigate the effect at DICs. Pol2 binding before E2 stimulation was

significantly decreased by siRad21 (site 3 vs. 1 in Fig. 3d, KD_Ctrl as compared with WT_Ctrl in Figs.

3e-f) and had a similarly low level of binding in E2-treated wild-type cells (site 2 in Fig. 3d, WT_E2 in Figs.

3e-f). Thus, KD of cohesin indeed decreased Pol2 on LC-DICs, indicating that cohesin binding on

LC-DICs is not passive but is required for Pol2 stalling. Pol2 binding in KD_Ctrl cells was not affected by

E2-stimulation (site 4 vs. 3 in Fig. 3d, KD_E2 as compared with KD_Ctrl in Figs. 3e-f), possibly because

Pol2 that was paused in WT_Ctrl cells had already been released in KD_Ctrl cells. Importantly, the effect

of siRad21 on the Pol2 signal at TSSs of E2-responsive genes was distinct from LC-DICs, in which Pol2

binding did not change significantly after E2 stimulation in WT cells (from WT_Ctrl to WT_E2 in Figs. 3d

and 3f) but decreased after siRad21 in stimulated cells (from WT_E2 to KD_E2 in Figs. 3d and 3f). These

results also supported the model that the loss of cohesin binding causes the release of paused Pol2 on

LC-DICs.

Interestingly, siRad21 did not largely affect Pol2ser2 binding. Pol2ser2 levels on LC-DICs showed

no obvious difference between WT and siRad21 cells (Figs. 3d-f). In KD_Ctrl cells, there was no more

stalling at LC-DICs, but there was also no stimulating effect of E2; thus Pol2ser2 did not change from

WT_Ctrl to KD_Ctrl. In KD_E2 cells, transcription was activated by E2 stimulation but was limited by the

loss of cohesin on TSSs, and thus Pol2ser2 binding changed slightly from WT_E2 to KD_E2. To explore

changes in the expression level of genes that harbor LC-DICs after siRad21 treatment, we conducted

RNA-seq with siRad21 (Supplementary Fig. 6f). Without E2 treatment, siRad21 did not significantly affect

gene expression (p=0.23, KD_Ctrl as compared with WT_Ctrl). After E2 treatment, siRad21 moderately

affected gene expression (p=0.005, KD_E2 as compared with WT_E2), possibly because only a subset

of Pol2 that had paused on LC-DICs represented productive Pol2. We also quantitatively compared Pol2

and Pol2ser2 signals under four different conditions on various cohesin sites (Fig. 3f, Supplementary Fig.

6g). The results confirmed the significantly reduced bindings of Pol2 in WT_E2, KD_Ctrl and KD_E2 as

compared with WT_Ctrl cells (Fig. 3f, Mann–Whitney U-test, one-sided). Such a tendency was distinct

from those involving the TSSs of E2-responsive genes, up-regulated and non-changed intragenic

cohesin, other cohesin sites, and also other enhancer sites. Our results suggested that the role of

LC-DICs involves Pol2 pausing, which is different from the known roles of other cohesin sites.

In Figs. 1g-h, we showed the elevated binding of many TFs on DICs. To investigate whether the

increased binding of multiple TFs is caused by a decrease in cohesin binding, we generated ChIP-seq

data for CBP, P300 and Mau2 from siRad21 cells. Remarkably, a cohesin deficiency resulted in stronger

binding of those TFs at LC-DICs, which surpassed the level in ER-stimulated WT cells (dashed arrow in

Fig. 3g, Supplementary Fig. 7a-b). In contrast, there was little effect at the other intragenic enhancer site

(Fig. 3g). Considering that E2 stimulation recruits TFs by estrogen responsive elements in WT cells, the

increased binding of TFs in non-E2-stimulated siRad21 cells suggested that cohesin suppresses TF

binding at LC-DICs in some way, and this suppression is removed by the loss of cohesin, which is

followed by increased TF localization. Considering the chromatin de-compaction by E2 stimulation

shown in Fig. 1e, the increased binding of TFs at LC-DICs can be, at least in part, explained by a more

accessible chromatin structure near the LC-DICs, which is induced by the disruption of cohesin-mediated

interactions.

Machine learning analysis of DIC features

To confirm our findings, we implemented machine learning (ML) approaches (Supplementary Fig.

8a). We generated an integrated data matrix consisting of 175 features from genomic, transcriptomic and

epigenomic data for all cohesin sites (Supplementary Table 5; Methods). Especially, this matrix includes

features related to genomic location (e.g., intragenic or TSS) and to perturbation by E2 stimulation such

as Mvalue and ΔDLR. Supplementary Fig. 8b showed a Pearson correlation heatmap followed by

hierarchical clustering between all-by-all features for DICs or all cohesin sites. The 175 features resulted

in clear clusters both among DICs and among all cohesin sites (dashed boxes of different colors). We

annotated the clusters as promoter, enhancer, enhancer-promoter interaction (E-P), insulator and

chromatin architecture. As compared with all cohesin sites, DICs showed lower co-binding tendency in

the promoter cluster and higher co-binding in the enhancer and E-P clusters. This showed the

effectiveness of our matrix in distinguishing DICs from other cohesin sites.

We then applied unsupervised k-means clustering (k = 10) to the matrix and obtained 10 clusters for

all cohesin sites (cluster 0–9, Fig. 4a, Supplementary Fig. 10), among which only cluster 4 and cluster 7

showed intragenic cohesin binding that decreased after E2 stimulation, indicating the DIC-like clusters

(Fig. 4b). We identified the following characteristics of cluster 4 (Fig. 4c, upper): 1) co-binding with

tissue-specific TFs (e.g., ER and FOXA1), 2) enrichment of enhancer markers and Pol2, 3) relatively low

intensity of cohesin and CTCF peaks and 4) chromatin de-compaction. Therefore, cluster 4 represented

the LC-DIC-like cluster. In contrast, cluster 7 (Fig. 4c, lower) showed the following characteristics: 1) lack

of TF co-binding, 2) high intensity of CTCF peaks and 3) highly related to topological boundaries and

chromatin architecture features (e.g., TAD borders, Hi-C loops). Therefore, cluster 7 represented the

HC-DIC-like cluster. Compared with "CNC-like" intragenic cohesin sites[5, 6] (clusters 2, 3 and 8;

Supplementary Fig. 10), cluster 4 (LC-DICs) co-localized only with enhancer markers and several master

regulators (FOXA1, ER and GATA3), and therefore it is distinct from typical *cis*-regulatory modules

(CRMs) at which many TFs co-localize. In contrast, cluster 7 (HC-DICs) consists of a cluster of intragenic

cohesin sites that tend to be localized to open chromatin, are highly de-compacted and contain loops but

are strongly negatively correlated with TFs. Therefore, they may be associated with a more universal

chromatin structure that is required for proper gene transcription.

To further explore the characteristics of DICs, we applied modeling of supervised ML (logistic

regression, support vector machine, and random forest) to predict LC- and HC-DICs from all cohesin

sites in a binary manner (labeled by 0 or 1). In this analysis, the input matrix consisted of 168 features

(features related to genomic location, cohesin changes and CTCF signal were excluded). We selected

chromosomes 16 to 22 for testing, and the remaining chromosomes were divided into training and

validation by five-fold cross-validation (see Methods). Because DICs are a small subset of all cohesin

sites, we used SMOTE over-sampling[28] to deal with such "imbalanced" classifications. The trained model

that was based on logistic regression achieved the best performance overall as compared with the others

(Supplementary Figs. 8c-d) and also performed adequately with the test data (Supplementary Fig. 8e).

Finally, we identified important features for the prediction of LC-DICs and HC-DICs by calculating the

relative feature importance from the trained model (Figs. 4d-e). LC-DICs were positively associated with

1) enhancer markers (H3K27ac, H3K4me1, P300); 2) Pol2 peaks, the Pol2-pausing regulator (LARP7)[29]

and a transcriptional repressor (ZBTB1)[30]; 3) tissue-specific regulators (FOXA1[26], HSF1[31], ER[18]). Both

LC-DICs and HC-DICs were positively associated with open chromatin (FAIRE-open, DNase-open) and

chromatin de-compaction (ΔDLR), and were negatively associated with H3K27me3 and CpG island

levels. HC-DICs, in particular, showed positive features of Hi-C loops but negative features of Pol2 loops

and TF binding, which is consistent with our analysis above, indicative of the TF-independent chromatin

de-compaction. Taken together, the application of machine learning successfully isolated special subsets

of cohesin sites (DICs) from all cohesin sites, which also provided us with additional characteristics by

which they can be identified.

Characterization of DIC tissue specificity

As DICs were enriched by many tissue-specific factors, we wondered whether our observations

about DICs were consistent with other tissues or cell types. We generated Rad21 ChIP-seq for 293T

cells (kidney), B-cells (lymphocytes), human skin fibroblast cells, RPE (retinal pigmented epithelium)

cells, and HeLa cells (cervical cancer). Cohesin peaks at MCF-7 derived LC-DICs were more specific in

MCF-7 cells, whereas cohesin peaks at HC-DICs were more ubiquitous across cell types (Figs. 5a-b).

We then asked whether HC-DICs play a role in tissue-specific transcription. Across cell types, the peak

intensities for Rad21 at HC-DICs were negatively correlated with transcription levels of their host genes

(Fig. 5c, Supplementary Fig. 9a), suggesting that genes with stronger HC-DIC binding had lower

transcription activities. Therefore, HC-DICs also showed the ability to impede transcription, which is

consistent with our motif analysis above.

To confirm whether DICs in other cell types also exhibit similar characteristics, we performed

ChIP-seq experiments on RPE cells with FBS (fetal bovine serum) and DRB (5,6-dichloro-1-β-D-

ribofuranosylbenzimidazole), which function as a stimulator and inhibitor of transcription[32], respectively.

First, we tested if the ML model trained by MCF-7 data was applicable to the RPE data. We used 25

features that were available for both MCF-7 and RPE cells to predict whether binding of an intragenic

cohesin was decreased or not after transcription stimulation (Fig. 5d). The predicted DICs overlapped

extensively with the experimentally determined ones (p < 10^{-161}, hypergeometric test), indicating that

DICs exhibited some common rules across cell types. Then we identified DICs of stimulation-responsive

genes in RPE cells (Supplementary Fig. 9b). Similar to DICs in MCF-7 cells, RPE derived DICs also

showed tissue-specific binding patterns (Supplementary Fig. 9c). FBS stimulation decreased the

intensity of cohesin (Rad21 and SA1) at DICs (position H1 to H2, L1 to L2 in Fig. 5e; Supplementary Fig.

9d), whereas transcriptional inhibition by DRB increased it. In addition, further treatment with DRB (i.e.,

FBS+DRB) reverted the decrease in cohesin binding caused by FBS stimulation (H2 to H5, L2 to L5 in

Fig. 5e; Supplementary Fig. 9d). Moreover, RNA Pol2 stalling and the release of paused Pol2 were also

observed at LC-DICs in RPE cells (Fig. 5e, Supplementary Fig. 9e). In addition, LC-DICs, but not

HC-DICs, co-bound with enhancer marks and several TFs. Thus, DICs are not a phenomenon

associated with only breast cancer cells but also are found in normal human cell lines.

Analysis of DICs in CdLS and CHOPS cells

Finally, we attempted to examine the participation of DICs in the observed phenotypes in individuals

with CdLS and CHOPS. To this end, we generated ChIP-seq data from patient-derived fibroblast cells,

together with the data that were generated in our previous study[17]. We overlapped the binding sites of

intragenic cohesin in different cell types (Fig. 5f). Whereas most sites were shared among samples, there

were 332 Rad21 sites that were absent in both CdLS and CHOPS cells, which we defined as DICs.

RNA-seq analysis showed that host genes of DICs (both LC- and HC-DICs) in both CdLS and CHOPS

cells showed a significant increase in transcription (Fig. 5g, Supplementary Fig. 9f; paired t-test),

suggesting that the decreased cohesin binding at DICs was correlated with expression upregulation in

both CdLS and CHOPS. We also assessed the binding of TFs at 185 LC-DICs and 147 HC-DICs (Fig. 5h,

Supplementary Fig. 9g). Interestingly, at LC-DICs, the peak intensity of AFF4 (causative gene of CHOPS)

increased in both CdLS and CHOPS cells, whereas lower binding of NIPBL (causative gene of CdLS)

was observed only in CdLS cells (Fig. 5h). Enhancer marker H3K27ac was highly enriched but did not

change among the three cell types, whereas Pol2 and Pol2ser5 (RNA pol II CTD phospho Ser5, which

represents paused Pol2) decreased in both CdLS and CHOPS cells, which is consistent with our

observations in MCF-7 and RPE cells. Taken together, this result suggests that DICs, especially

LC-DICs, are involved in abnormal transcription associated with both CdLS and CHOPS. As both CdLS

and CHOPs are involved in abnormal Pol2 regulation[33], LC-DICs might offer a common pathogenetic

mechanism. Based on the observations, we concluded that intragenic cohesin sites can be a good

candidate to investigate and link the phenotypes of these two cohesinopathy disorders.

Discussion

Cohesin is thought to be responsible for transcription regulation and chromatin folding. Several

models have been proposed to explain its functions. Cohesin can mediate enhancer-promoter loops with

the mediator complex or can function as a blocker between enhancer and promoter in conjunction with

insulator factor CTCF[34]. Cohesin also participates in the formation of chromatin topological structures via

the loop extrusion model[8]. A recent paper reported that transcription stimuli such as IFN-beta in THP-1

cells can displace cohesin from chromatin[10], which attracted our interest. Here, we focused on intragenic

cohesin, a subset of cohesin that was scarce, if not none, discussed in previous research. Of note, we

emphasized the effect of negative regulation of gene expression by cohesin-mediated chromatin loops,

whereas most of the previous studies implicitly assumed the positive regulation of expression by these

loops. DICs were negatively associated with activated transcription and chromatin compaction. Whereas

LC-DICs were highly enriched with enhancer markers and paused Pol2, HC-DICs were more involved in

the features of chromatin architectures. Importantly, DICs could be found in multiple cell types, especially

in CdLS and CHOPS cells, which partly supported the similarities between CdLS and CHOPS.

Chromatin interactions are required not only to facilitate transcription but also for Pol2 pausing[35]. By

using siRad21 cells, we confirmed that the release of Pol2 was related to the loss of cohesin at LC-DICs,

which supported our model that intragenic loops formed by cohesin paused Pol2 and that transcription

elongation from TSSs could remove such cohesin and then release the paused Pol2 (Fig 5i). Notably, we

did not observe significant changes in the expression of genes that host LC-DICs by siRad21

(Supplementary Fig. 6f). As most LC-DICs are located in intronic regions, Pol2 released from LC-DICs

might be involved in accurate RNA splicing. Notably, a recent study suggested that intragenic enhancers,

in addition to activating genes, also attenuate the transcription of their host genes during productive

elongation[12], which evokes the functional link between LC-DICs and Pol2 pausing. In contrast, HC-DICs

showed a high preference for loop occurrence mediated by CTCF, possibly to play a role in topological

boundaries (e.g., sub-TADs). Across different cell types and genes, the Rad21 signal at HC-DICs was

negatively correlated with the expression of host genes, indicating the role of HC-DICs in restraining

transcription. Whereas we observed that more than half of HC-DIC–mediated loops anchored intronic

regions of two genes, it was difficult to infer the biological meaning of this because HC-DICs scarcely

overlapped with any other TFs. Further biological approaches such as genome editing of HC-DICs of

activated genes could be promising in the future.

Modifications at intragenic regions affect transcription events. For instance, intragenic methylation

can prevent spurious transcription initiation[11]; Intragenic microRNAs affect the expression of their host

genes[13]. Here we present the first study that focused on intragenic cohesin sites. We also used penalty

linear regression followed by univariate linear regression to better understanding the changes of cohesin

bindings in intragenic regions (see Method and Supplementary Fig. 8f-h). Apart from the decreased ones,

the increased intragenic cohesin sites seemed to be also correlated with many important features, as

they are positively predicted by ER and several TFs. Although we characterized intragenic cohesin sites

that showed decreased bindings in this study, all types of intragenic cohesin might have a role in

transcription regulation. In addition, Kowalczyk et al. point out that intragenic enhancers can act as

alternative promoters[36]. Our DICs did not overlap with any known alternative promoters. Even though we

know little about the molecular mechanism, our results strongly suggest a previously undescribed

function of cohesin in intragenic regions with respect to gene expression regulation.

In summary, large-scale multi-omics enabled us to identify a cluster of cohesin DICs in MCF-7 and

other cell types. Some tissue-specific DICs (LC-DICs) were related to enhancers and Pol2

pausing/release, whereas others (HC-DICs) contributed to chromatin architecture and might attenuate

transcription. Our integrated analysis and machine learning approaches indicated distinct characteristics

that distinguish DICs from other cohesin binding sites. Based on these genomic, epigenomic and

transcriptomic characteristics, we can infer that DICs have distinct functions as compared with other

cohesin sites.

Methods

Cell culture and treatment

RPE cells[37] and MCF-7 cells (JCRB Cell Bank) were cultured in DMEM containing 10% FBS and 1%

penicillin/streptomycin. Before subsequent treatments, RPE cells were cultured in serum-free medium for

48 hours and then were incubated in DMEM containing 10% FBS for 30 min. MCF-7 cells were

maintained in phenol red–free medium containing charcoal-dextran–stripped FBS (Life Technologies) at

70–80% confluency for 2 days before treatment with 50 nM E2 (SIGMA) for the indicated times. Rad21

siRNA oligos[4] were transfected using Lipofectamine RNAiMax (Life Technologies) according to the

manufacturer's instructions at 40 hours before treatment with E2. DRB (TCI) was added at 1.5 hours

before treatment with E2. The effect of cohesin (Rad21)-deficiency were verified by western blot as

shown in Supplementary Fig. 6e.

ChIP and antibodies

Cells were fixed in medium or phosphate buffered saline with 1% formaldehyde at room

temperature for 10 min. ChIP experiments were performed as described[38]. ChIP-seq libraries were

prepared using NEBNext ChIP-seq Library Prep Master Mix Set for Illumina (New England BioLabs).

Rabbit polyclonal antibody for Rad21 has been described in[39]. Antibody for Pol2 (#14958) was from Cell

Signaling Technology. Antibodies for Mau2 (ab46906) and SA1 (ab4457) were from Abcam. Antibodies

for TAF (A303-505A) and AFF4 (A302-538A) were from Bethyl Laboratory. CTCF (07-729) antibody was

from Merck Millipore. Antibodies for Pol2ser2 and H3K27ac were kindly provided by Dr. H Kimura

(TITech). Antibodies for P300 (sc-585) and Med1 (sc-5334) were from Santa Cruz Biotechnology.

Antibody for CBP (606402) was from BioLegend.

ChIP-seq analysis

After quality check by FastQC and SSP[40], ChIP-seq reads were aligned to the human reference

genome (hg38) using Bowtie[41] version 1.2.2 with "-n2 -m1" parameters, by which we considered only

uniquely mapped reads and allowed two mismatches in the first 28 bases per read. Peak calling was

performed using MACS2[42] version 2.2.6 with default settings. We used DROMPA[43] version 3.7.2 to

conduct statistical analysis and visualization. For visualization of ChIP-seq binding to particular

chromatin regions, reads were normalized relative to total read number, and gene annotation was

obtained from NCBI reference sequences (RefSeq; hg38). Read profiles around the sites of interest were

plotted with the PROFILE mode of DROMPA, whereas the heatmap of target sites (2.5 kb around the

peak summit) was plotted using HEATMAP mode. Genomic distribution in Fig. 2e was plotted by

ChIPseeker[44]. Downstream analysis, such as peak overlap, was performed by Bedtools[45] version 2.29.2

and Samtools[46] version 1.9. Sources for all ChIP-seq data and other next-generation sequencing data

(including our data and public data) are listed in Supplementary Tables S2-3.

Hi-C analysis

All in-situ Hi-C data (control or E2-treated MCF-7 cells with two replicates) were aligned to the hg38

human reference genome. Further analysis was carried out mainly by Juicer[22] version 1.11.04. All

contact matrices were normalized by the KR method in Juicer. Chromatin loops were annotated using the

HiCCUPS algorithm with default parameters[22]. The loop regions we used were merged from the results

of 5-kb,10-kb and 25-kb resolutions. Aggregate peak analysis (APA) was performed using the 'apa'

mode of Juicer (5-kb resolution), to measure the enrichment of the Hi-C signal around a set of peaks.

The visualization of the contact matrix on the *MREG* locus was accomplished by Matplolib. After

correction and normalization, comparable contact matrices were plotted at a 5-kb resolution. We merged

two adjacent bins for smoothing. Other Hi-C analyses were performed using HOMER[10]. We made the

Tag directory with the "GATC" restriction site sequence. Chromatin compaction scores ΔDLR and ΔICF

were calculated for each 5-kb region across the genome (-res 5000) from a 15-kb window size (-window

15000). Other metrics including PC1, insulation score and TAD boundaries were obtained using HOMER

with default parameters. We used the WashU epigenome browser[47] to visualize Supplementary Fig. 3b.

ChIA-PET analysis

RNA polymerase II−bound chromatin interactions in MCF-7 cells were extracted from ChIA-PET

data (GSE33664). All fastq files were applied to the published pipeline Mango[48] with default parameters,

based on the hg38 reference genome. ChIA-PET interactions were visualized by DROMPA with the

parameter '-inter'.

RNA-seq and GRO-seq analysis

Using HISAT2[49] version 2.2.0, we aligned paired-end RNA-seq reads to the index established from

the hg38 reference genome. The output SAM files were converted to BAM files by Samtools. Htseq[50]

version 0.11.3 was then used with default parameters to generate a count table, which describes the

number of reads on each gene. We used a GTF file (GRCh38.p12) from GENCODE for gene annotation.

Subsequent differential expression analysis was achieved using DESeq2[51], with its internal normalization.

For GRO-seq, alignment was carried out using Bowtie with "-n2 -m1" parameters. The output was

preprocessed and visualized using DROMPA.

Data collection and machine learning

All datasets used in machine learning are listed in Table S5. Apart from our data, public omics data

in wild-type MCF-7 cells were downloaded mainly from the GEO database, ENA database, ENCODE

project, FANTOM5 project, UCSC genome browser and GWAS Catalog database. These data were then

overlapped with all 184,140 cohesin peaks. As a result, we obtained 15 continuous features and 160

binary features, the latter of which indicated whether a kind of data was co-localized (1) or not

co-localized (0) at a cohesin site. After normalization of continuous features, the big matrix, which

consisted of 184,140 rows (cohesin sites) and 175 features, was imported into for different analyses. The

features correlation heatmap for all cohesin sites and DICs was made with the R package *corrplot*. We

used scikit-learn version 0.22.1 to perform machine learning. Overall, the parameters used in scikit-learn

were optimized by grid search with 5-fold cross-validation. For unsupervised learning (k-means), all 175

features were used to fit models. For supervised learning (logistic regression, support vector machine,

random forest), we omitted Mvalue, cohesin location and CTCF signal information and then used the

remaining 168 features as independent variables $X_i = (X_{i1}, X_{i2}, ..., X_{ij})$, for $i = 1, 2, ..., 184140$ and $j =$

$1, 2, ..., 168$. Based on whether they were DICs or not, we labeled each cohesin site as 1 or 0 and then

used it as a dependent variable $Y_i \in \{0, 1\}$. The conditional probability of logistic regression was

calculated as follows:

510
$$P(Y_i = 1 | X = X_i) = \frac{1}{1 + e^{-(\beta_0 + \sum_{j=1}^{168} \beta_j X_{ij})}}$$

where β_j is the regression coefficient of each feature. We used training data to do model fitting and

used test data to validate model performance. To apply the MCF-7−derived ML model to RPE cells, we

used 25 features that were available in both MCF-7 and RPE cells. We used logistic regression with L1

penalty to decide whether each intragenic cohesin site had decreased binding (1) or not (0). Then the

MCF-7−fitted model was applied to the RPE features to predict DICs.

We also applied penalized regression followed by univariate linear regression as described[52], to

reveal which features contributed to negative or positive Mvalue (log ratio of cohesin peak intensity

between E2 and control) in intragenic cohesin (26066 sites). We used 169 features (of the 175 features,

6 were excluded: 5 features related to cohesin position and the Mvalue feature) as independent variables

$X_i = (X_{i1}, X_{i2}, ..., X_{ij})$, for $i = 1, 2, ..., 26066$ and $j = 1, 2, ..., 169$, whereas the Mvalue was the dependent

variable Y_i. Instead of using the ordinary least squares approach, we used the elastic net loss function:

522
$$L_{enet}(\hat{\beta}) = \frac{\sum_{i=1}^{n}(y_i - x_i'\hat{\beta})^2}{2n} + \lambda(\frac{1-\alpha}{2}\sum_{j=1}^{169}\hat{\beta_j}^2 + \alpha\sum_{j=1}^{169}|\hat{\beta_j}|)$$

to the linear model $Y_i = \beta_0 + \sum_{j=1}^{169} \beta_j X_{ij}$, where $n = 26066$ and $\hat{\beta}$ was the estimation of β. λ was

chosen by cross-validation, and $\alpha = 0.5$ was used to consider both the L1 and L2 penalty. Feature

selection with such regularization was useful for filtering out non-significant or redundant features. The

remaining features were applied to univariate linear regression $Y = a + bX$ to calculate the regression

coefficient (Supplementary Figs. 8f–h).

Extraction of DICs

Quantitative comparison of Rad21 binding events was performed using MAnorm[19] version 1.3.0

with default parameters. This results in the normalized Mvalue, a quantitative measure of differential

binding for all cohesin sites. To acquire more convincing sites, we combined our results with high-quality

ChIP-seq data from E-TABM-828[5]. Criteria for selecting decreased peaks were peak width < 3 kb and

Mvalue < −0.5. Next, we used RefSeq as the reference genome to obtain intragenic regions. As

described in Supplementary Fig. 1e, we excluded 10kb flanking regions around TSS and TES. Only large

genes (gene length > 20 kb) were considered. E2-responsive genes were defined as genes with an

increased Pol2ser2 ChIP-seq signal (ratio > 1.2) in the presence of E2 relative to control and that were

validated by RNA-seq data. Decreased peaks at intragenic region of 499 E2-responsive gene were

defined as DICs. Next, to quantify CTCF read density on DICs, we used MULTICI options in DROMPA

software. Peaks with extremely low Rad21 signals were omitted. Finally, we separated the DICs into 141

high-CTCF DICs (HC-DICs) and 417 low-CTCF DICs (LC-DICs).

Motif analysis

Motifs were analyzed using HOMER. Briefly, peak files in standard bed format were converted to

HOMER peak files, and then the command *findMotifsGenome.pl* was used to discover the motif. The

results included known motifs as well as *de novo* discovered motifs. The size of the region used for motif

finding was set to 200 bp. The top 10 motifs with the lowest *q*-values (Benjamini-Hochberg) are shown.

Software environment

All analyses were based on Ubuntu 18.04.4 with Python 3.6.9 and R 3.6.3. Data were processed

using R base package or Numpy (v 1.17.2) as well as Pandas (v 0.25.1) in Python. Figures were drawn

with DROMPA (v 3.7.2), Matplotlib (v 3.1.1), ggplot2 and R base plotting.

Abbreviations:

ChIP-seq: chromatin immunoprecipitation followed by sequencing; CNCs: cohesin-non-CTCF sites; RNA-seq: RNA sequencing; Hi-C: high-throughput chromosome conformation capture; ChIA-PET: chromatin interaction analysis by paired-end tag sequencing; DIC: decreased intragenic cohesin; LC-DIC: low-CTCF decreased intragenic cohesin; HC-DIC: high-CTCF decreased intragenic cohesin. Pol2: RNA polymerase II; Pol2ser2: RNA pol II CTD serine-2 phosphorylation; TSS: transcription start site; TES: transcription end site; TAD: topologically associating domain.

Competing interests:

The authors declare that they have no competing interest.

Author contributions

J.W. performed all bioinformatics analysis. R.N. conceived this project. J.W. and R.N. drafted the manuscript. M.B. prepared ChIP-seq and RNA-seq samples. K.S. supervised the sample preparation and sequencing, suggested ways to improve the analysis, and improved the manuscript.

Data availability

The raw sequencing data and processed files are available at the GEO

References:

1. Kim Y, Shi Z, Zhang H, Finkelstein IJ, Yu H. Human cohesin compacts DNA by loop extrusion. *Science* **366**, 1345-1349 (2019).

2. Nishiyama T. Cohesion and cohesin-dependent chromatin organization. *Current opinion in cell biology* **58**, 8-14 (2019).

3. Bloom MS, Koshland D, Guacci V. Cohesin Function in Cohesion, Condensation, and DNA Repair Is Regulated by Wpl1p via a Common Mechanism in Saccharomyces cerevisiae. *Genetics* **208**, 111-124 (2018).

4. Wendt KS*, et al.* Cohesin mediates transcriptional insulation by CCCTC-binding factor. *Nature* **451**, 796-801 (2008).

5. Schmidt D*, et al.* A CTCF-independent role for cohesin in tissue-specific transcription. *Genome research* **20**, 578-588 (2010).

6. Faure AJ*, et al.* Cohesin regulates tissue-specific expression by stabilizing highly occupied cis-regulatory

587   modules. *Genome research* **22**, 2163-2175 (2012).

588   7.   Kagey MH*, et al.* Mediator and cohesin connect gene expression and chromatin architecture. *Nature* **467**,

589   430-435 (2010).

590   8.   Fudenberg G, Imakaev M, Lu C, Goloborodko A, Abdennur N, Mirny LA. Formation of Chromosomal

591   Domains by Loop Extrusion. *Cell reports* **15**, 2038-2049 (2016).

592   9.   Dixon JR, Gorkin DU, Ren B. Chromatin Domains: The Unit of Chromosome Organization. *Molecular cell* **62**,

593   668-680 (2016).

594   10.   Heinz S*, et al.* Transcription Elongation Can Affect Genome 3D Structure. *Cell* **174**, 1522-1536 e1522

595   (2018).

596   11.   Neri F*, et al.* Intragenic DNA methylation prevents spurious transcription initiation. *Nature* **543**, 72-77 (2017).

597   12.   Cinghu S*, et al.* Intragenic Enhancers Attenuate Host Gene Expression. *Molecular cell* **68**, 104-117 e106

598   (2017).

599   13.   Hinske LC, Galante PA, Kuo WP, Ohno-Machado L. A potential role for intragenic miRNAs on their hosts'

600   interactome. *BMC genomics* **11**, 533 (2010).

601   14.   Krantz ID*, et al.* Cornelia de Lange syndrome is caused by mutations in NIPBL, the human homolog of

602   Drosophila melanogaster Nipped-B. *Nature genetics* **36**, 631-635 (2004).

603   15.   van der Lelij P*, et al.* Synthetic lethality between the cohesin subunits STAG1 and STAG2 in diverse cancer

604   contexts. *eLife* **6**,   (2017).

605   16.   Kon A*, et al.* Recurrent mutations in multiple components of the cohesin complex in myeloid neoplasms.

606   *Nature genetics* **45**, 1232-1237 (2013).

607   17.   Izumi K*, et al.* Germline gain-of-function mutations in AFF4 cause a developmental syndrome functionally

608   linking the super elongation complex and cohesin. *Nature genetics* **47**, 338-344 (2015).

609   18.   Theodorou V, Stark R, Menon S, Carroll JS. GATA3 acts upstream of FOXA1 in mediating ESR1 binding by

610   shaping enhancer accessibility. *Genome research* **23**, 12-22 (2013).

611   19.   Shao Z, Zhang Y, Yuan GC, Orkin SH, Waxman DJ. MAnorm: a robust model for quantitative comparison of

612   ChIP-Seq data sets. *Genome biology* **13**, R16 (2012).

613   20.   Bowman EA, Kelly WG. RNA polymerase II transcription elongation and Pol II CTD Ser2 phosphorylation: A

614   tail of two kinases. *Nucleus* **5**, 224-236 (2014).

615   21.   Li X*, et al.* A unified mechanism for intron and exon definition and back-splicing. *Nature* **573**, 375-380

616   (2019).

617   22.   Durand NC*, et al.* Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments.

618   *Cell Syst* **3**, 95-98 (2016).

23.   Ren G*, et al.* CTCF-Mediated Enhancer-Promoter Interaction Is a Critical Regulator of Cell-to-Cell Variation

of Gene Expression. *Molecular cell* **67**, 1049-1058 e1046 (2017).

24.   Lalmansingh AS, Karmakar S, Jin Y, Nagaich AK. Multiple modes of chromatin remodeling by Forkhead box

proteins. *Biochimica et biophysica acta* **1819**, 707-715 (2012).

25.   Fournier M*, et al.* FOXA and master transcription factors recruit Mediator and Cohesin to the core

transcriptional regulatory circuitry of cancer cells. *Scientific reports* **6**, 34962 (2016).

26.   Hurtado A, Holmes KA, Ross-Innes CS, Schmidt D, Carroll JS. FOXA1 is a key determinant of estrogen

receptor function and endocrine response. *Nature genetics* **43**, 27-33 (2011).

27.   Andersson R*, et al.* An atlas of active enhancers across human cell types and tissues. *Nature* **507**, 455-461

(2014).

28.   Chawla NV, Bowyer KW, Hall LO, Kegelmeyer WP. SMOTE: synthetic minority over-sampling technique.

Journal of artificial intelligence research **16**, 321-357 (2002).

29.   Frilander MJ, Barboric M. The Interlocking Lives of LARP7: Fine-Tuning Transcription, RNA Modification,

and Splicing through Multiple Non-coding RNAs. *Molecular cell* **78**, 5-8 (2020).

30.   Matic I*, et al.* Site-specific identification of SUMO-2 targets in cells reveals an inverted SUMOylation motif

and a hydrophobic cluster SUMOylation motif. *Molecular cell* **39**, 641-652 (2010).

31.   Santagata S*, et al.* High levels of nuclear heat-shock factor 1 (HSF1) are associated with poor prognosis in

breast cancer. *Proceedings of the National Academy of Sciences of the United States of America* **108**,

18378-18383 (2011).

32.   Busslinger GA*, et al.* Cohesin is positioned in mammalian genomes by transcription, CTCF and Wapl.

Nature **544**, 503-507 (2017).

33.   Piche J, Van Vliet PP, Puceat M, Andelfinger G. The expanding phenotypes of cohesinopathies: one ring to

rule them all! *Cell Cycle* **18**, 2828-2848 (2019).

34.   Merkenschlager M, Nora EP. CTCF and Cohesin in Genome Folding and Transcriptional Gene Regulation.

Annual review of genomics and human genetics **17**, 17-43 (2016).

35.   Ong CT, Corces VG. Enhancer function: new insights into the regulation of tissue-specific gene expression.

Nature reviews Genetics **12**, 283-293 (2011).

36.   Kowalczyk MS*, et al.* Intragenic enhancers act as alternative promoters. *Molecular cell* **45**, 447-458 (2012).

37.   Gallego-Paez LM*, et al.* Smc5/6-mediated regulation of replication progression contributes to chromosome

assembly during mitosis in human cells. *Molecular biology of the cell* **25**, 302-317 (2014).

38.   Komata M, Katou Y, Tanaka H, Nakato R, Shirahige K, Bando M. Chromatin immunoprecipitation protocol

for mammalian cells. *Methods in molecular biology* **1164**, 33-38 (2014).

39.   Minamino M*, et al.* Esco1 Acetylates Cohesin via a Mechanism Different from That of Esco2. *Current*

biology : CB **25**, 1694-1706 (2015).

40. Nakato R, Shirahige K. Sensitive and robust assessment of ChIP-seq read distribution using a strand-shift
profile. *Bioinformatics* **34**, 2356-2363 (2018).

41. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of short DNA
sequences to the human genome. *Genome biology* **10**, R25 (2009).

42. Zhang Y*, et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137 (2008).

43. Nakato R, Itoh T, Shirahige K. DROMPA: easy-to-handle peak calling and visualization software for the
computational analysis and validation of ChIP-seq data. *Genes Cells* **18**, 589-601 (2013).

44. Yu G, Wang LG, He QY. ChIPseeker: an R/Bioconductor package for ChIP peak annotation, comparison
and visualization. *Bioinformatics* **31**, 2382-2383 (2015).

45. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**,
841-842 (2010).

46. Li H*, et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079 (2009).

47. Zhou X*, et al.* Exploring long-range genome interactions using the WashU Epigenome Browser. *Nature*
methods **10**, 375-376 (2013).

48. Phanstiel DH, Boyle AP, Heidari N, Snyder MP. Mango: a bias-correcting ChIA-PET analysis pipeline.
Bioinformatics **31**, 3092-3098 (2015).

49. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and genotyping with
HISAT2 and HISAT-genotype. *Nature biotechnology* **37**, 907-915 (2019).

50. Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-throughput sequencing data.
Bioinformatics **31**, 166-169 (2015).

51. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with
DESeq2. *Genome biology* **15**, 550 (2014).

52. Hu S*, et al.* ncHMR detector: a computational framework to systematically reveal non-classical functions of
histone modification regulators. *Genome biology* **21**, 48 (2020).

Figure 1

Fig. 1 Classification of decreased intragenic cohesin (DIC) sites. a. Quantitative comparison of
cohesin peak intensity between control (Ctrl) and estrogen (E2)-treated MCF-7 cells. Mvalue is the
$\log_2($ fold change) of normalized read densities under comparison. Avalue is the average signal strength
of each peak. The lower panel shows the proportion of cohesin binding that decreased or increased. **b.**
Average binding profiles on different cohesin sites. Red, control. Blue, E2 treatment. **c.** Genomic
distribution of all DICs suggested most of DICs were located in introns. **d.** Aggregate peak analysis
around DICs or all cohesin peaks at a 5-kb resolution. **e.** Chromatin compaction scores around DICs, all
cohesin sites and all enhancers (summit ± 500 kb). ΔDLR (E2 vs. control) was calculated at a 25-kb
resolution. **f.** Heatmap of ChIP-seq reads at DICs (peak summit ± 2.5 kb). CTCF (E2+) signal was used
for sorting the order. DICs were divided into HC-DICs (gray bar) and LC-DICs (black bar). E2−, control;

E2+, estrogen treated. **g.** Average binding profiles (summit ± 1.5 kb) of Rad21 and TFs on HC-DICs

(upper) and LC-DICs (lower). Reads were normalized relative to the whole genome. The same *y*-axis

scale was used for each protein. **h.** Read distribution of Rad21 and TFs around *MREG* and *PAK4* loci.

Areas shaded in pink and light yellow indicate LC-DICs and HC-DICs, respectively. All reads were

normalized relative to the whole genome. **i.** Hi-C contact map (5-kb resolution) around *MREG*. Arrows

show the disappearance of chromatin interactions after E2 stimulation. The loop anchor is shown as a

black bar in 1h.

Figure 2

Fig. 2 Enhancers and loops on DICs. a. The proportion of cohesin overlap with CBP peaks. Fisher's exact test was used between LC-DICs and other cohesin sites. **p < 10^{-5}, ***p < 10^{-10}. **b.** Genomic binding of multiple histone markers. Enhancer markers H3K27ac and H3K4me1 (black dashed box) were enriched at LC-DICs. Gray dashed boxes indicate histone markers for promoters. Red arcs show Pol2-mediated loops. **c.** Loop occurrence probabilities on DICs, calculated based on Pol2 ChIA-PET or Hi-C data. We used Fisher's exact test to do statistical comparisons. *p < 0.05, **p < 0.01, ***p < 0.001. **d.** The percentage of HiC loops (LC-DICs and HC-DICs) that interacted with cohesin, enhancers, promoters or the promoter of their host gene. **e.** Genomic distribution of the other end of DIC-loops as generated by ChIPseeker. **f.** Insulation scores (ISs) for each 25-kb genomic region around DICs or all cohesin sites (summit ± 500 kb). DICs were further divided (dashed line) into HC-DICs and LC-DICs.

Figure 3

Fig. 3 Pol2 pauses on LC-DICs. a. Genomic binding of Pol2 and Pol2ser2 and GRO-seq data for

MREG and *PAK4* loci. **b.** Binding profiles of Pol2 and Pol2ser2 on LC-DICs or HC-DICs, under control

and E2 conditions. Reads were normalized relative to the whole genome. **c.** The $\log_{10}$ value of the ratio

E2/Ctrl with respect to the Pol2 signal on various cohesin sites; *p*-values (Mann–Whitney *U*-test) were

calculated between LC-DICs and other cohesin sites. Up-IC, up-regulated intragenic cohesin; stable-IC,

unchanged intragenic cohesin; TSS_{E2res}, TSS of E2-responsive gene. **d.** Visualization of Pol2 and

Pol2ser2 ChIP-seq data around *MREG* and *PAK4*. MCF-7 cells were treated as indicated. WT, wild type;

KD, Rad21 knockdown; Ctrl, control; E2, estrogen. **e.** Pol2 and Pol2ser2 binding profiles around LC-DICs

(summit ± 1.5 kb) under four different conditions. Reads were normalized relative to the whole genome. **f.**

Quantitative comparison (Mann–Whitney *U*-test) of Pol2 and Pol2ser2 signals on LC-DICs or on TSSs of

E2-responsive genes (TSS_E2res) under four different conditions. **g.** ChIP-seq results for CBP, P300

744 and Mau2 on *PAK4* gene in WT and Rad21 knockdown MCF-7 cells with the absence and presence of

745 E2 stimulation. Dashed arrows show increased binding events in cohesin-deficient cells.

Figure 4

Fig. 4 Machine learning methods to classify DICs. a. Cluster heatmap by k-means (k = 10) methods.
The 175 features were roughly classified into different groups. **b.** Magnified view of the heatmap area
shown in A indicates three important features for these 10 clusters: "aroundTSS" indicates whether a
cohesin site was near TSS; "intragenic" indicates whether a cohesin site was within a gene body;
"Mvalue" is the $\log_2$ ratio of Rad21 peak density between the E2 and control group. **c.** The mean of each
feature for cluster 4 and 7. The *y* axis shows the normalized (*z*-score) values of binary and continuous
features. Error bars represent 95% confidence intervals. **d-e.** The importance of individual features from
the logistic regression model for (**d**) LC-DICs and (**e**) HC-DICs. Top positive and negative features are
listed inside the plots. Model performance is indicated in the bottom right.

Figure 5

Fig. 5 DICs in other cell types. a. Rad21 ChIP-seq of the *MREG* locus in various cell lines (MCF-7, 293T, B-cells, Fibroblasts, RPE, HeLa). HC-DICs were observed in other cells but not LC-DICs. **b.** The percentage of MCF-7 DICs that could be found in other cell types. **c.** Relationships between peak density of HC-DICs and the expression of their host genes. Gene expression data for the different cell types were obtained from the GTEx database. **d.** Overlap between DICs that were predicted relative to those that were experimentally isolated; *p*-values were calculated with the hypergeometric test. **e.** ChIP-seq data of the example locus in RPE cells. L1-5 indicated LC-DIC sites and H1-5 indicated HCDIC sites. **f.** Overlap of intragenic cohesin sites in normal and CdLS- and CHOPs-derived fibroblast cells. **g.** Expression of genes associated with DICs (paired t-test) in Normal, CdLS and CHOPs cells. **h.** Binding of Rad21, NIPBL, AFF4, H3K27ac, Pol2 and Pol2ser5 on LC-DICs. Dashed lines represent the average binding signal of each TF on all cohesin sites. Wilcoxon signed-rank test was used to compute *p*-values. **i.**

809   Proposed model for our DICs. Briefly, intragenic loops formed by LC-DICs pause Pol2, whereas

810   HC-DICs form CTCF-mediated loops. Transcription stimulation could displace DICs and disrupt DIC

811   loops, consequently releasing the Pol2 pause. The exposed enhancer sites on LC-DICs then bind their

812   respective TFs and interact with promoters.