Trypanosome morphogenesis involves recruitment of a pleckstrin homology domain protein by an orphan kinesin to the microtubule quartet

Corinna Benz[1], Nora Müller[2], Sabine Kaltenbrunner[1,2,†], Hana Váchová[3], Marie Vancová[1,2], Julius Lukeš[1,2], Vladimír Varga[3], Hassan Hashimi[1,2,*]

[1]Institute of Parasitology, Biology Center, Czech Academy of Sciences, České Budějovice, Czechia

[2] Faculty of Science, University of South Bohemia, České Budějovice, Czechia

[3]Laboratory of Cell Motility, Institute of Molecular Genetics of the Czech Academy of Sciences, Prague, Czechia

[†]Current address: Johannes Kepler University, Medical Faculty, Linz, Austria

*Corresponding author: hassan@paru.cas.cz

Running title: TbKifX2 and TbPH1 association with the trypanosomal microtubule quartet

ABSTRACT

Kinesins are motor proteins found in all eukaryotic lineages that move along microtubule tracks to mediate numerous cellular processes such as mitosis and intracellular transport of cargo. In trypanosomatids, the kinesin superfamily has undergone a prominent expansion, giving these protists one of the most diverse kinesin repertoires including two trypanosomatid-restricted groups. Here, we characterize in *Trypanosoma brucei* TbKifX2, a hitherto orphaned trypanosomatid-specific kinesin. TbKifX2 tightly interacts with TbPH1, a kinesin-like protein with an inactive motor domain, a rare instance of a kinesin interacting with a protein bearing a catalytically dead kinesin motor domain. TbKifX2 recruits TbPH1 to the microtubule quartet (MtQ), a characteristic but poorly understood cytoskeletal structure that is part of the multipartite flagellum attachment zone (FAZ) and extends from the basal body to the anterior of the cell body. The proximal proteome of TbPH1 is comprised of four proteins that localize to the FAZ, consistent with the notion that the TbKifX2/TbPH1 complex bind the MtQ, a FAZ component. Simultaneous ablation of TbKifX2/TbPH1 leads to formation of prominent protrusions from the cell posterior. Thus, these two trypanosomatid-restricted proteins appear to contribute to morphogenesis in *T. brucei*. Furthermore, TbKifX2 is the first motor protein to be localized to the MtQ.

Keywords: Trypanosoma, kinesin, cytoskeleton, microtubules, microtubule quartet, morphogenesis

INTRODUCTION

The vermiform morphology of the unicellular parasite *Trypanosoma brucei* that causes African
trypanosomiasis in humans and nagana in cattle is defined by a subpellicular array of microtubules
(MTs) (Robinson *et al.*, 1995; Gull, 1999; Wheeler *et al.*, 2013). These MTs are cross-linked to each
other and the plasma membrane via regularly spaced fibrils (Sherwin *et al.*, 1987). A high order is
maintained within the corset with the MT plus ends converging at the parasite's posterior end and
the minus ends located at the anterior of the cell (Robinson *et al.*, 1995).

The only place where the continuity of the *T. brucei* MT array is interrupted is at the flagellar pocket,
the deep invagination where the flagellum exits the cell body and the only place where endo- and
exocytoses occur (Field and Carrington, 2009). The opening of the flagellar pocket is encircled by a
collar, which influences the overall shape of this cavity. Distal to the collar, the so-called hook
complex covers the flagellum as it exits the flagellar pocket, with two lateral arms also flanking the
flagellum attachment zone (FAZ) filament and the MT quartet (MtQ) inside the cell (Albisetti *et al.*,
2017). The FAZ filament and MtQ are among the interconnected fibres, filaments and junctional
complexes of the multipartite FAZ (Lacomble *et al.*, 2009), which together form a lateral attachment
for the flagellum as it follows a helical path along the cell body, connecting the flagellar skeleton to
the cytoskeleton through both the flagellar and plasma membranes (Sunter and Gull, 2016). The
MtQ originates close to the basal bodies, wraps itself around the flagellar pocket and interdigitates
within the MT corset next to the FAZ filament. The orientation of MTs in the MtQ is antiparallel to
the one found within the MT corset, thus their minus ends are proximal to the basal bodies
(Robinson *et al.*, 1995).

The trypanosome MT corset is exceptionally stable and only reorganised during cell division and life
cycle transitions (Sinclair and de Graffenried, 2019). Cytoskeletal structures (MtQ, MT corset,
flagellum, basal bodies, which anchor the flagella within the cell, and FAZ) need to be duplicated and
faithfully segregated during the cell division cycle, just like other single-copy organelles. In addition,
this has to be coordinated with replication and division of the nucleus and the single mitochondrial
DNA network termed the kinetoplast, which is located on the opposite side of the basal bodies
relative to the flagellum (Robinson *et al.*, 1995; Wheeler *et al.*, 2019).

Basal body duplication and segregation are the first visible steps in the cell division cycle and
essential for subsequent cytokinesis (Woodward and Gull, 1990). The new MtQ starts growing just
prior to basal body maturation and once the new flagellum has grown enough to invade the flagellar
pocket, morphogenetic changes result in two separate pockets enveloping both the new and old
flagella (Lacomble *et al.*, 2010). Once the new flagellum has been nucleated and elongated, new MTs
are integrated into the array between the old and new flagella, thereby increasing the overall width
of the cell (Wheeler *et al.*, 2013).

With its antiparallel orientation, the MtQ together with the FAZ filament create an asymmetric seam
within the MT corset, which works as a 'cellular ruler' and helps to determine the site of future
furrow ingression during cytokinesis (Sunter and Gull, 2016). Any insults to the integrity of these
structures result in cytokinesis defects and it is thus tempting to envision the MtQ as a potential
specialised 'highway' in charge of transport and delivery of cargo to the site of important cytokinesis
events. Furthermore, the endplasmic reticulum associates with parts of the MtQ, although the
importance of this association remains mysterious. The first protein localized to this structure,
TbSpef1, is found at the proximal end of the MtQ, which wraps around the flagellar pocket between
basal bodies and the flagellar pocket collar (Gheiratmand *et al.*, 2013). Its depletion causes problems
with the biogenesis of the MtQ and associated structures as well as motility defects. The minus end

of the MtQ MTs is anchored to the basal bodies by a protein named TbSAF1, found in close proximity
to TbSpef1 but localized between the pro and mature basal bodies (Dong *et al.*, 2020). TbSAF1 was
found in a proximal proteomics screen of proteins adjoining to and/or directly interacting with
TbSpef1, which found three additional proteins that localize to the MtQ between the basal body and
flagellar pocket collar (Dong *et al.*, 2020). However, no motor proteins that localise to the MtQ, and
which may potentially bring cargo to the cytokinesis site, have thus far been identified.

Kinesins are well-studied motor proteins that regulate MT dynamics, organize MT networks, and
transport cargo along MTs. Transporting kinesins can form homo- and heterodimers and consist of a
head region containing the kinesin motor domain, which is responsible for ATP hydrolysis and MT
binding, and a tail region important for dimerization with a kinesin/kinesin-like partner and cargo
binding (Hirokawa and Noda, 2008). In trypanosomatid genomes, kinesins and kinesin-like proteins
constitute one of the largest protein superfamilies (Berriman *et al.*, 2005; El-Sayed *et al.*, 2005; Ivens
et al., 2005). To date, 47 genes encoding high-likelihood kinesins have been identified in the *T. brucei*
genome (Wickstead and Gull, 2006; Wickstead *et al.*, 2010b). Moreover, there is a similar number of
kinesin-related proteins, which did not make the threshold due to sequence variance (Berriman *et
al.*, 2005). Interestingly, numerous kinesins are members of two major clades named X-1 and X-2
that are restricted to the trypanosomatid lineage (Wickstead and Gull, 2006; Wickstead *et al.*,
2010b). Why the kinesin repertoire has expanded so prominently in trypanosomatids remains
mysterious.

Hitherto studied kinesins of *T. brucei* have been shown to be important for various processes
occurring in different cellular compartments. KinA and KinB are involved in cell cycle progression and
localise to the central spindle during mitosis (Li *et al.*, 2008) while TbKif13-1 localizes either to the
nucleus or the mitotic spindle, where it meditates chromosome segregation during mitosis
(Wickstead *et al.*, 2010a). KinC and KinD are distributed throughout the cytoskeleton, where they are
involved in subpellicular MT organization, basal body segregation and cytokinesis (Hu *et al.*, 2012;
Wei *et al.*, 2013).

Some kinesins are associated with the flagellum and adjacent FAZ. KIN2A and KIN2B are kinesins
involved in intraflagellar transport, a process that directly affects flagellar assembly and in turn
affects cell motility and cytokinesis (Aslett *et al.*, 2009; Bertiaux *et al.*, 2020). The orphan kinesin KIN-
E predominantly localises to the flagellar tip and is required for elongation of the new FAZ (An and Li,
2018). Another orphan kinesin named KLIF initially localises to the tip of the new, elongating FAZ and
associates with the old FAZ as the cytokinetic furrow ingresses towards the posterior end, thus
mediating the final stages of cytokinesis (Hilton *et al.*, 2018). Other kinesins localising to specific
cytoskeletal structures include FCP2/TbKinX1 and FCP4/TbKin15, which are found at the flagella
connector (FC), a trypanosome-specific structure that connects the outgrowing new flagellum to the
old one (Varga *et al.*, 2017). Interestingly, FCP2/TbKinX1 belongs to kinesin clade X-1, one of the two
trypanosomatid-restricted kinesin families (Wickstead and Gull, 2006; Wickstead *et al.*, 2010b).

Here we present data on the localisation and function of two members of the kinesin superfamily,
TbKifX2 and TbPH1, with the former protein belonging to clade X-2. Functional characterization and
the unique localization of these kinesins further expands our knowledge on how these diverse and
dexterous proteins affect the biology of these extremely diverse and successful parasites.

RESULTS

TbPH1 is a kinesin-like protein that fractionates with the *T. brucei* cytoskeleton

TbPH1 (Tb927.3.2490) is a 110 kDa protein named after a C-terminal pleckstrin homology (PH)
domain, which is followed by a proximate homeodomain-like (HDL) fold and preceded by a coiled
coil (CC) domain (Fig. 1). The protein undergoes post-translational modifications, with
phosphorylation of a serine at the start of the HDL (Urbaniak *et al.*, 2012) and a methylarginine in
the middle of the primary structure (Lott *et al.*, 2013). While TbPH1 has an N-terminal kinesin motor
domain, two substitutions of highly conserved amino acids within the Walker A motif (G93P and
T/S95R) likely ablate the ATP hydrolase activity. Thus, TbPH1 is unlikely to be a *bona fide* kinesin
motor protein.

TbPH1 also belongs to the ~2000 proteins found in the granule-enriched fraction (containing cellular
aggregates, nuclei, kinetoplasts and flagella as well as associated cytoskeletal structures) obtained
by 'MT sieving', a method originally conceived to purify stress granules from *T. brucei* (Fritz *et al.*,
2015). In brief (see S1 Fig. for 'MT sieving' pipeline), cells are lysed by non-ionic detergent in low
ionic strength buffer, which maintains the cytoskeleton, including the MT corset that in turn cages
predominantly detergent insoluble cellular structures with at least one dimension >20-30 nm. The
MT corset is then disrupted by 300 mM NaCl to release these elements for subsequent analysis.

To verify TbPH1's apparent association with cytoskeletal fractions, we subjected a *T. brucei* cell line
in which the endogenous *TbPH1* open reading frame was appended with a C-terminal V5 tag (Dean
et al., 2017) to cell fractionation by a procedure identical to 'MT sieving'. The cell equivalents of the
ensuing fractions were inspected for the presence of TbPH1-V5 and SCD6, a stress granule marker
that served as a control for this method (Fig. 2A). SCD6 was mostly found in the soluble fractions as
previously reported for unstressed *T. brucei* (Fritz *et al.*, 2015), with a distribution across the
analysed fractions very distinct from that of TbPH1-V5. TbPH1-V5 was well retained in all
cytoskeletal fractions P1 and P2, although a fraction was released in the SN1 with other soluble
cytosolic proteins upon initial lysis (Fig. 2A). Importantly, an aliquot of P2 fraction that was fixed on
microscopic slides for indirect immunofluorescence (IFA) using an anti-V5 antibody shows that
TbPH1-V5 is localized as a line extending from the DAPI-stained kinetoplast (Fig. 2B) that decreases
in signal intensity distally, suggestive of a cytoskeletal localization. This was confirmed by the
observation that the vast majority of TbPH1 was released into the soluble SN3 fraction after high salt
disintegration of the cytoskeleton. A small amount of TbPH1-V5 was still found in the final pellet P3,
which is consistent with its detection in the cohort of proteins retained by 'MT sieving' (Fritz *et al.*,
2015). Thus, TbPH1 is a kinesin-like protein that fractionates with the trypanosomal cytoskeleton.

TbPH1 strongly interacts with the trypanosomatid-restricted orphan kinesin TbKifX2

Because kinesins tend to form dimers (Marx *et al.*, 2009) and kinesin-like TbPH1 contains a CC
domain (Fig. 1), which usually mediates protein-protein interactions (Burkhard *et al.*, 2001), we
proceeded to see if TbPH1-V5 has an interaction partner by immunoprecipitation (IP) via its C-
terminal epitope tag. To enrich for soluble TbPH1-V5 and augment IP stringency, we followed the
previously described fractionation technique and used fraction SN3 as the input (Fig. 3A). A mock IP
control using the parental cell line without an expressed V5 epitope was performed in parallel.

After confirming immunocapture of TbPH1-V5 in the first eluate (Fig. 3A), the IP eluate from this and
the mock control were resolved on a SDS PAGE gel. Sypro Ruby protein gel staining showed two
prominent bands of ~117 kDa (Fig. 3B, B1), the expected size for TbPH1-V5, and of ~80 kDa (Fig. 3B,
B2). Importantly, these bands were not observed in the mock control, indicating their presence is
not due to unspecific binding to the magnetic beads used to immobilize the anti-V5 antibody. Mass
spectrometry analysis of these two excised bands identified TbPH1 itself as expected and a
previously uncharacterised kinetoplastid-specific kinesin (Tb927.9.15470) (S1 Dataset), which we call

TbKifX2 based on a previously proposed kinesin nomenclature (Wickstead *et al.*, 2010b). In contrast
to TbPH1, TbKifX2's kinesin motor domain contains a conventional Walker A motif (Fig. 1). This
motor domain is located on the N-terminus, indicating that TbKiFX2 actively moves toward the plus-
end of MTs (Hirokawa *et al.*, 2009). It also contains a CC domain and two serine phosphorylation
sites (Urbaniak *et al.*, 2012), similarly to TbPH1.

Interaction between these two proteins was further verified by tagging TbKifX2 with a C-terminal HA
tag in the TbPH1-V5 cell line. Indeed, TbKifX2 mirrored the fractionation pattern of TbPH1-V5 (Fig.
3C). As before, the SN3 fraction was used as an input for TbKifX2-HA IP, which co-
immunoprecipitated TbPH1-V5. Thus, we verified that TbPH1 and TbKifX2 exhibit a strong
interaction that persists even in 300 mM NaCl.

Whole cells expressing TbPH1-V5 and TbKifX2-HA were fixed and permeabilized prior to incubation
with antibodies recognizing either epitope-tag (Figs. 3D and E). We observed that both proteins
possess a considerable cell body signal that appears to be nucleus-excluded, with a prominent signal
along the proximal region of the flagellum. This pattern is similar to the TrypTag localization data,
which visualized mNeonGreen-tagged proteins in immobilized live cells (Dean *et al.*, 2017). The cell
body signal likely represents the cytosolic population of both proteins that was also observed in our
fractionation data (Figs. 2A and 3C). It should be also noted that TbPH1-V5 only also displays an
increased signal on the anterior tip of the cell body (Fig. 3D, asterisks).

Proximity proteomics reveals TbPH1 and TbKifX2 interactors that are associated with the flagellum
attachment zone

Since conventional IP of TbPH1-V5 under high salt conditions identified only TbKifX2 as its major
interaction partner, we sought a different method to identify potential cargo and/or interactors of
this putative kinesin complex. To do this, we tagged TbPH1 with a C-terminal biotin ligase 'BioID2'
(Kim *et al.*, 2016), appended with an HA tag (Pyrih *et al.*, 2020), to facilitate proximity-dependent
biotin identification (BioID) of neighbouring proteins. First, we determined by IFA on permeabilized
whole cells using anti-HA antibody that TbPH1-BioID2-HA localizes adjacent to the kinetoplast, which
was also observed for TbPH1-V5 (*c.f.* Figs. 2B and 4A). TbPH1-BioID2-HA also exhibited biotinylation
activity, as visualized by streptavidin-conjugated to Alexa Fluor 488 (Fig. 4A), in contrast to the
parental cell line lacking BioID2 (Fig. 4B). Thus, we confirmed that the chimeric protein is properly
localized and appears to biotinylate TbPH1 itself or proximal proteins.

Purification of biotinylated proteins was performed as described elsewhere (Morriswood *et al.*,
2013) (S2 Fig.). Western blotting revealed that the vast majority of TbPH1-BioID2-HA was present in
the soluble S1 fraction (Fig. 4C), which was used for subsequent analysis. At first glance, this result
appears to disagree with previous findings from the 'MT sieving' fractionation, which employs
comparable amounts of non-ionic detergent and buffers of low ionic strength; however, we should
emphasise that 'MT sieving' fractionation uses a different detergent for a shorter incubation time
and at a lower temperature in comparison to the BioID purification (see Methods), which likely
explains this difference.

Next, the S1 fraction was incubated with streptavidin-coated magnetic beads to capture biotinylated
proteins, which were subsequently trypsinized on beads prior to liquid chromatography-tandem
mass spectroscopy (LC-MS/MS). A mock control using the parental cell line lacking BioID2 expression
was processed in parallel. A total of 1,131 proteins were detected with high confidence in at least
one of three biological replicates (Andromeda Protein Score ≥20; >1 unique peptide per protein) (S2
dataset). Following imputation of missing values, seven proteins were shown to be preferentially

biotinylated by TbPH1-BioID2-HA, as visualised by plotting the -10 log t-test p value versus the t test
difference in a volcano plot (Fig. 4D). Statistically significant hits are found above the cut-off curve in
this plot.

TbKifX2 was the most enriched protein based on both parameters, consistent with its strong
interaction with TbPH1. As expected, TbPH1 is also among the enriched proteins, indicating that
TbPH1-BioID2-HA biotinylated itself and/or a nearby untagged, endogenous form. The sub-cellular
location of the five novel proteins was determined using the TrypTag protein localization resource
(Dean et al., 2017). This allowed us to already exclude one of them (Tb927.10.12110) as a
contaminant, given its nuclear localization. A putative kinesin with a C-terminal PH domain
(Tb927.6.2880) is enriched in the basal body. Two proteins are enriched in the flagellar pocket: a
dynamin-like protein 1/2 (DLP2, Tb927.3.4760; DLP1, Tb927.3.4720) and a flagellar pocket protein
FP45 (Tb927.9.9730). DLP1/2 is involved in clathrin-mediated endocytosis (Morgan et al., 2004; Benz
et al., 2017) as well as in mitochondrial scission (Chanez et al., 2006; Benz et al., 2017). Interestingly,
FP45 has also been demonstrated to be a soluble protein that associates with the membrane
surrounding the flagellar pocket by immunogold labelling in an independent study (Gheiratmand et
al., 2013). Furthermore, TrypTag localization of FP45 demonstrates a line emanating from the
flagellar pocket toward the anterior in many cells, similar to TbPH1. Only a putative flavin-trafficking
protein (Tb927.8.1950) does not have a TrypTag localization entry at this time. However, this protein
was found among those that interact with TOEFAZ1 (alias CIF1, (Zhou et al., 2018)), a protein that
associates with the tip of the extending FAZ (Fig. 4E) (Hilton et al., 2018). Furthermore, Hilton and
colleagues N-terminally tagged Tb927.8.1950 to determine its localization by IFA, which exhibited an
enriched signal in the vicinity of the kinetoplast. Also, both TbPH1 and TbKifX2 were found within
the TOEFAZ1-BioID proximity proteome with high statistical significance (P=0.001138 and 0.0032861,
respectively), along with two other proteins biotinylated by TbPH1-BioID2-HA (Fig. 4E-F) (Hilton et
al., 2018).

We also compared the TbPH1 proximity proteome to those of TbMORN1, a protein that is part of the
hook complex (Morriswood et al., 2013), and TbSpef1, a protein localized to the MtQ (Dong et al.,
2020) (Fig. 4F). Tellingly, there was no overlap among any of these proteomes. We conclude that
none of members of the TbPH1 proximity proteome are part of the hook complex and that TbSpef1
and TbPH1 are not near each other given that the radius of biotinylation of adjacent proteins by the
BioID biotin ligase is approximately 10 nm (Kim et al., 2014).

These observations confirm the tight interaction between TbPH1 and TbKifX2. They also suggest that
these proteins associate with the basal body, a cytoskeletal element near the flagellar pocket, as
well as the FAZ given the localization of the biotinylated proteins. Thus, we proceeded to test the
basal body and FAZ localization suggested by the TbPH1-BioID (Fig. 4) and IFA (Figs. 2B, 3D and E)
results.

TbPH1 and TbKifX2 localize to the microtubule quartet within the cytoskeleton

We next decided to address whether TbPH1 and TbKifX2 localize to or near the FAZ, as was
suggested by the TbPH1 proximity proteome. Cytoskeletons were detergent-extracted from the
TbPH1-V5 and TbKifX2-HA expressing cells to visualize any potential localization to discrete
cytoskeletal structures. Indeed both proteins co-localized to a long structure running along the long-
axis of the cytoskeleton adjacent to the flagellum (Figs. 5A and S3A). Furthermore, detergent
extraction solubilized the cell body signal (Figs. 3D and E), confirming it represented the cytosolic
population of TbPH1 and TbKifX2 (Figs 2A and 3C). The specificity of the antibodies recognizing V5

and HA on extracted cytoskeletons was verified by a lack of signal in the parental cell line lacking
either of these epitopes (Fig. S3B).

We proceeded to identify specifically which cytoskeletal element TbPH1 and TbKifX2 associate with.
We used antibody recognizing TbKifX2-HA as a proxy for the TbKifX2-TbPH1 complex because of
their demonstrated tight interaction (Fig 3), and their co-localization on detergent-extracted
cytoskeletons. We note that in contrast to the TbPH1 co-localization assay depicted in Figs. 5A and
S3A, we used rabbit antibody recognizing the HA tag to allow us to counterstain with antibodies
raised in mouse and rat.

We found TbKifX2-HA exhibited a strong signal further toward the proximal region of the flagellum
that tapered in signal intensity in a line juxtaposed with the FAZ filament (Figs. 5B and S3D). This
region of strong TbKifX2-HA signal co-localized with the basal bodies (Figs. 5C and S3C). This pattern
led us to investigate whether the kinesin was bound to the MtQ, as it originates close to the basal
body and is considered a part of the FAZ (Lacomble *et al.*, 2009; Sunter and Gull, 2016). To this end,
we used the monoclonal antibody 1B41 recognizing the MtQ (Gallo *et al.*, 1988), a somewhat
contentious claim that is belied by its use as a MtQ marker in subsequent studies (Rotureau *et al.*,
2011) (see Experimental Procedures for more details about the 1B41 batch used here). Indeed,
TbKifX2-HA shows co-localization with 1B41 (Figs. 5D and S3E), suggesting that TbKifX2 may
associate with the MtQ.

To verify that TbKifX2 indeed binds the MtQ, we employed expansion microscopy (ExM), which
enlarges cells by embedding them within a swellable polyacrylamide matrix prior to immunolabelling
to allow examination of cellular structures in greater detail (Gambarotto *et al.*, 2019). ExM has
recently been implemented in *T. brucei* (Gorilak *et al.*, 2021; Amodeo *et al.*, 2021), and accentuates
the part of MtQ that wraps around the flagellar pocket upon immunolabeling with antibody
recognizing MT-incorporated, acetylated α-tubulin (Fig 6A) (Gorilak *et al.*, 2021). Additionally, this
procedure allows the proximal MtQ to be distinguished from the nearby hook complex, which lacks
MTs (Esson *et al.*, 2012).

Performing ExM allowed us to localize TbPH1-V5 or TbKifX2-HA to the part of the MtQ that spirals
around the flagellar pocket , as demonstrated by the co-localization of each epitope tag with
acetylated α-tubulin in both 3D reconstruction of expanded cells and individual confocal focal planes
(Figs. 6A-F). The associated V5 or HA signal was not discernible beyond the point of the MtQ joining
the corset MTs, where the signal of FAZ constituents, such as ClpGM6 (Hayes *et al.*, 2014), starts (*c.f.*
Figs. 6A and C with Fig. S4). The fact, that the signal of TbPH1-V5 or TbKifX2-HA was apparent only in
the part of the MtQ around the flagellar pocket is consistent with this being the region of the highest
concentration of both proteins in the cell based on IFA (Figs. 3D-E), as well as with a lower sensitivity
of ExM due to dilution of epitopes stemming from specimen expansion. Taken altogether, ExM
reveals that TbPH1 and TbKifX2 localize to the MtQ element of the *T. brucei* cytoskeleton.
Additionally, we observed a punctate signal of an anti-V5 or anti-HA antibody in the expanded cell
body (Figs. 6C-F), consistent with the significant cytosolic fraction of both proteins (Figs. 2A, 3C-E).

297 Depletion of TbPH1 and TbKifX2 causes prominent morphological defects to the posterior end of *T.*
298 brucei*

To assay the function of the novel kinesin TbKifX2 and its associated kinesin-like protein TbPH1, we
depleted their expression by doxycycline-induced RNAi in the cell line expressing TbPH1-V5 and
TbKifX2-HA. The downregulation of either protein alone only marginally affected the growth of *T.*
brucei in comparison to the non-induced cells, despite efficient depletion of each of the respective

tagged proteins (Fig. 7A, Fig. S5A). After 6 days of RNAi-silencing of TbKifX2, we fractionated the
cytoskeleton from the cytosol under low salt conditions (Fig. 7B). Consistent with 'MT sieving'
fractionation (Figs. 2A and 3C) and IFA (Figs 3D-E and 5A), western blot detection revealed TbKifX2-
HA and TbPH1-V5 in both cytoskeletal and cytosolic fractions in cells grown in the absence of
tetracycline, in which neither protein was downregulated (Fig. 7B). However, TbPH1 was
preferentially depleted from the cytoskeleton fraction upon TbKifX2 silencing. This result indicates
that TbKifX2 recruits TbPH1-V5 to the MtQ (Fig. 7B).

To get a more robust RNAi phenotype for phenotypic analysis, we generated cell lines in which both
proteins were downregulated, which showed more pronounced growth inhibition than the single
knock-downs (Fig. 7A). Growth slowed after 3 days of RNAi induction and did not reach levels of the
uninduced counterparts at any point during the 7 day time course (Fig. S5B). Depletion of both
proteins was confirmed by western blotting (Fig. S5B, inset), although neither protein was decreased
below the detection limits of this assay, perhaps explaining the lack of a more severe growth
phenotype.

Because both TbPH1 and TbKifX2 are modified in a way suggestive of post-translational regulation
(Fig. 1), plus both associate with proteins that may regulate the cell cycle (*e.g.* FP45) (Gheiratmand
et al., 2013)**,** we wondered whether the growth inhibition of the double RNAi knockdown is due to a
cell cycle defect. Thus, we assayed the proportion of cell cycle stages present over a course of 7 days
of TbPH1/TbKifX2 depletion, sorting individual DAPI-stained cells into different cell cycle stages by
scoring their number of nuclei (N) and kinetoplasts (K) (Robinson *et al.*, 1995). The proportion of
1N1K cells decreased while 2N2K cells doubled at the same time from around 10% of the population
to about 20% (Fig. S5C). The number of zoids (cells with 1 kinetoplast and lacking a nucleus) reached
a peak at day 5, making up about 7% of the population, which may suggest problems with
cytokinesis. However, we did not observe the consequent emergence of 2N1K or multinucleated
cells, arguing against a specific cytokinesis defect (Robinson *et al.*, 1995). Thus, we conclude that
either TbKifX2 and TbPH1 do not play a direct role in cell cycle progression and cytokinesis, or the
partial depletion of these proteins did not reach the threshold needed to affect cytokinesis. Thus we
looked for other phenotypes that would explain the moderately decreased fitness of these cells.

We assayed the MtQ and FAZ by IFA after 5 days of RNAi depletion and observed no defects in either
of these structures, indicating that TbPH1 and TbKifX2 do not play a role in their biogenesis (Fig.
S5D). However, by phase contrast microscopy we noticed the emergence of two morphological
phenotypes, namely multiple cells joined together on their posterior end, a phenotype resembling
abscission defects (Ralston *et al.*, 2006; Benz *et al.*, 2012) and single cells with protrusions from the
posterior end (Fig. 7C).

To better visualize these phenotypes, we observed these cells by scanning electron microscopy (Fig.
7D). While control cells showed normal trypomastigote morphology (Fig. 7D), we regularly observed
cells with multiple protrusions (arrows) originating from the posterior end in the induced population
(Fig. 7D). The occurrence of these cell types was also corroboratively found in transmission electron
microscopy, which in addition provided evidence that these protrusions appear to contain normal
cellular contents and were surrounded by subpellicular MTs (Fig. S6A-B).

This effect of TbPH1/TbKifX2-depletion resembled the RNAi-phenotype of two tubulin-tyrosine-
ligase-like (TTLL) proteins, which post-translationally modify the C-terminal tails of α- and β-tubulin
by addition of polyglutamate side chains (Jentzsch *et al.*, 2020). Upon depletion of either TTLL,
protrusions were observed emanating from the posterior end, which prompted the authors to name
this morphological change the 'glove' phenotype. To determine whether TbPH1/TbKifX2

downregulation truly phenocopies that of each TTLL, we immunolabelled cells 5 days post RNAi-
induction with antibodies recognizing glutamyl side chains (GT335 antibody) and tyrosinated α-
tubulin (YL 1/2 antibody) (Fig. 5C and S3C); the latter antibody was used previously to visualize the
basal bodies, where it additionally cross-reacts with TbRP2 (Andre *et al.*, 2014). Upon depletion of
either TTLL, polyglutamylation was decreased and α-tubulin tyrosination increased in the posterior
end, from which the protrusions emerged (Jentzsch *et al.*, 2020). We did not observe either of these
changes in post-translational modifications upon TbPH1/TbKifX2 silencing by observing them on
individual cells by IFA (Fig. S6C) or the whole cell population by western blot analysis (Fig. S6D),
suggesting the mechanism underlying the emergence of posterior-end protrusions differs from that
of the TTLLs.

DISCUSSION

Kinesins and related kinesin-like proteins constitute one of the largest protein families encoded by
the *T. brucei* genome (Berriman *et al.*, 2005) and are represented by diverse gene repertoires in
other trypanosomatids as well (El-Sayed *et al.*, 2005; Ivens *et al.*, 2005; Wickstead and Gull, 2006;
Wickstead *et al.*, 2010b). Among the eukaryotes, trypanosomatids have experienced one of the
largest expansions of the kinesin motor superfamily, suggesting these proteins endow trypanosomes
with unique biological properties, many of which still await elucidation (Berriman *et al.*, 2005;
Wickstead and Gull, 2006; Wickstead *et al.*, 2010b). Indeed, this expansion has resulted in the
emergence of two trypanosomatid-specific clades of kinesins, named X-1 and X-2. A kinesin
belonging to the former, named FCP2/TbKinX1, has been shown to maintain a connection of a new
flagellum's distal tip along the side of the old flagellum, thus facilitating the unique mechanism of
templated flagellar replication in *T. brucei* (Varga *et al.*, 2017).

In this work, we have shed light on the function of a member of the trypanosomatid-specific X-2
clade, which we named TbKifX2. We show that TbKifX2 is localized to the MtQ component of the
cytoskeleton while also maintaining a cytosolic fraction. Thus, TbKifX2 is the first motor protein that
has been demonstrated to localize to the MtQ, a cytoskeletal component that is considered to be
part of the FAZ and restricted to trypanosomatids (Sunter and Gull, 2016).

TbKifX2 recruits TbPH1, a likely catalytically-dead, kinesin-like protein with a C-terminal PH domain,
to the MtQ. Given that TbPH1 remains in the cytosolic fraction upon TbKifX2 depletion argues that
both proteins do not always interact. This notion is also consistent with the enrichment of TbPH1 on
the anterior tip of the cell, which was not observed in KifX2. However, when TbKifX2 and TbPH1 do
interact on the MtQ, the part of the cytoskeleton where both proteins co-localize, they interact with
a strong affinity that persists in high salt concentrations, a condition that usually disrupts non-
covalent interactions between the CC domains (Grigoryan and Keating, 2008).

As far as we know, there are not many reports of kinesin interactions with a protein with a kinesin-
like structure, but still being catalytically inactive. The most extensively characterized example being
the budding yeast Kar3 kinesin and Vik1, which confers to the Kar3/Vik1 complex a strong
interaction with a microtubule (Allingham *et al.*, 2007). We hypothesize that the TbKifX2/TbPH1
complex may represent an adaptation for binding the MtQ. How the TbKifX2/TbPH1 complex
preferentially interacts with the MtQ over the corset microtubules is an intriguing question. It has
been suggested that the MtQ contains β-tubulin with a unique but still unknown modification that is
recognized by the 1B41 monoclonal antibody (Gallo *et al.*, 1988; Gallo and Precigout, 1988), and
could have an increased affinity to the kinesin complex investigated here. Alternatively, the kinesins

could be activated at a specific site, for example at the proximal end of the MtQ, similar to the yeast
kinesin Kip2 being recruited to the minus ends of microtubules by the activity of the spindle pole
bodies (Chen *et al.*, 2019). However, at this juncture, we do not know whether TbKifX2 and TbPH1
interaction occurs prior to or after the proteins bind the MtQ.

The TbPH1 proximity proteome obtained *via* BioID labelling reassuringly revealed a protein cohort
that is enriched around the proximal end of the MtQ. Among these were another kinesin with a PH
domain that was localized to the basal bodies. Also, two proteins were found to be associated with
the flagellar pocket, including FP45 (Gheiratmand *et al.*, 2013). The other protein is interestingly a
putative flavin-trafficking protein, containing an ApbE-like domain, which is common in prokaryotes
but a rare occurrence in eukaryotic proteins (Schenk *et al.*, 2021). What an enzyme potentially
influencing redox is doing in association with the MtQ is another open question that has been
uncovered in this study.

The TbPH1 proximity proteome was also informative in terms of proteins that were missing. One
group of absent proteins are those identified to be part of the hook complex (Morriswood *et al.*,
2013), allowing us to conclude that TbPH1 does not associate with this cytoskeletal structure which
lacks MTs (Esson *et al.*, 2012). Also, the eleven high confidence proteins found to be in the TbSpef1
proximity proteome were not found in our study (Dong *et al.*, 2020). This is quite surprising given
that TbSpef1 also localized to the proximal MtQ (Gheiratmand *et al.*, 2013). This may indicate that
these two proteins maintain a constant spatial separation, even if they both seemingly localize to the
same, constrained part of the cell. Indeed the biotin ligase fused to TbPH1 to mediate BioID
biotinylates proteins within a 10 nm radius (Kim *et al.*, 2014), suggesting that TbSpef1 and TbPH1,
and perhaps other MtQ-binding proteins, may be partitioned at nanoscales. Consistent with this
notion, only one of the eleven proteins in the TbSpef1 proximity proteome was found in that of
TOEFAZ, namely the known FAZ protein named FAZ 10 (Moreira *et al.*, 2017). The higher degree of
overlap between the proximity proteomes of TOEFAZ, which is enriched at the distal tip of the FAZ
(Hilton *et al.*, 2018; Zhou *et al.*, 2018), and TbPH1 in comparison to the TbSpef1 proximity proteome
indicates that perhaps TbPH1 is more distal to TbSpef1 in the MtQ.

Simultaneous depletion of TbPH1 and TbKifX2 does not affect the overall morphology of MtQ or FAZ,
indicating they are likely not involved in the biogenesis of these cellular structures. As these single
copy organelles must be faithfully replicated during cell proliferation, it is consistent with the double
knockdown having a minimal impact on the cell cycle. Nevertheless, TbKifX2/ TbPH1 downregulation
has a striking morphogenesis phenotype, namely the accumulation of protrusions from the posterior
end of *T. brucei*. The intracellular contents of these protrusions appear to be comparable to the rest
of the cell and they are also surrounded by subpellicular MTs. Interestingly, these protrusions are
reminiscent of the 'glove phenotype' observed upon depletion of two TTLL enzymes, which are
responsible for post-translational modifications appending the C-terminal tails of α- and β-tubulin
(Jentzsch *et al.*, 2020).

However, we can now only speculate about the nature of these posterior extrusions. For one,
simultaneous TbPH1 and TbKifX2 depletion did not affect polyglutamylation or tyrosination of
tubulins, indicating that these proteins do not play a role in regulating either of these post-
translational modifications as is the case for either TTLL (Jentzsch *et al.*, 2020). Malfunction of
regulated polyglutamylation and tyrosination led to mislocalization of EB1, a MT-plus-end-binding
protein that binds to the MT corset at the posterior tip (Sheriff *et al.*, 2014; Jentzsch *et al.*, 2020). It
was proposed that EB1 proteins may dock onto tubulins with the aforementioned post-translational
modifications (Jentzsch *et al.*, 2020). So, the malformations seen in TbPH1/TbKifX2 may affect EB1 or
other such MT-plus-end-binding proteins, leading to uncontrolled MT growth or aberrant MT

organization. However, it should be noted that EB1 was not identified in the TbPH1 proximity
proteome, so any effect on this protein may be an indirect consequence of RNAi-silencing. The
protrusions may also be a consequence of delayed or uncoordinated cytokinesis or represent
mechanical damage to the posterior ends due to faulty separation of daughter cells at the final stage
of cytokinesis. However, arguing against protrusions arising from a putative cytokinesis defect is that
they were not observed in *T. brucei* unable to pull apart due to motility defects (Ralston *et al.*, 2006).

The finding that the depletion of proteins that associate with the MtQ can affect the gross
morphology of *T. brucei* was unexpected. Unfortunately, the proteins that were identified in the
TbPH1 proximity proteome do not provide an obvious explanation that would link the phenotype to
the localization of TbKifX2 and TbPH1. Nevertheless, we have uncovered an intriguing relationship
between the MtQ and *T. brucei* morphogenesis that is worthy of further exploration.

EXPERIMENTAL PROCEDURES

Cells

Procyclic SmOxP9 cells (Poon *et al.*, 2012) were grown in SDM79 with 1 µg/ml of puromycin. Cell
lines bearing epitope tags were grown with 50 µg /ml hygromycin (TbPH1-V5), 15 µg /ml G418
(TbKifX2-HA) and 10 µg /ml blasticidin (TbPH1-BioID2-HA). Selection for the RNAi constructs was
with 2.5 µg /ml phleomycin (TbKifX2 RNAi, TbPH1 RNAi and TbKifX2+TbPH1 RNAi).

Plasmids and generation of cell lines

For epitope tagging at the endogenous locus with HA and V5, long PCR products were generated
with primers TbPH1V5 FW (TTTCTTCCGGTCTTGACTTATTCCGCTCAGGGAAGCTTTATGATTTCTTATGCGA
AAAGAGAGTCATACCGCTCCCGTACggttctggtagtggttcc) and TbPH1V5 RV (ATAATAAAGAGGAAGGGAAG
GTAAAGTTCAGAAACAAATTCTGTTGGCTCCTGATAACACTCTCATATTTCCTTCACCGCccaatttgagagacctgtg
c) and primer TbKifX2HA FW (AACTACAGTTGCAAATGTGCATGATAGGAAGAAGTTGCCTCACGATGGAGC
GTTGGAAAGCGGAACATGTCGGAACCGCAggttctggtagtggttcc) and TbKifX2HA RV (CATTTTAAACAATTTC
TCCT TCCCTCTTTTTTCACTCATTCATTCATTCATTCACCACCTTCTTTTTTTTTTTGCCTcctgtgc) using
pPOT-V5-HygR and pPOT-HA-NeoR vectors as templates (based on (Kaurov *et al.*, 2018) and
modified from pPOTv4 (Dean *et al.*, 2017)). For tagging with the BioID2 ligase (38), a modified
version of pPOTv7-Blast-mNG (Pyrih *et al.*, 2020) was used with the aforementioned primers for
TbPH1 endogenous locus tagging. 50 µl PCR reactions were directly transfected into procyclic
SmOxP9 cells using the Amaxa Nucleofactor electroporator and selection occurred approximately 16
hours after electroporation.

To create a vector to simultaneously downregulate both TbPH1 and TbKifX2, a fragment of each (as
suggested by RNAit2 software (https://dag.compbio.dundee.ac.uk/RNAit/) (Redmond *et al.*, 2003))
was PCR amplified using primers TbPH1-RNAi-Fw (TAATtctagaGTTGCGTGATGAGCTTCAAA), TbPH1-
RNAi-Rv (TATAaagcttCTCCTCTAGCACCGTTTTGC), TbKifX2-RNAi-Fw (TAATggatccCACCGCCATTCTGCA
AACAA) and TbKifX2-RNAi-Rv (TATAtctagaGTCATTTGCGTGGGCGATAC) and cloned into BamHI-HindIII
sites of p2T7-177 (Wickstead *et al.*, 2002) in a three way ligation using XbaI as the restriction site in
between the two RNAi fragments. The plasmid was linearised with NotI before electroporation.

RNAi and growth curves

Procyclic form cell lines for depletion of either TbPH1 or TbKifX2 were induced at a density of $2x10^6$
cells/ml with 1 µg/ml doxycycline and growth monitored daily with a Z2 Coulter Counter Analyzer
(Beckman Coulter). Cells were diluted back to the initial cell density every day. Cell lines carrying the

vector for simultaneous depletion of TbPH1 and TbKifX2 were induced at a density of $5x10^5$ cells/ml
with 1 µg/ml doxycycline and growth monitored daily with a Neubauer counting chamber. Cells were
diluted back to the initial cell density every two days.

DAPI counts

485 For cell cycle analysis, cells either induced or not for depletion of TbKifX2 and TbPH1 were spread on
486 glass slides, the cell suspension allowed to air-dry and the slides then fixed in methanol at -20 °C.
487 Slides were rehydrated for 5 minutes in PBS before a drop of ProLong™ Gold antifade reagent with
488 DAPI (ThermoScientific) was added, a coverslip applied and the slide sealed. At least 200 cells were
489 scored according to the number of nuclei and kinetoplasts per slide.

Cell fractionation via 'microtubule sieving'

491 To entrap non-soluble and cytoskeletal proteins within the MT corset, a MT sieving technique as
492 described in (Fritz *et al.*, 2015) was employed. The experiment was conducted exactly as described
493 before in (Fritz *et al.*, 2015) (see also S1 Fig.).

V5 immunoprecipitation

For immunoprecipitation (IP) of TbPH1-V5 from procyclic trypanosome lysates, the V5 antibody
(Thermo Fisher) was coupled and cross-linked to protein G coated dynabeads (Thermo Scientific)
using 6 mg/ml dimethyl pimelidate dihydrochloride in 0.2M triethanolamine. Beads were then
washed in IP wash buffer (20 mM Tris-HCl, pH 7.7, 100 mM KCl, 3 mM $MgCl_2$, 0.5% TritonX-100,
cOmplete™ EDTA-free protease inhibitor (Roche)) and stored at 4 °C before further usage. For
proteomic analysis of TbPH1 interactions, $1x10^9$ PCF cells were pre-fractionated using the MT sieving
technique described above. The SN3 lysate was incubated with V5-coupled dynabeads for 24 hours
at 4 °C. Following several washes in IP wash buffer, proteins were eluted using low pH glycine buffer.
Aliquots were taken of the different steps for western blot analysis.

Western blots

Cell lysates were prepared by boiling in 1x SDS sample buffer and the equivalent of $2x10^6$ cells (RNAi
lines) or $1x10^7$ cells (IP samples) was loaded per well. Following western blotting, PVDF membranes
were blocked for 1 hour in 5% milk-PBS. Incubation with primary antibodies (anti-V5 at 1:1,000;
ThermoScientific, anti-HA at 1:1,000; ThermoScientific, anti-tubulin at 1:5,000; Sigma) was
performed at 4 °C overnight. Secondary antibodies (HRP-coupled anti-mouse or –rabbit IgG; Sigma)
were used at a concentration of 1:2,000 and signals were visualised using Clarity Western ECL
substrate (Bio-Rad) on a ChemiDoc MP (Bio-Rad).

SYPRO Ruby staining

513 Following electrophoresis, polyacrylamide gels were placed into fixing solution (7% glacial acetic
514 acid, 50% methanol) for 30 minutes at room temperature. Following another, identical fixation step,
515 the gel was incubated with SYPRO Ruby solution overnight at room temperature. A washing step in
516 washing solution (7% glacial acetic acid, 10% methanol) for 30 minutes at room temperature was
517 followed by three 5 minute washes in milliQ water before imaging on a ChemiDoc MP (Bio-Rad).

Immunofluorescence and cytoskeleton extraction

Approximately $2x10^6$ cells were used per slide. Cells were washed in PBS and fixed in 2.3% PFA in PBS
for approximately 30 minutes before being transferred to microscopy slides (ThermoScientific).
Following neutralisation in 0.1M glycine in PBS, slides were incubated in methanol at -20° C

overnight for permeabilisation. Slides were then rehydrated in PBS and blocked in 1% BSA in PBS for
1 hour followed by incubation with primary antibody for 1 hour in a humid chamber. The slides were
then washed three times with PBS before incubation in AlexaFluor-conjugated secondary antibody
(goat anti-rabbit/mouse, used at 1:1,000; Invitrogen). Following three further washes in PBS, a drop
of ProLong Gold Antifade reagent with DAPI (ThermoScientific) was added, a coverslip applied and
the slide sealed with nail polish. Slides were imaged with a Zeiss Axioscope or an Olympus FluoView
Fv1000 confocal microscope.

For immunofluorescence analysis of cells extracted using the 'MT sieving' fractionation method
described above the pellet fraction P2 was fixed in 2.3% PFA as described. All subsequent steps were
the same as for whole cell immunofluorescence.

To extract cytoskeletons conventionally, approximately $1x \, 10^7$ PCF cells were collected and
centrifuged at room temperature for 5 min at 800 x g and the supernatant discarded. Thereafter, the
cell pellet was washed once in PBS and after resuspension in PBS applied in a drop-wise fashion to
Superfrost plus® slides (ThermoScientific). The cells were left to settle before the addition of PEME
buffer (100 mM Pipes, pH 6.9, 1 mM $MgSO_4$, 2 mM EGTA, 0.1 mM EDTA) containing 0.5% (v/v) NP-40
(Igepal) for 10 seconds. This cytoskeleton extraction was followed by a 10 minute fixation step in 4%
PFA. All subsequent steps were as described above for whole cell immunofluorescence.

<u>Antibodies used in immunofluorescence</u>

Antibody	Dilution	Origin
anti-HA (produced in rabbit)	1:1,000	Sigma, Catalog #: H6908
anti-V5 (produced in mouse)	1:1,000	ThermoScientific, Catalog #: 377500
1B41 (MtQ)[1]	1:1,000	(Gallo et al., 1988)
L3B2	1:10	(Kohl et al., 1999)
YL1/2	1:50	(Sherwin and Gull, 1989)
GT335	1:2000	Adipogen Life Sciences, Catalog #: AG-20B-0020
[1]IgM purified from hybridoma supernatant and not mouse ascitic fluid as in original report.		

Scanning and transmission electron microscopy

For TEM, cell pellets were high-pressure frozen, freeze substituted in the presence of 2% OsO4 in
acetone, and embedded into Spi-Pon812 resin (SPI) as described previously (Kaurov et al., 2018). For
SEM of the TbPH1/TbKifX2 RNAi cell line, cells were fixed in 2.5% glutaraldehyde in 100 mM PBS
overnight at 4°C and then spotted onto poly-l-lysine-coated glass cover slips. The cells were post-
fixed in 2% osmium tetroxide in 100 mM PBS for 1 h at room temperature and finally washed in the
same buffer. After dehydration cells were critical-point dried, coated with gold palladium and
imaged.

Expansion microscopy

Expansion microscopy samples were prepared as described previously (Gorilak et al., 2021). In brief,
$1x10^6$ cells were fixed overnight in a solution containing 4% formaldehyde and 4% acrylamide in PBS
and adhered to a poly-L-lysine (Sigma, P4707) coated coverslip. The cells were gelated for 30 min at
37°C in PBS supplemented with 19% sodium acrylate, 10% acrylamide, and 0.1% N, N'-
Methylenebisacrylamide, 0.5% N, N, N′, N′-tetramethylethylenediamine, and 0.5% ammonium
persulfate. The specimens were further denatured by a 2 hour incubation at 95°C in a buffer

consisting of 50 mM tris(hydroxymethyl)aminomethane-hydrochloride, 200 mM sodium chloride,
200 mM sodium dodecyl sulphate, pH 9.0, and expanded by three 20 min incubations with 15 ml of
ultrapure water.

Thereafter, the gels were incubated overnight first with a mixture of anti-HA (Cell Signaling
Technology, 3724S; used at 1:250) and C3B9 ((Woods *et al.*, 1989), used at 1:10), anti-V5 (Sigma,
V8137-.2MG; used at 1:250) and C3B9 or anti-ClpGM6 ((Hayes *et al.*, 2014), used at 1:200) and C3B9
antibodies, followed by washes and an overnight incubation with a mixture of secondary antibodies
(Invitrogen, A11001 and A21428; used at 1:500). Both primary and secondary antibodies were
diluted in PBS supplemented with 2% bovine serum albumin. Finally, the gels were washed with
ultrapure water.

Prior to imaging pieces of gels containing stained expanded cells was transferred to a glass-bottom
dish coated with poly-L-lysine, and imaged using a Leica TCS SP8 confocal microscope with an HC PL
apochromatic 63x/NA 1.40 oil immersion objective. Z-stacks were acquired with the step size of 100
nm. Confocal z-stacks were processed in Fiji (Schindelin *et al.*, 2012). 3D reconstructions were
performed in the 3D viewer plugin.

BioID

PCF cells (TbPH1-BioID2-HA and SmOxP9 as controls) were grown in the presence of 50 µM biotin for
24 hours. For proximity-dependent <u>bio</u>tin <u>id</u>entification (BioID), 10^9 PCF cells were extracted in PEME
buffer (100 mM Pipes, pH 6.9, 1 mM $MgSO_4$, 2 mM EGTA, 0.1 mM EDTA) containing 0.5% (v/v) NP-40
(Igepal) for 15 minutes at room temperature resulting in extract E1. Following centrifugation at
3,400 x g for 2 minutes, supernatant S1 was created and pellet P1 was further processed by
extraction in lysis buffer (0.4% SDS, 500 mM NaCl, 5 mM EDTA, 1 mM DTT, 50 mM Tris-HCl, pH7.4).
Another centrifugation step at 16,000 x g for 10 minutes created supernatant S2. Both supernatants,
S1 and S2, were then incubated with streptavidin-conjugated Dynabeads (company) for 4 hours at 4
°C. An aliquot of flow through samples F1 and F2 were retained for western blotting and the
dynabeads were washed five times with PBS. A small sample of the beads was then resuspended in
2x SDS PAGE buffer and boiled, while the remainder of the beads was stored at -80 °C until further
processing for mass spectrometry analysis.

Mass spectroscopy analysis of captured biotinylated proteins

Trypsin-digestion of captured biotinylated proteins was performed on bead prior to liquid
chromatography-tandem mass spectroscopy (LC-MS/MS) as previously described (Pyrih *et al.*, 2020).
Data was processed using MaxQuant (Cox and Mann, 2008) version 1.6.14 which incorporates the
Andromeda search engine (Cox *et al.*, 2011). Proteins were identified by searching a protein
sequence database containing *T. brucei brucei* 927 annotated proteins (Version 51, TriTrypDB (Aslett
et al., 2009), http://www.tritrypdb.org/) supplemented with frequently observed contaminants.
Search parameters specified an MS tolerance of 6 ppm, an MS/MS tolerance at 0.5 Da and full
trypsin specificity, allowing for up to two missed cleavages. Carbamidomethylation of cysteine was
set as a fixed modification and oxidation of methionine and N-terminal protein acetylation were
allowed as variable modifications. The experimental design included matching between runs for
biological replicates. Peptides were required to be at least 7 amino acids in length, with false
discovery rates (FDRs) of 0.01 calculated at the levels of peptides, proteins and modification sites
based on the number of hits against the reversed sequence database. The obtained data was
subsequently processed in Perseus version 1.6.14 as described in (Zoltner *et al.*, 2020).

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium
via the PRIDE partner repository with the dataset identifier PXD025802.

ACKNOWLEDGEMENTS

We thank Keith Gull (University of Oxford), Linda Kohl (Muséum National d'Histoire Naturelle), Bill
Wickstead (University of Nottingham), and Ziyin Li (University of Texas Medical School at Houston)
for helpful comments, Susanne Kramer (University of Würzburg), Cynthia He (National University of
Singapore), and Chris de Graffenried (Brown University) for antibodies, Peter Gorilak (Institute of
Molecular Genetics) for assistance with imaging, as well as Karel Harant and Pavel Talacko (BIOCEV,
Prague) and Zbyněk Zdráhal and David Potěšil (CEITEC, Brno) for mass spectroscopy analysis on BioID
samples and excised bands, respectively. This work was supported by the Czech Science Foundation
grants 20-23513S to HH, and 21-09283S and ERC CZ LL1601 to JL, the ERD funds of the Czech
Ministry of Education 16_019/0000759 to HH and JL, and BioImaging grant LM2018129 to MV. Work
in the VV laboratory was supported by an EMBO Installation Grant and by the J. E. Purkyne
Fellowship of the Czech Academy of Sciences.

Author contributions

Conceptualization (CB, NM, SK, VV, HH), formal analysis (CB, NM, SK, HV, MV, VV, HH), investigation
(CB, NM, SK, HV, MV), visualization (CB, VV, HH), writing – original draft (CB, HV, JL, VV, HH), funding
acquisition (JL, VV, HH), project administration (HH).

REFERENCES

Albisetti, A., Florimond, C., Landrein, N., Vidilaseris, K., Eggenspieler, M., Lesigang, J., *et al.* (2017)
Interaction between the flagellar pocket collar and the hook complex via a novel microtubule-
binding protein in *Trypanosoma brucei*. *PLoS Pathog* **13**: e1006710
https://pubmed.ncbi.nlm.nih.gov/29091964/.

Allingham, J.S., Sproul, L.R., Rayment, I., and Gilbert, S.P. (2007) Vik1 Modulates Microtubule-Kar3
Interactions through a Motor Domain that Lacks an Active Site. *Cell* **128**: 1161–1172
https://pubmed.ncbi.nlm.nih.gov/17382884/.

Amodeo, S., Kalichava, A., Fradera-Sola, A., Bertiaux-Lequoy, E., Guichard, P., Butter, F., and
Ochsenreiter, T. (2021) Characterization of the novel mitochondrial genome segregation factor
TAP110 in *Trypanosoma brucei*. *J Cell Sci* **134**: jcs.254300
http://jcs.biologists.org/lookup/doi/10.1242/jcs.254300.

An, T., and Li, Z. (2018) An orphan kinesin controls trypanosome morphology transitions by targeting
FLAM3 to the flagellum. *PLOS Pathog* **14**: e1007101
https://dx.plos.org/10.1371/journal.ppat.1007101.

Andre, J., Kerry, L., Qi, X., Hawkins, E., Drižyte, K., Ginger, M.L., and McKean, P.G. (2014) An
alternative model for the role of RP2 protein in flagellum assembly in the African trypanosome. *J Biol
Chem* **289**: 464–475.

Aslett, M., Aurrecoechea, C., Berriman, M., Brestelli, J., Brunk, B.P., Carrington, M., *et al.* (2009)
TriTrypDB: A functional genomic resource for the Trypanosomatidae. *Nucleic Acids Res* **38**: D457–

D462 http://tritrypdb.org.

Benz, C., Clucas, C., Mottram, J.C., and Hammarton, T.C. (2012) Cytokinesis in bloodstream stage
Trypanosoma brucei requires a family of katanins and spastin. *PLoS One* **7**: e30367.

Benz, C., Stříbrná, E., Hashimi, H., and Lukeš, J. (2017) Dynamin-like proteins in *Trypanosoma brucei*:
A division of labour between two paralogs? *PLoS One* **12**: e0177200
https://pubmed.ncbi.nlm.nih.gov/28481934/.

Berriman, M., Ghedin, E., Hertz-Fowler, C., Blandin, G., Renauld, H., Bartholomeu, D.C., *et al.* (2005)
The genome of the African trypanosome *Trypanosoma brucei*. *Science* **309**: 416–422
http://dx.doi.org/10.1126/science.1112642.

Bertiaux, E., Mallet, A., Rotureau, B., and Bastin, P. (2020) Intraflagellar transport during assembly of
flagella of different length in *Trypanosoma brucei* isolated from tsetse flies. *J Cell Sci* **133**
https://pubmed.ncbi.nlm.nih.gov/32843573/.

Burkhard, P., Stetefeld, J., and Strelkov, S. V. (2001) Coiled coils: A highly versatile protein folding
motif. *Trends Cell Biol* **11**: 82–88 https://pubmed.ncbi.nlm.nih.gov/11166216/.

Chanez, A.-L., Hehl, A.B., Engstler, M., and Schneider, A. (2006) Ablation of the single dynamin of *T.*
brucei blocks mitochondrial fission and endocytosis and leads to a precise cytokinesis arrest. *J Cell Sci*
119: 2968–74 http://www.ncbi.nlm.nih.gov/pubmed/16787942.

Chen, X., Widmer, L.A., Stangier, M.M., Steinmetz, M.O., Stelling, J., and Barral, Y. (2019) Remote
control of microtubule plus-end dynamics and function from the minus-end. *Elife* **8**: 1–32.

Cox, J., and Mann, M. (2008) MaxQuant enables high peptide identification rates, individualized
p.p.b.-range mass accuracies and proteome-wide protein quantification. *Nat Biotechnol* **26**: 1367–
1372 https://www.nature.com/articles/nbt.1511.

Cox, J., Neuhauser, N., Michalski, A., Scheltema, R.A., Olsen, J. V., and Mann, M. (2011) Andromeda:
A peptide search engine integrated into the MaxQuant environment. *J Proteome Res* **10**: 1794–1805
https://pubmed.ncbi.nlm.nih.gov/21254760/.

Dean, S., Sunter, J.D., and Wheeler, R.J. (2017) TrypTag.org: A Trypanosome Genome-wide Protein
Localisation Resource. *Trends Parasitol* **33**: 80–82 https://pubmed.ncbi.nlm.nih.gov/27863903/.

Dong, X., Lim, T.K., Lin, Q., and He, C.Y. (2020) Basal Body Protein TbSAF1 Is Required for
Microtubule Quartet Anchorage to the Basal Bodies in *Trypanosoma brucei*. *mBio* **11**: e00668-20
http://mbio.asm.org/content/11/3/e00668-20.abstract.

El-Sayed, N.M., Myler, P.J., Bartholomeu, D.C., Nilsson, D., Aggarwal, G., Tran, A.N., *et al.* (2005) The
genome sequence of *Trypanosoma cruzi*, etiologic agent of Chagas disease. *Science* **309**: 409–415
https://pubmed.ncbi.nlm.nih.gov/16020725/.

Esson, H.J., Morriswood, B., Yavuz, S., Vidilaseris, K., Dong, G., and Warren, G. (2012) Morphology of
the trypanosome bilobe, a novel cytoskeletal structure. *Eukaryot Cell* **11**: 761–772.

Field, M.C., and Carrington, M. (2009) The trypanosome flagellar pocket. *Nat Rev Micro* **7**: 775–786
http://dx.doi.org/10.1038/nrmicro2221.

Fritz, M., Vanselow, J., Sauer, N., Lamer, S., Goos, C., Siegel, T.N., *et al.* (2015) Novel insights into RNP
granules by employing the trypanosome's microtubule skeleton as a molecular sieve. *Nucleic Acids*
Res **43**: 8013–8032 https://pubmed.ncbi.nlm.nih.gov/26187993/.

Gallo, J.M., and Precigout, E. (1988) Tubulin expression in trypanosomes. *Biol Cell* **64**: 137–143
https://pubmed.ncbi.nlm.nih.gov/3067795/.

Gallo, J.M., Précigout, E., and Schrével, J. (1988) Subcellular sequestration of an antigenically unique
beta-tubulin. *Cell Motil Cytoskeleton* **9**: 175–183 https://pubmed.ncbi.nlm.nih.gov/2452022/.

Gambarotto, D., Zwettler, F.U., Guennec, M. Le, Schmidt-Cernohorska, M., Fortun, D., Borgers, S., *et*
al. (2019) Imaging cellular ultrastructures using expansion microscopy (U-ExM). *Nat Methods* **16**: 71–
74 https://pubmed.ncbi.nlm.nih.gov/30559430/.

Gheiratmand, L., Brasseur, A., Zhou, Q., and He, C.Y. (2013) Biochemical characterization of the bi-
lobe reveals a continuous structural network linking the bi-lobe to other single-copied organelles in
Trypanosoma brucei. J Biol Chem **288**: 3489–3499 https://pubmed.ncbi.nlm.nih.gov/23235159/.

Gorilak, P., Pružincová, M., Vachova, H., Olšinová, M., and Varga, V. (2021) Expansion microscopy
facilitates quantitative super-resolution studies of cytoskeletal structures in kinetoplastid parasites.
Open Biol **11**: 210131 https://doi.org/10.1101/2021.04.20.440601.

Grigoryan, G., and Keating, A.E. (2008) Structural specificity in coiled-coil interactions. *Curr Opin*
Struct Biol **18**: 477–483 https://pubmed.ncbi.nlm.nih.gov/18555680/.

Gull, K. (1999) The cytoskeleton of trypanosomatid parasites. *Annu Rev Microbiol* **53**: 629–655
https://pubmed.ncbi.nlm.nih.gov/10547703/.

Hayes, P., Varga, V., Olego-Fernandez, S., Sunter, J., Ginger, M.L., and Gull, K. (2014) Modulation of a
cytoskeletal calpain-like protein induces major transitions in trypanosome morphology. *J Cell Biol*
206: 377–384.

Hilton, N.A., Sladewski, T.E., Perry, J.A., Pataki, Z., Sinclair-Davis, A.N., Muniz, R.S., *et al.* (2018)
Identification of TOEFAZ1-interacting proteins reveals key regulators of *Trypanosoma brucei*
cytokinesis. *Mol Microbiol* **109**: 306–326 http://doi.wiley.com/10.1111/mmi.13986.

Hirokawa, N., and Noda, Y. (2008) Intracellular transport and kinesin superfamily proteins, KIFs:
Structure, function, and dynamics. *Physiol Rev* **88**: 1089–1118.

Hirokawa, N., Noda, Y., Tanaka, Y., and Niwa, S. (2009) Kinesin superfamily motor proteins and
intracellular transport. *Nat Rev Mol Cell Biol* **10**: 682–696
https://www.nature.com/articles/nrm2774.

Hu, L., Hu, H., and Li, Z. (2012) A kinetoplastid-specific kinesin is required for cytokinesis and for
maintenance of cell morphology in *Trypanosoma brucei. Mol Microbiol* **83**: 565–578
https://pubmed.ncbi.nlm.nih.gov/22168367/.

Ivens, A.C., Peacock, C.S., Worthey, E.A., Murphy, L., Aggarwal, G., Berriman, M., *et al.* (2005) The
genome of the kinetoplastid parasite, *Leishmania major. Science* **309**: 436–442
https://pubmed.ncbi.nlm.nih.gov/16020728/.

Jentzsch, J., Sabri, A., Speckner, K., Lallinger-Kube, G., Weiss, M., and Ersfeld, K. (2020) Microtubule
polyglutamylation is important for regulating cytoskeletal architecture and motility in *Trypanosoma*
brucei. J Cell Sci **133**.

Kaurov, I., Vancová, M., Schimanski, B., Cadena, L.R., Heller, J., Bílý, T., *et al.* (2018) The Diverged
Trypanosome MICOS Complex as a Hub for Mitochondrial Cristae Shaping and Protein Import. *Curr*
Biol **28**: 3393-3407.e5.

Kim, D.I., Birendra, K.C., Zhu, W., Motamedchaboki, K., Doye, V., and Roux, K.J. (2014) Probing
nuclear pore complex architecture with proximity-dependent biotinylation. *Proc Natl Acad Sci U S A*
111: 2453–2461.

Kim, D.I., Jensen, S.C., Noble, K.A., Kc, B., Roux, K.H., Motamedchaboki, K., and Roux, K.J. (2016) An
improved smaller biotin ligase for BioID proximity labeling. *Mol Biol Cell* **27**: 1188–1196.

Kohl, L., Sherwin, T., and Gull, K. (1999) Assembly of the paraflagellar rod and the flagellum
attachment zone complex during the *Trypanosoma brucei* cell cycle. *J Eukaryot Microbiol* **46**: 105–
109 https://pubmed.ncbi.nlm.nih.gov/10361731/.

Kramer, S., Queiroz, R., Ellis, L., Webb, H., Hoheisel, J.D., Clayton, C., and Carrington, M. (2008) Heat
shock causes a decrease in polysomes and the appearance of stress granules in trypanosomes
independently of eIF2α phosphorylation at Thr169. *J Cell Sci* **121**: 3002–3014
https://pubmed.ncbi.nlm.nih.gov/18713834/.

Lacomble, S., Vaughan, S., Gadelha, C., Morphew, M.K., Shaw, M.K., McIntosh, J.R., and Gull, K.
(2009) Three-dimensional cellular architecture of the flagellar pocket and associated cytoskeleton in
trypanosomes revealed by electron microscope tomography. *J Cell Sci* **122**: 1081–1090
http://jcs.biologists.org/cgi/content/full/122/8/1081/DC1.

Lacomble, S., Vaughan, S., Gadelha, C., Morphew, M.K., Shaw, M.K., McIntosh, J.R., and Gull, K.
(2010) Basal body movements orchestrate membrane organelle division and cell morphogenesis in
Trypanosoma brucei. *J Cell Sci* **123**: 2884–2891.

Li, Z., Umeyama, T., and Wang, C.C. (2008) The chromosomal passenger complex and a mitotic
kinesin interact with the tousled-like kinase in trypanosomes to regulate mitosis and cytokines. *PLoS*
One **3** https://pubmed.ncbi.nlm.nih.gov/19043568/.

Lott, K., Li, J., Fisk, J.C., Wang, H., Aletta, J.M., Qu, J., and Read, L.K. (2013) Global proteomic analysis
in trypanosomes reveals unique proteins and conserved cellular processes impacted by arginine
methylation. *J Proteomics* **91**: 210–225 https://pubmed.ncbi.nlm.nih.gov/23872088/.

Marx, A., Hoenger, A., and Mandelkow, E. (2009) Structures of kinesin motor proteins. *Cell Motil*
Cytoskeleton **66**: 958–966 https://pubmed.ncbi.nlm.nih.gov/19530174/.

Moreira, B.P., Fonseca, C.K., Hammarton, T.C., and Baqui, M.M.A. (2017) Giant FAZ10 is required for
flagellum attachment zone stabilization and furrow positioning in *Trypanosoma brucei*. *J Cell Sci* **130**:
1179–1193.

Morgan, G.W., Goulding, D., and Field, M.C. (2004) The single dynamin-like protein of *Trypanosoma*
brucei regulates mitochondrial division and is not required for endocytosis. *J Biol Chem* **279**: 10692–
701 http://dx.doi.org/10.1074/jbc.M312178200.

Morriswood, B., Havlicek, K., Demmel, L., Yavuz, S., Sealey-Cardona, M., Vidilaseris, K., *et al.* (2013)
Novel bilobe components in *Trypanosoma brucei* identified using proximity-dependent biotinylation.
Eukaryot Cell **12**: 356–367.

Poon, S.K., Peacock, L., Gibson, W., Gull, K., and Kelly, S. (2012) A modular and optimized single
marker system for generating *Trypanosoma brucei* cell lines expressing T7 RNA polymerase and the
tetracycline repressor. *Open Biol* **2**: 110037 https://pubmed.ncbi.nlm.nih.gov/22645659/.

Pyrih, J., Rašková, V., Škodová-Sveráková, I., Pánek, T., and Lukeš, J. (2020) ZapE/Afg1 interacts with
Oxa1 and its depletion causes a multifaceted phenotype. *PLoS One* **15**: e0234918
https://pubmed.ncbi.nlm.nih.gov/32579605/.

Ralston, K.S., Lerner, A.G., Diener, D.R., and Hill, K.L. (2006) Flagellar motility contributes to
cytokinesis in *Trypanosoma brucei* and is modulated by an evolutionarily conserved dynein
regulatory system. *Eukaryot Cell* **5**: 696–711 https://pubmed.ncbi.nlm.nih.gov/16607017/.

Redmond, S., Vadivelu, J., and Field, M.C. (2003) RNAit: An automated web-based tool for the
selection of RNAi targets in *Trypanosoma brucei*. *Mol Biochem Parasitol* **128**: 115–118.

Robinson, D.R., Sherwin, T., Ploubidou, A., Byard, E.H., and Gull, K. (1995) Microtubule polarity and

dynamics in the control of organelle positioning, segregation, and cytokinesis in the trypanosome
cell cycle. *J Cell Biol* **128**: 1163–1172 https://pubmed.ncbi.nlm.nih.gov/7896879/.

Rotureau, B., Subota, I., and Bastin, P. (2011) Molecular bases of cytoskeleton plasticity during the
Trypanosoma brucei parasite cycle. *Cell Microbiol* **13**: 705–716.

Schenk, R., Bachmaier, S., Bringaud, F., and Boshart, M. (2021) Efficient flavinylation of glycosomal
fumarate reductase by its own ApbE domain in *Trypanosoma brucei*. *FEBS J*
https://doi.org/10.1111/febs.15812 https://pubmed.ncbi.nlm.nih.gov/33755328/.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., *et al.* (2012) Fiji: An
open-source platform for biological-image analysis. *Nat Methods* **9**: 676–682
https://www.nature.com/articles/nmeth.2019.

Sheriff, O., Lim, L.F., and He, C.Y. (2014) Tracking the biogenesis and inheritance of subpellicular
microtubule in *Trypanosoma brucei* with inducible YFP-α-tubulin. *Biomed Res Int* **2014**.

Sherwin, T., and Gull, K. (1989) Visualization of detyrosination along single microtubules reveals
novel mechanisms of assembly during cytoskeletal duplication in trypanosomes. *Cell* **57**: 211–221
https://pubmed.ncbi.nlm.nih.gov/2649249/.

Sherwin, T., Schneider, A., Sasse, R., Seebeck, T., and Gull, K. (1987) Distinct localization and cell
cycle dependence of COOH terminally tyrosinolated α-tubulin in the microtubules of *Trypanosoma*
brucei brucei. *J Cell Biol* **104**: 439–446 https://pubmed.ncbi.nlm.nih.gov/3546334/.

Sinclair, A.N., and Graffenried, C.L. de (2019) More than Microtubules: The Structure and Function of
the Subpellicular Array in Trypanosomatids. *Trends Parasitol* **35**: 760–777
https://pubmed.ncbi.nlm.nih.gov/31471215/.

Sunter, J.D., and Gull, K. (2016) The Flagellum Attachment Zone: 'The Cellular Ruler' of Trypanosome
Morphology. *Trends Parasitol* **32**: 309–324 https://pubmed.ncbi.nlm.nih.gov/26776656/.

Urbaniak, M.D., Guther, M.L.S., and Ferguson, M. a J. (2012) Comparative SILAC proteomic analysis
of *Trypanosoma brucei* bloodstream and procyclic lifecycle stages. *PLoS One* **7**: e36619
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3344917&tool=pmcentrez&rendertype
=abstract.

Varga, V., Moreira-Leite, F., Portman, N., and Gull, K. (2017) Protein diversity in discrete structures at
the distal tip of the trypanosome flagellum. *Proc Natl Acad Sci U S A* **114**: E6546–E6555
https://www.pnas.org/content/114/32/E6546.

Wei, Y., Hu, H., Lun, Z.-R., and Li, Z. (2013) The Cooperative Roles of Two Kinetoplastid-Specific
Kinesins in Cytokinesis and in Maintaining Cell Morphology in Bloodstream Trypanosomes. *PLoS One*
8: e73869 https://dx.plos.org/10.1371/journal.pone.0073869.

Wheeler, R.J., Gull, K., and Sunter, J.D. (2019) Coordination of the cell cycle in trypanosomes. *Annu*
Rev Microbiol **73**: 133–154 https://pubmed.ncbi.nlm.nih.gov/31500537/.

Wheeler, R.J., Scheumann, N., Wickstead, B., Gull, K., and Vaughan, S. (2013) Cytokinesis in
Trypanosoma brucei differs between bloodstream and tsetse trypomastigote forms: implications for
microtubule-based morphogenesis and mutant analysis. *Mol Microbiol* **90**: 1339–1355
http://www.ncbi.nlm.nih.gov/pubmed/24164479.

Wickstead, B., Carrington, J.T., Gluenz, E., and Gull, K. (2010a) The expanded kinesin-13 repertoire of
trypanosomes contains only one mitotic kinesin indicating multiple extra-nuclear roles. *PLoS One* **5**:
e0015020 https://pubmed.ncbi.nlm.nih.gov/21124853/.

Wickstead, B., Ersfeld, K., and Gull, K. (2002) Targeting of a tetracycline-inducible expression system

to the transcriptionally silent minichromosomes of *Trypanosoma brucei*. *Mol Biochem Parasitol* **125**:
211–6 http://www.ncbi.nlm.nih.gov/pubmed/12467990.

Wickstead, B., and Gull, K. (2006) A "Holistic" Kinesin Phylogeny Reveals New Kinesin Families and
Predicts Protein Functions. *Mol Biol Cell* **17**: 1734–1743.

Wickstead, B., Gull, K., and Richards, T.A. (2010b) Patterns of kinesin evolution reveal a complex
ancestral eukaryote with a multifunctional cytoskeleton. *BMC Evol Biol* **10**:
https://doi.org/10.1186/1471-2148-10-110.

Woods, A., Sherwin, T., Sasse, R., MacRae, T.H., Baines, A.J., and Gull, K. (1989) Definition of
individual components within the cytoskeleton of *Trypanosoma brucei* by a library of monoclonal
antibodies. *J Cell Sci* **93 (Pt 3)**: 491—500 http://europepmc.org/abstract/MED/2606940.

Woodward, R., and Gull, K. (1990) Timing of nuclear and kinetoplast DNA replication and early
morphological events in the cell cycle of *Trypanosoma brucei*. *J Cell Sci* **95 (Pt 1)**: 49–57
http://www.ncbi.nlm.nih.gov/pubmed/2190996.

Zhou, Q., An, T., Pham, K.T.M., Hu, H., and Li, Z. (2018) The CIF1 protein is a master orchestrator of
trypanosome cytokinesis that recruits several cytokinesis regulators to the cytokinesis initiation site.
J Biol Chem **293**: 16177–16192 https://pubmed.ncbi.nlm.nih.gov/30171070/.

Zoltner, M., Pino, R.C. del, and Field, M.C. (2020) Sorting the Muck from the Brass: Analysis of
Protein Complexes and Cell Lysates. In *Methods in Molecular Biology*. Humana Press Inc., pp. 645–
653 https://pubmed.ncbi.nlm.nih.gov/32221947/.

FIGURE LEGENDS

Fig. 1: Domain architecture of TbPH1 and TbKifX2. Schematic depiction includes phosphorylation and
arginine methylation sites. The consensus Walker A motif is shown in between the two schemata for
comparison of the respective motifs within the kinesin motor domains of TbPH1 and TbKifX2. Key
residues in ATP hydrolysis shown in red and mutations in purple. The likely inactive state of the
TbPH1 'kinesin motor domain' is highlighted by striped pattern. CC: coiled coil domain, PH: pleckstrin
homology domain, HDL: homeodomain-like fold.

Fig. 2: TbPH1-V5 fractionates with the cytoskeleton. **(A)** Western blot of-trypanosome cell fractions.
Fractionation procedure is summarized in S1 Fig. Antibodies used were anti-V5 and anti-SCD6
(Kramer *et al.*, 2008). T: total lysate, SN: supernatant, P: pellet. **(B)** IFA of on an aliquot of fraction P2
placed on slides for fluorescence microscopy. Cells were labelled with anti-V5 antibody (white) and
DAPI (magenta). The scale bar corresponds to 5 µm.

Fig. 3: TbPH1 interacts with TbKifX2, a kinetoplastid-specific kinesin. **(A)** Western blot verifying
immunoprecipitation of TbPH1-V5. The blot was probed with anti-V5 and anti-α-tubulin antibodies.
The parental cell line SmOxP9 was used as a control. T: total lysate, SN3: supernatant 3, FT: flow
through, E: elution. **(B)** SyproRuby-stained gel of elution fractions of TbPH1-V5 IP and mock control
(SmOxP9). B1 and B2: bands excised and by mass spectrometry identified to be TbPH1 and TbKifX2,
respectively. Asterisks: band corresponding to denatured immunoglobulin light chain that was
incidentally eluted. **(C)** Western blot showing interaction of TbPH1-V5 and TbKifX2-HA. TbKifX2-HA
was immunoprecipitated in the same way as PH1-V5. T: total lysate, SN: supernatant, P: pellet, FT:
flow through, E: elution. **(D)** IFA on whole cells detecting TbPH1-V5 (white) and counterstained with
DAPI to visualize DNA (magenta). Arrowheads denote increased signal in the area along the proximal
part of the flagellum. Asterisks demark an increased signal at the anterior cell end. Merge of DAPI

and Phase contrast images shown on bottom. Scale bar, 10 μm. **(E)** IFA on whole cells detecting
TbKifX2-HA (white). Images depicted as in **D**.

Fig. 4: The proximal proteome of TbPH1. **(A)** IFA on whole cells showing localisation of TbPH1-
BioID2-HA (white) and biotinylated proteins (white) and counterstained with DAPI (magenta).
Arrowheads denote signal in vicinity of kinetoplast, near the proximal part of the flagellum. DIC on
right to show observed cell. Scale bar, 5 μm. **(B)** IFA as in **A** of parental cell line lacking TbPH1-BioID2-
HA. **(C)** Western blot (top panel) and fluorescently-detected proteins resolved on SDS-PAGE as a
loading control (bottom panel, stain-free gel) of a representative BioID experiment. The blot was
probed with anti-HA to confirm the presence of TbPH1-BioID2-HA in the respective fractions. E:
extract, SN: supernatant, F: flow through, B: boiled beads. **(D)** Volcano plot with -10 log t-test p value
plotted against the t-test difference between TbPH1-BioID2-HA sample and mock (SmOxP9 cells)
from three biological replicates. Statistically significant hits are found above the curve in the top
right quadrant and their identity is indicated. Accession numbers are given with additional
annotations in parentheses; these are color coded based on their localization determined by TrypTag
(Dean *et al.*, 2017). **(E)** Heat map of SAINTq score (*i.e.* P value) of proteins interacting with TOEFAZ1
(Hilton *et al.*, 2018), the relative enrichment of proteins also found in the TbPH1 proximal proteome
are indicated. a:Tb927.3.2490 (TbPH1), b: Tb927.8.1950 (ApbE-like domain), c: Tb927.9.15470
(TbKifX2), d: Tb927.3.4720/60 (DLP1/2). PG, protein groups. **(F)** Venn diagram comparing the
proximity proteome of TbPH1 with those of the TOEFAZ (as depicted in **E**), the hook complex protein
TbMorn1 (all 79 biotinylated proteins (Morriswood *et al.*, 2013)), and the 11 high confidence
proteins in the TbSpef1, a MtQ binding protein (Dong *et al.*, 2020). Proteins exclusive to the TbPH1
proximity proteome are highlighted in green while magenta highlights those that are found to also
be in proximity with TOEFAZ.

Fig. 5: Localization of cytoskeleton-associated TbKifX2 and TbPH1. **(A)** IFA on extracted cytoskeletons
probing localisation of TbPH1-V5 (green) and TbKifX2-HA (white). **(B)** IFA on extracted cytoskeletons
probing co-localisation of TbKifX2-HA (white) with the FAZ (green), the latter labelled with L3B2
antibody. **(C)** IFA on extracted cytoskeletons probing co-localisation of TbKifX2-HA (white) with basal
bodies (green), the latter labelled with YL1/2 antibody **(D)** IFA on extracted cytoskeletons probing
co-localisation of TbKifX2-HA (white) with the MtQ (green), the latter labelled with 1B41 antibody. In
all panels, merge images depict co-localization in magenta and DIC on right showing observed
extracted cytoskeleton. Scale bar, 5 μm.

Fig 6: TbKifX2 and TbPH1 localize to the MtQ. Ultrastructure expansion microscopy images depicted
as 3D projections (A-D) and representative confocal image sections used to make projections (E-F) of
T. brucei expressing TbPH1-V5 and TbKifX2-HA. (A) 3D reconstruction of the area of the cell in the
vicinity of the basal body and flagellum exit from the flagellar pocket looking from outside the MT
corset. Top shows signal from anti-V5 (green) and anti-acetylated α-tubulin (white) antibodies.
Bottom is the same image of anti-acetylated α-tubulin signal only with relevant structures marked.
Tick marks at 5 μm increments (A-C). (B) Same as A but showing anti-HA antibody signal (green) and
a view shown from inside the MT corset. (C) 3D reconstruction of the posterior part of the cell.
Signal colors correspond to A. (D) 3D reconstruction of whole dividing *T. brucei* cell. Signal colors
correspond to B. Box highlights region of interest shown in F. Tick marks at 10 μm increments. (E)
Confocal images from the vicinity of the basal body showing from left to right signal of anti-V5 and
anti-acetylated α-tubulin antibodies(both white) plus an overlay of both signals with V5 signal in
green. Left image shows 5 μm scale bar. Middle image marks relevant structures. (F) Same as in E
except two sections being shown, focusing on new (Top) and old MtQ (Bottom), and the left images

showing anti-HA antibody signal. Solid arrowheads point to overlay of epitope antibody signal and
MtQ. Hollow arrowheads point to representative epitope antibody signals found in the cytosol.

Fig. 7: Depletion of TbPH1 and TbKifX2 affects posterior end remodelling. **(A)** Bar chart showing
population doubling times for TbPH1, TbKifX2 and TbPH1/TbKifX2 depletion cell lines. -/+: -/+
doxycycline to induce RNAi, the average difference in hours from three experiments is shown above
the respective bars. **(B)** Western blot of cellular fractionation experiment following TbKifX2 RNAi.
The blot was probed with anti-HA (to detect TbKifX2), anti-V5 (to detect TbPH1) and anti-α-tubulin
as a loading control. WCL: whole cell lysate. **(C)** Phase contrast images of TbPH1/TbKifX2 RNAi cells. +
tet: induced for 5 days. The scale bar corresponds to 5 μm. Arrows indicate morphological defects.
(D) Scanning electron microscopy of TbPH1/TbKifX2 RNAi cells. + tet: induced for 5 days. Arrows
pointing to posterior end protrusions. The scale bar corresponds to 10 μm.

S1 Fig.: Schematic depiction of microtubule sieving technique. According to (Fritz *et al.*, 2015).

S2 Fig.: Schematic depiction of proximity-dependent biotinylation (BioID). According to (Morriswood
et al., 2013). Only steps in black were performed in our study.

S3 Fig.: Wider field of view of multiple extracted cytoskeletons co-localizing TbKifX2 and TbPH1 to
basal body **(C)**, FAZ **(D)** and MtQ **(E)**. Images depicted as in Fig 5 with scale bar corresponding to 5
μm. **(A)** Co-localization (merge, magenta) of TbPH1-V5 (green) and TbKifX2-HA (white). **(B)** Parental
cell line lacking TbPH1-V5 and TbKifX2-HA, incubated with antibodies recognizing V5 and HA
epitopes used to acquire other IFA images.

S4 Fig.: Ultrastructure expansion microscopy of a cell labelled with anti-ClpGM6 (to detect the FAZ,
green) and anti-acetylated α tubulin antibody C3B9 (grey). Only the region around the pocket and
flagellum exit shown.

S5 Fig.: TbPH1/TbKifX2 RNAi. **(A)** Growth curves of TbPH1 and TbKlfX2 RNAi cell lines. Filled circles:
TbKifX2 − tet, open circles: TbKlfX2 + tet, filled triangles: TbPH1 − tet, open triangles: TbPH1 + tet.
Standard deviation of measurements at each time point (n=3) is not visible at this scale. Insets show
western blots probed with anti-HA (to detect TbKifX2), anti-V5 (to detect TbPH1) and anti-α-tubulin
as a loading control. **(B)** Growth curve of TbPH1/TbKifX2 RNAi cell line. Filled circles: TbPH1/TbKifX2
− tet, open circles: TbPH1/TbKifX2 + tet. Standard deviation of measurements at each time point
(n=3) is not visible at this scale. Inset depicts western blot showing depletion of TbPH1/TbKifX2.
Samples were taken at the same time as the growth curve in B was performed. The blot was probed
with anti-HA (to detect TbKifX2), anti-V5 (to detect TbPH1) and anti-α-tubulin as a loading control.
(C) Bar chart depicting different cell cycle phases during TbPH1/TbKifX2 RNAi time course. DAPI
counts were performed in triplicate with at least 200 cells counted per time point. N: nucleus, K:
kinetoplast. **(D)** IFA of TbPH1/TbKifX2 RNAi cells. + tet: induced for 5 days. Extracted cytoskeletons
were labelled with L3B2 (recognising the FAZ filament, white), mAB 1B41 (recognising the MtQ,
white) and DAPI (magenta). Scale bar, 5 μm.

S6 Fig.: Transmission electron microscopy and IFA of TbPH1/TbKifX2 RNAi cell line. Cells were
induced for 5 days. **(A)** Image depicting posterior end protrusions (arrow). Scale bar, 1 μm. **(B)** Close-
up of protrusion encapsulated by microtubule corset (arrowheads). The scale bar corresponds to 500
nm. **(C)** IFA on cytoskeletons probing effects of TbPH1/TbKifX2 depletion post-translational
modifications of tubulin. Cells were counterstained with GT335 (red, staining glutamyl side chains)
and YL1/2 (green, staining tyrosinated α-tubulin). The scale bar corresponds to 5 μm. **(D)** Western
blots taken after 3 to 5 days (d) of growth in presence (+) or absence (-) of doxycycline probed with
antibodies recognizing GT335, YL1/2 and α-tubulin, the last as a loading control.

S1 Dataset: Mass spectrometry analysis of excised bands B1 and B2 from Fig. 3B.

S2 Dataset: Mass spectrometry data of TbPH1 BioID experiment. Data with and without imputation
and filtered for Andromeda scores >20 and presence of >1 peptide, which was used to
generate the volcano plot in Fig. 4D.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

1044

Fig. 5

Fig. 6

1045

Fig. 7

Suppl. Fig. 1

1066

1067

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

Suppl. Fig. 2

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

Suppl. Fig. 3

Suppl. Fig. 4

1100

1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126

Suppl. Fig. 5

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

Suppl. Fig. 6

1148
1149