Enhancement of Chikungunya virus genome replication in

mammalian cells at a sub-physiological temperature

Jinchao Guo[1,2], Aaron E. Lin[2], Chetan Aditya[2#], Alexander Ploss[2]*

and Mark Harris[1]*

[1] School of Molecular and Cellular Biology, Faculty of Biological Sciences and

Astbury Centre for Structural Molecular Biology, University of Leeds, Leeds, LS2 9JT,

United Kingdom.

[2] Department of Molecular Biology, Lewis Thomas Laboratory, Princeton University,

Washington Road, Princeton, NJ, USA.

Present address: Department of Bioengineering and Therapeutic Science,

University of California San Francisco, San Francisco, CA 94143, USA

* Joint corresponding authors, e-mail: m.harris@leeds.ac.uk, aploss@princeton.edu

Word counts: Abstract: 250, main text: 8166

9 figures, 7 supplementary figures.

Keywords: Chikungunya virus, alphavirus; subgenomic replicon; mammalian cells;

sub-physiological temperature; interferon-stimulated genes, stress granules

Author contributions: JG: conceptualization, formal analysis, investigation,

visualization and writing.   AEL: formal analysis, investigation, visualization and

writing. CA: formal analysis, investigation, visualization.   AP: funding acquisition,

supervision, visualization, writing. MH: conceptualization, funding acquisition,

supervision, visualization and writing.

Abstract

Chikungunya virus (CHIKV) is an alphavirus transmitted by *Aedes* mosquitoes, causing fever, rash and arthralgia in mammals. The function of the CHIKV non-structural protein 3 (nsP3) remains enigmatic. Building on previous studies (Gao *et al*, 2019) [1], we generated a panel of mutants in a conserved and surface-exposed cluster in the nsP3 alphavirus unique domain (AUD) and tested their replication using a sub-genomic replicon (SGR). Three of these SGR mutants replicated well in mosquito cells but poorly in mammalian cells. We observed that this difference was due to culture temperature (mosquito cells: 28°C; mammalian cells: 37°C), as the mutants exhibited no replication defect in mammalian cells grown at a sub-physiological temperature (28^{o}C). Similar phenotypes were observed for infectious CHIKV and the closely related O'Nyong Nyong virus. Intriguingly, the wildtype SGR replicated more efficiently in mammalian cells at 28^{o}C compared to 37^{o}C. To explore the mechanism behind this difference, we focused on two known antiviral pathways: interferon-stimulated genes (ISGs) and stress granules (SGs). SGR replication was concomitant with increased expression of ISGs at 37°C, but this response was impaired at 28^{o}C. We also observed enhanced recruitment of the SG component G3BP1 into cytoplasmic sites of viral genome replication at 28^{o}C. These findings may have real-world implications as when a mosquito bites a mammal, the virus first infects cells in peripheral tissues which are often at sub-physiological temperatures. We propose that alphaviruses such as CHIKV have evolved mechanisms to both promote viral genome replication and concomitantly limit antiviral responses in these cells.

Significance statement

Chikungunya virus (CHIKV) is a re-emerging arbovirus that is transmitted by *Aedes*

mosquitos and poses epidemic threats. Arboviruses must be able to grow efficiently

in both the mosquito vector and the mammalian host, which have different body

temperatures. Following a mosquito bite, the first cells infected are typically in the

skin and are at sub-physiological temperatures, ie lower than the internal body

temperature (37^{o}C). Here we show that CHIKV grows more efficiently in mammalian

cells cultured at 28°C, compared to 37°C. We investigated the mechanism behind

this increase in growth efficiency and demonstrated that at the lower temperature the

production of antiviral genes was impaired. Additionally, proteins that respond to

cellular stress and are needed for virus replication were more abundantly recruited to

sites in the cell where the virus was replicating. We propose that CHIKV has

evolved mechanisms to promote its replication in mammalian cells at sub-

physiological temperatures to facilitate infection of mammals via a mosquito bite.

Introduction

Chikungunya virus (CHIKV) belongs to the *alphavirus* genus in the *Togaviridae*

family [2]. It is an arbovirus that is transmitted by mosquitoes and causes fever, rash

and arthralgia with infrequent long-term morbidity and, in rare cases, mortality in

mammals [3]. CHIKV was first isolated in Tanzania in 1952-1953 [4, 5], and has

subsequently spread across the world, including Africa, Europe, Asia and America [6,

7]. CHIKV is mainly transmitted by the mosquito species *Aedes (Ae.) aegypti* and *Ae.*

albopictus, and the latter has become the main vector for virus spread in Asia [8].

Both field and laboratory experiments demonstrated that these two mosquito species

only survive between lower and upper temperature thresholds [9]. Models and

human cases showed that transmission occurs between 18°C and 34°C with

maximal transmission occurring in a range from 26-29°C [10].

CHIKV has a positive-sense, single-stranded RNA genome, 11.5 kb in length, which

is both capped and polyadenylated (Fig. S1A). It contains two open reading frames

(ORFs). The first ORF encodes the non-structural proteins (nsP1-nsP4) and is

translated directly from full-length genomic RNA. ORF2 encodes the structural

proteins (capsid, E1-E3 and 6K) and is translated from a sub-genomic RNA. The

non-structural proteins are required for the synthesis of positive and negative sense

full-length genomic RNA as well as the sub-genomic RNA. Biochemical analysis

indicates specific functions of the non-structural proteins: nsP1 has methyl- and

guanyl-transferase activities, nsP2 exhibits helicase and protease activities and nsP4

is the RNA-dependent RNA polymerase [11].

The function of nsP3 remains poorly defined [12], despite being essential for genome

replication. CHIKV nsP3 consists of three domains: a macro-domain at the N-

terminus, a central alphavirus unique domain (AUD) and a hypervariable domain

(HVD) at the C-terminus (Fig. S1A). The macro-domain exhibits RNA binding, ADP-

ribose binding and ADP-ribosylhydrolase activities [13-15].   The HVD is highly

phosphorylated and interacts with a number of host proteins [16].   Despite high

levels of conservation across alphaviruses, AUD function remains elusive.   In this

context, we showed that the AUD binds to viral RNA and plays a role in virus

assembly by stimulating transcription of the sub-genomic RNA and concomitant

translation of ORF2 [1].   In addition, recent cryo-EM analysis revealed that nsP3

assembles into tubular structures, driven by a helical arrangement of the AUD [17].

AUD mutations that disrupted these assemblies also impaired virus replication.

nsP3 also plays multiple roles in rewiring host cell biology, especially altering stress

granules (SGs) [18]. SGs are cytoplasmic, non-membranous aggregates formed by

eukaryotic cells in response to cell stress and translation inhibition. Typical SGs

contain silent mRNAs, translation factors, small ribosomal subunits, and various

mRNA-binding proteins.   Despite varied compositions resulting from different

stresses [19], the RasGAP SH3-domain binding proteins (G3BP1/2) are key

components required for formation of all SGs [20-22]. However, alphavirus infection

blocks SG formation: the CHIKV nsP3 hypervariable domain interacts with G3BP1/2

and sequesters them into cytoplasmic foci which are the sites of genome replication

[23, 24] [18]. Recent work also demonstrated that the CHIKV nsP3 macro-domain

can alter the composition of SGs or disassemble them entirely [25].   Therefore,

CHIKV nsP3 plays a key role in both blocking SG formation and altering their

composition in order to promote genome replication.

In addition to altering SGs, CHIKV must also limit the interferon (IFN)-mediated

innate immunity, which provides the first line against invading pathogens. As with

many RNA viruses, CHIKV genome replication produces double-stranded RNA

which is sensed by infected cells, which in turn produce and secrete type I IFNs to

induce an antiviral state in neighboring cells [26]. Cells that sense type I IFNs initiate

the JAK-STAT signaling pathway and the downstream production of IFN-stimulated

genes (ISGs) [27]. This leads to antiviral states which are effective in restricting

replication of both DNA and RNA viruses [28]. A number of ISGs have been shown

to suppress infection by the model alphavirus Sindbis virus [29]. Additionally, it has

been shown that in a mouse model production of type I IFN was impaired at sub-

physiological temperatures, and thus CHIKV replication and arthritis was more

pronounced [30, 31].

In this study we sought to investigate the function of the nsP3 AUD in CHIKV

replication in cells from both the vertebrate host and the mosquito vector. Three

AUD mutants (W220A, D249A and Y324A) replicated poorly in mammalian cells but

efficiently in mosquito cells. However, culture temperature confounded this result,

since mammalian cells are typically cultured at 37°C while mosquito cells are

cultured at 28°C. We demonstrate that a wildtype CHIKV sub-genomic replicon

(SGR) replicated more efficiently in mammalian cells at a sub-physiological

temperature, suggesting that the original AUD mutant phenotype was temperature-

dependent rather than species-specific. To explore the mechanism behind

differential replication at 28°C or 37°C, we focused on two known antiviral pathways:

ISGs and SGs. Transcriptomic analysis revealed that ISG expression was impaired

at the lower temperature. Furthermore, analysis of SGs revealed that G3BP1 was

more efficiently recruited to cytoplasmic replication sites at the sub-physiological

temperature. Taken together these observations may explain, at least in part, why

replication of CHIKV SGR in mammalian cells is enhanced at a lower temperature.

Results

Construction and analysis of AUD mutants

In this study we set out to build on our previous work which had identified a role for

two residues (P247 and V248) in the function of the CHIKV nsP3 AUD [1]. Alanine

substitution of both residues disrupted the binding of the AUD to the sub-genomic

promoter, reducing transcription of the sub-genomic RNA and translation of

structural proteins, with concomitant effects on virus assembly. Interestingly, the

P247A/V248A mutation also disrupted the helical arrangement of nsP3 observed by

cryo-EM [17]. To further investigate the phenotype that we observed, we first

generated a series of mutations in surface-exposed and conserved residues

proximal to P247 and V248. To identify such residues, AUD amino acid sequences

from both Old and New World alphaviruses were aligned (Fig. S1B). The nsP2/nsP3

protein structure of Sindbis virus (SINV) has been determined [32], and the AUD

sequences between SINV and CHIKV share high similarity (118 out of 243 residues

are identical). A panel of conserved and surface exposed residues in the AUD were

therefore selected for mutation based on the SINV nsP2/nsP3 protein structure [32]

(Figs. S1C). Seven residues were chosen for further study as they were either

identical or similar between the two viruses. The C-terminal residue of nsP3 (Y324

in nsP3 amino acid numbering) was selected as a control as it is distal to P247/V248

and highly conserved among alphaviruses (Fig. S1B).

These mutations were introduced into a CHIKV dual luciferase sub-genomic replicon

(CHIKV-SGR-D-Luc) (Fig. S1A), which was derived from the ECSA strain LR2006

OPY1 isolate [33, 34]. In this construct, renilla luciferase (R-Luc) is fused within the

nsP3 hypervariable domain in ORF1, and firefly luciferase (FF-Luc) replaces the

structural proteins in ORF2.   The activity of R-Luc is thus a marker of ORF1

translation from the full-length RNA: at early time points R-Luc levels reflect

transfection efficiency and input RNA translation, at later times R-Luc is also

dependent upon genome replication.   As ORF2 is translated from a sub-genomic

RNA (sgRNA) which is templated from the negative strand replicative intermediate,

FF-Luc expression is dependent upon genome replication but will also reflect

translation from the sgRNA. A polymerase-inactive mutant (GDD>GAA in nsP4) was

used as a negative control.

AUD mutants exhibit different replication phenotypes in mammalian and

mosquito cells

CHIKV mainly induces arthritis and joint pain [3] and infects muscle cells. We

therefore initially chose to investigate the effects of the AUD mutants on CHIKV

genome replication in C2C12 cells, an immortalized mouse myoblast cell line widely

used in the study of CHIKV [35]. As expected, wildtype (WT) CHIKV-SGR-D-Luc

showed a very high level of replication in C2C12 cells, with FF-Luc levels peaking at

12 h post transfection (h.p.t) and increasing over 1000-fold between 4-12 h.p.t. (Fig

S2A). AUD mutants T218Q, M219Q, P221A, C246A and A251Q exhibited similar

levels of replication as compared to WT (Fig S2A), however, W220A, D249A and

Y324A showed significantly lower levels of replication compared to WT (Fig S2A).

As CHIKV is transmitted by mosquitoes, we then evaluated the replication capacities

of the AUD mutants in C6/36 and U4.4 cells, both derived from *Ae. albopictus*.

C6/36 cells harbor a frameshift mutation in the Dicer-2 (Dcr2) gene, which leads to

the translation of a truncated Dcr2 protein and a defect in the RNAi response [36].

Consistent with this, WT CHIKV-SGR-D-Luc showed robust replication in both cell

lines, with higher levels of replication in C6/36 than U4.4 cells (Fig. S2B/C).

Interestingly, unlike the replication defects observed in C2C12 cells, all AUD mutants

showed similar levels of replication compared to WT in both C6/36 and U4.4 cells

(Fig. S2B/C). In C6/36 cells levels of FF-luc continued to increase up to 48 h.p.t.

whereas they peaked at 24 h.p.t. in U4.4 cells.

The striking phenotypic differences between mammalian and mosquito cell lines for

the three AUD mutants (W220A, D249A and Y324A) led us to investigate further. A

simple explanation might be that these mutants had reverted to WT in mosquito cells,

as we had previously observed this for the AUD mutant R243A/K245A [1]. This

mutant replicated poorly in mammalian cells but efficiently in mosquito cells, and this

was explained by the fact that R243A/K245A had reverted to WT in mosquito cells

allowing efficient replication. To test this hypothesis, total RNA from U4.4 and C6/36

cells at 72 h.p.t was extracted, the nsP3 coding sequence was amplified by RT-PCR

and Sanger sequenced. Reassuringly, none of the AUD mutants had reverted to WT,

and all three mutants retained the same sequence as the input RNA (Fig. S2D). The

results demonstrated that the ability of these three AUD mutants to replicate in

mosquito cells was not due to reversion. We conclude that the three residues W220,

D249 and Y324 within the CHIKV nsP3 AUD play a role in genome replication in

mammalian cells but not in mosquito cells.

The AUD mutants exhibit a temperature dependent phenotype

We reasoned that a simple explanation for the different phenotypes in mammalian

and mosquito cells might be the temperature difference, as mammalian cells are

cultured at 37°C whereas mosquito cells are cultured at 28°C [37]. To test this, the

replication of WT and mutant CHIKV-SGR-D-Luc were analysed in C2C12 cells

incubated at 28°C. Intriguingly, none of the three mutants exhibited any phenotype

at this lower temperature, and FF-luc (Fig. 1B) levels were indistinguishable from WT

at 28°C. In addition, we noted that at 37°C WT CHIKV-SGR-D-Luc FF-luc levels

reached a plateau at 12 h.p.t. (Fig 1A), but in contrast, at 28°C FF-luc levels

continued to increase (approximately 100-fold) from 12 to 24 h.p.t. (Fig 1B). As FF-

luc levels were similar at both temperatures up to 12 h.p.t. this suggested that initial

rates of replication and/or translation were temperature-independent, however after

12 h.p.t. these rates were limited at 37°C, but this limitation was relieved at the lower

temperature. Translation from the CHIKV SGR at the two temperatures in C2C12

cells was quantified by western blotting for nsP1 expression (Fig. 1C). Reassuringly,

none of the mutants exhibited detectable nsP1 at 37°C whereas a low level of nsP1

was detectable for the WT at this temperature. In contrast, WT and all mutants

exhibited high and equivalent levels of nsP1 at 28°C, consistent with the elevated

levels of replication and translation at this temperature.

To provide further evidence for the impaired replication at 37°C compared to 28°C,

we quantified both positive and negative strand CHIKV RNA over a period of 24 h.p.t.

in C2C12 cells (Fig. 1D). Although the high level of transfected input RNA hampered

quantification of genome replication at early time points (e.g. 4 h.p.t.) a clear

distinction in RNA levels could be seen at later time points. Specifically, the levels of

both positive and negative strand RNA declined at 37°C between 4-24 h.p.t. whereas

at 28°C they continued to increase. By 24 h.p.t. there was at least a 10-fold higher

level of CHIKV RNA at 28°C compared to 37°C.

To confirm that this observation was not restricted to C2C12 cells, we also tested the

replication of WT CHIKV-SGR-D-luc and the three mutants in Vero (African Green

Monkey kidney epithelial) cells. Vero cells are defective in IFN α/β production due to

a 9 Mb deletion in chromosome 12 but express the type 1 IFNAR1/2 receptor and

can therefore respond to exogenous IFN [38, 39]. We again observed increased

reporter activity of both WT and mutants in these cells at the lower temperature (Fig.

S2E) suggesting that this temperature-driven difference in replication did not depend

on intact IFN production.

We conclude that the three AUD mutants (W220A, D249A and Y324A) are

temperature sensitive, exhibiting no defective phenotype in mammalian cells at a

sub-physiological temperature (28°C). However, these experiments also revealed

the novel observation that replication and/or translation of the WT CHIKV SGR was

significantly enhanced at a sub-physiological temperature, and we sought to further

investigate this intriguing phenomenon.

Enhanced replication at sub-physiological temperature is not dependent on

the reporter protein and is not observed for other arboviruses

As the CHIKV-SGR-D-Luc construct contains two reporters, we considered it prudent

to confirm that the phenotype we observed was not influenced by these reporters, or

by the insertion of a reporter into nsP3. To test this, two alternative CHIKV SGRs

were utilized, namely CHIKV-SGR-nsP3-mCh-FF-Luc (in which the R-Luc reporter

was replaced with mCherry) and CHIKV-SGR-FF-Luc (lacking R-Luc and thus

expressing native nsP3) (Fig. S3A). These two SGRs were assayed in parallel with

CHIKV-SGR-D-luc in C2C12 cells at the two temperatures (28°C or 37°C). All three

showed enhanced levels of replication at 28°C compared to 37°C (Fig. S3B). As

described previously (Fig. 1B), this effect was most apparent between 12-24 h.p.t.

These data confirm that enhanced genome replication at sub-physiological

temperature is not dependent on the nature of the nsP3-fused reporter protein.

We reasoned that the enhancement of genome replication at sub-physiological

temperature might be a common feature in other arbovirus species that share with

alphaviruses a mode of transmission from mosquitoes to humans via a peripheral

bite. We therefore tested replication of mini-genome constructs derived from

Bunyamwera virus (BUNV), the type species of the genus Orthobunyavirus

(negative-sense, single-stranded RNA viruses) [40] and an SGR derived from Zika

virus, a positive-sense, single-stranded RNA virus in the *Flaviviridae* family [41]. As

shown in Fig. S3C/D, although both the BUNV mini-genome [42] and the Zika nano-

luc SGR [43] replicated poorly in BSR-T7 and BHK-21 cells respectively, they

exhibited similar levels of genome replication at 28°C or 37°C , indicating that this

phenotype is not a common feature in other arboviruses.

To determine if the temperature sensitivity of the AUD mutants was unique to CHIKV

or conserved within other alphaviruses, we generated the same panel of mutants in

the context of an infectious clone of the closely related alphavirus, O'Nyong Nyong

virus (ONNV). In this clone, the fluorescent reporter ZsGreen was expressed from a

sgRNA transcribed from a second copy of the sub-genomic promoter (ONNV-2SG-

ZsGreen) (Fig. S4A) (kind gift from Andres Merits, University of Tartu).   This

construct enabled rapid assessment of genome replication following transfection, by

quantification of both ZsGreen fluorescence per cell and the number of ZsGreen

positive cells. Although expression of ZsGreen was dependent on genome

replication as production of the sgRNA is templated from the negative strand, the

levels of ZsGreen expression are representative of both replication and translation of

the sgRNA. As shown in Fig. S4B, the three AUD mutants (W220A, D249A and

Y324A) showed a defect in replication at 37°C as judged by a reduction in the

number of ZsGreen positive cells, whereas at 28°C the replication of all three was

indistinguishable from WT. In addition, the absolute levels of ZsGreen per cell for

WT was approximately 2-fold higher at 28°C compared to 37°C, consistent with

enhanced genome replication at the sub-physiological temperature.

AUD mutants remain temperature sensitive in the context of infectious CHIKV

As we had previously identified a novel function of the CHIKV AUD in promoting

virus assembly, we wished to determine whether the phenotypes we had observed in

the context of CHIKV SGR were also seen in the context of infectious virus. The

three AUD mutations (W220A, D249A and Y324A) were therefore generated in the

context of the 'Integration of Chikungunya virus research' (ICRES) CHIKV infectious

clone, also derived from the ECSA strain LR2006 OPY1 isolate [34]. *In vitro*

transcribed RNA for each construct was transfected into C2C12 cells, cells were

grown at either 28°C or 37°C, and released virus was harvested at 48 h post

infection (h.p.i.). All supernatants were titrated by plaque assay on BHK-21 cells

grown at 32°C. As shown in Fig. 2A, the three mutants produced significantly less

infectious virus than WT at 37°C (between 10^3 and 10^4 reduction in titre). At 28°C,

unlike the SGR data where the mutants were indistinguishable from WT, the mutants

exhibited significantly reduced titres although this phenotype was much less

pronounced than at 37°C (between 5-60 fold reduction at 28°C). Sequence analysis

of ICRES-CHIKV AUD mutants at 28°C or 37°C indicated that no reversion had

occurred at either temperature (data not shown). Unlike the marked differences in

genome replication observed using SGRs, there was only a modest increase (3-fold)

in virus production at 28°C compared to 37°C. This suggested that the higher level

of genome replication at 28°C did not lead to a concomitant increase in assembly

and release of infectious virus, however, we noted that this result was obtained

following transfection of *in vitro* transcribed RNA. We therefore performed a one-

step growth curve analysis, evaluating virus production and genome replication over

a time-course of 96 h following infection of C2C12 cells at a low MOI. As shown in

Fig. 2B, the production of infectious virus was delayed at 28°C compared to 37°C,

however, the maximal titre reached was 5-fold higher (1.8 x 10^8 at 72 h.p.i.

compared to 3.6 x 10^7 at 24 h.p.i.). A similar delay was observed when levels of

intracellular positive and negative strand CHIKV RNA were quantified (Fig. 2C). As

there was no marked difference in the maximal level of positive strand RNA at both

temperatures this may be due to a higher level of packaging of the positive strand

RNA into nascent infectious virus and release from the cell at the lower temperature

(Fig. 2B). In contrast, the level of negative strand RNA did attain a higher value.

These data are consistent with the results obtained using SGRs, namely that at later

time points after infection the rate of genome replication was limited at 37°C but this

limitation was relieved at the lower temperature. However, it should be noted that

the differences observed in the context of infectious virus were not as marked as

those seen in the SGR experiments, perhaps reflecting other rate limiting cell factors

or processes involved in assembly and release of infectious virus.

Temperature shift assay reveals that early events in SGR-D-Luc replication are

not enhanced at sub-physiological temperature

Previous studies have investigated temperature-sensitive mutants of alphaviruses by

use of a temperature shift assay [44-46]. To further explore the role of temperature

in controlling CHIKV genome replication, a temperature shift assay was conducted in

C2C12 cells. WT CHIKV-SGR-D-Luc RNA was transfected into C2C12 cells and

incubated either at 37°C or 28°C for 12 h, with and without a subsequent

temperature shift from 12-24 h.p.t. (28°C to 37°C (designated S*), or 37°C to 28°C

(designated S#)) (Fig. 3A). As seen previously (Fig. 1A), replication of WT CHIKV-

SGR-D-Luc (as assessed by FF-Luc levels) up to 12 h.p.t. proceeded at broadly

similar rates at both temperatures (Fig. 3B), and between 12-24 h.p.t. replication

increased ~100-fold at 28°C but did not increase at 37°C. However, when the

temperature was shifted from 28°C to 37°C at 12 h.p.t. (S*), the replication increased

only ~20-fold between 12-24 h.p.t. (Fig. 3B, left). The opposite effect was seen

when transfected cells were first incubated at 37°C: replication reduced slightly

between 12-24 h.p.t. at 37°C, however, when the temperature was shifted from 37°C

to 28°C at 12 h.p.t., replication increased 5-fold between 12-24 h.p.t. (Fig. 3B, right).

Importantly, both cell growth assessed by quantification of viable cells (Fig. 3C), and

metabolic activity assessed by MTT assay (Fig. 3D) were higher at 37°C compared

to 28°C. Temperature shifts elicited the expected changes in both cell number and

metabolic activity. These data confirmed that the enhanced replication observed at

sub-physiological temperature cannot be explained by increased cell growth and

metabolic activity. In fact, we observed an inverse correlation between virus genome

replication and cell growth and metabolic activity. These data were also confirmed

by western blotting for nsP1 (Fig. 3E), demonstrating that translation of the 5' ORF

was also enhanced at the sub-physiological temperature.

Enhanced genome replication at a sub-physiological temperature is correlated

with impaired expression of ISGs

We considered that one reason for the increased level of CHIKV genome replication

at 28°C might be due to a sub-optimal antiviral transcriptional response, even in the

absence of exogenous IFN stimulation. It has previously been documented that IFN

responses are impaired at sub-physiological temperatures, and this has been

proposed to explain the enhanced CHIKV replication and pathogenicity observed in

mice housed at 22°C compared to 30°C [31].

We first asked if the transcriptional response of C2C12 cells to stimulation with the

synthetic dsRNA analogue poly I:C was affected by temperature. C2C12 cells at

either 28°C or 37°C were mock treated or stimulated with poly I:C for 24 h prior to

analysis of the transcription of three well characterized ISGs by qRT-PCR.   As

shown in Fig 4A, at 37°C poly I:C treatment significantly enhanced the transcription

of *Ifi44i*, *Cmpk2* and *Uba7* compared to cells treated at 28°C, or mock treated cells.

This is consistent with impairment of the ISG response at 28°C.

To test whether this poly I:C response involved canonical IFN signalling via the JAK-

STAT pathway, we treated cells with ruxolitinib (Ruxo: a JAK-STAT inhibitor).   We

first determined that the maximal non-cytotoxic concentration of Ruxo in C2C12 cells

was 10 µM (Fig S5). As expected, treatment of cells with 10 µM Ruxo blocked the

poly I:C stimulated transcription of these 3 genes (Fig 4A), suggesting that this was

acting via the JAK-STAT pathway presumably in an autocrine fashion following the

poly I:C stimulation of IFN expression.

We next tested whether there was a temperature sensitive effect of JAK-STAT

signalling on CHIKV replication.   Specifically, whether inhibition of JAK-STAT

signalling at 37°C could relieve the block to genome replication between 12-24 h.

However, as shown in Fig 4B, Ruxo treatment had no significant effect on the

replication of the CHIKV WT D-luc SGR at either temperature. We also examined

the effect of active CHIKV genome replication on the transcription of the 3 known

ISGs: *Ifi44l*[47], *Cmpk2* [48], and *Uba7* [49]. As shown in Fig 4C, induction of ISG

transcription by CHIKV was higher at 37°C than at 28°C. Notably, with the exception

of *Cmpk2*, the transcriptional upregulation was not affected by Ruxo treatment.

Taken together these data indicate that CHIKV replication induced much higher

levels of these 3 ISGs at 37°C than at 28°C, and that unusually this ISG induction

was not halted by pharmacologic inhibition of JAK-STAT signalling, hinting at non-

canonical ISG activation.

Therefore, we next wished to determine whether only these 3 ISGs were induced, or

whether a global ISG induction also occurred during CHIKV infection. To test this, we

conducted a global transcriptomic analysis. We performed bulk RNA-Seq on total

RNA from C2C12 cells transfected with either WT CHIKV-SGR-D-luc, or the non-

replicative GAA mutant, harvested at either 12 h or 24 h.p.t. We performed pairwise

differential gene expression analysis at the two temperatures, or between WT and

GAA transfected cells. As expected, this analysis revealed many differentially

expressed genes (defined as p adj. < 0.05, and [log2 fold change] >1) both in

response to CHIKV genome replication, but also between the two temperatures.

In this study, to assess effects on CHIKV genome replication, we focused our

analysis on known ISGs (annotated from the Molecular Signatures database

(MSigDB) HALLMARK_INTERFERON_ALPHA_RESPONSE dataset) [50]. In

agreement with our prior RT-qPCR data (Fig. 4C), at 12 h.p.t., *Ifi44l*, *Cmpk2*, and

Uba7 were upregulated in WT- versus GAA-transfected cells cultured at 37°C but not

at 28°C, along with a number of other prominent ISGs, such as *Batf2, Cxcl11,* and

Oasl1, as well as immunoproteosome components such as *Psmb8* (Figs. 5A-B).

Across all 94 annotated ISGs, this amounted to 32 ISGs that were potently (up to

512-fold) up-regulated at 37°C, versus only 2 ISGs (*Ifit3*, *Ccrl2*) that were modestly

(up to 8-fold) up-regulated at 28°C (Figs. 5A-D, 6). More ISGs were up-regulated at

24 h.p.t. in comparison to 12 h.p.t at 37°C or 28°C; nevertheless we observed the

same trend: both the number of ISGs, and the level of upregulation, were markedly

greater at 37°C compared to 28°C at 24 h.p.t. (Figs. 5E-H and 7). At 37°C, 52 out of

94 ISGs were up-regulated but only 11 of these ISGs were up-regulated at 28°C (Fig.

7). When we ranked the fold-changes of gene expressions between WT-transfected

cells at 37°C versus 28°C, the 94 annotated ISGs were strongly enriched among the

highest ranks (Fig. 5I, Normalized enrichment score (NES) -2.09, q-value 0.00 at 12

h.p.t.; Fig. 5J, NES -2.85, q-value 0.00 at 12 h.p.t.), but not when the replication-

deficient GAA was transfected (in this case the data failed to generate a valid score,

likely due to insufficient enrichment or technical limitations in the dataset, data not

shown). To validate these data we undertook qRT-PCR for 6 selected genes (*Uba7,*

Stat2, Ifi44I, Isg20, Oas1a and *Cmpk2*) for which well-characterised primers were

available in house. As shown in Fig. S6 the overall levels of these 6 genes were

higher at 37°C compared to 28°C, at either 12 h.p.t or 24 h.p.t. Overall, these data

support the conclusion that global ISG expression in response to CHIKV genomic

replication is significantly impaired at the sub-physiological temperature.

Enhanced recruitment of the key SG protein G3BP1 to replication sites at a

sub-physiological temperature.

In addition to the ISG response, other cellular pathways may act in an antiviral

fashion. Cell stress triggers SG assembly, which halts translation and therefore may

be detrimental to viral replication. During alphavirus infection, nsP3 sequesters key

SG proteins such as G3BP1/2, and recruits them to cytoplasmic foci where viral

genome replication occurs, leading to inhibition of the formation of *bona fide* SG [23-

25, 51]. We considered that this function of nsP3 might play a role in the enhanced

replication observed in mammalian cells at a sub-physiological temperature. SG

formation can be induced by treatment with sodium arsenite ($NaAsO_2$) which results

in oxidative stress, and they can be disassembled by cycloheximide (CHX) treatment.

Importantly, it has been previously shown that SG formation may be temperature-

dependent, as $NaAsO_2$ was only able to induce SG at 37°C but not 28°C [52]. Using

G3BP1 (detected using a pan-G3BP antibody) as a marker for the formation of bona

fide SG we confirmed that this was also the case in our hands (Fig. S7), as no

G3BP1-positive SG were observed following $NaAsO_2$ treatment of C2C12 cells at

28°C. As expected, at 37°C SG were efficiently induced by $NaAsO_2$ treatment and

disassembled by CHX (Fig. S7).

To investigate this further, we exploited the CHIKV SGR-nsP3-mCh-FF-luc construct

in which nsP3 was fused to mCherry (see Fig. S3A), and we analysed the

distribution and abundance of nsP3-mCherry and G3BP1 in transfected C2C12 cells

(Fig. 8). As a further marker for sites of active genome replication (cytoplasmic foci)

we also stained cells with an antibody to visualize dsRNA (J2), previously shown by

us [1] and others [24, 53] to colocalise with nsP1, nsP3 and G3BP1. Consistent with

previous data in this manuscript (eg Fig. S3B), at 24 h.p.t., both dsRNA and nsP3-

mCherry accumulated to a higher level at 28°C compared to 37°C, and this

enhancement was not seen at earlier timepoints (Fig.8B). In contrast, the amount of

G3BP1 in transfected cells increased over time and was higher at 37°C when

compared to 28°C (Fig. 8B).

We then quantified the colocalisation of nsP3, G3BP1 and dsRNA (Fig. 9A).

Quantitative analysis (Fig. 9B) showed that at 12 h.p.t. there were no differences in

the amount of colocalized nsP3/G3BP1, nsP3/dsRNA or G3BP1/dsRNA between the

two temperatures, however, at 24 h.p.t., the levels of colocalisation for all three pairs

were significantly higher at 28°C compared to 37°C. Taken together, these data

demonstrate that there are more replication sites, higher levels of dsRNA, and more

recruitment of the SG protein G3BP1 to replication sites at 28°C than at 37°C.

Discussion

This study set out to build on prior work regarding the functions of the CHIKV AUD

[1], and analyzed the phenotype of a panel of alanine substitutions. Our study took

an intriguing turn when we observed that 3 of these mutants (W220A, D249A and

Y324A) exhibited a defective replication phenotype in mammalian cells at 37^{o}C, but

no defect in either mosquito cells, or in mammalian cells at a sub-physiological

temperature (28^{o}C), defining these as temperature-sensitive (ts) mutants (Figs 1B

and S2B/C). Although ts mutations in alphavirus nsP3 have been previously

identified, for example F312S and A268V in Sindbis virus, these resulted in a defect

in negative-strand RNA synthesis at the non-permissive temperature (40^{o}C) [54].

These data point to a critical role of nsP3, and the AUD in particular, in genome

replication. Further biochemical analysis of these ts mutants in nsP3 may shed light

on the specific role of nsP3 but is not the focus of this manuscript.

This analysis led us to the serendipitous discovery that wildtype CHIKV SGR

replicated more efficiently at sub-physiological temperatures in mammalian cells.

The novelty of this intriguing and arguably counter-intuitive observation led us to

focus our attention on the underlying mechanisms. We note that, although a 100-

fold enhancement of CHIKV genome replication at a sub-physiological temperature

was observed in the context of CHIKV SGRs (Fig. 1B), the enhancement in the

context of infectious virus was less dramatic (Fig. 2A). We observed only a modest

(5-fold), but reproducible increase in virus production with delayed kinetics at sub-

physiological temperatures, suggesting that other processes during the virus lifecycle

(e.g. virus assembly and release) are also affected by temperature. This is

consistent with the modest increase in maximum virus titre in mice infected with

CHIKV sub-cutaneously through the feet and housed at 22°C compared to 30°C [31].

In considering why CHIKV genome replication in mammalian cells might be

enhanced at sub-physiological temperatures, we reasoned that the first cells infected

in a mammalian host via a mosquito bite would be in the periphery and thus not at

37^oC. It would be reasonable to expect that other arboviruses would also exhibit the

same differential, however, that was not the case as both BUNV and ZIKV replicated

to similar levels at 28^oC and 37^oC (Fig S3C/D) This suggests that this phenotype is

driven by alphavirus specific processes, and we note that neither the nsP3 macro-

domain nor the AUD have functional homologues in either bunyaviruses or

flaviviruses. Analysis of the timescale of genome replication at different

temperatures also revealed that the phenotype was most apparent after 12 h.p.t. In

most cases, replication rates up to 12 h.p.t. did not differ between the two

temperatures (e.g. see Fig. 1A/B), however between 12-24 h.p.t. replication

continued to increase at 28^oC, but plateaued at 37^oC.

An explanation for our observation was that it reflected an aspect of the cellular

response to virus genome replication that was perhaps impaired at the sub-

physiological temperature and only manifest after 12 h.p.t. An obvious candidate

would be the type I IFN response, which has indeed been shown to be reduced at

lower temperatures promoting CHIKV arthritis in a mouse model [31]. Consistent

with this, the response to the synthetic dsRNA analog poly I:C was impaired at 28^oC

and blocked by the JAK-STAT inhibitor Ruxo. Similarly, the ISG transcriptional

response to CHIKV genome replication (in SGR transfected cells) was severely

impaired at the sub-physiological temperature (Figs. 5-7). At the sub-physiological

temperature only a few ISGs from the total of 94 were up-regulated, while at 37^{o}C

many more ISGs were up-regulated (Fig 5C, D, G and H, and Fig 6). This data is

also consistent with previous research [30], in which transcription of ISGs following

IFN treatment was reduced at a sub-physiological temperature.

The exact antiviral mechanisms of ISGs against CHIKV, and whether reduced ISG

expression fully explains temperature-dependent replication remains unclear. Here

we attempted to study the antiviral effect of ISGs in two systems. First, we employed

Vero cells, which cannot produce IFNα/β, and therefore we would expect these cells

to have impaired ISG responses. In Vero cells, we found that CHIKV still replicates

better at 28°C than at 37°C, suggesting that intact type I IFN production is not

required for temperature-driven differences in replication. Second, we treated mouse

C2C12 cells with Ruxo to inhibit JAK-STAT signaling (which is triggered by IFN

signaling) and to thereby block ISG induction. Under canonical conditions, IFN

production triggers a well-characterized JAK1/2-dependent phosphorylation cascade,

which can be inhibited by Ruxo. This phosphorylation leads to the formation of

ISGF3 (p-STAT1, p-STAT2, and IRF9), which subsequently binds to ISG promoters

to activate transcription at later stages. However, under non-canonical conditions

[55], antiviral responses can still be mediated by non-phosphorylated forms of these

factors or alternative transcriptional regulators, bypassing JAK-STAT activation.

One key player in non-canonical signaling is IRF3, a major regulator of type I IFN

production, which can also induce a subset of ISGs independently of IFN signaling.

For example, IRF3 activation upregulates ISG54, ISG20, and IFP35 [56]. Similarly,

during influenza virus infection, several ISGs, are induced without IFN production,

but relies on activation of IRF3 and transcription induction of p56 gene [57]. Likewise,

HCV infection triggers direct recruitment of NF-κB and IRF3 to the CXCL10 promoter

via RIG-I and TLR3 sensing, bypassing type I/III IFN pathways [58]. In HEV infection,

RIG-I plays a dual role, activating IRF3 and IRF7 via JAK-STAT pathway, while also

inducing ISGs such as IFIT1, RANTES, and CXCL10 independently of STAT1

phosphorylation [59]. Similarly, during Coxsackievirus B3 infection, MDA5, rather

than RIG-I, drives an IFN-independent antiviral response in inhibiting viral replication

[60].

More recently, studies on CHIKV infection suggest that CHIKV actively subverts

canonical IFN responses, enhancing its reliance on non-canonical signaling

pathways to restrict viral replication [61]. These observations highlight the critical role

of both canonical and non-canonical IFN signaling in antiviral immunity. A deeper

understanding of these mechanisms could provide novel therapeutic strategies

against CHIKV and other viral infections.

An alternate hypothesis is that reduced ISG expression is only one factor that allows

for increased CHIKV replication at reduced temperatures, and other mechanisms

may play roles in the phenotype. To further investigate the mechanisms

underpinning the enhanced CHIKV replication at the sub-physiological temperature

we analysed the production of SGs, cytoplasmic protein-mRNA aggregates formed

in response to stress and believed to play an antiviral role [18]. Alphavirus infection

has previously been documented to disrupt formation of *bona fide* SG by recruiting

key SG components such as G3BP1/2 to cytoplasmic foci where genome replication

occurs [24], as judged by positivity for both nsP3 and dsRNA. G3BP1/2 have been

shown to be critical for alphavirus genome replication [21]. Sequestration of

G3BP1/2 away from *bona fide* SG will have the added benefit of limiting the antiviral

effects of SG [18].

Interestingly, as seen previously [52], treatment with NaAsO$_2$ (which induces

oxidative stress) did not result in SG formation at 28^{o}C (Fig S6), suggesting that

there was a defect in this process at the sub-physiological temperature. However,

following transfection with a CHIKV SGR (expressing an mCherry-nsP3 fusion) we

observed that, whereas total levels of G3BP1 were higher at 37^{o}C (Fig 8), it was

more efficiently recruited into nsP3/dsRNA positive cytoplasmic foci at 28^{o}C (Fig 9).

Specifically, at 24 h.p.t. not only were there more nsP3/dsRNA positive cytoplasmic

foci at 28^{o}C (Fig. 8), but also the recruitment of G3BP to nsP3/dsRNA foci was

significantly higher at 28^{o}C (Fig 9).

These data are consistent with the hypothesis that the enhanced level of CHIKV

genome replication in mammalian cells at a sub-physiological temperature is

associated with more efficient recruitment of cellular factors to cytoplasmic sites of

replication. A likely explanation is that the absence of *bona fide* SG formation at

28^{o}C (Fig. S7) means that SG components are more abundant in the cytoplasm, and

this facilitates recruitment by nsP3 (or other nsPs). In addition, our data suggest that

the availability of SG components is not a limiting factor for genome replication until

after 12 h.p.t. as there are no differences in replication or recruitment of G3BP1 to

cytoplasmic foci at this timepoint at either temperature (see Figs. 1A, 9B).

Why the process of recruiting SG components to cytoplasmic sites of replication

should differ at different temperatures remains to be determined. Importantly the

ADP-ribosylhydrolase activity of the nsP3 macro-domain has recently been shown to

play a key role in SG formation, disassembly and composition [25]. It is possible that

the enzymatic activity and/or substrate specificity of the macro-domain could be

modulated by temperature. Although the AUD *per se* has not been directly shown to

play a role in SG formation, it is flanked by the macro-domain and HVD, both of

which have now been implicated in SG formation. Localization of CHIKV nsP3 to

cytoplasmic foci in invertebrate cells was shown to depend on both the AUD and the

HVD [62], although it should also be noted that the interaction with G3BP1/2 (or its

insect homologue, Rasputin) involves specific sequences within the HVD [63]. It is

thus possible that all three domains of nsP3 coordinate to control formation of *bona*

fide SG, and recruit SG components to cytoplasmic sites of genome replication.

In conclusion, our data have revealed a hitherto unrecognized enhancement of

CHIKV genome replication in mammalian cells at sub-physiological temperature that

may play a role in the early stages of transmission of virus to the mammalian host

via a mosquito bite. Our observations point to a critical role of the AUD in this

difference, and further work seeking to understand the effects of temperature on

nsp3 interactions with host cell proteins would benefit greatly from a proteomic

approach.

Materials and Methods

Sequence alignment. AUD amino acid sequences from a selection of alphaviruses

(Sindbis, Semliki, Ross River, Venezuelan, Eastern and Western Equine

Encephalitis, ONNV and CHIKV) were obtained from NCBI and aligned by Clustal

Omega. The conserved AUD residues were identified by Pymol and their locations

were predicted from the Sindbis nsP2/3 protein structure (PDB ID code 4GUA) [32].

Cell culture. Vero and BHK-21 cells were grown in DMEM supplied with 10% foetal

calf serum (FCS), 0.5 mM non-essential amino acids and penicillin-streptomycin

(100 units/mL). C2C12 cells were grown as above but with 20% FCS. All

mammalian cells were maintained at 37°C or 28°C with 5% CO_2. BHK-21 cells were

obtained from John Barr and C2C12 cells from Michelle Peckham (University of

Leeds). U4.4 and C6/36 cell lines (*Ae. albopictus*) were originally obtained from

Susan Jacobs (The Pirbright Institute) and were cultured in Leibovitz's L-15 media

containing 10% FCS, 10% tryptose phosphate broth and penicillin-streptomycin (100

units/mL), at 28°C without CO_2. Infection and viral RNA transfection experiments

were carried in accordance with protocols reviewed and approved by the Institutional

Biosafety Committees of the University of Leeds and Princeton University (# 1175)

Plasmid constructs. CHIKV-SGR-D-Luc was used as a template for site-directed

mutagenesis using specific primers to generate each of the AUD mutations. The

AUD mutated fragments were excised and ligated back into the CHIKV-SGR-D-Luc

backbone, or into the infectious ICRES-CHIKV clone [34]. For construction of the

ONNV AUD mutants, a fragment including the AUD coding sequence was excised

from the ONNV-2SG-ZsGreen (donated by Andres Merits, University of Tartu), and

ligated into pcDNA3.1 to generate pcDNA3.1-ONNV-AUD. This was used as a

template for site-directed mutagenesis using specific primers to generate ONNV

AUD mutations. The ONNV AUD mutated fragments were excised from pcDNA3.1-

ONNV-AUD and ligated into the ONNV-2SG-ZsGreen backbone. CHIKV-SGR-

nsP3-mCh-FF-Luc was generated by SpeI digestion of the FF-Luc fragment from

SGR-D-Luc and insertion of the mCherry coding sequence. CHIKV-SGR-FF-Luc

was constructed by SpeI digestion of the R-Luc fragment from SGR-D-Luc and re-

ligation.

Transfection and dual luciferase assay. Capped RNAs of WT and mutant CHIKV-

SGR-D-Luc, ICRES-CHIKV, ONNV-2SG-ZsGreen, or WT CHIKV-SGR-nsP3-mCh-

FF-Luc and WT CHIKV-SGR-FF-Luc were synthesized and purified using the

mMACHINE™ SP6 Transcription Kit (Thermo Fisher Scientific). In brief, 250 ng of

each of the RNAs was transfected into 5×10^4 cells using Lipofectamine 2000 (Life

Technologies) according to the manufacturer's instructions. Cells were harvested

and both FF-Luc and R-Luc activities were measured using the Dual-luciferase Kit

(Promega) according to the manufacturer's instructions. All experiments were

repeated three times independently.

Sequence analysis. Total RNA from transfected or infected cells was extracted with

Trizol, followed by reverse transcription to generate cDNA using the SuperScript IV

Reverse Transcriptase (Invitrogen) according to the manufacturer's instructions. The

expected DNA fragments of the CHIKV or ONNV genome were amplified by PCR

(primer sequences available upon request) and analysed by Sanger sequencing.

MTT assay. C2C12 cells were seeded into 96 well plates. The next day, cells were

mock transfected and incubated at 37°C or 28°C. At indicated time points, medium

was removed and replaced with 150 µl phenol-red free DMEM and 30 µl 6 mM MTT.

Cells were incubated at 37°C for 2 h before media was removed and replaced with

100 µl DMSO. Plates were covered with foil and agitated until all MTT granules had

dissolved, prior to reading the absorbance at 570 nm.

ICRES-CHIKV one-step growth curve. C2C12 cells were seeded into 12 well

plates. The next day, cells were infected with ICRES-CHIKV at MOI 0.1 pfu/cell and

incubated at 37°C for 1 h. The cells were then washed with PBS and replaced with

fresh complete media, followed by incubation at 37°C or 28°C respectively. At

indicated time points cell supernatants were harvested for plaque assay, and cell

were lysed and extracted with TRIzol solution (ThermoFisher) for qRT-PCR analysis.

Plaque assay. Cell supernatants were stored at -80°C until required, thawed,

serially diluted and applied to monolayers of BHK-21 cells for 1 h at 37°C. The

inocula were aspirated and cells were overlaid with 0.8% methylcellulose for 72 h at

37°C or 96 h at 28°C for ONNV infection, and 48 h at 37°C for ICRES-CHIKV.

Plaques were either visualized by EVOS microscopy (ONNV) or stained with 0.05%

crystal violet for 30 min (CHIKV).

qRT-PCR assay. Total RNA from transfected or infected cells was extracted with

Trizol. Strand-specific CHIKV RNA primers were chosen from previous work [64] to

amplify nucleotides 1-1560 of the CHIKV genome and used in reverse transcription

reactions with a LunaScript8482 RT SuperMix Kit (NEB), followed by qPCR using

the 2x qPCRBIO SyGreen Blue Mix (PCRBIOSYSTEMS). Positive and negative

standards were generated using capped ICRES-CHIKV RNA with strand-specific

RNA primers and were *in vitro* transcribed as described before.

Western blotting. Cells were lysed in PLB or Glasgow Lysis Buffer (GLB) (10mM

PIPES-NaOH pH7.2, 120mM KCl, 30mM NaCl, 5mM $MgCl_2$, 1% Triton X-100, 10%

glycerol) supplemented with protease inhibitors (Roche Diagnostics), protein

samples were quantified by BCA assay (Pierce). 10 or 20 μg of each sample was

heat denatured at 95°C for 5 min and separated by SDS-PAGE, proteins were then

transferred to polyvinylidene fluoride (PVDF) membrane and blocked with 50% (v/v)

Odyssey blocking buffer (Li-Cor) diluted in 1X Tris-buffered saline Tween-20 (TBS-T)

(50 mM Tris-HCl pH 7.4, 150 mM NaCl, 0.1% Tween-20). The membrane was then

incubated with primary antibodies (1:1000 dilution) at 4°C overnight and the next day

incubated with fluorescently labelled anti-mouse (700 nm) and anti-rabbit (800 nm)

secondary antibodies for 1 h at room temperature (RT). Membranes were then

imaged on a Li-Cor Odyssey Sa Imager (Li-cor).

Immunofluorescence assay. C2C12 cells were seeded into 12 well plates with

coverslips. The next day, cells were transfected with WT CHIKV-SGR-nsP3-mCh-

FF-Luc as described before. At indicated time points cells were fixed with 4%

formaldehyde, washed with PBS, permeabilized with ice-cold methanol at -20 °C for

10 min and blocked with 2% BSA for 1 h. The cells were incubated with primary

antibody to G3BP1 (Abcam) (1:200) at 4°C overnight. Cells were washed with

DEPC-PBS, prior to incubation with J2 dsRNA antibody (Scicons) (1:100 dilution) at

RT for 1 h. All cells were stained with Alexa Fluor-594 chicken anti-rabbit and Alexa

Fluor-647 goat anti-mouse secondary antibodies (1:500 dilution) at RT for 1 h. Cells

were mounted onto glass slides with Prolong Gold antifade reagent (Invitrogen)

containing 4',6'-diamidino-2-phenylindole dihydrochloride (DAPI) and sealed with nail

varnish. Confocal microscopy images were acquired using a Zeiss LSM880 upright

microscope. Post-acquisition analysis of images was performed using Zen software

(Zen version 2018 black edition 2.3, Zeiss), or Fiji ImageJ (v1.49) software.

Bulk RNA sequencing (RNA-Seq). Capped RNAs of WT and GAA CHIKV-SGR-D-

Luc were synthesized and purified as described above. In brief, 250 ng of each of

the RNAs was transfected into 5×10^4 cells using Lipofectamine 2000 (Life

Technologies) according to the manufacturer's instructions. Cells were harvested at

12 or 24 h.p.t. Starting from 50 ng total RNA per sample, bulk RNA barcoding and

sequencing (BRB-Seq) was performed as described [65] with minor modifications.

Briefly, reverse transcription was performed with Template Switching RT Enzyme

Mix (NEB) and a uniquely barcoded oligo(dT)30 primer for each sample, modified to

use Illumina TruSeq Read 1 priming site instead of Nextera Read 1 [66]. Following

the remainder of the standard BRB-Seq protocol, up to 24 barcoded first-strand

cDNAs were pooled into a single tube, second-strand cDNA was synthesized with

Gubler-Hoffman nick translation, and tagmented cDNA pools with Tn5 produced in-

house [67]. Fourteen cycles of PCR using a P5-containing primer and a unique

multiplexed i7 indexing primer (Chromium i7 Multiplex Kit, 10X Genomics) were

performed.   Libraries were purified and size-selected with 0.65X SPRIselect

(Beckman Coulter) and then 0.7X SPRIselect.   All pools were sequenced on a

NovaSeq SP v1.5 flowcell (Illumina) with 28 cycles of Read 1, 8 cycles of Read i7,

and 101 cycles of Read 2.

RNA-Seq analysis. Reads were demultiplexed with Picard v2.25.6 (from within viral-

core v2.1.33) using the read structure "5S8B15M8B101T" and Q20M1 mismatch

tolerance.   Reads were mapped to the mouse genome mm39 with STAR v2.7.10a,

and those reads mapping to genes on the primary assembly were counted with

htseq-count v1.99.2. Pairwise differential gene expression between conditions was

performed using DESeq2 (1.40.2) [68]. Genes whose expression changed 2-fold

between conditions with adjusted *p*-value <0.05 (Wald test, Benjamini-Hochberg

procedure for multiple hypothesis testing) were considered differentially expressed.

The ISGs (dataset from HALLMARK_ALPHA_RESPONSE [50]) from Mus musculus

species that were significantly expressed were labeled as triangles while the others

were as dots. Heatmaps of the differential expressed ISGs at 37°C or 28°C were

generated using the R package (R-studio v4.3.1).

Data and code availability. Raw RNA-Seq reads and the processed counts matrix

can be accessed at NCBI GEO accession: GSE264494. Reviewers may access this

submission using the token 'epglissetbkldmj', which will be made public following

publication.   The   analysis   scripts   used   in   this   study   are   available   at:

[#https://github.com/path/to/repo#].

Gene set enrichment analysis (GSEA). GSEA was performed by using GSEA

software version (4.3.2), downloaded from Broad Institute Gene Set Enrichment

Analysis website (www.broad.mit.edu/gsea). The enrichment gene sets used in the

analysis were selected in MSigDB (HALLMARK_ALPHA_RESPONSE) from Mus

musculus species. Significance of enrichment magnitude was set at a False

Discovery Rate (FDR) of 5% for GSEA.

Statistical analysis. Statistical analysis was performed using the unpaired Student's

t tests (two-tailed). Data in bar graphs represent statistically significant differences

and displayed as the means ± S.E. of three experimental replicates (n = 3).

Acknowledgments

We thank the following people for their invaluable assistance in this work: Prof. Andres Merits (Institute of Technology, University of Tartu, Estonia) for the CHIKV SGRs, ICRES constructs, ONNV-2SG-ZsGreen construct and antibody to nsP1. Dr. Andrew Tuplin for the Zika virus replicon, Dr John Barr for the BUNV mini-genome construct, Drs. Niluka Goonawardane and Ruth Hughes for help in Fiji ImageJ analysis, Dr Sreenivasan Ponnambalam for access to the EVOS microscope (all at the University of Leeds). Dr. Wei Wang, Jennifer Miller, and Jean Arly Volmar of the Genomics Core Facility at the Princeton University Lewis-Sigler Institute for Integrative Genomics for next-generation sequencing assistance, Drs. Abhishek Biswas and Lance R. Parsons for the help with the RNA-seq analysis.

Funding

This work was in part supported by a Wellcome Investigator Award to MH (grant number 096670) and grants from the National Institutes of Health (R01 AI138797, R01 AI153236, R01 AI146917, R01 AI168048, R01 AI107301, and U19A171401 to AP), Open Philanthropy (to AP) and Princeton University (to AP). JG was supported by a University of Leeds/China Scholarship Council PhD studentship, and subsequently by a New Jersey Commission on Cancer Research (NJCCR) postdoctoral fellowship COCR23PDF016. AEL is supported by Damon Runyon Postdoctoral Fellowship DRG-2432-21. The Zeiss LSM880 confocal microscope at the University of Leeds was funded by a Wellcome multi-user equipment grant (WT104818MA). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

[1] Y. Gao, N. Goonawardane, J. Ward, A. Tuplin, M. Harris, Multiple roles of the non-structural protein 3 (nsP3) alphavirus unique domain (AUD) during Chikungunya virus genome replication and transcription, PLoS Pathog 15(1) (2019) e1007239.
[2] R. Chen, S. Mukhopadhyay, A. Merits, B. Bolling, F. Nasar, L.L. Coffey, A. Powers, S.C. Weaver, C. Ictv Report, ICTV Virus Taxonomy Profile: Togaviridae, J Gen Virol 99(6) (2018) 761-762.
[3] O. Schwartz, M.L. Albert, Biology and pathogenesis of chikungunya virus, Nat Rev Microbiol 8(7) (2010) 491-500.
[4] M.C. Robinson, An epidemic of virus disease in Southern Province, Tanganyika Territory, in 1952-53. I. Clinical features, Trans R Soc Trop Med Hyg 49(1) (1955) 28-32.
[5] W.H. Lumsden, An epidemic of virus disease in Southern Province, Tanganyika Territory, in 1952-53. II. General description and epidemiology, Trans R Soc Trop Med Hyg 49(1) (1955) 33-57.
[6] I. Schuffenecker, I. Iteman, A. Michault, S. Murri, L. Frangeul, M.C. Vaney, R. Lavenir, N. Pardigon, J.M. Reynes, F. Pettinelli, L. Biscornet, L. Diancourt, S. Michel, S. Duquerroy, G. Guigon, M.P. Frenkiel, A.C. Brehin, N. Cubito, P. Despres, F. Kunst, F.A. Rey, H. Zeller, S. Brisse, Genome microevolution of chikungunya viruses causing the Indian Ocean outbreak, PLoS Med 3(7) (2006) e263.
[7] K. Sergon, C. Njuguna, R. Kalani, V. Ofula, C. Onyango, L.S. Konongoi, S. Bedno, H. Burke, A.M. Dumilla, J. Konde, M.K. Njenga, R. Sang, R.F. Breiman, Seroprevalence of Chikungunya virus (CHIKV) infection on Lamu Island, Kenya, October 2004, Am J Trop Med Hyg 78(2) (2008) 333-7.
[8] A.J. Tatem, S.I. Hay, D.J. Rogers, Global traffic and disease vector dispersal, Proc Natl Acad Sci U S A 103(16) (2006) 6242-7.
[9] O.J. Brady, M.A. Johansson, C.A. Guerra, S. Bhatt, N. Golding, D.M. Pigott, H. Delatte, M.G. Grech, P.T. Leisnham, R. Maciel-de-Freitas, L.M. Styer, D.L. Smith, T.W. Scott, P.W. Gething, S.I. Hay, Modelling adult Aedes aegypti and Aedes albopictus survival at different temperatures in laboratory and field settings, Parasit Vectors 6 (2013) 351.
[10] E.A. Mordecai, J.M. Cohen, M.V. Evans, P. Gudapati, L.R. Johnson, C.A. Lippi, K. Miazgowicz, C.C. Murdock, J.R. Rohr, S.J. Ryan, V. Savage, M.S. Shocket, A. Stewart Ibarra, M.B. Thomas, D.P. Weikel, Detecting the impact of temperature on transmission of Zika, dengue, and chikungunya using mechanistic models, PLoS Negl Trop Dis 11(4) (2017) e0005568.
[11] T. Ahola, A. Merits, Functions of Chikungunya Virus Nonstructural Proteins, in: C.M. Okeoma (Ed.), Chikungunya Virus: Advances in Biology, Pathogenesis, and Treatment, Springer International Publishing, Cham, 2016, pp. 75-98.
[12] J.C. Rupp, K.J. Sokoloski, N.N. Gebhart, R.W. Hardy, Alphavirus RNA synthesis and non-structural protein functions, J Gen Virol 96(9) (2015) 2483-2500.
[13] H. Malet, B. Coutard, S. Jamal, H. Dutartre, N. Papageorgiou, M. Neuvonen, T. Ahola, N. Forrester, E.A. Gould, D. Lafitte, F. Ferron, J. Lescar, A.E. Gorbalenya, X. de Lamballerie, B. Canard, The crystal structures of Chikungunya and Venezuelan equine encephalitis virus nsP3 macro domains define a conserved adenosine binding pocket, J Virol 83(13) (2009) 6534-45.
[14] L. Eckei, S. Krieg, M. Butepage, A. Lehmann, A. Gross, B. Lippok, A.R. Grimm, B.M. Kummerer, G. Rossetti, B. Luscher, P. Verheugd, The conserved

macrodomains of the non-structural proteins of Chikungunya virus and other
pathogenic positive strand RNA viruses function as mono-ADP-ribosylhydrolases,
Sci Rep 7 (2017) 41746.
[15] R.L. McPherson, R. Abraham, E. Sreekumar, S.E. Ong, S.J. Cheng, V.K. Baxter,
H.A. Kistemaker, D.V. Filippov, D.E. Griffin, A.K. Leung, ADP-ribosylhydrolase
activity of Chikungunya virus macrodomain is critical for virus replication and
virulence, Proc Natl Acad Sci U S A 114(7) (2017) 1666-1671.
[16] N.J. Foy, M. Akhrymuk, A.V. Shustov, E.I. Frolova, I. Frolov, Hypervariable
domain of nonstructural protein nsP3 of Venezuelan equine encephalitis virus
determines cell-specific mode of virus replication, J Virol 87(13) (2013) 7569-84.
[17] V. Kril, M. Hons, C. Amadori, C. Zimberger, L. Couture, Y. Bouery, J. Burlaud-
Gaillard, A. Karpov, D. Ptchelkine, A.L. Thienel, B.M. Kummerer, A. Desfosses, R.
Jones, P. Roingeard, L. Meertens, A. Amara, J. Reguera, Alphavirus nsP3 organizes
into tubular scaffolds essential for infection and the cytoplasmic granule architecture,
Nat Commun 15(1) (2024) 8106.
[18] J.J. Fros, N.E. Domeradzka, J. Baggen, C. Geertsema, J. Flipse, J.M. Vlak, G.P.
Pijlman, Chikungunya virus nsP3 blocks stress granule assembly by recruitment of
G3BP into cytoplasmic foci, J Virol 86(19) (2012) 10873-9.
[19] N. Kedersha, P. Anderson, Mammalian stress granules and processing bodies,
Methods Enzymol 431 (2007) 61-81.
[20] N. Kedersha, M.D. Panas, C.A. Achorn, S. Lyons, S. Tisdale, T. Hickman, M.
Thomas, J. Lieberman, G.M. McInerney, P. Ivanov, P. Anderson, G3BP-Caprin1-
USP10 complexes mediate stress granule condensation and associate with 40S
subunits, J Cell Biol 212(7) (2016) 845-60.
[21] H. Matsuki, M. Takahashi, M. Higuchi, G.N. Makokha, M. Oie, M. Fujii, Both
G3BP1 and G3BP2 contribute to stress granule formation, Genes Cells 18(2) (2013)
135-46.
[22] H. Tourriere, K. Chebli, L. Zekri, B. Courselaud, J.M. Blanchard, E. Bertrand, J.
Tazi, The RasGAP-associated endoribonuclease G3BP assembles stress granules,
J Cell Biol 160(6) (2003) 823-31.
[23] M.D. Panas, T. Schulte, B. Thaa, T. Sandalova, N. Kedersha, A. Achour, G.M.
McInerney, Viral and cellular proteins containing FGDF motifs bind G3BP to block
stress granule formation, PLoS Pathog 11(2) (2015) e1004659.
[24] F.E. Scholte, A. Tas, I.C. Albulescu, E. Zusinaite, A. Merits, E.J. Snijder, M.J.
van Hemert, Stress granule components G3BP1 and G3BP2 play a proviral role
early in Chikungunya virus replication, J Virol 89(8) (2015) 4457-69.
[25] A.K. Jayabalan, S. Adivarahan, A. Koppula, R. Abraham, M. Batish, D.
Zenklusen, D.E. Griffin, A.K.L. Leung, Stress granule formation, disassembly, and
composition are regulated by alphavirus ADP-ribosylhydrolase activity, Proc Natl
Acad Sci U S A 118(6) (2021).
[26] E.V. Mesev, R.A. LeDesma, A. Ploss, Decoding type I and III interferon
signalling during viral infection, Nat Microbiol 4(6) (2019) 914-924.
[27] G.R. Stark, J.E. Darnell, Jr., The JAK-STAT pathway at twenty, Immunity 36(4)
(2012) 503-14.
[28] W.M. Schneider, M.D. Chevillotte, C.M. Rice, Interferon-stimulated genes: a
complex web of host defenses, Annu Rev Immunol 32 (2014) 513-45.
[29] Y. Zhang, C.W. Burke, K.D. Ryman, W.B. Klimstra, Identification and
characterization of interferon-induced proteins that inhibit alphavirus replication, J
Virol 81(20) (2007) 11246-55.

[30] W.C. Lane, M.D. Dunn, C.L. Gardner, L.K.M. Lam, A.M. Watson, A.L. Hartman,
K.D. Ryman, W.B. Klimstra, The Efficacy of the Interferon Alpha/Beta Response
versus Arboviruses Is Temperature Dependent, mBio 9(2) (2018).
[31] N.A. Prow, B. Tang, J. Gardner, T.T. Le, A. Taylor, Y.S. Poo, E. Nakayama,
T.D.C. Hirata, H.I. Nakaya, A. Slonchak, P. Mukhopadhyay, S. Mahalingam, W.A.
Schroder, W. Klimstra, A. Suhrbier, Lower temperatures reduce type I interferon
activity and promote alphaviral arthritis, PLoS Pathog 13(12) (2017) e1006788.
[32] G. Shin, S.A. Yost, M.T. Miller, E.J. Elrod, A. Grakoui, J. Marcotrigiano,
Structural and functional insights into alphavirus polyprotein processing and
pathogenesis, Proc Natl Acad Sci U S A 109(41) (2012) 16534-9.
[33] L. Pohjala, A. Utt, M. Varjak, A. Lulla, A. Merits, T. Ahola, P. Tammela, Inhibitors
of alphavirus entry and replication identified with a stable Chikungunya replicon cell
line and virus-based assays, PLoS One 6(12) (2011) e28923.
[34] K. Tsetsarkin, S. Higgs, C.E. McGee, X. De Lamballerie, R.N. Charrel, D.L.
Vanlandingham, Infectious clones of Chikungunya virus (La Reunion isolate) for
vector competence studies, Vector Borne Zoonotic Dis 6(4) (2006) 325-37.
[35] G.C. Roberts, C. Zothner, R. Remenyi, A. Merits, N.J. Stonehouse, M. Harris,
Evaluation of a range of mammalian and mosquito cell lines for use in Chikungunya
virus research, Sci Rep 7(1) (2017) 14641.
[36] E.M. Morazzani, M.R. Wiley, M.G. Murreddu, Z.N. Adelman, K.M. Myles,
Production of virus-derived ping-pong-dependent piRNA-like small RNAs in the
mosquito soma, PLoS Pathog 8(1) (2012) e1002470.
[37] G.F. Zhu, Z.H. Liu, J. Wang, Improved technic for dengue virus micro cell culture,
Chin Med J (Engl) 97(1) (1984) 73-4.
[38] J. Desmyter, J.L. Melnick, W.E. Rawls, Defectiveness of interferon production
and of rubella virus interference in a line of African green monkey kidney cells (Vero),
J Virol 2(10) (1968) 955-61.
[39] J.M. Emeny, M.J. Morgan, Regulation of the interferon system: evidence that
Vero cells have a genetic defect in interferon production, J Gen Virol 43(1) (1979)
247-52.
[40] V.H. Leonard, A. Kohl, T.J. Hart, R.M. Elliott, Interaction of Bunyamwera
Orthobunyavirus NSs protein with mediator protein MED8: a mechanism for
inhibiting the interferon response, J Virol 80(19) (2006) 9667-75.
[41] R.W. Malone, J. Homan, M.V. Callahan, J. Glasspool-Malone, L. Damodaran, B.
Schneider Ade, R. Zimler, J. Talton, R.R. Cobb, I. Ruzic, J. Smith-Gagen, D. Janies,
J. Wilson, G. Zika Response Working, Zika Virus: Medical Countermeasure
Development Challenges, PLoS Negl Trop Dis 10(3) (2016) e0004530.
[42] F. Weber, E.F. Dunn, A. Bridgen, R.M. Elliott, The Bunyamwera virus
nonstructural protein NSs inhibits viral RNA synthesis in a minireplicon system,
Virology 281(1) (2001) 67-74.
[43] M. Mutso, S. Saul, K. Rausalu, O. Susova, E. Zusinaite, S. Mahalingam, A.
Merits, Reverse genetic system, genetically stable reporter viruses and packaged
subgenomic replicon based on a Brazilian Zika virus isolate, J Gen Virol 98(11)
(2017) 2712-2724.
[44] J. Saraste, L. Kaaiainen, H. Soderlund, S. Keranen, RNA synthesis directed by
a temperature-sensitive mutant of Semliki Forest virus, J Gen Virol 37(2) (1977) 399-
406.
[45] L. Kaariainen, D. Sawicki, P.J. Gomatos, Cleavage defect in the non-structural
polyprotein of Semliki Forest virus has two separate effects on virus RNA synthesis,
J Gen Virol 39(3) (1978) 463-73.

[46] J. Suopanki, D.L. Sawicki, S.G. Sawicki, L. Kaariainen, Regulation of alphavirus 26S mRNA transcription by replicase component nsP2, J Gen Virol 79 (Pt 2) (1998) 309-19.

[47] J. Du, H. Luo, S. Ye, H. Zhang, Z. Zheng, K. Liu, Unraveling IFI44L's biofunction in human disease, Front Oncol 14 (2024) 1436576.

[48] J.H. Lai, D.W. Wu, C.H. Wu, L.F. Hung, C.Y. Huang, S.M. Ka, A. Chen, Z.F. Chang, L.J. Ho, Mitochondrial CMPK2 mediates immunomodulatory and antiviral activities through IFN-dependent and IFN-independent pathways, iScience 24(6) (2021) 102498.

[49] S. Liu, G. Hu, S. Luo, W. Wu, Q. Zhou, R. Jin, Y. Zhang, H. Ruan, H. Huang, H. Li, Insights into the evolution of the ISG15 and UBA7 system, Genomics 114(2) (2022) 110302.

[50] A. Liberzon, C. Birger, H. Thorvaldsdottir, M. Ghandi, J.P. Mesirov, P. Tamayo, The Molecular Signatures Database (MSigDB) hallmark gene set collection, Cell Syst 1(6) (2015) 417-425.

[51] M.D. Panas, M. Varjak, A. Lulla, K.E. Eng, A. Merits, G.B. Karlsson Hedestam, G.M. McInerney, Sequestration of G3BP coupled with efficient translation inhibits stress granules in Semliki Forest virus infection, Mol Biol Cell 23(24) (2012) 4701-12.

[52] J.R. Wheeler, T. Matheny, S. Jain, R. Abrisch, R. Parker, Distinct stages in stress granule assembly and disassembly, Elife 5 (2016).

[53] D.Y. Kim, J.M. Reynaud, A. Rasalouskaya, I. Akhrymuk, J.A. Mobley, I. Frolov, E.I. Frolova, New World and Old World Alphaviruses Have Evolved to Exploit Different Components of Stress Granules, FXR and G3BP Proteins, for Assembly of Viral Replication Complexes, PLoS Pathog 12(8) (2016) e1005810.

[54] Y.S. Hahn, A. Grakoui, C.M. Rice, E.G. Strauss, J.H. Strauss, Mapping of RNA-temperature-sensitive mutants of Sindbis virus: complementation group F mutants have lesions in nsP4, J Virol 63(3) (1989) 1194-202.

[55] S. Swaraj, S. Tripathi, Interference without interferon: interferon-independent induction of interferon-stimulated genes and its role in cellular innate immunity, mBio 15(10) (2024) e0258224.

[56] J. Andersen, S. VanScoy, T.F. Cheng, D. Gomez, N.C. Reich, IRF-3-dependent and augmented target genes during viral infection, Genes Immun 9(2) (2008) 168-75.

[57] M.J. Kim, A.G. Latham, R.M. Krug, Human influenza viruses activate an interferon-independent transcription of cellular antiviral genes: outcome with influenza A virus is unique, Proc Natl Acad Sci U S A 99(15) (2002) 10096-101.

[58] J. Brownell, J. Bruckner, J. Wagoner, E. Thomas, Y.M. Loo, M. Gale, Jr., T.J. Liang, S.J. Polyak, Direct, interferon-independent activation of the CXCL10 promoter by NF-kappaB and interferon regulatory factor 3 during hepatitis C virus infection, J Virol 88(3) (2014) 1582-90.

[59] L. Xu, W. Wang, Y. Li, X. Zhou, Y. Yin, Y. Wang, R.A. de Man, L.J.W. van der Laan, F. Huang, N. Kamar, M.P. Peppelenbosch, Q. Pan, RIG-I is a key antiviral interferon-stimulated gene against hepatitis E virus regardless of interferon production, Hepatology 65(6) (2017) 1823-1839.

[60] E. Francisco, M. Suthar, M. Gale, Jr., A.B. Rosenfeld, V.R. Racaniello, Cell-type specificity and functional redundancy of RIG-I-like receptors in innate immune sensing of Coxsackievirus B3 and encephalomyocarditis virus, Virology 528 (2019) 7-18.

[61] Y. Suzuki, Interferon-induced restriction of Chikungunya virus infection, Antiviral Res 210 (2023) 105487.

[62] J.J. Fros, C. Geertsema, K. Zouache, J. Baggen, N. Domeradzka, D.M. van
Leeuwen, J. Flipse, J.M. Vlak, A.B. Failloux, G.P. Pijlman, Mosquito Rasputin
interacts with chikungunya virus nsP3 and determines the infection rate in Aedes
albopictus, Parasit Vectors 8 (2015) 464.
[63] G. Nowee, J.W. Bakker, C. Geertsema, V.I.D. Ros, G.P. Goertz, J.J. Fros, G.P.
Pijlman, A Tale of 20 Alphaviruses; Inter-species Diversity and Conserved
Interactions Between Viral Non-structural Protein 3 and Stress Granule Proteins,
Front Cell Dev Biol 9 (2021) 625711.
[64] N.E. Plaskon, Z.N. Adelman, K.M. Myles, Accurate strand-specific quantification
of viral RNA, PLoS One 4(10) (2009) e7468.
[65] D. Alpern, V. Gardeux, J. Russeil, B. Mangeat, A.C.A. Meireles-Filho, R.
Breysse, D. Hacker, B. Deplancke, BRB-seq: ultra-affordable high-throughput
transcriptomics enabled by bulk RNA barcoding and sequencing, Genome Biol 20(1)
(2019) 71.
[66] E.V. Mesev, A.E. Lin, E.G. Guare, B.L. Heller, F. Douam, B. Adamson, J.E.
Toettcher, A. Ploss, Membrane-proximal motifs encode differences in signaling
strength between type I and III interferon receptors, Sci Signal 16(806) (2023).
[67] S. Picelli, A.K. Björklund, B. Reinius, S. Sagasser, G. Winberg, R. Sandberg,
Tn5 transposase and tagmentation procedures for massively scaled sequencing
projects, Genome Res 24(12) (2014) 2033-2040.
[68] M.I. Love, W. Huber, S. Anders, Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2, Genome Biol 15(12) (2014) 550.

Figure legends

Figure 1. Replication and translation of CHIKV AUD mutants in C2C12 cells at

28°C or 37°C. C2C12 cells were transfected with WT and mutant CHIKV-SGR-D-

Luc RNAs and cultured at 37°C (A) or 28°C (B). Cells were harvested at indicated

time points. FF-Luc activities of WT and mutants were measured and normalized to

their respective 4 h values, which thus represent fold-change compared to the 4 h

values. (**C**) Protein expression of nsP1 for the WT and mutant CHIKV-SGR-D-Luc

at 28°C or 37°C in C2C12 cells. Cells were lysed at 24 h.p.t., 10 µg of each sample

was loaded and analysed by western blotting. Actin was also detected by western

blot as a loading control. (**D**) Quantification of CHIKV positive and negative strand

RNA from C2C12 cells transfected with CHIKV-SGR-D-luc. Transfected C2C12

cells from (**A**) were lysed with TRIzol solution, and total RNA was purified by phenol-

chloroform extraction. cDNAs of CHIKV positive or negative strand RNA were

synthesized as described before [64].

Figure 2. Phenotype of WT and AUD mutants in the context of ICRES-CHIKV at

28°C or 37°C. (**A**) Phenotype of WT and AUD mutants in ICRES-CHIKV production

at 28°C or 37°C. C2C12 cells were transfected with WT and mutant ICRES-CHIKV

RNA and incubated at 28°C or 37°C. Supernatants were collected 48 h post

infection (h.p.i.) from both temperatures and titrated at 37°C by plaque assay.

Significant differences denoted by * (P<0.05), ** (P<0.01) or **** (P<0.0001). Data

are displayed as the means ±S.E. of three experimental replicates. (**B**) One step

growth curve of ICRES-CHIKV in C2C12 cells at 28°C or 37°C. C2C12 cells were

infected with ICRES-CHIKV at MOI 0.1 and incubated at 37°C or 28°C respectively.

Cell supernatants were collected at indicated time points and virus titers were

determined by plaque assay. (**C**) CHIKV RNA quantification from infected C2C12

cells. C2C12 cells collected at indicated time points were lysed by TRIzol solution,

and total RNA were purified by phenol-chloroform extraction. cDNAs of CHIKV

positive or negative strand RNA were synthesized as described before [27].

Figure 3. Temperature shift assay of WT SGR-D-Luc. (**A**) Schematic of

temperature shift assay. Transfected C2C12 cells were incubated at either 28°C or

37°C prior to a temperature shift from 12-24 h.p.t. (28°C to 37°C (designated 24 h

S*), or 37°C to 28°C (designated 24 h S#). Controls maintained the same

temperature from 0-24 h.p.t. (**B**) Replication assay. WT CHIKV-SGR-D-Luc RNA

was transfected into C2C12 cells and incubated at 28°C or 37°C according to the

schematic in **(A)**. FF-Luc activities were measured and normalized to their

respective 4 h values. Significant differences denoted by ** (P<0.01) or ****

(P<0.0001), compared to replication at 12 h.p.t. Data are displayed as the means

±S.E. of three experimental replicates. Cell numbers were enumerated (**C**) and cell

metabolic activity **(D)** determined by MTT assay at 28°C or 37°C. **(E)** Protein

expression of nsP1 at 28°C or 37°C. Cells were lysed at 24 h.p.t. and 10 µg of each

sample was analysed by western blot. Actin was also detected by western blot as a

loading control.

Figure 4. Effect of ruxolitinib (Ruxo) treatment on ISG expression in cells

transfected with poly I:C or WT CHIKV-SGR-D-Luc. (**A**) Effect of Ruxo on ISG

expression in response to poly I:C treatment. C2C12 cells in 24 well plates were pre-

treated with 10 uM Ruxo for 12 h before transfection with 1 ug poly I:C per well and

ISG expression was measured by qRT-PCR at 24 hpt. (**B**) Replication of WT and

GAA CHIKV-SGR-D-Luc in response to Ruxo at 28°C or 37°C. C2C12 cells were

first pre-treated with 10 uM Ruxo 12 h before transfection of WT and GAA CHIKV-

SGR-D-Luc. Cells were then incubated at 37°C or 28°C and harvested at 4, 12 and

24 hpt. (**C**) Fold change analysis of ISGs for WT and GAA at 37°C or 28°C

compared to mock control. Fifty µl of lysed samples was subjected to RNA extraction,

followed by elution in RNase-free H_2O. Ct values for three ISGs were measured

alongside the housekeeping gene β-actin across different samples, and fold changes

were calculated accordingly. Significant differences denoted by n.s. (P>=0.05), *

(P<0.05) or ** (P<0.001) compared to WT are only shown for the 24 h time point for

clarity. Data are displayed as the means ±S.E. of three experimental replicates.

Figure 5. Volcano plots of differential expressed genes in response to WT

CHIKV-SGR-D-Luc replication at 28°C or 37°C. C2C12 cells were transfected with

WT CHIKV-SGR-D-Luc at 28°C or 37°C as described above. At 12 h.p.t or 24 h.p.t.,

total RNA was harvested and subjected to bulk RNA-Seq. (**A-H**) Volcano plots of

pairwise differential gene expression between the different conditions. The IFN-

stimulated genes (ISGs) that were significantly expressed are labeled as triangles

while non-ISGs are labeled as dots. (**I and J**) Waterfall plots of gene set enrichment

analysis (GSEA) of differentially expressed genes within ISGs. ISGs were defined as

the 94 genes within the *M. musculus* Molecular Signatures Database (MSigDB)

HALLMARK_ALPHA_RESPONSE [50].

Figure 6. Heatmap of ISGs in response to WT CHIKV-SGR-D-Luc replication at

28°C or 37°C at 12 h.p.t. The differentially expressed ISGs from the volcano plots at

12 h.p.t. were selected and grouped by R-studio. The heatmap plot was generated

using Graphpad Prism version 9.0.

Figure 7. Heatmap of ISGs in response to WT CHIKV-SGR-D-Luc replication at

28°C or 37°C at 24 h.p.t. The differentially expressed ISGs from the volcano plots at

24 h.p.t. were selected and grouped by R-studio. The heatmap plot was generated

using Graphpad Prism version 9.0.

Figure 8. Formation of cytoplasmic nsP3/dsRNA positive foci at 28°C or 37°C.

(**A**) C2C12 cells were transfected with WT CHIKV-SGR-nsP3-mCh-FF-Luc and

incubated at 28°C or 37°C respectively.   Cells were then fixed with 4%

paraformaldehyde and stained with antibodies to G3BP1 (Green) or dsRNA (gray).

nsP3-mCherry was expressed as a fused protein (Red).   Scale bar 10 µm. (**B**)

Quantification of the overall dsRNA, nsP3-mCherry and G3BP1 expression at 28°C

or 37°C. Protein expression levels were determined for 10 cells and calculated by

Fiji ImageJ. Significant differences denoted by * (P<0.05), *** (P<0.001) and ****

(P<0.0001), compared to values at 37°C.

Figure 9. Colocalisation of G3BP1 with nsP3 and dsRNA at 28°C or 37°C. (**A**)

Detection of dsRNA, nsP3 and G3BP1 in WT CHIKV-SGR-nsP3-mCh-FF-Luc

transfected C2C12 cells. Immunofluorescence assays were performed as before.

Colocalisation of dsRNA, nsP3 and G3BP1 is shown. All image analysis and

processing was performed using Fiji software. Fluorescence plot profile of nsP3,

dsRNA and G3BP1 signal intensities along the indicated white line is shown. (**B**)

Quantitative analysis of the colocalization between dsRNA, nsP3 and G3BP1 signals.

Pearson's correlation coefficient between dsRNA and G3BP1, nsP3 and G3BP1,

dsRNA and nsP3 were determined for 10 cells and calculated by Fiji ImageJ.

Significant differences denoted by **** ($P<0.0001$), compared to values at 37°C.

1068

Figure 1

Figure 2

Figure 3

A

Fold change in ISGs over mock control at 37°C and 28°C

B

Viral replication for WT and GAA with and without Ruxolitinib at 37° C and 28°C

C

Fold change of ISGs for WT and GAA at 37° C and 28°C over mock control

Figure 4

Figure 5

Heatmap pairwise comparison 12 h.p.t

Figure 6

Heatmap pairwise comparison 24 h.p.t

Figure 7

A Immunostaining of infected cells

B Quantification of fluorescence intensity

Figure 8

Figure 9