Balancing selection at a wing pattern locus is associated with major shifts in genome-wide patterns of diversity and gene flow in a butterfly

María Ángeles Rodríguez de Cara[1*$], Paul Jay[1*$], Quentin Rougemont[1*$], Mathieu Chouteau[1,2], Annabel Whibley[3,4], Barbara Huber[5], Florence Piron-Prunier[3], Renato Rogner Ramos [6], André V. L. Freitas[6], Camilo Salazar[7], Karina Lucas Silva-Brandão[8], Tatiana Texeira Torres[9], Mathieu Joron[1$]

* contributed equally

[1]Centre d'Ecologie Fonctionnelle et Evolutive (CEFE), Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France
[2]Laboratoire Ecologie, Evolution, Interactions Des Systèmes Amazoniens (LEEISA), Université de Guyane, IFREMER, CNRS, Cayenne, Guyane Française
[3]Institut de Systématique Evolution Biodiversité (ISYEB), Museum National d'Histoire Naturelle, CNRS, Sorbonne-Université, EPHE, Université des Antilles, Paris, France
[4]School of Biological Sciences, University of Auckland, Auckland, New Zealand
[5]Instituto de Ciencias Ecológicas y Ambientales (ICAE), Univ de los Andes, Mérida, Venezuela
[6]Departamento de Biologia Animal, Instituto de Biologia, Unicamp, Campinas, São Paulo, Brazil
[7]Department of Biology, Faculty of Natural Sciences, Universidad del Rosario, Carrera 24 No 63C-69, Bogotá 111221, Colombia.
[8]Centro de Biologia Molecular e Engenharia Genética, Universidade Estadual de Campinas. Av. Candido Rondom 400. Campinas, São Paulo, Brazil
[9]Department of Genetics and Evolutionary Biology, Institute of Biosciences, University of São Paulo (USP), São Paulo, Brazil

$ Corresponding authors: angeles.decara@gmail.com, paul.yann.jay@gmail.com, mathieu.joron@cefe.cnrs.fr, quentinrougemont@orange.fr

Abstract: Selection shapes genetic diversity around target mutations, yet little is known about how selection on specific loci affects the genetic trajectories of populations, including their genome-wide patterns of diversity and demographic responses. Here we study the patterns of genetic variation and geographic structure in a neotropical butterfly, *Heliconius numata*, and its closely related allies in the so-called melpomene-silvaniform clade. *H. numata* is known to have evolved an inversion supergene which controls variation in wing patterns involved in mimicry associations with distinct groups of co-mimics. Butterflies show disassortative mate preferences and heterozygote advantage at this locus. We contrasted patterns of genetic diversity and structure 1) among extant polymorphic and monomorphic populations of *H. numata*, 2) between *H. numata* and its close relatives, and 3) between ancestral lineages. We show that *H. numata* populations which carry the inversions as a balanced polymorphism show markedly distinct patterns of diversity compared to all other taxa. They show the highest genetic diversity and effective population size estimates in the entire clade, as well as a low level of geographic structure and isolation by distance across the entire Amazon basin. By contrast, monomorphic populations of *H. numata* as well as its sister species and their ancestral lineages all show lower effective population sizes and genetic diversity, and higher levels of geographical structure across the continent. One hypothesis is that the large effective population size of polymorphic populations could be caused by the shift to a regime of balancing selection due to the genetic load and disassortative preferences associated with inversions. Testing this hypothesis with forward simulations supported the observation of increased diversity in populations with the supergene. Our results are consistent with the hypothesis that the formation of a supergene triggered a change in gene flow, causing a general increase in genetic diversity and the homogenisation of genomes at the continental scale.

Introduction: Genetic diversity is shaped by selective processes such as stabilizing or disruptive selection, and by demographic processes such as fluctuations in effective population size. Empirical studies on genetic diversity within and among populations abound, fuelled by an increasing availability of whole genome data, and spurred by our interest in understanding the underlying causes of variation in diversity (e.g. Beichmann 2018, Muers 2009; Murray 2017; Nielsen et al. 2009). At the locus scale, strong directional or disruptive selection tends to reduce diversity within populations (Mitchell-Olds et al. 2007), while balancing selection tends to enhance diversity (Charlesworth 2006). Genome-wide factors reducing diversity include low effective population sizes, generating drift, while high genetic diversity is enhanced by large population sizes and gene flow. Overall, it is well recognised that demographic changes should have a genome-wide effect on diversity, while positive selection is expected to play a role on the sites within and around the genes involved in trait variation (Glinka et al. 2003, Muers 2009, Nielsen et al. 2009).

Variation in behaviour and life-history traits, for instance involving changes in offspring viability or dispersal distance, may also affect species demography, and thus whole genome genetic diversity. However, whether and how genetic variability in a population may be driven by phenotypic evolution at certain traits is poorly understood, and confounding effects may affect patterns of genomic diversity, such as variation in census population size or colonization history. Dissecting how selection on a trait may affect genome-wide diversity can be tackled by comparing closely-related populations differing at this trait coupled with knowledge of when the differences evolved. Here, we took advantage of the dated introgressive origin of a chromosomal inversion associated with major life-history variation to study the demographics and whole genome consequences of changes in the selection regime at a major-effect locus.

Heliconius butterflies are aposematic, chemically-defended butterflies distributed over the American tropics from Southern Brazil to Southern USA (Emsley 1965; Brown 1979) (Fig 1A). *Heliconius* butterflies are well-known for visual resemblance among coexisting species, a relationship called Müllerian mimicry which confers increased protection from bird predators through the evolution of similar warning signals (Sheppard et al. 1985). Most species are locally monomorphic, but their mimicry associations vary among regions, and most species display a geographic mosaic of distinct mimetic "races" through their range. In contrast to most *Heliconius* species, the tiger-patterned *Heliconius numata* is well-known for maintaining both mimicry polymorphism within localities, with up to seven differentiated coexisting forms, and extensive geographic variation in the distribution of wing phenotypes (Brown & Benson 1974; Joron et al. 1999). Forms of *H. numata* combine multiple wing characters conveying resemblance to distinct sympatric species in the genus *Melinaea* and other local Ithomiini species (Nymphalidae: Danainae). Polymorphism in *H. numata* is controlled by a supergene, i.e. a group of multiple linked functional loci segregating together as a single Mendelian locus, coordinating the variation of distinct elements of phenotype (Brown & Benson 1974; Joron et al. 2006). Supergene alleles are characterized by rearrangements of the ancestral chromosomal structure, forming three distinct chromosomal forms with zero (ancestral type, Hn0), one (Hn1) or three chromosomal rearrangements (Hn123) (Fig 1B). The ancestral arrangement, Hn0, devoid of inversions, is fixed in most *Heliconius* species (although an inversion in the same region evolved independently in a distantly-related *Heliconius* lineage (Edelman et al. 2019)). Arrangement Hn1 contains a 400kb inversion called P_1 originating from an introgression event about 2.2 My ago from *H. pardalinus,* in which P_1 is fixed (Jay et al. 2018). This introgression is thought to be the founding event triggering the formation of the supergene and the maintenance of polymorphism in *H. numata* (Jay et al. 2018). Arrangement Hn123 displays two additional inversions, P_2 and P_3, in linkage with P_1, and therefore originated after the introgression of P_1 into the *H. numata* lineage (Jay et al. 2021).

Heliconius numata is widespread in the lowland and foothill tropical forests of the Amazon basin, the Guianas, and the Brazilian Atlantic Forest (Mata Atlântica), but the frequencies of the three chromosome arrangements vary across the range. Ancestral type Hn0 is fixed in the Atlantic Forest populations of Brazil (forms *robigus* or *ethra)*, but segregates at intermediate frequencies in all other *H. numata* populations throughout the range (forms *silvana* and *laura*) (Fig 1C). Chromosome type Hn1 is associated with the Andean mimetic form *bicoloratus* and is found in the Eastern Andean foothills of Ecuador, Peru, and Bolivia. Chromosome type Hn123 is associated with a large diversity of wing-pattern forms of intermediate allelic dominance, including *tarapotensis*, *arcuella* and *aurora*, and is reported from Andean, lowland Amazonian and Guianese populations. Inversion polymorphism is therefore structured across the range, with populations being fixed for the ancestral chromosome (Atlantic Forest, see Text S1 & Table S1-2), or displaying a polymorphism with two (Amazon-Guiana) or three (Andes) chromosomal types in coexistence (Joron et al. 2011). Monomorphic populations of the Atlantic forest, devoid of rearrangements at the supergene locus, might represent the ancestral state displayed by *H. numata* populations before the evolution of the supergene via introgression (Fig 1C).

The wing patterns of *H. numata* are subject to selection on their resemblance to local co-mimics (Chouteau et al. 2016), but the polymorphism is maintained by balancing selection on the chromosome types. Balancing selection is indeed mediated by disassortative mating favouring

mixed-form mating (Chouteau et al. 2017) and is likely to have evolved in the response to the deleterious mutational load carried by inversions, which causes heterozygous advantage in *H. numata* (Jay et al. 2021, Faria et al. 2019, Maisonneuve et al. 2019). The introgression of P_1 and the formation of a supergene were associated with a major shift in the selection regime (Jay et al. 2018). The mating system was also changed during or after introgression. These events may therefore have profoundly affected the population biology of the recipient species, *H. numata*. We investigate here whether the adaptive introgression of a balanced inversion is associated with a signature in the genetic diversity and geographic structure. In particular, we predict that genetic diversity should be higher in *H. numata* than in closely related taxa. Similarly nucleotide diversity should be higher in all polymorphic populations carrying either one segment (Hn1) or two (Hn1,Hn123) compared to the population that is monomorphic and only carries the non-inverted segment (Hn0) in the Brazilian Atlantic Forest. We analyse changes in the demographic history of the clade containing *H. numata* and closely related taxa, as well as their current patterns of diversity and demography, using three well separated populations of *H. numata* representing different states of inversion polymorphism. Our results suggest that following supergene formation, a change in the selection regime and mating system may have facilitated gene flow among morphs and had key consequences in current patterns of genetic structure.

Material and Methods

We used here whole genome resequencing from 137 specimens of *Heliconius,* including 68 *H. numata*. Sampling included specimens from populations in the Andean foothills (3 chromosome types), from the upper Amazon (2 chromosome types), from French Guiana (2 chromosome types) and from the Brazilian Atlantic Forests (1 chromosome type) (Fig 1C; Table S3). Related taxa were represented by the sister species *H. ismenius*, found west of the Andes (parapatric to *H. numata*), by Amazonian representatives of the lineage *H. pardalinus* (donor of the inversion), *H. elevatus, H. ethilla, H. besckei* as well as *H. hecale,* and by *H. melpomene* and *H. cydno* as outgroups. Only Andean, Amazonian and Guianese populations of *H. numata* display chromosomal polymorphism, all other taxa being fixed for the standard gene arrangement (Hn0), or for the inverted arrangement Hn1 (*H. pardalinus*) (Jay et al. 2018). Hereafter, *H. numata* populations from the Andes, Amazon and French Guiana will be collectively referred to as "Amazonian", and populations from the Atlantic Forest as "Atlantic". Butterfly bodies were preserved in NaCl saturated DMSO solution at 20°C and DNA was extracted using QIAGEN DNeasy blood and tissue kits according to the manufacturer's instructions with RNase treatment. Illumina Truseq paired-end whole genome libraries were prepared and 2x100bp reads were sequenced on the Illumina HiSeq 2000 platform. Reads were mapped to the *H. melpomene* Hmel2 reference genome (Davey et al., 2016) using Stampy (version 1.0.28; Lunter and Goodson, 2011) with default settings except for the substitution rate which was set to 0.05 to allow for the expected divergence from the reference of individuals in the so-called silvaniform clade (*H. numata, H. pardalinus, H. elevatus, H. hecale, H. ismenius, H. besckei* and *H. ethilla*). *H. melpomene* and *H. cydno* belonging to the so-called *melpomene* clade, their genomes were mapped with a substitution rate of 0.02. Alignment file manipulations were performed using SAMtools v0.1.3 (Li et al. 2009). After mapping, duplicate reads were excluded using the *MarkDuplicates* tool in Picard (v1.1125; http://broadinstitute.github.io/picard) and local indel realignment using IndelRealigner was performed with GATK (v3.5; DePristo et al. 2011). Invariant and polymorphic sites were called with GATK HaplotypeCaller, with options --min_base_quality_score 25 --min_mapping_quality_score 25 -stand_emit_conf 20 --heterozygosity 0.015.

F_{ST}, d_{XY} and π, were calculated in overlapping windows of 25 kb based on linkage disequilibrium
decay (*Heliconius* Genome Consortium 2012) using custom scripts provided by Simon H. Martin
(https://github.com/simonhmartin), and the genome-wide average was calculated using our own
scripts (available from https://github.com/angelesdecara). Distance in km between sampling sites
was measured along a straight line, not taking into account potential physical barriers. Following
Rousset (1997), in a 2-dimensional habitat, under a model of isolation by distance (IBD)
differentiation, measured as $F_{ST}/(1-F_{ST})$, should increase as a function of the logarithm of distance.
Therefore, we tested for the existence and intensity of an IBD signal among species and between
populations of *H. numata* using a linear model. If IBD is stronger in species not polymorphic for the
inversion we should observed significantly steeper slopes in these species. To test this, we measured
IBD (I) within populations of each species separately, (ii) for all *H. numata* within the Amazonian
region (excluding Atlantic forest populations) and (iii) for all *H. numata* including the Atlantic
region. The slopes of $F_{ST}/1- F_{ST}$ versus log(distance) were calculated using the R package *lsmeans*
(Lenth 2016); the slope difference among species or between populations within species was
estimated with an ANOVA and its significance evaluated with function pairs of this package (Text
S1 and see example script on github.com/angelesdecara/).

Admixture (Alexander et al. 2009) analyses were run on a subset of the 68 *H. numata* genomes,
keeping only 15 individuals from Peru to have a more balanced representation of individuals across
the geographic distribution . Filters were applied to keep biallelic sites with minimum mean depth
of 8, maximum mean depth of 200 and at most 50% genotypes missing. We only kept 1 SNP per
kilobase to remove linked variants using the thinning function in vcftools, and we obtained the
optimal number of clusters using cross-validation for values of K from 1 to 10 (Alexander et Lange,
2011). Principal component analyses (PCA) were performed with the same filters as for admixture,
using the same *H. numata* genomes as for the admixture analyses, using plink2 (Chang et al. 2015).

In order to estimate demographic parameters independently of the effect of selection on diversity,
we performed stringent filtering on the dataset. We removed all predicted genes and their 10,000
base-pair flanking regions, before performing G-PhoCS (Gronau et al. 2011) analyses as detailed
below. Repetitive regions were masked using RepeatMasker and Tandem Repeat Finder (Benson
1999). GC islands detected with CpGcluster.pl with parameters 50 and 1E-5 (Hackenberg et al.,
2006) were also masked. Scaffolds carrying the supergene rearrangements (Hmel215006 to
Hmel215028) were excluded, as were scaffolds from the sex chromosome (Z) and mtDNA, since
those are expected to show unusual patterns of diversity due to selection and different effective
population sizes.

We analysed the demographic history of *H. cydno, H. numata, H. ismenius, H. pardalinus* and *H.*
elevatus with G-PhoCS, which allows for the joint inference of divergence time, effective
population sizes and gene flow. In order to detect differences in demography correlating with the
presence of the supergene in *H. numata,* we conducted analyses separating the Atlantic population
of *H. numata* from Amazonian populations. G-PhoCS is an inference method based on a full
coalescent isolation-with-migration model. Inferences are conditioned on a given population
phylogeny (based on Kozak et al. 2015) with migration bands (i.e. priors in the migration rates) that
describe allowed scenarios of post-divergence gene flow. The model assumes distinct migration rate
parameters associated with each pair of populations, and allows for asymmetric gene flow. Given

the computational burden of G-PhoCS, we selected two individuals per taxon or population, retaining those with the highest sequencing depth (see Table S3). The input dataset consisted of 4092 genomic regions, each 1kb in length and spaced at approximately 30kb intervals (above the value at which LD decay at more than half of its value) and with genotypes in at least one of the two samples of each taxon We used as priors for coalescence times (τ) and genetic diversity (θ), Gamma functions with α=1 and β=100, and for migration rates α=0.002 and β=0.00001. These priors were chosen to allow good convergence while also ensuring non informativity. In order to calculate the highest posterior density interval, we used the library HDInterval in R, and to integrate such posterior densities we used the library sfsmisc in R. We rescaled the results using a mutation rate of 1.9E-9 (Martin et al. 2016) and 4 generations per year (i.e., g=0.25). Migration bands were considered significant following the criteria of Freedman et al. (2012): if the 95% HPD interval did not include 0 or if the total migration rate (migration rate times the duration of the migration band) was larger than 0.03 with posterior probability larger than 0.5.

Demographic Reconstruction of population size changes, split and mixtures

The G-phocs method provided useful information across all species but *i*) do not allow to quantify the time scale of population size change, *ii*) is limited in the number of individuals it can handle and *iii*) displayed limited accuracy to distinguish Ne and m in a simulation study (Gronau et al. 2011). We thus constructed additional models to test the hypothesis that *H. numata* populations with inversion polymorphism display an increased effective population size due to disassortative mating To test this, we used ∂a∂i to reconstruct the demographic history of *H. numata* individuals from the Amazonian forest, quantify their historical changes in effective population size and test their divergence history from 1) *H. numata* from the Brazilian area, which do not carry the inversion; and 2) *H. pardalinus* individuals. We allowed for change in effective population size in both the ancestral populations. Theoretically, the change in effective population size in *H. numata* associated with the change in mating system should be more recent than the time of introgression of the inversion into *H. numata*. To verify this hypothesis we allowed for change in *Ne* of the daughter population at any time after the split. We tested different models of divergence with and without (asymmetric) migration and included the effect of linked selection.

Since, the conditions of historical divergence are not known, we tested a model of divergence with ongoing migration (IM) a model of divergence with ancient migration if gene-flow has stopped recently (AM) and, in the case of divergence into multiple refugia, a model of secondary contact (SC). We also included a model of Strict Isolation (SI) as a null model.

The models shared the following parameters: the ancestral populations of size N_{anc} can grow or shrink to a size N_{anc2} between T_{anc} and up until its splits at time T_{split} into two daughter populations of size N_1 and N_2. Under the SI model, no gene flow occurs between the two populations. Under AM, gene flow occurred between T_{split} and T_{am} and is followed by a period of strict isolation. Under IM, gene flow occurs at a constant rate at each generation between the two populations. Gene flow can be asymmetric, so that two independent migration rates m_{12} (from population 2 to 1) and m_{21} (from population 1 to 2) were modeled. Under the SC model, the population evolved in strict isolation between T_{split} and until T_{sc} where a secondary contact occurs continuously up to present time. Gene flow is modeled as $M = 2N_{REF}.m$. In ∂a∂i, heterogeneity in effective population size was used to account for linked selection by defining two categories of loci with varying effective population sizes (proportion 1-Q of loci with a "neutral Ne" and a proportion Q of loci with a reduced effective population size due to either selection at linked site). To quantify how linked selection affects reduced N_e, we used a Hill-Robertson scaling factor (Hrf) to relate the effective

population size of loci influenced by selection (Nr = Hrf * N_e) to that of neutral loci (N_e). A
hierarchical approach was used to avoid over-fitting: first we compared models assuming constant
effective population size. Second, the best identified models were modified to incorporate
population expansion or decline, as expected given the observed distribution of genetic diversity.
Population expansion was implemented using two additional parameters for population 1 and
population 2, allowing each population to either grow or decline exponentially at any time after
their split from the ancestral population (controlled by parameters s1 and s2 for population 1 and 2
respectively).
Models were fitted using the diffusion theory implemented in $\partial a \partial i$ (Gutenkunst et al. 2009) and
includes the effect of linked selection. $\partial a \partial i$ uses the SFS as a summary of the data. For a given
demographic model, the SFS is computed using diffusion approximation and compared to the
empirical SFS using AIC.
We used stringent filtering (GQ>30, 4 < mean depth < 80) and no missing data to keep high quality
sites and remove potential paralogs or PCR duplicates exhibiting excessive read depth. To minimize
linkage we subset our data to keep one SNP every 5kb. No MAF filter was used and singletons
were kept to avoid ascertainment bias in estimates of demographic parameters. For each model, 32
independent replicate runs were performed and only models with the lowest AIC and ΔAIC were
kept.
276
Forward Simulations
In order to better understand the nature of the processes that generate higher genetic diversity in *H.*
numata compared to closely related taxa, we used simulations to test the hypothesis that
disassortative mating generates an increase in levels of genetic diversity at a genome-wide scale.
We hypothesized that such level of genetic diversity is higher than expected under i) random mating
(a model similar to panmixia) or ii) assortative mating, as commonly observed in other *Heliconius*
species. To test this hypothesis we run forward simulation under disassortative, assortative and
random matting using slim v3.6 (Messer et al. 2013).
We simulated a stepping stone model with 10 demes, each composed of 1,000 diploid individuals
and connected by a (symmetric) migration parameter (m). Each individual received neutral and
deleterious (ratio 16:6) mutations at a rate μ = 1e-8 μ/bp/generation (rescaled to μ = 1e-6 for faster
simulation of a larger population). We simulated an individual with a pair of 1Mb chromosome,
including a single locus with 5 alleles with perfect dominance (allele 1 > allele 2 > allele 3 > allele
4 > allele 5) given 5 possible alternative phenotypes (referred hereafter as "morph"). Each allele
was fully linked (no recombination) with a given deleterious recessive mutation, generating
overdominance at this loci so that polymorphism is always maintained. Local adaptation was
introduced in the model through a single parameter defining randomly which morphs were favored
in each population. In each population, either 2 or 3 morphs benefited from a fitness advantage
compared to the others. The fitness reduction varied between 0 (= fitness of zero for migrants in a
demes) and 1 (no reduction of fitness). We tested 3 possible values for this parameter (0, 0.25 and
0.5).
Finally, disassortative mating was controlled by a mate choice parameter defining whether a morph
would reproduce with another morph. The strength of the parameter varied between 0 (= complete
disassortative mating) and 1 (= no mating weight). We tested 3 possible values for this parameter
(0, 0.25 and 0.5).
We run the model for 80,000 generations to reach demographic equilibrium and assessed levels of
synonymous diversity (π_s). We tested all combinations of the 3 values for levels of disassortative

mating and local adaptation and ran 10 replicates per combination in order to estimate the variance
around π_s.

Similarly, we run a model with strict assortative mating, controlled by a parameter defining whether
similar morphs reproduced together. The strength of the parameter varied between 0 (complete
assortative mating where a given individual mate only with an identical morph) and 1 (where
individual mate randomly with regards to the morph). We tested 3 possible values for this parameter
(0, 0.25 and 0.5). As for disassortative mating, all combinations of assortative mating and local
adaptation values were tested. For each model we tested 3 values for the migration rate, m = 1e-4,
1e-6 and 1e-8, resulting in a total of 54 comparisons.

For graphical display in Figure 4, the values of assortative/disassortative mating were rescaled on a
scale between (0 and 1) with 0 indicating no disassortative mating but complete assortative mating
and 1 complete disassortative mating (or no assoartive mating). A value of 0.5 was equivalent to
random mating.

Results
Using cross validation error as a measure of the optimal number of clusters with Admixture, we
found that K=2 was the optimal cluster number describing within-species genetic variation in *H.*
numata (Fig 2A). One cluster corresponds to the Atlantic population, forming a well-differentiated
genetic entity compared to all other *H. numata* populations. All Amazonian populations of *H.*
numata showed a remarkable uniformity. This pattern is consistent with the population structure
inferred using microsatellite markers (Fig S1). Population structure revealed by PCA is in line with
the admixture analysis (Fig 2B). Individuals from the Atlantic populations of *H. numata* clustered
together to one side of the first PCA axis, whereas all other individuals from all other populations
clustered to the other side. The second axis of the PCA separates individuals from French Guiana
from the other samples of the upper Amazon. This separation was not found with Admixture (i.e.
with K=3) from the complete dataset. To better investigate the existence of a hierarchical
population structure, we excluded individuals from the Atlantic populations and compared
individuals from French Guiana to a randomly sampled set of Peruvian individuals. In this case we
found a clear separation in two groups corresponding to French Guiana and Peru (**Fig S2A**). The
same pattern was observed when replacing Peru by Colombia or Ecuador (**Fig S2B,C**). In
accordance, pairwise genome-wide estimates of differentiation (F_{ST}) between *H. numata*
populations showed elevated values when comparing the Atlantic population to other populations,
low values between French Guiana and other Amazonian population, and were the lower when
comparing pairs of Amazonian populations outside of French Guiana. (Fig 2C, Table S4). For
instance, the population from La Merced in Peru shows an $F_{ST} = 0.032$ with the population from
French Guiana at a distance of 3019km, but an $F_{ST} = 0.311$ (an order of magnitude higher) with the
Atlantic population at a similar distance. The comparison between La Merced and Ecuador was
even lower (Fst = 0.0159). Isolation by distance among Amazonian populations of *H. numata*,
estimated using the proxy $F_{ST}/1\text{-Fst} \sim \log10(km)$ was significant ($R^2 = 0.41$, p = 1.61e-06, slope =
0.02). An analysis of the slope revealed a lower rate of increase in F_{ST} with distance in *H. numata*
compared to all other taxa (Fig 2C, Table S4, Supp Text S1). By contrast, IBD between Atlantic
and Amazonian populations of *H. numata* is close to what is observed in other species, and not
significantly different (see Supp. Text S1).

Analyses of genetic diversity show that all populations of *H. numata,* except those from the Atlantic
Forest, have a similarly high genetic diversity (Fig 3A). By contrast, closely related *Heliconius* taxa

show significantly lower genetic diversity (Fig 3A). These patterns are similar to those obtained using G-PhoCS to analyse the demographic histories in a phylogenetic context, where Amazonian populations of *H. numata* show higher population sizes compared to Atlantic populations (Fig 3B, Table S5). G-PhoCS analyses also show a demographic history in which gene flow plays a crucial role (Table S6). For instance, our analyses show significant gene flow right at the beginning of the divergence between *H. ismenius* and the other silvaniforms, as well as in the divergence between *H. pardalinus* and *H. elevatus.* The effective population sizes inferred from Atlantic genomes are one order of magnitude lower than those obtained using *H. numata* populations from other localities (Fig 3A and Table S5). In our cladogram, the increase in *H. numata* population size is restricted to the Amazonian branch, excluding Atlantic populations.

Demographic reconstruction from $\partial a \partial i$

The model selection procedure based on AIC gave higher support for a model of secondary contact (SC) in the pairwise comparison between *H. numata* from Peru and *H. numata* from Brazil. The pairwise comparison between *H. numata* and *H. pardalinus* supported a model of divergence with continuous gene-flow (IM) (**Table S7, Figure S3**). All models supported an expansion occuring in the ancestral population, followed by further growth of the *H. numata* carrying the inversion supergene to reach a size of several millions, which was by far the largest effective size compared to all other species. This stands in stark contrast with the results observed in the samples from Brazil (which do not harbour the inversion) (**Table 2**). Accordingly, *H. numata* populations from the Atlantic forests of Brazil appear to have been subject to a bottleneck at the start of their divergence from Amazonian populations, followed by exponential growth, suggesting a strong (and recent) founding event, leading to a comparatively smaller population size than that observed in the rest of *H. numata.* It is worth noting however that effective population size was hard to estimate in pairwise comparisons between *H. numata* from Peru and SE Brazil. Indeed, parameter uncertainty was large, and model residuals (**Figure S3**) were also large. Our results indicated that *H. pardalinus* displayed an initially large population size followed by a comparatively smaller size expansion than *H. numata* (**Table 2**). Estimates of current effective population sizes are therefore qualitatively similar to those from G-phocs.

Forward simulations

Forward simulations under different levels of local adaptation (controlled by the strength of divergent selection), disassortative mating and migration are displayed in **Figure 4B**. The same results under a model of assortative mating involving different levels of selection and migration are displayed in **Figure 4A**. Overall, synonymous genome-wide nucleotide diversity (π_S) was higher in 73 % of the models including disassortative mating (average $\pi_S = 0.0145$) when compared to their equivalent under assortative mating (average $\pi_S = 0.011$), a weak but significant difference (p <0.01, see **Figure S5**). In summary, modest differences were observed among models with different strengths of divergent selection or disassortative mating, the most influential variable being the rate of migration (**Figure 4)**.

Discussion

Our results suggest that populations displaying inversion polymorphism in the *P* supergene in *H. numata* also display distinctive population demography and gene flow. Differences in demographic and differentiation regimes associated with structural variation at this locus are revealed when comparing polymorphic populations of *H. numata* to closely-related monomorphic taxa, such as (1)

peripheral populations of *H. numata,* (2) sister taxa, and (3) inferred ancestral lineages. This suggests that the existence of a mimicry supergene controlling polymorphism in *H. numata* may be associated, in time and in space, with major differences in population biology. We hypothesize this to be due to a change in the balancing selection regime due to heterozygote advantage (Jay et al. 2021) and in the associated evolution of disassortative mating (Chouteau et al. 2017) following the onset of inversion polymorphism, causing direct effects on ecological parameters such as gene flow, immigration success and effective population size. Testing this hypothesis through forward simulation yielded mixed evidence for a genome-wide effect of this disassortative mating, especially when compared to a simple model of random mating.

Our analyses show large-scale variation in genetic diversity among closely related taxa in this clade of *Heliconius* butterflies. Within *H. numata*, genetic diversity found in polymorphic Amazonian populations is ~4 times higher thanin populations from the Atlantic Forest. Generally, Amazonian populations of *H. numata* harbour the highest genetic diversity in the entire *melpomene*/silvaniform clade, which contrasts with the low diversity found in the most closely related taxa such as *H. ismenius* or *H. besckei*. Inferring historical demography during the diversification of the *H. numata* lineage reveals that the large effective population size in that species is only associated with the branch representing polymorphic, Amazonian *H. numata* populations, while internal branches all show very low diversity estimates. This suggests that ancestral and putatively monomorphic populations of *H. numata* were similar in their diversity parameters to current sister species *H. ismenius* populations, or to current peripheral Atlantic *H. numata* populations. Although low-diversity lineages could have lost diversity due to recent events such as strong bottlenecks, as estimates of effective population size from dadi indicated for the Atlantic population. Nevertheless our dadi estimates do suggest that the Amazonian populations of *H. numata* underwent a dramatic increase in effective population size posterior to their split with Central American (*H. ismenius*), *H. pardalinus* and the Atlantic populations. Those findings are in agreement with G-Phocs analyses. The Amazonian branch of the *H. numata* radiation is characterized by the long-term maintenance of inversion polymorphism, triggered by the introgression of a chromosomal inversion about 2.2 Ma ago. Therefore, the major shift in demography between Amazonian and Atlantic populations indeed appears to coincide, at least in the broad sense, with the occurrence of inversion polymorphism, even though the lack of replication of this event impedes firmly establishing causality.

Another striking result is the lack of genetic structure displayed by *H. numata* across the Amazon, with all Amazonian populations forming a single genetic cluster. Only Atlantic populations stand out and display high differentiation with other *H. numata* from the rest of the range. French Guiana and Peruvian populations, separated by over 3000 km across the Amazon, are weakly genetically differentiated compared to pairs of populations at comparable distances in other species, and show only modestly stronger differentiation than pairs of *H. numata* populations taken at short distances. *H. numata* populations from the Amazon show significantly lower isolation by distance than all other taxa, as measured by the change in F_{ST} across distance (F_{ST}/km) (Fig. 2C), with a very distinctive, flat slope of isolation by distance. The only exception is found when comparing Amazonian populations with Atlantic populations of Brazil, displaying a level of differentiation in line with that of pairs of populations at similar distances within other taxa.

Effective population size is affected by census size, mating system, and the force and type of selection acting on traits (Charlesworth 2009). Selection is often viewed as a force only affecting

the genetic variation around specific, functional loci in the genome, but it may also affect whole genome diversity, for instance when its action is sufficient to modify local demography or mating patterns. In *H. numata*, morphs and therefore inversion genotypes show disassortative mate preferences, i.e., they preferentially mate with individuals carrying different chromosome types (Chouteau et al. 2017). Disassortative mating enhances heterozygosity and the mating success of individuals expressing rare alleles (negative frequency dependence) (Knoppien 1985; Hedrick et al. 2018). Consequently, immigrants expressing rare, recessive alleles have a mating advantage in *H. numata*. Disassortative mating associated with the supergene should therefore bring an advantage to immigrant genomes in LD with recessive supergene alleles, possibly enhancing genome-wide gene flow.Supergenes are also characterised by single-locus Mendelian inheritance, by which mimicry phenotypes are maintained in the face of recombination, even after immigration.Effective migration regime in populations harbouring a mimicry supergene is therefore likely to be quite different to that observed in other mimetic taxa such as *Heliconius melpomene* or *H. erato*, in which mimicry variation is controlled by multiple loci with diverse dominance patterns. In those taxa, hybrid offspring display recombinant patterns breaking down mimicry, even after multiple generations of backcrossing, and pure forms mate assortatively with respect to wing pattern (McMillan et al. 1997, Mallet et al. 1998, Jiggins et al. 2001); both processes select against mimetic variants migrating from adjacent areas with distinct warning patterns. The expectation is that immigrant genomes should be consistently associated with mimicry breakdown in the case of multilocus architectures, which should translate into an effect on effective migration genomewide, compared to situations with polymorphic mimicry supergenes. In *H. numata*, the evolution of a polymorphic mimicry supergene and disassortative mate preferences could therefore explain the relative lack, compared to other *Heliconius* taxa, of differentiation among polymorphic populations, even across large distances. Furthermore, enhanced gene flow could also cause an increase in effective population size estimates (Slatkin 1987), putatively explaining why polymorphic populations of *H. numata* harbour the highest genetic diversity, and display the highest *Ne* estimates in the entire *melpomene-silvaniform* clade of *Heliconius*. These hypotheses are also supported by our forward simulation which suggests that indeed, disassortative mating resulted in enhanced genetic diversity compared with assortative mating, although the effect was small. In addition, our results also suggests that a simple model of random mating may explain well the data, thus purely demographic expansions may also generate high genetic diversity and high effective population size, as observed from our ∂a∂i demographic modelling.

Alternative processes may of course contribute to the observed patterns. Amazonian and Atlantic populations may differ in other aspects that could also result in differences in genetic diversity. Habitat availability and structure may be different, possibly entailing differences in the maintenance of diversity. The Atlantic Forest is vast in area, but may represent a smaller biome compared to the Amazon, and is isolated from the bulk of the range of *H. numata*, which could result in populations displaying characteristics of peripheral populations with smaller effective population sizes (Eckert et al. 2008). Reduced effective population size is supported by our data. One major caveat associated to our inference remains the small number of individuals (n = 12) from the Atlantic forest. Genetic diversity might be underestimated, notably if populations have a history of fragmentation in this area. The other *Heliconius* species in the clade have much in common with *H. numata* in terms of habitat and general ecology, yet their niche and life-history specificities and their phylogenetic histories may result in consistent differences with the polymorphic *H. numata* populations. All those specificities may contribute to the observed pattern in which polymorphic

Amazonian populations of *H. numata* display high effective population size and a weak geographic structure in genome-wide genetic variation. Yet this pattern of variation correlates parsimoniously with the evolution of a supergene causing disassortative mating and single-locus control of mimicry variation in Amazonian *H. numata* populations, which provides an elegant mechanism explaining their differences with extant and ancestral closely-related lineages. However, we cannot rule out a role for conjectural differences in ecology and geography with all other taxa.

In conclusion, our results show a remarkable contrast in the demography and differentiation of populations within the Amazonian range of *H. numata* compared to closely related taxa and ancestral lineages, as well as with other taxa in the *melpomene*/silvaniform clade. The distinctiveness of this widely polymorphic species in the clade is consistent with the hypothesis that the evolution of a supergene maintained by balancing selection represents a major transition in this lineage, triggering changes in genome-wide patterns of diversity and population ecology over the last 2 million years since its formation. If this hypothesis is correct, the evolution of a locus under balancing selection may therefore feed-back on population ecology and diversification, and consequently on speciation.

Eco-evolutionary feedbacks between changes in genomic architecture and the ecological parameters of populations are still not well understood and few cases have been studied. The evolution of self-incompatibility in plants, affecting the rules of mating and feeding back on population ecology, connectivity, and demography, may be one example, but effects of the evolution of trait genetic architecture on population ecology may be more common than previously thought. In our study, more work on the determinants of variation in effective population sizes in the genus *Heliconius* is needed to determine the precise impact of the supergene on demography in *H. numata*. We believe that our results emphasize a potential link between genomic architecture, selection and demography, and should inspire future theoretical and modelling studies. Overall our result suggests that balancing selection maintaining structural polymorphisms affecting life-history traits may have a profound influence on species ecology.

Contributions:
MARdC, PJ, QR and MJ designed the study and wrote the manuscript. BH, AVLF, TTT, RRR, KLSB provided the Atlantic samples. CS provided the Colombian samples. MARdC, PJ and QR performed genomic analyses and simulations with input from AW. MARdC, PJ, MJ, FPP and MC collected the Peruvian and Ecuatorian samples. MC performed microsatellite analyses and organized fieldworks and butterfly rearing. All authors contributed to editing the manuscript.

Aknowledgements:
This work was funded by grants Hybevol (ANR-12-JSV7-0005) and Supergene (ANR-18-CE02-0019-01) from the Agence Nationale de la Recherche and European Research Council Grant MimEvol (StG-243179). We acknowledge the Genotoul and the Montpellier Bioinformatics Biodiversity (MBB) platforms for providing us with calculation time. We thank Dr. Vitor Becker, at the Serra Bonita Reserve (Bahia), Alexandre Soares, at the MN/UFRJ (Rio de Janeiro) and Dr. Marcelo Duarte at the MZ/USP (Sao Paulo) for their contribution to the collection of butterflies in Brazil. Field collections in Colombia were conducted under permit no. 530 issued by the Autoridad Nacional de Licencias Ambientales (ANLA).We are grateful to Marianne Elias and Violaine Llaurens for comments and discussions. AVLF acknowledges support from Fundação de Amparo à Pesquisa do Estado de São Paulo – (FAPESP) (Biota-Fapesp grants 2011/50225-3, 2013/50297-0)

and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) (421248/2017-3 and 304291/2020-0). KLSB acknowledges the financial support of FAPESP Process # 2012/16266-7. Brazilian specimens are registered under SISGEN (A701768).

Data availability:

The raw sequence data were deposited in NCBI SRA and accession numbers are indicated in Supplementary table 3.

References

1. Alexander DH, Novembre J, Lange K (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research* **19**:1655-1664.

2. Alexander, D.H., Lange, K. Enhancements to the ADMIXTURE algorithm for individual ancestry estimation. BMC Bioinformatics 12, 246 (2011). https://doi.org/10.1186/1471-2105-12-246

3. Beichmann AC, Huerta-Sanchez E, Lohmueller KE (2018). Using Genomic Data to Infer Historic Population Dynamics of Nonmodel Organisms. *Annual Review of Ecology, Evolution, and Systematics* **49**:433–56

4. Benson G (1999) Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Research* **27**:573-580.

5. Brown KS (1979). Ecologia Geográfica e Evolução nas Florestas Neotropicais. – Univ. Estadual de Campinas, Campinas, Brazil.

6. Brown KS, Benson WW (1974). Adaptive polymorphism associated with multiple müllerian mimicry in *Heliconius numata* (Lepid.: Nymph.). *Biotropica* **6**:205–228

7. Brown KS, Mielke OHH. 1972. The Heliconians of Brazil (Lepidoptera: Nymphalidae). Part II. Introduction and general comments, with a supplementary revision of the tribe. *Zoologica, New York*, **57**:1–40.

8. Chang CC, Chow CC, Tellier LCAM, Vattikuti S, Purcell SM, Lee JJ (2015) Second-generation PLINK: rising to the challenge of larger and richer datasets. GigaScience, **4**:s13742–015–0047–8.

9. Charlesworth B (2009) Fundamental concepts in genetics: effective population size and patterns of molecular evolution and variation. *Nature Reviews Genetics* **10**:195-205.

10. Chouteau M, Arias M, Joron M (2016). Warning signals are under positive frequency-dependent selection in nature. *Proceedings of the National Academy of Sciences of the USA* **113**:2164–2169.

11. Chouteau M, Llaurens V, Piron-Prunier F, Joron M. (2017). Polymorphism at a mimicry supergene maintained by opposing frequency-dependent selection pressures. *Proceedings of the National Academy of Sciences of the USA* **114**: 8325–8329.

12. Davey JW, Chouteau M, Barker SL, Maroja L, Baxter SW, Simpson F, et al. (2016). Major Improvements to the *Heliconius melpomene* Genome Assembly Used to Confirm 10 Chromosome Fusion Events in 6 Million Years of Butterfly Evolution. *G3* **6**:695–708. doi:10.1534/g3.115.023655

13. Danecek P,Adam Auton, Goncalo Abecasis, Cornelis A. Albers, Eric Banks, Mark A. DePristo, Robert Handsaker, Gerton Lunter, Gabor Marth, Stephen T. Sherry, Gilean McVean, Richard Durbin and 1000 Genomes Project Analysis Group, The Variant Call Format and VCFtools, Bioinformatics, 2011

14. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis AA, del
Angel G, Rivas MA, Hanna M, McKenna A, Fennell TJ, Kernytsky AM, Sivachenko AY,
Cibulskis K, Gabriel SB, Altshuler D, Daly MJ (2011). A framework for variation discovery
and genotyping using next-generation DNA sequencing data. *Nature Genetics* **43**:491–498.

15. Eckert CG, Samis KE, Lougheed SC (2008). Genetic variation across species' geographical
ranges: the central–marginal hypothesis and beyond. *Molecular Ecology* **17**:1170–1188.

16. Edelman NB, Frandsen PB, Miyagi M, Clavijo B, Davey J, Dikow RB, García-Accinelli G,
Van Belleghem SM, Patterson N, Neafsey DE, Challis R, Kumar S, Moreira GRP, Salazar C,
Chouteau M, Counterman BA, Papa R, Blaxter M, Reed RD, Dasmahapatra KK, Kronforst
M, Joron M, Jiggins CD, McMillan WO, Di Palma F, Blumberg AJ, Wakeley J, Jaffe D,
Mallet J (2019). Genomic architecture and introgression shape a butterfly radiation. *Science*
366:594-599.

17. Emsley MG 1965. Speciation in *Heliconius* (Lep., Nymphalidae): morphology and
geographic distribution. *Zoologica, New York* **50**:191–254.

18. Faria R, Johannesson K, Butlin RK, Westram AM (2019). Evolving inversions. *Trends in*
Ecology & Evolution **34**:239-248.

19. Freedman AH, Gronau I, Schweizer RM, Ortega-Del Vecchyo D, Han E, et al. (2012)
Genome Sequencing Highlights the Dynamic Early History of Dogs. *PLoS Genetics*
10:e1004016.

20. Glinka S, Ometto L, Mousset S, Stephan W, De Lorenzo D (2003) Demography and natural
selection have shaped genetic variation in Drosophila melanogaster: a multi-locus approach.
Genetics **165**:1269-1278.

21. Gronau I, Hubisz MJ, Gulko B, Danko CG, Siepel A (2011). Bayesian inference of ancient
human demography from individual genome sequences. *Nature Genetics* **43**:1031-1034.

22. Hackenberg M, Previti C, Luque-Escamilla PL, Carpena P, Martínez-Aroza J, Oliver JL.
(2006) CpGcluster: a distance-based algorithm for CpG-island detection. *BMC*
Bioinformatics **7**:446.

23. Hedrick PW, Tuttle EM, Gonser RA (2018) Negative-Assortative Mating in the White-
Throated Sparrow. *Journal of Heredity* **109**:223-231.

24. *Heliconius* Genome Consortium (2012). Butterfly genome reveals promiscuous exchange of
mimicry adaptations among species. *Nature* **487**: 94–8.

25. Jay P, Whibley A, Frézal L, Rodríguez de Cara MÁ, Nowell RW, Mallet J, Dasmahapatra
KK, Joron M. (2018). Supergene evolution triggered by the introgression of a chromosomal
inversion. *Current Biology* **28**:1839-1845.

26. Jay P, Chouteau M, Whibley A, Bastide H, Parrinello H, Llaurens V, Joron M. (2021).
Mutation load at a mimicry supergene sheds new light on the evolution of inversion
polymorphisms. *Nature Genetics* **53**:288-293.

27. Jiggins C, Naisbit R, Coe R, Mallet J 2001. Reproductive isolation caused by colour pattern
mimicry. *Nature* **411**:302–305.

28. Joron M, Wynne IR, Lamas G, Mallet J (1999) Variable selection and the coexistence of
multiple mimetic forms of the butterfly *Heliconius numata*. *Evol Ecol* **13**: 721–754.

29. Joron M, Papa R, Beltran M, Chamberlain N, Mavarez J, et al. (2006) A conserved
supergene locus controls colour pattern diversity in *Heliconius* butterflies. *PLoS Biology*

621 **4**:e303

622 30. Joron M, Frezal L, Jones RT, Chamberlain NL, Lee SF, Haag CR, Whibley A, Becuwe M,
623  Baxter SW, Ferguson L, Wilkinson PA, Salazar C, Davidson C, Clark R, Quail MA,
624  Beasley H, Glithero R, Lloyd C, Sims S, Jones MC, Rogers J, Jiggins CD, ffrench-Constant
625  RH (2011). Chromosomal rearrangements maintain a polymorphic supergene controlling
626  butterfly mimicry. *Nature* **477**:203–206.

627 31. Knoppien P (1985) Rare male mating advantage: a review. *Biological Reviews* **60**:81-117.

628 32. Lenormand T (2002) Gene flow and the limits to natural selection. *Trends in Ecology and*
629  *Evolution* **17**:183-189.

630 33. Lenth RV (2016). Least-Squares Means: The R Package lsmeans. *Journal of Statistical*
631  *Software* **69**:1-33.

632 34. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin
633  R; 1000 Genome Project Data Processing Subgroup (2009). The Sequence Alignment/Map
634  format and SAMtools. *Bioinformatics* **25**:2078-2079.

635 35. Lunter G, Goodson M (2011). Stampy: a statistical algorithm for sensitive and fast mapping
636  of Illumina sequence reads. *Genome Research* **21**:936-939.

637 36. Maisonneuve L., Chouteau M, Joron M, Llaurens V (2021). Evolution and genetic
638  architecture of disassortative mating at a locus under heterozygote advantage. *Evolution*
639  **75**:149-165.

640 37. Mallet J, McMillan W, Jiggins C (1998). Estimating the mating behavior of a pair of
641  hybridizing *Heliconius* species in the wild. *Evolution* **52**:503–510.

642 38. Martin SH, Möst M, Palmer WJ, Salazar C, McMillan WO, Jiggins FM, Jiggins CD (2016).
643  Natural Selection and Genetic Diversity in the Butterfly *Heliconius melpomene*. *Genetics*
644  **203**:525-541.

645 39. Mitchell-Olds T, Willis JH, Goldstein DB (2007). Which evolutionary processes influence
646  natural genetic variation for phenotypic traits? *Nature Reviews Genetics* **8**:845–856.

647 40. Muers, M (2009) Separating demography from selection, *Nature Reviews Genetics* **10**:280–
648  281.

649 41. Murray GGR, Soares AER, Novak BJ, Schaefer NK, Cahill JA, Baker AJ, Demboski JR,
650  Doll A, Da Fonseca RR, Fulton TL, Gilbert MTP, Heintzman PD, Letts B, McIntosh G,
651  O'Connell BL, Peck M, Pipes ML, Rice ES, Santos KM, Sohrweide AG, Vohr SH, Corbett-
652  Detig RB, Green RE, Shapiro B (2017). Natural selection shaped the rise and fall of
653  passenger pigeon genomic diversity. *Science* **358**:951-954.

654 42. McMillan W, Jiggins C, Mallet J (1997). What initiates speciation in passion-vine butterflies?
655  *Proceedings of the National Academy of Sciences of the USA* **94**:8628–8633.

656 43. Nadeau NJ, Pardo-Diaz C, Whibley A, Supple M A, Saenko SV, Wallbank RWR *et al*.
657  (2016). The gene *cortex* controls mimicry and crypsis in butterflies and moths. *Nature*,
658  **534**:106–110.

659 44. Nielsen, R., Hubisz, M.J., Hellmann, I., Torgerson, D., Andres, A.M., Albrechtsen, A.,
660  Gutenkunst R, Adams MD, Cargill M, Boyko A, Indap A, Bustamante CD, and Clark AG
661  (2009). Darwinian and demographic forces affecting human protein coding genes. *Genome*
662  *Research* **19**:838–849.

663 45.

46. Rosser N, Phillimore AB, Huertas B, Willmott KR, Mallet J (2012) Testing historical
explanations for gradients in species richness in heliconiine butterflies of tropical America.
Biological Journal of the Linnean Society **105**:479–497.

47. Schiffels S, Durbin R (2014) Inferring human population size and separation history from
multiple genome sequences. *Nature Genetics* **46**:919-925.

48. Saenko SV, Chouteau M, Piron-Prunier F, Blugeon C, Joron M, Llaurens V (2019)
Unravelling the genes forming the wing pattern supergene in the polymorphic butterfly
Heliconius numata. *EvoDevo* **10**:1-12.

49. Sheppard PM, Turner JRG, Brown KS, Benson WW, Singer MC (1985) Genetics and the
evolution of Muellerian mimicry in *Heliconius* butterflies. *Philosophical Transactions of the*
Royal Society of London, B Biological Sciences **308**: 433–610

50. Slatkin M (1987) Gene flow and the geographic structure of natural populations. *Science*
236:787-792

Figure 1 | Genetic and population structure at the P supergene.

A. Schematic phylogeny of the sampled species. It includes all members of the silvaniform clade and two outgroups, *H. melpomene* and *H. cydno*.

B. Schematic description of the genetic structure of the P supergene. Three chromosomal arrangements coexist in *H. numata* and are associated with different morphs.

C. Origin of *H. numata* specimens used for analyses and distribution of chromosome arrangements across the neotropics. Numbers in brackets indicate sampled specimens in each locality (the Tarapoto population lumps several neighbouring subsamples on the map)

Figure 2 | *H. numata* is characterised by low population structure.

A. Admixture plot for *H. numata*. The optimal cluster number for *H. numata* is two, and it splits *H. numata* into two categories, whereas they come from Atlantic forest or the Amazon. BR=Brazil (Atlantic), PR=Peru, VE=Venezuela, CO=Colombia, EC=Ecuador, FG=French Guiana. B. Principal component analysis computed on whole genome SNP. Color code match those given in the asmixture label of panel A. C. Relationship between genetic differentiation (F_{ST}/(1-F_{ST})) and logarithm of geographical distance. F_{ST} is measured between morphs/populations of the same species.

Figure 3 | Variation in present and past effective population size in *Heliconius* species

A. Variation in Pi in several *Heliconius* populations, showing higher genetic diversity in *H. numata* populations from the Amazon than other taxa. Population names indicates their origin as in Figure 2 (e.g. PR=Peru), with the addition of PA=Panama. The *H. numata* population with a lowest diversity is the one from the Atlantic forest (Brazil). **B.** Schematic representation of Gphocs results (presented in table S5-6). Gene flow was modelled but not represented graphically for clarity. showing that Amazonian populations of *H. numata*, which have the P supergene, show a dramatic increase in population size posterior to their split with the Atlantic populations of Brazil, which lack the supergene.

Figure 4 | Differences in synonymous nucleotide diversity (π_S) emerged at a genome-wide scale under divergent selection and mating region. Results from forward simulations of 10 populations undergoing local adaptation and different mating strategy are presented. Shown are levels of synonymous diversity obtained under assortative (A) versus disassortative mating (B) under different rates of migration and different local adaptation fitness. Each combination of parameters in brackets display the (dis)-assortative mating weight and the fitness value for local adaptation respectively. A left value of 0 in the bracket means complete assortative mating and 0.5 means no assortative mating or disassortative mating. Value of 1 means complete disassortative mating. A right value of 0 in the bracket means a fitness of 0 for non locally adapted individuals in a demes, A value of 0.5 means a reduced fitness of 0.5 relative to the maximum value.

Table 1: Estimates of demographic parameters for each best model in each comparison.
Biological parameters assumied a mutation rates of 2.9e-09 µ/bp/generation. NeAnc, = Effective
population sizes for the ancestral population, and descending population 1 and 2. NeParda, NePeru
and NeBrazil corresponds to effective population size of *H. pardalinus H. numata* in Peru and *H.*
numata in Brazil respectively. NeB1Parda and NeB2Peru, NeB1Brasil provides effective
population size for the population *H. pardalinus H. numata* in Peru and *H. numata* in Brazil
respectively after their expansion. Hrf = Hill-Robertson factor reflecting the reduction in effective
population size due to linked selection. Q = proportion of the genome with a reduced effective
population size. m12 = migration from population 2 into population 1. m21 = migration from
population 1 into population 2 (scaled according to 2N*ref*m_{ij}). T_{ANC} = Time of population size
change in the ancestral population, Tp1 = Time of size change in population 1 and Tp2 = Time of
size change in population 2. T_{ANC}, T_{SPLIT} and Tsc, Tp1, Tp2 are provided assuming for generations
per year.

H. Numata Perou - *H. pardalinus*		*H. numata* Perou - *H. numata* Brazil	
Best model	**IMA2NG**	Best. Model	**SCA2NG**
Log Lik.	-634.48	Log. Lik	-1430.36
Theta	16,757.39	Theta	3481.23
Nref	6,829,000	Nref	1,036,000
NeAnc	538,000 [0 – 1,281,000]	NeAnc	12,912,000[702,000 – 32,850,000]
NeParda	9,209,000 [3,163,000 – 15,254,000]	NeBrasil	83,000 [74,000 – 93,000]
NePeru	650,000 [625,000 – 676,000]	NePeru	60,373,000 [6,969,000 – 113,775,000]
NeB1Parda	25,233,000 [7,282,000 – 48,477,000]	NeB1Brasil	292,000 [147,000 – 465,000]
NeB2Peru	11,746,000 [6,511,000 – 17,368,000]	NeB2Peru	373,573,000 [13,310,000 – 1,190,748,000]
m12	1.99e-08 [0 – 8.59e-08]	m12	4.71E-08 [0 – 1.25e-07]
m21	1.37e-07 [9.84e-8 – 1.79e-7]	m21	3.42E-09 [0 – 2.19e-07]
T_{ANC}	3,237,000 [3,039,000 – 6,169,000]	T_{ANC}	501,000 [254,000 – 748,000]
T_{SPLIT}	1,892,000 [1,674,000 – 2,111,000]	T_{SPLIT}	110,000 [69,000 – 151,000]
$Tp1$	108,000 [662,000 – 1,500,000]	$Tp1$	11,000 [6,000 – 17,000]
$Tp2$	964,000 [779,000 – 1,150,000]	$Tp2$	55,000 [0 -157,000]
		T_{SC}	41,000 [7,000 – 75,000]
H_{RF}	0.10 [0.03 – 0.17]	H_{RF}	0.47 [0.43 – 0.50]
Q	0.37 [0 – 0.99]	Q	0.47 [0.39 – 0.56]

728 **List of Supplementary Materials:**

729 Table S1 to S7

730 Fig S1 to S5

731 Text S1