Coral assemblages at higher latitudes favour short-term potential over long-term performance

James Cant[1,2,*], James D. Reimer[3,4], Brigitte Sommer[5,6], Katie Cook[1], Sun W. Kim[7], Carrie A. Sims[7], Takuma Mezaki[8], Cliodhna O'Flaherty[1], Maxime Brooks[1], Hamish A. Malcolm[9], John M. Pandolfi[7], Roberto Salguero-Gómez[10,11,12 *,**], Maria Beger[1,11, *,**]

1. School of Biology, Faculty of Biological Sciences, University of Leeds, Leeds, LS2 9JT United Kingdom.
2. Department of Biology, Faculty of Science, University of Southern Denmark, 5230 Odense, Denmark.
3. Molecular Invertebrate Systematics and Ecology Laboratory, Graduate School of Engineering and Science, University of the Ryukyus, Nishihara, Okinawa, Japan.
4. Tropical Biosphere Research Centre, University of the Ryukyus, Nishihara, Okinawa, Japan.
5. School of Life Sciences, University of Technology Sydney, Ultimo, NSW 2007, Australia
6. School of Life and Environmental Science, The University of Sydney, Camperdown, NSW 2006, Australia.
7. Australian Research Council Centre of Excellence for Coral Reef Studies, School of Biological Sciences, The University of Queensland, Brisbane, QLD 4072, Australia.
8. Kuroshio Biological Research Foundation, Nishidomari, Otsuki-cho, Kochi, Japan.
9. Solitary Islands Marine Park, Department of Primary Industries, Coffs Harbour, NSW 2450, Australia.
10. Department of Zoology, University of Oxford, Oxford, 11a Mansfield Rd, OX1 3SZ United Kingdom.
11. Centre for Biodiversity and Conservation Science, School of Biological Sciences, University of Queensland, Brisbane, QLD, 4072 Australia.
12. Max Planck Institute for Demographic Research, Konrad Zuße Straße 1, 18057 Rostock, Germany.

[**]Share equal contribution

[*]Corresponding authors:

James Cant bsjic@leeds.ac.uk, Rob Salgeuro-Gómez rob.salguero@zoo.ox.ac.uk, Maria Beger M.Beger@leeds.ac.uk.

Author contributions: JC, RS-G and MB developed the research ideas. JC, JDR, BS, KC, SWK, CS, TM, CF, HAM, and MB collected field data, with JC, CF, MBr analysing the data. JC, RS-G, and MB led the writing of the manuscript with all authors contributing critically to the writing and giving final approval for publication.

Competing interests statement: All authors declare no competing interests

Keywords: amplification, demography, integral projection model (IPM), partial least squares analysis, subtropical, transient dynamics.

Abstract

The current exposure of species assemblages to high environmental variability may grant them

resilience to future increases in climatic variability. In globally threatened coral reef

ecosystems, management seeks to protect resilient reefs within variable environments. Yet, our

lack of understanding for the determinants of coral population performance within variable

environments hinders forecasting the future reassembly of coral communities. Here, using

Integral Projection Models, we compare the short- (*i.e.*, transient) and long-term (*i.e.*,

asymptotic) demographic characteristics of tropical and subtropical coral assemblages to

evaluate how thermal variability influences the structural composition of coral communities

over time. Exploring spatial variation across the dynamics of functionally different

competitive, stress-tolerant, and weedy coral assemblages in Australia and Japan, we show that

coral assemblages trade-off long-term performance for transient potential in response to

thermal variability. We illustrate how coral assemblages can reduce their susceptibility towards

environmental variation by exploiting volatile short-term demographic strategies, thus

enhancing their persistence within variable environments. However, we also reveal

considerable variation across the vulnerability of competitive, stress-tolerant, and weedy coral

assemblages towards future increases in thermal variability. In particular, stress-tolerant and

weedy corals possess an enhanced capacity for elevating their transient potential in response to

environmental variability. Accordingly, despite their current exposure to high thermal

variability, future climatic shifts threaten the structural complexity of coral assemblages,

derived mostly from competitive coral taxa within highly variable subtropical environments,

emulating the degradation expected across global coral communities.

Introduction

Coral reefs worldwide are threatened by unprecedented climatic and anthropogenic disruption (1, 2), with their conservation now reliant on enhancing the resilience of coral communities, specifically their capacity to resist and recover from increased climatic variability (3, 4). Changes in environmental regimes provoke spatial shifts in the performance and distribution of populations, which ultimately upscale to the compositional reassembly of biological communities (5, 6). Exposure to increased environmental variability is, however, expected to indirectly augment the capacity for populations to resist and recover from repeated disturbances (7, 8). Yet, nuanced relationships between population characteristics and biophysical conditions ensure inconsistent responses towards climate shifts, even across populations within the same region (9, 10). For instance, differential population sensitivities to habitat change can accelerate or reverse expected poleward range shifts in response to climate warming (10). Thus, anticipating the future resilience of natural communities requires understanding the collective vulnerabilities of their constituent populations (11) and the determinants underpinning their resilience to increasingly recurrent disturbances (12, 13). However, regional variations in the response of coral assemblages to climatic disturbances have generated misconceptions in our interpretation of the status of coral communities worldwide, with global population sizes of many species perhaps larger than previously anticipated (14). Only by linking the mechanisms underpinning heterospecific variation across the responses of populations to environmental variability can one predict the resilience of biological communities to increased climatic variability (11, 15, 16). Evidently, to accurately forecast the ongoing reassembly of global coral communities we must improve our understanding for how environmental variability shapes coral population performance across community- and regional-scales (7).

To explore the performance of populations exposed to recurrent disturbances within variable environments, their transient (*i.e.*, short-term) dynamics must be considered (17–20).

Yet, asymptotic (*i.e.*, long-term) population growth rate (λ), a metric that describes temporal

changes in population size at stationary equilibrium (21), is the predominant metric used to

quantify population performance (21, 22). In reality, though, natural populations are exposed

to repeated disturbances that favour the persistence of transient conditions, preventing the

emergence of stationary equilibria (17, 20, 23). Within variable environments, repeated

disturbances impose short-term changes upon the structure of populations that can elevate

(*amplify*) or diminish (*attenuate*) their growth rates, resulting in population performance

characteristics deviating from long-term expectations (18, 24). Quantifying how transient

population performance deviates from long-term expectations (henceforth *transient potential*)

is therefore crucial for predicting the success or failure of natural populations (25); an approach

that remains neglected within coral research (19).

Located at the intersection of tropical and temperate ecoregions, subtropical coral

communities provide an opportunity for evaluating the abiotic determinants of coral population

performance (26–28). Over recent decades, subtropical coral communities have undergone

transformation with various coral taxa undergoing poleward range expansions in response to

shifting thermal regimes (29–33). However, at higher latitudes, coral communities are exposed

to enhanced seasonality and cooler winter temperatures, and thus experience greater abiotic

variability relative to their tropical reef counterparts (34). Over time, the persistence of

populations within stable environments diminishes their capacity for tolerating novel

environmental states, thus reducing their resilience towards future climatic shifts (8).

Consequently, the endurance of subtropical coral communities within variable high-latitude

environments is expected to enhance their abiotic resilience (7). Indeed, as seen across other

communities (e.g., plants; 35), enhanced transient potential can elevate the performance of

acroporid coral populations within variable subtropical environments (36). Yet, without

knowledge for how differing coral assemblages utilise demographic strategies to mediate their

performance in response to environmental variability, we will be unable to forecast the impacts

of increasing climatic variability on the condition of global coral reef ecosystems.

In species rich communities, evaluating ecological dynamics requires a trait-based

approach to condense vast quantities of demographic detail (37). Given the diversity of coral

reef ecosystems, exploring patterns across the demographic characteristics of co-occurring

coral species presents a logistical challenge (38). Yet, this is a challenge that can be navigated

by pooling individuals based on shared trait characteristics. Morphological, physiological and

phenological functional traits influence the fitness of individuals and thus determine the

demographic characteristics of their populations (39), their responses to disturbances (40), and

subsequently the assembly of biological communities (41–43). Indeed, functional trait

characteristics impact upon the demographic properties of coral populations (e.g., colony

growth and reproduction; 44, 45), mediating their ability to respond to local abiotic patterns

(46). Given such strong links between coral traits and demographic performance, the

categorisation of coral taxa into competitive, stress tolerant, generalist and weedy life history

assemblages (sensu 47) is used to evaluate broadscale patterns in coral community reassembly

(48–50). The trait-based assessment of coral community assembly also offers greater insight

into the wider implications of ongoing community shifts than taxonomic-based assessments,

thereby aiding the management of coral reef ecosystems (50).

Here, we investigate how the performance characteristics of tropical and subtropical

coral populations map onto patterns of thermal variability across assemblages of competitive,

stress-tolerant, and weedy coral taxa. Using Integral Projection Models (IPMs; 51), we quantify

the association between different dimensions of thermal variability (monthly mean sea surface

temperature [SST], monthly SST variance, and monthly SST frequency spectrum) and the

transient potential and long-term performance characteristics of tropical and subtropical coral

assemblages in southern Japan and eastern Australia (Fig. 1). Specifically, we anticipate that,

in accordance with their exposure to elevated thermal variability, subtropical coral assemblages

will possess greater transient potential in comparison to tropical assemblages. We expect that

this pattern will persist irrespective of functional strategy, corresponding with the need for

subtropical coral populations to exploit periodically disturbed environments.

Results and Discussion

Our analyses reveal a trade-off between long-term performance and transient potential across

the examined coral assemblages. We find that this trade-off corresponds with the exposure of

coral populations to thermal variability along a gradient from warmer, more stable

environments to cooler, more variable conditions (Fig. 2). Using partial least squares

regression, we evaluated how patterns in the long-term performance, demographic recovery,

and transient potential, of coral populations conform with their exposure to abiotic variability.

We obtained estimates of long-term population performance (asymptotic population growth

rate, λ), demographic recovery (damping ratio [ρ], *i.e.*, a relative measure of the time needed

for a population to converge to a stable equilibrium; 21), and transient potential (transient

envelope [*TE*], *i.e.*, the difference between maximum and minimum population size following

disturbance; 24, 52) from IPMs depicting the dynamics of tropical and subtropical assemblages

of competitive, stress-tolerant, and weedy coral taxa in Japan and Australia (Fig. 1;

Supplementary S1 & S2). Furthermore, we quantified the exposure of these assemblages to

thermal variability using three measures of local SST regimes: monthly mean SST ($\bar{x}_{sst}$),

monthly SST variance (cv_{sst}), and monthly SST frequency spectrum (β_{sst}; Supplementary S3).

Notably, the coral assemblages exposed to more variable thermal conditions display

enhanced transient potential. Our partial least squares regression explains 92.17% of the

variance in the three measures of thermal exposure ($\bar{x}_{sst}$, cv_{sst}, and β_{sst}), whilst also capturing

37.43% of the variance in long-term performance (λ), demographic recovery (ρ), and transient

potential (*TE*; Fig. 2, $r^2_{[y]}$). The first partial least squares regression component reflects a

gradient in SST variability, capturing 60.97% of the variance in thermal conditions experienced

by our examined coral assemblages. It is along this component that divergent patterns within

estimates of λ and *TE* are most pronounced. Subsequently, estimates of *TE* are positively

correlated with the measures of thermal variability (cv_{sst}) and frequency spectrum (β_{sst}), whilst

higher λ estimates associate with warmer mean monthly SSTs ($\bar{x}_{sst}$; Fig. 2). Meanwhile,

damping ratio (ρ) estimates are aligned with the second partial least squares regression

component describing secondary patterns in the mean SST ($\bar{x}_{sst}$) and frequency (β_{sst}) variables.

Enhanced transient potential is thought to buffer the performance of populations in response to

elevated abiotic variability, thereby underpinning their capacity to exploit more variable

environments (35, 53). However, variation in transient potential across the assemblages of

differing coral taxa, in response to increased thermal variability (Fig. 3), suggests that exposure

to abiotic variability alone does not assure resilience towards future climatic variability.

The trade-off between long-term performance and transient potential does not manifest

consistently across the examined tropical and subtropical coral assemblages (Fig. 3A & Table

1), with inter-specific variation mediated instead by characteristics of population turnover (Fig.

3B & C). We explore inter-assemblage variation across estimates of long-term performance

and transient population growth potential. Again, we quantified long-term performance using

λ, whilst the *demographic stability index* (*DSI*) calculated from our IPMs provided a measure

of transient growth potential. A three-way ANOVA reveals significant interactions between

the three factors of assemblage classification (competitive, stress-Tolerant or weedy),

ecoregion (tropical *vs*. subtropical), and country (Australia vs. Japan; ANOVA$_\lambda$: $F_{2,11562}$ =

5698.47, $p < 0.001$; ANOVA$_{DSI}$: $F_{2,11581} = 589.8$, $p < 0.001$). Despite this, the tropical

assemblages routinely possess higher estimates of λ relative to their corresponding subtropical

counterparts (Tukey: $p < 0.001$ in all cases; Table 1). The one exception were weedy corals in

Japan, where λ is highest in the subtropics ($\lambda_{[t]}$ = 0.760 [95% CI: 0.750, 0. 770], $\lambda_{[s]}$ = 0.807

[0.802, 0.812]; p < 0.001). Alternatively, the subtropical coral assemblages typically possess a

greater capacity for amplifying population growth following a disturbance than the tropical

assemblages (Fig. 3A). However, this pattern is not consistent across life history strategies,

with competitive assemblages exhibiting the opposite trend in Australia (p < 0.001), and no

variation in Japan (p = 0.999).

The long-term performance and transient potential of the coral assemblages

corresponds with patterns in their generation time (Fig. 3B & C). To further evaluate the drivers

mediating coral population performance within variable environments we used Type 2 linear

regression (54) to explore the relationship between estimates of generation time (T, *i.e.*, the

time needed for individuals of a population to be replaced; 55), long-term performance (λ), and

transient potential (TE) calculated from our IPMs. Generation time is a strong predictor of long-

term population growth rate (r^2 = 0.704), with long-term performance increasing with

generation time (Fig. 3B). Conversely, longer generation times are associated with reduced

transient potential (Fig. 3C; r^2 = 0.409). Hence, our observed trade-off between long-term

performance and transient potential, in response to thermal variability, manifests inconsistently

across our examined tropical and subtropical coral assemblages, due to variation in their

characteristics of temporal population turnover.

Transient buffering in variable environments

Principally, a trade-off between long-term performance and transient potential implies that

long-term performance does not predict the capacity for populations to endure repeated

disturbances. Simultaneously, however, it also suggests that whilst enhanced transient potential

may enable natural populations to persist within variable environments, it comes at a cost to

their long-term performance. Historically, variability in population growth rate was thought to

diminish individual fitness (56), thus hindering the persistence of populations (57). This

understanding formed the basis of the demographic buffering hypothesis, whereby populations

can minimise the influence of environmental variability on their long-term performance by

limiting temporal variability in crucial vital rates (e.g., survival, development and reproduction;

58). Thus, variable environments were assumed to select for populations with the ability to

buffer key vital rates, thereby reducing temporal variation in performance characteristics (56,

58, 59). More recently, however, enhanced transient potential has been presented as an adaptive

mechanism that allows populations to exploit regions with high environmental variability (35).

Indeed, Ellis & Crone (53) demonstrated how increased transient potential can buffer the

effects of stochastic conditions on population growth rates, an effect that was increasingly

evident in populations possessing lower λ estimates. Thus, it is not unexpected, that coral

assemblages established within variable environments, would possess enhanced transient

potential (Fig. 2). Yet, if the vital rate schedules of these assemblages have indeed evolved

specifically to maximise their short-term performance, this would likely carry an energetic cost

to their long-term performance characteristics.

Our finding that transient potential is greatest in coral assemblages displaying reduced

long-term performance contrasts with previous work on mammals and plants showing a

positive association between population growth rates and transient potential (e.g., 60, 61).

Faster population growth rates are assumed of populations characterised by faster individual

development and high fecundity (62), with these populations also expected to exhibit greater

variability in size following disturbances (61). Whilst it is concerning that each of our surveyed

assemblages are in, or close to, a state of long-term decline ($\lambda < 1$; Table 1), projected long-

term performance is greatest in the tropical assemblages which also typically display lower

transient amplification in population growth (Fig. 3A). These findings agree with evidence that

transient potential is mediated by population turnover. Populations exhibiting longer

generation times typically display reduced temporal variability in size corresponding with the

fact that higher individual survival reduces the need to counteract disturbances (60); a pattern

that we show to be evident in our examined coral assemblages (Fig. 3C). Accordingly, our

observation that long-term performance is positively correlated with generation time in coral

populations (Fig. 3B), perhaps presents a more intriguing result. Although, with the long- and

short-term performance characteristics of coral assemblages corresponding with their relative

exposure to thermal variability (Fig. 2), this finding further implies that trading off long-term

performance for transient potential is an adaptive characteristic in coral populations.

Forecasting community reassembly

Here we show that stress-tolerant and weedy coral taxa possess more pronounced transient

amplification, highlighting a potential mechanism supporting their persistence at higher

latitudes. Using a subset of the dataset presented here, Cant *et al.* (36) recently demonstrated

how short-term increases in population growth following disturbance could support the

establishment of a subtropical coral assemblage of competitive *Acropora* spp. in southern

Japan. However, subtropical-tropical variation in the amplification capacity of competitive

coral assemblages appears minimal in comparison to the variation we observe across the stress-

tolerant and weedy assemblages in both Australia and Japan (Fig. 3A). Weedy corals comprise

species that exhibit small colony sizes, fast growth rates, and internal fertilisation, producing

larvae that settle quickly after release (47, 63). Together, these strategies support faster

population turnover, enabling weedy coral species to proliferate within highly disturbed

environments (64). Conversely, stress-tolerant corals display slower growth rates, longer life

expectancies, high fecundity, and broadcast spawning strategies (47, 65). The larger, more

robust, morphologies associated with stress-tolerant coral taxa maximise energy storage,

promoting their persistence within challenging environments (66). Meanwhile, longer lifespans

and elevated fecundity allow stress-tolerant corals to endure abiotic variation by taking

advantage of sporadic improvements in local conditions (47). Consequently, our findings

support existing projections that weedy and stress-tolerant coral taxa are likely to become

increasingly prevalent throughout disturbed coral communities (67, 68). However, these

projections herald the future loss of the structural complexity considered essential to the

functioning of reef ecosystems (69).

Crucially, our findings do not reflect the current reality for many coral assemblages

within regions of high environmental variability, suggesting that the composition of coral

communities is not solely mediated by the interplay between transient dynamics and abiotic

variability. Despite the lower amplificatory capacity we report for subtropical competitive

corals compared to the subtropical weedy and stress-tolerant assemblages, competitive coral

taxa dominate many subtropical coral assemblages (70–72). Utilising fast growth strategies,

colonies of competitive coral taxa are capable of rapidly colonising available substrate, quickly

outcompeting heterospecific colonies for both space and light (47). Whilst this competitive

nature perhaps explains the enhanced amplificatory capacity of the tropical competitive

assemblages relative to the tropical stress-tolerant and weedy assemblages (Fig. 3A), the

sensitivity of many competitive coral taxa to environmental change means that these

assemblages are often regarded as early successional, dominating only within optimal

environments and receding as reef ecosystems approach climax states (73, 74). Within

subtropical environments, however, coral community composition is mediated by

environmental pressures and dispersal barriers that filter the occurrence of species according

to their trait characteristics (46, 75). As a result, subtropical coral assemblages typically consist

of a subset of tropical species found on tropical coral reefs (46), as well as subtropical

specialists and endemics. The dominance of competitive coral taxa within subtropical coral

assemblages, despite their reduced transient performance relative to other coral taxa, therefore,

implies that competitive interactions profoundly influence the performance of coral

populations (76, 77). Certainly, further investigation into the influence of competitive

interactions on the transient dynamics of coral populations is needed to disentangle how the

dynamics of coexistence between coral populations facilitate their persistence within variable

environments.

Conclusions

Limitations in our understanding for the abiotic determinants driving the dynamics of coral

assemblages inhibits our capacity to forecast their future performance and, therefore, manage

global coral community reassembly (78–80). Here, we demonstrate how coral assemblages

within regions of high environmental variability exhibit demographic strategies associated with

enhanced transient potential, but at a cost to their long-term performance (Fig. 2). Climatic

change is exposing coral communities worldwide to increased abiotic variability. Crucially,

our findings here emphasize that whilst coral assemblages can adopt demographic strategies to

enhance their viability when exposed to abiotic variability, the winners and losers within future,

more variable environments cannot be predicted from existing measures of long-term

performance. However, the relationship that we observed between transient potential and

thermal variability was not universal across coral taxa, nor did it manifest identically across

hemispheres. Subtle patterns in the association between population dynamics and their climate

drivers hinder predictions of the consequences of environmental change within biological

communities (81). Nevertheless, relative to competitive coral taxa, weedy and stress-tolerant

corals appear to possess a greater capacity for enduring within environments characterised by

repeated abiotic disturbances. Yet, competitive coral taxa are often associated with more

complex morphologies and therefore support the structural complexity critical to the wider

functioning of coral associated ecosystems (69). Accordingly, future increases in abiotic

variability threaten the viability of coral associated ecosystems.

Methods

Modelling population dynamics

Integral Projection Models (IPMs) capture the influences of the state composition of

individuals on the performance of populations over discrete time periods (*t* to *t+1*; 51). Here,

to quantify the long-term performance characteristics and transient (*i.e.*, short-term) potential

of coral populations, we used IPMs describing patterns in colony survival (σ), transitions in

size (growth and shrinkage, γ), fragmentation probability (κ), fecundity (φ), and recruitment

(ϕ), each as a function of colony size (*z*; visible horizontal surface area, cm^2). Specifically, our

IPMs took the form

311
$$n[z', t + 1] = \int_{L}^{U} (P_{z'z} + F_{z'z}\, n[z, t]\, \delta z) \tag{1}$$

312
$$P_{z'z} = (1 - \kappa_z)\, \sigma_z\, \gamma_{z'z} + (\kappa_z\, \kappa b_z\, \kappa_z^0) \tag{2}$$

313
$$F_{z'z} = \varphi_z\, \phi C_0 \tag{3}$$

with [L, U] representing the range of possible colony sizes; calculated as 10% above and below

observed maximum and minimum colony sizes to avoid accidental exclusion (82).

Accordingly, the structure of a population at time *t+1* (*n[z', t+1]*) is a product of its structure

at time *t* (*n[z', t]*) subject to the survival (σ_z) and transition of individual colonies from size *z*

to size *z'* ($\gamma_{z'z}$); the probability of colony fragmentation (κ_z) and the number (κb_z) and size

distribution of any colony remnants produced (κ_z^0); and colony fecundity (φ_z) combined with

the probability of successful recruitment (ϕ) and the size distribution of surviving recruits (C$_0$).

Data Collection

We parameterised our IPMs using data collected during repeated annual surveys of 3171 tagged

colonies within tropical and subtropical coral communities in southern Japan and eastern

Australia, conducted between 2016 and 2019 (Fig. 1; Supplementary S1). We tagged individual

colonies using permanent plots arranged haphazardly throughout four focal coral communities

(Australian subtropics [AS], Australian tropics [AT], Japanese subtropics [JS], Japanese

tropics [JT]) and demarcated with numbered tags (36, 68). All tagged colonies were identified

in situ to the lowest possible taxonomic level (either genus or species). No samples were taken

from tagged colonies, as although this would have allowed us to resolve species identity, we

wanted to avoid any lasting interference with the processes of colony survival, growth, and

fragmentation.

To facilitate comparing population characteristics observed across spatially distinct

regions in Australia and Japan with varying degrees of species overlap (83), we grouped tagged

colonies across each region according to shared life-history-strategies (sensu 47–49), primarily

delineated based on their morphology, growth rate and reproductive mode (47). Specifically,

we categorised colonies as 'competitive', 'weedy', 'stress-tolerant' or 'generalist' following

the genera classifications of Darling *et al*. (47), with minor adaptions made based on local

expertise (see supplementary S2 for a detailed list). In the event that genera represented species

classified across different categories (19 cases), we randomly assigned individuals across the

relevant categories in proportion with the number of species within each category known to

occur in the area (sensu 49). Following the pooling of colonies according to their life-history-

strategies, we omitted all individuals defined as generalists from subsequent analyses due to

their limited representation across our regional samples (n: AS = 22 colonies; AT = 31; JS =

17; JT = 65). Consequently, we constructed IPMs concerning the dynamics of 12 coral

assemblages corresponding with competitive, stress-tolerant, and weedy coral taxa across four

geographical locations (Fig. 1).

Photographs capturing the visible horizontal extent of tagged colonies were used to

follow individuals over successive surveys and obtain longitudinal records of colony surface

area (cm^2; transformed to a $\log_{10}$ scale) over time. Using generalised linear mixed models

(GLMMs), we estimated size-specific patterns in colony survival (σ), transitions in size (γ),

and fragmentation probability (κ) for each population (Supplementary S1). In each case, our

GLMMs included random effects (colony identity and survey location) to account for any

autocorrelation between observations and within-subject variability associated with our

pooling of data recorded from individuals followed across multiple years, and at different sites.

Colony survival (σ) reflected the continued presence of tagged individuals across survey

intervals (*t* to *t+1*) and was modelled as a logistic function of colony surface area at time *t*.

Colony size transitions (γ), representing both growth through colony extension, and shrinkage

through partial mortality (84), were modelled using the polynomial relationship between initial

colony surface area at time *t* and subsequent surface area at time *t+1*. Colony fragmentation

probability (κ) was then modelled as a polynomial logistic function of colony size at time *t*.

During our surveys, we recorded fragmentation in the event of observed colony breakage,

recording the size (surface area, cm^2) of all remnants produced in each case. Subsequently, we

also modelled the number (κb_z) and size (κ_Z^0) of remnant colonies produced during

fragmentation as a function of colony size at time *t*, using Poisson and polynomial GLMMs,

respectively.

Alongside our surveys of tagged individual colonies, we also monitored colony

recruitment within our permanent coral plots. During each annual survey, we recorded the

number and size of new colonies appearing within each plot. These recruitment counts enabled

us to quantify annual and regional variability in recruit densities (Table S2), as well as estimate

population-specific recruit size distributions (C_0; Supplementary S1). However, prior to

parameterising recruitment dynamics within our IPMs, we determined patterns in colony

fecundity (φ). This approach was necessary because evaluating population performance

requires an explicit consideration of fecundity to link the dynamics of existing individuals with

the introduction of new, genetically distinct individuals (21). Using data relating colony size

and larval output (larval density, cm^3) extracted from the Coral Trait Database (85, 86), we

calculated colony fecundity (φ) as the polynomial relationship between colony size at *t* and

expected larval output (Supplementary S1). Next, to ensure our modelled recruitment dynamics

accurately reflected our empirical observations we parameterised a recruit survival function (ϕ)

within our IPMs. Here, recruit survival (ϕ) serves as a probability function converting expected

larval output into a proportional contribution of observed recruit counts as a function of colony

size, which we calculated by dividing total expected larval output in any given year by the

corresponding annual recruitment count (Supplementary S1, *sensu* 68, 87).

Quantifying population characteristics

From our IPMs, we obtained estimates of long-term performance (asymptotic population

growth, λ), generation time (*T*), and transient potential (damping ratio [ρ], maximal

amplification [$\bar{\rho}_{max}$] & transient envelope [*TE*]) for each tropical and subtropical coral

assemblage (21, 24, 52, 55, 88). Estimates of λ are typically used as a measure of long-term

population viability (22), and reflect whether a population is expected to grow ($\lambda > 1$) or decline

($\lambda < 1$) when at stationary equilibrium (21). Generation time is a measure of population

turnover, describing the time needed for individuals of a population to be replaced (55).

Alternatively, the measures of transient potential describe the expected characteristics of

populations following their displacement from stationary equilibrium due to disturbances. The

damping ratio constitutes a measure of demographic recovery (52, 89), describing the rate at

which a population perturbed from its stationary equilibrium converges back to its asymptotic

growth trajectory (21). Meanwhile, maximal amplification quantifies the greatest increase in

population size following a disturbance, relative to its asymptotic growth trajectory (24, 88).

Finally, the transient envelope quantifies the magnitude by which the transient dynamics of a

population deviates from its long-term trajectory (52).

To calculate the aforementioned demographic characteristics, we discretised our IPMs

into large matrices. We attained these matrices by applying the 'midpoint rule' to integrate

each IPM into a high-dimension matrix (200×200 cells), with the probability of transitioning

from one cell to the next approximated at the cell midpoint and multiplied by the cell width as

per (90). Estimates of λ were then identified as the dominant eigenvalue of each discretised

matrix, whilst we estimated damping ratios as the ratio between the subdominant and dominant

eigenvalues. With the *R* package *Rage* (91) we then calculated generation time using estimates

of net reproductive rate (R_0) and λ obtained from each matrix,

$$T = \log(R_0) - \log(\lambda). \tag{4}$$

408   Next, we determined the transient envelope of each assemblage using their associated Kreiss

409   bounds of amplification ($\overline{K}_\lambda^*$) and attenuation ($\underline{K}_\lambda^*$),

$$TE = \overline{K}_\lambda^* - \underline{K}_\lambda^*. \tag{5}$$

Respectively, the Kreiss bounds of amplification and attenuation reflect the largest and smallest

expected long-term density of a population following the dissipation of transient conditions,

relative to its asymptotic growth trajectory (92–94). We acknowledge here that this definition

is more commonly applied to measures of population inertia (24), which are more typically

used in estimating transient envelopes (52). However, Kreiss bound estimates have been

demonstrated to align with corresponding estimates of population inertia and, unlike estimates

of population inertia, are not sensitive to imprimitive population models (*i.e.*, non-negative

models permitting transitions between all state classes, but with transitions between certain

stages occurring only at periodic intervals; 21, 24); hence their selection here. We derived these

Kreiss bounds, alongside estimates of maximal amplification, using their corresponding

functions in the *R* package *popdemo* (95).

Across each demographic measure, we determined the variance in our assemblage-

specific estimates through Jack-knife resampling. During resampling, we generated 1,000 IPM

variants for each assemblage, each time using 95% of our original data sample without

replacement, whilst permitting recruit survival probabilities (ϕ) to vary within observed limits.

Finally, prior to their inclusion in further analyses, the jack-knifed distributions of the λ,

generation time, transient envelope, and maximal amplification variables required

transforming to ensure approximate normality. We omitted 26 variants for which $\lambda > 2$, as these

presented unrealistic illustrations of population performance (*i.e.*, more than doubling

population size every year), before applying a log transformation to the generation time

variable and a power transformation (y^x) across the damping ratio ($y^{-2.0}$), transient envelope ($y^{-0.1}$)

$^{0.1}$) and maximal amplification variables ($y^{-0.5}$).

Evaluating spatial trends in population characteristics

To test for patterns in the spatial variation of long-term performance and transient potential

across tropical and subtropical coral assemblages, we utilised partial least squares regression,

ANOVA, and Type 2 linear regression. Initially, we applied a partial least squares regression

to test whether trade-offs between the long-term performance characteristics and transient

potential of coral assemblages align with their exposure to abiotic variability. Partial least

squares regression quantifies the association between multiple predictor variables and one or

more dependant variables (96). Subsequently, using this technique we simultaneously

evaluated the relationships between mean estimates of λ, damping ratio, and transient envelope

obtained for each assemblage, and their correlation with patterns in thermal conditions to

provide an insight into the demographic trade-offs of coral assemblages and their mechanistic

drivers.

To evaluate how abiotic variability mediates the trade-off between the short- and long-

term performance characteristics of coral assemblages, within our partial least squares

regression we quantified the abiotic conditions experienced by each coral assemblage using

three measures of local sea surface temperature (SST) regimes: mean monthly SST ($\bar{x}_{sst}$),

monthly SST variance (cv_{sst}), and monthly SST frequency spectrum (β_{sst}; Supplementary S3).

Focusing on the four geographical regions in which our focal coral assemblages were surveyed

(GPS: AS = -30.3°, 153.1°; AT = -23.4°, 151.9°; JS = 32.8°, 132.6°; JT = 26.5°, 128.1°; Fig.

1), we extracted high resolution monthly SST readings (°C; overlaid on a 1° latitude-longitude

grid) taken between January 1950 and December 2019, inclusive, from the HadISST dataset

(97). Arranging these SST records into 69-year timeseries for each location, we then calculated

the mean ($\bar{x}_{sst}$) and coefficient of variance (cv_{sst}) for each timeseries. Next, we estimated the

frequency spectrum of each time series. Spectral analysis is used to quantify the periodicity of

recurrent variability within a timeseries, with higher frequencies associated with shorter-term

fluctuations (98). The frequency spectrum of a time series is represented by its spectral

exponent (β) and equal to the negative slope between its log spectral density and log frequency

(99), which we calculated using the package *stats* (100). After testing these abiotic predictor

variables for collinearity (Supplementary S3), we performed our partial least square regression

analyses using the *R* package *plsdepot* (101).

Next, we assessed how patterns in the long-term performance, and capacity for coral

assemblages to benefit from recurrent disturbance vary between tropical and subtropical

regions, and how this variation manifests across coral taxa. Using a three-way ANOVA, we

separately investigated variation in estimates of λ and maximal amplification across the three

factors of country (Australia *vs*. Japan), ecoregion (tropical *vs*. subtropical), and assemblage

classification (competitive, stress-tolerant or weedy). With maximal amplification estimates

inverted during transformation, larger values subsequently reflect reduced amplification

potential. For the purposes of clarity in this analysis, we will henceforth refer to this reversed

scale as a *demographic stability index* (*DSI*), with lower values corresponding with enhanced

amplification. Finally, we evaluated drivers of long- and short-term performance, by using

Type 2 linear regression to separately evaluate the relationship between generation time (*T*)

and estimates of λ and transient envelope (*TE*). Type 2 linear regression is an approach for

quantifying the relationship between two non-independent variables, such that both variables

include an element of error (54). Here, due to differences in the magnitude of the variance (σ^2)

across our variables of generation time, λ, and transient envelope (σ^2: *T* = 1.139; λ = 0.009; *TE*

= 0.016) we performed a Ranged Major Axis regression using the *R* package *lmodel2* (102).

Acknowledgements

The authors would like to thank S. Dalton, L. Lachs, C. Kim, R. Edgar, K.-L. Gomez-Cabrera,

N. Kyriacou, I. Mizukami, H. Kise, C. Fourreau, G. Masucci, P. Biondi, S. Nishihira, M.

Tamae, H. Nakakoji, C. Tan, L. Lawrence, T. Hofmann, I. Montero-Serra, and all staff at Dive

Quest, Pacific Marine and SeaAir for their assistance with field data collection. Additionally,

aspects of this research would not have been possible without the research permits provided by

the Solitary Islands Marine Park branch of the NSW Department of Primary Industries (SIMP

2016/002V2, MEAA 20/45) and the Great Barrier Reef Marine Park Authority (G19/42221.1).

Funding for this research was provided by a Natural Environment Research Council (NERC)

Doctoral Training Programme Scholarship and a Royal Geographical Society Ralph Brown

Expedition Award (RBEA 03/19) to JC, the Australian Research Council Centre of Excellence

for Coral Reef Studies (CE140100020) to JMP and others, the Australian Research Council

Centre of Excellence for Environmental Decisions (CE110001014), a British Ecological

Society small grant, the Winifred Violet Scott Charitable Trust, and the European Union's

Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant

agreement TRIM-DLV-747102 to MB. BS was supported by a Chancellor's Postdoctoral

Research Fellowship from the University of Technology Sydney and a University of Sydney

Fellowship.

References

1.   T. P. Hughes, *et al.*, Spatial and temporal patterns of mass bleaching of corals in the

Anthropocene. *Science* **359**, 80–83 (2018).

2.   T. P. Hughes, *et al.*, Global warming impairs stock–recruitment dynamics of corals.

Nature **568**, 387–390 (2019).

3.   J. Kleypas, *et al.*, Designing a blueprint for coral reef survival. *Biological*

Conservation **257**, 1–7 (2021).

4.   E. McLeod, E. C. Shaver, M. Beger, J. Koss, G. Grimsditch, Using resilience

assessments to inform the management and conservation of coral reef ecosystems.

Journal of Environmental Management **277**, 1–8 (2021).

5.   G. Pecl, M. B. Araujo, J. D. Bell, J. Blanchard, T. C. Bonebrake, Biodiversity

redistribution under climate change: Impacts on ecosystems and human well-being

Publication Details. *Science* **355**, 1–9 (2017).

6.   Ø. Totland, J. Nyléhn, Assessment of the effects of environmental change on the

performance and density of Bistorta vivipara: The use of multivariate analysis and

experimental manipulation. *Journal of Ecology* **86**, 989–998 (1998).

7.   E. B. Rivest, S. Comeau, C. E. Cornwall, The role of natural variability in shaping the

response of coral reef organisms to climate change. *Current Climate Change Reports*

3, 271–281 (2017).

8.   P. W. Boyd, *et al.*, Biological responses to environmental heterogeneity under future

ocean conditions. *Global change biology* **22**, 2633–2650 (2016).

9.   C. Parmesan, G. Yohe, A globally coherent fingerprint of climate change. *Nature* **421**,

37–42 (2003).

10.   I. C. Chen, J. K. Hill, R. Ohlemüller, D. B. Roy, C. D. Thomas, Rapid range shifts of

species associated with high levels of climate warming. *Science* **333**, 1024–1026

(2011).

11.   S. E. Williams, L. P. Shoo, J. L. Isaac, A. A. Hoffmann, G. Langham, Towards an

integrated framework for assessing the vulnerability of species to climate change.

PLoS biology **6** (2008).

12.   D. P. Vázquez, E. Gianoli, W. F. Morris, F. Bozinovic, Ecological and evolutionary

impacts of changing climatic variability. *Biological Reviews* **92**, 22–42 (2017).

13.   D. T. Iles, R. F. Rockwell, D. N. Koons, Shifting vital rate correlations alter predicted

population responses to increasingly variable environments. *American Naturalist* **193**,

E57–E64 (2019).

14.   A. Dietzel, M. Bode, S. R. Connolly, T. P. Hughes, The population sizes and global

extinction risk of reef-building coral species at biogeographic scales. *Nature Ecology*

and Evolution **5**, 663–669 (2021).

15.   W. B. Foden, *et al.*, Identifying the world's most climate change vulnerable species: A

537   systematic trait-based assessment of all birds, amphibians and corals. *PLoS ONE* **8**

538   (2013).

539   16.   T. P. Dawson, S. T. Jackson, J. I. House, I. C. Prentice, G. M. Mace, Beyond

540   predictions: Biodiversity conservation in a changing climate. *Science* **332**, 53–58

541   (2011).

542   17.   A. Hastings, Transients: The key to long-term ecological understanding? *Trends in*

543   *Ecology and Evolution* **19**, 39–45 (2004).

544   18.   T. H. G. Ezard, *et al.*, Matrix models for a changeable world: The importance of

545   transient dynamics in population management. *Journal of Applied Ecology* **47**, 515–

546   523 (2010).

547   19.   J. Cant, R. Salguero-Gómez, M. Beger, Transient demographic approaches can

548   drastically expand the toolbox of coral reef science. *bioRxiv*, 2021.04.08.438926

549   (2021).

550   20.   A. Hastings, *et al.*, Transient phenomena in ecology. *Science* **361**, 1–9 (2018).

551   21.   H. Caswell, *Matrix population models: Construction, analysis and interpretation*, 2nd

552   Edition (Oxford University Press Inc, 2001).

553   22.   E. E. Crone, *et al.*, How do plant ecologists use matrix population models? *Ecology*

554   *Letters* **14**, 1–8 (2011).

555   23.   A. Hastings, Transient dynamics and persistence of ecological systems. *Ecology*

556   *Letters* **4**, 215–220 (2001).

557   24.   I. Stott, S. Townley, D. Hodgson, A framework for studying transient dynamics of

558   population projection matrix models. *Ecology Letters* **14**, 959–970 (2011).

25.   D. N. Koons, J. B. Grand, B. Zinner, R. F. Rockwell, Transient population dynamics:

Relations to life history and initial population state. *Ecological Modelling* **185**, 283–

297 (2005).

26.   E. F. Camp, *et al.*, The future of coral reefs subject to rapid climate change: Lessons

from natural extreme environments. *Frontiers in Marine Science* **5**, 1–21 (2018).

27.   J. A. Burt, *et al.*, Insights from extreme coral reefs in a changing world. *Coral Reefs*

39, 495–507 (2020).

28.   M. Beger, B. Sommer, P. L. Harrison, S. D. A. Smith, J. M. Pandolfi, Conserving

potential coral reef refuges at high latitudes. *Diversity and Distributions* **20**, 1–13

(2014).

29.   W. F. Precht, R. B. Aronson, Climate flickers and range shifts of reef corals. *Front*

Ecol Environ **2**, 307–314 (2004).

30.   C. A. Tuckett, T. de Bettignies, J. Fromont, T. Wernberg, Expansion of corals on

temperate reefs: Direct and indirect effects of marine heatwaves. *Coral Reefs* **36**, 947–

956 (2017).

31.   H. Yamano, K. Sugihara, K. Nomura, Rapid poleward range expansion of tropical reef

corals in response to rising sea surface temperatures. *Geophysical Research Letters* **38**,

1–6 (2011).

32.   A. H. Baird, B. Sommer, J. S. Madin, Pole-ward range expansion of Acropora spp.

along the east coast of Australia. *Coral Reefs* **31**, 1063 (2012).

33.   D. J. Booth, J. Sears, Coral expansion in Sydney and associated coral-reef fishes.

Coral Reefs **37**, 995 (2018).

34.   B. Sommer, M. Beger, P. L. Harrison, R. C. Babcock, J. M. Pandolfi, Differential

582   response to abiotic stress controls species distributions at biogeographic transition

583   zones. *Ecography* **41**, 478–490 (2018).

584   35.   J. L. McDonald, I. Stott, S. Townley, D. J. Hodgson, Transients drive the demographic

585   dynamics of plant populations in variable environments. *Journal of Ecology* **104**, 306–

586   314 (2016).

587   36.   J. Cant, *et al.*, Transient amplification enhances the persistence of tropicalising coral

588   populations in marginal high latitude environments. *BioRxiv*, 2021.04.23.441133

589   (2021).

590   37.   L. Chalmandrier, *et al.*, Linking functional traits and demography to model species-

591   rich communities. *Nature Communications* **12**, 1–9 (2021).

592   38.   J. S. Madin, *et al.*, A trait-based approach to advance coral reef science. *Trends in*

593   *Ecology and Evolution* **31**, 419–428 (2016).

594   39.   C. Violle, *et al.*, Let the concept of trait be functional! *Oikos* **116**, 882–892 (2007).

595   40.   J. P. Grime, S. Pierce, *The evolutionary strategies that shape ecosystems* (John Wiley

596   and Sons Ltd., 2012).

597   41.   B. J. McGill, B. J. Enquist, E. Weiher, M. Westoby, Rebuilding community ecology

598   from functional traits. *Trends in Ecology and Evolution* **21**, 178–185 (2006).

599   42.   M. W. Cadotte, K. Carscadden, N. Mirotchnick, Beyond species: Functional diversity

600   and the maintenance of ecological processes and services. *Journal of Applied Ecology*

601   **48**, 1079–1087 (2011).

43.   D. S. Falster, Å. Brännström, M. Westoby, U. Dieckmann, Multitrait successional

forest dynamics enable diverse competitive coexistence. *Proceedings of the National*

Academy of Sciences of the United States of America **114**, E2719–E2728 (2017).

44.   J. S. Madin, M. O. Hoogenboom, S. R. Connolly, Integrating physiological and

biomechanical drivers of population growth over environmental gradients on coral

reefs. *Journal of Experimental Biology* **215**, 968–976 (2012).

45.   M. Álvarez-Noriega, *et al.*, Fecundity and the demographic strategies of coral

morphologies. *Ecological Society of America* **97**, 3485–3493 (2016).

46.   B. Sommer, P. L. Harrison, M. Beger, J. M. Pandolfi, Trait-mediated environmental

filtering drives assembly at biogeographic transition zones. *Ecology* **95**, 1000–1009

(2014).

47.   E. S. Darling, L. Alvarez-Filip, T. A. Oliver, T. R. Mcclanahan, I. M. Côté, Evaluating

life-history strategies of reef corals from species traits. *Ecology Letters* **15**, 1378–1386

(2012).

48.   E. S. Darling, T. R. McClanahan, I. M. Côté, Life histories predict coral community

disassembly under multiple stressors. *Global Change Biology* **19**, 1930–1940 (2013).

49.   J. Zinke, *et al.*, Gradients of disturbance and environmental conditions shape coral

community structure for south-eastern Indian Ocean reefs. *Diversity and Distributions*

24, 605–620 (2018).

50.   E. S. Darling, *et al.*, Social–environmental drivers inform strategic management of

coral reefs in the Anthropocene. *Nature Ecology and Evolution* **3**, 1341–1350 (2019).

51.   M. R. Easterling, S. P. Ellner, P. M. Dixon, Size-specific sensitivity: applying a new

structured population model. *Ecology* **81**, 694–708 (2000).

52.   P. Capdevila, I. Stott, M. Beger, R. Salguero-Gómez, Towards a comparative

626   framework of demographic resilience. *Trends in Ecology & Evolution* **35**, 776–786

627   (2020).

628   53.   M. M. Ellis, E. E. Crone, The role of transient dynamics in stochastic population

629   growth for nine perennial plants. *Ecology* **94**, 1681–1686 (2013).

630   54.   R. R. Sokal, F. J. Rohlf, *Biometry: The principles and practice of statistics in*

631   *biological research*, 3rd Editio (W. H. Freeman & Co, 1995).

632   55.   J. Gaillard, *et al.*, Generation Time: A reliable metric to measure life-history

633   variation among mammalian populations. *The American Naturalist* **166**, 119–123

634   (2005).

635   56.   C. A. Pfister, Patterns of variance in stage-structured populations: Evolutionary

636   predictions and ecological implications. *Proceedings of the National Academy of*

637   *Sciences* **95**, 213–218 (1998).

638   57.   R. Lande, Risks of population extinction from demographic and environmental

639   stochasticity and random catastrophes. *American Naturalist* **142**, 911–927 (1993).

640   58.   W. F. Morris, D. F. Doak, Buffering of life histories against environmental

641   stochasticity: Accounting for a spurious correlation between the variabilities of vital

642   rates and their contributions to fitness. *American Naturalist* **163**, 579–590 (2004).

643   59.   C. H. Hilde, *et al.*, The demographic buffering hypothesis: Evidence and challenges.

644   *Trends in Ecology and Evolution* **35**, 523–538 (2020).

645   60.   W. F. Morris, *et al.*, Longevity can buffer plant and animal populations against

646   changing climatic variability. *Ecology* **89**, 19–25 (2008).

647   61.   M. Gamelon, *et al.*, Influence of life-history tactics on transient dynamics: A

648   comparative analysis across mammalian populations. *American Naturalist* **184**, 673–

649   683 (2014).

650   62.   M. K. Oli, The fast-slow continuum and mammalian life-history patterns: An empirical

651   evaluation. *Basic and Applied Ecology* **5**, 449–463 (2004).

652   63.   N. Knowlton, The future of coral reefs. *PNAS* **98**, 5419–5425 (2001).

653   64.   M. Adjeroud, *et al.*, Recovery of coral assemblages despite acute and recurrent

654   disturbances on a South Central Pacific reef. *Scientific Reports* **8** (2018).

655   65.   C. N. Klepac, D. J. Barshis, Reduced thermal tolerance of massive coral species in a

656   highly variable environment: Reduced heat tolerance of massive corals. *Proceedings of*

657   *the Royal Society B: Biological Sciences* **287**, 19–21 (2020).

658   66.   R. van Woesik, A. Irikawa, R. Anzai, T. Nakamura, Effects of coral colony

659   morphologies on mass transfer and susceptibility to thermal stress. *Coral Reefs* **31**,

660   633–639 (2012).

661   67.   Y. Loya, *et al.*, Coral bleaching: the winners and the losers. *Ecology Letters* **4**, 122–

662   131 (2001).

663   68.   J. Cant, *et al.*, The projected degradation of subtropical coral assemblages by recurrent

664   thermal stress. *Journal of Animal Ecology* **90**, 233–247 (2021).

665   69.   N. A. J. Graham, K. L. Nash, The importance of structural complexity in coral reef

666   ecosystems. *Coral Reefs* **32**, 315–326 (2013).

667   70.   V. J. Harriott, P. L. Harrison, S. A. Banks, The coral communities of Lord Howe

668   Island. *Marine and Freshwater Research* **46**, 457–465 (1995).

71.   Y. Nozawa, M. Tokeshi, S. Nojima, Structure and dynamics of a high-latitude

scleractinian coral community in Amakusa, southwestern Japan. *Marine Ecology*

Progress Series **358**, 151–160 (2008).

72.   K. Sugihara, *et al.*, Latitudinal changes in hermatypic coral communities from west

Kyushu to Oki Islands in Japan. *Journal of the Japanese Coral Reef Society* **11**, 51–67

(2009).

73.   S. K. Wilson, J. P. W. Robinson, K. Chong-Seng, J. Robinson, N. A. J. Graham, Boom

and bust of keystone structure on coral reefs. *Coral Reefs* **38**, 625–635 (2019).

74.   H. Ohba, K. Hashimoto, K. Shimoike, T. Shibuno, Y. Fujioka, Secondary succession

of coral reef communities at Urasoko Bay , Ishigaki Island , the Ryukyus (southern

Japan) in *Proceedings of the 11th International Coral Reef Symposium*, (2008), pp.

321–325.

75.   T. L. Mizerek, *et al.*, No evidence for tropicalization of coral assemblages in a

subtropical climate change hot spot. *Coral Reefs* **40**, 1451–1461 (2021).

76.   J. A. Idjadi, R. H. Karlson, Spatial arrangement of competitors influences coexistence

of reef-building corals. *Ecology* **88**, 2449–2454 (2007).

77.   M. Brito-Millán, B. T. Werner, S. A. Sandin, D. E. McNamara, Influence of

aggregation on benthic coral reef spatio-temporal dynamics. *Royal Society Open*

Science **6**, 1–14 (2019).

78.   P. J. Edmunds, *et al.*, Evaluating the causal basis of ecological success within the

scleractinia: an integral projection model approach. *Marine Biology* **161**, 2719–2734

(2014).

79.   P. J. Edmunds, B. Riegl, Urgent need for coral demography in a world where corals
692   are disappearing. *Marine Ecology Progress Series* **635**, 233–242 (2020).

693   80.   P. J. Edmunds, Vital rates of small reef corals are associated with variation in climate.
694   *Limnology and Oceanography*, 1–13 (2020).

695   81.   T. Coulson, *et al.*, Modeling effects of environmental change on wolf population
696   dynamics, trait evolution, and life history. *Science* **334**, 1275–1278 (2011).

697   82.   J. L. Williams, T. E. X. Miller, S. P. Ellner, Avoiding unintentional eviction from
698   integral projection models. *Ecology* **93**, 2008–2014 (2012).

699   83.   J. E. N. Veron, M. G. Stafford-Smith, E. Turak, L. M. DeVantier, Corals of the World
700   (2016) (November 12, 2019).

701   84.   J. S. Madin, A. H. Baird, M. L. Baskett, S. R. Connolly, M. A. Dornelas, Partitioning
702   colony size variation into growth and partial mortality. *Biology Letters* **16**, 1–5 (2020).

703   85.   J. S. Madin, *et al.*, The Coral Trait Database, a curated database of trait information for
704   coral species from the global oceans. *Scientific Data* **3**, 1–21 (2016).

705   86.   V. R. Hall, T. P. Hughes, Reproductive strategies of modular organisms: Comparative
706   studies of reef-building corals. *Ecology* **77**, 950–963 (1996).

707   87.   L. Bramanti, M. Iannelli, T. Y. Fan, P. J. Edmunds, Using demographic models to
708   project the effects of climate change on scleractinian corals: Pocillopora damicornis as
709   a case study. *Coral Reefs* **34**, 505–515 (2015).

710   88.   I. Stott, M. Franco, D. Carslake, S. Townley, D. Hodgson, Boom or bust? A
711   comparative analysis of transient population dynamics in plants. *Journal of Ecology*
712   **98**, 302–311 (2010).

89.   D. Hodgson, J. L. McDonald, D. J. Hosken, What do you mean, "resilient"? *Trends in*

Ecology and Evolution **30**, 503–506 (2015).

90.   P. A. Zuidema, E. Jongejans, P. D. Chien, H. J. During, F. Schieving, Integral

projection models for trees: A new parameterization method and a validation of model

output. *Journal of Ecology* **98**, 345–355 (2010).

91.   O. R. Jones, *et al.*, Rcompadre and Rage - two R packages to facilitate the use of the

COMPADRE and COMADRE databases and calculation of life history traits from

matrix population models. *bioRxiv*, 2021.04.26.441330 (2021).

92.   S. Townley, D. Carslake, O. Kellie-Smith, D. Mccarthy, D. Hodgson, Predicting

transient amplification in perturbed ecological systems. *Journal of Applied Ecology* **44**

(2007).

93.   S. Townley, D. J. Hodgson, Erratum et addendum: Transient amplification and

attenuation in stage-structured population dynamics. *Journal of Applied Ecology* **45**,

1836–1839 (2008).

94.   H. O. Kreiss, Über Die Stabilitätsdefinition Für Differenzengleichungen Die Partielle

Differentialgleichungen Approximieren. *BIT* **2**, 153–181 (1962).

95.   I. Stott, D. J. Hodgson, S. Townley, Popdemo: An R package for population

demography using projection matrix analysis. *Methods in Ecology and Evolution* **3**,

797–802 (2012).

96.   L. M. Carrascal, I. Galván, O. Gordo, Partial least squares regression as an alternative

to current regression methods used in ecology. *Oikos* **118**, 681–690 (2009).

97. N. A. Rayner, *et al.*, Global analyses of sea surface temperature, sea ice, and night

marine air temperature since the late nineteenth century. *Journal of Geophysical*

Research **108**, 1–37 (2003).

98. J. V. Greenman, T. G. Benton, The frequency spectrum of structured discrete time

population models : its properties and their ecological implications. *Oikos* **110**, 369–

389 (2005).

99. D. A. Vasseur, P. Yodzis, The color of environmental noise. *Ecology* **85**, 1146–1152

(2004).

100. R Core Team, R: A language and environment for statistical computing. (2019).

101. G. Sanchez, plsdepot: Partial least squares (PLS) data analysis methods (2012).

102. P. Legendre, lmodel2: Model II regression (2018).

Tables and Figures

Figure 1. Using repeated annual surveys of tagged individual colonies, conducted between

2016 and 2019, we quantified the influence of environmental variability on the long-term

performance and transient potential of tropical and subtropical coral populations in

southern Japan and eastern Australia. **(A)** As climate shifts induce range expansions in

many coral species worldwide, their populations are increasingly exposed to a gradient in

thermal regimes, illustrated here by mean monthly sea surface temperatures ($\bar{x}_{sst}$; °C) recorded

between 1950 and 2019 (97). **(B)** To explore spatial patterns in the long-term performance and

transient (short-term) potential of coral populations exposed to varying thermal regimes, we

constructed Integral Projection Models (IPMs) describing the dynamics of tropical and

subtropical assemblages of competitive, stress-tolerant, and weedy coral taxa. To parameterise

these models, between 2016 and 2019 we documented the survival, growth, fragmentation, and

recruitment patterns of 3171 tagged individual colonies within the tropical reef communities

(▲) of Okinawa (Japan) and Heron Island (Australia), and within the subtropical coral

communities (●) of Kochi (Japan) and the Solitary Islands Marine Park (Australia).

Figure 2. A trade-off exists between long-term performance and transient potential across

our examined coral populations, corresponding with their relative exposure to thermal

variability. Partial least squares regression **s**core plot illustrating the association between

thermal conditions, and the long-term performance (λ) and transient potential (transient

envelope [*TE*] & damping ratio [ρ]) of tropical (▲) and subtropical (●) populations of

competitive (blue), stress-tolerant (yellow), and weedy (red) coral taxa. To quantify the thermal

conditions experienced by each coral population, we used sea surface temperatures (SST)

recorded between 1950 and 2019 to calculate regional estimates of mean monthly SST ($\bar{x}_{sst}$),

monthly SST variance (cv_{sst}), and monthly SST frequency spectrum (β_{sst}). Component scores

illustrate the relative degree of variance explained in the thermal predictor variables, whilst $r^2_{[y]}$

reflects the cumulative variance explained across the demographic characteristics. The shaded

polygons reflect the clustering of tropical and subtropical populations, whilst the dotted lines

delineate regions of association to facilitate the visualisation of patterns in correlation between

the abiotic and demographic variables.

Figure 3. Inter-specific variation within the trade-off observed between long-term

performance and transient potential across tropical and subtropical correlates with

patterns in population turnover rate (A) Interaction plot showcasing how estimates of

demographic stability index (DSI) vary between associated tropical (▲) and subtropical (●)

populations of competitive (blue), stress-tolerant (yellow), and weedy (red) coral taxa in

Australia and Japan. We present DSI, as an inverse measure of maximal amplification ($\bar{\rho}_{max}$),

describing the ability for populations to undergo elevated growth following disturbance. Thus,

lower DSI estimates correspond with enhanced amplification capacity. We also applied Type

2 linear regression to separately explore the association of population turnover characteristics

with **(B)** long-term performance (asymptotic population growth rate; λ), and **(C)** transient

potential (transient envelope, *TE*) across tropical and subtropical populations of competitive,

stress-tolerant, and weedy coral taxa in Australia and Japan. We note here that transient

envelope estimates were reversed during transformation to achieve normality, thus higher

values reflected diminished transient potential. We have therefore displayed transient potential

on a reversed scale to facilitate comparisons with patterns in long-term performance (λ). We

used generation time (displayed here on the log scale) as a measure of population turnover rate,

with higher estimates reflecting slower rates of population turnover. Across panels B and C r^2

values are provided as measure of model fit. Across all panels error is displayed using 95% CI.

803 **Figure 1**

813   **Figure 2**

824   **Figure 3**

830   **Table 1.** Population growth rates (λ) obtained from corresponding tropical and subtropical

831   populations of competitive, stress-tolerant, and weedy coral taxa in Australia and Japan.

832   Shading used to highlight the highest estimate of population growth across each tropical-

833   subtropical pairing. Error displayed using 95% CI.

Country	Life-history group	Tropical	Subtropical
Australia	Competitive	0.983 [0.981, 0.984]	0.958 [0.957, 0.959]
	Stress-tolerant	0.983 [0.980, 0.985]	0.899 [0.898, 0.899]
	Weedy	0.981 [0.980, 0.982]	0.686 [0.684, 0.687]
Japan	Competitive	1.001 [0.999, 1.004]	0.640 [0.639, 0.641]
	Stress-tolerant	0.913 [0.909, 0.917]	0.885 [0.877, 0.894]
	Weedy	0.760 [0.750, 0.770]	0.807 [0.802, 0.812]

835