Coral assemblages at higher latitudes favour short-term potential over long-term performance

James Cant[1, 2,*], James D. Reimer[3,4], Brigitte Sommer[5,6], Katie M. Cook[2], Sun W. Kim[7], Carrie A. Sims[8], Takuma Mezaki[9], Cliodhna O'Flaherty[2], Maxime Brooks[2], Hamish A. Malcolm[10], John M. Pandolfi[7], Roberto Salguero-Gómez[11,12,13 *,**], Maria Beger[2,12, *,**]

Running title: Transient dynamics in subtropical corals

1. Centre for Biological Diversity, St Andrews, Fife, KY16 9TH United Kingdom.
2. School of Biology, Faculty of Biological Sciences, University of Leeds, Leeds, LS2 9JT United Kingdom.
3. Molecular Invertebrate Systematics and Ecology Laboratory, Graduate School of Engineering and Science, University of the Ryukyus, Nishihara, Okinawa, Japan.
4. Tropical Biosphere Research Centre, University of the Ryukyus, Nishihara, Okinawa, Japan.
5. School of Life Sciences, University of Technology Sydney, Ultimo, NSW 2007, Australia
6. School of Life and Environmental Science, The University of Sydney, Camperdown, NSW 2006, Australia.
7. Australian Research Council Centre of Excellence for Coral Reef Studies, School of Biological Sciences, The University of Queensland, Brisbane, QLD 4072, Australia.
8. Smithsonian Tropical Research Institute, Balboa, Ancón, Panama City, Republic of Panama.
9. Kuroshio Biological Research Foundation, Nishidomari, Otsuki-cho, Kochi, Japan.
10. Fisheries Research, Department of Primary Industries, Coffs Harbour, NSW 2450, Australia.
11. Department of Zoology, University of Oxford, Oxford, 11a Mansfield Rd, OX1 3SZ United Kingdom.
12. Centre for Biodiversity and Conservation Science, School of Biological Sciences, University of Queensland, Brisbane, QLD, 4072 Australia.
13. Max Planck Institute for Demographic Research, Konrad Zuße Straße 1, 18057 Rostock, Germany.

**Share equal contribution

*Corresponding authors:

James Cant jic2@st-andrews.ac.uk, Rob Salguero-Gómez rob.salguero@zoo.ox.ac.uk, Maria Beger M.Beger@leeds.ac.uk.

Statement of Authorship: JC, RS-G and MB developed the research ideas. JC, JDR, BS, KC, SWK, CS, TM, CF, HAM, and MB collected field data, with JC, CF, MBr analysing the data. JC, RS-G, and MB led the writing of the manuscript with all authors contributing critically to the writing and giving final approval for publication.

Competing interests: All authors declare no competing interests

Abstract

The persistent exposure of coral communities to more variable abiotic regimes is assumed to

augment their resilience to future climatic variability. Yet, while the determinants of coral

population resilience across species remain unknown, we are unable to predict the winners

and losers across reef ecosystems exposed to increasingly variable conditions. Using annual

surveys of 3171 coral individuals across Australia and Japan (2017-2019), we explore spatial

variation across the short- and long-term dynamics of competitive, stress-tolerant, and weedy

assemblages to evaluate how thermal variability mediates the structural composition of coral

communities. We illustrate how, by promoting short-term potential over long-term

performance, coral assemblages can reduce their vulnerability to stochastic environments.

However, compared to stress-tolerant, and weedy assemblages, competitive coral taxa display

a reduced capacity for elevating their short-term potential. Accordingly, future climatic shifts

threaten the structural complexity of coral assemblages in variable environments, emulating

the degradation expected across global tropical reefs.

Keywords: amplification, demography, integral projection model (IPM), resilience,

subtropical, transient dynamics.

Introduction

Anticipating the resilience of natural communities requires an in-depth understanding for the

determinants underpinning their constituent populations' responses to recurrent

disturbances[1,2]. Changes in environmental regimes provoke spatial shifts in the performance

and distribution of populations, which upscale to the compositional reassembly of biological

communities[3,4]. Exposure to more variable environments is expected to indirectly augment

community resilience[5,6]. However, nuanced relationships between population characteristics

and biophysical conditions ensure inconsistent responses to climate shifts[7]; with differential

population sensitivities to habitat change having both accelerated and reversed expected

poleward range shifts in response to climate warming[8]. By linking the mechanisms driving

heterospecific variation across population responses to environmental change one can predict

the resilience of whole communities to increased climatic variability[2,9,10].

Located at the interface between tropical and temperate ecoregions, subtropical coral

communities provide an opportunity for exploring the determinants of population

resilience[11–13]. Recently, subtropical coral communities have undergone transformation with

tropical coral taxa undergoing poleward range expansions in response to shifting thermal

regimes[14–18]. At higher latitudes, however, coral communities are exposed to enhanced

seasonality and cooler temperatures, and thus experience greater abiotic variability relative to

their tropical counterparts[19]. Subsequently, subtropical coral communities offer insight into

how differing coral assemblages utilise strategies to mediate their performance in response to

environmental stochasticity across community- and regional-scales.

Exploring the performance of populations exposed to environmental stochasticity

requires a consideration of their transient (*i.e.*, short-term) dynamics[20–23]. Asymptotic (*i.e.*,

long-term) population growth rate (λ), which describes temporal changes in population size at

stationary equilibrium[24], is the predominant metric used to quantify population

performance[24,25]. However, stochastic conditions maintain natural populations within a

transient state, preventing the emergence of stationary equilibria[20,23,26]. Within stochastic

environments, recurrent disturbances impose short-term changes upon the structure of

populations that can elevate (*amplify*) or diminish (*attenuate*) their growth rates, resulting in

population performance characteristics deviating from long-term expectations[21,27].

Quantifying how transient population performance deviates from long-term expectations

(henceforth *transient potential*) is therefore crucial for predicting the success or failure of

natural populations [28], an approach that remains neglected within coral research[22].

In species rich communities, evaluating ecological dynamics requires a trait-based

approach to condense vast quantities of demographic detail[29]. Given the diversity of coral

communities, exploring patterns across the demographic characteristics of co-occurring

species presents a logistical challenge[30]. Yet, this is a challenge that can be navigated by

pooling individuals based on shared trait characteristics. Morphological, physiological and

phenological functional traits influence the fitness of individuals and thus determine the

demographic characteristics of their populations[31], their responses to disturbances[32], and

subsequently the assembly of biological communities[33–35]. Indeed, functional trait

characteristics impact upon the demographic properties of coral populations (e.g., colony

growth and reproduction[36,37]), mediating their ability to respond to local abiotic patterns[38].

Given such strong links between coral traits and demographic performance, the categorisation

of coral taxa into competitive, stress tolerant, generalist and weedy life history assemblages

(*sensu* [39]) can be used to evaluate broadscale patterns in coral community reassembly[40–42].

Trait-based assessments of coral community assembly also better inform upon the wider

implications of ongoing community shifts than taxonomic-based assessments, thereby aiding

the management of coral reef ecosystems[42].

Here, we investigate how the performance characteristics of tropical and subtropical

coral populations map onto patterns of thermal variability across assemblages of competitive,

stress-tolerant, and weedy taxa. Using Integral Projection Models (IPMs)[43], we quantify the

association between different dimensions of thermal variability (monthly mean sea surface

temperature [SST], monthly SST variance, and monthly SST frequency spectrum) and the

transient potential and long-term performance characteristics of tropical and subtropical coral

assemblages across both the northern and southern hemispheres (Fig. 1). Specifically, we

anticipate that, compared to their tropical counterparts, subtropical coral assemblages will

prioritise transient potential over long-term performance, corresponding to the need for

subtropical coral populations to endure periodically disturbed environments. Thus, we expect

that characteristics associated with amplification capacity and transient potential will align

with cooler, more variable environments, whilst measures of long-term performance will be

greater in warmer, more consistent environments; a pattern that will persist irrespective of

functional strategy and geographic location.

Results and Discussion

Our analyses reveal contrasting patterns in long-term performance and transient potential

corresponding with the exposure of coral populations to thermal variability along a gradient

from warmer, more stable environments to cooler, more variable conditions (Fig. 2). Using

partial least squares regression (PLSR), we evaluated how patterns in the long-term

performance, demographic recovery, and transient potential, of coral populations conform

with their exposure to thermal variability. We obtained estimates of long-term population

performance (asymptotic population growth rate, λ), demographic recovery (damping ratio

[ρ], *i.e.*, a relative measure of the time needed for a population to converge to a stable

equilibrium[24], and transient potential (transient envelope [*TE*], *i.e.*, the difference between

maximum and minimum population size following disturbance[27,45]) from IPMs depicting the

dynamics of tropical and subtropical assemblages of competitive, stress-tolerant, and weedy

coral taxa in Japan and Australia (Fig. 1; Supplementary S1 & S2). Meanwhile, we quantified

the exposure of these assemblages to thermal variability using three measures of local SST

regimes: monthly mean SST ($x\bar{}_{sst}$), monthly SST variance (cv_{sst}), and monthly SST

frequency spectrum (β_{sst}; Supplementary S3).

Notably, we illustrate how coral assemblages exposed to more variable thermal

conditions display enhanced transient potential. Using data focused on a single taxon, Cant et

al. [44] suggested that a capacity for short-term increases in population growth observed in a

subtropical *Acropora* spp. assemblage may underpin its viability in more variable high-

latitude environments. Here, we present evidence that this compensatory strategy is not only

isolated in competitive coral taxa and does indeed correspond with the exposure of

subtropical coral assemblages to more variable environments. Explaining 92.17% of the

variance across our three measures of thermal exposure ($x\bar{}_{sst}$, cv_{sst}, and β_{sst}), our PLSR

captures 37.43% of the variance in long-term performance (λ), demographic recovery (ρ), and

transient potential (*TE*; Fig. 2, $r^2_{[y]}$). The first PLSR component depicts a gradient in SST

variability, describing 60.97% of the variance in thermal conditions experienced by our

examined coral assemblages. It is along this component that divergent patterns within

estimates of λ and *TE* are most pronounced. Consequently, estimates of *TE* are positively

correlated with the measures of thermal variability (cv_{sst}) and frequency spectrum (β_{sst}),

whilst higher λ estimates associate with warmer mean monthly SSTs ($x\bar{}_{sst}$; Fig. 2).

Meanwhile, damping ratio (ρ) estimates are aligned with the second PLSR component that

describes patterns in mean SST ($x\bar{}_{sst}$) and SST frequency (β_{sst}). Enhanced transient potential

is thought to buffer the performance of populations exposed to elevated abiotic variability,

elevating their capacity to exploit more variable environments[46,47]. However, variation in

transient potential across our assemblages of differing coral taxa (Fig. 3), suggests that

exposure to abiotic variability alone does not assure resilience towards future climatic

variability.

Although evident across taxa, the contrasting pattern we observe between long-term

performance and transient potential does not manifest consistently between paired tropical

and subtropical coral assemblages (Fig. 3A & Table 1). We explored inter-assemblage

variation across estimates of long-term performance and transient potential. Again, we

quantified long-term performance using λ, whilst a *demographic stability index* (*DSI*)

calculated from our IPMs provided a measure of transient potential. A three-way ANOVA

revealed significant interactions between the factors of assemblage classification

(competitive, stress-tolerant, or weedy), ecoregion (tropical *vs*. subtropical), and country

(Australia vs. Japan; $ANOVA_{\lambda}$: $F_{2,11562} = 5698.47$, $p < 0.001$; $ANOVA_{DSI}$: $F_{2,11581} = 589.8$, p

< 0.001). Despite this, the tropical assemblages routinely possess higher estimates of λ

relative to their subtropical counterparts (Tukey: $p < 0.001$ in all cases; Table 1). The one

exception were weedy corals in Japan, where λ is highest in the subtropics ($\lambda_{[t]} = 0.760$ [95%

CI: 0.750, 0. 770], $\lambda_{[s]} = 0.807$ [0.802, 0.812]; $p < 0.001$). Alternatively, our subtropical coral

assemblages typically possess a greater capacity for amplifying population growth following

a disturbance than our tropical assemblages (Fig. 3A). Yet, this pattern is not consistent

across life history strategies, with competitive assemblages exhibiting the opposite trend in

Australia ($p < 0.001$) and no variation in Japan ($p = 0.999$).

The relative long-term performance and transient potential of our coral assemblages

corresponds with patterns in their generation time (Fig. 3B & C). We used Type 2 linear

regression[48] to explore the relationship between estimates of generation time (*T*, *i.e*., the time

needed for individuals of a population to be replaced[49], long-term performance (λ), and

transient potential (*TE*) calculated from our IPMs. Generation time is a strong predictor of

long-term population growth rate ($r^2 = 0.704$), with long-term performance increasing with

generation time (Fig. 3B). Conversely, longer generation times are associated with reduced

transient potential (Fig. 3C; $r^2 = 0.409$). Hence, shifts between long-term performance and

transient potential, in response to thermal variability, manifests differently across our

examined tropical and subtropical coral assemblages, due to variation in their characteristics

of temporal population turnover.

Transient buffering in variable environments

Principally, contrasting patterns between long-term performance and transient potential imply

that long-term performance does not predict the capacity for populations to endure repeated

disturbances. Also, whilst enhanced transient potential may enable natural populations to

persist within variable environments, it comes at a cost to their long-term performance.

Historically, variability in population growth rate was thought to diminish individual

fitness[50], thus hindering the persistence of populations[51]. This understanding formed the basis

of the demographic buffering hypothesis, whereby populations can minimise the influence of

environmental stochasticity on their long-term performance by limiting temporal variability

in crucial vital rates (e.g., survival, development, and reproduction[52]). Thus, variable

environments were assumed to select for populations with the ability to buffer key vital rates,

thereby reducing temporal variation in performance characteristics[50,52,53]. More recently,

however, enhanced transient potential has been presented as an adaptive mechanism that

allows populations to exploit more stochastic environments[46]. Ellis & Crone [47] demonstrated

how increased transient potential can buffer the effects of stochastic conditions on population

growth rates, an effect that was increasingly evident in populations possessing lower λ

estimates. Thus, it is not unexpected that coral assemblages established within variable

environments would possess enhanced transient potential (Fig. 2), but the energetic cost

associated with this strategy would likely inhibit their long-term performance characteristics.

Our finding that transient potential is greatest in coral assemblages displaying reduced

long-term performance contrasts with previous work on mammals and plants showing a

positive association between population growth rates and transient potential (e.g., [54,55]).

Faster population growth rates are assumed of populations characterised by faster individual

development and high fecundity[56], with these populations also expected to exhibit greater

variability in size following disturbances[55]. While each of our surveyed assemblages are in,

or close to, a state of long-term decline ($\lambda < 1$; Table 1), projected long-term performance was

highest in the tropics, where relative transient amplification was at its lowest (Fig. 3A).

Populations exhibiting longer generation times typically display reduced temporal variability

in size due to higher investment in individual survival reducing the need to counteract

disturbances[54]; a pattern that we show to be evident in our examined coral assemblages (Fig.

3C).

Interspecific variation in transient characteristics

Here we show that stress-tolerant and weedy coral taxa possess more pronounced transient

amplification at higher latitudes, highlighting a potential mechanism supporting their

persistence. Short-term increases in population growth following disturbance have been

demonstrated in a representative subtropical competitive coral assemblage (*Acropora* spp.)[44].

However, subtropical-tropical variation in the amplification capacity of competitive coral

assemblages appears minimal in comparison to the variation observed here across stress-

tolerant and weedy assemblages (Fig. 3A). Weedy corals typically exhibit smaller colony

sizes, faster growth rates, and brooding reproductive strategies, producing larvae that settle

quickly after release[39,57]. Together, these strategies support faster population turnover,

enabling weedy coral species to proliferate within highly disturbed environments[58].

Conversely, stress-tolerant corals display slower growth rates, longer life expectancies, high

fecundity, and broadcast spawning strategies[39,59]. The larger, more robust, morphologies

associated with stress-tolerant coral taxa maximise energy storage, promoting their

persistence within challenging environments[60]. Longer lifespans and elevated fecundity allow

stress-tolerant corals to endure stochastic conditions by taking advantage of sporadic

improvements in local conditions[39]. Consequently, our findings support existing projections

that weedy and stress-tolerant coral taxa are likely to become increasingly prevalent

throughout disturbed coral communities[61,62]. However, these projections herald the future loss

of the structural complexity considered essential to the functioning of reef ecosystems[63].

Crucially, our findings do not reflect the current reality for many coral assemblages

within regions of high abiotic variability, suggesting that the composition of coral

communities is not solely mediated by the interplay between transient dynamics and abiotic

variability. Despite the lower amplificatory capacity of subtropical competitive corals

compared to subtropical weedy and stress-tolerant assemblages, competitive coral taxa

dominate many subtropical coral assemblages[64–66]. Utilising fast growth strategies, colonies

of competitive coral taxa are capable of rapidly colonising available substrate, quickly

outcompeting heterospecifics for both space and light[39]. Whilst this competitive nature

explains their dominance across contemporary subtropical communities, the sensitivity of

many competitive coral taxa to environmental shifts means that these assemblages are often

regarded as early successional, dominating only within optimal environments, and receding

as reef ecosystems approach climax states[67,68]. Within subtropical environments coral

community composition is mediated by environmental pressures and dispersal barriers that

filter the occurrence of species according to their trait characteristics[38,69]. As a result,

subtropical coral assemblages typically consist of a subset of tropical species found on

tropical coral reefs[38], as well as subtropical specialists and endemics. The dominance of

competitive coral taxa within subtropical coral assemblages, despite their reduced transient

performance relative to other coral taxa, may therefore, imply that competitive interactions

profoundly influence the performance of coral populations[70,71]. Certainly, further

investigation into the influence of competitive interactions upon the transient dynamics of

coral populations is needed to disentangle how coexistence between coral populations

facilitates their persistence within variable environments.

Conclusions

A limited understanding for the abiotic determinants driving the dynamics of coral

assemblages inhibits our capacity to predict their future performance and, therefore, manage

global coral community reassembly[72–74]. Here, we demonstrate how coral assemblages within

regions of high environmental stochasticity exhibit demographic strategies associated with

enhanced transient potential, but at a cost to their long-term performance. Climatic change is

exposing coral communities worldwide to increased abiotic variability. Crucially, our

findings here emphasize that whilst coral assemblages can adopt demographic strategies

enhancing their viability when exposed to abiotic variability, the winners and losers within

future, more variable environments cannot be predicted from existing measures of long-term

performance. Equally, the relationship that we observed between transient potential and

thermal variability was not universal across coral taxa, nor did it manifest identically across

hemispheres. Subtle patterns in the association between population dynamics and their

climate drivers hinder predictions of the consequences of environmental change within

biological communities[75]. Nevertheless, relative to competitive coral taxa, weedy and stress-

tolerant corals appear to possess a greater capacity for enduring within environments

characterised by repeated abiotic disturbances. Yet, competitive coral taxa are often

associated with more complex morphologies and therefore support the structural complexity

critical to the wider functioning of coral associated ecosystems[63]. Accordingly, future

increases in abiotic variability threaten the viability of coral associated ecosystems.

Methods

Modelling population dynamics

Integral Projection Models (IPMs) capture how the state composition of individuals

influences the performance of populations over discrete time periods (*t* to *t+1*)[43]. Here, to

quantify the long-term performance characteristics and transient (*i.e.*, short-term) potential of

coral populations, we used IPMs describing patterns in colony survival (σ), transitions in size

(growth and shrinkage, γ), fragmentation probability (κ), fecundity (φ), and recruitment ($\square$),

each as a function of colony size (*z*; visible horizontal surface area, cm^2). Specifically, our

IPMs took the form

286
$$n[z', t + 1] = \int_L^U (P_{z'z} + F_{z'z}\, n[z, t]\, \delta z) \tag{1}$$

287
$$P_{z'z} = (1 - \kappa_z)\, \sigma_z\, \gamma_{z'z} + (\kappa_z\, \kappa b_z\, \kappa_z^0) \tag{2}$$

288
$$F_{z'z} = \varphi_z\, \phi C_0 \tag{3}$$

with [L, U] representing the range of possible colony sizes; calculated as 10% above and

below observed maximum and minimum colony sizes to avoid accidental exclusion[76].

Accordingly, the structure of a population at time *t+1* (*n[z', t+1]*) is a product of its structure

at time *t* (*n[z', t]*) subject to the survival (σ_z) and transition of individual colonies from size *z*

to size *z'* ($\gamma_{z'z}$); the probability of colony fragmentation (κ_z) and the number (κb_z) and size

distribution of any colony remnants produced (κ_z^0); and colony fecundity (φ_z) combined with

the probability of successful recruitment ($\square$) and the size distribution of surviving recruits

(C$_0$).

Data collection

We parameterised our IPMs using data collected during repeated annual surveys of 3171

tagged colonies within tropical and subtropical coral communities in southern Japan and

eastern Australia, conducted between 2016 and 2019 (Fig. 1; Supplementary S1). We tagged

individual colonies using permanent plots arranged haphazardly throughout four focal coral

communities (Australian subtropics [AS], Australian tropics [AT], Japanese subtropics [JS],

Japanese tropics [JT]) and demarcated with numbered tags[44,62]. All tagged colonies were

identified *in situ*, or from photographs, to the lowest possible taxonomic level (either genus or

species). No samples were taken from tagged colonies, as although this would have allowed

us to resolve species identity, we wanted to avoid any lasting interference with the processes

of colony survival, growth, and fragmentation.

To facilitate comparing population characteristics observed across spatially distinct

regions in Australia and Japan with varying degrees of species overlap[77], we grouped tagged

colonies across each region according to shared life-history-strategies (*sensu* [39–41]).

Specifically, we categorised colonies as 'competitive', 'weedy', 'stress-tolerant' or

'generalist' based on their morphology, growth rate and reproductive mode, following the

genera classifications of Darling et al. [39], with minor adaptions made based on local expertise

(see supplementary S2 for a detailed list). In the event that genera represented species

classified across multiple categories (19 cases), we randomly assigned individuals across the

relevant categories in proportion with the number of species within each category known to

occur in the area (*sensu* [41]). Following the pooling of colonies according to their life-history-

strategies, we omitted all individuals defined as generalists from subsequent analyses due to

their limited representation across our regional samples (n: AS = 22 colonies; AT = 31; JS =

17; JT = 65). Consequently, we constructed IPMs concerning the dynamics of each functional

coral assemblage (competitive, stress-tolerant, and weedy; Fig 1) at each of the four

geographical locations.

Photographs capturing the visible horizontal extent of tagged colonies were used to

follow individuals over successive surveys and obtain longitudinal records of colony surface

area (cm^2; transformed to a $\log_{10}$ scale) over time. Using generalised linear mixed models

(GLMMs), we estimated size-specific patterns in colony survival (σ), transitions in size (γ),

and fragmentation probability (κ) for each population (Supplementary S1). In each case, our

GLMMs included random effects (colony identity and survey location) to account for any

autocorrelation between observations and within-subject variability associated with our

pooling of data recorded from individuals followed across multiple years, and at different

sites. Colony survival (σ) reflected the continued presence of tagged individuals across

survey intervals (*t* to *t+1*) and was modelled as a logistic function of colony surface area at

time *t*. Colony size transitions (γ), representing both growth through colony extension, and

shrinkage through partial mortality[78], were modelled using the polynomial relationship

between initial colony surface area at time *t* and subsequent surface area at time *t+1*. Colony

fragmentation probability (κ) was then modelled as a polynomial logistic function of colony

size at time *t*. During our surveys, we recorded fragmentation in the event of observed colony

breakage, recording the size (surface area, cm^2) of all remnants produced in each case.

Subsequently, we also modelled the number (κb_z) and size (κ_Z^0) of remnant colonies produced

during fragmentation as a function of colony size at time *t*, using Poisson and polynomial

GLMMs, respectively.

Alongside our surveys of tagged individual colonies, we also monitored colony

recruitment within our permanent coral plots. During each annual survey, we recorded the

number and size of new colonies appearing within each plot. These recruitment counts

enabled us to quantify annual and regional variability in recruit densities (Table S2), as well

as estimate population-specific recruit size distributions (C_0; Supplementary S1). Prior to

incorporating recruitment dynamics into our IPMs, however, we first determined patterns in

colony fecundity (φ). Using data relating colony size and larval output (larval density, cm^3)

extracted from the Coral Trait Database[79,80], we calculated colony fecundity (φ) as the

polynomial relationship between colony size at *t* and expected larval output (Supplementary

S1). We emphasise here that this approach enabled us to complete the life cycle loop, linking

the dynamics of existing individuals with the introduction of new, genetically distinct

individuals within our IPMs; a necessary step when evaluating population performance[24].

However, to ensure our modelled recruitment dynamics, and therefore, our modelling

framework, reflected empirical observations, we parameterised a recruit settlement function

(□) into our IPMs. As a probability-based function, this recruit settlement function converts

modelled larval outputs into proportional recruit densities corresponding with our empirical

recruitment counts. We determined this recruit settlement function by dividing total expected

larval output in any given year by the corresponding annual recruitment count

(Supplementary S1, *sensu* [62,81]).

Quantifying population characteristics

From our IPMs, we obtained estimates of long-term performance (asymptotic population

growth, λ), generation time (*T*), and transient potential (damping ratio [ρ], maximal

amplification [$\bar{\rho}_{max}$] & transient envelope [*TE*]) for each tropical and subtropical coral

assemblage[24,27,45,49,82]. Estimates of λ are typically used as a measure of long-term population

viability[25], and reflect whether a population is expected to grow ($\lambda > 1$) or decline ($\lambda < 1$)

when at stationary equilibrium[24]. Generation time is a measure of population turnover,

describing the time needed for individuals of a population to be replaced[49]. Alternatively, the

measures of transient potential describe the expected characteristics of populations following

their displacement from stationary equilibrium due to disturbances. The damping ratio

constitutes a measure of demographic recovery[45,83], describing the rate at which a population

perturbed from its stationary equilibrium converges back to its asymptotic growth

trajectory[24]. Meanwhile, maximal amplification quantifies the greatest increase in population

size following a disturbance, relative to its asymptotic growth trajectory[27,82]. Finally, the

transient envelope quantifies the magnitude by which the transient dynamics of a population

deviates from its long-term trajectory[45].

To calculate the aforementioned demographic characteristics, we discretised our IPMs

into large matrices. Applying the 'midpoint rule' we integrated each IPM into a high-

dimension matrix (200×200 cells), with the probability of transitioning from one cell to the

next approximated at the cell midpoint and multiplied by the cell width as per Zuidema et al.

[84]. Estimates of λ were then identified as the dominant eigenvalue of each discretised matrix,

whilst we estimated damping ratios as the ratio between the subdominant and dominant

eigenvalues. With the *R* package *Rage*[85] we then calculated generation time using estimates

of net reproductive rate (R_0) and λ obtained from each matrix,

385
$$T = \frac{\log (R_0)}{\log (\lambda)}. \tag{4}$$

386   Next, we determined the transient envelope of each assemblage using their associated Kreiss

387   bounds of amplification ($\overline{K}_\lambda^*$) and attenuation ($\underline{K}_\lambda^*$),

388
$$TE = \overline{K}_\lambda^* - \underline{K}_\lambda^*. \tag{5}$$

Respectively, the Kreiss bounds of amplification and attenuation reflect the largest and

smallest expected long-term densities of a population following the dissipation of transient

conditions, relative to its asymptotic growth trajectory[86–88]. We acknowledge here that this

definition is more commonly applied to measures of population inertia[27], which are more

typically used in estimating transient envelopes[45]. However, Kreiss bound estimates have

been demonstrated to align with corresponding estimates of population inertia and, unlike

estimates of population inertia, are not sensitive to imprimitive population models (*i.e.*, non-

negative models permitting transitions between all state classes, but with transitions between

certain stages occurring only at periodic intervals[24,27]), hence their selection here. We derived

these Kreiss bounds, alongside estimates of maximal amplification, using their corresponding

functions in the *R* package *popdemo*[89].

Across each demographic measure, we determined the variance in our assemblage-

specific estimates through Jack-knife resampling. During resampling, we generated 1,000

IPM variants for each assemblage, each time using 95% of our original data sample without

replacement, whilst permitting recruit survival probabilities ($\square$) to vary within observed

limits. Finally, prior to their inclusion in further analyses, the jack-knifed distributions of λ,

generation time, transient envelope, and maximal amplification each required transforming to

ensure approximate normality. We omitted 26 variants for which $\lambda > 2$, as these presented

unrealistic illustrations of population performance (*i.e.*, more than doubling population size

every year), before applying a log transformation to the generation time variable and a power

transformation (y^x) across the damping ratio ($y^{-2.0}$), transient envelope ($y^{-0.1}$) and maximal

amplification variables ($y^{-0.5}$).

Evaluating spatial trends in population characteristics

To test for patterns in the spatial variation of long-term performance and transient potential

across tropical and subtropical coral assemblages, we utilised partial least squares regression

(PLSR), ANOVA, and Type 2 linear regression. Initially, we applied PLSR to test whether

contrasting patterns in the long-term performance characteristics and transient potential of

coral assemblages align with their exposure to abiotic variability. A PLSR regression

quantifies the association between multiple predictor variables and one or more dependant

variables[90]. Subsequently, using this technique we simultaneously evaluated the relationships

between mean estimates of λ, damping ratio, and transient envelope obtained for each

assemblage, and their correlation with patterns in thermal conditions, to provide an insight

into the demographic trade-offs of coral assemblages and their mechanistic drivers.

To evaluate how abiotic variability mediates the selection of short- and long-term

performance characteristics in coral assemblages, within our PLSR analyses we quantified

the abiotic conditions experienced by each coral assemblage using three measures of local sea

surface temperature (SST) regimes: mean monthly SST ($\bar{x}_{sst}$), monthly SST variance (cv_{sst}),

and monthly SST frequency spectrum (β_{sst}; Supplementary S3). Focusing on the four

geographical regions in which our focal coral assemblages were surveyed (GPS: AS = -30.3°,

153.1°; AT = -23.4°, 151.9°; JS = 32.8°, 132.6°; JT = 26.5°, 128.1°; Fig. 1), we extracted

monthly SST readings (°C; overlaid on a 1° latitude-longitude grid) taken between January

1950 and December 2019, inclusive, from the HadISST dataset[91]. Arranging these SST

records into 69-year timeseries for each location, we then calculated the mean ($\bar{x}_{sst}$) and

coefficient of variance (cv_{sst}) for each timeseries. Next, we estimated the frequency spectrum

of each time series. Spectral analysis is used to quantify the periodicity of recurrent

variability within a timeseries, with higher frequencies associated with shorter-term

fluctuations[92]. The frequency spectrum of a time series is represented by its spectral exponent

(β) and equal to the negative slope between its log spectral density and log frequency[93],

which we calculated using the package *stats*[94]. After testing these abiotic predictor variables

for collinearity (Supplementary S3), we performed our PLSR analyses using the *R* package

plsdepot[95].

Finally, we assessed how patterns in the long-term performance, and capacity for

coral assemblages to benefit from recurrent disturbance vary between tropical and subtropical

regions, and how this variation manifests across coral taxa. Using a three-way ANOVA, we

separately investigated variation in estimates of λ and maximal amplification across the three

factors of country (Australia *vs*. Japan), ecoregion (tropical *vs*. subtropical), and assemblage

classification (competitive, stress-tolerant, or weedy). With maximal amplification estimates

inverted during transformation, larger values subsequently reflect reduced amplification

potential. For the purposes of clarity in this analysis, we refer to this reversed scale as a

demographic stability index (*DSI*), with lower values corresponding with enhanced

amplification. We evaluated drivers of long- and short-term performance, by using Type 2

linear regression to separately evaluate the relationship between generation time (*T*) and

estimates of λ and transient envelope (*TE*). Type 2 linear regression is an approach for

quantifying the relationship between two non-independent variables, such that both variables

include an element of error[48]. Here, due to differences in the magnitude of variance (σ^2)

across our variables of generation time, λ, and transient envelope (σ^2: *T* = 1.139; λ = 0.009;

TE = 0.016) we performed a Ranged Major Axis Type 2 regression using the *R* package

lmodel2[96].

Acknowledgements

The authors would like to thank S. Dalton, L. Lachs, C. Kim, R. Edgar, K.-L. Gomez-

Cabrera, N. Kyriacou, I. Mizukami, H. Kise, C. Fourreau, G. Masucci, P. Biondi, S.

Nishihira, M. Tamae, H. Nakakoji, C. Tan, L. Lawrence, T. Hofmann, I. Montero-Serra, and

all staff at Dive Quest, Marine Space, Pacific Marine, and SeaAir, for their assistance with

field data collection. Additionally, aspects of this research would not have been possible

without the research permits provided by the Solitary Islands Marine Park branch of the NSW

Department of Primary Industries (SIMP 2016/002V2, MEAA 20/45) and the Great Barrier

Reef Marine Park Authority (G19/42221.1). Funding for this research was provided by a

Natural Environment Research Council (NERC) Doctoral Training Programme Scholarship

to JC, a Royal Geographical Society Ralph Brown Expedition Award (RBEA 03/19) to MB

and JC, the Australian Research Council Centre of Excellence for Coral Reef Studies

(CE140100020) to JMP and others, the Australian Research Council Centre of Excellence for

Environmental Decisions (CE110001014), a British Ecological Society small grant, the

Winifred Violet Scott Charitable Trust, and the European Union's Horizon 2020 research and

innovation programme under the Marie Skłodowska-Curie grant agreement TRIM-DLV-

747102 to MB. BS was supported by a Chancellor's Postdoctoral Research Fellowship from

the University of Technology Sydney and a University of Sydney Fellowship.

References

1.　Vázquez, D.P., Gianoli, E., Morris, W.F., and Bozinovic, F. (2017). Ecological and

evolutionary impacts of changing climatic variability. Biological Reviews *92*, 22–42.

10.1111/brv.12216.

2.　Williams, S.E., Shoo, L.P., Isaac, J.L., Hoffmann, A.A., and Langham, G. (2008).

Towards an integrated framework for assessing the vulnerability of species to climate

change. PLoS Biol *6*. 10.1371/journal.pbio.0060325.

3.　Pecl, G., Araujo, M.B., Bell, J.D., Blanchard, J., and Bonebrake, T.C. (2017).

Biodiversity redistribution under climate change: Impacts on ecosystems and human

well-being Publication Details. Science (1979) *355*, 1–9.

4.　Totland, Ø., and Nyléhn, J. (1998). Assessment of the effects of environmental change

on the performance and density of Bistorta vivipara: The use of multivariate analysis

and experimental manipulation. Journal of Ecology *86*, 989–998. 10.1046/j.1365-

2745.1998.00318.x.

5.　Rivest, E.B., Comeau, S., and Cornwall, C.E. (2017). The role of natural variability in

shaping the response of coral reef organisms to climate change. Curr Clim Change Rep

3, 271–281. 10.1007/s40641-017-0082-x.

6.　Boyd, P.W., Cornwall, C.E., Davison, A., Doney, S.C., Fourquez, M., Hurd, C.L.,

Lima, I.D., and McMinn, A. (2016). Biological responses to environmental

heterogeneity under future ocean conditions. Glob Chang Biol *22*, 2633–2650.

10.1111/gcb.13287.

7.   Parmesan, C., and Yohe, G. (2003). A globally coherent fingerprint of climate change.

Nature *421*, 37–42.

8.   Chen, I.C., Hill, J.K., Ohlemüller, R., Roy, D.B., and Thomas, C.D. (2011). Rapid

range shifts of species associated with high levels of climate warming. Science (1979)

333, 1024–1026. 10.1126/science.1206432.

9.   Foden, W.B., Butchart, S.H.M., Stuart, S.N., Vié, J.C., Akçakaya, H.R., Angulo, A.,

DeVantier, L.M., Gutsche, A., Turak, E., Cao, L., et al. (2013). Identifying the world's

most climate change vulnerable species: A systematic trait-based assessment of all

birds, amphibians and corals. PLoS One *8*. 10.1371/journal.pone.0065427.

10.   Dawson, T.P., Jackson, S.T., House, J.I., Prentice, I.C., and Mace, G.M. (2011).

Beyond predictions: Biodiversity conservation in a changing climate. Science (1979)

332, 53–58.

11.   Camp, E.F., Schoepf, V., Mumby, P.J., Hardtke, L.A., Rodolfo-Metalpa, R., Smith,

D.J., and Suggett, D.J. (2018). The future of coral reefs subject to rapid climate

change: Lessons from natural extreme environments. Front Mar Sci *5*, 1–21.

10.3389/fmars.2018.00004.

12.   Burt, J.A., Camp, E.F., Enochs, I.C., Johansen, J.L., Morgan, K.M., Riegl, B., and

Hoey, A.S. (2020). Insights from extreme coral reefs in a changing world. Coral Reefs

39, 495–507. 10.1007/s00338-020-01966-y.

13.   Beger, M., Sommer, B., Harrison, P.L., Smith, S.D.A., and Pandolfi, J.M. (2014).

516   Conserving potential coral reef refuges at high latitudes. Divers Distrib *20*, 1–13.

517   10.1111/ddi.12140.

518   14.   Precht, W.F., and Aronson, R.B. (2004). Climate flickers and range shifts of reef

519   corals. Front Ecol Environ *2*, 307–314.

520   15.   Tuckett, C.A., de Bettignies, T., Fromont, J., and Wernberg, T. (2017). Expansion of

521   corals on temperate reefs: Direct and indirect effects of marine heatwaves. Coral Reefs

522   *36*, 947–956. 10.1007/s00338-017-1586-5.

523   16.   Yamano, H., Sugihara, K., and Nomura, K. (2011). Rapid poleward range expansion

524   of tropical reef corals in response to rising sea surface temperatures. Geophys Res Lett

525   *38*, 1–6. 10.1029/2010GL046474.

526   17.   Baird, A.H., Sommer, B., and Madin, J.S. (2012). Pole-ward range expansion of

527   Acropora spp. along the east coast of Australia. Coral Reefs *31*, 1063.

528   10.1007/s00338-012-0928-6.

529   18.   Booth, D.J., and Sears, J. (2018). Coral expansion in Sydney and associated coral-reef

530   fishes. Coral Reefs *37*, 995. DOI 10.1007/s00338-018-1727-5.

531   19.   Sommer, B., Beger, M., Harrison, P.L., Babcock, R.C., and Pandolfi, J.M. (2018).

532   Differential response to abiotic stress controls species distributions at biogeographic

533   transition zones. Ecography *41*, 478–490. 10.1111/ecog.02986.

534   20.   Hastings, A. (2004). Transients: The key to long-term ecological understanding?

535   Trends Ecol Evol *19*, 39–45. 10.1016/j.tree.2003.09.007.

536   21.   Ezard, T.H.G., Bullock, J.M., Dalgleish, H.J., Millon, A., Pelletier, F., Ozgul, A., and

537   Koons, D.N. (2010). Matrix models for a changeable world: The importance of

538 transient dynamics in population management. Journal of Applied Ecology *47*, 515–

539 523. 10.1111/j.1365-2664.2010.01801.x.

540 22. Cant, J., Salguero-Gómez, R., and Beger, M. (2021). Transient demographic

541 approaches can drastically expand the toolbox of coral reef science. bioRxiv *41*,

542 2021.04.08.438926. 10.1101/2021.04.08.438926.

543 23. Hastings, A., Abbott, K.C., Cuddington, K., Francis, T., Gellner, G., Lai, Y.C.,

544 Morozov, A., Petrovskii, S., Scranton, K., and Zeeman, M.L. (2018). Transient

545 phenomena in ecology. Science (1979) *361*, 1–9. 10.1126/science.aat6412.

546 24. Caswell, H. (2001). Matrix population models: Construction, analysis and

547 interpretation 2nd Edition. (Oxford University Press Inc).

548 25. Crone, E.E., Menges, E.S., Ellis, M.M., Bell, T., Bierzychudek, P., Ehrlén, J., Kaye,

549 T.N., Knight, T.M., Lesica, P., Morris, W.F., et al. (2011). How do plant ecologists use

550 matrix population models? Ecol Lett *14*, 1–8. 10.1111/j.1461-0248.2010.01540.x.

551 26. Hastings, A. (2001). Transient dynamics and persistence of ecological systems. Ecol

552 Lett *4*, 215–220.

553 27. Stott, I., Townley, S., and Hodgson, D. (2011). A framework for studying transient

554 dynamics of population projection matrix models. Ecol Lett *14*, 959–970.

555 10.1111/j.1461-0248.2011.01659.x.

556 28. Koons, D.N., Grand, J.B., Zinner, B., and Rockwell, R.F. (2005). Transient population

557 dynamics: Relations to life history and initial population state. Ecol Modell *185*, 283–

558 297. 10.1016/j.ecolmodel.2004.12.011.

559   29.   Chalmandrier, L., Hartig, F., Laughlin, D.C., Lischke, H., Pichler, M., Stouffer, D.B.,

560   and Pellissier, L. (2021). Linking functional traits and demography to model species-

561   rich communities. Nat Commun *12*, 1–9. 10.1038/s41467-021-22630-1.

562   30.   Madin, J.S., Hoogenboom, M.O., Connolly, S.R., Darling, E.S., Falster, D.S., Huang,

563   D., Keith, S.A., Mizerek, T., Pandolfi, J.M., Putnam, H.M., et al. (2016). A trait-based

564   approach to advance coral reef science. Trends Ecol Evol *31*, 419–428.

565   10.1016/j.tree.2016.02.012.

566   31.   Violle, C., Navas, M., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., and Garnier, E.

567   (2007). Let the concept of trait be functional! Oikos *116*, 882–892.

568   10.1111/j.2007.0030-1299.15559.x.

569   32.   Grime, J.P., and Pierce, S. (2012). The evolutionary strategies that shape ecosystems

570   (John Wiley and Sons Ltd.).

571   33.   McGill, B.J., Enquist, B.J., Weiher, E., and Westoby, M. (2006). Rebuilding

572   community ecology from functional traits. Trends Ecol Evol *21*, 178–185.

573   10.1016/j.tree.2006.02.002.

574   34.   Cadotte, M.W., Carscadden, K., and Mirotchnick, N. (2011). Beyond species:

575   Functional diversity and the maintenance of ecological processes and services. Journal

576   of Applied Ecology *48*, 1079–1087. 10.1111/j.1365-2664.2011.02048.x.

577   35.   Falster, D.S., Brännström, Å., Westoby, M., and Dieckmann, U. (2017). Multitrait

578   successional forest dynamics enable diverse competitive coexistence. Proc Natl Acad

579   Sci U S A *114*, E2719–E2728. 10.1073/pnas.1610206114.

36.   Madin, J.S., Hoogenboom, M.O., and Connolly, S.R. (2012). Integrating physiological

and biomechanical drivers of population growth over environmental gradients on coral

reefs. Journal of Experimental Biology *215*, 968–976. 10.1242/jeb.061002.

37.   Álvarez-Noriega, M., Baird, A.H., Dornelas, M., Madin, J.S., Cumbo, V.R., and

Connolly, S.R. (2016). Fecundity and the demographic strategies of coral

morphologies. Ecological Society of America *97*, 3485–3493.

38.   Sommer, B., Harrison, P.L., Beger, M., and Pandolfi, J.M. (2014). Trait-mediated

environmental filtering drives assembly at biogeographic transition zones. Ecology *95*,

1000–1009. 10.1890/13-1445.1.

39.   Darling, E.S., Alvarez-Filip, L., Oliver, T.A., Mcclanahan, T.R., and Côté, I.M.

(2012). Evaluating life-history strategies of reef corals from species traits. Ecol Lett

15, 1378–1386. 10.1111/j.1461-0248.2012.01861.x.

40.   Darling, E.S., McClanahan, T.R., and Côté, I.M. (2013). Life histories predict coral

community disassembly under multiple stressors. Glob Chang Biol *19*, 1930–1940.

10.1111/gcb.12191.

41.   Zinke, J., Gilmour, J.P., Fisher, R., Puotinen, M., Maina, J., Darling, E., Stat, M.,

Richards, Z.T., McClanahan, T.R., Beger, M., et al. (2018). Gradients of disturbance

and environmental conditions shape coral community structure for south-eastern

Indian Ocean reefs. Divers Distrib *24*, 605–620. 10.1111/ddi.12714.

42.   Darling, E.S., McClanahan, T.R., Maina, J., Gurney, G.G., Graham, N.A.J.,

Januchowski-Hartley, F., Cinner, J.E., Mora, C., Hicks, C.C., Maire, E., et al. (2019).

Social–environmental drivers inform strategic management of coral reefs in the

Anthropocene. Nat Ecol Evol *3*, 1341–1350. 10.1038/s41559-019-0953-8.

43. Easterling, M.R., Ellner, S.P., and Dixon, P.M. (2000). Size-specific sensitivity:

applying a new structured population model. Ecology *81*, 694–708.

44. Cant, J., Cook, K., Reimer, J.D., Mezaki, T., Nakamura, M., O'Flaherty, C., Salguero-

Gómez, R., and Beger, M. (2022). Transient amplification enhances the persistence of

tropicalising coral populations in marginal high latitude environments. Ecography.

10.1101/2021.04.23.441133.

45. Capdevila, P., Stott, I., Beger, M., and Salguero-Gómez, R. (2020). Towards a

comparative framework of demographic resilience. Trends Ecol Evol *35*, 776–786.

10.1016/j.tree.2020.05.001.

46. McDonald, J.L., Stott, I., Townley, S., and Hodgson, D.J. (2016). Transients drive the

demographic dynamics of plant populations in variable environments. Journal of

Ecology *104*, 306–314. 10.1111/1365-2745.12528.

47. Ellis, M.M., and Crone, E.E. (2013). The role of transient dynamics in stochastic

population growth for nine perennial plants. Ecology *94*, 1681–1686.

48. Sokal, R.R., and Rohlf, F.J. (1995). Biometry: The principles and practice of statistics

in biological research 3rd Editio. (W. H. Freeman & Co).

49. Gaillard, J., Yoccoz, N.G., Lebreton, J., Bonenfant, C., Devillard, S., Loison, A.,

Pontier, D., and Allaine, D. (2005). Generation Time: A reliable metric to measure

life-history variation among mammalian populations. Am Nat *166*, 119–123.

50. Pfister, C.A. (1998). Patterns of variance in stage-structured populations: Evolutionary

predictions and ecological implications. Proceedings of the National Academy of

Sciences *95*, 213–218.

51.   Lande, R. (1993). Risks of population extinction from demographic and environmental

stochasticity and random catastrophes. American Naturalist *142*, 911–927.

10.1086/285580.

52.   Morris, W.F., and Doak, D.F. (2004). Buffering of life histories against environmental

stochasticity: Accounting for a spurious correlation between the variabilities of vital

rates and their contributions to fitness. American Naturalist *163*, 579–590.

10.1086/382550.

53.   Hilde, C.H., Gamelon, M., Sæther, B.E., Gaillard, J.M., Yoccoz, N.G., and Pélabon, C.

(2020). The demographic buffering hypothesis: Evidence and challenges. Trends Ecol

Evol *35*, 523–538. 10.1016/j.tree.2020.02.004.

54.   Morris, W.F., Pfister, C.A., Tuljapurkar, S., Haridas, C. V., Boggs, C.L., Boyce, M.S.,

Bruna, E.M., Church, D.R., Coulson, T., Doak, D.F., et al. (2008). Longevity can

buffer plant and animal populations against changing climatic variability. Ecology *89*,

19–25.

55.   Gamelon, M., Gimenez, O., Baubet, E., Coulson, T., Tuljapurkar, S., and Gaillard,

J.M. (2014). Influence of life-history tactics on transient dynamics: A comparative

analysis across mammalian populations. American Naturalist *184*, 673–683.

10.1086/677929.

56.   Oli, M.K. (2004). The fast-slow continuum and mammalian life-history patterns: An

empirical evaluation. Basic Appl Ecol *5*, 449–463. 10.1016/j.baae.2004.06.002.

57.   Knowlton, N. (2001). The future of coral reefs. Proceedings of the National Academy

of Sciences *98*, 5419–5425.

58.   Adjeroud, M., Kayal, M., Iborra-Cantonnet, C., Vercelloni, J., Bosserelle, P., Liao, V.,

648   Chancerelle, Y., Claudet, J., and Penin, L. (2018). Recovery of coral assemblages

649   despite acute and recurrent disturbances on a South Central Pacific reef. Sci Rep *8*.

650   10.1038/s41598-018-27891-3.

651   59.   Klepac, C.N., and Barshis, D.J. (2020). Reduced thermal tolerance of massive coral

652   species in a highly variable environment: Reduced heat tolerance of massive corals.

653   Proceedings of the Royal Society B: Biological Sciences *287*, 19–21.

654   10.1098/rspb.2020.1379rspb20201379.

655   60.   van Woesik, R., Irikawa, A., Anzai, R., and Nakamura, T. (2012). Effects of coral

656   colony morphologies on mass transfer and susceptibility to thermal stress. Coral Reefs

657   *31*, 633–639. 10.1007/s00338-012-0911-2.

658   61.   Loya, Y., Sakai, K., Yamazato, K., Nakano, Y., Sambali, H., and van Woesik, R.

659   (2001). Coral bleaching: the winners and the losers. Ecol Lett *4*, 122–131.

660   62.   Cant, J., Salguero☐Gómez, R., Kim, S.W., Sims, C.A., Sommer, B., Brooks, M.,

661   Malcolm, H.A., Pandolfi, J.M., and Beger, M. (2021). The projected degradation of

662   subtropical coral assemblages by recurrent thermal stress. Journal of Animal Ecology

663   *90*, 233–247. 10.1111/1365-2656.13340.

664   63.   Graham, N.A.J., and Nash, K.L. (2013). The importance of structural complexity in

665   coral reef ecosystems. Coral Reefs *32*, 315–326. 10.1007/s00338-012-0984-y.

666   64.   Harriott, V.J., Harrison, P.L., and Banks, S.A. (1995). The coral communities of Lord

667   Howe Island. Mar Freshw Res *46*, 457–465.

65.   Nozawa, Y., Tokeshi, M., and Nojima, S. (2008). Structure and dynamics of a high-

latitude scleractinian coral community in Amakusa, southwestern Japan. Mar Ecol

Prog Ser *358*, 151–160. 10.3354/meps07342.

66.   Sugihara, K., Sonoda, N., Imafuku, T., Nagata, S., Ibusuki, T., and Yamano, H.

(2009). Latitudinal changes in hermatypic coral communities from west Kyushu to Oki

Islands in Japan. Journal of the Japanese Coral Reef Society *11*, 51–67.

67.   Wilson, S.K., Robinson, J.P.W., Chong-Seng, K., Robinson, J., and Graham, N.A.J.

(2019). Boom and bust of keystone structure on coral reefs. Coral Reefs *38*, 625–635.

10.1007/s00338-019-01818-4.

68.   Ohba, H., Hashimoto, K., Shimoike, K., Shibuno, T., and Fujioka, Y. (2008).

Secondary succession of coral reef communities at Urasoko Bay, Ishigaki Island, the

Ryukyus (southern Japan). In Proceedings of the 11th International Coral Reef

Symposium, pp. 321–325.

69.   Mizerek, T.L., Madin, J.S., Benzoni, F., Huang, D., Luiz, O.J., Mera, H., Schmidt-

Roach, S., Smith, S.D.A., Sommer, B., and Baird, A.H. (2021). No evidence for

tropicalization of coral assemblages in a subtropical climate change hot spot. Coral

Reefs *40*, 1451–1461. 10.1007/s00338-021-02167-x.

70.   Idjadi, J.A., and Karlson, R.H. (2007). Spatial arrangement of competitors influences

coexistence of reef-building corals. Ecology *88*, 2449–2454. 10.1890/06-2031.1.

71.   Brito-Millán, M., Werner, B.T., Sandin, S.A., and McNamara, D.E. (2019). Influence

of aggregation on benthic coral reef spatio-temporal dynamics. R Soc Open Sci *6*, 1–

14. 10.1098/rsos.181703.

72.   Edmunds, P.J., Burgess, S.C., Putnam, H.M., Baskett, M.L., Bramanti, L., Fabina,

691   N.S., Han, X., Lesser, M.P., Madin, J.S., Wall, C.B., et al. (2014). Evaluating the

692   causal basis of ecological success within the scleractinia: an integral projection model

693   approach. Mar Biol *161*, 2719–2734. 10.1007/s00227-014-2547-y.

694   73.   Edmunds, P.J., and Riegl, B. (2020). Urgent need for coral demography in a world

695   where corals are disappearing. Mar Ecol Prog Ser *635*, 233–242. 10.3354/meps13205.

696   74.   Edmunds, P.J. (2020). Vital rates of small reef corals are associated with variation in

697   climate. Limnol Oceanogr, 1–13. 10.1002/lno.11650.

698   75.   Coulson, T., Macnulty, D.R., Stahler, D.R., VonHoldt, B., Wayne, R.K., and Smith,

699   D.W. (2011). Modeling effects of environmental change on wolf population dynamics,

700   trait evolution, and life history. Science (1979) *334*, 1275–1278.

701   76.   Williams, J.L., Miller, T.E.X., and Ellner, S.P. (2012). Avoiding unintentional eviction

702   from integral projection models. Ecology *93*, 2008–2014.

703   77.   Veron, J.E.N., Stafford-Smith, M.G., Turak, E., and DeVantier, L.M. (2016). Corals of

704   the World. http://www.coralsoftheworld.org/page/home/.

705   78.   Madin, J.S., Baird, A.H., Baskett, M.L., Connolly, S.R., and Dornelas, M.A. (2020).

706   Partitioning colony size variation into growth and partial mortality. Biol Lett *16*, 1–5.

707   10.1098/rsbl.2019.0727.

708   79.   Madin, J.S., Anderson, K.D., Andreasen, M.H., Bridge, T.C., Cairns, S.D., Connolly,

709   S.R., Darling, E.S., Diaz, M., Falster, D.S., Franklin, E.C., et al. (2016). The Coral

710   Trait Database, a curated database of trait information for coral species from the global

711   oceans. Sci Data *3*, 1–21. 10.1038/sdata.2016.17.

712   80.   Hall, V.R., and Hughes, T.P. (1996). Reproductive strategies of modular organisms:

713   Comparative studies of reef-building corals. Ecology *77*, 950–963.

714   81.   Bramanti, L., Iannelli, M., Fan, T.Y., and Edmunds, P.J. (2015). Using demographic

715   models to project the effects of climate change on scleractinian corals: Pocillopora

716   damicornis as a case study. Coral Reefs *34*, 505–515. 10.1007/s00338-015-1269-z.

717   82.   Stott, I., Franco, M., Carslake, D., Townley, S., and Hodgson, D. (2010). Boom or

718   bust? A comparative analysis of transient population dynamics in plants. Journal of

719   Ecology *98*, 302–311. 10.1111/j.1365-2745.2009.01632.x.

720   83.   Hodgson, D., McDonald, J.L., and Hosken, D.J. (2015). What do you mean,

721   "resilient"? Trends Ecol Evol *30*, 503–506. 10.1016/j.tree.2015.06.010.

722   84.   Zuidema, P.A., Jongejans, E., Chien, P.D., During, H.J., and Schieving, F. (2010).

723   Integral projection models for trees: A new parameterization method and a validation

724   of model output. Journal of Ecology *98*, 345–355. 10.1111/j.1365-2745.2009.01626.x.

725   85.   Jones, O.R., Barks, P., Stott, I., James, T.D., Levin, S., Petry, W.K., Capdevila, P.,

726   Che-Castaldo, J., Jackson, J., Römer, G., et al. (2021). Rcompadre and Rage - two R

727   packages to facilitate the use of the COMPADRE and COMADRE databases and

728   calculation of life history traits from matrix population models. BioRxiv,

729   2021.04.26.441330.

730   86.   Townley, S., Carslake, D., Kellie-Smith, O., Mccarthy, D., and Hodgson, D. (2007).

731   Predicting transient amplification in perturbed ecological systems. Journal of Applied

732   Ecology *44*. 10.1111/j.1365-2664.2007.01333.x.

733   87.   Townley, S., and Hodgson, D.J. (2008). Erratum et addendum: Transient amplification

734   and attenuation in stage-structured population dynamics. Journal of Applied Ecology

735   *45*, 1836–1839. 10.1111/j.1365-2664.2008.01562.x.

736   88.   Kreiss, H.O. (1962). Über Die Stabilitätsdefinition Für Differenzengleichungen Die

737   Partielle Differentialgleichungen Approximieren. BIT *2*, 153–181.

738   10.1007/BF01957330.

739   89.   Stott, I., Hodgson, D.J., and Townley, S. (2012). Popdemo: An R package for

740   population demography using projection matrix analysis. Methods Ecol Evol *3*, 797–

741   802. 10.1111/j.2041-210X.2012.00222.x.

742   90.   Carrascal, L.M., Galván, I., and Gordo, O. (2009). Partial least squares regression as

743   an alternative to current regression methods used in ecology. Oikos *118*, 681–690.

744   10.1111/j.1600-0706.2008.16881.x.

745   91.   Rayner, N.A., Parker, D.E., Horton, E.B., Folland, C.K., Alexander, L. V., Rowell,

746   D.P., Kent, E.C., and Kaplan, A. (2003). Global analyses of sea surface temperature,

747   sea ice, and night marine air temperature since the late nineteenth century. J Geophys

748   Res *108*, 1–37. 10.1029/2002jd002670.

749   92.   Greenman, J. V., and Benton, T.G. (2005). The frequency spectrum of structured

750   discrete time population models : its properties and their ecological implications.

751   Oikos *110*, 369–389.

752   93.   Vasseur, D.A., and Yodzis, P. (2004). The color of environmental noise. Ecology *85*,

753   1146–1152.

94.   R Core Team (2019). R: A language and environment for statistical computing.

95.   Sanchez, G. (2012). plsdepot: Partial least squares (PLS) data analysis methods.

96.   Legendre, P. (2018). lmodel2: Model II regression.

Tables and Figures

Figure 1. Using repeated annual surveys of tagged individual colonies, conducted

between 2016 and 2019, we quantified the influence of environmental stochasticity on

the long-term performance and transient potential of tropical and subtropical coral

populations in southern Japan and eastern Australia. **(A)** As climate shifts induce range

expansions in many coral species worldwide, their populations are increasingly exposed to a

gradient in thermal regimes, illustrated here by mean monthly sea surface temperatures

($\bar{x}_{sst}$; °C) recorded between 1950 and 2019[91]. **(B)** Between 2016 and 2019, we documented

the survival, growth, fragmentation, and recruitment patterns of 3171 tagged coral individuals

within the tropical reef communities (▲) of Okinawa (Japan) and Heron Island (Australia),

and within the subtropical coral communities (●) of Kochi (Japan) and the Solitary Islands

Marine Park (Australia). **(C)** Using this data, we parameterised Integral Projection Models

(IPMs) describing the dynamics of tropical and subtropical assemblages of competitive,

stress-tolerant, and weedy coral taxa. Combining outputs obtained from these models with

measures of the thermal regimes experienced by each population we then explored the

relationships between the long-term performance and transient (short-term) potential of coral

populations, and their thermal exposure.

Figure 2. Contrasting patterns in long-term performance and transient potential across

our examined coral populations, corresponding with their relative exposure to thermal

variability. Partial least squares regression score plot illustrating the association between

thermal conditions, and the long-term performance (λ) and transient potential (transient

envelope [*TE*] & damping ratio [ρ]) of tropical (▲) and subtropical (●) populations of

competitive (blue), stress-tolerant (yellow), and weedy (red) coral taxa. To quantify the

thermal conditions experienced by each coral population, we used sea surface temperatures

(SST) recorded between 1950 and 2019 to calculate regional estimates of mean monthly SST

($x\bar{}_{sst}$), monthly SST variance (cv_{sst}), and monthly SST frequency spectrum (β_{sst}). Component

scores illustrate the relative degree of variance explained in the thermal predictor variables,

whilst $r^2_{[y]}$ reflects the cumulative variance explained across the demographic characteristics.

The shaded polygons reflect the clustering of tropical and subtropical populations, whilst the

dotted lines delineate regions of association to facilitate the visualisation of patterns in

correlation between the abiotic and demographic variables.

Figure 3. Inter-specific variation within the contrasting patterns observed between long-term performance and transient potential across tropical and subtropical correlates with patterns in population turnover rate (A) Interaction plot showcasing how estimates of demographic stability index (DSI) vary between associated tropical (▲) and subtropical (●) populations of competitive (blue), stress-tolerant (yellow), and weedy (red) coral taxa in Australia and Japan. We present DSI, as an inverse measure of maximal amplification ($\bar{\rho}_{max}$), describing the ability for populations to undergo elevated growth following disturbance. Thus, lower DSI estimates correspond with enhanced amplification capacity. We also applied Type 2 linear regression to separately explore the association of population turnover characteristics with **(B)** long-term performance (asymptotic population growth rate; λ), and **(C)** transient potential (transient envelope, *TE*) across tropical and subtropical populations of competitive, stress-tolerant, and weedy coral taxa in Australia and Japan. We note here that transient envelope estimates were reversed during transformation to achieve normality, thus higher values reflected diminished transient potential. We have therefore displayed transient potential on a reversed scale to facilitate comparisons with patterns in

805   long-term performance (λ). We used generation time (Years; displayed here on the log scale)

806   as a measure of population turnover rate, with higher estimates reflecting slower rates of

807   population turnover. Across panels B and C r^2 values are provided as measure of model fit.

808   Across all panels error is displayed using 95% CI.

814   **Figure 1**

815   **Figure 2**

826   **Figure 3**

832   **Table 1.** Population growth rates (λ) obtained from corresponding tropical and subtropical

833   populations of competitive, stress-tolerant, and weedy coral taxa in Australia and Japan.

834   Shading used to highlight the highest estimate of population growth across each tropical-

835   subtropical pairing. Error displayed using 95% CI.

Country	Life-history group	Tropical	Subtropical
Australia	Competitive	0.983 [0.981, 0.984]	0.958 [0.957, 0.959]
	Stress-tolerant	0.983 [0.980, 0.985]	0.899 [0.898, 0.899]
	Weedy	0.981 [0.980, 0.982]	0.686 [0.684, 0.687]
Japan	Competitive	1.001 [0.999, 1.004]	0.640 [0.639, 0.641]
	Stress-tolerant	0.913 [0.909, 0.917]	0.885 [0.877, 0.894]
	Weedy	0.760 [0.750, 0.770]	0.807 [0.802, 0.812]

846